Title: Evaluating the Roles of Drift and Selection in Trait Loss along an Elevational Gradient

Authors: Sophia F. Buysse[1,2,3], Samuel G. Pérez[1,2,3], Joshua R. Puzey[4], Ava Garrison[1,2,3], Gideon S. Bradburd[5],

Christopher G. Oakley[6], Stephen J. Tonsor[7], F. Xavier Picó[8], Emily B. Josephs[1,2,9]*, Jeffrey K Conner[1,2,3]*

Affiliations

[1]Plant Biology Department, Michigan State University, East Lansing, MI 48824, USA

[2]Ecology, Evolution, and Behavior Program, Michigan State University, East Lansing, MI 48824, USA

[3]Kellogg Biological Station, Hickory Corners, MI 49060, USA

[4]Department of Biology, William & Mary, Williamsburg, VA 23187, USA

[5]Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI 48109, USA

[6]Department of Botany and Plant Pathology, and the Center for Plant Biology, Purdue University, West

Lafayette, IN 47907, USA

[7]Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA 15260 (deceased)

[8]Departamento de Ecología y Evolución, Estación Biológica de Doñana, Consejo Superior de

Investigaciones Científicas, Sevilla 41092, Spain

[9]Plant Resilience Institute, Michigan State University, East Lansing, MI 48824, USA

*Co-Corresponding Authors:

Jeff Conner

3700 E. Gull Lake Drive

Hickory Corners, MI 49060

connerj@msu.edu

Emily Josephs

612 Wilson Rd, Rm 166

East Lansing MI 48824 USA

Josep993@msu.edu

Author Contributions: SGP collected the phenotypic data and conceived the research with JKC. JRP

oversaw the sequencing. AG and GSB did preliminary analyses and contributed to conceptual

development. CGO, SJT and FXP provided seed material and guidance in project development. SFB

analyzed the data and drafted the manuscript with guidance from EBJ and JKC. JKC managed the project.

All authors provided comments on the manuscript.

Funding: Research reported was supported by the National Institute of General Medical Sciences of the

National Institutes of Health (NIH) under award R35GM137919 (awarded to GSB) and R35GM142829

(awarded to EBJ), and by National Science Foundation awards DEB 0919452 (to JKC) and DEB 2223962

(to JKC and GSB). The content is solely the responsibility of the authors and does not necessarily

represent the official views of the NIH or NSF.

Data Availability: These sequence data will be submitted to the GenBank database upon manuscript

acceptance. All phenotypic data and code will be made available on GitHub upon acceptance. GWAS

output files will be made available on Dryad upon acceptance.

Conflict of Interest: The authors declare no conflicts of interest.

Acknowledgements:

We thank D. Schemske for guidance throughout the initial stages of the project, N. Zhang for library

preparation, and A. Platts for bioinformatics assistance. This manuscript was improved by comments

from Conner lab and Josephs lab members. This work is not possible without the foundational work of

Dr. Steven Tonsor; all authors agree on his inclusion in this manuscript and JKC accepts all responsibility

for his contribution. We dedicate this paper to his memory.

Abstract

Traits that have lost function sometimes persist through evolutionary time. These traits may

persist if there is not enough standing genetic variation for the trait to allow a response to

selection, if selection against the trait is weak relative to drift, or if the trait has a residual

function. To determine the evolutionary processes shaping whether nonfunctional traits are

retained or lost, we investigated short stamens in 16 populations of *Arabidopsis thaliana* along

an elevational cline in northeast Spain. We found a cline in short stamen number from retention

of short stamens in high elevation populations to incomplete loss in low elevation populations.

We did not find evidence that limited genetic variation constrains the loss of short stamens at

high elevations, nor evidence for divergent selection on short stamens between high and low

elevations. Finally, we identified loci associated with short stamens in northeast Spain that are

different from loci associated with variation in short stamen number across latitudes from a

previous study. Overall, we did not identify the evolutionary mechanisms contributing to an

elevational cline in short stamen number but did identify different genetic loci underlying

variation in short stamen along similar phenotypic clines.

Keywords (provide 3 to 6): – genetic drift, trait loss, natural selection, effective population size,

stamen, elevational cline

Teaser text

The evolutionary mechanisms underlying loss or retention of traits that have lost function are

poorly understood. Short stamens in *Arabidopsis thaliana* provide a compelling system to

investigate the roles of genetic drift and selection in trait loss across latitudinal and elevational

clines. This study investigates how drift and selection shape short stamen loss in 16 populations

of *A. thaliana* along an elevational gradient in Northeast Spain. An investigation of the loci

underlying variation in short stamen number suggests variants in different genes may cause trait

loss in similar phenotypic clines within a species.

Introduction

Traits that have lost function are often lost through evolutionary time, yet some persist.
Nonfunctional traits could be lost through direct selection against the trait (Dorken et al., 2004;
Lahti et al., 2009), correlated responses to selection on other traits caused by pleiotropy or
linkage disequilibrium (Yoshizawa et al., 2012), or the accumulation of selectively neutral
mutations (Fong et al., 1995). In contrast, nonfunctional traits may be retained by evolutionary
constraint if the nonfunctional trait is genetically correlated with a different, functional trait
(Lande, 1979; Walsh & Blows, 2009). Nonfunctional trait loss could also be prevented if there is
not enough standing genetic variation for selection to act on the trait (Lahti et al., 2009) or if
weak selection against the trait is unable to overcome drift (Charlesworth, 2009). Both processes
may be pronounced in species with small effective population sizes. Determining the
evolutionary processes shaping whether nonfunctional traits are retained or lost will have broad
implications for our understanding of the interplay of direct selection, correlated responses to
selection, and genetic drift (reviewed in Futuyma, 2010).

Here, we use an elevational cline in short stamen number in *A. thaliana* from Northeast Spain to
investigate how evolutionary processes interact to shape variation in a trait that has lost function.
Flowers across almost all of the 3,700 species in the Brassicaceae family have four long and two
short stamens (Zomlefer, 1994). This tetradynamous stamen arrangement is strongly
phylogenetically conserved. The function of short stamens in Brassicaceae is unknown,
especially in self-pollinating species. In outcrossing species, the presence of short stamens
increased the duration of pollinator visits (Kudo, 2003) and short stamen anthers produce more
pollen though less of it is removed than pollen from long stamen anthers during pollinator visits
(Conner et al., 1995). There is also selection to maintain both long and short stamens in *R.*

raphanistrum, an outcrossing species (Conner et al., 2003; Waterman et al., 2023), but the

function of short stamens even in outcrossing species is unknown. Loss of short stamens,

resulting in flowers with four long stamens and either one or zero short stamens (Fig. 1A), has

been observed in self-pollinating *Arabidopsis thaliana* and a few other Brassicaceae genera

(Bowman et al., 1999; Matsuhashi et al., 2012; Müller, 1961) but is not common across the

family and has not been observed in *Arabidopsis lyrata* or *Capsella bursa-pastoris*, close

relatives of *A. thaliana*. See Royer et al. (2016) for a review of stamen number variation.

A. thaliana evolved to be almost entirely self-fertilizing from an outcrossing ancestor between

0.5 – 1 million years ago (Durvasula et al., 2017; Tang et al., 2007). In flowering plants, a

transition from outcrossing to self-pollination and subsequent relaxation of selection for

pollination often results in a suite of trait changes, called the 'selfing syndrome', that includes

smaller flowers and reduced distance between the anthers and stigma (Sicard & Lenhard, 2011).

Self-pollination also decreases local effective population size (Caballero, 1994), increasing the

effects of genetic drift and thus reducing within-population genetic variation.

Because short stamens in *A. thaliana* are below the stigma, we expect them to have little or no

function in self-pollination. Prior work confirmed these expectations: short stamens in *A.*

thaliana do not contribute significantly to self-fertilized seed number and many populations in

the native range have flowers with zero or one short stamens (Royer et al., 2016). Almost all

flowers have four long stamens, and the short stamens are not reduced but rather lost entirely.

Fixation for zero short stamens has not been observed in all *A. thaliana* natural populations

studied to date (Royer et al., 2016). However, variation between populations is common. For

example, southern European populations are more likely to lose short stamens while northern

European populations typically retain both short stamens. Royer et al. (2016) identified three

QTL underlying the variation in short stamen number across Europe.

Short stamen loss in *A. thaliana* is an excellent model to study the evolutionary mechanisms

influencing variation in a trait that has lost most or all function. Because northern European

populations of *A. thaliana* have undergone repeated bottlenecks that reduced effective population

size, and thus standing genetic variation (Beck et al., 2008; François et al., 2008; Lewandowska-

Sabat et al., 2010), short stamens may be retained in these populations because there is

insufficient genetic variation for stamen number and/or weak selection against short stamens

cannot overcome drift. Alternatively, the latitudinal cline of the European populations could

result from local adaptation for short stamen number or another correlated trait.

There are many evolutionary scenarios that could result in phenotypic clines, but the same

evolutionary processes occurring across latitudes may shape elevational clines in short stamen

number. High elevations were glaciated, even at southern latitudes (Hughes et al., 2006) and

while Northeast Spain does include putative refugia regions (Médail & Diadema, 2009), prior

work in Northeast Spain has shown high elevation montane populations have less genetic

variation and smaller effective population sizes than lower elevation coastal populations (Gomaa

et al., 2011; Montesinos et al., 2009). Further, because environmental factors vary similarly with

latitude and elevation, selection may result in similar phenotypic clines. Yet, latitudinal and

elevational phenotypic clines are not always parallel (Daco et al., 2021; Kooyers et al., 2015).

Even if the same evolutionary processes are resulting in parallel phenotypic clines, they may not

have a parallel genetic basis (Fulgione et al., 2022; Gamba et al., 2024). If drift contributes to the

phenotypic clines, a parallel genetic basis is unlikely as drift acts equally and randomly across

the genome and any gene in the short stamen developmental pathway could be impacted. If

selection contributes to the phenotypic clines, a parallel genetic basis is still unlikely to result

from standing genetic variation for short stamens in *A. thaliana* due to small effective population

sizes, no evidence for strong selection on short stamens, and the multiple loci involved in the

latitudinal cline (MacPherson & Nuismer, 2017). Thus, we do not expect a parallel genetic basis

for short stamen loss across latitude and across elevation despite a strong expectation for parallel

phenotypic clines.

To understand the processes shaping variation in short stamen loss, we assessed 16 populations

of *A. thaliana* along an elevational gradient in Northeast Spain. We counted short stamen number

on plants grown in a growth-chamber common garden and sequenced multiple individuals per

population to quantify genetic variation. We identified an elevational cline in short stamen

number in Northeast Spain, with more stamen loss at low elevation, similar to the previously

described latitudinal cline (Royer et al., 2016). We then used multiple regression, polygenic

adaptation detection, and genome wide association studies to address possible evolutionary

mechanisms contributing to the cline:

1) Does variation in effective population size contribute to the short stamen loss cline?

Less stamen loss in populations with less genetic variation after accounting for

variation due to elevation would be consistent with the hypothesis that stamen loss is

constrained by a lack of genetic variation and less effective selection.

2) Does local adaptation contribute to the short stamen loss cline? Evidence for

divergent selection on short stamen number between high and low elevation

populations would support local adaptation contributing to the cline.

3) Are the same loci associated with variation in short stamen number across elevation

as across latitude? Identifying loci within the three previously identified QTL (Royer

et al., 2016) would support parallel genomic evolution between the latitudinal and

elevational clines in short stamen number. Alternatively, we may detect different

large-effect loci or small effect loci, many of which will be below our detection

threshold.

Materials and Methods

Seed Collection

Population collection sites span a continuous elevation gradient from 61m to 1706m above sea

level along the Mediterranean coast and into the Pyrenees (Fig. 1B; Fig. 2-3 legend; Table S1).

Seeds from five to nine individuals were haphazardly collected from multiple patches in each of

16 populations ($N = 112$ genotypes) in northeast Spain (Montesinos-Navarro et al., 2011). These

lineages are also used in Montesinos-Navarro et al. (2011), Montesinos-Navarro et al. (2012),

and Wolfe and Tonsor (2014). They include the 10 populations from Montesinos et al. (2009)

and the 9 populations from Gomaa et al. (2011).

Phenotyping

To assess genetic differentiation for short stamen number, plants were grown in growth-chamber

common gardens in three blocks at Michigan State University, East Lansing, MI. Seeds were

stratified at 4°C for five days before we increased temperatures to 22°C:18°C for 16-hour days at

a constant 60% humidity for four weeks. After emergence, plants were thinned to one seedling

per cell in a 200 cell plug tray. One to four plants from each of the 112 genotypes (median = 2;

$N = 230$ plants) were vernalized at 4°C with 10-hour days for six weeks before returning to

22°C:18°C for 16-hour days through flowering.

On each plant, we counted stamen number (Fig. 1A) on up to three flowers at each of up to three

timepoints throughout the duration of flowering to estimate short stamen number (1 to 9 flowers

per plant; median = 6; $N = 1,423$ flowers). Short stamen number varied among flowers within all

but three of the 58.2% of plants with stamen loss; the average standard error of short stamen

number within individuals was 0.16. Population mean short stamen number was calculated as the

arithmetic mean from all flowers phenotyped in a population (42 to 169 flowers per population;

mean = 88.9; median = 99.5; average standard error within population = 0.06). The arithmetic

population means were highly correlated with estimated marginal means for each population

calculated with the *emmeans* package (Lenth, 2021) in R v4.2.2 (R Core Team, 2021) from a

model with population as a fixed effect, random effects of flowering timepoint nested within

plant nested within genotype which is nested within population, and random effect for block ($r =$

0.975; $p < 0.001$).

We sequenced the entire genomes of a subset of 61 genotypes (see below). The 61 genotypes

include representatives from all 16 populations (3 or 4 different genotypes per population); we

elected not to sequence all genotypes as prior work indicated low within-population variation

(Gomaa et al., 2011; Montesinos et al., 2009). In the sequenced genotypes, stamen number was

counted on up to four plants per genotype (median = 2; $N = 141$) and up to nine flowers per plant

(median = 7; $N = 971$). For all analyses that incorporate both phenotypic and SNP information,

arithmetic population means were recalculated as the mean from flowers scored for short stamen

number from sequenced genotypes (18 to 118 flowers per population; median = 63). The

population means from all flowers and the population means from only the sequenced genotypes

are highly correlated ($r = 0.985$; $p < 0.001$) indicating the sequenced genotypes are

representative of all plants scored for short stamen number.

Testing for an elevational cline

We calculated both linear and quadratic regressions of short stamen number on elevation to

identify if there is an elevational cline in short stamen number parallel to the latitudinal cline

(Table S3). The quadratic regression was included because it increased the adjusted R^2

substantially even though the quadratic term was not significant.

Sequencing

A subset of 61 genotypes (3-4 per population) were chosen for Ilumina paired-end whole-

genome 150bp sequencing. Nextera adapter sequences were trimmed from the raw sequence data

with Trim Galore (Krueger, 2019). We also clipped the first 15 bp of each read because quality

checks with FastQC (Andrews, 2019) and MultiQC (Ewels et al., 2016) showed an identical,

unusual, pattern in the first 10 bases of each read: GTTTTAAACT. Reads were mapped to the

TAIR10 reference genome (Berardini et al., 2015) using BWA mem with default settings (Li &

Durbin, 2009). The mean mapping rate for properly paired reads was 96.6% with a mean of

15,224,790 properly paired and mapped reads per genotype and a total of 943,936,960 mapped

and paired reads in the dataset. The median depth across genotypes was low but acceptable at

8X. There is variation in coverage, missing data, and quality scores between genotypes, but no

genotypes were excluded on this basis in order to maximize sample size (Table S2). Duplicate

reads were marked with the Genome Analysis Toolkit (GATK) v4.1.4.1 (Van der Auwera &

O'Connor, 2020) MarkDuplicates Spark. After an initial round of Haplocaller and

GenotypeGVCF, the dataset was filtered with the parameters suggested by GATK best practices

(Van der Auwera & O'Connor, 2020): QD <2, FS > 60, MQ < 40, ReadPosRankSum < -8, and

MQRankSum < -12.5. The filtered file was used as a known variants file for base quality score

recalibration (BQSR). Variants were then called again with HaplotypeCaller using the GVCF

flag to keep all sites before combining samples for GenotypeGVCF with the all-sites flag to

create a dataset that includes both variant and invariant sites (116,855,685 total sites).

Testing the drift hypotheses

To estimate the effects of past drift and gene flow in each population, genome-wide pairwise

nucleotide diversity (pi) was calculated for each population as an approximate measure of

effective population size with pixy v1.0.4 (Korunes & Samuk, 2021) from a filtered dataset

containing variant and invariant sites based on the pixy protocol. We used quality filters to

remove all indels and low-quality sites that met the following criteria: quality score less than 20,

mean depth less than 3, and more than 25% missing data. The filtered dataset contains

99,202,614 sites. A peak in nucleotide diversity was observed in the centromere of each

chromosome, likely caused by fewer mapped sites (Korunes and Samuk, *pers. com.*). However,

excluding previously published centromeres (Clark et al., 2007) did not meaningfully change

genome-wide nucleotide diversity ($r = 0.998$; $p < 0.001$), so centromeres are included in all of the

following analyses (results excluding the centromeres can be found in Figs. S5-S7).

To test if variation in effective population size is contributing to a cline in short stamen number,

we used multiple regression to identify how short stamen number was predicted by nucleotide

diversity and elevation. While elevation is correlated with climatic variables such as precipitation

and temperature (Montesinos-Navarro et al., 2011) that could be selective agents for plant traits,

we used elevation because previous work in these same 16 populations demonstrated that

elevation explained 54% of the variance in trait principal component 1 (traits include: phenology,

water use efficiency, instantaneous CO_2 and H_2O exchange, PSII quantum efficiency, and

specific leaf area) while the climate PC1 only explained 36% of the trait variation (Wolfe &

Tonsor, 2014). We then visualized these results by using the residuals of single regression

models. The visualized results underestimate the confidence interval because they do not account

for uncertainty in calculating the residuals in the single regression models; the reported statistics

are from the full model. We used *lme4* in R for all models (Bates et al., 2015).

A negative relationship between nucleotide diversity and short stamen number after correcting

for elevation would support the hypothesis that smaller effective population sizes are

constraining trait loss due to low standing variation and less effective selection. A relationship

between short stamen number and elevation after correcting for nucleotide diversity could be

evidence for divergent selection because it indicates the short stamen number cline persists

beyond the variation explained by nucleotide diversity. However, a relationship between short

stamen number and elevation could also result from shared evolutionary history, sometimes

called shared genetic drift, if relatedness among populations is associated with elevation

(Colautti & Lau, 2015). We cannot distinguish divergent selection from relatedness in the

multiple regression analysis. Therefore, we further investigated evidence for divergent selection

by testing for local adaptation in short stamen number.

Testing for local adaptation

We used plink v1.9 (Chang et al., 2015) to filter the all-sites output from GATK to a "variant

sites only" dataset for conducting genetic principal component analysis (PCA). The "variant sites

only" dataset was also used for a genome-wide association study (GWAS; see below). We used

quality filters to remove all indels and non-biallelic sites in addition to low quality sites that met

the following criteria: minor allele frequency less than 5%, quality score less than 25, mean

depth less than 5, and more than 25% missing data. The variant sites dataset contains 1,858,706

SNPs. Filtering parameters were chosen to maintain the same percent nonsynonymous sites,

annotated by snpEff (Cingolani et al., 2012), as stricter parameters that removed sites with

greater than 15% missing data and a quality score less than 60 while retaining more variants in

the dataset. Genetic PCA was calculated for the first 20 PCs with plink v1.9 (Chang et al., 2015)

to characterize genetic relatedness within, and differentiation between, populations. We looked

for correlations between the population average PC value and elevation for each of the first 4

PCs to identify population structure associated with elevation.

We tested for divergent selection across populations on short stamen number, i.e., selection for

more stamens at high elevation and/or fewer stamens at low elevation than expected due to

genetic drift, with Qpc using the *quaint* R package (Josephs et al., 2019). Qpc is an extension of

Qst-Fst analysis that uses genetic PCs to estimate additive genetic variance within and between

populations (Josephs et al., 2019), and tests for phenotypic differentiation due to selection

beyond that expected from neutral evolution. This differs from the multiple regression test for

the drift hypothesis described above because Qpc explicitly considers among population

differentiation. Qpc incorporates genetic relatedness within and among populations through a

kinship matrix, rather than a single measure of population differentiation as in Qst-Fst. Qpc then

tests for excess phenotypic divergence along major axes of genetic relatedness (principal

components of the kinship matrix) rather than excess phenotypic divergence between

populations as in Qst-Fst (Josephs et al., 2019). The input kinship matrix was generated from a

random subset of 50,000 SNPs that had no missing data. This kinship matrix is slightly different

from the kinship matrix used for genetic PCA because Qpc requires a kinship matrix

standardized across all loci, not each locus individually as in plink, and *quaint* is not capable of

dealing with missing data.

Using genome wide association studies (GWAS) to identify loci associated with short stamen loss

We performed GWAS to find genomic regions associated with short stamen number. Mean short

stamen number was not normally distributed (Shapiro-Wilk $w = 0.85$; $p = 3.27\text{x}10^{-6}$; Fig. S1).

Arcsine transformation made the distribution closer to normal but still skewed (Shapiro-Wilk w

$= 0.91$; $p = 2.72\text{x}10^{-4}$; Fig. S1). Phenotypic and genotypic data were merged in plink v1.9 (Chang

et al., 2015). GWAS was conducted in gemma v0.98.4 with the Wald hypothesis test for each

SNP against the alternate allele and a centered kinship matrix to account for population structure

(Zhou & Stephens, 2012). The output was visualized with the *qqman* (Turner, 2018) and *ggplot2*

(Wickham et al., 2019) R packages. SNPs were further investigated if they passed a significance

threshold specified using a false discovery rate (FDR) < 0.05 determined with p.adjust in R.

Because the arcsine-transformed stamen number still showed some skew, we conducted GWAS

using additional strategies for quantifying short stamen number. We ran a GWAS with

untransformed mean short stamen number and another with mean short stamen number coded as

a binary trait. In the latter, genotypes with no short stamen loss (mean short stamen number $= 2$)

were coded as controls and genotypes with any amount of short stamen loss (mean short stamen

number < 2) were coded as cases. Finally, a fourth GWAS was conducted on the subset of

genotypes that experience short stamen loss (i.e., GWAS on only the genotypes with mean short

stamen number < 2; Shapiro-Wilk $w = 0.93$; $p = 0.0132$) to ameliorate the zero-inflated-like

distribution in the other continuous GWAS caused by excess genotypes with a mean short

stamen number equal to 2 (Fig. S1). In total we ran four GWAS on different ways of quantifying

short stamen number: arcsine transformed, untransformed, binary coded, and the subset with

short stamen number less than two.

We identified genetic regions with SNPs associated with short stamen number in at least two

GWAS to find candidate regions for further study ("shared" SNPs). We identified overlapping

regions associated with stamen loss among the four GWAS analyses by creating a 1kb window

centered on each SNP that had an FDR adjusted p-value below 0.10 in any GWAS and searching

for SNPs within that window that were also below an FDR adjusted p-value of 0.10 in at least

one other GWAS. We chose a more lenient FDR of 0.10 for this analysis because we have higher

confidence SNPs are associated with short stamen number if they pass a significance threshold in

multiple analyses. We chose a window of 1kb to include regions flanking the associated SNP,

although we recognize that *A. thaliana* genes can be larger than 1kb and that linkage

disequilibrium (LD) begins to drop off around 50kb in our individuals (Fig. S2), so this is a

conservative overlap criterion. SNPs in overlapping stamen loss associated regions were

investigated with the TAIR10 genome browser.

All figures were created with *ggplot2* (Wickham et al., 2019) and *ggpubr* (Kassambara, 2023)

unless otherwise noted.

Results

Short stamen loss is more common at low elevation

Mean short stamen number, when measured in a common garden, increases with the elevation of

the source population until reaching close to two stamens in plants from ~1300m (quadratic

regression: $\beta = 4.84 \times 10^{-4}$; $p = 0.004$; $\gamma = -4.73 \times 10^{-7}$; $p = 0.137$; Fig. 2A; Table S3). Thus, short

stamen retention is more common in populations from higher elevations and loss is more

common at low elevations in northeast Spain. This elevational cline is similar to the previously

identified latitudinal cline where short stamen loss is more common at southern latitudes (Royer

et al., 2016).

Nucleotide diversity decreases with elevation, but there is no evidence that variation in effective population size contributes to the cline in short stamen number

Consistent with expectations that *A. thaliana* in the Pyrenees experienced repeated founder effects after the last glaciation, high elevation populations have less nucleotide diversity than low elevation populations ($\beta = -1.81\text{x}10^{-6}$; $p < 0.001$; Fig. 2B; Table S3). The three populations with the lowest nucleotide diversity are high elevation populations BIS, VIE, and PAN. The results are consistent with prior findings that genetic diversity and effective population size decrease with elevation in this region (Gomaa et al., 2011; Montesinos et al., 2009). Population nucleotide diversity is comparable to nucleotide diversity of *A. thaliana* across the Iberian Peninsula and the European range (The 1001 Genomes Consortium, 2016).

We hypothesized a negative relationship between nucleotide diversity and short stamen number if variation in effective population size was contributing to the short stamen cline, but nucleotide diversity did not significantly predict short stamen number when accounting for elevation ($\beta = -138.5$; $p = 0.167$; Fig. 2D; Table S3). Further, the relationship between short stamen number and elevation is similar whether nucleotide diversity is included or not (Fig. 2C, compare to Fig. 2A; Table S3). These results suggest that variation in effective population size is not contributing significantly to the short stamen number cline. Rather, local adaptation, correlated responses to selection, or shared evolutionary history may be contributing to the cline.

There is no evidence for divergent selection on short stamen number along the elevational cline

An elevational cline in short stamen number could result from local adaptation to different optima at high and low elevations. To demonstrate evidence of local adaptation resulting from divergent selection, we would need to show that the cline in short stamen number is stronger

than what could be generated by neutral evolution alone. Genetic structure, here measured using

genetic PCA, shows a strong pattern of genetic differentiation by elevation; both PC1 ($r = -0.75$;

$p = 9.22\text{x}10^{-4}$) and PC2 ($r = -0.54$; $p = 0.03$) are correlated with elevation (Fig. 3A). Neither PC3

nor PC4 are correlated with elevation (Fig. S3). The BOS population is an outlier along both PC1

and PC2 (Fig. 3A). This aligns with geographic information because BOS is located on a

northern face of the Spanish Pyrenees while the other populations are on southern faces, and

prior work identified BOS in the Northwestern genetic cluster of the Iberian Peninsula while the

other 15 populations belong to the Northeast cluster or are classified as mixed (Castilla et al.,

2020). The PCA was conducted a second time after removing the BOS population (Fig. 3B).

These results continue to show an elevational cline along PC1 ($r = -0.92$; $p = 1.15\text{x}10^{-6}$), though

PC2 ($r = -0.17$; $p = 0.55$) separates the four highest elevation populations from each other with

the closest clustering of PAN and VIE. As before, neither PC3 nor PC4 are correlated with

elevation (Fig. S3).

The strong genetic differentiation across elevation we observed means that we have low power to

test for even greater differentiation in stamen loss as evidence for divergent selection. Thus it is

not very surprising that we did not find evidence for divergent selection on short stamen number

with Qpc. Differences in short stamen number between high and low elevation populations were

not larger than could be explained by neutral evolution in any of the first four PCs, which

together explain 40.2% of the genetic variation (Fig. 3C; $p = 0.257, 0.914, 0.802, 0.795$ in order).

Therefore, the Qpc analysis does not support local adaptation contributing to the cline, but we

cannot definitively rule out local adaptation because of the low power of our analysis.

Parallel clines in short stamen have different genetic bases

We conducted GWAS to identify loci associated with short stamen number. Because short

stamen number was not normally distributed (Fig. S1), we conducted GWAS with four strategies

for quantifying short stamen number: arcsine-transformed mean short stamen number,

untransformed mean short stamen number, a binary trait of whether or not individuals lacked any

short stamens, and an untransformed subset of only genotypes that lack some short stamens.

Each GWAS found different SNPs associated with the trait. The qqplot for the arcsine-

transformed GWAS shows most p-values close to the 1:1 line between Expected and Observed

(Fig. 4B) but the p-values in the binary GWAS deviate from the 1:1 line (Fig. S4B), indicating a

high false positive rate in identifying SNPS associated with any short stamen loss. In accordance

with this observation, the binary GWAS had 1,707 SNPs associated with short stamen loss at

FDR < 0.05 while the other analyses only had two or three (Figs. 4, S4). Only two SNPs are

associated with arcsine-transformed short stamen number (FDR < 0.05; Fig. 4). The SNP at

Chr3:2942726 is within a FAD/NAD(P)-binding oxidoreductase family protein (AT3G09580)

that localizes in the chloroplast (Tomizioli et al., 2014). The SNP at Chr5:13458838 is not within

a gene; the closest gene, *PICALM3* (AT5G35200), is just over 3kb away.

We identified top candidates for loci associated with short stamen number by identifying 1kb

windows that have SNPs associated with short stamen number in multiple GWAS ("shared"

SNPs). To do so, we calculated 1kb windows centered on all SNPs associated with short stamen

number (FDR < 0.10) and searched for other SNPs that fit the same criteria and fell within each

window. Twenty SNPs were associated with short stamen number in more than one GWAS from

9 different 1kb windows (Table S4). None of the SNPs fall within genes with known stamen

function. However, a shared SNP at Chr3:2253161 is approximately 40kb away, thus within LD

(Fig. S2), from *FHA2* (AT3G07220), a SMAD/FHA domain containing protein involved in

stamen development (Ahn et al., 2013; Gu et al., 2020). Mutating *FHA2* proteins can cause

plants to have fewer stamens, though flowers sometimes lose long stamens and sometimes lose

short stamens (Ahn et al., 2013). The two SNPs associated with arcsine-transformed short

stamen number (FDR < 0.05) are included in the shared SNPs (Fig. 4; Table S4).

All of the SNPs associated with variation in short stamen number fall outside the 95% credible

intervals of the previously identified QTL for short stamen number that used recombinant inbred

lines with parents from the extremes of the latitudinal short stamen loss gradient (Royer et al.,

2016). The closest intersection is on chromosome 5, where SNPs associated with short stamens

are approximately 2,000kb further into the chromosome than the previously identified short

stamen loss QTL. This is beyond the start of LD decay we estimated of 50kb (Fig. S2). Thus, the

parallel phenotypic clines in short stamen number across elevation and latitude are not parallel

genetically.

Discussion

In this study, we tested for an elevation cline in short stamen loss. We found that short stamen

number increased with elevation (Fig. 2A), similar to the cline of more short stamens at higher

latitudes observed by Royer et al. (2016). Short stamen number is one of many traits that show

an elevational cline in this region (Montesinos et al., 2009; Montesinos-Navarro et al., 2011;

Wolfe & Tonsor, 2014). Of these traits, days to bolting shows similar elevational and latitudinal

clines with delayed bolting at high elevations and northern latitudes (Montesinos-Navarro et al.,

2011; Stinchcombe et al., 2004). Thus, short stamen number joins a growing number of traits

with an elevational cline in northeast Spain and a much smaller set of traits with parallel clines in

elevation and latitude. These clines could be caused by effective population sizes because both

high elevation populations and northern latitude populations may have undergone repeated

bottlenecks that reduced effective population size since the last glaciation (Beck et al., 2008;

François et al., 2008; Hughes et al., 2006; Lewandowska-Sabat et al., 2010; Oakley et al., 2019).

While northeast Spain does include putative glacial refugia regions (Médail & Diadema, 2009),

none of the 16 populations analyzed here have been classified as relict lineages that predate the

last glaciation (Castilla et al., 2020). Selection could also contribute to similar latitudinal and

elevational clines because environmental factors such as temperature often vary in the same way

across latitude and elevation.

We then tested if the cline in short stamen loss was caused by variation in effective

population size if a combination of low genetic variation and less effective selection in

high elevation populations hinders a response to selection against short stamens. We

found variation in genetic diversity across elevations consistent with expectations from

repeated founder effects during range expansion after the last glaciation and/or low

gene flow (Fig. 2B). However, the multiple regression suggests that variation in genetic

diversity is not contributing to the short stamen number cline (Fig. 2C and 2D).

Instead, the multiple regression suggests that elevation can explain both variation in

genetic diversity and in stamen number. It is important to note that our phenotypic

clines show populations with low genetic diversity retain two short stamens, consistent

with our hypothesis. However, studies that identify a role of genetic drift in trait loss

have hypothesized the opposite, where populations with low genetic diversity have trait

loss caused by drift (Eckert et al., 1996; Lahti et al., 2009).

Next, we tested if local adaptation was contributing to the cline in short stamen loss using Qpc,

an extension of Qst-Fst (Josephs et al., 2019). We found no evidence of local adaptation

contributing to the cline in short stamen number (Fig. 3C). One caveat to our findings is that Qst-

Fst struggles to identify selection when there is large genetic differentiation between populations

or weak selection that results in similar values for Qst and Fst (Whitlock & Guillaume, 2009).

This is also potentially an issue for Qpc in this study as there is a great deal of genetic

differentiation between high and low elevation populations (Fig. 3A). However, the Qst-Fst

approach has identified local adaptation contributing to elevational clines in leaf succulence and

specific leaf area in 14 populations of *A. thaliana* from the Swiss Alps where there is also a great

deal of genetic differentiation (Luo et al., 2015). Qpc has also previously been used to identify

local adaptation in 249 natural *A. thaliana* accessions from across the European native range;

local adaptation was identified in initial size, growth rate at 16°C, and temperature response

(Clauw et al., 2022), so there is still potential to use Qpc to identify local adaptation in systems

with strong genetic differentiation. Ultimately, reciprocal transplant studies across elevations in

the field will provide the strongest evidence for or against local adaptation, but genomic

approaches like Qpc will still be necessary when reciprocal transplants are not possible or when

many populations are being studied as we do here.

Finally, we used GWAS to identify loci associated with short stamen number. We identified 20

SNPs associated with short stamen number in multiple GWAS (FDR < 0.10). These SNPs lie

outside the QTL for short stamens that were previously identified in a cross between a northern

European and southern European accession (Royer et al., 2016). The untransformed effect sizes

of our loci (Table S4) range from an absolute value of 0.23 to 0.33 and are much larger than the

effect sizes of the QTL identified in Royer et al. (2016), untransformed effect sizes range from

0.05 to 0.17, suggesting that we may be missing the previously identified QTL due to lower

power in the GWAS. Alternatively, the populations studied here may be fixed for the southern

allele identified in Royer et al (2016). However, our GWAS effect sizes may be inflated due to

winner's curse (Göring et al., 2001; Josephs et al., 2017) which causes effect sizes to be

overestimated in studies with small sample sizes such as ours (Capen et al., 1971). In addition,

the QTL identified in Royer et al. (2016) were epistatic, which could make them harder to detect

through standard GWAS approaches. Further, many of the loci underlying short stamen loss

likely have effect sizes too small to detect by the methods we used and by the previous QTL

analysis. For example, a QTL study in maize identified 50 QTL associated with a trait they

artificially selected for, yet these QTL explained only half the genetic divergence for the trait

indicating many other QTL were involved but not detected (Laurie et al., 2004). Overall, our

results suggest that the latitudinal cline in short stamen number across Europe has a different

genetic basis than the elevational cline in short stamen number observed in northeast Spain.

While the short stamen loss clines are similar between latitude and elevation, the lack of overlap

between the associated loci suggests that short stamen loss is caused by variants at different

genes in different geographic regions. These results are consistent with other findings that the

chromosome regions underlying the same phenotypic cline can differ by region. For example,

elevational clines of *A. thaliana* in different geographic regions show delayed flowering at higher

elevations but the genetic basis of this cline varies by global region (Gamba et al., 2024).

Additionally, parallel clines in flowering time in *A. thaliana* in the Cape Verde Islands arose

from mutations at different genes, *FRI* and *FLC* (Fulgione et al., 2022). Even in outcrossing

Arabidopsis species, functional parallel evolution is more common than genetic parallel

evolution (Bohutínská et al., 2021). The lack of genetic parallelism is not surprising but may

complicate future work to identify the developmental pathways resulting in loss of short stamens.

In conclusion, short stamen loss has occurred in low elevation populations more than in high

elevation populations. Our results suggest that retention of two short stamens in high elevation

populations is not explained by reduced effective population sizes or by local adaptation to

different optima in high and low elevation populations. Thus, the evolutionary mechanisms

underlying variation in short stamen number are excitingly complex and deserve further study. A

third hypothesis is that correlated responses to selection may contribute to the cline in short

stamen number (Futuyma, 2010; Lande, 1979). However, correlated responses to selection on

other locally adapted traits could result in weak evidence for divergent selection; we did not find

this. Future work should better characterize direct selection and correlated responses to selection

on short stamens in the field by measuring fitness of plants with natural variation in short stamen

number and experimental manipulation of stamens, like that done by Royer et al. (2016), in the

field. Further, refining the short stamen number QTL to candidate genes and comparing to

genetic regions underlying locally adapted traits could uncover the role of linked or pleiotropic

genes leading to correlated responses to selection. These additional studies across both latitudinal

and elevational gradients will characterize the interplay of direct selection, correlated responses

to selection, and genetic drift in trait loss and identify parallel evolutionary forces at play across

environmental contexts.

Figures:

Figure 1: Populations studied across an elevation gradient. A) Natural variation in short

stamen number. Flowers have 2 (left), 1 (center), or 1(right) short stamens. All flowers have four

long stamens. Photos by Frances Whalen. **B)** Each point represents the seed collection site for

16 populations of *A. thaliana* across northeast Spain. The map coloring indicates elevation from

green at low elevation to white at high elevation.

Alt text: Photos of three Arabidopsis flowers and a map showing seed collections sites in

Northeastern Spain.

Figure 2: Effective population size does not explain retention of short stamens at high elevation. A) Mean short stamen number shows a quadratic elevational cline. **B)** Elevation strongly predicts population mean pairwise nucleotide diversity, our measure of effective population size. **C)** The residuals of the model of short stamen number regressed on nucleotide diversity then regressed on elevation and **D)** the residuals of the model of short stamen number regressed on quadratic elevation then regressed on nucleotide diversity. C and D visually underestimate the confidence intervals because they do not include uncertainty in calculating the

residuals; the interpretation remains the same because statistics displayed on the figures are from

the full model ($r^2 = 0.50$; Table S3). In all panels, the color of each point represents the

population elevation, black lines are the regression, and grey lines are 95% confidence intervals.

Alt text: Graphs depicting relationships between short stamen number, elevation, and nucleotide

diversity. Short stamen number increases with elevation and the linear term is significant but the

quadratic term is not. Nucleotide diversity decreases with elevation. After accounting for

nucleotide diversity, short stamen number still increases with elevation. After accounting for

elevation, nucleotide diversity now increases with elevation though the slope is not significant.

Figure 3: Genetic variation is correlated with elevation, but there is no evidence for divergent selection on short stamen number. A) Genetic PCA for PC1 and PC2 from 1,858,706 SNPs. **B**) Genetic PCA for PC1 and PC2 with BOS individuals excluded. **C**) Qpc results showing the relationship between mean short stamen number and the greatest axis of genetic differentiation, PC1, (black line, *p*=0.26) is within expectations due to neutral evolution (grey shading), as were PC2-4 (see text). The fill color of each point represents the population elevation. Panels A and C used slightly different input data (see Methods).

547   **Alt text:** Graphs depicting genetic relatedness from genetic principal component analysis in

548   panels A and B. In both panels, principal component 1 explains 13 percent of genetic variation

549   and principal component 2 explains 10 percent. Panel C shows a positive relationship between

550   mean short stamen number and principal component 1 but it is within expectations due to neutral

551   evolution.

Figure 4. Few SNPs are associated with short stamen number. Manhattan plot (A) and QQ plot (B) for arcsine-transformed mean short stamen number. The yellow dashed line indicates significance at p=0.05 after Bonferroni correction. Blue points are significant below a FDR of 0.10. Points labeled with their chromosomal location are significant below a FDR of 0.05. Points with a green outline are shared between at least two short stamen GWAS below FDR 0.10 (n = 20). The black bars on the x axis are Bayes 95% credible intervals for short stamen number QTL identified by Royer et al. (2016).

Alt text: Graphs showing results of arcsine-transformed genome wide association study. The two labeled points in panel A are at chromosome 3 position 2942726 and on chromosome 5 at 13458838. There are eight points significant below a false discovery rate of 0.10, most of them are outlined in green. Half of the shared SNPs are not significant in this GWAS.

References

Ahn, E.-R., Cho, H.-K., & Pai, H.-S. (2013). The forkhead-associated domain 2 (FHA2) in Arabidopsis plays a role in plant fertility by regulating stamen development. *Planta*, *237*(4), 1015–1023. https://doi.org/10.1007/s00425-012-1815-7

Andrews, S. (2019). *FastQC* (Version 0.11.9) [Computer software]. Babraham Bioinformatics. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*(1), 1–48. https://doi.org/10.18637/JSS.V067.I01

Beck, J. B., Schmuths, H., & Schaal, B. A. (2008). Native range genetic variation in Arabidopsis thaliana is strongly geographically structured and reflects Pleistocene glacial dynamics. *Molecular Ecology*, *17*(3), 902–915. https://doi.org/10.1111/j.1365-294X.2007.03615.x

Berardini, T. Z., Reiser, L., Li, D., Mezheritsky, Y., Muller, R., Strait, E., & Huala, E. (2015). The arabidopsis information resource: Making and mining the "gold standard" annotated reference plant genome. *Genesis*, *53*(8), 474–485. https://doi.org/10.1002/dvg.22877

Bohutínská, M., Vlček, J., Yair, S., Laenen, B., Konečná, V., Fracassetti, M., Slotte, T., & Kolář, F. (2021). Genomic basis of parallel adaptation varies with divergence in Arabidopsis and its relatives. *Proceedings of the National Academy of Sciences*, *118*(21), e2022713118. https://doi.org/10.1073/pnas.2022713118

Bowman, J. L., Brüggemann, H., Lee, J., & Mummenhoff, K. (1999). Evolutionary Changes in Floral Structure within Lepidium L. (Brassicaceae). *International Journal of Plant Sciences*, *160*(5), 917–929. https://doi.org/10.1086/314194

Caballero, A. (1994). Developments in the prediction of effective population size. *Heredity*,

73(6), Article 6. https://doi.org/10.1038/hdy.1994.174

Capen, E. C., Clapp, R. V., & Campbell, W. M. (1971). Competitive Bidding in High-Risk

Situations. *Journal of Petroleum Technology*, *23*(06), 641–653.

https://doi.org/10.2118/2993-PA

Castilla, A. R., Méndez-Vigo, B., Marcer, A., Martínez-Minaya, J., Conesa, D., Picó, F. X., &

Alonso-Blanco, C. (2020). Ecological, genetic and evolutionary drivers of regional

genetic differentiation in Arabidopsis thaliana. *BMC Evolutionary Biology*, *20*(1).

https://doi.org/10.1186/s12862-020-01635-2

Chang, C. C., Chow, C. C., Tellier, L. C., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015).

Second-generation PLINK: rising to the challenge of larger and richer datasets.

GigaScience, *4*(1), 7. https://doi.org/10.1186/S13742-015-0047-8

Charlesworth, B. (2009). Effective population size and patterns of molecular evolution and

variation. *Nature Reviews Genetics*, *10*(3), 195–205. https://doi.org/10.1038/nrg2526

Cingolani, P., Platts, A., Wang, L. L., Coon, M., Nguyen, T., Wang, L., Land, S. J., Lu, X., &

Ruden, D. M. (2012). A program for annotating and predicting the effects of single

nucleotide polymorphisms, SnpEff. *Fly*, *6*(2), 80–92. https://doi.org/10.4161/fly.19695

Clark, R. M., Schweikert, G., Toomajian, C., Ossowski, S., Zeller, G., Shinn, P., Warthmann, N.,

Hu, T. T., Fu, G., Hinds, D. A., Chen, H., Frazer, K. A., Huson, D. H., Schölkopf, B.,

Nordborg, M., Rätsch, G., Ecker, J. R., & Weigel, D. (2007). Common sequence

polymorphisms shaping genetic diversity in Arabidopsis thaliana. *Science*, *317*(5836),

338–342. https://doi.org/10.1126/science.1138632

Clauw, P., Kerdaffrec, E., Gunis, J., Reichardt-Gomez, I., Nizhynska, V., Koemeda, S., Jez, J., &
    Nordborg, M. (2022). Locally adaptive temperature response of vegetative growth in
    Arabidopsis thaliana. *eLife*, *11*, e77913. https://doi.org/10.7554/eLife.77913

Colautti, R. I., & Lau, J. A. (2015). Contemporary evolution during invasion: Evidence for
    differentiation, natural selection, and local adaptation. *Molecular Ecology*, *24*(9), 1999–
    2017. https://doi.org/10.1111/mec.13162

Conner, J. K., Davis, R., & Rush, S. (1995). The effect of wild radish floral morphology on
    pollination efficiency by four taxa of pollinators. *Oecologia*, *104*(2), 234–245.
    https://doi.org/10.1007/BF00328588

Conner, J. K., Rice, A. M., Stewart, C., & Morgan, M. T. (2003). Patterns and mechanisms of
    selection on a family-diagnostic trait: Evidence from experimental manipulation and
    lifetime fitness selection gradients. *Evolution*, *57*(3), 480–486.
    https://doi.org/10.1111/j.0014-3820.2003.tb01539.x

Daco, L., Colling, G., & Matthies, D. (2021). Altitude and latitude have different effects on
    population characteristics of the widespread plant Anthyllis vulneraria. *Oecologia*,
    197(2), 537–549. https://doi.org/10.1007/s00442-021-05030-6

Dorken, M. E., Neville, K. J., & Eckert, C. G. (2004). Evolutionary vestigialization of sex in a
    clonal plant: Selection versus neutral mutation in geographically peripheral populations.
    Proceedings of the Royal Society B: Biological Sciences, *271*(1555), 2375–2380.
    https://doi.org/10.1098/rspb.2004.2875

Durvasula, A., Fulgione, A., Gutaker, R. M., Alacakaptan, S. I., Flood, P. J., Neto, C.,
    Tsuchimatsu, T., Burbano, H. A., Picó, F. X., Alonso-Blanco, C., & Hancock, A. M.
    (2017). African genomes illuminate the early history and transition to selfing in

Arabidopsis thaliana. Proceedings of the National Academy of Sciences, *114*(20), 5213–5218. https://doi.org/10.1073/pnas.1616736114

Eckert, C. G., Manicacci, D., & Barrett, S. C. H. (1996). Genetic Drift and Founder Effect in Native Versus Introduced Populations of an Invading Plant, Lythrum Salicaria (lythraceae). *Evolution*, *50*(4), 1512–1519. https://doi.org/10.1111/j.1558-5646.1996.tb03924.x

Ewels, P., Magnusson, M., Lundin, S., & Käller, M. (2016). MultiQC: Summarize analysis results for multiple tools and samples in a single report. *Bioinformatics*, *32*(19), 3047–3048. https://doi.org/10.1093/bioinformatics/btw354

Fong, D. W., Kane, T. C., & Culver, D. C. (1995). Vestigialization and loss of nonfunctional characters. *Annual Review of Ecology and Systematics*, *26*(1), 249–268. https://doi.org/10.1146/annurev.es.26.110195.001341

François, O., Blum, M. G. B., Jakobsson, M., & Rosenberg, N. A. (2008). Demographic History of European Populations of Arabidopsis thaliana. *PLOS Genetics*, *4*(5), e1000075. https://doi.org/10.1371/journal.pgen.1000075

Fulgione, A., Neto, C., Elfarargi, A. F., Tergemina, E., Ansari, S., Göktay, M., Dinis, H., Döring, N., Flood, P. J., Rodriguez-Pacheco, S., Walden, N., Koch, M. A., Roux, F., Hermisson, J., & Hancock, A. M. (2022). Parallel reduction in flowering time from de novo mutations enable evolutionary rescue in colonizing lineages. *Nature Communications*, *13*(1), 1461. https://doi.org/10.1038/s41467-022-28800-z

Futuyma, D. J. (2010). Evolutionary constraint and ecological consequences. *Evolution*, *64*(7), 1865–1884. https://doi.org/10.1111/J.1558-5646.2010.00960.X

Gamba, D., Lorts, C. M., Haile, A., Sahay, S., Lopez, L., Xia, T., Takou, M., Kulesza, E.,

Elango, D., Kerby, J., Yifru, M., Bulafu, C. E., Wondimu, T., Glowacka, K., & Lasky, J.

R. (2024). The genomics and physiology of abiotic stressors associated with global

elevational gradients in Arabidopsis thaliana. *New Phytologist*.

https://doi.org/10.1111/nph.20138

Gomaa, N. H., Montesinos-Navarro, A., Alonso-Blanco, C., & Picó, F. X. (2011). Temporal

variation in genetic diversity and effective population size of Mediterranean and

subalpine Arabidopsis thaliana populations. *Molecular Ecology*, *20*(17), 3540–3554.

https://doi.org/10.1111/j.1365-294X.2011.05193.x

Göring, H. H. H., Terwilliger, J. D., & Blangero, J. (2001). Large Upward Bias in Estimation of

Locus-Specific Effects from Genomewide Scans. *The American Journal of Human*

Genetics, *69*(6), 1357–1369. https://doi.org/10.1086/324471

Gu, B.-W., Tan, L.-M., Zhang, C.-J., Hou, X.-M., Cai, X.-W., Chen, S., & He, X.-J. (2020).

FHA2 is a plant-specific ISWI subunit responsible for stamen development and plant

fertility. *Journal of Integrative Plant Biology*, *62*(11), 1703–1716.

https://doi.org/10.1111/jipb.12945

Hughes, P. D., Woodward, J. C., & Gibbard, P. L. (2006). Quaternary glacial history of the

Mediterranean mountains. *Progress in Physical Geography: Earth and Environment*,

30(3), 334–364. https://doi.org/10.1191/0309133306pp481ra

Josephs, E. B., Berg, J. J., Ross-Ibarra, J., & Coop, G. (2019). Detecting Adaptive Differentiation

in Structured Populations with Genomic Data and Common Gardens. *Genetics*, *211*, 989–

1004. https://doi.org/10.1534/genetics.118.301786

Josephs, E. B., Stinchcombe, J. R., & Wright, S. I. (2017). What can genome-wide association

studies tell us about the evolutionary forces maintaining genetic variation for quantitative

traits? *New Phytologist*, *214*(1), 21–33. https://doi.org/10.1111/nph.14410

Kassambara, A. (2023). *ggpubr: "ggplot2" Based Publication Ready Plots* (Version R package

version 0.6.0) [Computer software]. <https://CRAN.R-project.org/package=ggpubr>.

Kooyers, N. J., Greenlee, A. B., Colicchio, J. M., Oh, M., & Blackman, B. K. (2015). Replicate

altitudinal clines reveal that evolutionary flexibility underlies adaptation to drought stress

in annual Mimulus guttatus. *New Phytologist*, *206*(1), 152–165.

https://doi.org/10.1111/nph.13153

Korunes, K. L., & Samuk, K. (2021). pixy: Unbiased estimation of nucleotide diversity and

divergence in the presence of missing data. *Molecular Ecology Resources*.

https://doi.org/10.1111/1755-0998.13326

Krueger, F. (2019). *Trim Galore* (Version 0.6.5) [Computer software]. Babraham

Bioinformatics. https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/

Kudo, G. (2003). Anther arrangement influences pollen deposition and removal in hermaphrodite

flowers. *Functional Ecology*, *17*(3), 349–355. https://doi.org/10.1046/j.1365-

2435.2003.00736.x

Lahti, D. C., Johnson, N. A., Ajie, B. C., Otto, S. P., Hendry, A. P., Blumstein, D. T., Coss, R.

G., Donohue, K., & Foster, S. A. (2009). Relaxed selection in the wild. *Trends in*

Ecology and Evolution, *24*(9), 487–496. https://doi.org/10.1016/j.tree.2009.03.010

Lande, R. (1979). Quantitative Genetic Analysis of Multivariate Evolution , Applied to Brain:

Body Size Allometry. *Evolution*, *33*(1), 402–416.

Laurie, C. C., Chasalow, S. D., LeDeaux, J. R., McCarroll, R., Bush, D., Hauge, B., Lai, C.,

Clark, D., Rocheford, T. R., & Dudley, J. W. (2004). The genetic architecture of response

to long-term artificial selection for oil concentration in the maize kernel. *Genetics*,

168(4), 2141–2155. https://doi.org/10.1534/genetics.104.029686

Lenth, R. V. (2021). *emmeans: Estimated marginal Means, aka Least Squares Means* (Version R

package version 1.6.2-1) [Computer software]. https://CRAN.R-

project.org/package=emmeans

Lewandowska-Sabat, A. M., Fjellheim, S., & Rognli, O. A. (2010). Extremely low genetic

variability and highly structured local populations of Arabidopsis thaliana at higher

latitudes. *Molecular Ecology*, *19*(21), 4753–4764. https://doi.org/10.1111/j.1365-

294X.2010.04840.x

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics*, *25*(14), 1754–1760.

https://doi.org/10.1093/bioinformatics/btp324

Luo, Y., Widmer, A., & Karrenberg, S. (2015). The roles of genetic drift and natural selection in

quantitative trait divergence along an altitudinal gradient in Arabidopsis thaliana.

Heredity, *114*(2), 220–228. https://doi.org/10.1038/hdy.2014.89

MacPherson, A., & Nuismer, S. L. (2017). The probability of parallel genetic evolution from

standing genetic variation. *Journal of Evolutionary Biology*, *30*(2), 326–337.

https://doi.org/10.1111/jeb.13006

Matsuhashi, S., Sakai, S., & Kudoh, H. (2012). Temperature-Dependent Fluctuation Of Stamen

Number In Cardamine hirsuta (Brassicaceae). *International Journal of Plant Sciences*,

173(4), 391–398. https://doi.org/10.1086/663966

Médail, F., & Diadema, K. (2009). Glacial Refugia Influence Plant Diversity Patterns in the

Mediterranean Basin. *Journal of Biogeography*, *36*(7), 1333–1345.

Montesinos, A., Tonsor, S. J., Alonso-Blanco, C., & Picó, F. X. (2009). Demographic and

genetic patterns of variation among populations of Arabidopsis thaliana from contrasting

native environments. *PLoS ONE*, *4*(9). https://doi.org/10.1371/journal.pone.0007213

Montesinos-Navarro, A., Wig, J., Xavier Pico, F., & Tonsor, S. J. (2011). Arabidopsis thaliana

populations show clinal variation in a climatic gradient associated with altitude. *New*

Phytologist, *189*(1), 282–294. https://doi.org/10.1111/j.1469-8137.2010.03479.x

Müller, A. (1961). Zur Charakterisierung der Blüten und Infloreszenzen vonArabidopsis thaliana

(L.) Heynh. *Die Kulturpflanze*, *9*(1), 364–393. https://doi.org/10.1007/BF02095757

Oakley, C. G., Lundemo, S., Ågren, J., & Schemske, D. W. (2019). Heterosis is common and

inbreeding depression absent in natural populations of *Arabidopsis thaliana*. *Journal of*

Evolutionary Biology, *32*(6), 592–603. https://doi.org/10.1111/jeb.13441

R Core Team. (2021). *R: A language and environment for statistical computing* [Computer

software]. R Foundation for Statistical Computing. https://www.R-project.org/.

Royer, A. M., Kremer, C., George, K., Pérez, S. G., Schemske, D. W., & Conner, J. K. (2016).

Incomplete loss of a conserved trait: Function, latitudinal cline, and genetic constraints.

Evolution, *70*(12), 2853–2864. https://doi.org/10.1111/evo.13096

Sicard, A., & Lenhard, M. (2011). The selfing syndrome: A model for studying the genetic and

evolutionary basis of morphological adaptation in plants. *Annals of Botany*, *107*(9),

1433–1443. https://doi.org/10.1093/aob/mcr023

Stinchcombe, J. R., Weinig, C., Ungerer, M., Olsen, K. M., Mays, C., Halldorsdottir, S. S.,

Purugganan, M. D., & Schmitt, J. (2004). A latitudinal cline in flowering time in

742   *Arabidopsis thaliana* modulated by the flowering time gene *FRIGIDA*. *Proceedings of*

743   *the National Academy of Sciences*, *101*(13), 4712–4717.

744   https://doi.org/10.1073/pnas.0306401101

745 Tang, C., Toomajian, C., Sherman-Broyles, S., Plagnol, V., Guo, Y. L., Hu, T. T., Clark, R. M.,

746   Nasrallah, J. B., Weigel, D., & Nordborg, M. (2007). The evolution of selfing in

747   Arabidopsis thaliana. *Science*, *317*(5841), 1070–1072.

748   https://doi.org/10.1126/science.1143153

749 The 1001 Genomes Consortium. (2016). 1,135 Genomes Reveal the Global Pattern of

750   Polymorphism in Arabidopsis thaliana. *Cell*, *166*(2), 481–491.

751   https://doi.org/10.1016/j.cell.2016.05.063

752 Tomizioli, M., Lazar, C., Brugière, S., Burger, T., Salvi, D., Gatto, L., Moyet, L., Breckels, L.

753   M., Hesse, A.-M., Lilley, K. S., Seigneurin-Berny, D., Finazzi, G., Rolland, N., & Ferro,

754   M. (2014). Deciphering Thylakoid Sub-compartments using a Mass Spectrometry-based

755   Approach *. *Molecular & Cellular Proteomics*, *13*(8), 2147–2167.

756   https://doi.org/10.1074/mcp.M114.040923

757 Turner, S. D. (2018). qqman: An R package for visualizing GWAS results using Q-Q and

758   manhattan plots. *Journal of Open Source Software*, *3*(25), 731.

759   https://doi.org/10.21105/JOSS.00731

760 Van der Auwera, G., & O'Connor, B. (2020). *Genomics in the Cloud: Using Docker, GATK, and*

761   *WDL in Terra (1st Edition)*. O'Reilly Media.

762 Walsh, B., & Blows, M. W. (2009). Abundant genetic variation + strong selection = multivariate

763   genetic constraints: A geometric view of adaptation. *Annual Review of Ecology,*

Evolution, and Systematics, *40*, 41–59.

https://doi.org/10.1146/annurev.ecolsys.110308.120232

Waterman, R., Sahli, H., Koelling, V. A., Karoly, K., & Conner, J. K. (2023). Strong evidence

for positive and negative correlational selection revealed by recreating ancestral

variation. *Evolution*, *77*(1), 264–275. https://doi.org/10.1093/evolut/qpac001

Whitlock, M. C., & Guillaume, F. (2009). Testing for Spatially Divergent Selection: Comparing

QST to FST. *Genetics*, *183*(3), 1055–1063. https://doi.org/10.1534/genetics.108.099812

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G.,

Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller,

K., Ooms, J., Robinson, D., Seidel, D., Spinu, V., … Yutani, H. (2019). Welcome to the

Tidyverse. *Journal of Open Source Software*, *4*(43), 1686.

https://doi.org/10.21105/joss.01686

Wolfe, M. D., & Tonsor, S. J. (2014). Adaptation to spring heat and drought in northeastern

Spanish Arabidopsis thaliana. *New Phytologist*, *201*(1), 323–334.

https://doi.org/10.1111/nph.12485

Yoshizawa, M., Yamamoto, Y., O'Quin, K. E., & Jeffery, W. R. (2012). Evolution of an

adaptive behavior and its sensory receptors promotes eye regression in blind cavefish.

BMC Biology, *10*(108). https://doi.org/10.1186/1741-7007-11-82

Zhou, X., & Stephens, M. (2012). Genome-wide efficient mixed-model analysis for association

studies. *Nature Genetics 2012 44:7*, *44*(7), 821–824. https://doi.org/10.1038/ng.2310

Zomlefer, W. B. (1994). *Guide to flowering plant families*. University of North Carolina Press.

785