Xenometabolome of Early-Life Stage Salmonids Exposed to 6PPD-Quinone

Phillip J. Ankley[1], Francisco C. da Silva Junior[1], Catherine Roberts[1], Andreas N. M. Eriksson[2],

Evan Kohlman[1], Justin Dubiel[2], Blake Hunnie[1], Katherine Anderson-Bain[2], Rayen M. Urrutia[2],

Natacha Hogan[1,3], John P. Giesy[1,5,6], Ed Krol[4], Steve Wiseman[2], Markus Hecker[1,7], and

Markus Brinkmann[1,7,8*]

[1]Toxicology Centre, University of Saskatchewan, SK, S7N 0H5, Canada

[2]Dept of Biological Science, University of Lethbridge, Lethbridge, AB, SA9 202, Canada

[3]Dept of Animal and Poultry Science, University of Saskatchewan, Saskatoon, SK, S7N 5A8, Canada

[4]Drug Discovery and Development Research Group, College of Pharmacy and Nutrition, University of

Saskatchewan, Saskatoon, SK, S7N 5E5, Canada

[5]Dept of Integrative Biology and Center for Integrative Toxicology, Michigan State University, East

Lansing, MI, 48824, USA

[6]Dept of Environmental Sciences, Baylor University, Waco, TX, 76798, USA

[7]School of Environment and Sustainability, University of Saskatchewan, Saskatoon, SK, S7N 5C8,

Canada

[8]Global Institute for Water Security, University of Saskatchewan, Saskatoon, SK, S7N 1K2, Canada

Corresponding author:

Markus Brinkmann

44 Campus Drive, Saskatoon, SK, S7N 0H5, Canada

markus.brinkmann@usask.ca

Abstract:

N-(1,3-Dimethylbutyl)-*N′*-phenyl-*p*-phenylenediamine-quinone (6PPD-Q) is a ubiquitous and acutely toxic

transformation product (TP) derived from the rubber tire antioxidant *N*-(1,3-Dimethylbutyl)-*N′*-phenyl-*p*-

phenylenediamine (6PPD). While not all salmonids are sensitive to acute lethality caused by 6PPD-Q, its

potency can vary by several orders of magnitude among fish species. The main driver(s) of species

sensitivity differences is a pressing question, with one area of interest examining whether differences in

their ability to biotransform and detoxify 6PPD-Q could be a driving factor. This study utilized liquid-

chromatography high-resolution mass spectrometry (LC-HRMS) to assess biotransformation and

metabolome-wide effects of 6PPD-Q on early-life stage salmonids, including two sensitive species,

rainbow trout (*Oncorhynchus mykiss*) and lake trout (*Salvelinus namaycush*), and one tolerant species,

brown trout (*Salmo trutta*). Three phase I TPs and seven phase II TPs were detected, with differences in

peak areas revealing that brown trout had the greatest ability to detoxify 6PPD-Q. TP-OH1, an alkyl

sidechain monohydroxylated TP, was detected in both rainbow and lake trout but not in brown trout, with

more research needed to understand potential TP-OH1 mediated toxicity. Several endogenous metabolites

were found to be dysregulated in rainbow and lake trout, indicative of mitochondrial dysfunction and altered

metabolism. Results of this study indicate a difference in the biotransformation capability of 6PPD-Q

among salmonid fish species and subsequent unique metabolome responses.

Keywords: Tire-derived chemical; urban runoff; high-resolution mass spectrometry; biotransformation;

cross-species sensitivity

INTRODUCTION

N-(1,3-Dimethylbutyl)-*N*′-phenyl-*p*-phenylenediamine-quinone (6PPD-Q) is a ubiquitous and acutely toxic transformation product derived from the rubber tire antioxidant *N*-(1,3-Dimethylbutyl)-*N*′-phenyl-*p*-phenylenediamine (6PPD).[1] 6PPD is released through diffusion from and wear of tire surfaces, then released into the environment, where it can react with oxidants, such as ozone, to form 6PPD-Q.[2] Urban runoff after rainfall events is the main source of 6PPD-Q into receiving aquatic ecosystems.[3] The total global amount of 6PPD-Q generated per year is estimated to fall between 26 to 1,900 tons,[4] with this chemical being detected globally up to levels of 2.29 µg/L in aquatic environments.[1] The amount of this contaminant being released into aquatic environments and concentrations at which it has been detected have been shown to significantly adversely impact fish populations, in particular, coho salmon (*Oncorhynchus kisutch*) and other select salmonid fishes.

6PPD-Q is the leading cause of urban runoff mortality syndrome (URMS) in U.S. Pacific Northwest coho salmon,[1] with a 24-h median lethal concentration (LC_{50}) of 0.095 µg/L.[5] 6PPD-Q has also been shown to be acutely toxic to other salmonids, including adult brook trout (*Salvelinus fontinalis*; 24-h $LC_{50} = 0.59$ µg/L),[6] rainbow trout (*Oncorhynchus mykiss*; 24-h $LC_{50} = 1.96$ µg/L),[6] and white-spotted char (*Salvelinus leucomaenis pluvius*; 24-h $LC_{50} = 0.8$ µg/L).[7] Compared to other fish species, such as white sturgeon (*Acipenser transmontanus*; 24-h $LC_{50} > 12.7$ µg/L) and zebrafish (*Danio rerio*; 24-h $LC_{50} = 308.7$ µg/L),[8] some salmonids are several orders of magnitude more sensitive to aqueous exposure of 6PPD-Q. What may be driving these large species differences in sensitivity is a pressing question for ecotoxicologists and risk assessors.

One area of investigation regarding the principal determinant of differences in sensitivity among fish species is their ability to biotransform and detoxify 6PPD-Q upon exposure. Previous research has indicated that fish differ in the levels of transformation products (TPs) from 6PPD-Q exposure, including increased levels of phase I hydroxylated 6PPD-Q and phase II glucuronidated 6PPD-Q in the bile of more tolerant species (i.e., chinook salmon (*Oncorhynchus tshawytscha*), westslope cutthroat trout (*Oncorhynchus clarkii lewisi*), and white sturgeon (*A. transmontanus*)).[9] Zebrafish embryos have also been

shown to have multiple biotransformation pathways, including sulfonate-, N-acetylcysteine-, and

glutathione-conjugation, that could reduce sensitivity to 6PPD-Q by detoxifying the chemical.[10]

Hydroxylated TPs are a result of phase I biotransformation, including an alkyl sidechain monohydroxylated

TP (TP-OH1), a phenyl ring monohydroxylated TP (TP-OH2),[11] and a dihydroxylated TP (TP-2OH).[11,12]

Additional phase II TPs occurring in 6PPD-Q-exposed salmonids must be better characterized to understand

differences in detoxification upon aqueous exposure.

Nontargeted liquid-chromatography high-resolution mass spectrometry (LC-HRMS) provides a

powerful method to characterize the xenometabolome and endometabolome of tissue of larval fishes.[13,14]

Xenometabolome analysis profiles TPs from the biotransformation of contaminants within organisms from

environmental exposure,[15] allowing the understanding of exposure and metabolism of contaminants that

could lead to potential toxicity (reactive metabolite) or detoxification (inactive metabolite).[16] Endogenous

metabolites are involved in the growth and homeostasis of fish, and the metabolome profile is the ultimate

measurement of cellular and tissue responses to toxicant exposure and a key link associating genes with

observable phenotypes.[17,18] Nontargeted LC-HRMS has the capability to simultaneously profile both the

xenometabolome and the endometabolome of fishes for assessment of contaminant contribution to toxicity

and changes in fish health from exposure.[19–21]

This study utilized LC-HRMS to assess biotransformation and metabolome-wide effects of 6PPD-

Q on the early-life stages of three salmonids, including two sensitive species, rainbow trout and lake trout

(45-d alevin $LC_{50} = 0.39$ µg/L; *Salvelinus namaycush*),[22] and one tolerant species, brown trout (48-h alevin

$LC_{50} = 12.2$ µg/L; *Salmo trutta*).[23] Specific objectives of this study included: (1) profile the

xenometabolome of 6PPD-Q within the salmonids; (2) compare detected TPs and levels among species;

and (3) assess if and how 6PPD-Q affects the endometabolome of each species.

MATERIALS AND METHODS

Chemicals and Fish Culture

6PPD-Q and deuterium-labeled 6PPD-Q-d5 were purchased from Toronto Research Chemicals

(Canada) and had purities of $\geq$ 97%, with standard solutions being prepared in HPLC-grade methanol

(MeOH). Stock solutions for fish exposures were prepared in fish facility water and dimethyl sulfoxide

(DMSO) to achieve a final solvent concentration of 0.1% (v/v).

Lake trout (LKT) embryos were obtained following field spawning from White Swan Lake,

Saskatchewan, and transported to the Aquatic Toxicology Research Facility (ATRF) at the University of

Saskatchewan (SK, Canada) under the Government of Saskatchewan Fish Transport/Stocking permit

FTP02-2022 as described previously.[24] Eyed embryos were moved into glass aquaria, held under a 12:12 h

light: dark photoperiod, and gradually acclimated to 9 $\pm$ 0.5 °C until hatch. Rainbow trout (RBT) triploid

eyed embryos were obtained from Lyndon Fish Hatcheries (ON, Canada). Eyed embryos were kept in an

environmentally controlled exposure chamber in glass aquaria with aerated dechlorinated facility water and

acclimated as described above using temperatures of 14 $\pm$ 0.5 °C. Brown trout (BRT) were obtained from

the Allison Creek Brood Trout Station (Coleman) in Alberta, Canada. Approximately 4000 newly fertilized

embryos were transported to the Aquatic Research Facility (ARF) at the University of Lethbridge (AB,

Canada) under dark and cool conditions and maintained in a flow-through system at a temperature of 10 °C

and 16:8 h light: dark photoperiod until hatching.

Exposure Experiments

All fish exposures were conducted independently, and varying nominal concentrations were used

for exposure (Supplementary Table S1). LKT and RBT exposures were approved by the University of

Saskatchewan Animal Research Ethics Board (Animal Use Protocol 2022-0002) following Canadian

Council on Animal Care guidelines, and BRT exposures were approved by the University of Lethbridge

Animal Welfare Committee (Animal Use Protocol 2324).

Newly hatched LKT, RBT, and BRT were exposed to nominal concentrations (ranging from 0.125

µg/L – 10 µg/L) of 6PPD-Q (Supplementary Table S1) until two weeks post-swim up following OECD

guidelines for the Early-life Stage Toxicity Test No. 210.[25] Upon hatch, RBT and LKT both had 18 alevins

placed in each tank (2.5 L) with five replicate tanks for each concentration (RBT – 0.125, 0.25, 0.5, 1, 2,

and 4 µg/L; LKT – 1.25 and 2.5 µg/L) as well as a solvent control (SC) of 0.01% DMSO, and BRT had 30

alevins in each tank (3.5 L) with five replicate tanks per chemical treatment (2.5, 5, and 10 µg/L). Fish were

maintained at an average daily temperature of 9 °C for LKT, 14 °C for RBT, and 10 °C for BRT, with a

photoperiod of 12:12 h light: dark for RBT and LKT, and a photoperiod of 16:8 h light: dark for BRT. All

exposures were conducted under static-renewal conditions, with 70% replacement of water and chemicals

daily throughout the entire exposure. Water quality parameters were monitored daily in randomly selected

aquaria, including temperature, dissolved oxygen, pH, ammonia, nitrates, nitrites, and hardness.[22] Samples

of water were collected into LC vials throughout the exposure and stored at –20 °C to confirm

concentrations of chemicals using LC-HRMS as described previously,[22] spiking 950 µL of water with 50

µL of 1000 µg/L 6PPD-Q-d5 for a final internal standard concentration of 50 µg/L. Greater details of the

LC-HRMS analysis of water samples are found in Supplementary Text S1. Salmonids were fed with

hatched brine shrimp (*Artemia* spp. nauplii) once daily approaching and through the first week of swim-up

and twice daily for the remainder of the exposure until 24 hours prior to termination. At 28 days post-hatch

(dph) for RBT and BRT and 45 dph for LKT, fry were euthanized in 150 mg/L buffered tricaine

methanesulfonate (MS-222). Individual fish were weighed and measured for length (standard and total).

Two to three fish from each tank were randomly collected for xenometabolome analysis of whole larval

tissue.

Mass Spectrometry Analysis

Whole larval fish were weighed before and after freeze-drying for 48 h. Twelve volumes of MeOH

per mg of dried mass (dm) were added with the inclusion of two 5-mm steel beads and 25 µL of 6PPD-Q-

d5 dissolved in MeOH (1,000 µg/L) before complete homogenization using Tissue Lyzer II at 20 Hz for 3

minutes. Samples were then centrifuged at 4,000 × g at 4 ºC for 30 min with the supernatant transferred to

a new 2.0 mL microcentrifuge tube with 6 volumes of ACN and vortexed for 1 min before storage at –20

ºC for 12 h. After overnight salt and protein precipitation, samples were centrifuged at 4,000 × g at 4 ºC for

30 min before transferring the supernatant to a new 2.0-mL glass vial to undergo nitrogen drying. Samples

were then reconstituted in 500 µL of 50:50 MeOH: H_2O and filtered using a syringe + PTFE 0.22 µM

syringe filter (Agilent, USA) into a new 2-mL glass vial before LC-HRMS analysis with a final internal

standard concentration of 50 µg 6PPD-Q-d5/L.

Nontargeted analyses were conducted utilizing a data-dependent acquisition (DDA) method.

Details on the separation of analytes using ultra-high performance liquid chromatography (UHPLC)

coupled with a heated electrospray ionization (HESI) ion source to a Q-Exactive™ HF Quadrupole-Orbitrap

(Thermo Scientific) are provided in Supplementary Text S2 and Table S3. The following parameters for

nontargeted analysis of extracts of larval fish were used to acquire the ddMS2 top 5 positive mode scans:

sheath gas flow = 35, aux gas flow = 10, sweep gas flow = 1, aux gas heater temperature = 300 °C, spray

voltage = 3.00 kV, S-lens RF = 60.0, and capillary temperature = 350 °C. A Full MS method used scan

settings of resolution = 60,000, positive ion mode, AGC target = 5e5, maximum injection time = 100 ms,

and Full MS scan range of 70-1,000 *m/z*. The ddMS2 used the following scan settings: resolution = 15,000,

AGC target = 1e5, maximum injection time = 100 ms, loop count = 5, TopN = 5, an isolation window of

2.0 *m/z*, and stepped normalized collision energy (NCE) of 15, 30 and 45. Blanks and a quality control (QC)

pool of all samples for each species were run throughout the sequence, and a 4-point standard curve was

run at the beginning of each run (0.05, 2.5, 5, and 25 µg/L) for semi-quantification of 6PPD-Q body

burdens.

Nontargeted Data Analysis

Thermo *.raw files were converted to mzML format using MSConvert with the peak picking filter

before downstream processing using MZmine 3.[26] Spectral preprocessing, feature detection, and

preliminary compound identification using MoNA LC-MS-MS Positive Mode database (Accessed August

23, 2023, https://mona.fiehnlab.ucdavis.edu/) were conducted in MZmine 3 (version 4.0.3) using

parameters outlined in supporting information (Supplementary File S1). To eliminate background, features

were removed if detected in two or more blanks and were less than 10-fold more abundant in samples. After

processing, mgf files were exported using the export feature lists command for feature-based molecular

networking (FBMN) using GNPS web server and molecular formula identification and compound

annotation using SIRIUS (version 5.8.5).[27,28] FBMN was run using default settings, and the resulting graph

was exported and visualized in Cytoscape to detect and prioritize TPs.[29] Several new TPs were observed

and identified using Biotransformer 3.0 and CFM-ID for spectral verification, along with review of

literature.[11] SIRIUS GUI was used to analyze the feature data using the SIRIUS algorithm with Orbitrap-

based default settings. Features were additionally analyzed using CSI:FingerID with Fingerprint Prediction

using Fallback Adducts of [M+H]+, [M]+, [M+K]+, and [M+Na]+, and Structure Database Search using

Bio Database, HMDB, and KEGG.[30]

The resulting data was filtered and merged to detect TPs and compare differences among species

for the 2-2.5 µg/L 6PPD-Q exposure bin. TPs were kept if they were detected in QC samples, omitting TPs

detected at low abundance and in a small number of samples. TPs that shared identical *m/z* were merged if

retention times (RTs) were ± 0.5 min of the average and had the same fragmentation pattern of precursor,

indicating they were the same TP with a slight shift in RT, which can occur at greater concentrations when

conducting a DDA experiment. Details of precursor and product ions used for verification are provided in

the supplementary information (Supplementary Table S4). If merged, the peak area was summed, and the

RT was averaged for the respective TPs. Peak areas were normalized to measured dm of larval fish to

compute the normalized relative abundance of TPs, as fish species differed in size upon take-down

(Supplementary Table S1). To investigate the relative bioaccumulation between salmonid species, the 4-

point standard curve was used to estimate 6PPD-Q body burden from peak areas, with concentrations being

normalized to dm of fish.

Statistics

Data analyses and visualizations of 6PPD-Q body burden and normalized peak areas of TPs were

performed using GraphPad Prism software (version 10.1.1). MetaboAnalyst (version 6.0) was used to

analyze larval fish metabolome response to 6PPD-Q.[31] All features passing filtering were used to compute

a Partial Least-Squares Discriminant Analysis (PLS-DA) to discern potential differences between

treatments (Supplementary Text S3). A volcano plot was used to differentiate dysregulated metabolites

between SC and 2-2.5 µg/L exposure bin using a false-discovery rate (FDR) *p*-value of 0.05 and fold change

(FC) of 2.0. Significant features that could be annotated by having a compound name derived from SIRIUS

and mapped to MetaboAnalyst common metabolite name database were used to further interpret the

biological response. Concentration-response analysis computed metabolite benchmark doses (BMDs) for

metabolomic points-of-departure (mPOD) using default settings while including Exp5 and Hill statistical

models with Exp2, Exp3, Exp4, Linear, Poly2, and Power models.[32] Briefly, an ANOVA was performed

using a limma R package, with metabolites having an FDR < 0.05 fitted to statistical models. BMDs were

calculated using a lack-of-fit p-value of 0.10, benchmark response factor of 1.0, and control abundance

using the mean of control samples. mPODs were estimated from the metabolomic BMD analysis as: (1) the

BMD of the 20th most sensitive metabolite, (2) the max 1st peak of metabolite BMD distribution, and (3)

the 10th percentile of all metabolite BMDs.

RESULTS AND DISCUSSION

Levels of 6PPD-Q Within Fish Samples

Verification of nominal exposure concentrations for 2-2.5 µg/L exposure bin indicated time-

weighted averaged concentrations of 1.30 ± 0.250 µg/L (average ± standard deviation) (65.0% of nominal

exposure concentration), 1.30 ± 0.167 µg/L (52.0%),[22] and 0.615 ± 0.550 µg/L (24.6%) for RBT, LKT, and

BRT, respectively. Recovery of 6PPD-Q-d5 from larval fish was $37.0 \pm 7.84\%$, $68.9 \pm 18.4\%$, and $47.1 \pm$

8.11%. for RBT, LKT, and BRT, respectively. Methanol was used to prioritize polar metabolites and TPs;

therefore, some loss of 6PPD-Q and 6PPD-Q-d5 (RT = 15.7) was expected.[33] 6PPD-Q body burdens were

33.2 ± 24.1 ng/g (RBT), 41.7 ± 13.0 ng/g (LKT), and 44.9 ± 25.5 ng/g (BRT) for the 2-2.5 µg/L exposure

bin. Concentrations of 6PPD-Q body burdens increased among fish species with increasing exposure

concentrations (Figure 1A). At the 2-2.5 µg/L exposure, LKT and BRT tended towards slightly greater

body burdens of 6PPD-Q compared to RBT (Figure 1B; ANOVA, *p*-value = 0.746), but these likely resulted

from the difference in nominal exposure concentrations (2 µg/L – RBT vs. 2.5 µg/L – LKT and BRT).

More research that incorporates time-course sampling of larval fish from aqueous exposure is needed to

understand the rate and magnitude of 6PPD-Q bioaccumulation between species.

Figure 1. Concentrations and normalized peak areas of 6PPD-Q and phase I and II transformation products

(TPs), respectively, in extracts of larval fish tissue. A) Concentrations of 6PPD-Q (ng/g) in extracted fish

tissue within nominal exposure bins for lake trout (LKT), rainbow trout (RBT), and brown trout (BRT). B)

Concentrations of 6PPD-Q (ng/g) in extracts of fish tissue for 2-2.5 µg/L nominal exposure. C) Ratio of the

summed normalized peak areas for phase I and II transformation products (TPs) in 2-2.5 µg/L exposed

fishes. D) Phase I 6PPD-Q TPs normalized peak areas in 2-2.5 µg/L exposed fishes. E) Phase II TPs

normalized peak areas in 2-2.5 µg/L exposed fishes.

Detection and Comparison of 6PPD-Q Transformation Products Among Fishes

TPs from biotransformation pathways were detected using suspect screening in 6PPD-Q-exposed

larval fish tissue. Three phase I TPs from CYP (Cytochromes P450) biotransformation[11,34] were detected in

fish exposed to 2-2.5 µg 6PPD-Q/L. These TPs included two monohydroxylated TPs and one

dihydroxylated TP (Figure 2D). The two monohydroxylated TPs (m/z = 315.17) included an alkyl sidechain

hydroxylation (TP-OH1, RT = 13.45) and a phenyl ring hydroxylation (TP-OH2, RT = 14.56), both of

which have been previously described.[11] TP-OH1 was only detected in the more sensitive fish species for

the 2-2.5 µg/L exposure bin (Figure 1D), with BRT only having TP-OH2 present. The dihydroxylated TP

(m/z = 331.17) showed identical fragmentation patterns for all species, indicating a phenyl ring

hydroxylation and alkyl sidechain hydroxylation, and was found to be greatest in RBT (averaged sum;

3.30×10^4 arbitrary units, au), followed by LKT (5.74×10^2 au) and BRT (4.31×10^2 au) (Figure 1D). RBT

had the greatest amounts of phase I TPs (7.36×10^4 au; 53.5% peak area of all summed TPs) compared to

both LKT (6.44×10^3 au; 28.3%) and BRT (5.64×10^3 au; 11.1%) (Figure 1D). Two additional phase I

phenyl ring monohydroxylated TPs were detected in BRT exposed to 10 µg/L. Having a similar

fragmentation profile as the phenyl ring hydroxylation suggests that hydroxylation might have occurred at

positions other than the para position (i.e., ortho and meta) at these higher concentrations, leading to

different log K_{ow} and retention times (RT = 16.48 and 17.83) (Supplementary Figure S1). This provides

additional evidence that the preferred phenyl ring monohydroxylation occurs at the para position due to

increased availability for CYP biotransformation and decreased steric hindrance.[11] Phase I to phase II

normalized relative abundance ratio values indicated that RBT had the lowest capacity to conjugate

hydroxylated TPs (Figure 1C), followed by LKT. BRT had the greatest ability to biotransform hydroxylated

TPs further through phase II conjugation, which could have important implications for the differences in

the reported species-specific sensitivities to 6PPD-Q.

A total of seven phase II TPs were detected for 2-2.5 µg/L exposure bin, with presence and relative

amounts differing among salmonids (Figure 1E). The seven phase II TPs included two glucuronide-

conjugated monohydroxylated TPs (m/z = 491.202), four glucuronide-conjugated dihydroxylated TPs (m/z

= 507.198), and one sulfonate-conjugated dihydroxylated TP ($m/z = 411.122$). In total, five, three, and seven

phase II TPs were detected for RBT, LKT, and BRT, respectively (Figure 1D). Glucuronide-conjugated

TPs dominated in both RBT (5.53×10^4 au; 40.2%) and BRT (3.94×10^4 au; 67.0%), while sulfonate-

conjugated TP dominated in LKT (1.24×10^4 au; 62.6%), indicating differences in the biotransformation

pathways that predominate among these three salmonids. RBT had the overall greatest amounts of phase II

TPs (6.40×10^4 au; 46.5%), followed by BRT (4.50×10^4 au; 88.9%), with LKT having the lowest relative

amounts of phase II TPs (1.63×10^4 au; 71.7%). However, RBT was the only species where phase II TP

peak areas were in deficit compared to phase I peak areas (Figure 1C).

271   The two glucuronide-conjugated monohydroxylated TPs were both *O*-glucuronidated at either the

alkyl sidechain hydroxylation (TP-*O*-Gluc1) or phenyl ring hydroxylation (TP-*O*-Gluc2) position (Figure

2, Supplementary Figure S2) with matching fragmentation (MS^2) to the two monohydroxylated TPs after

neutral loss of glucuronide acid ($m/z = 176.032$). The four glucuronide-conjugated dihydroxylated TPs

could be either *N*-glucuronidation or *O*-glucuronidation, attributed to four different TPs (Figure 2,

Supplementary Figure S3). The one sulfonate-conjugated dihydroxylated TP was likely *O*-sulfonated at the

phenyl ring hydroxylation, with an MS^2 fragment of $m/z = 337.05$, indicating that the other hydroxylation

event occurred on the methyl carbon adjacent to the amine, as previously predicted (Figure 2,

Supplementary Figure S4).[11]

Figure 2. Biotransformation pathways for 6PPD-Q phase I and phase II TPs. 6PPD-Q is biotransformed by CYP to form two different monohydroxylated TPs, TP-OH1 and TP-OH2. Monohydroxylated 6PPD-Q can then undergo glucuronide conjugation *via* UDP-glucuronosyltransferase (UGT) to form two TPs (TP-*O*-Gluc1 and TP-*O*-Gluc2). 6PPD-Q can be further biotransformed by CYP to form a dihydroxylated TP (TP-2-OH). The dihydroxylated TP can be further biotransformed by sulfate (Sulfotransferase; SULT) or glucuronide conjugation (UGT). The probable location of the sulfate conjugation is on the phenyl ring hydroxylation and does not occur on the alkyl sidechain hydroxylation (TP-2-*O*-Sulf; Supplementary

Figure S4). Glucuronide conjugation of the dihydroxylated TP can form four different TPs by *N*-

glucuronidation or *O*-glucuronidation (TP-2-*O*-Gluc1-4).

Metabolome Response of 6PPD-Q-Exposed Fishes

Differences between the greater exposure concentrations and SC for RBT and LKT were discerned

using PLS-DA (Figure 3A, B), reflecting alterations in the metabolome due to 6PPD-Q exposure. No

significant differences were detected for BRT, as all treatments overlapped in 95% confidence intervals

(Supplementary Figure S5). RBT and LKT treatments were largely separated by component 1 of the PLS-

DA, accounting for 26.5 and 49.1 of the variability, respectively. Volcano plot for the 2-2.5 µg/L exposure

bin, relative to SC, revealed a total of 153 (146 increased in the 6PPD-Q treatment group and seven

decreased compared to the SC) and 170 (nine increased and 161 decreased) dysregulated metabolites for

RBT and LKT, respectively (Figure 3C, D), with no dysregulated metabolites for BRT. The shift in

dysregulation of metabolites between RBT and LKT could reflect the differences in exposure length

between the species (28 d vs. 45 d), with differences in the initial and subsequent compensatory responses.

LKT was also shown to be more sensitive to 6PPD-Q-exposure, with a 45-d LC_{50} of 0.39 µg/L,[22] while

RBT displayed compensatory mechanisms towards chemical stressor response (28-d LC_{50} of 0.56 µg/L)

(unpublished), potentially reflecting the greater number of metabolites increased in abundance.

Xanthine was the annotated metabolite with the largest FC ($\log_2 FC = 6.46$) between SC and 6PPD-

Q group in RBT, while xanthine was decreased ($\log_2 FC = -6.78$) in the exposed LKT (Supplementary File

S2). Xanthine, an alkaloid, has a wide array of functions within organisms while also being imperative in

the catabolism of nucleotides and nucleic acids, being a precursor of uric acid,[35] indicating response to

stress in RBT and lethal disruption from stress in LKT. Xanthine is also an antagonist of adenosine

receptors,[36] with adenosine having the largest FC ($\log_2 FC = 7.17$) in exposed LKT. Adenosine is an

endogenous modulator of neuronal excitability,[37] and the increased abundance of the metabolite could be

potentially tied to some URMS-related symptoms (e.g., pectoral fin splaying and circling).[38] However, most

of the behaviors are likely tied to loss of respiration capability (e.g., surface swimming and mouth

gaping)[9,39]. Additional research is needed to directly link changes in metabolome profiles with URMS

behaviors and neurotoxicity. Several acylcarnitines of mid- to long-chain length were increased ($n = 6$) in

RBT from 6PPD-Q exposure, including stearoylcarnitine, valeryl-carnitine, 3-

hydroxyhexadecanoylcarnitine, 3-hydroxyhexanoylcarnitine, 3-hydroxyoctadecanoylcarnitine, and

butyrylcarnitine, indicating mitochondrial fatty acid oxidation was impacted within this species (perhaps

indicative of uncoupling of the mitochondrial electron transport chain).[39,40] Acylcarnitines are fatty acid

metabolites that transport acyl groups from the cytosol into the mitochondrial matrix for β-oxidation and

subsequent energy production,[41] and increases in acylcarnitines can be due to mitochondrial dysfunction

and alterations in metabolism. LKT had two acylcarnitines decreased in abundance, oleoylcarnitine and L-

acetylcarnitine. L-acetylcarnitine, a short-chain acylcarnitine, has a wide range of neuroprotective and

neurotrophic effects[42,43] while also facilitating the transfer of acetyl-Coenzyme A across mitochondrial

membranes for energy production. A decrease in L-acetylcarnitine could indicate mitochondrial and

neuroprotection dysfunction. No increased or decreased metabolites overlapped between RBT and LKT,

indicating that these two species differed in their physiological response to 6PPD-Q at the time of sampling.

Results indicate that RBT was trying to manage stressor impacts further, whereas LKT was ultimately dying

from exposure, also reflecting the differences in the biotransformation capability between the two species.

Using dose-response analysis, an mPOD was computed for the RBT metabolome (Supplementary

Figure S6), while LKT and BRT metabolome data were not conducive to mPOD analysis due to the limited

number of doses or a low number of dysregulated metabolites, respectively.[32,44] A total of 213 metabolites

were fitted with models, with 203 metabolites having BMDs for mPOD computation. The RBT mPOD had

a 0.44 µg/L feature-level BMD for the 20th feature and 10th percentile, which closely aligns with the 28-d

LC_{50} of larval RBT (unpublished) and is protective of the adult RBT 24-h LC_{50} of 1.96 µg/L.[7] Dose-response

analysis of metabolomics datasets is an emerging area of research in aquatic toxicology that could be widely

protective of aquatic health through predictive means, as portrayed here using 6PPD-Q as a study chemical.

Figure 3. RBT and LKT metabolome responses to 6PPD-Q. PLS-DA of A) RBT metabolome and B) LKT

metabolome response to 6PPD-Q. Individual points represent each sample collected. Shapes and colors

represent the different treatment groups. Volcano plot of dysregulated metabolites for C) RBT and D) LKT.

A false-discovery rate (FDR) *p*-value of 0.05 and fold change (FC) of 2.0 was used for discerning

dysregulated metabolites. A negative fold change is displayed in blue, and a positive fold change is

displayed in red.

OUTLOOK AND FUTURE RESEARCH

We profiled the xenometabolome of three different salmonids exposed to 6PPD-Q, revealing that

three phase I and seven phase II TPs dominated at lower exposure to 6PPD-Q. Different biotransformation

pathways can dominate among salmonids, with RBT and BRT having greater levels of glucuronide-

conjugated TPs than the sulfate-conjugated TP found to dominate in LKT. This can have important implications regarding species-specific biotransformation and subsequent response to contaminants, including 6PPD-Q. More research is needed to understand the main mechanisms of toxicological responses to 6PPD-Q and its phase I TPs, including TP-OH1, which was detected in both RBT and LKT but not BRT. A time-course metabolome response to 6PPD-Q is also needed to understand how larval fish initially respond to the contaminant and subsequent compensatory responses to have a greater mechanistic understanding of 6PPD-Q toxicity. Our results indicate that a difference in biotransformation capability and pathways is observed among salmonid fish species, and subsequent metabolome response to 6PPD-Q exposure can help reveal the mode of toxic action.

Acknowledgment

This project was supported partially by a financial contribution from Fisheries and Oceans Canada. Additional funding was provided to M.H., S.W., N.H., and M.B. through the Discovery Grants program of the Natural Sciences and Engineering Research Council of Canada (NSERC). Instrumentation used for chemical analyses within this research was obtained with funding from Western Economic Diversification Canada (WED) and the Canadian Foundation for Innovation (CFI). P.J.A. was supported through a Toxicology Centre Devolved Scholarship. M.B. is currently a faculty member of the Global Water Futures (GWF) program, which received funds from the Canada First Research Excellence Funds (CFREF).

References

(1) Tian, Z.; Zhao, H.; Peter, K. T.; Gonzalez, M.; Wetzel, J.; Wu, C.; Hu, X.; Prat, J.; Mudrock, E.; Hettinger, R.; Cortina, A. E.; Biswas, R. G.; Kock, F. V. C.; Soong, R.; Jenne, A.; Du, B.; Hou, F.; He, H.; Lundeen, R.; Gilbreath, A.; Sutton, R.; Scholz, N. L.; Davis, J. W.; Dodd, M. C.; Simpson, A.; McIntyre, J. K.; Kolodziej, E. P. A Ubiquitous Tire Rubber–Derived Chemical Induces Acute Mortality in Coho Salmon. *Science* **2021**, *371* (6525), 185–189. https://doi.org/10.1126/science.abd6951.

(2) Zoroufchi Benis, K.; Behnami, A.; Minaei, S.; Brinkmann, M.; McPhedran, K. N.; Soltan, J. Environmental Occurrence and Toxicity of 6PPD Quinone, an Emerging Tire Rubber-Derived Chemical: A Review. *Environ. Sci. Technol. Lett.* **2023**, *10* (10), 815–823. https://doi.org/10.1021/acs.estlett.3c00521.

(3) Rauert, C.; Charlton, N.; Okoffo, E. D.; Stanton, R. S.; Agua, A. R.; Pirrung, M. C.; Thomas, K. V. Concentrations of Tire Additive Chemicals and Tire Road Wear Particles in an Australian Urban Tributary. *Environ. Sci. Technol.* **2022**, *56* (4), 2421–2431. https://doi.org/10.1021/acs.est.1c07451.

(4) Hu, X.; Zhao, H. N.; Tian, Z.; Peter, K. T.; Dodd, M. C.; Kolodziej, E. P. Transformation Product Formation upon Heterogeneous Ozonation of the Tire Rubber Antioxidant 6PPD (N-(1,3-Dimethylbutyl)-N′-Phenyl-p-Phenylenediamine). *Environ. Sci. Technol. Lett.* **2022**, *9* (5), 413–419. https://doi.org/10.1021/acs.estlett.2c00187.

(5) Tian, Z.; Gonzalez, M.; Rideout, C. A.; Zhao, H. N.; Hu, X.; Wetzel, J.; Mudrock, E.; James, C. A.; McIntyre, J. K.; Kolodziej, E. P. 6PPD-Quinone: Revised Toxicity Assessment and Quantification with a Commercial Standard. *Environ. Sci. Technol. Lett.* **2022**, *9* (2), 140–146. https://doi.org/10.1021/acs.estlett.1c00910.

(6) Brinkmann, M.; Montgomery, D.; Selinger, S.; Miller, J. G. P.; Stock, E.; Alcaraz, A. J.; Challis, J. K.; Weber, L.; Janz, D.; Hecker, M.; Wiseman, S. Acute Toxicity of the Tire Rubber-Derived Chemical 6PPD-Quinone to Four Fishes of Commercial, Cultural, and Ecological Importance. *Environ. Sci. Technol. Lett.* **2022**, *9* (4), 333–338. https://doi.org/10.1021/acs.estlett.2c00050.

(7) Hiki, K.; Yamamoto, H. The Tire-Derived Chemical 6PPD-Quinone Is Lethally Toxic to the White-Spotted Char Salvelinus Leucomaenis Pluvius but Not to Two Other Salmonid Species. *Environ. Sci. Technol. Lett.* **2022**, *9* (12), 1050–1055. https://doi.org/10.1021/acs.estlett.2c00683.

(8) Varshney, S.; Gora, A. H.; Siriyappagouder, P.; Kiron, V.; Olsvik, P. A. Toxicological Effects of 6PPD and 6PPD Quinone in Zebrafish Larvae. *J Hazard Mater* **2022**, *424* (Pt C), 127623. https://doi.org/10.1016/j.jhazmat.2021.127623.

(9) Montgomery, D.; Ji, X.; Cantin, J.; Philibert, D.; Foster, G.; Selinger, S.; Jain, N.; Miller, J.; McIntyre, J.; de Jourdan, B.; Wiseman, S.; Hecker, M.; Brinkmann, M. Interspecies Differences in 6PPD-Quinone Toxicity Across Seven Fish Species: Metabolite Identification and Semiquantification. *Environ Sci Technol* **2023**, *57* (50), 21071–21079. https://doi.org/10.1021/acs.est.3c06891.

(10) Grasse, N.; Seiwert, B.; Massei, R.; Scholz, S.; Fu, Q.; Reemtsma, T. Uptake and Biotransformation of the Tire Rubber-Derived Contaminants 6-PPD and 6-PPD Quinone in the Zebrafish Embryo (Danio Rerio). *Environ. Sci. Technol.* **2023**, *57* (41), 15598–15607. https://doi.org/10.1021/acs.est.3c02819.

(11) Ankley, P. J.; da Silva Junior, F. C.; Montgomery, D.; Schultz, M.; Krol, E. S.; Hecker, M.; Brinkmann, M. Biotransformation of 6PPD-Quinone In Vitro Using RTL-W1 Cell Line. *Environ. Sci. Technol. Lett.* https://doi.org/10.1021/acs.estlett.4c00342.

(12) Zhang, J.; Cao, G.; Wang, W.; Qiao, H.; Chen, Y.; Wang, X.; Wang, F.; Liu, W.; Cai, Z. Stable Isotope-Assisted Mass Spectrometry Reveals In Vivo Distribution, Metabolism, and Excretion of Tire Rubber-Derived 6PPD-Quinone in Mice. *Science of The Total Environment* **2024**, *912*, 169291. https://doi.org/10.1016/j.scitotenv.2023.169291.

(13) Ribbenstedt, A.; Ziarrusta, H.; Benskin, J. P. Development, Characterization and Comparisons of Targeted and Non-Targeted Metabolomics Methods. *PLOS ONE* **2018**, *13* (11), e0207082. https://doi.org/10.1371/journal.pone.0207082.

(14) Evich, M. G.; Mosley, J. D.; Ntai, I.; Cavallin, J. E.; Villeneuve, D. L.; Ankley, G. T.; Collette, T. W.; Ekman, D. R. Untargeted MSn-Based Monitoring of Glucuronides in Fish: Screening Complex Mixtures for Contaminants with Biological Relevance. *ACS ES T Water* **2022**, *2* (12), 2481–2490. https://doi.org/10.1021/acsestwater.2c00310.

(15) Al-Salhi, R.; Abdul-Sada, A.; Lange, A.; Tyler, C. R.; Hill, E. M. The Xenometabolome and Novel Contaminant Markers in Fish Exposed to a Wastewater Treatment Works Effluent. *Environ. Sci. Technol.* **2012**, *46* (16), 9080–9088. https://doi.org/10.1021/es3014453.

(16) Nair, P.; Sun, J.; Xie, L.; Kennedy, L.; Kozakiewicz, D.; Kleywegt, S.; Hao, C.; Byun, H.; Barrett, H.; Baker, J.; Monaghan, J.; Krogh, E.; Song, D.; Peng, H. Synthesis and Toxicity Evaluation of Tire Rubber-Derived Quinones. ChemRxiv June 20, 2023. https://doi.org/10.26434/chemrxiv-2023-pmxvc.

(17)Fiehn, O. Metabolomics--the Link between Genotypes and Phenotypes. *Plant Mol Biol* **2002**, *48* (1–
2), 155–171.
(18)Bonvallot, N.; Tremblay-Franco, M.; Chevrier, C.; Canlet, C.; Debrauwer, L.; Cravedi, J.-P.; Cordier,
S. Potential Input from Metabolomics for Exploring and Understanding the Links between
Environment and Health. *J Toxicol Environ Health B Crit Rev* **2014**, *17* (1), 21–44.
https://doi.org/10.1080/10937404.2013.860318.
(19)Xu, M.; Legradi, J.; Leonards, P. A Comprehensive Untargeted Metabolomics Study in Zebrafish
Embryos Exposed to Perfluorohexane Sulfonate (PFHxS). *Science of The Total Environment* **2023**,
887, 163770. https://doi.org/10.1016/j.scitotenv.2023.163770.
(20)Magnuson, J. T.; Giroux, M.; Cryder, Z.; Gan, J.; Schlenk, D. The Use of Non-Targeted
Metabolomics to Assess the Toxicity of Bifenthrin to Juvenile Chinook Salmon (*Oncorhynchus
Tshawytscha*). *Aquatic Toxicology* **2020**, *224*, 105518.
https://doi.org/10.1016/j.aquatox.2020.105518.
(21)Ortiz-Villanueva, E.; Jaumot, J.; Martínez, R.; Navarro-Martín, L.; Piña, B.; Tauler, R. Assessment of
Endocrine Disruptors Effects on Zebrafish (Danio Rerio) Embryos by Untargeted LC-HRMS
Metabolomic Analysis. *Science of The Total Environment* **2018**, *635*, 156–166.
https://doi.org/10.1016/j.scitotenv.2018.03.369.
(22) Roberts, C.; Lin, J.; Kohlman, E.; Jain, N.; Amekor, M.; Alcaraz, A. J.; Hogan, N.; Hecker, M.;
Brinkmann, M. Acute and Sub-Chronic Toxicity of 6PPD-Quinone to Early-Life Stage Lake Trout
(Salvelinus Namaycush). bioRxiv March 29, 2024, p 2024.03.26.586843.
https://doi.org/10.1101/2024.03.26.586843.
(23)Foldvik, A.; Kryuchkov, F.; Sandodden, R.; Uhlig, S. Acute Toxicity Testing of the Tire Rubber–
Derived Chemical 6PPD-Quinone on Atlantic Salmon (Salmo Salar) and Brown Trout (Salmo
Trutta). *Environmental Toxicology and Chemistry* **2022**, *41* (12), 3041–3045.
https://doi.org/10.1002/etc.5487.
(24)Ankley, P.; Amekor, M.; Kohlman, E.; Roberts, C.; Zhou, Y.; Hecker, M.; Giesy, J.; Brinkmann, M.;
Hogan, N. Effects of Antimicrobial Exposure on the Gut Microbiome of Early-Life Stage Fish: A
Chemical and Species Comparison. Rochester, NY March 18, 2024.
https://doi.org/10.2139/ssrn.4763861.
(25)OECD. Test No. 210: Fish, Early-Life Stage Toxicity Test, 2013. https://www.oecd-
ilibrary.org/content/publication/9789264203785-en.
(26)Schmid, R.; Heuckeroth, S.; Korf, A.; Smirnov, A.; Myers, O.; Dyrlund, T. S.; Bushuiev, R.; Murray,
K. J.; Hoffmann, N.; Lu, M.; Sarvepalli, A.; Zhang, Z.; Fleischauer, M.; Dührkop, K.; Wesner, M.;
Hoogstra, S. J.; Rudt, E.; Mokshyna, O.; Brungs, C.; Ponomarov, K.; Mutabdžija, L.; Damiani, T.;
Pudney, C. J.; Earll, M.; Helmer, P. O.; Fallon, T. R.; Schulze, T.; Rivas-Ubach, A.; Bilbao, A.;
Richter, H.; Nothias, L.-F.; Wang, M.; Orešič, M.; Weng, J.-K.; Böcker, S.; Jeibmann, A.; Hayen,
H.; Karst, U.; Dorrestein, P. C.; Petras, D.; Du, X.; Pluskal, T. Integrative Analysis of Multimodal
Mass Spectrometry Data in MZmine 3. *Nat Biotechnol* **2023**, *41* (4), 447–449.
https://doi.org/10.1038/s41587-023-01690-2.
(27)Dührkop, K.; Fleischauer, M.; Ludwig, M.; Aksenov, A. A.; Melnik, A. V.; Meusel, M.; Dorrestein,
P. C.; Rousu, J.; Böcker, S. SIRIUS 4: A Rapid Tool for Turning Tandem Mass Spectra into
Metabolite Structure Information. *Nat Methods* **2019**, *16* (4), 299–302.
https://doi.org/10.1038/s41592-019-0344-8.
(28)Nothias, L.-F.; Petras, D.; Schmid, R.; Dührkop, K.; Rainer, J.; Sarvepalli, A.; Protsyuk, I.; Ernst, M.;
Tsugawa, H.; Fleischauer, M.; Aicheler, F.; Aksenov, A. A.; Alka, O.; Allard, P.-M.; Barsch, A.;
Cachet, X.; Caraballo-Rodriguez, A. M.; Da Silva, R. R.; Dang, T.; Garg, N.; Gauglitz, J. M.;
Gurevich, A.; Isaac, G.; Jarmusch, A. K.; Kameník, Z.; Kang, K. B.; Kessler, N.; Koester, I.; Korf,
A.; Le Gouellec, A.; Ludwig, M.; Martin H, C.; McCall, L.-I.; McSayles, J.; Meyer, S. W.;
Mohimani, H.; Morsy, M.; Moyne, O.; Neumann, S.; Neuweger, H.; Nguyen, N. H.; Nothias-
Esposito, M.; Paolini, J.; Phelan, V. V.; Pluskal, T.; Quinn, R. A.; Rogers, S.; Shrestha, B.; Tripathi,

A.; van der Hooft, J. J. J.; Vargas, F.; Weldon, K. C.; Witting, M.; Yang, H.; Zhang, Z.; Zubeil, F.; Kohlbacher, O.; Böcker, S.; Alexandrov, T.; Bandeira, N.; Wang, M.; Dorrestein, P. C. Feature-Based Molecular Networking in the GNPS Analysis Environment. *Nat Methods* **2020**, *17* (9), 905–908. https://doi.org/10.1038/s41592-020-0933-6.

(29) Shannon, P.; Markiel, A.; Ozier, O.; Baliga, N. S.; Wang, J. T.; Ramage, D.; Amin, N.; Schwikowski, B.; Ideker, T. Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks. *Genome Res* **2003**, *13* (11), 2498–2504. https://doi.org/10.1101/gr.1239303.

(30) Dührkop, K.; Shen, H.; Meusel, M.; Rousu, J.; Böcker, S. Searching Molecular Structure Databases with Tandem Mass Spectra Using CSI:FingerID. *Proceedings of the National Academy of Sciences* **2015**, *112* (41), 12580–12585. https://doi.org/10.1073/pnas.1509788112.

(31) Pang, Z.; Zhou, G.; Ewald, J.; Chang, L.; Hacariz, O.; Basu, N.; Xia, J. Using MetaboAnalyst 5.0 for LC–HRMS Spectra Processing, Multi-Omics Integration and Covariate Adjustment of Global Metabolomics Data. *Nat Protoc* **2022**, *17* (8), 1735–1761. https://doi.org/10.1038/s41596-022-00710-w.

(32) Yao, C.-H.; Wang, L.; Stancliffe, E.; Sindelar, M.; Cho, K.; Yin, W.; Wang, Y.; Patti, G. J. Dose-Response Metabolomics To Understand Biochemical Mechanisms and Off-Target Drug Effects with the TOXcms Software. *Anal Chem* **2020**, *92* (2), 1856–1864. https://doi.org/10.1021/acs.analchem.9b03811.

(33) Hu, X.; Zhao, H. (Nina); Tian, Z.; Peter, K. T.; Dodd, M. C.; Kolodziej, E. P. Chemical Characteristics, Leaching, and Stability of the Ubiquitous Tire Rubber-Derived Toxicant 6PPD-Quinone. *Environ. Sci.: Processes Impacts* **2023**, *25* (5), 901–911. https://doi.org/10.1039/D3EM00047H.

(34) Zhang, Y.-Y.; Huang, J.-W.; Liu, Y.-H.; Zhang, J.-N.; Huang, Z.; Liu, Y.-S.; Zhao, J.-L.; Ying, G.-G. *In Vitro* Metabolism of the Emerging Contaminant 6PPD-Quinone in Human and Rat Liver Microsomes: Kinetics, Pathways, and Mechanism. *Environmental Pollution* **2024**, *345*, 123514. https://doi.org/10.1016/j.envpol.2024.123514.

(35) Singh, N.; Shreshtha, A. K.; Thakur, M. S.; Patra, S. Xanthine Scaffold: Scope and Potential in Drug Development. *Heliyon* **2018**, *4* (10), e00829. https://doi.org/10.1016/j.heliyon.2018.e00829.

(36) Jacobson, K. A.; Gao, Z.-G. Adenosine Receptors as Therapeutic Targets. *Nat Rev Drug Discov* **2006**, *5* (3), 247–264. https://doi.org/10.1038/nrd1983.

(37) Boison, D. Adenosine and Seizure Termination: Endogenous Mechanisms. *Epilepsy Curr* **2013**, *13* (1), 35–37. https://doi.org/10.5698/1535-7511-13.1.35.

(38) Scholz, N. L.; Myers, M. S.; McCarthy, S. G.; Labenia, J. S.; McIntyre, J. K.; Ylitalo, G. M.; Rhodes, L. D.; Laetz, C. A.; Stehr, C. M.; French, B. L.; McMillan, B.; Wilson, D.; Reed, L.; Lynch, K. D.; Damm, S.; Davis, J. W.; Collier, T. K. Recurrent Die-Offs of Adult Coho Salmon Returning to Spawn in Puget Sound Lowland Urban Streams. *PLOS ONE* **2011**, *6* (12), e28013. https://doi.org/10.1371/journal.pone.0028013.

(39) Mahoney, H.; da Silva Junior, F. C.; Roberts, C.; Schultz, M.; Ji, X.; Alcaraz, A. J.; Montgomery, D.; Selinger, S.; Challis, J. K.; Giesy, J. P.; Weber, L.; Janz, D.; Wiseman, S.; Hecker, M.; Brinkmann, M. Exposure to the Tire Rubber-Derived Contaminant 6PPD-Quinone Causes Mitochondrial Dysfunction In Vitro. *Environ. Sci. Technol. Lett.* **2022**, *9* (9), 765–771. https://doi.org/10.1021/acs.estlett.2c00431.

(40) Reuter, S. E.; Evans, A. M. Carnitine and Acylcarnitines. *Clin Pharmacokinet* **2012**, *51* (9), 553–572. https://doi.org/10.1007/BF03261931.

(41) Indiveri, C.; Iacobazzi, V.; Tonazzi, A.; Giangregorio, N.; Infantino, V.; Convertini, P.; Console, L.; Palmieri, F. The Mitochondrial Carnitine/Acylcarnitine Carrier: Function, Structure and Physiopathology. *Molecular Aspects of Medicine* **2011**, *32* (4), 223–233. https://doi.org/10.1016/j.mam.2011.10.008.

(42) Ferreira, G. C.; McKenna, M. C. L-Carnitine and Acetyl-l-Carnitine Roles and Neuroprotection in Developing Brain. *Neurochem Res* **2017**, *42* (6), 1661–1675. https://doi.org/10.1007/s11064-017-2288-7.

(43) Smeland, O. B.; Meisingset, T. W.; Borges, K.; Sonnewald, U. Chronic Acetyl-l-Carnitine Alters
Brain Energy Metabolism and Increases Noradrenaline and Serotonin Content in Healthy Mice.
Neurochemistry International **2012**, *61* (1), 100–107. https://doi.org/10.1016/j.neuint.2012.04.008.
(44) Ewald, J. D.; Basu, N.; Crump, D.; Boulanger, E.; Head, J. Characterizing Variability and
Uncertainty Associated with Transcriptomic Dose–Response Modeling. *Environ. Sci. Technol.*
2022, *56* (22), 15960–15968. https://doi.org/10.1021/acs.est.2c04665.
539