**Xenometabolome of Early-Life Stage Salmonids Exposed to 6PPD-Quinone**

Phillip J. Ankley[1], Francisco C. da Silva Junior[1], Blake Hunnie[1], Catherine Roberts[1], Andreas N. M.

Eriksson[2], Evan Kohlman[1], Jennifer Corker[3], Jodi Westcott[3], Bradlee Germain[3], Michael J. Martin[3], Justin

Dubiel[2], Katherine Anderson-Bain[2], Rayen M. Urrutia[2], Natacha Hogan[1,4], John P. Giesy[1,5,6], Steve

Wiseman[2], Ed Krol[8], Markus Hecker[1,7,9], and Markus Brinkmann[1,7,9*]

*[1]Toxicology Centre, University of Saskatchewan, SK, S7N 0H5, Canada*

*[2]Dept of Biological Science, University of Lethbridge, Lethbridge, AB, SA9 202, Canada*

*[3]Cayman Chemical Company, Ann Arbor Michigan, 48108, USA*

*[4]Dept of Animal and Poultry Science, University of Saskatchewan, Saskatoon, SK, S7N 5A8, Canada*

*[5]Dept of Integrative Biology and Center for Integrative Toxicology, Michigan State University, East*

*Lansing, MI, 48824, USA*

*[6]Dept of Environmental Sciences, Baylor University, Waco, TX, 76798, USA*

*[7]School of Environment and Sustainability, University of Saskatchewan, Saskatoon, SK, S7N 5C8,*

*Canada*

*[8]Drug Discovery and Development Research Group, College of Pharmacy and Nutrition, University of*

*Saskatchewan, Saskatoon, SK, S7N 5E5, Canada*

*[9]Global Institute for Water Security, University of Saskatchewan, Saskatoon, SK, S7N 1K2, Canada*

Corresponding author:

Markus Brinkmann

44 Campus Drive, Saskatoon, SK, S7N 0H5, Canada

markus.brinkmann@usask.ca

**Abstract**

*N*-(1,3-Dimethylbutyl)-*N'*-phenyl-*p*-phenylenediamine-quinone (6PPD-Q) is a ubiquitous transformation

product (TP) derived from the rubber tire antioxidant *N*-(1,3-Dimethylbutyl)-*N'*-phenyl-*p*-

phenylenediamine (6PPD) and is acutely toxic to certain species of *Salmonidae*. Not all salmonids are

sensitive to acute lethality caused by 6PPD-Q, with 6PPD-Q potency varying by several orders of

magnitude among teleosts. The main driver(s) of species sensitivity differences is (are) a pressing

question, with one area of interest examining whether differences in teleosts ability to biotransform and

detoxify 6PPD-Q could be a key factor. This study utilized liquid-chromatography high-resolution mass

spectrometry (LC-HRMS) to assess biotransformation and metabolome-wide effects of 6PPD-Q on early-

life stage salmonids, including two sensitive species, rainbow trout (*Oncorhynchus mykiss*) and lake trout

(*Salvelinus namaycush*), and one tolerant species, brown trout (*Salmo trutta*). Three phase I TPs and

seven phase II TPs were detected, with differences in peak areas revealing that brown trout had the

greatest ability to detoxify 6PPD-Q. Monohydroxylated TPs were verified using co-developed analytical

standards that will be of use for future biomonitoring and exposure assessment. Several endogenous

metabolites were found to be dysregulated in rainbow and lake trout, indicative of mitochondrial

dysfunction, altered metabolism, and disrupted membrane permeability. Results of this study indicate a

difference in the biotransformation capability of 6PPD-Q among *Salmonidae* fish species and subsequent

unique metabolome responses.

**Keywords**

Tire-derived chemical; urban runoff; high-resolution mass spectrometry; biotransformation; cross-species

sensitivity

**Graphical Abstract**

**Synopsis**

Inter-species salmonid differences in the biotransformation capability of 6PPD-Q leads to sensitivity

variability and subsequent unique metabolome responses.

## INTRODUCTION

*N*-(1,3-Dimethylbutyl)-*N'*-phenyl-*p*-phenylenediamine-quinone (6PPD-Q) is a ubiquitous and *Salmonidae*-specific acutely toxic transformation product (TP) derived from the rubber tire antioxidant *N*-(1,3-Dimethylbutyl)-*N'*-phenyl-*p*-phenylenediamine (6PPD).[1] 6PPD is released into the environment through diffusion from and wear of tire surfaces, where it can react with oxidants, such as ozone, to form 6PPD-Q.[2] Urban runoff after rainfall events is the main source of 6PPD-Q into receiving aquatic ecosystems.[3] The total global amount of 6PPD-Q generated per year is estimated to fall between 26 to 1,900 tons,[4] with this chemical being detected globally up to levels of 2.29 µg/L in aquatic environments.[1] The amount of this contaminant being released into aquatic environments, and concentrations at which it has been detected *in situ*, have been shown to impact fish populations adversely. In particular, the coho salmon (*Oncorhynchus kisutch*), and other select *Salmonidae*, have been shown to be especially sensitive to 6PPD-Q exposure.[1,5]

6PPD-Q is the leading cause of urban runoff mortality syndrome (URMS) in U.S. Pacific Northwest coho salmon,[1] with a 24-h median lethal concentration ($LC_{50}$) of 0.095 µg/L.[6] 6PPD-Q has also been shown to be acutely toxic to other salmonids, including adult brook trout (*Salvelinus fontinalis*; 24-h $LC_{50}$ = 0.59 µg/L),[6] rainbow trout (*Oncorhynchus mykiss*; 24-h $LC_{50}$ = 1.96 µg/L),[5] white-spotted char (*Salvelinus leucomaenis pluvius*; 24-h $LC_{50}$ = 0.8 µg/L),[7] and juvenile lake trout (*Salvelinus namaycush*; 96-h $LC_{50}$ = 0.50 µg/L).[8] In contrast, most other fish species tested to date, such as white sturgeon (*Acipenser transmontanus*; 24-h $LC_{50}$ > 12.7 µg/L)[5] and zebrafish (*Danio rerio*; 24-h $LC_{50}$ = 308.7 µg/L),[9] as well as other salmonids including arctic char (*Salvelinus alpinus*; $LC_{50}$ > 14.2 µg/L)[5] and brown trout (*Salmo trutta*; 48-h alevin $LC_{50}$ > 12.2 µg/L),[10] are several orders of magnitude less sensitive to aqueous exposure of 6PPD-Q. The specific factors driving these large species differences in sensitivity are a pressing question for ecotoxicologists and risk assessors.

One area of investigation regarding a potential determinant of differences in sensitivity among fish species is their ability to biotransform and detoxify 6PPD-Q upon exposure. Previous research has indicated that fish differ in the levels of TPs from 6PPD-Q exposure, including increased levels of phase I

hydroxylated 6PPD-Q and phase II glucuronidated 6PPD-Q in the bile of more tolerant species, i.e.,

chinook salmon (*Oncorhynchus tshawytscha*), westslope cutthroat trout (*Oncorhynchus clarkii lewisi*),

and white sturgeon.[11] Zebrafish embryos have also been shown to have multiple biotransformation

pathways, including sulfonate-, N-acetylcysteine-, and glutathione-conjugation, that could reduce

sensitivity to 6PPD-Q by effectively detoxifying the chemical.[12] Hydroxylated 6PPD-Q TPs are a result of

phase I biotransformation, including an alkyl sidechain monohydroxylated TP (TP-OH1), a phenyl ring

monohydroxylated TP (TP-OH2),[13] and a dihydroxylated TP (TP-2-OH).[13,14] However, little is known

about additional phase II TPs occurring in 6PPD-Q-exposed salmonids, which should be better

characterized to understand differences in detoxification upon aqueous exposure.

Non-targeted liquid-chromatography high-resolution mass spectrometry (LC-HRMS) provides a

powerful method to characterize the xenometabolome and endometabolome of tissue of larval fishes.[15,16]

Xenometabolome analysis profiles TPs from the biotransformation of contaminants within organisms

from environmental exposure,[17] allowing the understanding of exposure and metabolism of contaminants

that could lead to potential toxicity (reactive metabolite) or detoxification (inactive metabolite).

Endogenous metabolites are involved in the growth and homeostasis of fish, and the metabolome profile

is the ultimate measurement of cellular and tissue responses to toxicant exposure and a key link

associating genes with observable phenotypes.[18,19] Non-targeted LC-HRMS has the capability to

simultaneously profile both the xenometabolome and the endometabolome of fishes for assessment of

contaminant contribution to toxicity and changes in fish health from exposure.[20–22]

This study utilized LC-HRMS to assess biotransformation and metabolome-wide effects of

6PPD-Q on the early-life stages of three salmonids, including two sensitive species, rainbow trout (28-d

alevin $LC_{50} = 0.56$ µg/L)[23] and lake trout (45-d alevin $LC_{50} = 0.39$ µg/L),[8] and one tolerant species, brown

trout.[23] Specific objectives of this study included: (1) profile the xenometabolome of 6PPD-Q within the

three selected salmonids; (2) compare and verify detected TPs and levels among the species; and (3)

assess      if      and      how      6PPD-Q      affects      the      endometabolome      of      each      species.

## MATERIALS AND METHODS

### Chemicals and Fish Culture

6PPD-Q (CAS #: 2754428-18-5) and deuterium-labeled 6PPD-Q-d5 (CAS #: 2750119-14-1) were purchased from Toronto Research Chemicals (Canada) and had purities of $\geq$ 97%, with standard solutions being prepared in HPLC-grade methanol (MeOH). Stock solutions for fish exposures were prepared in carbon- and bio-filtered City of Saskatoon water and dimethyl sulfoxide (DMSO) to achieve a final solvent concentration of 0.1% (v/v).

Lake trout (LKT) embryos were obtained following field spawning from White Swan Lake, Saskatchewan, and transported to the Aquatic Toxicology Research Facility (ATRF) at the University of Saskatchewan (SK, Canada) under the Government of Saskatchewan Fish Transport/Stocking permit FTP02-2022, as described previously.[24] Eyed embryos were moved into glass aquaria, held under a 12:12 h light: dark photoperiod, and gradually acclimated to 10 $\pm$ 0.5 °C until hatch. Triploid rainbow trout (RBT) eyed embryos were obtained from Lyndon Fish Hatcheries (ON, Canada). Eyed embryos were kept in an environmentally controlled exposure chamber in glass aquaria with aerated dechlorinated facility water and acclimated as described above using temperatures of 14 $\pm$ 0.5 °C until hatch. Brown trout (BRT) embryos were obtained from the Allison Creek Brood Trout Station (Coleman) in Alberta, Canada. Approximately 4000 newly fertilized embryos were transported to the Aquatic Research Facility (ARF) at the University of Lethbridge (AB, Canada) under dark and cool conditions and maintained in a flow-through system at a temperature of 10 °C and 16:8 h light: dark photoperiod until hatching.

### Exposure Experiments

All fish experiments were conducted independently, and varying nominal concentrations were used for exposure (Supplementary Table S1). LKT and RBT studies were approved by the University of Saskatchewan Animal Research Ethics Board (Animal Use Protocol 2022-0002) following Canadian Council on Animal Care guidelines and greater details of the experiments can be found in previous

publications.[8,23] BRT exposures were approved by the University of Lethbridge Animal Welfare

Committee (Animal Use Protocol 2324).

Newly hatched LKT, RBT, and BRT were exposed to nominal concentrations (ranging from

0.125 µg/L – 10 µg/L) of 6PPD-Q (Supplementary Table S1) until two weeks post-swim up following

OECD guidelines for the Early-life Stage Toxicity Test No. 210.[24] Upon hatch, RBT and LKT had 18 and

15 alevins, respectively, placed in each tank (2.5 L) with five replicate tanks for each concentration (RBT

– 0.125, 0.25, 0.5, 1, 2, and 4 µg/L; LKT – 1.25 and 2.5 µg/L) as well as a solvent control of 0.01%

DMSO. BRT had 30 alevins in each tank (3.5 L) with five replicate tanks per chemical treatment (2.5, 5,

and 10 µg/L). Fish were maintained at an average daily temperature of 10 °C for LKT, 14 °C for RBT,

and 10 °C for BRT, with a photoperiod of 12:12 h light: dark, gaining 1 hr light per week for a final

photoperiod of 16:8 h light: dark for LKT, RBT, and BRT All exposures were conducted under

semistatic-renewal conditions, with 70% replacement of water and chemicals daily throughout the entire

exposure. Water quality parameters were monitored daily in randomly selected aquaria, including

temperature and dissolved oxygen, and incoming processed facility water for ammonia, nitrates, and

nitrites (Supplementary Table S2).[8] Samples of water were collected into LC vials throughout the

exposure and stored at –20 °C to confirm concentrations of chemicals using LC-HRMS as described

previously,[8] spiking 950 µL of water with 50 µL of 1000 µg/L 6PPD-Q-d5 for a final internal standard

concentration of 50 µg/L (Supplementary Text S1). Salmonids were fed with hatched brine shrimp

(*Artemia* spp. nauplii) once daily approaching and through the first week of swim-up and twice daily for

the remainder of the exposure until 24 hours prior to termination. At 28 days post-hatch (dph) for RBT

and BRT and 45 dph for LKT, fry were euthanized in 150 mg/L buffered tricaine methanesulfonate (MS-

222). Individual fish were weighed and measured for length (standard and total). Two to three fish from

each tank were randomly collected for xenometabolome analysis of whole larval tissue.

**Mass Spectrometry Analysis**

Whole larval fish were weighed before and after freeze-drying, a process that lasted for 48 h.

Twelve volumes of MeOH (µL) per mg of dried mass (dm) were added with the inclusion of two 5-mm

steel beads and 25 µL of 6PPD-Q-d5 dissolved in MeOH (1,000 µg/L) before complete homogenization

using Tissue Lyzer II at 20 Hz for 3 minutes. Samples were then centrifuged at 4,000 × g at 4 ℃ for 30

min with the supernatant transferred to a new 2.0 mL microcentrifuge tube with 6 volumes of ACN and

vortexed for 1 min before storage at –20 ℃ for 12 h. After overnight salt and protein precipitation,

samples were centrifuged at 4,000 × g at 4 ℃ for 30 min before transferring the supernatant to a new 2.0-

mL glass vial to undergo nitrogen drying. Samples were then reconstituted in 500 µL of 50:50 MeOH:

$H_2O$ and filtered using a syringe + PTFE 0.22 µM syringe filter (Agilent, USA) into a new 2-mL glass

vial before LC-HRMS analysis with a final internal standard concentration of 50 µg 6PPD-Q-d5/L.

Non-targeted analyses were conducted utilizing a data-dependent acquisition (DDA) method.

Details on the separation of analytes using ultra-high performance liquid chromatography (UHPLC)

coupled with a heated electrospray ionization (HESI) ion source to a Q-Exactive[TM] HF Quadrupole-

Orbitrap (Thermo Scientific) are provided in Supplementary Text S3 and Table S4. The following

parameters for non-targeted analysis of extracts of larval fish were used to acquire the ddMS2 top 5

positive mode scans: sheath gas flow = 35, aux gas flow = 10, sweep gas flow = 1, aux gas heater

temperature = 300 °C, spray voltage = 3.00 kV, S-lens RF = 60.0, and capillary temperature = 350 °C. A

Full MS method used scan settings of resolution = 60,000, positive ion mode, AGC target = 5e5,

maximum injection time = 100 ms, and Full MS scan range of 70-1,000 *m/z*. The ddMS2 used the

following scan settings: resolution = 15,000, AGC target = 1e5, maximum injection time = 100 ms, loop

count = 5, TopN = 5, an isolation window of 2.0 *m/z*, and stepped normalized collision energy (NCE) of

15, 30 and 45. Blanks and a quality control (QC) pool of all samples for each species were run throughout

the sequence, and a 4-point standard curve was run at the beginning of each run (0.05, 2.5, 5, and 25

µg/L) for semi-quantification of 6PPD-Q body burdens. Monohydroxylated TPs were confirmed and

distinguished using co-developed analytical standards from Cayman Chemical using a paired full scan

and parallel reaction monitoring (PRM) method using the same LC-HRMS setup (Supplementary Text S3

and S4). These synthesized analytical standards were compared to biologically occurring

monohydroxylated TPs for verification (Supplementary Text S3).

**Non-targeted Data Analysis**

Thermo *.raw files were converted to mzML format using MSConvert with the peak picking

filter before downstream processing using MZmine 3.[25] Spectral preprocessing, feature detection, and

preliminary compound identification using MoNA LC-MS-MS Positive Mode database (Accessed

August 23, 2023) were conducted in MZmine 3 (version 4.0.3) using parameters outlined in supporting

information (Supplementary File S1). To eliminate background, features were removed if detected in two

or more blanks and were less than 10-fold more abundant in samples. After processing, mgf files were

exported using the export feature lists command for feature-based molecular networking (FBMN) using

GNPS web server and molecular formula identification and compound annotation using SIRIUS (version

5.8.5).[26,27] FBMN was run using default settings, and the resulting graph was exported and visualized in

Cytoscape to detect and prioritize TPs.[28] Several new TPs were observed and identified using

Biotransformer 3.0 and CFM-ID for spectral verification, along with review of literature.[13,29] SIRIUS GUI

was used to analyze the feature data using the SIRIUS algorithm with Orbitrap-based default settings.

Features were additionally analyzed using CSI:FingerID with Fingerprint Prediction using Fallback

Adducts of [M+H]+, [M]+, [M+K]+, and [M+Na]+, and Structure Database Search using Bio Database,

HMDB, and KEGG.[30]

The resulting data was filtered and merged to detect TPs and compare differences among species

for the 2-2.5 µg/L 6PPD-Q exposure bin. TPs were kept if they were detected in QC samples, omitting

TPs detected at low abundance and in a small number of samples. TPs that shared identical *m/z* were

merged if retention times (RTs) were ± 0.5 min of the average and had the same fragmentation pattern of

precursor, indicating they were the same TP with a slight shift in RT, which can occur at greater

concentrations when conducting a DDA experiment and when running multiple experiments. Details of

precursor and product ions used for verification are provided in the supplementary information

(Supplementary Table S5). If merged, the peak area was summed, and the RT was averaged for the

respective TPs. Peak areas were normalized to measured dm of larval fish to compute the normalized

relative abundance of TPs, as fish species differed in size upon take-down (Supplementary Table S1). To

investigate the relative bioaccumulation between salmonid species, the 4-point standard curve was used to

estimate 6PPD-Q body burden from peak areas, with concentrations being normalized to dm of fish.

**Statistics**

Data analyses and visualizations of 6PPD-Q body burden and normalized peak areas of TPs were

performed using GraphPad Prism software (version 10.1.1). MetaboAnalyst (version 6.0) was used to

analyze larval fish metabolome response to 6PPD-Q.[31] All features passing filtering were used to compute

a Partial Least-Squares Discriminant Analysis (PLS-DA) to discern potential differences between

treatments (Supplementary Text S5). A volcano plot was used to differentiate dysregulated metabolites

between solvent control and 2-2.5 µg/L exposure bin using a false-discovery rate (FDR) *p*-value of 0.05

and fold change (FC) of 2.0. Significant features that could be annotated by having a compound name

derived from SIRIUS and mapped to MetaboAnalyst common metabolite name database were used to

further interpret the biological response (Supplementary File S2). Concentration-response analysis

computed metabolite benchmark doses (BMDs) for metabolomic points-of-departure (mPOD) using

default settings while including Exp5 and Hill statistical models with Exp2, Exp3, Exp4, Linear, Poly2,

and Power models.[31] Briefly, an ANOVA was performed using a limma R package, with metabolites

having an FDR < 0.05 fitted to statistical models. BMDs were calculated using a lack-of-fit p-value of

0.10, benchmark response factor of 1.0, and control abundance using the mean of control samples.

mPODs were estimated from the metabolomic BMD analysis as: (1) the BMD of the 20[th] most sensitive

metabolite, (2) the max 1st peak of metabolite BMD distribution, and (3) the 10[th] percentile of all

metabolite                                                         BMDs.

## RESULTS AND DISCUSSION

### Levels of 6PPD-Q Within Fish Samples

Verification of nominal exposure concentrations for 2-2.5 µg/L exposure bin indicated time-weighted averaged concentrations of $1.30 \pm 0.250$ µg/L (average ± standard deviation) (65.0% of nominal exposure concentration),[23] $1.30 \pm 0.167$ µg/L (52.0%),[8] and $0.615 \pm 0.550$ µg/L (24.6%) for RBT, LKT, and BRT, respectively. Recovery of 6PPD-Q-d5 from larval fish extracts was $37.0 \pm 7.84\%$, $68.9 \pm 18.4\%$, and $47.1 \pm 8.11\%$. for RBT, LKT, and BRT, respectively. Methanol was used to prioritize polar metabolites and TPs; therefore, some loss of 6PPD-Q and 6PPD-Q-d5 (RT = 15.7) was expected.[32] 6PPD-Q body burdens were $33.2 \pm 24.1$ ng/g (RBT), $41.7 \pm 13.0$ ng/g (LKT), and $44.9 \pm 25.5$ ng/g (BRT) for the 2-2.5 µg/L exposure bin. Concentrations of 6PPD-Q body burdens increased among fish species with increasing exposure concentrations (Figure 1A). At the 2-2.5 µg/L exposure, LKT and BRT tended towards slightly greater body burdens of 6PPD-Q compared to RBT (Figure 1B; ANOVA, *p*-value = 0.746), but these likely resulted from the difference in nominal exposure concentrations (2 µg/L – RBT vs. 2.5 µg/L – LKT and BRT). More research that incorporates time-course sampling of larval fish from aqueous exposure is needed to understand the rate and magnitude of 6PPD-Q bioaccumulation between species. Bioaccumulation is an important measurement for environmental risk assessment,[33] and exposure-time is a necessary endpoint to consider when assessing bioaccumulation of chemicals. Additionally, 6PPD-Q is known to have time-driven effects,[1,34] with research suggesting differences between acute and sub-chronic mode of toxic action.[8]

**Figure 1.** Concentrations and normalized peak areas of 6PPD-Q and phase I and II transformation products (TPs), respectively, in extracts of larval fish tissue. A) Concentrations of 6PPD-Q (ng/g) in extracted fish tissue within nominal exposure bins for lake trout (LKT), rainbow trout (RBT), and brown trout (BRT). B) Concentrations of 6PPD-Q (ng/g) in extracts of fish tissue for 2-2.5 µg/L nominal exposure. C) Ratio of the summed normalized peak areas for phase I and II transformation products (TPs) in 2-2.5 µg/L exposed fishes. D) Phase I 6PPD-Q TPs normalized peak areas in 2-2.5 µg/L exposed fishes. E) Phase II TPs normalized peak areas in 2-2.5 µg/L exposed fishes.

**Detection and Comparison of 6PPD-Q Transformation Products Among Fishes**

TPs from biotransformation pathways were detected using suspect screening in 6PPD-Q-exposed larval fish tissue. Three phase I TPs from CYP (Cytochromes P450) biotransformation[13,35] were detected in fish exposed to 2-2.5 µg 6PPD-Q/L. These TPs included two monohydroxylated TPs and one dihydroxylated TP (Figure 2D). The two monohydroxylated TPs ($m/z$ = 315.17) included an alkyl

sidechain hydroxylation (TP-OH1, RT = 13.45) and a phenyl ring hydroxylation (TP-OH2, RT = 14.56),

both of which have been previously described.[13] TP-OH1 was verified to be hydroxylated on the carbon

4-position of the alkyl sidechain using a synthesized standard (6-PPD-Q-4-OH, Cayman Chemical),[36] and

TP-OH2 was additionally verified to occur on para position of the phenyl ring using an analytical

standard (*p*-hydroxy 6PPD-Q, Cayman Chemical item number 40605) (Supplementary Text S3;

Supplementary Figure S8). TP-OH1 was only detected in the more sensitive fish species for the 2-2.5

µg/L exposure bin (Figure 1D), with BRT only having the more abundant TP-OH2 present.[13,37] The

dihydroxylated TP (*m/z* = 331.17) showed identical fragmentation patterns for all species, indicating a

phenyl ring hydroxylation and alkyl sidechain hydroxylation, and was found to be greatest in RBT

(averaged sum; $3.30 \times 10^4$ arbitrary units, au), followed by LKT ($5.74 \times 10^2$ au) and BRT ($4.31 \times 10^2$ au)

(Figure 1D). RBT had the greatest amounts of phase I TPs ($7.36 \times 10^4$ au; 53.5% peak area of all summed

TPs) compared to both LKT ($6.44 \times 10^3$ au; 28.3%) and BRT ($5.64 \times 10^3$ au; 11.1%) (Figure 1D). Two

additional phase I phenyl ring monohydroxylated TPs were detected in BRT exposed to 10 µg/L. Having

a similar fragmentation profile as the phenyl ring monohydroxylation suggests that hydroxylation might

have occurred at positions other than the para position (i.e., ortho and meta) at these higher

concentrations, leading to different log $K_{ow}$ and retention times (RT = 16.48 and 17.83) (Supplementary

Figure S1). This provided additional evidence that the preferred phenyl ring monohydroxylation occurred

at the para position due to increased availability for CYP biotransformation and decreased steric

hindrance.[11] Phase I to phase II normalized relative abundance ratio values indicated that RBT had the

lowest capacity to conjugate hydroxylated TPs (Figure 1C), followed by LKT. BRT had the greatest

ability to biotransform hydroxylated TPs further through phase II conjugation, with the phase 1/ phase 2

peak area ratio being lowest for BRT, suggesting overall biotransformation of 6-PPDQ is likely important

in *Salmonidae*-specific acute toxicity,[11] which could have important implications for differences in

reported species-specific sensitivities to 6PPD-Q. Affinity for P450 enzymes within *Salmonidae* could

play an important role in the enhanced ability of specific species to detoxify 6PPD-Q.[38] Fish species

ability to biotransform chemicals can have important implications in respective sensitivity and can drive

changes in endemic biodiversity assemblages.[39] There is likely a difference between salmonids ability to

detoxify 6PPD-Q driving differences in sensitivity, whereas other phylogenetic groups could also have

differences in detoxication pathways but additionally changes in target protein structure.[7]

A total of seven phase II TPs were detected for 2-2.5 µg/L exposure bin, with presence and

relative amounts differing among salmonids (Figure 1E). The seven phase II TPs included two

glucuronide-conjugated monohydroxylated TPs ($m/z$ = 491.202), four glucuronide-conjugated

dihydroxylated TPs ($m/z$ = 507.198), and one sulfonate-conjugated dihydroxylated TP ($m/z$ = 411.122). In

total, five, three, and seven phase II TPs were detected for RBT, LKT, and BRT, respectively (Figure

1D). Greater number of phase II TPs were also detected in the more tolerant zebrafish, a *Cyprinidae*,

reflecting a greater capacity to detoxify.[12] Glucuronide-conjugated TPs dominated in both RBT (5.53 $\times$

$10^4$ au; 40.2%) and BRT (3.94 $\times$ $10^4$ au; 67.0%), while sulfonate-conjugated TP dominated in LKT (1.24

$\times$ $10^4$ au; 62.6%), indicating differences in the biotransformation pathways that predominate among these

three salmonids. The most predominant TPs in mammalian models were also monohydroxylated 6PPD-Q

and glucuronide-conjugated 6PPD-Q,[40] similar to both RBT and BRT. RBT had the overall greatest

amounts of phase II TPs (6.40 $\times$ $10^4$ au; 46.5%), followed by BRT (4.50 $\times$ $10^4$ au; 88.9%), with LKT

having the lowest relative amounts of phase II TPs (1.63 $\times$ $10^4$ au; 71.7%). However, RBT was the only

species where phase II TP peak areas were in deficit compared to phase I peak areas (Figure 1C).

The two glucuronide-conjugated monohydroxylated TPs were both *O*-glucuronidated at either the

alkyl sidechain hydroxylation (TP-*O*-Gluc1) or phenyl ring hydroxylation (TP-*O*-Gluc2) position (Figure

2, Supplementary Figure S2) with matching fragmentation (MS$^2$) to the two monohydroxylated TPs after

neutral loss of glucuronide acid ($m/z$ = 176.032). The four glucuronide-conjugated dihydroxylated TPs

could be either *N*-glucuronidation or *O*-glucuronidation, attributed to four different TPs (Figure 2,

Supplementary Figure S3). The single sulfonate-conjugated dihydroxylated TP was likely *O*-sulfonated at

the phenyl ring hydroxylation, with an MS$^2$ fragment of $m/z$ = 311.034 indicating loss of water and an

alkyl sidechain (Figure 2, Supplementary Figure S4).

**Figure 2.** Biotransformation pathways for 6PPD-Q phase I and phase II TPs. 6PPD-Q is biotransformed

by CYP to form two different monohydroxylated TPs, TP-OH1 and TP-OH2. Monohydroxylated 6PPD-

Q can then undergo glucuronide conjugation *via* UDP-glucuronosyltransferase (UGT) to form two TPs

(TP-*O*-Gluc1 and TP-*O*-Gluc2). 6PPD-Q can be further biotransformed by CYP to form a dihydroxylated

TP (TP-2-OH). The dihydroxylated TP can be further biotransformed by sulfate (Sulfotransferase; SULT)

or glucuronide conjugation (UGT). The probable location of the sulfate conjugation is on the phenyl ring

hydroxylation and does not occur on the alkyl sidechain hydroxylation (TP-2-*O*-Sulf; Supplementary

Figure S4). Glucuronide conjugation of the dihydroxylated TP can form four different TPs by *N*-

glucuronidation or *O*-glucuronidation (TP-2-*O*-Gluc1-4).

**Metabolome Response of 6PPD-Q-Exposed Fishes**

Differences between the greater exposure concentrations and solvent control for RBT and LKT

were discerned using PLS-DA (Figure 3A, B), reflecting alterations in the metabolome due to 6PPD-Q

exposure. No significant differences were detected for BRT, as all treatments overlapped in 95%

confidence intervals (Supplementary Figure S5). RBT and LKT treatments were largely separated by

component 1 of the PLS-DA, accounting for 26.5 and 49.1 % of the variability, respectively. Volcano plot

for the 2-2.5 µg/L exposure bin, relative to solvent control, revealed a total of 153 (146 increased in the

6PPD-Q treatment group and seven decreased compared to the solvent control) and 170 (nine increased

and 161 decreased) dysregulated metabolites for RBT and LKT, respectively (Figure 3C, D), with no

dysregulated metabolites for BRT. The shift in dysregulation of metabolites between RBT and LKT could

reflect the differences in exposure length between the species (28 d vs. 45 d), with differences in the

initial and subsequent compensatory responses. 6PPD-Q toxicity is observed to have differences between

acute and sub-chronic toxicity, with a longer duration for LKT potentially driving differences in

dysregulated metabolites compared to RBT.[8] LKT was shown to be slightly more sensitive to 6PPD-Q-

exposure, with a 45-d $LC_{50}$ of 0.39 µg/L compared to RBT 28-d $LC_{50}$ of 0.56 µg/L,[8,23] which may reflect

compensatory mechanisms towards the chemical stressor for RBT and the greater number of increased

metabolites.

Xanthine, an alkaloid, was the annotated metabolite with the greatest enrichment between solvent

control and 6PPD-Q exposure group in RBT ($log_2FC = 6.46$; Supplementary Figure S6A; Supplementary

File S2), while xanthine was depleted in exposed LKT ($log_2FC = -6.78$; Supplementary Figure S6B).

Xanthine has a wide array of functions within organisms while also being imperative in the catabolism of

nucleotides and nucleic acids, being a precursor of uric acid,[41] indicating response to stress in RBT and

disruption from stress in LKT. Xanthine is also an antagonist of adenosine receptors,[42] with adenosine

having the greatest enrichment in exposed LKT ($\log_2$FC = 7.17), indicating a balanced relationship

between the two metabolites. Adenosine is an endogenous modulator of neuronal excitability,[43] and the

increased abundance of the metabolite could be potentially tied to some of the Urban Runoff Mortality

Syndrome (URMS)-related symptoms (e.g., pectoral fin splaying and circling).[44] Acute toxicity from

6PPD-Q can lead to behavioral abnormalities in exposed salmonids, including loss of equilibrium, erratic

swimming patterns, lethargy, and disorientation, with these phenotypes being designated as URMS.[44]

Several additional URMS behaviors from acute 6PPD-Q toxicity are also potentially tied to loss of

respiration capability (e.g., surface swimming and mouth gaping).[45] Several acylcarnitines of mid- to

long-chain length were increased ($n = 6$) in RBT from 6PPD-Q exposure, including stearoylcarnitine,

valeryl-carnitine, 3-hydroxyhexadecanoylcarnitine, 3-hydroxyhexanoylcarnitine, 3-

hydroxyoctadecanoylcarnitine, and butyrylcarnitine, indicating mitochondrial fatty acid oxidation was

impacted within this species (perhaps indicative of uncoupling of the mitochondrial electron transport

chain).[46,47] Acylcarnitines are fatty acid metabolites that transport acyl groups from the cytosol into the

mitochondrial matrix for $\beta$-oxidation and subsequent energy production[48] and increases in acylcarnitines

can be due to mitochondrial dysfunction and alterations in metabolism. LKT had two acylcarnitines

decreased in abundance, oleoylcarnitine and L-acetylcarnitine. L-acetylcarnitine, a short-chain

acylcarnitine, has a wide range of neuroprotective and neurotrophic effects[48,49] while also facilitating the

transfer of acetyl-Coenzyme A across mitochondrial membranes for energy production with a decrease in

L-acetylcarnitine indicating mitochondrial and neuroprotection dysfunction.

Increases in both lysophosphatidylcholines (LPCs; $n = 6$) and phosphatidylcholines (PCs; $n = 2$)

were observed in RBT with decreases observed in LKT indicating that there was a disruption in

phospholipid biosynthesis, leading to potentially altered membrane permeability.[50] Higher concentrations

of LPCs can also disrupt mitochondrial integrity and modulate inflammatory chemokine expression.[51,52]

Changing levels of PCs can lead to neurotoxicity from altered levels of acetylcholine leading to additional

neurological effects.[53,54] Greater amounts of dysregulated LPCs derived from the cleaving of PC indicate

activation of signaling pathways that are involved in oxidative stress and inflammatory responses.[52]

Disruption in blood-brain barrier (BBB) permeability was an observed transcriptome response in

developing coho salmon that was indicated to alter central nervous system function.[55] LPCs are important

for the transport of polyunsaturated fatty acids (PUFAs) across BBB via major-facilitator superfamily

transporter MFSD2A.[56] Several long-chain polyunsaturated fatty acids (LC-PUFAs) were seen to increase

in sub-chronically exposed RBT including arachidonic acid (AA; omega-6 fatty acid), 8,9-

epoxyeicosatrienoic acid (a metabolite of AA formed through P450 epoxygenases), Dihomo-gamma-

linolenic acid (omega-6 fatty acid), docosahexaenoic acid (DHA; omega-3 fatty acid), eicosapentaenoic

acid ethyl ester (omega-3 fatty acid that is an esterified form of eicosapentaenoic acid (EPA)), and

docosapentaenoic acid (DPA; omega-3 fatty acid that is an intermediate between EPA and DHA).

Increases in these LC-PUFAs could indicate alterations in the AA pathway, with importance to

inflammatory processes and disease,[57] while alterations in important omega-3 PUFAs could modify key

involvement in anti-inflammatory processes and cell membrane permeability.[58] Several LC-PUFAs were

observed to decrease in exposed LKT including AA, DPA, EPA, linoleic acid, and dihomo-alpha-

linolenic acid indicating deficiencies in essential fatty acids that could lead to altered metabolic and

membrane function, chronic inflammation, neurological dysregulation, and oxidative stress.[59] Additional

metabolites with the greatest increased FC in exposed RBT included 11-hydroxy DHA ($\log_2FC = 4.51$;

Supplementary Figure S6C), a marker of inflammation and the result of oxidation,[60] palmitoleic acid

($\log_2FC = 4.10$; Supplementary Figure S6D), 14S-hydroxy-4Z,7Z,10Z,12E,16Z,19Z-docosahexaenoic

acid ($\log_2FC = 4.08$), an additional marker of oxidative stress resulting from oxidation of DHA,[61] and

inosine ($\log_2FC = 3.51$), a nucleoside and versatile bioactive metabolite with increases in abundance

linked to regulating inflammation, immunomodulatory effects, and neuroprotection.[62,63]

No increased or decreased metabolites overlapped between RBT and LKT, indicating that these

two species differed in their physiological response to 6PPD-Q at the time of sampling. Results indicate

that RBT were trying to manage stressor impacts further by compensation with a shorter exposure time,

whereas LKT physiological processes were ultimately deteriorating from an extended exposure duration,

reflecting the differences in the biotransformation capability between the two species. More effort will be

needed to profile endogenous metabolite alterations, including temporal responses, using developed target

LC-HRMS methods for specific metabolite classes. Additional research is also needed to link changes in

metabolome profiles directly with URMS behaviors and neurotoxicity.

402           Using dose-response analysis, an mPOD was computed for the RBT metabolome (Supplementary

Figure S7), while LKT and BRT metabolome data were not conducive to mPOD analysis due to the

limited number of doses or a low number of dysregulated metabolites, respectively.[64,65] A total of 213

metabolites were fitted with models, with 203 metabolites having BMDs for mPOD computation. The

RBT mPOD had a 0.44 µg/L feature-level BMD for the $20^{th}$ feature and $10^{th}$ percentile, which closely

aligns with the 28-d $LC_{50}$ of larval RBT and is protective of the adult RBT 24-h $LC_{50}$ of 1.96 µg/L or 72-g

$LC_{50} = 1.00$ µg/L.[7] Dose-response analysis of metabolomics datasets is an emerging area of research in

aquatic toxicology that could be widely indicative of aquatic health through predictive means and used for

chemical risk assessment,[66] as portrayed here using 6PPD-Q as a study chemical. Utilizing generated non-

targeted metabolomics datasets can provide an efficient method for calculating mPODs from organismal

response to chemicals.

**Figure 3.** RBT and LKT metabolome responses to 6PPD-Q. PLS-DA of A) RBT metabolome and B)

LKT metabolome response to 6PPD-Q. Individual points represent each sample collected. Shapes and

colors represent the different treatment groups. Volcano plot of dysregulated metabolites for C) RBT and

D) LKT. A false-discovery rate (FDR) *p*-value of 0.05 and fold change (FC) of 2.0 was used for

discerning dysregulated metabolites. A negative fold change is displayed in blue, and a positive fold

change                       is                       displayed                       in                       red.

The xenometabolome of three different salmonids exposed to 6PPD-Q was profiled, revealing

that three phase I and seven phase II TPs dominated at lower exposure to 6PPD-Q. Different

biotransformation pathways can dominate among salmonids, with RBT and BRT having greater levels of

glucuronide-conjugated TPs than the sulfate-conjugated TP found to dominate in LKT. This can have

important implications regarding species-specific biotransformation and subsequent response to

contaminants, including 6PPD-Q. More research is needed to understand the main mechanisms of

toxicological responses to 6PPD-Q and its phase I TPs, including TP-OH1, which was detected in both

RBT and LKT but not BRT. Synthesized analytical standards were used to verify the structure of both

TP-OH1 and TP-OH2 and will be useful for future biomonitoring and exposure assessment. A time-

course metabolome response to 6PPD-Q is also needed to understand how larval fish initially respond to

the contaminant and subsequent compensatory responses to have a greater mechanistic understanding of

6PPD-Q toxicity. These results indicate that a difference in biotransformation capability and pathways is

observed among salmonid fish species, and subsequent metabolome response to 6PPD-Q exposure can

help            reveal            the            mode            of            toxic            action.

**Acknowledgment**

This project was supported partially by a financial contribution from Fisheries and Oceans Canada. Additional funding was provided to M.H., S.W., N.H., and M.B. through the Discovery Grants program of the Natural Sciences and Engineering Research Council of Canada (NSERC). M.H. was also supported by the Canada Research Chairs program. Instrumentation used for chemical analyses within this research was obtained with funding from Western Economic Diversification Canada (WED) and the Canadian Foundation for Innovation (CFI). P.J.A. was supported through a Toxicology Centre Devolved Scholarship. M.B. is currently a faculty member of the Global Water Futures (GWF) program, which received funds from the Canada First Research Excellence Funds (CFREF). The graphical abstract was created with Bioicons.com. Additional support came from the staff at Allison Creek Brood Trout Station, who provided the Brown trout embryos, as well as the technical staff at the ARF at the University of Lethbridge (Dr. S. Mamun and H. Shepherd).

**References**

(1)  Tian, Z.; Zhao, H.; Peter, K. T.; Gonzalez, M.; Wetzel, J.; Wu, C.; Hu, X.; Prat, J.; Mudrock, E.; Hettinger, R.; Cortina, A. E.; Biswas, R. G.; Kock, F. V. C.; Soong, R.; Jenne, A.; Du, B.; Hou, F.; He, H.; Lundeen, R.; Gilbreath, A.; Sutton, R.; Scholz, N. L.; Davis, J. W.; Dodd, M. C.; Simpson, A.; McIntyre, J. K.; Kolodziej, E. P. A Ubiquitous Tire Rubber–Derived Chemical Induces Acute Mortality in Coho Salmon. *Science* **2021**, *371* (6525), 185–189. https://doi.org/10.1126/science.abd6951.

(2)  Zoroufchi Benis, K.; Behnami, A.; Minaei, S.; Brinkmann, M.; McPhedran, K. N.; Soltan, J. Environmental Occurrence and Toxicity of 6PPD Quinone, an Emerging Tire Rubber-Derived Chemical: A Review. *Environ. Sci. Technol. Lett.* **2023**, *10* (10), 815–823. https://doi.org/10.1021/acs.estlett.3c00521.

(3)  Rauert, C.; Charlton, N.; Okoffo, E. D.; Stanton, R. S.; Agua, A. R.; Pirrung, M. C.; Thomas, K. V. Concentrations of Tire Additive Chemicals and Tire Road Wear Particles in an Australian Urban Tributary. *Environ. Sci. Technol.* **2022**, *56* (4), 2421–2431. https://doi.org/10.1021/acs.est.1c07451.

(4)  Hu, X.; Zhao, H. N.; Tian, Z.; Peter, K. T.; Dodd, M. C.; Kolodziej, E. P. Transformation Product Formation upon Heterogeneous Ozonation of the Tire Rubber Antioxidant 6PPD (N-(1,3-Dimethylbutyl)-N′-Phenyl-p-Phenylenediamine). *Environ. Sci. Technol. Lett.* **2022**, *9* (5), 413–419. https://doi.org/10.1021/acs.estlett.2c00187.

(5)  Brinkmann, M.; Montgomery, D.; Selinger, S.; Miller, J. G. P.; Stock, E.; Alcaraz, A. J.; Challis, J. K.; Weber, L.; Janz, D.; Hecker, M.; Wiseman, S. Acute Toxicity of the Tire Rubber-Derived Chemical 6PPD-Quinone to Four Fishes of Commercial, Cultural, and Ecological Importance. *Environ. Sci. Technol. Lett.* **2022**, *9* (4), 333–338. https://doi.org/10.1021/acs.estlett.2c00050.

(6)  Tian, Z.; Gonzalez, M.; Rideout, C. A.; Zhao, H. N.; Hu, X.; Wetzel, J.; Mudrock, E.; James, C. A.; McIntyre, J. K.; Kolodziej, E. P. 6PPD-Quinone: Revised Toxicity Assessment and Quantification with a Commercial Standard. *Environ. Sci. Technol. Lett.* **2022**, *9* (2), 140–146. https://doi.org/10.1021/acs.estlett.1c00910.

(7)  Hiki, K.; Yamamoto, H. The Tire-Derived Chemical 6PPD-Quinone Is Lethally Toxic to the White-Spotted Char Salvelinus Leucomaenis Pluvius but Not to Two Other Salmonid Species. *Environ. Sci. Technol. Lett.* **2022**, *9* (12), 1050–1055. https://doi.org/10.1021/acs.estlett.2c00683.

(8)  Roberts, C.; Lin, J.; Kohlman, E.; Jain, N.; Amekor, M.; Alcaraz, A. J.; Hogan, N.; Hecker, M.; Brinkmann, M. Acute and Subchronic Toxicity of 6PPD-Quinone to Early Life Stage Lake Trout (Salvelinus Namaycush). *Environ Sci Technol* **2025**, *59* (1), 791–797. https://doi.org/10.1021/acs.est.4c09090.

(9)  Varshney, S.; Gora, A. H.; Siriyappagouder, P.; Kiron, V.; Olsvik, P. A. Toxicological Effects of 6PPD and 6PPD Quinone in Zebrafish Larvae. *J Hazard Mater* **2022**, *424* (Pt C), 127623. https://doi.org/10.1016/j.jhazmat.2021.127623.

(10) Foldvik, A.; Kryuchkov, F.; Sandodden, R.; Uhlig, S. Acute Toxicity Testing of the Tire Rubber–Derived Chemical 6PPD-Quinone on Atlantic Salmon (Salmo Salar) and Brown Trout (Salmo Trutta). *Environmental Toxicology and Chemistry* **2022**, *41* (12), 3041–3045. https://doi.org/10.1002/etc.5487.

(11) Montgomery, D.; Ji, X.; Cantin, J.; Philibert, D.; Foster, G.; Selinger, S.; Jain, N.; Miller, J.; McIntyre, J.; de Jourdan, B.; Wiseman, S.; Hecker, M.; Brinkmann, M. Interspecies Differences in 6PPD-Quinone Toxicity Across Seven Fish Species: Metabolite Identification and Semiquantification. *Environ Sci Technol* **2023**, *57* (50), 21071–21079. https://doi.org/10.1021/acs.est.3c06891.

(12) Grasse, N.; Seiwert, B.; Massei, R.; Scholz, S.; Fu, Q.; Reemtsma, T. Uptake and Biotransformation of the Tire Rubber-Derived Contaminants 6-PPD and 6-PPD Quinone in the Zebrafish Embryo (Danio Rerio). *Environ. Sci. Technol.* **2023**, *57* (41), 15598–15607. https://doi.org/10.1021/acs.est.3c02819.

(13) Ankley, P. J.; da Silva, F. C. Jr.; Montgomery, D.; Schultz, M.; Ed S. Krol; Hecker, M.; Brinkmann,
M. Biotransformation of 6PPD-Quinone In Vitro Using RTL-W1 Cell Line. *Environ. Sci. Technol.*
*Lett.* **2024**. https://doi.org/10.1021/acs.estlett.4c00342.
(14) Zhang, J.; Cao, G.; Wang, W.; Qiao, H.; Chen, Y.; Wang, X.; Wang, F.; Liu, W.; Cai, Z. Stable
Isotope-Assisted Mass Spectrometry Reveals in Vivo Distribution, Metabolism, and Excretion of
Tire Rubber-Derived 6PPD-Quinone in Mice. *Science of The Total Environment* **2024**, *912*, 169291.
https://doi.org/10.1016/j.scitotenv.2023.169291.
(15) Ribbenstedt, A.; Ziarrusta, H.; Benskin, J. P. Development, Characterization and Comparisons of
Targeted and Non-Targeted Metabolomics Methods. *PLOS ONE* **2018**, *13* (11), e0207082.
https://doi.org/10.1371/journal.pone.0207082.
(16) Evich, M. G.; Mosley, J. D.; Ntai, I.; Cavallin, J. E.; Villeneuve, D. L.; Ankley, G. T.; Collette, T.
W.; Ekman, D. R. Untargeted MSn-Based Monitoring of Glucuronides in Fish: Screening Complex
Mixtures for Contaminants with Biological Relevance. *ACS ES T Water* **2022**, *2* (12), 2481–2490.
https://doi.org/10.1021/acsestwater.2c00310.
(17) Al-Salhi, R.; Abdul-Sada, A.; Lange, A.; Tyler, C. R.; Hill, E. M. The Xenometabolome and Novel
Contaminant Markers in Fish Exposed to a Wastewater Treatment Works Effluent. *Environ. Sci.*
*Technol.* **2012**, *46* (16), 9080–9088. https://doi.org/10.1021/es3014453.
(18) Fiehn, O. Metabolomics--the Link between Genotypes and Phenotypes. *Plant Mol Biol* **2002**, *48* (1–
2), 155–171.
(19) Bonvallot, N.; Tremblay-Franco, M.; Chevrier, C.; Canlet, C.; Debrauwer, L.; Cravedi, J.-P.; Cordier,
S. Potential Input from Metabolomics for Exploring and Understanding the Links between
Environment and Health. *J Toxicol Environ Health B Crit Rev* **2014**, *17* (1), 21–44.
https://doi.org/10.1080/10937404.2013.860318.
(20) Ortiz-Villanueva, E.; Jaumot, J.; Martínez, R.; Navarro-Martín, L.; Piña, B.; Tauler, R. Assessment of
Endocrine Disruptors Effects on Zebrafish (Danio Rerio) Embryos by Untargeted LC-HRMS
Metabolomic Analysis. *Science of The Total Environment* **2018**, *635*, 156–166.
https://doi.org/10.1016/j.scitotenv.2018.03.369.
(21) Xu, M.; Legradi, J.; Leonards, P. A Comprehensive Untargeted Metabolomics Study in Zebrafish
Embryos Exposed to Perfluorohexane Sulfonate (PFHxS). *Science of The Total Environment* **2023**,
*887*, 163770. https://doi.org/10.1016/j.scitotenv.2023.163770.
(22) Magnuson, J. T.; Giroux, M.; Cryder, Z.; Gan, J.; Schlenk, D. The Use of Non-Targeted
Metabolomics to Assess the Toxicity of Bifenthrin to Juvenile Chinook Salmon (*Oncorhynchus*
*Tshawytscha*). *Aquatic Toxicology* **2020**, *224*, 105518.
https://doi.org/10.1016/j.aquatox.2020.105518.
(23) Roberts, C.; Kohlman, E.; Jain, N.; Amekor, M.; Alcaraz, A. J.; Hogan, N.; Brinkmann, M.; Hecker,
M. Sub-Chronic and Acute Toxicity of 6PPD-Quinone to Early-Life Stage Rainbow Trout
(Oncorhynchus Mykiss). bioRxiv September 26, 2024, p 2024.09.25.614982.
https://doi.org/10.1101/2024.09.25.614982.
(24) OECD. Test No. 210: Fish, Early-Life Stage Toxicity Test, 2013. https://www.oecd-
ilibrary.org/content/publication/9789264203785-en.
(25) Schmid, R.; Heuckeroth, S.; Korf, A.; Smirnov, A.; Myers, O.; Dyrlund, T. S.; Bushuiev, R.; Murray,
K. J.; Hoffmann, N.; Lu, M.; Sarvepalli, A.; Zhang, Z.; Fleischauer, M.; Dührkop, K.; Wesner, M.;
Hoogstra, S. J.; Rudt, E.; Mokshyna, O.; Brungs, C.; Ponomarov, K.; Mutabdžija, L.; Damiani, T.;
Pudney, C. J.; Earll, M.; Helmer, P. O.; Fallon, T. R.; Schulze, T.; Rivas-Ubach, A.; Bilbao, A.;
Richter, H.; Nothias, L.-F.; Wang, M.; Orešič, M.; Weng, J.-K.; Böcker, S.; Jeibmann, A.; Hayen,
H.; Karst, U.; Dorrestein, P. C.; Petras, D.; Du, X.; Pluskal, T. Integrative Analysis of Multimodal
Mass Spectrometry Data in MZmine 3. *Nat Biotechnol* **2023**, *41* (4), 447–449.
https://doi.org/10.1038/s41587-023-01690-2.
(26) Dührkop, K.; Fleischauer, M.; Ludwig, M.; Aksenov, A. A.; Melnik, A. V.; Meusel, M.; Dorrestein,
P. C.; Rousu, J.; Böcker, S. SIRIUS 4: A Rapid Tool for Turning Tandem Mass Spectra into

546   Metabolite Structure Information. *Nat Methods* **2019**, *16* (4), 299–302.
547   https://doi.org/10.1038/s41592-019-0344-8.

548 (27)Nothias, L.-F.; Petras, D.; Schmid, R.; Dührkop, K.; Rainer, J.; Sarvepalli, A.; Protsyuk, I.; Ernst, M.;
549   Tsugawa, H.; Fleischauer, M.; Aicheler, F.; Aksenov, A. A.; Alka, O.; Allard, P.-M.; Barsch, A.;
550   Cachet, X.; Caraballo-Rodriguez, A. M.; Da Silva, R. R.; Dang, T.; Garg, N.; Gauglitz, J. M.;
551   Gurevich, A.; Isaac, G.; Jarmusch, A. K.; Kameník, Z.; Kang, K. B.; Kessler, N.; Koester, I.; Korf,
552   A.; Le Gouellec, A.; Ludwig, M.; Martin H, C.; McCall, L.-I.; McSayles, J.; Meyer, S. W.;
553   Mohimani, H.; Morsy, M.; Moyne, O.; Neumann, S.; Neuweger, H.; Nguyen, N. H.; Nothias-
554   Esposito, M.; Paolini, J.; Phelan, V. V.; Pluskal, T.; Quinn, R. A.; Rogers, S.; Shrestha, B.; Tripathi,
555   A.; van der Hooft, J. J. J.; Vargas, F.; Weldon, K. C.; Witting, M.; Yang, H.; Zhang, Z.; Zubeil, F.;
556   Kohlbacher, O.; Böcker, S.; Alexandrov, T.; Bandeira, N.; Wang, M.; Dorrestein, P. C. Feature-
557   Based Molecular Networking in the GNPS Analysis Environment. *Nat Methods* **2020**, *17* (9), 905–
558   908. https://doi.org/10.1038/s41592-020-0933-6.

559 (28)Shannon, P.; Markiel, A.; Ozier, O.; Baliga, N. S.; Wang, J. T.; Ramage, D.; Amin, N.; Schwikowski,
560   B.; Ideker, T. Cytoscape: A Software Environment for Integrated Models of Biomolecular
561   Interaction Networks. *Genome Res* **2003**, *13* (11), 2498–2504. https://doi.org/10.1101/gr.1239303.

562 (29)Wang, F.; Liigand, J.; Tian, S.; Arndt, D.; Greiner, R.; Wishart, D. S. CFM-ID 4.0: More Accurate
563   ESI MS/MS Spectral Prediction and Compound Identification. *Analytical chemistry* **2021**, *93* (34),
564   11692. https://doi.org/10.1021/acs.analchem.1c01465.

565 (30)Dührkop, K.; Shen, H.; Meusel, M.; Rousu, J.; Böcker, S. Searching Molecular Structure Databases
566   with Tandem Mass Spectra Using CSI:FingerID. *Proceedings of the National Academy of Sciences*
567   **2015**, *112* (41), 12580–12585. https://doi.org/10.1073/pnas.1509788112.

568 (31)Pang, Z.; Chong, J.; Zhou, G.; de Lima Morais, D. A.; Chang, L.; Barrette, M.; Gauthier, C.; Jacques,
569   P.-É.; Li, S.; Xia, J. MetaboAnalyst 5.0: Narrowing the Gap between Raw Spectra and Functional
570   Insights. *Nucleic Acids Research* **2021**, *49* (W1), W388–W396.
571   https://doi.org/10.1093/nar/gkab382.

572 (32)Hu, X.; Zhao, H. (Nina); Tian, Z.; Peter, K. T.; Dodd, M. C.; Kolodziej, E. P. Chemical
573   Characteristics, Leaching, and Stability of the Ubiquitous Tire Rubber-Derived Toxicant 6PPD-
574   Quinone. *Environ. Sci.: Processes Impacts* **2023**, *25* (5), 901–911.
575   https://doi.org/10.1039/D3EM00047H.

576 (33)van der Oost, R.; Beyer, J.; Vermeulen, N. P. E. Fish Bioaccumulation and Biomarkers in
577   Environmental Risk Assessment: A Review. *Environmental Toxicology and Pharmacology* **2003**,
578   *13* (2), 57–149. https://doi.org/10.1016/S1382-6689(02)00126-6.

579 (34)Chow, M. I.; Lundin, J. I.; Mitchell, C. J.; Davis, J. W.; Young, G.; Scholz, N. L.; McIntyre, J. K. An
580   Urban Stormwater Runoff Mortality Syndrome in Juvenile Coho Salmon. *Aquatic Toxicology* **2019**,
581   *214*, 105231. https://doi.org/10.1016/j.aquatox.2019.105231.

582 (35)Zhang, Y.-Y.; Huang, J.-W.; Liu, Y.-H.; Zhang, J.-N.; Huang, Z.; Liu, Y.-S.; Zhao, J.-L.; Ying, G.-G.
583   *In Vitro* Metabolism of the Emerging Contaminant 6PPD-Quinone in Human and Rat Liver
584   Microsomes: Kinetics, Pathways, and Mechanism. *Environmental Pollution* **2024**, *345*, 123514.
585   https://doi.org/10.1016/j.envpol.2024.123514.

586 (36)Nair, P.; Barrett, H.; Tanoto, K.; Xie, L.; Sun, J.; Yang, D.; Yao, H.; Song, D.; Peng, H. Structure and
587   Toxicity Characterization of Alkyl Hydroxylated Metabolites of 6PPD-Q. ChemRxiv October 30,
588   2024. https://doi.org/10.26434/chemrxiv-2024-s5rhc.

589 (37)Xie, L.; Yu, J.; Nair, P.; Sun, J.; Barrett, H.; Meek, O.; Qian, X.; Yang, D.; Kennedy, L.;
590   Kozakiewicz, D.; Hao, C.; Hansen, J. D.; Greer, J. B.; Abbatt, J. P. D.; Peng, H. Structurally
591   Selective Ozonolysis of P-Phenylenediamines and Toxicity in Coho Salmon and Rainbow Trout.
592   *Environ. Sci. Technol.* **2024**, *58* (49), 21423–21432. https://doi.org/10.1021/acs.est.4c04817.

593 (38)Song, Z.; Yu, X.; Zhu, M.; Wu, Z.; Fu, Z.; Chen, J. Distinct Species-Specific and Toxigenic
594   Metabolic Profiles for 6PPD and 6PPD Quinone by P450 Enzymes: Insights from In Vitro and In
595   Silico Studies. *Environ. Sci. Technol.* **2024**, *58* (34), 14994–15004.
596   https://doi.org/10.1021/acs.est.4c03361.

(39) Franco, M. E.; Hollender, J.; Schirmer, K. Differential Biotransformation Ability May Alter Fish Biodiversity in Polluted Waters. *Environment International* **2025**, *195*, 109254. https://doi.org/10.1016/j.envint.2025.109254.

(40) Deng, M.; Ji, X.; Peng, B.; Fang, M. In Vitro and In Vivo Biotransformation Profiling of 6PPD-Quinone toward Their Detection in Human Urine. *Environ. Sci. Technol.* **2024**, *58* (21), 9113–9124. https://doi.org/10.1021/acs.est.4c01106.

(41) Singh, N.; Shreshtha, A. K.; Thakur, M. S.; Patra, S. Xanthine Scaffold: Scope and Potential in Drug Development. *Heliyon* **2018**, *4* (10), e00829. https://doi.org/10.1016/j.heliyon.2018.e00829.

(42) Jacobson, K. A.; Gao, Z.-G. Adenosine Receptors as Therapeutic Targets. *Nat Rev Drug Discov* **2006**, *5* (3), 247–264. https://doi.org/10.1038/nrd1983.

(43) Boison, D. Adenosine and Seizure Termination: Endogenous Mechanisms. *Epilepsy Curr* **2013**, *13* (1), 35–37. https://doi.org/10.5698/1535-7511-13.1.35.

(44) Scholz, N. L.; Myers, M. S.; McCarthy, S. G.; Labenia, J. S.; McIntyre, J. K.; Ylitalo, G. M.; Rhodes, L. D.; Laetz, C. A.; Stehr, C. M.; French, B. L.; McMillan, B.; Wilson, D.; Reed, L.; Lynch, K. D.; Damm, S.; Davis, J. W.; Collier, T. K. Recurrent Die-Offs of Adult Coho Salmon Returning to Spawn in Puget Sound Lowland Urban Streams. *PLOS ONE* **2011**, *6* (12), e28013. https://doi.org/10.1371/journal.pone.0028013.

(45) Mahoney, H.; da Silva Junior, F. C.; Roberts, C.; Schultz, M.; Ji, X.; Alcaraz, A. J.; Montgomery, D.; Selinger, S.; Challis, J. K.; Giesy, J. P.; Weber, L.; Janz, D.; Wiseman, S.; Hecker, M.; Brinkmann, M. Exposure to the Tire Rubber-Derived Contaminant 6PPD-Quinone Causes Mitochondrial Dysfunction In Vitro. *Environ. Sci. Technol. Lett.* **2022**, *9* (9), 765–771. https://doi.org/10.1021/acs.estlett.2c00431.

(46) Indiveri, C.; Iacobazzi, V.; Tonazzi, A.; Giangregorio, N.; Infantino, V.; Convertini, P.; Console, L.; Palmieri, F. The Mitochondrial Carnitine/Acylcarnitine Carrier: Function, Structure and Physiopathology. *Molecular Aspects of Medicine* **2011**, *32* (4), 223–233. https://doi.org/10.1016/j.mam.2011.10.008.

(47) Reuter, S. E.; Evans, A. M. Carnitine and Acylcarnitines. *Clin Pharmacokinet* **2012**, *51* (9), 553–572. https://doi.org/10.1007/BF03261931.

(48) Ferreira, G. C.; McKenna, M. C. L-Carnitine and Acetyl-l-Carnitine Roles and Neuroprotection in Developing Brain. *Neurochem Res* **2017**, *42* (6), 1661–1675. https://doi.org/10.1007/s11064-017-2288-7.

(49) Smeland, O. B.; Meisingset, T. W.; Borges, K.; Sonnewald, U. Chronic Acetyl-l-Carnitine Alters Brain Energy Metabolism and Increases Noradrenaline and Serotonin Content in Healthy Mice. *Neurochemistry International* **2012**, *61* (1), 100–107. https://doi.org/10.1016/j.neuint.2012.04.008.

(50) Huang, F.; Subbaiah, P. V.; Holian, O.; Zhang, J.; Johnson, A.; Gertzberg, N.; Lum, H. Lysophosphatidylcholine Increases Endothelial Permeability: Role of PKCα and RhoA Cross Talk. *American Journal of Physiology-Lung Cellular and Molecular Physiology* **2005**, *289* (2), L176–L185. https://doi.org/10.1152/ajplung.00003.2005.

(51) Chang, M.-C.; Lee, J.-J.; Chen, Y.-J.; Lin, S.-I.; Lin, L.-D.; Jein-Wen Liou, E.; Huang, W.-L.; Chan, C.-P.; Huang, C.-C.; Jeng, J.-H. Lysophosphatidylcholine Induces Cytotoxicity/Apoptosis and IL-8 Production of Human Endothelial Cells: Related Mechanisms. *Oncotarget* **2017**, *8* (63), 106177–106189. https://doi.org/10.18632/oncotarget.22425.

(52) Law, S.-H.; Chan, M.-L.; Marathe, G. K.; Parveen, F.; Chen, C.-H.; Ke, L.-Y. An Updated Review of Lysophosphatidylcholine Metabolism in Human Diseases. *Int J Mol Sci* **2019**, *20* (5), 1149. https://doi.org/10.3390/ijms20051149.

(53) Blusztajn, J. K.; Liscovitch, M.; Richardson, U. I. Synthesis of Acetylcholine from Choline Derived from Phosphatidylcholine in a Human Neuronal Cell Line. *Proceedings of the National Academy of Sciences* **1987**, *84* (15), 5474–5477. https://doi.org/10.1073/pnas.84.15.5474.

(54) Magaquian, D.; Delgado Ocaña, S.; Perez, C.; Banchio, C. Phosphatidylcholine Restores Neuronal Plasticity of Neural Stem Cells under Inflammatory Stress. *Sci Rep* **2021**, *11* (1), 22891. https://doi.org/10.1038/s41598-021-02361-5.

(55) Greer, J. B.; Dalsky, E. M.; Lane, R. F.; Hansen, J. D. Tire-Derived Transformation Product 6PPD-
Quinone Induces Mortality and Transcriptionally Disrupts Vascular Permeability Pathways in
Developing Coho Salmon. *Environ. Sci. Technol.* **2023**, *57* (30), 10940–10950.
https://doi.org/10.1021/acs.est.3c01040.
(56) Martinez-Molledo, M.; Nji, E.; Reyes, N. Structural Insights into the Lysophospholipid Brain Uptake
Mechanism and Its Inhibition by Syncytin-2. *Nat Struct Mol Biol* **2022**, *29* (6), 604–612.
https://doi.org/10.1038/s41594-022-00786-8.
(57) Wang, B.; Wu, L.; Chen, J.; Dong, L.; Chen, C.; Wen, Z.; Hu, J.; Fleming, I.; Wang, D. W.
Metabolism Pathways of Arachidonic Acids: Mechanisms and Potential Therapeutic Targets. *Sig*
*Transduct Target Ther* **2021**, *6* (1), 1–30. https://doi.org/10.1038/s41392-020-00443-w.
(58) Swanson, D.; Block, R.; Mousa, S. A. Omega-3 Fatty Acids EPA and DHA: Health Benefits
Throughout Life. *Advances in Nutrition* **2012**, *3* (1), 1–7. https://doi.org/10.3945/an.111.000893.
(59) Calder, P. C. Functional Roles of Fatty Acids and Their Effects on Human Health. *Journal of*
*Parenteral and Enteral Nutrition* **2015**, *39* (1S), 18S-32S.
https://doi.org/10.1177/0148607115595980.
(60) Picq, M.; Chen, P.; Perez, M.; Michaud, M.; Véricel, E.; Guichardant, M.; Lagarde, M. DHA
Metabolism: Targeting the Brain and Lipoxygenation. *Mol Neurobiol* **2010**, *42* (1), 48–51.
https://doi.org/10.1007/s12035-010-8131-7.
(61) VanRollins, M.; Murphy, R. C. Autooxidation of Docosahexaenoic Acid: Analysis of Ten Isomers of
Hydroxydocosahexaenoate. *J Lipid Res* **1984**, *25* (5), 507–517.
(62) Prestwich, E. G.; Mangerich, A.; Pang, B.; McFaline, J. L.; Lonkar, P.; Sullivan, M. R.; Trudel, L. J.;
Taghizedeh, K.; Dedon, P. C. Increased Levels of Inosine in a Mouse Model of Inflammation. *Chem*
*Res Toxicol* **2013**, *26* (4), 538–546. https://doi.org/10.1021/tx300473n.
(63) Kim, I. S.; Jo, E.-K. Inosine: A Bioactive Metabolite with Multimodal Actions in Human Diseases.
*Front Pharmacol* **2022**, *13*, 1043970. https://doi.org/10.3389/fphar.2022.1043970.
(64) Ewald, J. D.; Basu, N.; Crump, D.; Boulanger, E.; Head, J. Characterizing Variability and
Uncertainty Associated with Transcriptomic Dose–Response Modeling. *Environ. Sci. Technol.*
**2022**, *56* (22), 15960–15968. https://doi.org/10.1021/acs.est.2c04665.
(65) Yao, C.-H.; Wang, L.; Stancliffe, E.; Sindelar, M.; Cho, K.; Yin, W.; Wang, Y.; Patti, G. J. Dose-
Response Metabolomics To Understand Biochemical Mechanisms and Off-Target Drug Effects
with the TOXcms Software. *Anal Chem* **2020**, *92* (2), 1856–1864.
https://doi.org/10.1021/acs.analchem.9b03811.
(66) Zhao, H.; Liu, M.; Lv, Y.; Fang, M. Dose-Response Metabolomics and Pathway Sensitivity to Map
Molecular Cartography of Bisphenol A Exposure. *Environment International* **2022**, *158*, 106893.
https://doi.org/10.1016/j.envint.2021.106893.
683