Anatomically distinct cortical tracking of music and speech by slow (1-

8Hz) and fast (70-120Hz) oscillatory activity

Sergio Osorio[1,2,*] & M. Florencia Assaneo[3,*]

1. Department of Neurology, Harvard Medical School, Massachusetts General Hospital,
Boston, MA
2. Martinos Center for Biomedical Imaging, Massachusetts General Hospital, Boston, MA
3. Instituto de Neurobiología, Universidad Autónoma de México, Santiago de Querétaro,
MX.

* Corresponding authors: *sosoriogaleano@mgh.harvard.edu, fassaneo@inb.unam.mx*

Abstract

Music and speech encode hierarchically organized structural complexity at the service of

human expressiveness and communication. Previous research has shown that populations

of neurons in auditory regions track the envelope of acoustic signals within the range of

slow and fast oscillatory activity. However, the extent to which cortical tracking is

influenced by the interplay between stimulus type, frequency band, and brain anatomy

remains an open question. In this study, we reanalyzed intracranial recordings from thirty

subjects implanted with electrocorticography (ECoG) grids in the left cerebral hemisphere,

drawn from an existing open-access ECoG database. Participants passively watched a

movie where visual scenes were accompanied by either music or speech stimuli. Cross-

correlation between brain activity and the envelope of music and speech signals, along with

density-based clustering analyses and linear mixed effect modeling, revealed both

anatomically overlapping and functionally distinct mapping of the tracking effect as a

function of stimulus type and frequency band. We observed widespread left-hemisphere

tracking of music and speech signals in the Slow Frequency Band (SFB, band-passed

filtered low-frequency signal between 1-8Hz), with near zero temporal lags. In contrast,

cortical tracking in the High Frequency Band (HFB, envelope of the 70-120Hz band-passed

filtered signal) was higher during speech perception, was more densely concentrated in

classical language processing areas, and showed a frontal-to-temporal gradient in lag values

that was not observed during perception of musical stimuli. Our results highlight a complex

interaction between cortical region and frequency band that shapes temporal dynamics

during processing of naturalistic music and speech signals.

Introduction

Music and speech encode hierarchically organized structural complexity and support emotional and semantic expressiveness in human communication (1–7). To achieve this goal, small structural units (e.g., notes or syllables) are linearly combined to build up longer units (e.g., phrases or sentences), establishing non-linear relationships between high-level abstract representations. Hence, understanding how the brain physiologically supports perception of music and speech is of great importance for cognitive and auditory neuroscience.

Because both signals deploy in time, hierarchical structure is physically embedded in the modulation of a signal's amplitude over time (i.e. the acoustic envelope), which can be quantitively assessed via power density and Fourier analyses. Across multiple languages and different musical genres, music and speech signals are characterized by spectral peaks at around 2Hz and 5Hz, respectively, which roughly correspond to the mean note rate (8,9) and the mean syllabic rate (9,10). Despite subtle differences in their temporal dynamics, perception of both music and speech requires the ability to track multiple structural units and their dependencies. This has led to propose that the neural architecture supporting perception of both signals must derive from similar evolutionary affordances and must be supported by shared neural mechanisms (2,11,12).

During the perception of music and speech, neuronal populations track temporal regularities across different hierarchical levels within the signal, a phenomenon known as cortical tracking. Cortical tracking of structural units, such as notes and phrases, and of acoustic features, such as rhythm, melody, and modulations in the signal's envelope, has been observed within the Slow Frequency Band (SFB, band-passed filtered low-frequency

signal between 1-8Hz) activity during perception of musical stimuli (13–17). This tracking

has been shown to be modulated by several factors, including note rate (14), musical

expertise (13), stimulus familiarity (18) and stimulus complexity (19). Similarly, during

speech perception, brain activity within the same frequency band (i.e., SFB) tracks

structural units, such as syllables, words and phrases, and acoustic features such as word

onsets and amplitude modulations in the signal's envelope (17,20–26). SFB tracking of

speech signals is modulated by factors such as syllabic rate (20,22), speech intelligibility

(22,27,28) and attention (29,30).

Neural tracking of music and speech has also been observed within the range of

High Frequency gamma Band (HFB, >70 Hz) activity, particularly through the modulation

of HFB amplitude (22,23,31–34). During perception of music, cortical tracking of the

acoustic envelope and various other acoustic features, including pitch, voice, and melody,

has been associated with both time-locked and sustained HFB neural responses in regions

such as the superior temporal, middle temporal, supramarginal, and dorsolateral prefrontal

cortices (32,33,35–38). Similarly, during speech perception, HFB tracking of structural

units, such as syllables, words and phrases, and of the acoustic features such as pitch, peak

rate, syllable onsets, and the signal's envelope, has been observed in the superior temporal

regions and ventrolateral prefrontal cortices (22,23,31,34,39,40), with a gradient of

increasing tracking magnitude from perisylvian and prefrontal areas toward primary

auditory regions (31). Despite their limited spatial resolution and signal-to-noise ratios

compared to intracranial recordings, MEG and EEG studies corroborate HFB tracking of

the speech envelope in temporal electrodes and source-reconstructed auditory areas (41,42).

Studies using invasive methods such as electrocorticography and stereotactic EEG

have demonstrated that neural tracking of acoustic signals is widely distributed in cortical

space, exhibiting little or no anatomical regional selectivity based on stimulus type

(35,36,39). This findings contrasts with findings from fMRI studies, which indicate

anatomical specialization for speech in ventrolateral prefrontal regions and for music in

dorsolateral prefrontal and inferior parietal regions (43–48). Despite this apparent

discrepancy, recent evidence points to selectivity depending on the type of information

being tracked. One previous study reported that posterior portions of the Superior Temporal

Gyrus (STG) are more sensitive to syntactic complexity in speech stimuli, whereas

posterior portions of the Middle Temporal Gyrus (MTG) show greater sensitivity to music

complexity (49). Furthermore, selectivity across frequency bands has also been observed:

while the envelope of the music and speech acoustic signals is similarly tracked by both

SFB and HFB activity, acoustic edges, which correspond to the peak rate of temporal

modulations in the acoustic signals, are more accurately predicted by activity in the SFB

(39). These findings therefore suggest a more intricate interplay between stimulus type,

frequency band, and brain anatomy, highlighting the need for further investigation to fully

characterize these interactions.

Here, we analyzed an open-access database of electrocortical recordings acquired

during naturalistic, multimodal perception of music and speech stimuli (50). Prior to

surgical procedures, patients passively watched 30-second videos containing either music

(multi-instrument clips) or speech (single voice and dialogues) audios, while brain activity

was recorded. Data from a subsample of 30 subjects with electrode grids located in left

prefrontal, motor, premotor, somatosensory and/or temporal regions was analyzed, thus

offering a coverage of most left-hemisphere cortical areas involved in music and speech

perception. Specifically, we aimed to understand the anatomical organization and temporal

dynamics underlying the neural tracking of music and speech, examining both slow (1-

8Hz) and fast (70-120Hz) oscillatory bands known to be involved in perception and

processing of both acoustic signals.

To quantify cortical tracking of music and speech in a straightforward way, we

conducted cross-correlation analyses between the cochlear envelope of acoustic signals and

the bandpass-filtered brain signals (SFB, the brain activity filtered between 1-8 Hz; and

HFB the amplitude modulation of the 70-120 Hz filtered brain signal). To investigate

whether frequency-band-dependent tracking effects can be mapped to similar or different

cortical areas, we conducted density-based clustering analyses. Finally, to investigate the

effect of condition and cortical region in the strength of the tracking effect and its temporal

dynamics, we conducted mixed effect modeling analyzes using maximum correlation and

temporal lags as the predicted variables, and frequency band (SFB, HFB), stimulus type

(music, speech), and anatomical location of electrodes (cortical region) as regressors, while

controlling for the within-subject nature of the data and differences in EcoG grid locations

across subjects as potential random effects.

Materials and methods

Data was obtained from a publicly available dataset (50) collected at the University

Medical Center Utrecht from 63 patients that underwent intracortical electrode implantation

as part of diagnosis procedures for resection surgery due to a clinical history of drug-

resistant epileptic seizures. For this study, we used data from a subsample of 30 subjects

(mean age = 27.33, SD=15.28, 19 females, see supplementary table s1), selected on the

basis of electrode grid coverage (i.e., only patients with grids in left prefrontal, motor,

premotor, somatosensory and/or temporal regions). Language dominance for all subjects

was localized to the left hemisphere (table s1). This resulted in a total number of 1,858

electrodes across all subjects (mean = 61.93, SD = 19.09, supplementary table s2)

distributed across left-hemisphere prefrontal, central and temporal regions (figure s1).

Participants were passively presented 30-second interleaved excerpts from the

movie "Pipi on the run" (Pårymmen med Pippi Långstrump, 1970) dubbed to the Dutch

language, where visual scenes were accompanied by music (7 blocks) or speech (6 blocks)

stimuli. All visual and auditory excerpts were different from one another. The task was

originally designed as part of a language mapping procedure prior to surgical intervention.

All participants and legal guardians, when applicable, provided their informed consent to

participate in the study and to make their de-identified data publicly available (50).

Procedures were revised and approved by the Medical Ethical Committee of the University

Medical Center Utrecht in accordance with the Declaration of Helsinki (2013).

Data analysis

Acoustic signals

Acoustic data was analyzed using MATLAB Version 9.6.0 (2019a). The acoustic

signal was extracted from the audiovisual stimulus and segmented according to stimulus

type (speech or music). The last musical segment was excluded from analyses to have a

balanced number of segments across the two conditions. The segmented signals were then

down-sampled to 16,000 Hz. Next, each segment's auditory representation in the human

cochlea was obtained by detrending, resampling (200hz), and filtering each signal in 128

logarithmically spaced frequencies within the range of human audition (180 to 7,246 Hz).

The cochlear spectrogram was obtained using the NSL toolbox (51). Finally, signals were

averaged across the 128 frequencies to obtain the cochlear envelope of each acoustic

segment. For each envelope, the Power Spectrum Density was obtained using Welch's

method.

EcoG and neuroanatomical data preprocessing

Electrophysiological and neuroanatomical data was preprocessed using Brainstorm

(52). Electrodes were first re-referenced to the local average of each individual EcoG grid.

Next, data was bandpass filtered between 0.5 and 120Hz using an FIR Keyser filter

(attenuation level = 60 dB, Order = 3714). A second-order IIR notch filter was applied to

remove line noise (50Hz) and harmonics (100Hz). Then, electrodes reported as bad in the

original dataset and additional noisy channels visually identified on the basis of their power

spectrum properties via Welch's PSD analysis were rejected from further analysis (mean =

3.5, SD = 4.6).

To inspect EcoG grid location, electrodes were plotted against each subject's

cortical surfaces, which are available in the original dataset. After replicating individual

electrode locations for each participant, we then projected all electrodes in a the MNI-

ICBM152 cortical template (figure s1). In a few cases, this procedure resulted in spatial

distortion of some electrodes due to conversion between coordinate systems. In such cases,

electrode location was corrected by visually comparing the electrode's position in the

subject's anatomy and relocating it if necessary to the corresponding anatomical area in the

common cortical space. For more details on structural MRI preprocessing and post-surgical

grid coregistration procedures in the original data, see Berezutskaya et al., 2022.

Electrophysiological signals were then bandpass-filtered and epoched using

Fieldtrip (53). A third order Butterworth IIR filter was used to obtain continuous brain

signals within the Slow Frequency Band (SFB, 1-8Hz) and High-Frequency Band (HFB,

70-120Hz). For the latter frequency band, the envelope was obtained by extracting the

absolute value of the signal's Hilbert transform. Finally, data was epoched into twelve 30-

second segments, corresponding to the six music and six speech clips.

Cross-correlation analysis

Cross-correlation analyses were conducted using MATLAB Version 9.6.0 (2019a)

and 9.14.0 (2023a). The cross-correlation function was estimated for each electrode

between the z-normalized brain signals and the z-normalized cochlear envelopes of the

music and speech segments. This was done per subject, for the two frequency bands of

interest (SFB and HFB). Previous studies have used small temporal windows of

approximately 500ms, which excludes the slowest temporal modulations in the acoustic

signals. Here, we used a maximum lag of $\pm$ 400 samples, corresponding to a four-second

window between -2000 and 2000ms. This is in line with findings that larger temporal

windows are associated with better model performance in decoding analyses of brain

signals using Artificial Neural Networks (54). In the current work, no analyses were

conducted between the brain signals and information in the visual modality.

The maximum correlation coefficient and its corresponding temporal lag were

extracted from the correlation function. The mean coefficient estimates and corresponding

temporal lags were obtained by averaging across the values for the six segments within

each condition at the subject level (figure 1b, for data distributions see supplementary

figure s2). Statistical significance was estimated for each individual by obtaining a null

distribution of cross-correlation coefficients via permutation (figure 1b). For each subject,

the cross-correlation function was estimated 1,000 times between the z-normalized brain

signals and the z-normalized cochlear envelopes of six randomly generated white-noise

signals per condition. P-values for the observed data were subsequently obtained by

estimating the probability of observing the same or more extreme cross-correlation

coefficients given the empirically-obtained null distributions. The threshold for statistical

significance was set at an alpha value of 0.001.

Density-based cluster analysis

All statistically significant electrodes across our 30 subjects were then plotted in a

normalized cortical space (MNI-ICBM152). Electrodes that significantly correlated with

the acoustic envelope were identified performing multiple comparisons (i.e., 1858

electrodes were studied). Accordingly, a high number of false positives are expected, even

if a more conservative alpha value was chosen a priori. Because the permutation procedure

is agnostic of the anatomical location of electrodes, and in order to identify cortical regions

where the group-level effect was localized, we additionally conducted a density-based

cluster classification analysis on the collapsed data across all subjects using the DBSCAN

algorithm (55) as implemented by MATLAB. The assumption was that the overall effect of

cortical tracking should be more pronounced in task-relevant cortical regions. DBSCAN

should therefore spatially constrain any statistical effect in a data-driven manner to

functionally relevant cortical areas, while taking care of isolated electrodes which would

otherwise be considered outliers.

DBSCAN requires only two initial parameters: the minimum number of data points

for a cluster to be identified and the minimum radius within which those data points should

be contained (i.e., the epsilon parameter). This means that no starting data point or number

of clusters need to be defined a priori, unlike other clustering methods. To derive the

optimal parameters for the DBSCAN algorithm, we conducted a procedure to obtain the

parameter combination that optimized clustering within each condition. We tested five

minimum points (12 to 20 in steps of 2) and a range of 21 epsilon parameters (0 to 0.04 in

steps of 0.002). A matrix of parameter optimization indices (*OI*) was obtained by the

following formula.

239
$$OI = \left(\frac{E}{\epsilon}\right) * (PC)$$

Where ϵ is the 1-by-21 vector containing the list of possible minimum radiuses, P is the 1-

by-5 vector of possible minimum data points, Where E is the 5-by-21 matrix containing the

percentage of electrodes obtained for each combination of initial parameters, and C is the 5-

by-21 matrix containing the number of clusters obtained by each combination of

parameters. Finally, we normalized *OI* so that values range between 0 and 1, with 0

indicating poor algorithm performance and 1 indicating optimal performance.

Supplementary figure s3 shows the normalized OI values as a function of different

parameter combinations for the SFB and HFB analyses. The ideal parameters are those

which maximize OI_{norm} with the lowest possible cluster radius and the highest possible

number of electrodes within such cluster. The final parameters used for DBSCAN

clustering analyses for each frequency band and stimulus type are provided in table 1.

251
252

	Frequency band	$\mathcal{E}$	Min points	OI_{norm}
Speech Music	HFB	0.014	12	0.94
	SFB	0.012	12	0.92
	HFB	0.006	12	1.00
	SFB	0.006	16	1.00

253
254
255
256   **Table 1.** Optimal parameters for density-based clustering analysis. The epsilon ($\mathcal{E}$)
257   parameter represents the radius of search, and the minimum points stands for the minimum
258   number of electrodes within the search area for cluster identification.
259

Electrode anatomical labeling

DBSCAN returns a numbered list of identified clusters under the specified

parameter combination. We therefore grouped all clusters resulting from DBSCAN

analyses as a single group and anatomically labeled each electrode using the Desikan-

Killiany anatomical atlas (56). We took this approach because clusters where the result of

an algorithmic operation performed in space that is agnostic of brain anatomy, which in

practice means that two small clusters could be identified within the same anatomical

region. Electrodes were therefore labeled according to the cortical area in which they were

located by plotting each electrode on the MNI-ICBM152 template against overlaid

parcellations from the Desikan-Killiany atlas. This resulted in nine cortical regions of

interest (table 2), which were used as categorical predictors in our linear mixed effect

modeling analyses.

Linear mixed effect modeling

Linear Mixed Effect (LME) modeling was conducted using the LME4 package (57)

in R (R Core Team, 2023) using R Studio version 2023.6.1 (R Studio Corte Team, 2023).

Data was modeled for the two response variables of interest: cross-correlation values and

temporal lags. For each LME analysis, the minimum number of datapoints within each

cortical region for inclusion in the model was set to nine. For music, electrodes within the

STG, MTG, PFC and Supramarginal Gyrus (SG) were included in the analysis. The IFG,

premotor cortex, precentral and postcentral gyri were excluded because not enough

electrodes were found within these regions across frequency bands. For speech, electrodes

within the STG, MTG, precentral and postcentral gyri, IFG, and SG were included in the

analysis. Not enough electrodes were found across frequency bands in the premotor or

prefrontal cortices, so these regions were excluded. We additionally performed a joint LME

analysis for music and speech including the STG, MTG and SG as cortical regions of

interest.

Region	Music		Speech	
	SFB	*HFB*	*SFB*	*HFB*
STG	37	16	159	123
MTG	38	33	198	43
PFC	15	28	36	2
IFG	4	4	118	53
IPL	0	12	0	0
Premotor	0	5	24	0
Precentral	5	1	137	46
Postcentral	15	4	95	23
SG	9	28	90	30

Table 2. Number of statistically significant electrodes per anatomical region. *STG*,
(Superior Temporal Gyrus), *MTG* (Middle Temporal Gyrus), PFC (Prefrontal Cortex) *IFG*
(Inferior Frontal Gyrus), *IPL* (Inferior Parietal Lobe), SG (Supramarginal Gyrus)

A forward stepwise model selection procedure using Maximum Likelihood

Estimation was conducted to identify the models that best fit our data (table 3 and

supplementary tables s3, s4 and s5). Single predictor models for frequency band and

cortical region including a random intercept for subject were tested against a null (random

intercept-only) model. If any single-predictor model was significantly better than the null

model, the effect of adding additional terms and interactions while keeping the random

intercept unchanged was tested. Once the optimal combination of fixed effect terms was

determined, the effect of adding other random effects was tested. This procedure was

conducted for music and speech separately. However, we additionally conducted a joint

LME analysis where condition (music and speech) was considered an additional fixed

effect. Table 3 shows the best models predicting cross-correlation coefficients and temporal

lags per condition with their corresponding Akaike Information Criterion (AIC) and Bayes

Information Criterion (BIC). For music, no model outperformed a null model in predicting

temporal lags. For more information on model selection procedures, see supplementary

materials.

	LME model	AIC	BIC
Music	r ~ frequency + (1|subject)	-1299,136	-1285,854
	lag ~ (1+subject)	277.78	287.73
Speech	r ~ frequency*region + (1|subject) + (1|subject:region)	-6351,467	-6276,218
	lag ~ frequency*region + (1|subject) + (1|subject:region)	13426,74	13501,99
Both	r ~ frequency*region*condition + (1|subject) + (1|subject:region)	-4379.39	-4309.04
	lag ~ frequency*region*condition + 1(subject)	9624.23	9689.89

305   **Table 3**. Best fit models for each condition and for the collapsed model and for each
306   response variable of interest. For music, no model was significantly better in predicting
307   temporal lags than a null model.

Results

We first studied the properties of our acoustic stimuli. Top panels in figure 1a show the soundwave of two representative music and speech acoustic segments (see also figure s4 in supplementary materials for all other segments). A cochlear filter was applied to obtain the approximate spectrotemporal representation of each music and speech segment in the human cochlea (figure 1a, middle panels, see methods). Next, the cochlear envelope of each segment was estimated by averaging across all the cochlear frequencies (figure 1a, bottom panels). Figure 1b shows the normalized spectral power for the six music (left) and speech (right) segments correspondingly, obtained via Welch´s Power Spectrum Density (PSD) method. PSD analyses show high variability in spectral peaks across the different music and speech segments, consistent with the uncontrolled nature of the stimuli. The mean spectrum across music cochlear envelopes, however, shows peaks at around 0.2, 2 and 5 Hz, suggesting that across segments, music signals are somewhat more rhythmic than speech signals. Analyses confirmed statistically significant weak-to-moderate positive correlations between the six acoustic envelopes of music ($\alpha = 0.05$, two-tailed, table 4, top panel). For speech, no clear dominant peak was observed when the cochlear envelopes were averaged, reflecting marked differences in the temporal structure across the six different speech segments. This is consistent with the nature of the speech stimuli, which included both monologues and conversations between multiple individuals. Correlation analysis showed weak negative correlations for the acoustic envelopes of speech (table 4, bottom panel).

2	0.397***				
3	0.232***	0.103***			
4	0.442***	0.560***	0.299***		
5	0.290***	0.253***	0.012_n.s.	0.225***	
6	0.029***	-0.131***	0.147***	-0.037***	-0.006_n.s.

1						
2	0.036_n.s.					*Speech*
3	0.030***	0.006_n.s.				
4	-0.022_n.s.	0.030***	-0.050***			
5	0.049***	-0.102***	-0.042***	-0.098***		
6	0.043***	0.164***	-0.059***	0.002_n.s.	-0.006 _n.s.	

*** p < 0.001, ** p < 0.01, * p < 0.05

Table 4. Correlation across the music and speech segments.

We then studied neural tracking of the music and speech envelopes. For this, we

estimated the cross-correlation function between the brain and acoustic signals. Figure 1c

shows a schematic representation of the cross-correlation analysis pipeline. For each

subject (n = 30), we obtained the cross-correlation function between the cochlear envelopes

of the music (n = 6) and speech (n = 6) segments, and the brain signal recorded at each

electrode (n = 1858, mean = 61.93, SD = 19.09) in the Slow Frequency Band (SFB, 1 to

8Hz bandpass filtered signal) and the High Frequency Band (HFB, envelope of the 70 to

120Hz bandpass filtered signal). For each cross-correlation, we then extracted the

maximum brain-to-stimulus coefficient and its corresponding temporal lag. We finally

averaged the coefficients across the six segments within each stimulus type. Statistical

significance was estimated using a null distribution of cross-correlation coefficients

obtained via permutation (n $_{permutations}$ = 1000, α = 0.001, figure 1c). Finally, we additionally

implemented a density-based clustering classification analysis using the DBSCAN

algorithm to remove spatial outliers and identify cortical regions where significant

electrodes were most densely grouped.

Figure 1. Analysis of acoustic signals. **a.** Sound waves of representative acoustic segments for music (top-left panel) and speech (top-right panel), their cochlear spectrograms (middle panels) and cochlear envelope (bottom panels). **b.** PSD for all music (left) and speech (right) segments. For normalization, power at each frequency bin was divided the sum of power values across all frequencies. The red line shows average power across segments. **c.** Schematic representation of the cross-correlation analysis. For brain signals (SFB, 1-8Hz, top left and envelope of HFB, 70-120Hz, top right), cross-correlation was estimated against the cochlear envelope of stimuli signals (middle left) to obtain the maximum correlation coefficient and its corresponding lag (bottom left). A permutation procedure was conducted by estimating the cross-correlation function between SFB and HFB brain signals and the cochlear envelope of simulated white-noise (middle right), to obtain a null distribution of maximum correlation coefficients which was used to estimate significance thresholds. Solid black arrows represent the pipeline for real data. Dotted black lines represent the permutation procedure.

In both frequency bands, permutation and density-based clustering analyses showed a higher number of statistically significant electrodes tracking speech compared to music (figure 2). Electrodes tracking the music envelope in the SFB concentrated in perisylvian, temporal and prefrontal regions, including the middle and posterior portions of the STG, anterior, middle, and posterior portions of the MTG and ITG, the lowermost portions of the

precentral and postcentral gyri, and the most anterior portion of the dorsolateral Prefrontal

Cortex (dlPFC, figure 2a, left). For speech, cortical tracking in the SFB was abundantly

found in perisylvian, ventral prefrontal, pre and postcentral, and temporal regions,

including the anterior, middle, and posterior portions of both the STG and MTG, the

lowermost portions of the SG, middle and lower portions of the precentral and postcentral

gyri, premotor areas, the IFG, and the dlPFC (figure 2a, middle). In the SFB, the number of

statistically significant electrodes tracking the speech signals (n = 823, percent from total =

44.22%) was 6.5 times higher than the number of electrodes tracking the music signals (n =

125, percent from total = 6.72%, figure 2b). Interestingly, within the SFB, 84% of the

electrodes tracking the music signals (n = 105, percent from total = 5.65%) also tracked the

envelope of the speech signals (figure 2a, right and figure 2b).

Electrodes tracking music stimuli in the HFB concentrated around PFC and

premotor areas, as well as in SG, IPL, STG and MTG regions (figure 2c, left). In turn,

electrodes tracking speech in the HFB range were more densely concentrated in perisylvian

regions, including the middle and posterior STG, lowermost portions of the precentral and

postcentral gyri, and the middle and posterior portions of the IFG (figure 2c, middle). There

was an approximate two-fold increase in the number of electrodes showing statistically

significant tracking of speech in the HFB (n = 320, percent from total = 17.22%) compared

to music (n = 136, percent from total = 7.32%, figure 2d). Only 18.32% of electrodes

tracking the music envelope also tracked the speech envelope in the HFB. These electrodes

were restricted to posterior portions of the STG (n = 25, percent from total = 1.34%, figure

2c, right). For a non-thresholded figure showing the raw distribution of cross-correlation

coefficients and temporal lags prior to statistical analyses, see supplementary figure s5.

Figure 2. Statistically significant electrodes and their anatomical localization in the MNI-ICBM152 cortical template. **a.** and **c.** Spatial distribution of significant electrodes in the SFB (**a**) and HFB (**c**) after density-based clustering analyses for music (orange) and speech (purple). Electrodes that show mixed-selectivity (i.e., respond to both stimuli) are presented in green. **b.** and **d.** Number of electrodes that survive permutation statistics and density-based clustering classification in the SFB (**b**) and HFB (**d**).

Next, we investigated the organization of cross-correlation values and temporal lags (figure 3). For tracking of music signals in the SFB, cross-correlation coefficients (mean = 0.09, SD = 0.006) showed a gradient of increasing values from association areas, including anterior portions of the ITG and MTG, towards primary auditory areas and posterior portions of the STG (figure 3a, top-left). Temporal lags were maximally concentrated around time zero (mean = 12ms, median = 50ms, SD = 470ms, figure 3a, top-right), following a similar gradient with more positive lags found in anterior and middle portions of the STG (figure 3c, left). In the HFB, coefficients (mean = 0. 096, SD = 0.013) showed a more spatially-distributed gradient, with increasing values towards association areas, including superior parietal, middle temporal, and prefrontal regions (figure 3a, bottom-left). Similar to the SFB, temporal lags for the HFB concentrated around time zero (mean = -

14ms, median = 57ms, SD = 480ms, figure 3a, bottom-right), with maximum lags in middle

and posterior portions of the STG (figure 3c, right).

The cortical tracking effect for the speech signal in the SFB (mean = 0.10, SD =

0.001) was particularly distributed in cortical space, with higher coefficients found along

the anterior-to-posterior axis of the STG and MTG, but also in portions of the IFG,

precentral gyrus and SG (figure 3b, top-left). Similar to music, temporal lags for speech in

the SFB band converged around zero (mean = -6ms, median = -29ms, SD = 470ms, figure

3b, top-right, figure 3d, left). In turn, cortical tracking of the speech signal in the HFB

(mean = 0.10, SD = 0.023) showed a densely-grouped distribution and sharp increase in

cross-correlation coefficient values from prefrontal to posterior temporal regions, with a

concentration of the highest cross-correlation coefficients in the middle and posterior STG

(figure 3b, bottom-left). Interestingly, unlike all other conditions, mean temporal lags in the

HFB band were biased towards positive values (mean = 440ms, median = 490ms, SD =

590ms, figure 3b, bottom-right). Temporal lags in the HFB followed a similar gradient as

cross-correlation coefficients, with more positive lags (i.e. higher delays) concentrating

around the middle portion of the STG (figure 3d, right). Finally, to investigate the

consistency of our results across stimuli type, we re-analyzed our data using the same

analytical pipeline, while randomly excluding two segments (corresponding to 33% of the

data) at each of ten iterations. Results of these analyses show that both the magnitude of the

effect as well as its spatial organization remain consistent within each stimulus type across

all iterations (figure s6, supplementary materials).

Figure 3. Cortical tracking of music and speech envelopes in the MNI-ICBM152 cortical template. **a.** and **b.** Spatial distribution of mean correlation values for music (a) and speech (b) in the SFB and HFB. **c.** and **d.** Spatial distribution and histograms (left) of temporal lags (right) in the SFB and HFB. Positive lags indicate that the acoustic signal precedes brain signals, whereas negative lags indicate that the brain signals precede the acoustic signal. All electrodes shown are statistically significant (p < 0.001, uncorrected) and survive clustering analyses.

To quantify the effect of cortical tracking more accurately across anatomical regions, electrodes were next labeled according to their anatomical location (see methods). Electrodes were classified within one of nine left-hemisphere cortical areas (figure 4a and table 2). Cortical tracking of music was more prominent within the MTG and STG regions in the SFB, and within the MTG, SG and PFC in the HFB (figure 4b). For speech stimuli, cortical tracking was higher within MTG, STG and postcentral regions for the SFB, and within STG and IFG for the HFB (figure 4c). Next, we resorted to mixed effect modeling to

investigate how anatomical location and frequency band influence cortical tracking. The

two response variables of interest were cross-correlation coefficients and temporal lags.

The fixed effects of interest were frequency band and cortical region. The random effects of

interest were subject (to account for the within subject nature of the task) and the

interaction between subject and cortical region (to account for the different electrode grid

locations across subjects). We performed a forward stepwise model selection procedure to

select the best model explaining the data (see supplementary materials).

For music, a model including frequency band as fixed effect and subject as random

effect best fit the observed cross-correlation coefficients (AIC = -1299.13, BIC = -1285.85,

$R^2_{Conditional}$ = 0.12, $R^2_{Marginal}$ = 0.04, table s3). This model showed a statistically significant

main effect of frequency (F (1, 184.87) = 9.00, p = 0.003, figure 4d). Estimated Marginal

Means (emmeans) suggested that cortical tracking of the acoustic signals was significantly

higher in the HFB (emmean = 0.094, C.I. = [0.092, 0.097]) than in the SFB (emmean =

0.089, C.I. = [0.086, 0.092]). However, no model was significantly better than a null model

in predicting temporal lags during cortical tracking of music.

For speech, a model considering the interaction between frequency band and

cortical region as fixed effect, and both subject and the subject-by-region interaction as

random effects best explained the cross-correlation coefficients (AIC = -6351.47, BIC = -

6276.22, $R^2_{Conditional}$ = 0.19, $R^2_{Marginal}$ = 0.08, table s4). This model showed a significant

main effect of frequency (F (1, 1066.49) = 25.73, p = 4.64e-07) and cortical region (F (5,

162.87) = 8.81, p = 2.01e-07), and a significant interaction between frequency and cortical

region (F (5, 1058.98) = 7.47, p = 6.71e-07). Post-hoc pairwise comparisons showed that

cortical tracking of the speech envelope was significantly higher in the SFB compared to

the HFB within the IFG (emmean $_{SFB}$ = 0.105, CI = [0.102, 0.108]; emmean $_{HFB}$ = 0.095, CI

=[0.091, 0.099], p < 0.0001), MTG (emmean $_{SFB}$ = 0.10, CI = [0.101, 0.107]; emmean $_{HFB}$

= 0.99, CI = [0.091, 0.010], p = 0.022) and precentral gyrus (emmean $_{SFB}$ = 0.104, CI =

[0.101, 0108]; emmean $_{HFB}$ = 0.0942, CI = [0.0897, 0.099], p < 0.0001, figure 4e), and in

the HFB compared to the SFB within the STG (emmean $_{SFB}$ = 0.107, CI = [0.105, 0.110];

emmean $_{HFB}$ = 0.102, CI = [0.108, 0.114], p = 0.016, figure 4e). In the HFB, cortical

tracking was higher within STG (emmean = 0.109, CI = [0.107, 0.112]) compared to the

IFG (emmean = 0.100, CI = [0.097, 0.103], p < 0.0001), MTG (emmean = 0.102, CI =

[0.099, 0.105], p = 0.0002), precentral gyrus (emmean = 0.993, CI = [0.096, 0.102], p <

0.0001), postcentral gyrus (emmean = 0.993, CI = [0.095, 0.103], p = 0.0002) and SG

(emmean = 0.104, CI = [0.100, 0.108], p = 0.040, figure 4e). No statistically significant

differences were observed in cross-correlation coefficients across the different anatomical

regions for the SFB.

Figure 4. Anatomical regions of interest and significant effects as per mixed effect modeling analyses. **a.** Anatomical location of statistically significant electrodes after density-based clustering analyses. **b.** and **c.** Number of electrodes per anatomical parcellation for music (**a**) and speech (**c**). **d.** Main effect of frequency band for mean cross-correlation values during cortical tracking of music. **e.** Main and interaction effects for cross-correlation values during cortical tracking of speech. **f.** Main and interaction effects for temporal lags during cortical tracking of speech. Whiskers represent the Standard Error of the Mean (SEM). For all panels, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

For temporal lags during tracking of speech (figure 4f), the best model also included

frequency, region and their interaction as fixed effects, and subject as well as the subject-

by-region interaction as random effects (AIC = 1611.49, BIC = 1686.74, $R^2_{Conditional} = 0.25$,

$R^2_{Marginal} = 0.17$, table s4). This model showed significant main effects of frequency (F (1,

1066.70) = 117.27, p < 2.20e-16), cortical region (F (5, 150.79) = 3.54, p = 0.0047) and the

interaction between frequency and region (F (5, 1060.53) = 7.11, p = 1.48e-06). Post-hoc

pairwise comparisons showed more positive lags for HFB compared to SFB within the IFG

(emmean $_{HFB}$ = 114ms, CI = [-33ms, 262ms]; emmean $_{SFB}$ = -42ms, CI = [-152ms, 67ms], p

= 0.050), precentral gyrus (emmean $_{HFB}$ = 260ms, CI = [99ms, 420ms]; emmean $_{SFB}$ =

18ms, CI = [9ms, 128ms], p = 0.004), MTG (emmean $_{HFB}$ = 380ms, CI = [220ms, 541ms];

emmean $_{SFB}$ = -23ms, CI = [-118ms, 72ms], p < 0.0001), postcentral gyrus (emmean $_{HFB}$ =

431ms, CI = [219ms, 643ms]; emmean $_{SFB}$ = 2ms, CI = [-122ms, 126ms], p = 0.0002), SG

(emmean $_{HFB}$ = 455ms, CI = [266ms, 643ms]; emmean $_{SFB}$ = 32ms, CI [-9ms, 154ms], p <

0.0001) and STG (emmean $_{HFB}$ = 618ms, CI = [510ms, 725ms]; emmean $_{SFB}$ = -59ms, CI =

[-158ms, 40ms], p < 0.0001), suggesting that delays between the acoustic envelope of

speech and brain signals are higher in the HFB across all cortical regions of interest. In the

HFB, tracking within the STG (emmean = 618ms) occurred 503ms later compared to the

IFG (emmean = 114ms, p < 0.0001) and 358ms later compared to the postcentral gyrus

(emmean = 260ms, CI = [100ms, 420ms], p = 0.002). No differences in temporal lags

across cortical regions were observed within the SFB range (figure 4f).

Finally, we investigated the overall effect of cortical tracking and temporal lags by

designing a joint model for music and speech. For this, we used a subsample of electrodes

corresponding to the cortical regions where a significant effect of cortical tracking was

observed for both music and speech. This included the STG, MTG and the SG (figure 5a).

For mean cross-correlation values, the model that best fit the data was one including the

interaction between condition, frequency band, and cortical region as fixed effects, and

subject and the interaction between subject and cortical region as random effects (AIC = -

4502.26, BIC = -4431.91, $R^2_{Conditional}$ = 0.26, $R^2_{Marginal}$ = 0.17, table s5). Results of this

model suggest a main effect of condition (F (1, 799.99) = 76.49, p = 2.0e-16, figure 5b),

where cortical tracking of speech (emmean = 0.105, C.I.= [0.103, 0.107]) is significantly

higher than cortical tracking of music (emmean = 0.092, C.I. = [0.089, 0.095]). No effects

were found for frequency band or cortical region. For mean temporal lags, the model

included the interaction between condition, frequency band and region as fixed effects, and

a random intercept for subject (AIC = 1191.96, BIC = 1257.61, $R^2_{Conditional}$ = 0.22, R^2

$_{Marginal}$ = 0.20, table s5). Results of this model (figure 5c) show statistically significant main

effects of frequency band (F (1, 787.56) = 14.16, p = 0.0002), condition (F (1, 776.45) =

13.27, p = 0.0003), and the interaction between frequency and condition (F (1, 749.70) =

35.98, p = 3.096e-09). Post-hoc pairwise comparisons showed that cortical tracking of

speech in the HFB (emmean = 484ms, CI = [394ms, 574ms]) occurs, on average, 510ms

later than tracking of speech in the SFB (emmean = -25ms, CI = [-85ms, 34ms], p <

0.0001), 504ms later than tracking of music in the HFB (emmean = -20ms, CI = [-147ms,

107ms], p < 0.0001), and 385ms later than tracking of music in the SFB (emmean = 99ms,

CI = [-39ms, 237ms], p < 0.0001). No differences in temporal lags were observed for

cortical tracking of music across frequency bands.

Figure 5. Number of electrodes per anatomical location and statistical effects of joint
mixed effect model. Purple bars represent an effect for music whereas orange lines
represent an effect for speech. **a.** Number of electrodes in the three anatomical location
where statistically significant electrodes were found for both conditions. **b.** Main effect of
condition for cross-correlation coefficients. **c.** Main effect and interaction for temporal lags.
Whiskers represent the SEM. For all panels, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Discussion

Cortical tracking of music and speech has been previously observed in the range of slow (1-
8Hz, SFB) and fast (>70Hz, HFB) oscillatory activity under carefully controlled conditions
(13–16,18–20,22,28,29) and under naturalistic perceptual scenarios (17,21,38,39,41,42,58–
63,23,25,26,30,31,33–35). Here, we asked whether complex interactions between
anatomical region, frequency band and stimulus type exist during perception of naturalistic
music and speech stimuli that might have not been previously observed due to sample size,
methodological or analytical limitations. For this, we leverage the high spatiotemporal
resolution of electrocorticography by reanalyzing a publicly available EcoG dataset
collected during passive perception of naturalistic music and speech stimuli. Our findings
highlight both shared and distinct neural patterns between music and speech processing
across different cortical regions, with marked differences in the spectral and temporal
dynamics of brain responses.

Acoustic Features of Music and Speech

The spectral power analysis of the cochlear envelopes revealed substantial variability
across the six acoustic segments of music and speech. This variability was expected given
the naturalistic nature of the stimuli. Despite this variability, the spectral power
distributions of the cochlear envelopes suggest that music tends to be more rhythmic, as

evidenced by the dominant peaks at approximately 0.2, 2, and 5 Hz in the music envelopes.

In contrast, the speech segments did not exhibit a prominent peak, indicating more complex

and variable temporal patterns, in line with the diverse types of speech stimuli employed in

the study. Correlation analyses showed weak-to-moderate positive correlations within

music envelopes, suggesting some degree of temporal coherence across music segments.

This is further corroborated by the results showing weak negative correlations within

speech envelopes, which reflects the highly variable temporal structures inherent to speech,

including shifts in rhythm due to conversational interactions or changes in speech style.

Cortical organization of the tracking effect in the slow and fast frequency bands

In line with similar intracortical studies (22,23,35,39), we found distributed cortical

tracking in left hemisphere regions classically involved in language and music perception.

Density-based clustering analyses showed overlapping tracking of music and speech in the

SFB and HFB within the MTG, STG, SG, precentral and postcentral gyri. These findings

are in line with previous research using source-localized MEG signals (20,21,62,64) and

intracortical data (16,31–33,35,39). Overlap of cortical tracking around perisylvian and

middle and superior temporal areas for both stimuli suggests that shared neuroanatomical

mechanisms are involved in the processing of the acoustic envelopes of music and speech

(2,11,12). This was particularly true for the SFB, where 84% of electrodes that showed a

statistically significant effect for music also showed a significant effect for speech, thus

suggesting not only high anatomical overlap, but also sensitivity to both types of stimuli

within the range of slow oscillations.

However, we also found differences that were both frequency- and stimulus-

dependent. Tracking of music in both the SFB and HFB, and of speech in the SFB,

extended beyond classic auditory regions to include prefrontal, middle temporal, and

parietal areas. In contrast, tracking of speech in the HFB was densely localized to

perisylvian and ventral prefrontal regions, indicating a more focused cortical engagement

for speech processing in the range of high gamma band activity. One previous EcoG study

found similar a spatial organization pattern of neural activations during speech perception,

with SFB activity showing a global distribution in cortical space, while HFB activity was

primarily localized to the STG (23). This frequency-dependent pattern of global versus

local spatial organization aligns with the idea that slow oscillatory activity facilitates

communication among globally distributed neuronal ensembles, whereas high frequency

activity reflects local activation of small neural populations in response to specific sensory

features (65–73). Notably, the importance of cross-frequency interactions between the SFB

and gamma band oscillations for speech perception and language processing has also been

well documented (74–77). These interactions likely support multiscale extraction of both

acoustic features and hierarchical structures that are essential for language comprehension.

However, such cross-frequency coupling has been primarily observed between the slow

oscillatory range (0.5–8 Hz) and the slower component of gamma-band activity (25–40

Hz), rather than the faster component (>70 Hz).

Functional anatomical selectivity during tracking of natural music and speech signals

In the SFB, electrodes tracking the envelope of speech were observed within the IFG,

which were nevertheless absent during tracking of music stimuli. In contrast, in the HFB,

electrodes tracking the envelope of music stimuli were observed within the dlPFC and IPL

regions, which were absent during tracking of speech. Both of these findings are in line

with the fMRI literature on the role of the dlPFC and parietal cortex during music

perception, and of the IFG during speech perception (43–48). In the EcoG literature,

selective dlPFC recruitment in response to music has been previously reported during

perception of musical stimuli, which has been attributed to attentional and emotional

processing, as well as recall of complex melodic features in music stimuli (32,33,78). The

involvement of the IFG during tracking of speech stimuli in both SFB and HFB has also

been extensively reported in previous studies using intracortical methods (23,54,78). This

suggests that in spite of partial anatomical overlap in perisylvian regions, cortical tracking

of music and speech signals can also be intracortically mapped to distinct parietal and

prefrontal brain regions in a stimulus-dependent manner.

Multiple roles have been previously attributed to the IFG during speech perception,

including syntactic processing, semantic processing and phonological working memory

(45,79–81). Admittedly, activation of the IFG has also been reported during music

perception, however, recent metanalyses and clinical studies suggest this effect is right

lateralized (82,83). Finally, increased BOLD activation of IPL regions has been observed in

response to music compared to speech (84), and increased musical training is associated

with increased IPL activity among musicians (85).

Not many EcoG studies have directly investigated the extent to which tracking of

music and speech can be mapped to distinct cortical areas and, more importantly, how

interactions between cortical regions, frequency bands and stimulus type influence this

mapping. A recent study reported no substantial differences in anatomical regional

selectivity in how brain signals track music and speech across delta (1-4Hz), theta (5-8Hz),

alpha (8-12), beta (18-30Hz), low gamma (30-50Hz) and high-gamma (80-150Hz)

frequency bands (39). In contrast, our findings suggest functional anatomical differences,

particularly within the high-frequency gamma band (HFB), with patterns resembling those

observed in fMRI research. One possible explanation for this discrepancy in results could

be the use of density-based clustering analyses, which restricted the cortical tracking effect

to regions with a higher concentration of significant electrodes. This approach may have

revealed subtle differences in the cortical organization of the tracking effect that may have

been missed without accounting for the spatial distribution of statistically significant

electrodes. Alternatively, the divergence in results may stem from differences in stimulus

design. The previous study used unimodal auditory stimuli, whereas our study employed

audiovisual stimuli. In our study, the inclusion of multimodal stimuli may have engaged

association cortices or introduced attentional competition between modalities, potentially

leading to differences in how music and speech signals were integrated and processed.

Increased cortical tracking of speech across frequency bands

The net effect of cortical tracking was higher for speech than for music. This was

reflected in a higher proportion of electrodes tracking the speech envelope in both

frequency bands, as well as in the results of our joint linear mixed effect models, which

show a main effect of stimulus type in predicting mean cross-correlation coefficient values.

Such findings are aligned with the recent literature using naturalistic stimuli, which have

also shown that brain signals recover the envelope of speech signals more faithfully than

that of music signals (17,35,39). Importantly, Zuk and colleagues have previously

demonstrated that increased tracking of speech compared to music cannot be attributed to

spectral differences in the envelope of both signals (17). Moreover, power spectra of our

stimuli show that, across the different acoustic segments, speech is less rhythmic than

music, showing no clear dominating peak. This suggests that increased tracking of speech

cannot be attributed to higher regularity in the amplitude modulations of the speech
envelope compared to the music envelope.

While our methods, as well as the lack of data about behavioral performance,
prevent us from investigating the role of higher order cognitive processes, we cannot rule
out the possibility that the observed effect might have a cognitive explanation. Both speech
intelligibility and musical training have been shown to increase cortical tracking of the
corresponding signals (13,16,22,28,86), which implies that previous experience processing
a particular stimulus can modulates the magnitude of the cortical tracking effect.
Additionally, attentional allocation has been associated with increased cortical tracking in
previous studies (29,30,87,88).

Cortical gradient in lag values during HFB tracking of speech signals

Our analyses also revealed interesting temporal dynamics during cortical tracking of speech
stimuli in the range of HFB activity. For music and speech in the SFB, mean and median
lag values were close to zero across all anatomical regions of interest, thus suggesting that
neural activity in the slow oscillatory range tracks fast temporal modulations in the
incoming acoustic signal in a relatively time-locked manner. Similarly, median temporal
lags during HFB tracking followed the music signal by approximately 60ms on average.
This could additionally reflect fast neural responses to specific acoustic features present in
the music signal. In line with this, previous EcoG research has demonstrated that HFB
power within prefrontal, pre/postcentral, and temporal regions encodes features such as
loudness, harmonization, and the presence or absence of lyrics (78). In the same study,
onset of lyrics was characterized by additional increases in HFB power, localized to
posterior portions of the STG.

For speech, estimated marginal means show that HFB tracking always occurred at

later lags compared to the SFB. Our results showed an increasing gradient of lag values

from prefrontal, parietal and middle temporal to superior temporal regions, suggesting a

highly localized effect of late-latency HFB tracking of speech within middle and posterior

portions of the STG. The joint LME model further confirmed that HFB tracking of speech

occurred significantly later compared to tracking of speech in the SFB, and to tracking of

music in both the SFB and HFB in the three anatomical regions included in the analysis

(STG, MTG and SG). This HFB gradient, which is observed for speech but not for music,

could reflect tracking of amplitude modulations in the acoustic envelope at different

timescales, perhaps reflecting the encoding of hierarchical structures at multiple levels. If

this was the case, a prediction would be that, among trained musicians, a similar gradient in

lag values across functionally relevant anatomical regions should be observed, reflecting

their ability to extract structural units embedded in music signals. However, further

research is needed to test this hypothesis.

Limitations

Our study has some important limitations. First, we lack any behavioral measure that

facilitates the interpretation of observed differences in cortical tracking. Without such data,

we cannot rule out potential confounding factors, such as differential attentional allocation

(29,30,88) due to differences in the semantic context of visual stimuli accompanying music

and speech segments. Additionally, the presentation of a concurrent visual stream during

passive auditory perception of music and speech segments may have introduced another

confound. Previous studies have shown that rhythmic modulations in visual stimuli can

crossmodally activate auditory areas (63,89,90). As a result, the effects we observed may

have been influenced by the integration of visual stimuli with the auditory stimuli,

potentially differing in how music and speech were processed.

Another limitation of our study is the use of a relatively broad temporal window

(4000ms) for cross-correlation analyses. While this approach may have reduced statistical

power and increased the likelihood of spurious results, it was motivated by two

considerations. First, previous research has focused primarily on shorter temporal

modulations, such as beat, notes, syllables, and words, which are often analyzed using

smaller windows (around 500ms). This approach overlooks slower temporal structures such

as musical and speech phrases, which occur at much slower rates. Second, a prior study

using the same dataset demonstrated that longer segments (~6 seconds) yielded better

performance in predicting brain signals using Artificial Neural Networks (ANN), with

improved generalization to new stimuli (54). Therefore, we chose a broader window in

hopes of revealing temporal dynamics that may have been overlooked in previous studies.

Our study was also limited by the localization of the ECoG grids, which were

predominantly placed in the left hemisphere. This constraint prevents us from exploring

potential differences in the cortical tracking of music and speech in right hemisphere

regions, which have previously been implicated in music perception and could partially

explain our results. Additionally, a technical limitation involves the re-referencing of the

ECoG electrodes to the average of each grid. While this is a common practice in ECoG

studies, it is susceptible to field spread, meaning that activity observed in central regions

could be influenced by activity from auditory areas.

We derive our interpretations from marginal mean estimates obtained from LME

models. While this allows us to draw group-level conclusions, it does not capture the

considerable intra-individual variability in lag estimates, which include both positive and

negative lags across regions and conditions. Negative lags might reflect both higher-order

and low-level predictive processing. Indeed, prior research suggests that morphosyntactic

information facilitates predictive processing during speech comprehension (22,91,92),

while neural mechanisms involved in rhythm perception contribute to temporal predictions

during processing of music stimuli (11,93–95). Unfortunately, our methods do not allow us

to determine the extent to which negative lag values reflect prediction, and any

interpretation we could offer would be speculative. However, they were included in our

analyses because there is enough theoretical ground to assume that they reflect relevant

neurobiological and cognitive processes that support the perception of music and speech.

Relatedly, the neurobiological processes behind cortical tracking of speech and

music signals remain a topic of ongoing debate. Two prominent perspectives suggest that

cortical tracking of acoustic envelopes either represents the entrainment of endogenous

oscillations or reflects a series of evoked responses to amplitude fluctuations within the

acoustic signal (25,40,96–101). While our methods are not suited to test either of these

hypotheses, our results provide evidence of differences in the temporal dynamics and

cortical organization of low-level processing of acoustic envelopes, regardless of the

underlying mechanism. Additionally, we do not directly address the specific information

being tracked within each temporal bin and anatomical region in this work. This is

important because different structural units, acoustic features, or information content can be

differentially tracked by the human brain. While this question has been addressed elsewhere

in the literature (38,39,49,78,86,102), we hope that the open-access nature of the data,

along with improved methods to disentangle the contribution of low-level features and

higher-order information to cortical tracking, will motivate others to address similar

questions using this dataset.

Finally, several limitations of this study stem from the adoption of a mathematically

simple approach to quantifying cortical tracking. This simplicity limits the method's ability

to disentangle the underlying neurobiological mechanisms or precisely identify the acoustic

features or information tracked by neural signals, which in turn limits the depth of

interpretation we can provide for our results. Nevertheless, the consistency of our findings

with those of prior studies employing more sophisticated methodologies underscores the

potential of our approach. Importantly, this simplicity may hold significant advantages for

translational applications, particularly in contexts where minimizing computational

demands is a priority or where theoretical assumptions are less relevant for practical

applications.

Conclusion

In summary, our results show widespread cortical tracking and reveal both functional

overlap and anatomical specialization in the spatial and temporal dynamics associated with

the tracking of naturalistic music and speech stimuli. Passive perception of music and

speech was associated with distributed tracking of acoustic signals in the SFB with near

zero delays in mostly overlapping temporal and perisylvian regions. While some overlap

was observed in the HFB for both stimulus types, marked anatomical and functional

selectivity emerged, with regions such as the dlPFC and IPL areas preferentially tracking

music, and the IFG preferentially tracking speech. Notably, the overall magnitude of the

tracking effect was higher for speech across both frequency bands. Additionally, a gradient

in lag values was observed during HFB tracking of speech, spanning from association areas

in prefrontal, middle temporal, and parietal regions towards STG areas, which was

nonetheless absent during HFB tracking of music.

Our findings further support previous research indicating that cortical tracking of

music and speech extends beyond controlled experimental settings to more complex

naturalistic signals, which are far less controlled, less rhythmic, and that are presented

under conditions of sensory multimodality. Moreover, our results highlight the brain's use

of both domain-general mechanisms and frequency-dependent anatomical specialization

when tracking natural acoustic signals. Finally, we highlight a complex interaction between

stimulus type, cortical region, and frequency band, which could underlie the brain's ability

to extract hierarchical structures from the envelope of natural acoustic signals. However,

future research is needed to empirically test this hypothesis.

Conflict of interest

The authors declare no conflict of interest.

Data availability statement

Data for this study is open and can be accessed at https://openneuro.org/datasets/ds003688.

Code used for data analysis can be made available upon request.

References

1.   Fujita H, Fujita K. Human language evolution: a view from theoretical linguistics on

how syntax and the lexicon first came into being. Primates [Internet].

2022;63(5):403–15. Available from: https://doi.org/10.1007/s10329-021-00891-0

2.   Asano R. The evolution of hierarchical structure building capacity for language and

music: a bottom-up perspective. Primates [Internet]. 2022;63(5):417–28. Available

793  from: https://doi.org/10.1007/s10329-021-00905-x

794 3. Miyagawa S, Berwick RC, Okanoya K. The emergence of hierarchical structure in
795  human language. Front Psychol. 2013;4(February):1–6.

796 4. McFee B, Nieto O, Farbood MM, Bello JP. Evaluating hierarchical structure in
797  music annotations. Front Psychol. 2017;8(AUG):1–17.

798 5. Lerdahl F, Jackendoff R. An overview of hierarchical structure in music. Music
799  Percept An Interdiscip J. 1983;1(2):229–52.

800 6. Schön D, Morillon B. Music and language. In: The Oxford handbook of music and
801  the brain. 2019. p. 391–416.

802 7. Aboitiz F. A brain for speech. Springer Nature; 2017. 1–469 p.

803 8. van Noorden L, Moelants D. Resonance in the perception of musical pulse. Int J
804  Phytoremediation. 1999;21(1):43–66.

805 9. Ding N, Patel AD, Chen L, Butler H, Luo C, Poeppel D. Temporal modulations in
806  speech and music. Neurosci Biobehav Rev [Internet]. 2017;81:181–7. Available
807  from: https://doi.org/10.1016/j.neubiorev.2017.02.011

808 10. Pellegrino F, Coupe C, Marisco E. A cross-language perspective on speech
809  information rate. Language (Baltim). 2011;87(3):539–58.

810 11. Tillmann B. Music and language perception: Expectations, structural integration, and
811  cognitive sequencing. Top Cogn Sci. 2012;4(4):568–84.

812 12. Fiveash A, Bedoin N, Gordon RL, Tillmann B. Processing Rhythm In Speech And
813  Music: Shared Mechanisms And Implications For Developmental Speech And

814   Language Disorders. Neuropsychology. 2021;35(8):771–91.

815   13.   Doelling KB, Poeppel D. Cortical entrainment to music and its modulation by

816   expertise. Proc Natl Acad Sci U S A. 2015;112(45):E6233–42.

817   14.   Nozaradan S, Peretz I, Mouraux A. Selective neuronal entrainment to the beat and

818   meter embedded in a musical rhythm. J Neurosci. 2012;32(49):17572–81.

819   15.   Nozaradan S. Exploring how musical rhythm entrains brain activity with

820   electroencephalogram frequency-tagging. Philos Trans R Soc B Biol Sci.

821   2014;369(1658).

822   16.   Harding EE, Sammler D, Henry MJ, Large EW, Kotz SA. Cortical tracking of

823   rhythm in music and speech. Neuroimage [Internet]. 2019;185(2019):96–101.

824   Available from: https://doi.org/10.1016/j.neuroimage.2018.10.037

825   17.   Zuk NJ, Murphy JW, Reilly RB, Lalor EC. Envelope reconstruction of speech and

826   music highlights stronger tracking of speech at low frequencies [Internet]. Vol. 17,

827   PLoS Computational Biology. 2021. 1–32 p. Available from:

828   http://dx.doi.org/10.1371/journal.pcbi.1009358

829   18.   Kumagai Y, Arvaneh M, Tanaka T. Familiarity affects entrainment of EEG in music

830   listening. Front Hum Neurosci. 2017;11(July):1–8.

831   19.   Wollman I, Arias P, Aucouturier JJ, Morillon B. Neural entrainment to music is

832   sensitive to melodic spectral complexity. J Neurophysiol. 2020;123(3):1063–71.

833   20.   Assaneo MF, Poeppel D. The coupling between auditory and motor cortices is rate-

834   restricted: Evidence for an intrinsic speech-motor rhythm. Sci Adv. 2018;4(2):1–10.

21.   Chalas N, Daube C, Kluger DS, Abbasi O, Nitsch R, Gross J. Speech onsets and

sustained speech contribute differentially to delta and theta speech tracking in

auditory cortex. Cereb Cortex. 2023;33(10):6273–81.

22. Ding N, Melloni L, Zhang H, Tian X, Poeppel D. Cortical tracking of hierarchical

linguistic structures in connected speech. Nat Neurosci. 2015;19(1):158–64.

23. Zion Golumbic EM, Ding N, Bickel S, Lakatos P, Schevon CA, McKhann GM, et al.

Mechanisms underlying selective neuronal tracking of attended speech at a "cocktail

party." Neuron [Internet]. 2013;77(5):980–91. Available from:

http://dx.doi.org/10.1016/j.neuron.2012.12.037

24. Gross J, Hoogenboom N, Thut G, Schyns P, Panzeri S, Belin P, et al. Speech

Rhythms and Multiplexed Oscillatory Sensory Coding in the Human Brain. PLoS

Biol. 2013;11(12).

25. Oganian Y, Kojima K, Breska A, Cai C, Findlay A, Chang E, et al. Phase Alignment

of Low-Frequency Neural Activity to the Amplitude Envelope of Speech Reflects

Evoked Responses to Acoustic Edges, Not Oscillatory Entrainment. J Neurosci.

2023;43(21):3909–21.

26. Giordano BL, Ince RAA, Gross J, Schyns PG, Panzeri S, Kayser C. Contributions of

local speech encoding and functional connectivity to audio-visual speech perception.

Elife. 2017;6:1–27.

27. Doelling KB, Arnal LH, Ghitza O, Poeppel D. Acoustic landmarks drive delta-theta

oscillations to enable speech comprehension by facilitating perceptual parsing.

Neuroimage [Internet]. 2014;85:761–8. Available from:

http://dx.doi.org/10.1016/j.neuroimage.2013.06.035

28.   Peelle JE, Gross J, Davis MH. Phase-locked responses to speech in human auditory
859   cortex are enhanced during comprehension. Cereb Cortex. 2013;23(6):1378–87.

860   29.   Vanthornhout J, Decruy L, Francart T. Effect of task and attention on neural tracking
861   of speech. Front Neurosci. 2019;13(September):1–11.

862   30.   Viswanathan V, Bharadwaj HM, Shinn-Cunningham BG. Electroencephalographic
863   signatures of the neural representation of speech during selective attention. eNeuro.
864   2019;6(5):1–14.

865   31.   Kubanek J, Brunner P, Gunduz A, Poeppel D, Schalk G. The Tracking of Speech
866   Envelope in the Human Cortex. PLoS One. 2013;8(1).

867   32.   Herff SA, Herff C, Milne AJ, Johnson GD, Shih JJ, Krusienski DJ. Prefrontal high
868   gamma in ecog tags periodicity of musical rhythms in perception and imagination.
869   eNeuro. 2020;7(4):1–11.

870   33.   Ding Y, Zhang Y, Zhou W, Ling Z, Huang J, Hong XB, et al. Neural Correlates
871   ofMusic Listening and Recall in the Human Brain. J Neurosci. 2019;39(41):8112–
872   23.

873   34.   Commuri V, Kulasingham JP, Simon JZ. Cortical responses time-locked to
874   continuous speech in the high-gamma band depend on selective attention. Front
875   Neurosci. 2023;17.

876   35.   Berezutskaya J, Freudenburg Z V., Güçlü U, van Gerven MAJ, Ramsey NF. Neural
877   tuning to low-level features of speech throughout the perisylvian cortex. J Neurosci.
878   2017;37(33):7906–20.

36.   Sankaran N, Leonard MK, Theunissen F, Chang EF. Encoding of melody in the

human auditory cortex. Sci Adv. 2024;10(7):1–16.

37.   Norman-Haignere S V., Feather J, Boebinger D, Brunner P, Ritaccio A, McDermott

JH, et al. A neural population selective for song in human auditory cortex. Curr Biol

[Internet]. 2022;32(7):1470-1484.e12. Available from:

https://doi.org/10.1016/j.cub.2022.01.069

38.   Omigie D, Lehongre K, Navarro V, Adam C, Samson S. Neuro-oscillatory tracking

of low- and high-level musico-acoustic features during naturalistic music listening:

Insights from an intracranial electroencephalography study. Psychomusicology

Music Mind, Brain. 2020;30(1):37–51.

39.   Te Rietmolen N, Mercier M, Trebuchon A, Morillon B, Schon D. Speech and music

recruit frequency-specific distributed and overlapping cortical networks. Elife

[Internet]. 2024;2022.10.08.511398. Available from:

https://www.biorxiv.org/content/10.1101/2022.10.08.511398v1%0Ahttps://www.bio

rxiv.org/content/10.1101/2022.10.08.511398v1.abstract

40.   Oganian Y, Chang EF. A speech envelope landmark for syllable encoding in human

superior temporal gyrus. Sci Adv. 2019;5(11):1–13.

41.   Kulasingham JP, Brodbeck C, Presacco A, Kuchinsky SE, Anderson S, Simon JZ.

High gamma cortical processing of continuous speech in younger and older listeners.

Neuroimage [Internet]. 2020;222(August):117291. Available from:

https://doi.org/10.1016/j.neuroimage.2020.117291

42.   Synigal SR, Teoh ES, Lalor EC. Including measures of high gamma power can

improve the decoding of natural speech From EEG. Front Hum Neurosci.

2020;14(April):1–12.

43.   Green DM, Swets JA. Signal Detection Theory and Psychophysics. New York: John
Wiley & Sons, Ltd; 1966.

44.   Ohnishi T, Matsuda H, Asada T, Aruga M, Hirakata M, Nishikawa M, et al.
Functional anatomy of musical perception in musicians. Cereb Cortex.
2001;11(8):754–60.

45.   Enge A, Friederici AD, Skeide MA. A meta-analysis of fMRI studies of language
comprehension in children. Neuroimage [Internet]. 2020;215(April):116858.
Available from: https://doi.org/10.1016/j.neuroimage.2020.116858

46.   Chan MMY, Han YMY. The Functional Brain Networks Activated by Music
Listening:A Neuroimaging Meta-Analysis and Implications for Treatment.
Neuropsychology. 2022;36(1):4–22.

47.   Tremblay P, Small SL. On the context-dependent nature of the contribution of the
ventral premotor cortex to speech perception. Neuroimage [Internet].
2011;57(4):1561–71. Available from:
http://dx.doi.org/10.1016/j.neuroimage.2011.05.067

48.   Lankinen K, Ahveninen J, Uluç I, Daneshzand M, Mareyam A, Kirsch JE, et al. Role
of articulatory motor networks in perceptual categorization of speech signals: a 7T
fMRI study. Cereb Cortex. 2023;33(24):11517–25.

49.   McCarty MJ, Murphy E, Scherschligt X, Woolnough O, Morse CW, Snyder K, et al.
Intraoperative cortical localization of music and language reveals signatures of
structural complexity in posterior temporal cortex. iScience [Internet].

2023;26(7):107223. Available from: https://doi.org/10.1016/j.isci.2023.107223

50. Berezutskaya J, Vansteensel MJ, Aarnoutse EJ, Freudenburg Z V., Piantoni G,
Branco MP, et al. Open multimodal iEEG-fMRI dataset from naturalistic stimulation
with a short audiovisual film. Sci Data. 2022;9(1):1–13.

51. Chi T, Ru P, Shamma SA. Multiresolution spectrotemporal analysis of complex
sounds. J Acoust Soc Am. 2005;118(887).

52. Tadel F, Baillet S, Mosher JC, Pantazis D, Leahy RM. Brainstorm: A user-friendly
application for MEG/EEG analysis. Comput Intell Neurosci. 2011;2011.

53. Oostenveld R, Fries P, Maris E, Schoffelen JM. FieldTrip: open source software for
advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput
Intell Neurosci. 2011;2011:156869.

54. Berezutskaya J, Freudenburg Z V., Güçlü U, van Gerven MAJ, Ramsey NF. Brain-
optimized extraction of complex sound features that drive continuous auditory
perception. Vol. 16, PLoS Computational Biology. 2020. 1–34 p.

55. Ester M, Kriegel H-P, Sander J, Xu X. A Density-Based Algorithm for Discovering
Clusters in Large Spatial Databases with Noise Martin. KDD-96 Proc. 1996;2:565–
80.

56. Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, et al. An
automated labeling system for subdividing the human cerebral cortex on MRI scans
into gyral based regions of interest. Neuroimage. 2006;31(3):968–80.

57. Bates D, Mächler M, Bolker BM, Walker SC. Fitting linear mixed-effects models
using lme4. J Stat Softw. 2015;

58. Kaneshiro B, Nguyen DT, Norcia AM, Dmochowski JP, Berger J. Natural music
evokes correlated EEG responses reflecting temporal structure and beat. Neuroimage
[Internet]. 2020;214(December 2019):116559. Available from:
https://doi.org/10.1016/j.neuroimage.2020.116559

59. Brodbeck C, Kandylaki KD, Scharenborg O. Neural representations of non-native
speech reflect proficiency and interference from native language knowledge. J
Neurosci. 2024;44(1).

60. Micheli C, Schepers IM, Ozker M, Yoshor D, Beauchamp MS, Rieger JW.
Electrocorticography reveals continuous auditory and visual speech tracking in
temporal and occipital cortex. Eur J Neurosci. 2020;51(5):1364–76.

61. Park H, Ince RAA, Schyns PG, Thut G, Gross J. Frontal Top-Down Signals Increase
Coupling of Auditory Low-Frequency Oscillations to Continuous Speech in Human
Listeners. Curr Biol [Internet]. 2015;25(12):1649–53. Available from:
http://dx.doi.org/10.1016/j.cub.2015.04.049

62. Park H, Thut G, Gross J. Predictive entrainment of natural speech through two
fronto-motor top-down channels. Lang Cogn Neurosci [Internet]. 2020;35(6):739–
51. Available from: https://doi.org/10.1080/23273798.2018.1506589

63. Park H, Ince RAA, Schyns PG, Thut G, Gross J. Representational interactions during
audiovisual speech entrainment: Redundancy in left posterior superior temporal
gyrus and synergy in left motor cortex. PLoS Biol. 2018;16(8):1–26.

64. Ahissar E, Nagarajan S, Ahissar M, Protopapas A, Mahncke H, Merzenich MM.
Speech comprehension is correlated with temporal response patterns recorded from

auditory cortex. Proc Natl Acad Sci U S A. 2001;98(23):13367–72.

65.   Fries P, Nikolić D, Singer W. The gamma cycle. Trends Neurosci. 2007;30(7):309–

16.

66.   Rieder MK, Rahm B, Williams JD, Kaiser J. Human gamma-band activity and

behavior. Int J Psychophysiol. 2011;79(1).

67.   Ray S, Niebur E, Hsiao SS, Sinai A, Crone NE. High-frequency gamma activity (80-

150 Hz) is increased in human cortex during selective attention. Clin Neurophysiol.

2008;119(1):116–33.

68.   Canolty RT, Edwards E, Dalal SS, Soltani M, Nagarajan SS, Kirsch HE, et al. High

gamma power is phase-locked to theta oscillations in human neocortex. Science (80-

). 2006;313(5793):1626–8.

69.   Fries P. Rhythms for Cognition: Communication through Coherence. Neuron

[Internet].   2015;88(1):220–35.   Available   from:

http://dx.doi.org/10.1016/j.neuron.2015.09.034

70.   Fujisawa S, Buzsáki G. A 4 Hz Oscillation Adaptively Synchronizes Prefrontal,

VTA, and Hippocampal Activities. Neuron. 2011;72(1):153–65.

71.   Buzśaki G, Wang XJ. Mechanisms of gamma oscillations. Annu Rev Neurosci.

2012;35:203–25.

72.   Sirota A, Buzsáki G. Interaction between neocortical and hippocampal networks via

slow oscillations. halamus Relat Syst. 2005;3(4):245–59.

73.   Buzsaki G. Rhythms of the brain. Oxford: Oxford University Press; 2006.

74.   Luo H, Poeppel D. Cortical oscillations in auditory perception and speech: Evidence

for two temporal windows in human auditory cortex. Front Psychol.

2012;3(MAY):1–10.

75.   Hyafil A, Fontolan L, Kabdebon C, Gutkin B, Giraud AL. Speech encoding by

coupled cortical theta and gamma oscillations. Elife. 2015;4(MAY):1–45.

76.   Lizarazu M, Lallier M, Molinaro N. Phase–amplitude coupling between theta and

gamma oscillations adapts to speech rate. Ann N Y Acad Sci. 2019;1453(1):140–52.

77.   Lizarazu M, Carreiras M, Molinaro N. Theta-gamma phase-amplitude coupling in

auditory cortex is modulated by language proficiency. Hum Brain Mapp.

2023;44(7):2862–72.

78.   Sturm I, Biankertz B, Potes C, Schaik G, Curio G. ECoG high gamma activity

reveals distinct cortical representations of lyrics passages, Harmonic and timbre-

related changes in a rock song. Front Hum Neurosci. 2014;8(OCT):1–14.

79.   Friederici AD. The brain basis of language processing: From structure to function.

Physiol Rev. 2011;91(4):1357–92.

80.   Bulut T. Domain-general and domain-specific functional networks of Broca's area

underlying language processing. Brain Behav. 2023;13(7):1–20.

81.   Perrachione TK, Ghosh SS, Ostrovskaya I, Gabrieli JDE, Kovelman I. Phonological

working memory for words and nonwords in cerebral cortex. J Speech, Lang Hear

Res. 2017;60(7):1959–79.

82.   Asano R, Lo V, Brown S. The Neural Basis of Tonal Processing in Music: An ALE

Meta-Analysis. Music Sci. 2022;5:1–15.

83.   Chen X, Affourtit J, Ryskin R, Regev TI, Norman-Haignere S, Jouravlev O, et al.

1012   The human language system, including its inferior frontal component in "Broca's

1013   area," does not support music perception. Cereb Cortex. 2023;33(12):7904–29.

1014   84.   Merrill J, Sammler D, Bangert M, Goldhahn D, Lohmann G, Turner R, et al.

1015   Perception of words and pitch patterns in song and speech. Front Psychol.

1016   2012;3(MAR):1–13.

1017   85.   Liu Y, Liu G, Wei D, Li Q, Yuan G, Wu S, et al. Effects of musical tempo on

1018   musicians' and non-musicians' emotional experience when listening to music. Front

1019   Psychol. 2018;9(NOV):1–11.

1020   86.   Di Liberto GM, Pelofi C, Shamma S, de Cheveigné A. Musical expertise enhances

1021   the cortical tracking of the acoustic envelope during naturalistic music listening.

1022   Acoust Sci Technol. 2020;41(1):361–4.

1023   87.   Ahmed F, Nidiffer AR, O'Sullivan AE, Zuk NJ, Lalor EC. The integration of

1024   continuous audio and visual speech in a cocktail-party environment depends on

1025   attention. Neuroimage [Internet]. 2023;274(August 2022):120143. Available from:

1026   https://doi.org/10.1016/j.neuroimage.2023.120143

1027   88.   Simon A, Loquet G, Ostergaard J, Bech S. Cortical auditory attention decoding

1028   during music and speech listening. IEEE Trans Neural Syst Rehabil Eng.

1029   2023;31:2903–11.

1030   89.   Besle J, Fischer C, Bidet-Caulet A, Lecaignard F, Bertrand O, Giard MH. Visual

1031   activation and audiovisual interactions in the auditory cortex during speech

1032   perception: Intracranial recordings in humans. J Neurosci. 2008;28(52):14301–10.

90. Mégevand P, Mercier MR, Groppe DM, Golumbic EZ, Mesgarani N, Beauchamp
MS, et al. Crossmodal phase reset and evoked responses provide complementary
mechanisms for the influence of visual speech in auditory cortex. J Neurosci.
2020;40(44):8530–42.

91. Willems RM, Frank SL, Nijhof AD, Hagoort P, Van Den Bosch A. Prediction during
natural language comprehension. Cereb Cortex. 2016;26(6):2506–16.

92. Patel AD, Morgan E. Exploring cognitive relations between prediction in language
and music. Cogn Sci. 2017;41:303–20.

93. Rohrmeier MA, Koelsch S. Predictive information processing in music cognition. A
critical review. Int J Psychophysiol [Internet]. 2012;83(2):164–75. Available from:
http://dx.doi.org/10.1016/j.ijpsycho.2011.12.010

94. Merchant H, Grahn J, Trainor L, Rohrmeier M, Fitch WT. Finding the beat: A neural
perspective across humans and non-human primates. Philos Trans R Soc B Biol Sci.
2015;370(1664).

95. Quiroga-Martinez DR, C. Hansen N, Højlund A, Pearce M, Brattico E, Vuust P.
Musical prediction error responses similarly reduced by predictive uncertainty in
musicians and non-musicians. Eur J Neurosci. 2020;51(11):2250–69.

96. Doelling KB, Florencia Assaneo M, Bevilacqua D, Pesaran B, Poeppel D. An
oscillator model better predicts cortical entrainment to music. Proc Natl Acad Sci U
S A. 2019;116(20):10113–21.

97. Meyer L, Sun Y, Martin AE. Synchronous, but not entrained: exogenous and
endogenous cortical rhythms of speech and language processing. Lang Cogn

1055  Neurosci [Internet]. 2020;35(9):1089–99. Available from:

1056  https://doi.org/10.1080/23273798.2019.1693050

1057 98. Zou J, Xu C, Luo C, Jin P, Gao J, Li J, et al. Θ-Band cortical tracking of the speech

1058  envelope shows the linear phase property. eNeuro. 2021;8(4):1–7.

1059 99. Zoefel B, ten Oever S, Sack AT. The involvement of endogenous neural oscillations

1060  in the processing of rhythmic input: More than a regular repetition of evoked neural

1061  responses. Front Neurosci. 2018;12(MAR):1–13.

1062 100. Giraud A, Kleinschmidt A, Poeppel D, Lund TE, Frackowiak RSJJ, Laufs H.

1063  Endogenous cortical rhythms determine cerebral specialization for speech perception

1064  and production. Neuron. 2007;56(6):1127–1134.

1065 101. Duecker K, Doelling KB, Breska A, Coffey EBJ, Sivarao D V., Zoefel B.

1066  Challenges and Approaches in the Study of Neural Entrainment. J Neurosci.

1067  2024;44(40):1–11.

1068 102. Tezcan F, Weissbart H, Martin AE. A tradeoff between acoustic and linguistic

1069  feature encoding in spoken language comprehension. Elife. 2023;12:1–24.

1070