A portable and dual-inducible control system for multistep biosynthetic pathways
in Gram-negative bacteria

Andrés Felipe Carrillo Rincón[1,2], Alexandra J. Cabral[1], Andras Gyorgy[2], Natalie G. Farny[1,3]*

1. Department of Biology and Biotechnology, Worcester Polytechnic Institute, Worcester, MA, USA
2. Division of Engineering, NYU Abu Dhabi, Abu Dhabi, UAE
3. Program in Bioinformatics and Computational Biology, Worcester Polytechnic Institute,
Worcester, MA, USA

*Correspondence: nfarny@wpi.edu

Keywords: inducible promoters, dual expression, reporter system, lycopene, β-carotene, luciferase,
Escherichia coli, *Pseudomonas putida*, *Vibrio natriegens*

Abstract

The successful production of industrially relevant natural products hinges on two key factors: the
cultivation of robust microbial chassis capable of synthesizing the desired compounds, and the availability
of reliable genetic tools for expressing target genes. The development of versatile and portable genetic
tools offers a streamlined pathway to efficiently produce a variety of compounds in well-established
chassis organisms. The σ^{70}*lac* and *tet* expression systems have shown effective regulation and robust
expression of recombinant proteins across various Gram-negative bacteria. To leverage their advantages,
here both expression systems were combined into a single plasmid and assessed for their performance in
producing fluorescent reporters as well as the terpenoids lycopene and β-carotene. This rapid approach
enabled the straightforward transformation of the well-established microorganisms *Escherichia coli*,
Pseudomonas putida, and *Vibrio natriegens* into efficient microbial cell factories. The dynamic range and
the basal expression levels of the σ^{70} expression systems were further enhanced through the incorporation
of translational control mechanisms via toehold switches. This improvement was assessed using the highly
sensitive luciferase reporter system. This study presents the development and remaining challenges of a
versatile genetic tool that is portable across well-established bacterial chassis and capable of controlling
the expression of multiple genes, thus facilitating the biosynthesis and study of natural products.

Graphical Abstract

Highlights

- A dual-inducible duet-expression system is described for Gram-negative organisms.
- Production of lycopene and β-carotene is demonstrated in *E. coli, P. putida,* and *V. natriegens*.
- Incorporation of a toehold switch effectively reduces leaky expression of target genes in the uninduced state.
- The σ^{70} duet *lac/tet* expression system is a versatile tool for multi-gene biosynthesis across different bacterial chassis.

1. Introduction

Modern sequencing techniques have unveiled a myriad of unknown compounds and proteins encoded by microorganisms with potential industrial applications ranging from biotherapeutics to sustainable manufacturing (Gomez-Escribano, Alt et al. 2016, Wu, Yin et al. 2022). However, these microorganisms are not always easy to cultivate, and manipulating their genes poses a central challenge in unlocking their full potential (Katz and Baltz 2016). *Escherichia coli* is typically the primary choice for expressing proteins from other organisms due to its robust genetic toolkit and well-established cultivation parameters (Baeshen, Al-Hejin et al. 2015). Unfortunately, *E. coli* may struggle to produce toxic genes or complex proteins necessary for the synthesis of natural products (Snijder and Hakulinen 2016, İncir and Kaplan 2024). To overcome these limitations of *E. coli*, alternative hosts like *Pseudomonas putida* and *Vibrio natriegens* have become appealing chassis for producing recombinant proteins. They offer standard laboratory techniques akin to *E. coli*, along with robust genetic and metabolic backgrounds, and rapid growth characteristics (Loeschcke and Thies 2015, Becker, Wimberger et al. 2019). These microorganisms have successfully produced natural products such as non-ribosomal peptides, polyketides, terpenoids, fatty acids, and polymers (Gemperlein, Zipf et al. 2016, Gemperlein, Hoffmann et al. 2017, Ellis, Tschirhart et al. 2019, Hernandez-Arranz, Perez-Gil et al. 2019, Wang, Tschirhart et al. 2020). However, their genetic toolkit is not as versatile and developed as that of *E. coli*.

The inducible *tet* and *lac* promoters, optimized for σ^{70} recognition, have shown precise regulation and robust gene overexpression in various Gram-negative bacteria. However, these genetic tools have only been assessed for their efficacy in producing the reporter protein mCardinal and the cocaine esterase CocE (Carrillo Rincón and Farny 2023). Multiple biological processes necessitate the coordinated activity of numerous genes, such as those involved in synthesizing natural products encoded by biosynthetic gene clusters (Jensen 2016, Meyer, Segall-Shapiro et al. 2019). To facilitate multi-gene expression, the pET system was modified to concurrently express various genes within the same *E. coli* BL21 strain using the

pETDuet system (Tolia and Joshua-Tor 2006). However, in pETDuet, both multiple cloning sites are induced

by IPTG, preventing independent or sequential control (Zhang, Liu et al. 2012). To overcome this

limitation, the *E. coli* "marionette" strain was developed containing twelve inducible systems, including

the well-known lactose, tetracycline and arabinose expression systems among others (Meyer, Segall-

Shapiro et al. 2019). Additionally, the synergistic effect of combined promoters to produce target

recombinant proteins in *E. coli* has demonstrated differing levels of success thus far (Öztürk, Ergün et al.

2017). A dual-inducible duet expression system was also engineered in *P. putida* utilizing the original *lac*,

tet and *bad* promoters, resulting in the successful production of fluorescent proteins and an inducible

CRISPR interference system (Gauttam, Mukhopadhyay et al. 2020, Gauttam, Mukhopadhyay et al. 2021).

Despite these significant achievements, the original promoters utilized in these studies lack the strength

and versatility of the optimized σ^{70} *lac* and *tet* promoter (Carrillo Rincón and Farny 2023).

Adapting the consensus sequences recognized by the σ^{70} factor to the *lac* and *tet* promoters

significantly enhanced their strength, surpassing the efficiency of the pET system in *E. coli* (Carrillo Rincón

and Farny 2023). Nevertheless, the uninduced state of the optimal σ^{70}V2LacI expression system was

scarcely repressed by its transcriptional regulator LacI, thereby negating the advantages of having an

inducible system (Carrillo Rincón and Farny 2023). One strategy to enhance the fold-change and minimize

the leaky expression of the target gene, thereby reducing unnecessary consumption of cellular resources

required for cell growth, involves the adaptation of toehold switch riboregulators (Hwang, Kim et al.

2021). In the toehold-mediated strand displacement (TMSD) strategy, the toehold sequesters the

ribosome binding site (RBS) of an mRNA molecule, and translation is activated in the presence of the

trigger RNA complementary to the toehold sequence (Wang and Simmel 2022). Toehold switches have

shown to significantly enhance the mean dynamic range by over 400-fold (Green, Silver et al. 2014). Thus,

incorporating them into strong inducible promoters, like the σ^{70}V2LacI, could effectively reduce leakage

in the uninduced state.

In this study, we showed that the strong inducible promoters σ^{70}V2Tc and σ^{70}V2/3Lac, recognized

by the σ^{70}, can be tailored for a dual-inducible duet-expression system, facilitating the overexpression of

multiple genes in well-studied microorganisms such as *E. coli*, *P. putida*, and *V. natriegens*. The efficacy of

this duet expression system was confirmed through the expression of reporter systems mCardinal and

sfGFP, as well as the production of terpenoids lycopene and β-carotene. Furthermore, we enhanced the

fold-change and minimized the leakiness of these expression systems by incorporating toehold switches,

a refinement validated by the sensitive reporter system luciferase. In summary, our findings illustrate that

the σ^{70} duet *lac/tet* expression system offers a versatile gene tool for activating multiple genes with

various inducers across different bacterial chassis. This approach simplifies the production of multiple

recombinant proteins or natural products. Additionally, the incorporation of a toehold switch effectively

reduces the expression of target genes in the uninduced state, a crucial feature for sensitive or toxic

applications.

2.   Materials and Methods

2. 1 Bacterial strains and culture conditions

E. coli DH10B, *P. putida* JE90 (Elmore, Furches et al. 2017) and *V. natriegens* V. max (Becker,

Wimberger et al. 2019) were cultured in LB media supplemented with Kanamycin when required (50

µg/mL for *E. coli* and *P. putida*, and 400 µg/mL for *V. natriegens*). *E. coli* and *P. putida* cells were

transformed by electroporation, and *V. natriegens* by chemical transformation with the plasmids listed in

Table 1. *E. coli* was cultivated at 37°C and *P. putida* and *V. natriegens* at 30°C at 220 rpm.

2.2 Plasmid construction

Phusion® High Fidelity Polymerase (Thermo Scientific) and primers synthesized by Integrated DNA

Technologies (IDT) were used in all PCR amplifications for plasmid construction. Plasmids were

constructed using Gibson Assembly EX Cloning Kit or T4 DNA ligase (Thermo Scientific) according to

manufacturer's instructions. Plasmids were transformed into ElectroMAX DH10B Cells (Thermo Scientific)

according to manufacturer's instructions. Transformants were selected on LB (Miller) agar plates

containing 50 mg/L kanamycin sulfate for selection and incubated at 37°C. Plasmids were constructed

using a combination of ligation of phosphorylated oligonucleotides, and Gibson Assembly. Sequences of

all plasmids were confirmed using Sanger sequencing performed by Quintarabio. The backbone of the

pJH0204 vector was modified via Gibson to incorporate in the multicloning site the restriction sites BamHI,

NdeI, HindIII, KpnI, EcoRI, PstI, XbaI, NcoI, XhoI and eliminate the EcoRI and NcoI sites from the kanamycin

cassette, thus generating the plasmid pJH0204F. Then, the σ^{70}V2TcR-mCardinal transcriptional unit was

cloned into pJH0204F using BamHI and PstI. The new plasmid was digested with PstI and XbaI to

accommodate the transcriptional unit placI-lacI amplified by PCR flanked by PstI and XbaI. Next, the

plasmid was digested with NcoI and XhoI to accommodate mCardinal previously amplified and flanked

with the same enzymes. Finally, the plasmid was opened with XbaI and NcoI and the V2Lac and V3Lac

promoters were cloned via Gibson generating the plasmid pσ^{70}V2TcR-mCardinal/V2LacI-mCardinal and

pσ^{70}V2TcR-mCardinal/V3LacI-mCardinal (Fig. 1). Next the sfGFP was cloned after the σ^{70}V2Tc via NdeI and

HindIII generating the plasmids pσ^{70}V2TcR-sfGFP/V2LacI-mCardinal and pσ^{70}V2TcR-sfGFP/V3LacI-

mCardinal. The crtEBI genes were PCR amplified adjusting an NdeI at the 5' of crtE and a HindIII at the 3'

of the crtI and assembled in the pUC19 vector digested with BamHI and HindIII generating the plasmid

pUC19-crtEBI. The crtEBI gene cluster was cloned into pσ^{70}V2TcR via NdeI and HindIII producing the

plasmid pσ^{70} V2TcR-crtEBI. Further, crtY was cloned via Gibson into pσ^{70}V2TcR-crtEBI digested with HindIII

generating the plasmid pσ^{70}V2TcR-crtEBIY. Finally, the crtY gene was PCR amplified flanked by NcoI and

XhoI and cloned to generate the plasmids pσ^{70}V2TcR-crtEBI/V2LacI-crtY and pσ^{70}V2TcR-crtEBI/V3LacI-

crtY. The DNA parts containing the σ^{70}V2Tc-THS-trigger/V2Tc-THS-target and σ^{70}V3Lac-THS-trigger/

V2Lac-THS-target were synthesized as gBlocks and incorporated via Gibson assembly into the plasmids

pJH0204-V2TcR-mCardinal and pJH0204- σ^{70}V3LacI-mCardinal previously digested with BamHI/NdeI and

NdeI, respectively. Further, the pσ^{70}V2TcR-Toehold-mCardinal and σ^{70}V3-2LacI-Toehold-mCardinal

plasmids were digested with NdeI/HindIII to replace the mCardinal with the luxCDABE operon, previously

PCR amplified with appropriate overlapping tails and incorporated into the plasmid via Gibson assembly.

To produce the pSC101 plasmids the ampicillin cassette was replaced via Golden Gate assembly. Briefly,

the pSC101 was PCR amplified in the outward direction with respect to the ampicillin cassette, creating

the backbone amplicon, flanked with BsaI cut sites. The kanamycin cassette was PCR amplified flanked

with BsaI cut sites that generate complementary sticky ends to the pSC101 backbone. Both DNA fragments

were assembled using the standard Golden Gate protocol generating pSC101-Kn. The cassettes containing

the luxCDABE operon with the different expression systems were digested with BamHI and XhoI and

cloned into pSC101-Kn previously amplified with primers that incorporated the BamhI/XhoI restriction

sites. The complete list of plasmids used and generated in this study is presented in Table 1. Primers

(Supplementary Table S1) and DNA sequences used in this study are presented in the supplementary

information.

2.3 Measurement of the reporter systems in the plate reader

Growth curves and fluorescence measurements were performed in 96-well clear bottom

microplates (Corning Incorporated, NY) using the BioTek Synergy H1 Hybrid Multi-Mode Reader following

the experimental approach described in (Krishnamurthi, Niyonshuti et al. 2021). Briefly, overnight cultures

of the recombinant strains were diluted into fresh LB medium supplemented with kanamycin to reach

OD_{600} 0.05. *E. coli* experiments were performed at 37^{o}C, and *P. putida* and *V. natriegens* at 30^{o}C, shaking

at 355 cycles per minute (cpm). Cell density (OD_{600}) and fluorescence (sfGFP excitation 485nm, emission

510nm; mCardinal excitation 604nm, emission 659nm) were monitored every hour for 16 hours. *E. coli*

cultures were induced at 3 hours with 0.2 mM IPTG and/or 0.1 µg/mL aTc, *P. putida* at 3 hours with 5 mM

IPTG and/or 1 µg/mL aTc, and *V. natriegens* at 2 hours with 0.2 mM IPTG and/or 0.1 µg/mL aTc when

required.

2.4 Lycopene and β-carotene determination

Lycopene and β-carotene were measured by UHPLC following the procedure described in

(Bohoyo-Gil, Dominguez-Valhondo et al. 2012) using a Shimadzu Nexera UHPLC. Lycopene and β-carotene

were extracted with acetone as described in (Xu, Xu et al. 2018) with the following modifications. 5% of

overnight cultures of the recombinant *E. coli*, *P. putida* and *V. natriegens* strains containing either the LYC

operon or the crtEBIY genes were inoculated in 5 mL of fresh LB media supplemented with kanamycin and

grown at 220 rpm and 37^{o}C for *E. coli* and 30^{o}C for *P. putida* and *V. natriegens*. The cultures were induced

at OD_{600} 0.7 with aTc, IPTG, or both accordingly. After the induction the cultures were grown for 4 hours,

2 mL samples were collected, centrifuged at 14,000 rpm, the supernatant was discarded, and the cell

pellets were stored at -80^{o}C. The cell pellets were resuspended with 200 µL acetone and incubated at 55^{o}C

for 15 min in the dark followed by centrifugation at 14,000 rpm, then the supernatant was filtered through

a 0.22 μm pore-size nylon membrane for UHPLC analysis. Peak values at 460nm were measured for

analysis. Standard curves were generated and used to calculate concentration, shown in Supplementary

Figures S1 and S2.

Bacterial cultures for UV-vis spectrophotometric analysis were backdiluted to OD600 = 0.05,

grown at 30°C or 37°C as appropriate for the strain, then induced with aTc at three hours, then grown for

an additional 17 hours. Carotenoids were extracted for UV-vis spectrophotometric analysis by acetone

extraction as described (Alper, Miyaoku et al. 2005). Briefly, 5mL of each culture was spun down, media

was discarded from the bacterial pellet, and the pellet was re-suspended in 2.5mL of acetone. Samples

were heated at 55ºC for 15 minutes with intermittent vortexing. Samples were spun and the lycopene or

β-carotene content in the supernatant was analyzed by UV-vis spectrophotometry. The determination of

carotenoid concentration by UV-vis spectrophotometry was performed by scanning from 300 – 600nm on

a Thermo Scientific Evolution 300 Spectrophotometer, essentially as described (Karnjanawipagul,

Nittayanuntawech et al. 2010), using measurements at 456nm for β-carotene and 474nm for lycopene.

Standard curves were generated and used to calculate concentration, shown in Supplementary Figures S3

and S4.

2.5 Bioluminescence assay

E. coli or *P. putida* overnight cultures supplemented with kanamycin (LB, 37°C with shaking, 220 rpm)

were diluted 1/200 in fresh LB in 96-well clear bottom microplates (Corning Incorporated, NY) shaking at

355 cycles per minute (cpm) and induced after three hours of growth with 0.2/ 5 mM IPTG and 0.1/1.0

μg/mL aTc accordingly. Bioluminescence (gain: 100) and cell density (OD_{600}) were monitored every 20

minutes with the BioTek Synergy H1 Hybrid Multi-Mode Reader instrument for 20 hours.

3.   Results and Discussion

3.1 Development of the dual expression system

The σ^{70}V2TcR and σ^{70}V3LacI expression systems exhibited tight regulation and robust induction in

E. coli in the presence of anhydrotetracycline (aTc) and IPTG, respectively, while in *P. putida* and *V.*

natriegens the σ^{70}V2TcR and σ^{70}V2LacI demonstrated superior performance as inducible promoters

(Carrillo Rincón and Farny 2023). Our previous findings indicated that multicopy plasmids like pUC19

cannot effectively support LacI expression, even with its native weak promoter. Therefore, we developed

the dual expression system using the pJH0204 vector as a backbone, which contains the pColE1 replication

origin and *attB* sites specific for the bxb1 recombinase (Elmore, Dexter et al. 2023). Each transcriptional

unit of the dual expression system was insulated by terminators (Kelly, Taylor et al. 2019), and the

transcriptional regulators *tetR* and *lacI* were positioned between and in opposite directions of the

inducible promoters σ^{70}V2Tc and σ^{70}V2Lac or σ^{70}V3Lac (σ^{70}V2/3Lac), blocking undesired transcription of

the open reading frames (ORF). Additionally, two different multiple cloning sites were placed after the

inducible promoters σ^{70}V2Tc and σ^{70}V2/3Lac to facilitate the incorporation of target genes (Fig. 1A).

The dual expression system was assessed in *E. coli*, *P. putida*, and *V. natriegens* to evaluate: i)

independent control of two different fluorescent proteins, ii) potential synergy when expressing the same

protein, and iii) control of the biosynthetic gene cluster (BGC) responsible for the biosynthesis of lycopene

and β-carotene. Expression of various recombinant proteins was achieved by placing sfGFP under the

control of σ^{70}V2TcR and mCardinal under the control of σ^{70}V2/3LacI (Fig. 1C). Synergy was gauged using

the reporter system mCardinal, which was incorporated into both inducible promoters of the dual

expression system (Fig. 1B). Finally, production of the terpenoids lycopene and β-carotene by the dual

expression system was achieved by the assembly of the crtEBIY genes from *Pantoea ananatis* (Hernandez-

Arranz, Perez-Gil et al. 2019), which were synthesized with the codon usage for *E. coli.* The crtEBI (LYC

operon) was assembled in the pUC19 vector via Gibson assembly, adapting the ribosome binding site

(RBS) of J23119 (Gong, Liu et al. 2016) to crtB and the RBS pJL1 (Wang, Li et al. 2018) to crtI, then the

synthetic construct was transferred to the σ^{70}V2TcR expression system to evaluate lycopene production

(Fig. 1D). β-carotene production was evaluated with the incorporation of the crtY gene with the J23119

RBS to σ^{70}V2TcR-crtEBI yielding σ^{70}V2TcR-crtEBIY (Fig. 1E). The dual control of the crtEBI operon together

with the crtY gene to produce β-carotene was achieved by adapting the crtEBI to the σ^{70}V2TcR expression

system and the crtY gene to σ^{70}V2/3LacI expression system present in the dual vector (Fig. 1F).

Figure 1. Genetic circuits used in this study. A. Design of the dual expression systems σ⁷⁰V2TcR-V2LacI and σ⁷⁰V2TcR-V3LacI. **B.** Dual expression system controlling expression of sfGFP from the σ⁷⁰V2Tc promoter and mCardinal from the σ⁷⁰V2Lac or σ⁷⁰V3Lac promoters. The σ⁷⁰V3Lac promoter constructs are used for expression in *E. coli,* and the σ⁷⁰V2Lac constructs are used in *P. putida* and *V. natriegens.* **C.** Dual expression system controlling expression of mCardinal under both the σ⁷⁰V2Tc promoter and the σ⁷⁰V2Lac or σ⁷⁰V3Lac promoter. **D.** σ⁷⁰V2TcR expression system controlling the LYC operon (crtEBI) for lycopene production. **E.** σ⁷⁰V2TcR expression system controlling expression of crtEBIY for β-carotene production. **F.** Dual expression system controlling expression of crtEBI from the σ⁷⁰V2Tc promoter and crtY from the σ⁷⁰V2LacI or σ⁷⁰V3LacI promoters.

231   **Table 1.** List of plasmids evaluated in this study.

Name	Expression system 1	ORF1	Expression system 2	ORF2	Source
$p\sigma^{70}$ V2TcR-mCardinal	V2TcR	mCardinal			(Carrillo Rincón and Farny 2023)
$p\sigma^{70}$ V2lacI-mCardinal	V2lacI	mCardinal			(Carrillo Rincón and Farny 2023)
$p\sigma^{70}$ V3LacI-mCardinal	V3LacI	mCardinal			(Carrillo Rincón and Farny 2023)
$p\sigma^{70}$ V2TcR-sfGFP	V2TcR	sfGFP			(Carrillo Rincón and Farny 2023)
$p\sigma^{70}$ V2lacI-sfGFP	V2LacI	sfGFP			(Carrillo Rincón and Farny 2023)
$p\sigma^{70}$ V3LacI-sfGFP	V3LacI	sfGFP			(Carrillo Rincón and Farny 2023)
pJH0204F					This study
$p\sigma^{70}$ V2TcR-mCardinal/V2LacI-mCardinal	V2TcR	mCardinal	V2LacI	mCardinal	This study
$p\sigma^{70}$ V2TcR-mCardinal/V3LacI-mCardinal	V2TcR	mCardinal	V3LacI	mCardinal	This study
$p\sigma^{70}$ V2TcR- sfGFP/V2LacI-mCardinal	V2TcR	mCardinal	V2LacI	sfGFP	This study
$p\sigma^{70}$ V2TcR- sfGFP/V3LacI-mCardinal	V2TcR	mCardinal	V3LacI	sfGFP	This study
pUC19-crtEBI	-	crtEBI			This study
$p\sigma^{70}$ V2TcR-crtEBI	V2TcR	crtEBI			This study
$p\sigma^{70}$ V2TcR-crtEBIY	V2TcR	crtEBIY			This study
$p\sigma^{70}$ V2TcR-crtEBI/V2LacI-crtY	V2TcR	crtEBI	V2LacI	crtY	This study
$p\sigma^{70}$ V2TcR-crtEBI/V3LacI-crtY	V2TcR	crtEBI	V3LacI	crtY	This study
$p\sigma^{70}$ V2TcR-Toehold-mCardinal	V2TcR	mCardinal			This study

pσ^{70} V3-2LacI-Toehold-mCardinal	V3-2LacI	mCardinal			This study
pσ^{70} V2TcR-Toehold-luxCDABE	V2TcR	luxCDABE			This study
pσ^{70} V3-2LacI-Toehold-luxCDABE	V3-2LacI	luxCDABE			This study
pσ^{70} V2TcR -luxCDABE	V2TcR	luxCDABE			This study
pσ^{70} V2LacI-luxCDABE	V2LacI	luxCDABE			This study
pSC101-Kn					This study
pSC101Kn σ^{70} V2TcR -luxCDABE	V2TcR	luxCDABE			This study
pSC101Kn σ^{70} V2LacI-luxCDABE	V2LacI	luxCDABE			This study
pSC101Kn σ^{70} V2TcR-Toehold-luxCDABE	V2TcR	luxCDABE			This study
pSC101Kn σ^{70} V3-2LacI-Toehold-luxCDABE	V3-2LacI	luxCDABE			This study

3.2 Independent and concurrent control of two fluorescent proteins with the dual expression system

Duet vectors are available for *E. coli* and *Pseudomonas fluorescens*, but they are solely inducible by either IPTG or arabinose (Tolia and Joshua-Tor 2006, Nakata 2017), whereas dual-inducible duet expression vectors have been developed specifically for *Corynebacterium glutamicum* (Gauttam, Desiderato et al. 2019) and *P. putida* (Gauttam, Mukhopadhyay et al. 2020, Gauttam, Mukhopadhyay et al. 2021). Unfortunately, there is currently no a versatile dual overexpression system that permits the differential expression of distinct sets of genes through two independent inducible promoters, which can be used across various Gram-negative species. We have previously demonstrated the strength and portability of the σ^{70}V2TcR and σ^{70}V2/3LacI promoters separately in *E. coli, P. putida*, and *V. natriegens* (Carrillo Rincón and Farny 2023). Thus, to test the independent and concurrent regulation of two proteins simultaneously from these promoters, we introduced two different reporter proteins into the dual-inducible σ^{70}V2TcR/V2LacI and σ^{70}V2TcR/V3LacI expression systems (Fig. 1B) and assessed their production in *E. coli*, *P. putida*, and *V. natriegens*. Specifically, the transcriptional unit V2TcR controlled the expression of sfGFP, while the V2LacI or V3LacI unit regulated mCardinal production (Fig. 2). This configuration was selected because the species under investigation are susceptible to autofluorescence in the yellow-green spectrum, which could interfere with the measurement of sfGFP (Carrillo Rincón and Farny 2023). The σ^{70}V2TcR promoter was therefore selected to regulate the expression of sfGFP due to its tighter control in the OFF state, facilitating a more sensitive estimation of the production of the reporter system. On the other hand, the σ^{70}V2/3LacI systems, which tend to leak considerably in *V. natriegens* (Carrillo Rincón and Farny 2023), control the expression of the far-red fluorescent protein mCardinal. Importantly, at this wavelength, the cells do not produce endogenous fluorescence (Canty, Hariharan et al. 2018).

To create a direct comparison, the induction of the individual expression systems σ^{70}V2TcR-sfGFP and σ^{70}V2/3LacI-mCardinal were measured in *E. coli* (Fig. 2A), *P. putida* (Fig. 2B) and *V. natriegens* (Fig.

2C). The dual-inducible σ^{70}V2TcR/V2LacI and σ^{70}V2TcR/V3LacI expression systems were also measured in

the same species (Fig. 2D-F). Under the control of σ^{70}V2TcR, sfGFP production was higher in the dual

system by 1.2-fold in *E. coli* and *P. putida*, and 1.7-fold in *V. natriegens* at 16 hours of induction (Fig. 2D-

F, left panels). However, an opposite trend was observed in the dual expression system for production of

mCardinal under the control of σ^{70}V2/3LacI, which was significantly reduced, resulting in 6.2-fold, 2.4-fold,

and 2.0-fold less red emission in *E. coli*, *P. putida*, and *V. natriegens*, respectively (Fig. 2D-F, right panels).

Interestingly, when the cultures containing the dual expression system were co-induced with aTc

and IPTG, both green and red fluorescence reached comparable levels compared to cultures induced with

only one inducer. So, while the maximum expression from each promoter was distinct from the maxima

achieved in the single promoter systems, concurrent induction of both promoters did not impact their

transcriptional output. This suggests that the promoters (σ^{70}V2TcR and σ^{70}V2/3LacI) are barely affected

by the activation of the other, reaching almost their maximum activity either upon sole activation or

concurrent activation (Fig. 2G). The three bacterial species investigated, each containing the dual

expression system, displayed barely detectable green fluorescence in the uninduced state, indicating tight

regulation of the σ^{70}V2TcR expression system (Fig. 2G). However, as expected, some mCardinal leakage

was observed originating from the σ^{70}V2LacI-mCardinal expression system in *V. natriegens* (Fig. 2F-G).

The dual expression system constitutively expresses the transcriptional regulators TetR and LacI,

both of which have been reported to exhibit toxicity at high concentrations (Lutz and Bujard 1997,

Romanuka, van den Bulke et al. 2009, Stetz, Carter et al. 2016), and each sensor requires cellular resources

(for example, ribosomes) to function, and the activation of one sensor can influence another indirectly

due to resource competition (Gyorgy, Jiménez et al. 2015). We anticipated a lower performance for both

promoters in the dual expression system, due to the metabolic burden imposed by the simultaneous

presence of TetR and LacI. However, this scenario was only observed with σ^{70}V2/3LacI, which exhibited

reduced production of mCardinal, which was particularly pronounced in *E. coli*, thus suggesting its

sensitivity to both transcriptional regulators. Conversely, the levels of sfGFP appeared to be enhanced by

σ^{70}V2TcR in the dual expression system. One hypothesis for this observation is that the stress response

triggered by the combination of expressing two toxic regulators in the presence of aTc results in the

induction of the RpoS regulon, causing increased expression of *rpoB* and several genes involved in

nucleotide salvage and purine biosynthesis (Bie, Zhang et al. 2023). Furthermore, it leads to the induction

of heat shock proteins (Bie, Zhang et al. 2023). Additionally, there is increased transcription of genes

participating in the SOS pathway, promotion of glucose metabolism, and induction of the oxidative stress

response (Bie, Zhang et al. 2023). To mitigate the adverse effect of these stresses, the cells produce

endogenous fluorophores and/or metabolic byproducts increasing the green emission of green light

(Monici 2005). For instance, *P. putida* secretes pyoverdine, a soluble fluorescent yellow-green pigment,

under iron-limiting conditions (Ringel and Brüser 2018). We predict that the σ^{70}V2TcR in the dual

expression system did not in fact display increased transcriptional activity. Instead, it is possible that the

presence of endogenous molecules interfering in the green spectrum, not originating from the

recombinant protein, amplified the sfGFP signal due to activation of cellular stress responses.

Figure 2. sfGFP and mCardinal production in *E. coli*, *P. putida*, and *V. natriegens* containing the dual expression. For all samples, the fluorescence mean of sfGFP signal (excitation 485, emission 510) and mCardinal signal (excitation 605, emission 659) was normalized by the cell density (OD$_{600}$). **A-C.** *E. coli* (A), *P. putida* (B) and *V. natriegens* (C) expressing the single reporter systems σ^{70}V2TcR-sfGFP (left panels) or σ^{70}V2/3LacI-mCardinal (right panels). **D-F.** *E. coli* (D), *P. putida* (E) and *V. natriegens* (F) expressing the dual expression systems σ^{70}V2TcR-sfGFP/σ^{70}V3LacI-mCardinal (in *E. coli*) and σ^{70}V2TcR-sfGFP/σ^{70}V2LacI-mCardinal (in *P. putida* and *V. natriegens*) and measuring sfGFP (left panels) and mCardinal (right panels). Error bars are +/- S.D., n=3. **G.** Matrix display of both sfGFP and mCardinal expression of individual and concurrent inductions of the dual expression systems.

3.3 Evaluating synergistic potential and metabolic cost of the dual expression system

From our first dual expression system experiments, we anticipated a metabolic cost to the cell.

We observed that while sfGFP fluorescence from the σ^{70}V2Tc promoter increased relative to a single

expression system, mCardinal fluorescence from the σ^{70}V2/3Lac promoters was significantly decreased.

If, as we hypothesized, the increased sfGFP was an artifact of cellular stress caused by the combination of

metabolic load and aTc exposure, then one would predict that driving the same protein from both

promoters would have no synergistic effect on expression over a single promoter system. We have

previously shown that increasing the copy number of the same promoter type in *E. coli* effectively

enhanced the production of mCardinal by the σ^{70}V2TcR expression system, while it had a detrimental

effect on the σ^{70}V3LacI expression system due to the toxicity associated with the transcriptional regulator

LacI (Carrillo Rincón and Farny 2023). Hence, to test the synergistic potential of the duet expression

system σ^{70}V2TcR/V3LacI, a version was made where both promoters regulate mCardinal (Fig 1C). The *E.*

coli strain harboring the duet expression system σ^{70}V2TcR/V3LacI exhibited 2-fold and 4-fold lower red

fluorescence compared (Fig. 3B) to strains containing σ^{70}V2TcR and σ^{70}V3LacI individually when induced

with aTc and IPTG, respectively (Fig. 3A). This observation supports the hypothesis that the increased

fluorescence signal in the σ^{70}V2TcR/V3LacI was an artifact of some kind, possibly a product of cellular

stress. Synergy was observed by the addition of IPTG and aTc to the σ^{70}V2TcR/V3LacI *E. coli* strain

improving mCardinal production by 1.5-fold against the induction obtained with aTc alone (Fig. 3B).

However, the mCardinal production in the duet system reached about the same total amount of

mCardinal as the σ^{70}V2TcR expression system alone (Fig. 3A,B). These results indicate that the tandem

version of the synthetic promoters σ^{70}V2TcR/V3LacI does not offer a real synergy in *E. coli*, but rather an

alternative to co-express different genes at high levels using different inducers.

In *P. putida* the σ^{70}V2TcR expression system exhibited similar performance in the single and dual

version (Fig. 3C & D). Conversely, induction of the dual system with IPTG resulted in a 3-fold decrease

relative to the σ^{70}V2lacI system alone (Fig. 3C & D). The simultaneous addition of both inducers (aTc and

IPTG) increased mCardinal expression by 1.1-fold over the σ^{70}V2TcR and σ^{70}V2LacI expression systems in

the dual system, demonstrating a minimal, statistically insignificant synergistic effect in this host. The duet

expression system σ^{70}V2TcR/V2LacI markedly reduced the efficiency of the synthetic promoters in *V.*

natriegens compared to their single expression system counterparts, with reductions of 12-fold and 2-fold

for σ^{70}V2LacI and σ^{70}V2TcR after induction with IPTG and aTc, respectively (Fig. 3E,F). Co-induction of the

dual expression system with IPTG and aTc did not yield a synergistic effect in mCardinal production, but

achieved similar yields as obtained by induction with aTc alone (Fig. 3F).

This information collectively demonstrates that synergy cannot be attained with the dual

expression system in *E. coli* and *V. natriegens*. In these two microorganisms, the pJH0204 plasmid

containing the pColE1 origin of replication was used, rendering 30 copies per cell in *E. coli* and 300 copies

per cell in *V. natriegens*. (Hershfield, Boyer et al. 1974, Tschirhart, Shukla et al. 2019). On the contrary,

the duet expression system appears to attenuate the activity of each promoter. Rewiring the cell

metabolism imposes additional stress which affects protein synthesis and growth rate (Snoeck, Guidi et

al. 2024). Consequently, the constitutive expression of the transcriptional regulators TetR and LacI, as well

as competition for the σ^{70} factor by both promoters, may pose a significant metabolic burden. This triggers

energetic inefficiency within the cell, leading to undesirable physiological changes and placing hidden

constraints on host productivity (Wu, Yan et al. 2016). In *P. putida*, the dual expression system was directly

integrated into the chromosome, thereby reducing the quantities of the transcriptional regulators TetR

and LacI and thus lowering the metabolic cost of the synthetic expression systems. This resulted in similar

mCardinal production with σ^{70}V2TcR, while the σ^{70}V2LacI expression system showed reduced strength.

Figure 3. Assessing potential synergy between *lac* and *tet* promoters. (**A, C, E**) Time course of mCardinal production from the single expression systems σ70V2TcR-mCardinal and σ70V3LacI-mCardinal (for *E. coli*) and σ70V2LacI-mCardinal (for *P. putida* and *V. natriegens*) in *E. coli* (**A**), *P. putida* (**C**) and *V. natriegens* (**E**). (**B, D, F**) Single and double inductions of the dual σ70V2TcR-mCardinal/V2LacI-mCardinal or σ70V2TcR-mCardinal/V2LacI-mCardinal expression system in *E. coli* (**B**), *P. putida* (**D**) and *V. natriegens* (**F**). For all samples, the fluorescence mean of mCardinal signal (excitation 605nm, emission 659nm) was normalized by the cell density (OD$_{600}$). Error bars are +/- S.D., n=3.

3.4 Production of lycopene and β-carotene in *E. coli*, *P. putida* and *V. natriegens* using the σ^{70}V2TcR single expression system

359
360

To assess the capability of the dual expression system to produce natural products, we tested the

ability of the *σ^{70}*V2TcR expression system to control the synthesis of multiple genes in a biosynthetic gene

cluster (BGC). To do so, we evaluated the production of lycopene and β-carotene. Lycopene production

has already been demonstrated in *E. coli* and *P. putida* through heterologous expression of the lycopene-

producing operon (LYC), which contains the crtEBI genes from *Pantoea ananatis* (Hernandez-Arranz,

Perez-Gil et al. 2019). Lycopene serves as the biosynthetic precursor of β-carotene. Therefore, the

expression of the LYC operon together with crtY leads to β-carotene production. This process has also

been demonstrated in *E. coli* and *V. natriegens* (Yoon, Park et al. 2007, Ellis, Tschirhart et al. 2019). The

LYC and crtEBIY operons were incorporated into the *σ^{70}*V2TcR expression system (Fig. 1D,E) and the

production of lycopene and β-carotene was quantified using UHPLC (Bohoyo-Gil, Dominguez-Valhondo et

al. 2012) after 4 hours of induction with aTc. In *E. coli*, production of both lycopene and β-carotene was

achieved. However, the induced culture only yielded 1.7-fold more lycopene and 1.2-fold more β-carotene

compared to the uninduced culture when induced for four hours (Fig. 4A, B). Upon a longer induction of

17 hours, there was no significant difference in lycopene or β-carotene yields between induced and

uninduced cultures (Fig. 4C, D, and Supplementary Figure S5). A decrease in lycopene production was

noted with longer induction times under these longer culture conditions (Fig. 4C), which may be due to

the documented toxicity of high lycopene production in *E. coli* (Miguez et al., 2018). The results

demonstrate that the tight control of the *σ^{70}*V2TcR system is still unable to completely repress the

expression of the crtEBI and crtEBIY BGCs in *E. coli*. A similar observation has been previously published,

where lycopene biosynthesis in *E. coli* was controlled using the pET expression system, and also showed

significant leakage, with production reaching only 1.1-fold more after 40 hours of induction than the

uninduced culture (Xu, Xu et al. 2018). Clearly, the tight regulation of terpenoid biosynthesis in *E. coli*

remains a significant challenge.

The σ^{70}V2TcR expression system produced a maximum 0.8 mg/L of lycopene in *E. coli* cultivated

in a culture tube after only 7 hours of growth (4 hours with the inducer aTc) (Fig. 4A), and a maximum of

0.46 mg/L β-carotene after 20 hours of growth (17 hours with the inducer aTc) (Fig. 4D). Notably, no media

optimizations were performed, nor any strain engineering beyond the addition of the crtEBI or crtEBIY

BGCs. While higher yields have been reported in *E. coli* in different strains, larger bioreactors, and

optimized media conditions (Xu, Xu et al. 2018), our results suggest that the σ^{70}V2TcR expression system

may contribute yet a further point of optimization to this system by increasing the strength and control

of the BGCs.

In recombinant strains of *P. putida* and *V. natriegens* containing the σ^{70}V2TcR-crtEBI and

σ^{70}V2TcR-crtEBIY transcriptional units, little to no lycopene or β-carotene production was detected

without induction. In both organisms, induction of the BGC by aTc led to the robust production of the

terpenoids (Fig. 4A-D). *P. putida* was reported to produce 0.0012 µg/L of lycopene after 24 hours of

growth with the pSEVA421 vector (Hernandez-Arranz, Perez-Gil et al. 2019). With the σ^{70}V2TcR expression

system, lycopene production reached 170 µg/L after only 4 hours of induction (Fig. 4A), and 1028 µg/L

after 17 hours of induction (Fig. 4C), marking a substantial increase in reported yield. Additionally, β-

carotene production by *P. putida* reached 761 µg/L, the highest yield among the three species in our study

(Fig. 4D). *V. natriegens* was reported to produce 500 µg/L of β-carotene cultivated in the rich medium

LBv2 for 20 hours (Ellis, Tschirhart et al. 2019), while in the σ^{70}V2TcR expression system achieved 286 µg/L

of lycopene (Fig. 4A) and 353 µg/L of β-carotene (Fig. 4B) in the standard LB medium and four hours of

induction with no particular optimization. These yields confirm the advantages of the σ^{70}V2TcR expression

system, and suggest that significantly increased yields would be possible in the future with the

combination of our expression system and other optimizations. The overall results demonstrate that the

σ^{70}V2TcR expression system can be efficiently used to drive production of BGCs across a range of Gram-

negative bacteria.

Figure 4. Production of lycopene and β-carotene with the σ⁷⁰V2TcR expression system in *E. coli, P. putida,* and *V. natriegens*.
Production of lycopene (**A**) and β-carotene (**B**) by the single σ⁷⁰V2TcR expression system expression system in E. *coli, P. putida*
and *V. natriegens* was measured by UHPLC by induction of OD_{600}= 0.7 cultures with aTc for 4 hours. Production of lycopene (**C**)
and β-carotene (**D**) by the single σ⁷⁰V2TcR expression system expression system in *E. coli* and *P. putida* was measured by UV-vis
spectrophotometry by induction of OD_{600}= 0.05 cultures with aTc for 17 hours. Production of lycopene (**E**) and β-carotene (**F**) by
the dual expression system in *E. coli, P. putida* and *V. natriegens*. Cultures were induced with aTc, IPTG or both inducers at $OD_{600.}$=
0.7 and cultured for 4 hours. Lycopene and β-carotene were extracted and measured by UHPLC. Error bars are +/- S.D., n=3.

3.5 Coordinated expression of the lycopene and β-carotene BGCs with the dual expression system

Natural products are often encoded by BGCs, and the ability to activate each gene at different

time points with different inducers can be advantageous, particularly when the genes are toxic or the

metabolites produced by the BGC affect the growth of the heterologous host (Kadjo and Eustáquio 2023).

In this study, we have demonstrated that the σ^{70}V2TcR expression system can effectively activate the

production of lycopene and β-carotene. As a next step, we attempted to exert control of β-carotene at

different points in the production pathway using two different inducers, aTc and IPTG. The LYC operon

was integrated into the σ^{70}V2TcR expression system, while the crtY gene was controlled by the σ^{70}V2/3LacI

expression system (Fig. 1F). The plasmid pσ^{70}V2TcR-crtEBI/V3LacI-crtY was transformed into *E. coli*, and

the plasmid pσ^{70}V2TcR-crtEBI/V2lacI-crtY was transformed into *P. putida* and *V. natriegens*. In all three

bacterial species, no lycopene accumulation was observed upon sole induction by aTc despite the crtEBI

genes being controlled by the σ^{70}V2TcR expression system (Fig. 4E). This result in *E. coli* can be explained

by the leakage of the crtY under the control of the synthetic promoter σ^{70}V3LacI. The high processivity of

the lycopene cyclase CrtY to metabolize lycopene directly into β-carotene requires tighter regulation of

the promoter to completely repress this gene, and the σ^{70}V3LacI expression systems lacks this control. On

the other hand, while the σ^{70}V2TcR-crtEBI system showed no measurable leakage in *P. putida* and *V.*

natriegens using fluorescent reporters, we previously showed that the σ^{70}V2lacI expression system does

leak (Carrillo Rincón and Farny 2023). Consequently, the lycopene produced after the addition of aTc was

directly processed into β-carotene by the CrtY present in the cells due to leakage of σ^{70}V2LacI.

Production of β-carotene in *E. coli* was observed regardless of the absence or presence of the

inducers aTc and IPTG (Fig. 4F), demonstrating that the dual σ^{70}V2TcR/ σ^{70}V3LacI expression system would

require tighter regulation in the OFF state to effectively downregulate the target gene in the absence of

the inducer. In *P. putida* and *V. natriegens*, production of β-carotene was only achieved when the cultures

were induced with aTc or with aTc+IPTG (Fig. 4F). IPTG induction alone did not lead to β-carotene

production because the crtEBI genes were not transcribed by the σ^{70}V2TcR expression system. Addition

of aTc resulted in β-carotene levels similar to the ones obtained when the cells were induced with

aTc+IPTG (Fig. 4F), thus indicating that the minimal leakage of the σ^{70}V2lacI-crtY in *P. putida* and *V.*

natriegens is sufficient to metabolize all lycopene produced by the σ^{70}V2TcR-crtEBI immediately into β-

carotene upon induction by aTc. The varying yields of β-carotene between *E. coli*, *P. putida*, and *V.*

natriegens highlight the differences in their suitability as hosts for natural product production. While *E.*

coli may struggle to efficiently produce complex products from BGCs, *P. putida* and *V. natriegens* appear

to be well-suited for this task, as evidenced by their superior performance in this study, almost doubling

the yields obtained from *E. coli*.

3.6 Adaptation of toehold switch to the synthetic σ^{70} expression systems and fold-change evaluation

with the reporter system mCardinal

The assessment of σ^{70} expression systems using the reporter mCardinal suggested that the

σ^{70}V2TcR promoter displays tight regulation, leading to undetectable levels of red fluorescence in the

uninduced state (Carrillo Rincón and Farny 2023). However, when the LYC operon and crtY were

incorporated into this synthetic promoter, lycopene and β-carotene were produced by *E. coli* in the

uninduced state, indicating that the promoter was not completely repressed. The processivity of enzymes,

unlike static fluorescent proteins, amplifies the effect of leaky transcription. In the σ^{70}V2/3LacI expression

systems the leakage was even more evident, as we were unable to repress the production of CrtY in the

dual expression system sufficiently to permit the intermediate accumulation of lycopene (Fig. 4E, F).

Maintaining low basal expression levels in the absence of the inducer is crucial to minimize unwanted

effects on cell growth and to prevent potential damage caused by the target protein, including issues like

self-aggregation or inclusion body formation (Schuster and Reisch 2022), or as evidenced in this study,

uncontrolled biosynthesis of a downstream product. If independent control of steps within a multistep

biosynthetic pathway is to be achieved, it will require significant improvements in control of inducible

promoter systems.

One solution to this problem involves the adaptation of RNA-based gene regulation to the gene

circuits to reducing the impact of basal expression (Hwang, Kim et al. 2021, Wang and Simmel 2022). The

current design of the synthetic σ^{70} expression system comprises a transcriptional control performed by

the transcriptional regulators TetR and LacI. Introducing an additional translational control mechanism,

such as a toehold switch (THS), alongside the existing transcriptional control, could enhance the precision

of the σ^{70} expression systems. This added layer of regulation could provide finer control over gene

expression, potentially reducing leakiness and improving inducibility. To this end, the σ^{70}V2TcR and

σ^{70}V2lacI expression systems were modified with the incorporation of the three components required to

add toehold-mediated translational control: the THS trigger, the THS target and the RBS (Green, Silver et

al. 2014). For the σ^{70}V2TcR expression system, two σ^{70}V2Tc promoters were incorporated into the system,

the first σ^{70}V2Tc promoter regulates the transcription of the THS trigger, while the second promoter is

fused to the THS target, which masks the RBS (Fig. 5A). The σ^{70}V2lacI follows the same architecture, except

the THS trigger is regulated by the V3Lac promoter due its reduced leakage compared to the V2Lac, which

controls the THS target that hides the RBS (Fig. 5B). This arrangement combines transcriptional and

translation regulation, therefore, the absence of the inducers (aTc or IPTG) results in both transcriptional

repression (via the transcriptional regulators TetR and LacI) and translational repression (via the THS

trigger) of mCardinal. Upon addition of the inducers, the constitutively expressed transcriptional

regulators are removed from the synthetic σ^{70} promoters, activating the transcription of the THS trigger,

which binds to the THS target inducing the conformational change that relieves the translational

repression of mCardinal resulting in its subsequent expression.

The comparison between the uninduced and induced states of the σ^{70}V2TcR expression system

with and without the toehold switch revealed very similar induction dynamics, as expected (Fig. 5C). It

was previously determined that the leakiness of σ^{70}V2TcR could not be observed with a fluorescent

reporter. The system may exhibit leakage even with the addition of the THS, but it would not be detectable

in this assay. The addition of aTc facilitated mCardinal production by both promoters. However, the

promoter with the THS exhibited a 2-fold reduction in the dynamic range compared to the promoter

without the switch, although it still maintained a satisfactory level of recombinant protein production (Fig.

5D). On the other hand, the addition of the THS had a significant effect on suppressing leaky transcription

from the σ^{70}V2lacI expression system (Fig. 5E). The dynamic range in the THS-σ^{70}V2lacI system improved

by 5.7-fold, but still, leaky transcription was significant. However, the riboswitch did reduce the leakiness

by 7-fold compared to the original version (Fig. 5F). Importantly, mCardinal production after IPTG

induction reached similar levels in both σ^{70}V2lacI expressions, with the variant lacking the THS producing

1.2-fold more mCardinal (Fig. 5).

These results demonstrate that THSs enhance the regulation of gene expression by modulating

translation initiation. This feature is particularly effective in inducible expression systems prone to

leakage. However, while fluorescent proteins like mCardinal offer valuable insights, they might not

capture the subtle nature of basal transcription. For instance, the evaluation of the σ^{70}V2TcR expression

system using fluorescent proteins indicated complete lack of fluorescent protein synthesis in the OFF state

when uninduced. However, the results presented for the production of lycopene in the dual expression

system revealed that the promoter leaks to some extent. Therefore, to truly assess any basal

transcriptional activity, the use of enzymatic reporter systems will be necessary, with higher sensitivity

for accurately evaluating gene expression dynamics.

Figure 5. Comparison of the σ⁷⁰ expression systems to the toehold versions controlling the expression of the reporter system

mCardinal in *E. coli*. A-B. Architecture of the synthetic gene circuits with the toehold switches (THSs) in the σ⁷⁰V2TcR system (A),

and in the σ⁷⁰V3-2LacI system (B). **C**. Time course of mCardinal production by pσ⁷⁰V2TcR-Toehold. **D**. Time course of mCardinal

production by σ⁷⁰V3-2LacI. **E.** Difference in mCardinal production between the ON and OFF states. **F.** Fold-change difference in

the red-light emission of the recombinant cells vs the wild type DH10B. The cultures were induced after 3 hours of growth with

aTc or IPTG accordingly. For all samples, the fluorescence mean of mCardinal signal (excitation 605, emission 659) was normalized

by the cell density (OD_{600}). Error bars are +/- S.D., n=3.

3.7 Evaluation of the σ⁷⁰ expression systems using luciferase as a reporter

The selection of the red fluorescent protein mCardinal to estimate the efficiency of the σ⁷⁰

expression systems was a strategic choice, to circumvent intrinsic issues associated with sfGFP and the

potential interference of green autofluorescence from cells, which could lead to inaccurate quantification

of the recombinant protein produced by the synthetic promoters (Carrillo Rincón and Farny 2023).

However, upon assessing the synthetic promoters for terpenoid production, it became evident that the

σ⁷⁰V2TcR expression system does display leaky transcription, a phenomenon not discernible with

mCardinal as the reporter. The integration of the THS into σ⁷⁰V2lacI effectively mitigated basal expression.

However, this effect was not evident in THS-σ^{70}V2TcR-mCardinal, as undetectable fluorescence in the OFF

state in this expression system was achieved even without the THS. The terpenoids lycopene and β-

carotene, aside from their nutraceutical properties, can serve as reporter systems due to their ability to

induce a color change in the cell biomass upon production (McNerney and Styczynski 2017). Indeed, the

biosynthesis of these pigments was instrumental in detecting the leakage of the σ^{70}V2TcR system.

However, accurate estimation of terpenoid yields necessitates laborious compound isolation techniques

and complex measurements. Consequently, their use as a reporter system may not be the ideal choice for

evaluating the metrics of an inducible system. To comprehensively grasp the behavior of the synthetic σ^{70}

expression systems and address the challenges posed by the limited sensitivity of mCardinal and the

intricate evaluation of terpenoid production, we opted to employ luciferase as a reporter.

The bacterial luciferase, encoded by the lux operon (luxCDABE), initiates the emission of

bioluminescence upon its expression, without requiring the addition of an external substrate (Hakkila,

Maksimow et al. 2002). The luciferase reporter was integrated into the σ^{70} expression systems and their

corresponding THS variants (Fig. 6A, B) to amplify the detection of basal expression levels, particularly in

the σ^{70}V2TcR system where the uninduced state appears completely OFF when using mCardinal (Carrillo

Rincón and Farny 2023), yet shows activity with lycopene and β-carotene. The time course evaluation

permitted the visualization of the leakage of the σ^{70}V2TcR, and also confirmed that the THS version of this

promoter diminished the leakage and amplified the signal of bioluminescence (Fig. 6C, E, G). Interestingly,

bioluminescence production with the riboswitch tends to diminish over time, suggesting that translational

regulation favors the repression of the lux operon after the mid-log phase of growth. In contrast, the

original σ^{70}V2TcR system maintains its pronounced leakage over time (Fig. 6G). In the σ^{70}V2lacI expression

system, the THS version exhibited a similar pattern to the THS- σ^{70}V2TcR system in both induced and

uninduced states, with higher basal levels, as expected (Fig 6H). The σ^{70}V2lacI variant, on the other hand,

exhibited higher leakage (Fig. 6D). Interestingly, induction with IPTG halted bioluminescence production

(Fig. 6D). This phenomenon may be attributed to the potency of this promoter, whose activation could

exceed the metabolic capacity of the cells. These results demonstrate that the reporter luciferase provides

a more sensitive assessment of the expression levels in the uninduced state compared to fluorescent

proteins like mCardinal. Importantly, the translational regulation provided by the toehold switches

efficiently amplifies the fold change and reduces the basal expression levels of the σ^{70} expression systems,

thereby improving the tight regulation of the promoters while maintaining their strength (Fig 6E, F).

However, total repression was not achieved by the adaptation of the toehold riboregulator as the

σ^{70}V2TcR-toehold expression system showed minimal (Fig 6G), and the σ^{70}V3-2LacI pronounced (Fig. 6H)

leakiness, thus prohibiting total control of the target recombinant genes.

Total gene expression control is crucial for regulating enzymatic reactions involved in the

biosynthesis of natural products, as demonstrated with the production of terpenes presented in this

study. The synthetic expression systems evaluated in this study were evaluated in the pJH0204 plasmid

which contains the pColE1 origin of replication (Carrillo Rincón and Farny 2023). This plasmid has the

advantage to integrate in the chromosome of *P. putida* and replicate in *V. natriegens*, therefore offers a

platform to use multiple chassis (Carrillo Rincón and Farny 2023). However, one potential disadvantage

of this plasmid in *E. coli* is the copy number. The pColE1 origin of replication produces 25–30 copies per

cell in *E. coli* (Hershfield, Boyer et al. 1974). To achieve tighter control over gene expression, we

transferred the synthetic expression systems to the low copy plasmid pSC101, which has approximately 3

copies per cell (Joshi, Yong et al. 2022). In this configuration, both σ^{70}V2TcR expression systems achieved

total repression of the lux operon in the uninduced state, thus indicating that in the σ^{70}V2TcR expression

system the transcriptional control provided by TetR is enough to shut down the σ^{70}V2Tc promoter (Fig.

6C, E, G). Activation by aTc successfully induced bioluminescence production; however, the THS-version

produced ∼2-fold less light, indicating that the translational control provided by the toehold affects the

activity of the promoter (Fig. 6C, E, G). A similar trend was observed with the σ^{70}V2LacI and σ^{70}V3-2LacI

expression systems in pSC101 (Fig 6D, F, H). Although total repression was not achieved, the translational

control provided by the toehold significantly reduced the basal expression levels and amplified the

dynamic range of the strong σ^{70}V2Lac promoter (Fig. 6E, H), the strongest promoter within the family of

σ^{70} expression systems.

In *P. putida*, the pJH0204 plasmid integrates at one specific location in the chromosome (Elmore,

Furches et al. 2017). In this configuration, using mCardinal as the reporter system, the σ^{70}V2TcR showed

no leakage, while the σ^{70}V2LacI exhibited minimal, barely detectable red fluorescence in the uninduced

state (Carrillo Rincón and Farny 2023). However, traces of lycopene and β-carotene were detected in the

uninduced state of the σ^{70}V2TcR (Fig. 4A, B), and only β-carotene was produced in the dual system (Fig.

4C &D), thus corroborating the leakage of σ^{70}V2LacI in this host. Since the transcriptional control provided

by the regulatory proteins TetR and/or LacI proved insufficient to fully repress the σ^{70} synthetic promoters,

even at one copy per cell, we decided to incorporate translation repression in *P. putida*. The luciferase

reporter system confirmed the leakage of the σ^{70} expression systems in this host (Fig. 7D), and the addition

of the toehold amplified the dynamic range and reduced the luminescence emission in the uninduced by

$\sim$2-fold in both expression systems (Fig. 7D). These results demonstrate that the transcriptional control

provided by the regulatory proteins can be enhanced by translational control via THS in *P. putida*, as noted

in the σ^{70}V2TcR system. However, these two layers of control are not sufficient to completely turn off the

σ^{70}V2LacI expression system. Unlike in *E. coli*, plasmid copy number is not a parameter that could be

further optimized in *P. putida. V. natriegens*, on the other hand, has demonstrated the ability to maintain

different plasmids with various origins of replication, including pUC, pSC101, p15a, and pBR1 (Tschirhart,

Shukla et al. 2019). The pJH0204, derivative of pColE1 ori, shows the highest copy number in this host

with 313 copies per cell (Tschirhart, Shukla et al. 2019). Therefore, we speculate that using a low-copy

plasmid in conjunction with the THS could also improve the control of the synthetic σ^{70} expression system

in this chassis bacteria.

Figure 6. Comparison of the σ⁷⁰ expression systems to the toehold versions controlling the expression of the lux operon in *E. coli* using pCOLE1 and pSC101 derived plasmids. **A-B**. Architecture of the synthetic gene circuits controlling the luxCDABE genes with and without the THS in σ⁷⁰V2TcR system (A), and in σ⁷⁰V3-2LacI system (B). **C-D**. Time course of bioluminescence production by the pσ⁷⁰V2TcR (C) and pσ⁷⁰V3-2LacI (D) expression systems; each bar represents the bioluminescence measurement after 3 hours of growth from time 0 to 18 hours. **E-F.** Difference in bioluminescence production between the ON and OFF states in pσ⁷⁰V2TcR (E) and pσ⁷⁰V3-2LacI (F) expression systems. **G-H**. Fold-change difference in the light emission of the recombinant cells containing the pσ⁷⁰V2TcR (G) and pσ⁷⁰V3-2LacI (H) expression systems vs the wild type DH10B. The cultures were induced after 3 hours of growth with aTc or IPTG accordingly. For all samples, the bioluminescence mean was normalized by the cell density (OD$_{600}$). Error bars are +/- S.D., n=3.

633
Figure 7. Comparison of the σ[70] expression systems to the toehold versions controlling the expression of the lux operon in *P.*
***putida*. A-B**. Time course of bioluminescence production by the pσ[70]V2TcR (A) and pσ[70]V3-2LacI (B) expression systems; each bar
represents the bioluminescence measurement after 3 hours of growth from time 0 to 18 hours. **C.** Difference in bioluminescence
production between the ON and OFF states in pσ[70]V2TcR and pσ[70]V3-2LacI expression systems. **D**. Fold-change difference in the
light emission of the recombinant cells containing the pσ[70]V2TcR or and pσ[70]V3-2LacI expression systems vs the wild type *P.*
putida. The cultures were induced after 3 hours of growth with aTc or IPTG accordingly. For all samples, the bioluminescence
mean was normalized by the cell density (OD_{600}). Error bars are +/- S.D., n=3.

Conclusions

The development of various bacterial chassis has garnered attention in pursuit of the goals of

next-generation industrial biotechnology (NGIB) (Yu, Wu et al. 2019). NGIB aims to utilize low-cost mixed

substrates, minimize freshwater consumption, save energy, and implement long-lasting, continuous,

intelligent processing methods (Yu, Wu et al. 2019). Despite the numerous advantages of *E. coli*, such as

its robust genetic toolbox and standardized cultivation parameters (Kent and Dixon 2020, Ruiz and Silhavy

2022), it falls short of achieving the goals of NGIB due to its sensitivity to harsh environments and limited

metabolic capabilities (De Wannemaeker, Bervoets et al. 2022, Zhang, Ottenheim et al. 2022). Alternative

hosts may be more attractive to accomplish complex tasks. For example, *P. putida* metabolic versatility

has a demonstrated ability to utilize lignin-derived compounds, sustain difficult redox reactions, tolerance

to organic solvents and stresses associated with industrial-scale cultivations, and shows a relatively high

GC-content similar to that of actinomycete-derived natural product BGCs (Nikel and de Lorenzo 2018,

Cook, Jacobson et al. 2021, Martin-Pascual, Batianis et al. 2021). The marine bacterium *V. natriegens*,

inherently halophilic, offers interesting advantages as well, such as rapid growth and a versatile

metabolism capable of utilizing a variety of substrates as carbon and energy sources (Hoff, Daniel et al.

2020, Thiele, Gutschmann et al. 2021, Thoma and Blombach 2021). Other non-model Gram-negative

bacterial chassis also demonstrate the potential to outperform *E. coli* (Calero and Nikel 2019, Ye and Chen

2021). However, each chassis necessitates the adaptation of distinct genetic tools. The σ^{70} expression

system showcased in this study, whether used individually or in combination, offers a promising solution

to the ongoing challenge of developing numerous host organisms. It provides a versatile and robust gene

expression platform with a broad host range. Consequently, proteins or BGCs of interest can be readily

cloned into each σ^{70} expression system and evaluated across various chassis, thereby expediting the

optimization process.

In this study, we selected chassis bacteria commonly used by the scientific community and,

through the straightforward incorporation of synthetic gene pathways for the biosynthesis of lycopene

and β-carotene, achieved high titers of these terpenoids using the σ^{70} expression system. Notably, no

media optimization was necessary to complete this task, highlighting a remarkable advantage offered by

the σ^{70} expression system. However, the production of terpenoids using the first generation of the σ^{70}

expression system revealed leakage in the OFF state, a feature imperceptible when using fluorescent

proteins as a reporter system. Approaches like decreasing plasmid copy number, weakening the RBS

sequences, and introducing mutations in the consensus promoter sequences have demonstrated the

ability to minimize leakage in the basal state, albeit at the cost of reducing the induced expression levels

(Calles, Goñi-Moreno et al. 2019). The strength and portability of the σ^{70} expression systems reside in the

specific nucleotide arrangement of the promoter and RBS DNA sequence (Carrillo Rincón and Farny 2023);

therefore, altering these sequences was not an option. Instead, the adaptation of the riboswitch toehold,

together with the use of a low-copy plasmid in *E. coli*, reduced the leakage of the σ^{70}V2TcR THS to non-

detectable levels and nearly abolished the basal expression levels of the σ^{70}V3-2LacI. This approach not

only amplified the dynamic range but also maintained the portability of the synthetic expression systems.

This was demonstrated in *P. putida*, where the DNA was integrated into the chromosome, and both THS

σ^{70} expression systems maintained strong induction, with the σ^{70}V2TcR THS reaching virtually zero

transcription. The second generation of the σ^{70} expression system (with THS) remains digital, with ON and

OFF state only, however, the emergence of plasmid like TULIP (TUnable Ligand Inducible Plasmid) with

controllable copy number (Joshi, Yong et al. 2022) could allow an analogue behavior of the versatile σ^{70}

expression systems.

In conclusion, the expression system developed in this study expands the genetic toolkit of various

promising Gram-negative chassis, thereby enhancing their capabilities to address fundamental biological

questions and bolster their biotechnological applications. Ultimately, the σ^{70} expression system has the

potential to facilitate the expression of biosynthetic gene clusters (BGCs) that have previously been

inaccessible or suboptimal due to the lack of appropriate gene tools or the limitations of model organisms

such as *E. coli*.

Declarations

Ethics approval and consent to participate: Not applicable.

Consent for publication: Not applicable.

Availability of data and materials: The raw datasets used and/or analyzed during the current study are

available from the corresponding author at reasonable request. All processed data generated or

analyzed during this study are included in this published article and its supplementary information files.

Competing interests: A.F.C.R and N.G.F have filed a U.S. patent application for the pσ^{70} *lac* and *tet*

promoters described in this work.

Funding: This work was supported by new faculty start-up funds from Worcester Polytechnic Institute to

N.G.F. & the Sandooq Al Watan grant PRJ-SWARD-256 to A.G.

Authors' contributions

Andres Felipe Carrillo Rincón: Conceptualization (lead); data curation (lead); formal analysis (lead);

investigation (lead); methodology (lead); resources (equal); validation (lead); visualization (lead); writing

– original draft (lead); writing – review and editing (equal).

Alexandra J. Cabral: Investigation (supporting); methodology (supporting); validation (supporting);

visualization (supporting); writing – review and editing (supporting).

Andras Gyorgy Conceptualization (supporting); funding acquisition (lead); project administration (lead);

resources (lead); supervision (supporting); writing – original draft (supporting); writing – review and

editing (supporting).

Natalie Gilks Farny: Conceptualization (supporting); funding acquisition (lead); project administration

(lead); resources (lead); supervision (lead); writing – original draft (equal); writing – review and editing

(equal).

Acknowledgements: We thank Adam Guss of Oak Ridge National Laboratory for *P. putida* strains and

plasmids; Kevin Keating, Emma Tobin, and Eric Young of WPI for assistance with UHPLC experiments and

for the gift of crtEBIY genes; Shimshon Belkin of Hebrew University of Jerusalem for the luxCDABE

operon.

References

Alper, H., K. Miyaoku and G. Stephanopoulos (2005). "Construction of lycopene-overproducing E. coli
strains by combining systematic and combinatorial gene knockout targets." <u>Nat Biotechnol</u> **23**(5): 612-
616.

Baeshen, M. N., A. M. Al-Hejin, R. S. Bora, M. M. Ahmed, H. A. Ramadan, K. S. Saini, N. A. Baeshen and E.
M. Redwan (2015). "Production of Biopharmaceuticals in E. coli: Current Scenario and Future
Perspectives." <u>J Microbiol Biotechnol</u> **25**(7): 953-962.

Becker, W., F. Wimberger and K. Zangger (2019). "Vibrio natriegens: An Alternative Expression System
for the High-Yield Production of Isotopically Labeled Proteins." <u>Biochemistry</u> **58**(25): 2799-2803.

Bie, L., M. Zhang, J. Wang, M. Fang, L. Li, H. Xu and M. Wang (2023). "Comparative Analysis of
Transcriptomic Response of Escherichia coli K-12 MG1655 to Nine Representative Classes of Antibiotics."
<u>Microbiol Spectr</u> **11**(2): e0031723.

Bohoyo-Gil, D., D. Dominguez-Valhondo, J. J. García-Parra and D. González-Gómez (2012). "UHPLC as a
suitable methodology for the analysis of carotenoids in food matrix." <u>European Food Research and</u>
<u>Technology</u> **235**(6): 1055-1061.

Calero, P. and P. I. Nikel (2019). "Chasing bacterial chassis for metabolic engineering: a perspective
review from classical to non-traditional microorganisms." <u>Microb Biotechnol</u> **12**(1): 98-124.

Calles, B., Á. Goñi-Moreno and V. de Lorenzo (2019). "Digitalizing heterologous gene expression in
Gram-negative bacteria with a portable ON/OFF module." <u>Mol Syst Biol</u> **15**(12): e8777.

Canty, L., S. Hariharan, Q. Liu, S. A. Haney and D. W. Andrews (2018). "Peak emission wavelength and
fluorescence lifetime are coupled in far-red, GFP-like fluorescent proteins." <u>PLoS One</u> **13**(11): e0208075.

Carrillo Rincón, A. F. and N. G. Farny (2023). "Unlocking the strength of inducible promoters in Gram-
negative bacteria." <u>Microb Biotechnol</u>.

Cook, T. B., T. B. Jacobson, M. V. Venkataraman, H. Hofstetter, D. Amador-Noguez, M. G. Thomas and B.
F. Pfleger (2021). "Stepwise genetic engineering of Pseudomonas putida enables robust heterologous
production of prodigiosin and glidobactin A." <u>Metab Eng</u> **67**: 112-124.

De Wannemaeker, L., I. Bervoets and M. De Mey (2022). "Unlocking the bacterial domain for industrial
biotechnology applications using universal parts and tools." <u>Biotechnol Adv</u> **60**: 108028.

Ellis, G. A., T. Tschirhart, J. Spangler, S. A. Walper, I. L. Medintz and G. J. Vora (2019). "Exploiting the
Feedstock Flexibility of the Emergent Synthetic Biology Chassis Vibrio natriegens for Engineered Natural
Product Production." <u>Mar Drugs</u> **17**(12).

Elmore, J. R., G. N. Dexter, H. Baldino, J. D. Huenemann, R. Francis, G. L. t. Peabody, J. Martinez-Baird, L.
A. Riley, T. Simmons, D. Coleman-Derr, A. M. Guss and R. G. Egbert (2023). "High-throughput genetic
engineering of nonmodel and undomesticated bacteria via iterative site-specific genome integration."
<u>Sci Adv</u> **9**(10): eade1285.

Elmore, J. R., A. Furches, G. N. Wolff, K. Gorday and A. M. Guss (2017). "Development of a high efficiency integration system and promoter library for rapid modification of Pseudomonas putida KT2440." Metab Eng Commun **5**: 1-8.

Gauttam, R., C. Desiderato, L. Jung, A. Shah and B. J. Eikmanns (2019). "A step forward: Compatible and dual-inducible expression vectors for gene co-expression in Corynebacterium glutamicum." Plasmid **101**: 20-27.

Gauttam, R., A. Mukhopadhyay, B. A. Simmons and S. W. Singer (2021). "Development of dual-inducible duet-expression vectors for tunable gene expression control and CRISPR interference-based gene repression in Pseudomonas putida KT2440." Microb Biotechnol **14**(6): 2659-2678.

Gauttam, R., A. Mukhopadhyay and S. W. Singer (2020). "Construction of a novel dual-inducible duet-expression system for gene (over)expression in Pseudomonas putida." Plasmid **110**: 102514.

Gemperlein, K., M. Hoffmann, L. Huo, P. Pilak, L. Petzke, R. Müller and S. C. Wenzel (2017). "Synthetic biology approaches to establish a heterologous production system for coronatines." Metab Eng **44**: 213-222.

Gemperlein, K., G. Zipf, H. S. Bernauer, R. Müller and S. C. Wenzel (2016). "Metabolic engineering of Pseudomonas putida for production of docosahexaenoic acid based on a myxobacterial PUFA synthase." Metab Eng **33**: 98-108.

Gomez-Escribano, J. P., S. Alt and M. J. Bibb (2016). "Next Generation Sequencing of Actinobacteria for the Discovery of Novel Natural Products." Mar Drugs **14**(4).

Gong, T., R. Liu, Y. Che, X. Xu, F. Zhao, H. Yu, C. Song, Y. Liu and C. Yang (2016). "Engineering Pseudomonas putida KT2440 for simultaneous degradation of carbofuran and chlorpyrifos." Microb Biotechnol **9**(6): 792-800.

Green, A. A., P. A. Silver, J. J. Collins and P. Yin (2014). "Toehold switches: de-novo-designed regulators of gene expression." Cell **159**(4): 925-939.

Gyorgy, A., J. I. Jiménez, J. Yazbek, H. H. Huang, H. Chung, R. Weiss and D. Del Vecchio (2015). "Isocost Lines Describe the Cellular Economy of Genetic Circuits." Biophys J **109**(3): 639-646.

Hakkila, K., M. Maksimow, M. Karp and M. Virta (2002). "Reporter genes lucFF, luxCDABE, gfp, and dsred have different characteristics in whole-cell bacterial sensors." Anal Biochem **301**(2): 235-242.

Hernandez-Arranz, S., J. Perez-Gil, D. Marshall-Sabey and M. Rodriguez-Concepcion (2019). "Engineering Pseudomonas putida for isoprenoid production by manipulating endogenous and shunt pathways supplying precursors." Microb Cell Fact **18**(1): 152.

Hershfield, V., H. W. Boyer, C. Yanofsky, M. A. Lovett and D. R. Helinski (1974). "Plasmid ColEl as a molecular vehicle for cloning and amplification of DNA." Proc Natl Acad Sci U S A **71**(9): 3455-3459.

Hoff, J., B. Daniel, D. Stukenberg, B. W. Thuronyi, T. Waldminghaus and G. Fritz (2020). "Vibrio natriegens: an ultrafast-growing marine bacterium as emerging synthetic biology chassis." Environ Microbiol **22**(10): 4394-4408.

Hwang, Y., S. G. Kim, S. Jang, J. Kim and G. Y. Jung (2021). "Signal amplification and optimization of
riboswitch-based hybrid inputs by modular and titratable toehold switches." J Biol Eng **15**(1): 11.

İncir, İ. and Ö. Kaplan (2024). "Escherichia coli as a versatile cell factory: Advances and challenges in
recombinant protein production." Protein Expr Purif **219**: 106463.

Jensen, P. R. (2016). "Natural Products and the Gene Cluster Revolution." Trends Microbiol **24**(12): 968-
977.

Joshi, S. H., C. Yong and A. Gyorgy (2022). "Inducible plasmid copy number control for synthetic biology
in commonly used E. coli strains." Nat Commun **13**(1): 6691.

Kadjo, A. E. and A. S. Eustáquio (2023). "Bacterial natural product discovery by heterologous
expression." Journal of Industrial Microbiology and Biotechnology **50**(1).

Karnjanawipagul, P., W. Nittayanuntawech, P. Rojsanga and L. Suntornsuk (2010). "Analysis of β-
carotene in carrot by spectrophotometry." Mahidol University Journal of Pharmaceutical Science **37**(8).

Katz, L. and R. H. Baltz (2016). "Natural product discovery: past, present, and future." J Ind Microbiol
Biotechnol **43**(2-3): 155-176.

Kelly, C. L., G. M. Taylor, A. Šatkutė, L. Dekker and J. T. Heap (2019). "Transcriptional Terminators Allow
Leak-Free Chromosomal Integration of Genetic Constructs in Cyanobacteria." Microorganisms **7**(8).

Kent, R. and N. Dixon (2020). "Contemporary Tools for Regulating Gene Expression in Bacteria." Trends
Biotechnol **38**(3): 316-333.

Krishnamurthi, V. R., Niyonshuti, II, J. Chen and Y. Wang (2021). "A new analysis method for evaluating
bacterial growth with microplate readers." PLoS One **16**(1): e0245205.

Loeschcke, A. and S. Thies (2015). "Pseudomonas putida-a versatile host for the production of natural
products." Appl Microbiol Biotechnol **99**(15): 6197-6214.

Lutz, R. and H. Bujard (1997). "Independent and tight regulation of transcriptional units in Escherichia
coli via the LacR/O, the TetR/O and AraC/I1-I2 regulatory elements." Nucleic Acids Res **25**(6): 1203-1210.

Martin-Pascual, M., C. Batianis, L. Bruinsma, E. Asin-Garcia, L. Garcia-Morales, R. A. Weusthuis, R. van
Kranenburg and V. A. P. Martins Dos Santos (2021). "A navigation guide of synthetic biology tools for
Pseudomonas putida." Biotechnol Adv **49**: 107732.

McNerney, M. P. and M. P. Styczynski (2017). "Precise control of lycopene production to enable a fast-
responding, minimal-equipment biosensor." Metab Eng **43**(Pt A): 46-53.

Meyer, A. J., T. H. Segall-Shapiro, E. Glassey, J. Zhang and C. A. Voigt (2019). "Escherichia coli
"Marionette" strains with 12 highly optimized small-molecule sensors." Nat Chem Biol **15**(2): 196-204.

Miguez, A.M., M.P. McNerney, and M. P. Styczynski (2018). "Metabolomics analysis of the toxic effects
of the production of lycopene and its precursors." Front Microbiol, **9**, p.324115.

Monici, M. (2005). "Cell and tissue autofluorescence research and diagnostic applications." Biotechnol
Annu Rev **11**: 227-256.

Nakata, P. A. (2017). "Construction of pDUO: A bicistronic shuttle vector series for dual expression of
recombinant proteins." Plasmid **89**: 16-21.

Nikel, P. I. and V. de Lorenzo (2018). "Pseudomonas putida as a functional chassis for industrial
biocatalysis: From native biochemistry to trans-metabolism." Metab Eng **50**: 142-155.

Öztürk, S., B. G. Ergün and P. Çalık (2017). "Double promoter expression systems for recombinant
protein production by industrial microorganisms." Appl Microbiol Biotechnol **101**(20): 7459-7475.

Ringel, M. T. and T. Brüser (2018). "The biosynthesis of pyoverdines." Microb Cell **5**(10): 424-437.

Romanuka, J., H. van den Bulke, R. Kaptein, R. Boelens and G. E. Folkers (2009). "Novel strategies to
overcome expression problems encountered with toxic proteins: application to the production of Lac
repressor proteins for NMR studies." Protein Expr Purif **67**(2): 104-112.

Ruiz, N. and T. J. Silhavy (2022). "How Escherichia coli Became the Flagship Bacterium of Molecular
Biology." J Bacteriol **204**(9): e0023022.

Schuster, L. A. and C. R. Reisch (2022). "Plasmids for Controlled and Tunable High-Level Expression in E.
coli." Appl Environ Microbiol **88**(22): e0093922.

Snijder, H. J. and J. Hakulinen (2016). "Membrane Protein Production in E. coli for Applications in Drug
Discovery." Adv Exp Med Biol **896**: 59-77.

Snoeck, S., C. Guidi and M. De Mey (2024). ""Metabolic burden" explained: stress symptoms and its
related responses induced by (over)expression of (heterologous) proteins in Escherichia coli." Microb
Cell Fact **23**(1): 96.

Stetz, M. A., M. V. Carter and A. J. Wand (2016). "Optimized expression and purification of biophysical
quantities of Lac repressor and Lac repressor regulatory domain." Protein Expr Purif **123**: 75-82.

Thiele, I., B. Gutschmann, L. Aulich, M. Girard, P. Neubauer and S. L. Riedel (2021). "High-cell-density
fed-batch cultivations of Vibrio natriegens." Biotechnol Lett **43**(9): 1723-1733.

Thoma, F. and B. Blombach (2021). "Metabolic engineering of Vibrio natriegens." Essays Biochem **65**(2):
381-392.

Tolia, N. H. and L. Joshua-Tor (2006). "Strategies for protein coexpression in Escherichia coli." Nat
Methods **3**(1): 55-64.

Tschirhart, T., V. Shukla, E. E. Kelly, Z. Schultzhaus, E. NewRingeisen, J. S. Erickson, Z. Wang, W. Garcia, E.
Curl, R. G. Egbert, E. Yeung and G. J. Vora (2019). "Synthetic Biology Tools for the Fast-Growing Marine
Bacterium Vibrio natriegens." ACS Synth Biol **8**(9): 2069-2079.

Wang, H., J. Li and M. C. Jewett (2018). "Development of a Pseudomonas putida cell-free protein
synthesis platform for rapid screening of gene regulatory elements." Synth Biol (Oxf) **3**(1): ysy003.

Wang, T. and F. C. Simmel (2022). "Riboswitch-inspired toehold riboregulators for gene regulation in
Escherichia coli." <u>Nucleic Acids Res</u> **50**(8): 4784-4798.

Wang, Z., T. Tschirhart, Z. Schultzhaus, E. E. Kelly, A. Chen, E. Oh, O. Nag, E. R. Glaser, E. Kim, P. F. Lloyd,
P. T. Charles, W. Li, D. Leary, J. Compton, D. A. Phillips, A. Dhinojwala, G. F. Payne and G. J. Vora (2020).
"Melanin Produced by the Fast-Growing Marine Bacterium Vibrio natriegens through Heterologous
Biosynthesis: Characterization and Application." <u>Appl Environ Microbiol</u> **86**(5).

Wu, C., Y. Yin, L. Zhu, Y. Zhang and Y. Z. Li (2022). "Metagenomic sequencing-driven multidisciplinary
approaches to shed light on the untapped microbial natural products." <u>Drug Discov Today</u> **27**(3): 730-
742.

Wu, G., Q. Yan, J. A. Jones, Y. J. Tang, S. S. Fong and M. A. G. Koffas (2016). "Metabolic Burden:
Cornerstones in Synthetic Biology and Metabolic Engineering Applications." <u>Trends Biotechnol</u> **34**(8):
652-664.

Xu, J., X. Xu, Q. Xu, Z. Zhang, L. Jiang and H. Huang (2018). "Efficient production of lycopene by
engineered E. coli strains harboring different types of plasmids." <u>Bioprocess Biosyst Eng</u> **41**(4): 489-499.

Ye, J. W. and G. Q. Chen (2021). "Halomonas as a chassis." <u>Essays Biochem</u> **65**(2): 393-403.

Yoon, S. H., H. M. Park, J. E. Kim, S. H. Lee, M. S. Choi, J. Y. Kim, D. K. Oh, J. D. Keasling and S. W. Kim
(2007). "Increased beta-carotene production in recombinant Escherichia coli harboring an engineered
isoprenoid precursor pathway with mevalonate addition." <u>Biotechnol Prog</u> **23**(3): 599-605.

Yu, L. P., F. Q. Wu and G. Q. Chen (2019). "Next-Generation Industrial Biotechnology-Transforming the
Current Industrial Biotechnology into Competitive Processes." <u>Biotechnol J</u> **14**(9): e1800437.

Zhang, C., L. Liu, L. Teng, J. Chen, J. Liu, J. Li, G. Du and J. Chen (2012). "Metabolic engineering of
Escherichia coli BL21 for biosynthesis of heparosan, a bioengineered heparin precursor." <u>Metab Eng</u>
14(5): 521-527.

Zhang, C., C. Ottenheim, M. Weingarten and L. Ji (2022). "Microbial Utilization of Next-Generation
Feedstocks for the Biomanufacturing of Value-Added Chemicals and Food Ingredients." <u>Front Bioeng</u>
<u>Biotechnol</u> **10**: 874612.

887