Assessing the effect of experimental evolution under combined thermal-

nutritional stress on larval thermotolerance and thermal plasticity in *Drosophila*

melanogaster

Keywords: experimental evolution, thermal tolerance, climate change, hardening,

heat, cold, acute shock, viability, phenotypic plasticity

Abstract

Animals commonly face combinations of thermal and nutritional stress in nature,

which will intensify under climate change. While genetic adaptation is necessary to

buffer long-term stress, it's unclear whether adaptation to combined stress can occur

without compromising viability and thermal plasticity. We tested larval

thermotolerance and thermal plasticity in *Drosophila melanogaster* selected under

different temperatures (18°C, 25°C, and 28°C) and diets (standard, diluted, and low-

protein:high-carbohydrate [P:C]). Basal larval cold tolerance was affected by both

protein concentration and temperature; larvae evolved higher basal cold tolerance on

the diluted and low P:C diets at 18°C and 28°C. Hardening increased cold tolerance for

most lines, except those selected at 18°C and 28°C on low P:C diets and at 25°C on

standard diets. Basal larval heat tolerance was affected by selection temperature;

selection at 25°C increased heat tolerance. An interaction between selection

temperature, selection diet, and hardening treatment affected larval heat tolerance;

hardening reduced heat tolerance in most lines, except those selected at 25°C on low

P:C diets and at 28°C on standard diets. Our results suggest that adaptation to

combined stress allows basal cold tolerance and its plasticity to co-evolve, but not

heat tolerance, highlighting ectotherm's vulnerability to long-term climate change.

Introduction

Anthropogenic climate change is increasing the frequency of climatic extremes (Thomas *et al.* 2004; Diffenbaugh and Field 2013; Wiens 2016). Combined shifts in atmospheric CO_2, precipitation, and temperature (Van Der Putten *et al.* 2010; Pecl *et al.* 2017; Masson-Delmotte *et al.* 2018) will alter plant growth, nutritional quality (Kreuzwieser and Gessler 2010; Lukac *et al.* 2010; Jin *et al.* 2019), and abundance (Van Der Putten *et al.* 2010; Pecl *et al.* 2017). These changes will likely impose increasing thermal and nutritional stress on organisms, both directly and indirectly through changes in microbial communities associated with plant decomposition (Bowes 1993; Richardson *et al.* 2002; Dwyer *et al.* 2007; DaMatta *et al.* 2010; Rosenblatt and Schmitz 2016). Ectothermic herbivores and frugivores, which rely on ambient temperature to maintain physiological performance and function (Hawlena and Schmitz 2010; Walters *et al.* 2012), will be particularly vulnerable to increased thermal and nutritional stress.

Organisms primarily respond to environmental stress by two means, phenotypic plasticity and genetic adaptation (Stillwell *et al.* 2007; Urban *et al.* 2014). Phenotypic plasticity, in which a single genotype produces different phenotypes depending on the environment, is often the initial response to environmental change (Huey *et al.* 2002; Urban *et al.* 2014; Seebacher *et al.* 2015; Sgrò *et al.* 2016). Plasticity allows for immediate (within-generation) responses to environmental change (Padilla and Savedo 2013; Sgrò *et al.* 2016) by shifting behaviours, physiologies, and morphologies that may contribute to population persistence (Chevin *et al.* 2010; Frankino *et al.* 2019; Noble *et al.* 2019), although it may not always be adaptive (Ghalambor *et al.* 2007; Leonard and Lancaster 2020). Nonetheless, an increasingly large body of work is focused on

understanding plastic responses to a range of stressors, particularly temperature and

nutrition, to better predict population persistence under climate change. Many of these

studies have focussed on plastic shifts in thermotolerance (van Heerwaarden *et al*.

2016; van Heerwaarden and Kellermann 2020; Rodrigues and Beldade 2020) since the

ability to withstand thermal extremes is a significant predictor of climate sensitivity

(Hoffmann *et al*. 2013; Noh *et al*. 2017; Kellermann and van Heerwaarden 2019; Ma *et*

al. 2021).

Plasticity in response to temperature and nutrition individually has been widely

reported, particularly in insects (Lee, 1989; Andersen *et al*. 2010; Sgrò *et al*. 2016;

Mitchell *et al*. 2017; Ma *et al*. 2021). For example, diets with higher protein increased

adult heat tolerance in insects (Andersen *et al*. 2010), while higher dietary

carbohydrates facilitate the synthesis of cold-resistant compounds, which in turn

enhance cold tolerance (Hazel 1995; Kostal and Simek 1998; Andersen *et al*. 2010;

Colinet *et al*. 2013).

Thermal plasticity can arise through acclimation (Rako and Hoffmann 2006; Andersen

et al. 2010; Kingsolver *et al*. 2016), which involves long-term exposure of days or weeks

to temperatures above or below the optimal temperatures but still within the normal

viable range of an organism. Acclimation to higher developmental temperatures often

increases basal heat tolerance (Kingsolver *et al*. 2016; Kellermann *et al*. 2017; Pumhan

et al. 2020; Jerbi-Elayed *et al*. 2021; Willot *et al*. 2021), while acclimation to lower

developmental temperatures can increase basal cold tolerance (Rako and Hoffmann

2006; Scharf *et al*. 2015; Noh *et al*. 2017).

Hardening, the short-term exposure of minutes or hours to sublethal temperatures, can

also induce plastic shifts in thermotolerance (Hoffmann *et al.* 2003; Bowler 2005;

Loeschcke and Sørensen 2005; Teets *et al.* 2020; Ma *et al.* 2021). In insects, high- and

low- temperature hardening treatments can improve subsequent heat (Chen *et al.*

1991; Sejerkilde *et al.* 2003; Bubliy *et al.* 2012; Manenti *et al.* 2015; Kalra *et al.* 2017) and

cold tolerance (Czajka and Lee 1990; Chen *et al.* 1991; Kelty and Lee 2001; Sejerkilde *et

al.* 2003; Jensen *et al.* 2007; Jakobs *et al.* 2015; Teets *et al.* 2020), respectively.

Plastic shifts in response to combinations of thermal and nutritional stress have been

documented, but to date, such studies have primarily focussed on life history traits

rather than thermal tolerance (Polak *et al.* 2004; Bubliy *et al.* 2012; Kutz *et al.* 2019;

Chakraborty *et al.* 2020). Interestingly, these studies suggest that plastic responses to

combinations of thermal and nutritional stress may actually be non-adaptive. For

instance, larvae of *Tenebrio molitor* and *Drosophila melanogaster* developed at higher

temperatures in combination with suboptimal diets, had decreased survival (Rho and

Lee 2017; Kutz *et al.* 2019).

Despite widespread evidence of thermal plasticity, studies suggest that plasticity is

unlikely to fully mitigate the negative consequences of ongoing temperature increases.

For example, plasticity was found to have a limited effect on increasing the upper

thermal limit of *D. melanogaster* (van Heerwaarden *et al.* 2016). There is also some

evidence suggesting that insects with higher basal thermotolerance may be less plastic

(Nyamukondiwa *et al.* 2011; Gerken *et al.* 2015; van Heerwaarden *et al.* 2016;

Kellermann *et al.* 2017; Noh *et al.* 2017; Kellermann and Sgrò 2018), implying a potential

trade-off between plasticity and basal thermotolerance (but see van Heerwaarden *et al.*

2024). Thus, in the context of ongoing climate change, genetic adaptation, while

occurring over longer timescales, will be key to the long-term persistence of

populations (Potvin and Tousignant 1996; Gienapp *et al.* 2008; Lande 2009).

Reliable predictions concerning whether and how populations might respond to long-

term selection imposed by climate change require answers to key questions (Cavicchi

et al. 1995): Do populations harbour the genetic variation necessary for rapid

evolutionary responses to combinations of stressors? Does adaptation to specific

thermal and nutritional conditions involve genetic changes in many traits? To what

extent does adaptation affect the ability of populations to mount plastic responses to

conditions other than those selected?

Experimental studies of evolution in the laboratory (Hoffmann and Harshman 1999;

Rose *et al.* 2005) have addressed these questions with respect to genetic adaptation to

thermal (Huey *et al.* 1991; Cavicchi *et al.* 1995; Bubliy and Loeschcke 2005) and

nutritional (Kolss *et al.* 2009; Kristensen *et al.* 2011; Leftwich *et al.* 2017) stress

individually. Cavicchi *et al.* (1995) and Huey *et al.* (1991) selected adult *D.*

melanogaster at different non-extreme temperatures and observed rapid evolution in

their thermotolerance. Lines adapted to higher temperatures evolved higher heat

tolerance (Huey *et al.* 1991; Cavicchi *et al.* 1995), while those adapted to cooler

temperatures had faster development at lower temperatures, but their cold tolerance

was not assessed (Huey *et al.* 1991). Interestingly, Cavicchi *et al.* (1995) found that

thermal adaptation also resulted in shifts in thermal plasticity. Lines evolved at 18°C

showed a positive acclimation response (heat tolerance increased) to development at

25°C and 28°C, while 25°C evolved lines showed a positive acclimation response to

25°C, but a negative acclimation response (heat tolerance decreased) to 28°C. Further,

the evolved lines also showed changes in their hardening responses, with the 25°C

evolved lines having a higher hardening threshold (the temperature that induces a

hardening response) compared to the 18°C evolved lines. Notably, the 28°C evolved

lines lost the ability to mount a hardening response, and exposure to the highest

hardening temperature reduced thermal tolerance. These results suggest that adapting

to elevated temperatures may come at the cost of reduced plasticity. However, the

effect of thermal adaptation on larval thermotolerance traits was not examined.

Experimental evolution has also been used to assess adaptation to changes in diet.

Kolss *et al*. (2009) examined adaptation to nutritional stress in larvae of *D.*

melanogaster by exposing them to poor (25% diluted) or control diets throughout

development; adults were placed on the control diet every generation. Evolved lines

were tested on both poor and control diets to assess whether adaptation was

associated with any trade-offs in life-history traits and whether nutritional plasticity had

evolved. When tested on the poor diet, the selected lines had higher larval viability than

the control lines, but viability remained the same for both lines when tested on the

control diet. However, adults of lines selected on poor diets were smaller, less fecund,

and less resistant to starvation than the controls. Interestingly, there was no effect of

selection or test diet on larval heat or cold tolerance. Similarly, Kristensen *et al*. (2011)

used experimental evolution to expose *D. melanogaster* larvae to standard or protein-

enriched diets. They found that adaptation to the high-protein diet resulted in reduced

larval-adult survival, suggesting that adaptation to changes in nutrition may be limited

by trade-offs between life-history traits. However, they did not examine larval or adult

thermotolerance. Finally, Leftwich *et al*. (2017) imposed nutritional selection on larvae

of the Mediterranean fruitfly and found that lines selected on the low-calorie diet

evolved higher egg-adult survival compared to those on the standard diet after 30

generations of selection, when tested on both diets. Once again, larval

thermotolerance was not assessed.

Only one study has examined adaptation to both nutritional stress and temperature.

Bochdanovits and De Jong (2003) exposed larvae to two temperatures (17.5°C and

27.5°C) and poor (50% diluted) and control diets. Overall, adaptation to the higher

temperature resulted in reduced larval survival on both diets. The 17.5°C selected lines

evolved larger body size when also adapted to a poor diet, while the 27.5°C evolved

lines were smaller on both diets. This suggests that adaptation to combinations of

temperature and nutrition can result in trade-offs between adult size and larval survival,

and potentially other traits that were not measured. However, the consequences of

adaptation to both temperature and nutrition on thermal tolerance and its plasticity

were not assessed.

In this study, we aimed to understand how long-term adaptation to combined thermal-

nutritional stress affects larval thermotolerance and thermal plasticity. We assessed

basal and induced thermotolerance of larvae from nine *D. melanogaster* experimental

evolution lines that were evolved under combinations of three temperatures (18°C,

25°C, and 28°C) and three diets (standard, diluted, and low-protein:high-carbohydrate

[low P:C] diets) using laboratory natural selection (Alton *et al*. 2024). These conditions

were chosen to reflect variation in the nutritional context that developing larvae

experience in rotting fruit, with the diluted diet representing reduced food availability

and the low P:C diet representing reduced food quality (Markow *et al*. 1999; Kristensen

et al. 2011; Almeida de Carvalho and Mirth 2017; Silva-Soares *et al.* 2017) and to

simulate future thermal and nutritional conditions in the context of long-term climate

change (DaMatta *et al.* 2010; Gowik and Westhoff 2011; IPCC 2018). The diluted and

low P:C diets contained the same concentration of protein, and the standard and low

P:C diets contained the same caloric concentration by replacing the calories from

protein with carbohydrates. This allowed us to examine the effect of protein and

carbohydrate concentrations of the selection diets independently of each other.

Only larvae were exposed to selection to reflect their limited capacity to avoid stress by

behaviourally choosing or moving to more favourable microhabitats in the wild (Austin

and Moehring 2019). Based on previous studies examining thermal and nutritional

adaptation individually, we first predicted that adaptation to 28°C would increase larval

heat tolerance (Huey et al. 1991; Cavicchi et al. 1995), and this effect would be

strongest for lines adapted to the standard diet, high in protein (Kolss et al. 2009).

Second, we predicted that adaptation to 18°C would result in increased cold tolerance,

and that this effect would be greater for lines adapted to the low P:C diet. Finally, if

basal thermotolerance trades off with thermal plasticity, we would expect plasticity to

be reduced in any lines that had evolved higher basal thermotolerance.

Our results provide insight into the complexity of evolutionary responses to

combinations of thermal and nutritional stress, and what this means for our ability to

predict responses to ongoing climate change. Previous work in microbes reveals that

adaptation to combinations of stressors can be complex and difficult to predict (Karve

et al. 2016). Consistent with this work, our results reveal that adaptation to

combinations of thermal and nutritional stress results in both genetic and plastic shifts

in pre-adult traits that are not easily predicted by adaptation to single stressors. Finally,

they highlight that studies that address the complexity of adaptation to multiple

stressors are needed to better understand responses to ongoing environmental

change.

Methods

Fly stock – Collection and initiation of experimental evolution lines

Two hundred female *D. melanogaster* were collected in Duranbah, Queensland (28.3°

S, 153° E) in 2018 to initiate isofemale lines (Alton *et al.* 2024). Flies were returned to the

laboratory and were treated with tetracycline to remove *Wolbachia* (Min and Benzer

1997). Two generations after tetracycline treatment, five males and five virgin females

from each line were used to create a mass-bred base population. A large number of

isofemale lines were used to maximise genetic variation in the founding mass-bred

population. The base population was maintained under control conditions — at 25°C

on the standard diet with 12:12 hour light:dark cycles (see below) — across 60 bottles

each containing approximately 750–1000 flies, for two generations before the initiation

of selection. It is worth noting that, because the mass-bred population was maintained

for only two generations before the selection experiment, it was most likely not fully

adapted to laboratory conditions and was therefore not at equilibrium. This, in turn,

may have contributed to linkage disequilibrium in the base (starting) population (e.g.

Kellermann *et al.* 2015). That said, we believe that the control lines at least would have

reached equilibrium by the time of the experiments described in this study, given

laboratory adaptation occurs relatively quickly, and is expected to have stabilised by 7–

10 generations (Harshman and Hoffmann 2000; Hoffmann *et al.* 2001). Because we

used laboratory natural selection (rather than directional selection), it is also likely that

the selected lines had reached equilibrium by the time the experiments were

conducted (Sgrò and Blows 2004).

Experimental evolution regimes and selection protocol

Eggs were collected from the base population and were divided among nine combined

thermal-nutritional selection treatments with five replicate lines per treatment (Alton *et*

al. 2024). These treatments consisted of three temperatures (18°C, 25°C, and 28°C)

and three diets (standard, diluted, and low P:C) (Table S1) under a 12:12 light:dark

cycle. Only larvae were exposed to selection. All diets contained varying amounts of

potato flake, inactive yeast, agar, and dextrose, with the addition of nipagin and

propionic acid to prevent bacterial and fungal growth (Holleley *et al.* 2008) (Table S1).

The standard diet had a 1:3 protein-to-carbohydrate (P:C) ratio (yeast 36.36 g/L) and

contained 320 kcal/L; the diluted diet had a 1:3 P:C ratio (yeast 9.09 g/L) and contained

80 kcal/L; and the low P:C diet had a 1:12 P:C ratio (yeast 8.18 g/L) and contained 320

kcal/L (Table S1).

These selection conditions reflect combinations of thermal and nutritional changes

that may be expected in the future under climate change; 18°C and 25°C represent the

current average Australian winter and summer temperatures respectively (Alton *et al.*

2024), while 28°C represents a 3°C increase of the average summer temperature

projected in twenty years (Australian Academy of Science 2021). The diluted diet and

the low P:C diet simulate a reduction of protein and varying carbohydrates in plant

composition projected under climate change (DaMatta *et al.* 2010), and span the range

of nutritional contexts experienced by developing larvae in rotting fruit (Matavelli *et al.*

2015; Silva-Soares *et al.* 2017). The 25°C standard diet selection regime was used as a

control and a reference point for further analysis.

At the start of each generation of selection, each selection replicate line was

established with three bottles containing 200–250 eggs on 70 mL of selection diets to

equalize density across all selection regimes. Animals were allowed to develop on their

respective diet and temperatures until adult eclosion. We avoided inadvertent selection

on development time by collecting all eclosing adults over a 5-day period across

selection lines (all adults had emerged over a 5-day period, even those lines that took

the longest to develop). Adults from the same replicate line were mixed and separated

into two bottles containing the standard diet at 25°C for three days to allow mating. On

the fourth day, adults were transferred to 250 mL egg-laying chambers containing 11

mL of egg-laying medium, the standard diet with double the concentration of agar of the

standard fly diet and a layer of autoclaved yeast for acclimation. We doubled the

concentration of agar to prevent females from embedding their eggs into the food,

making egg collection easier. After the 24-hour acclimation, fresh medium was

provided for egg-laying overnight. The eggs were collected the next morning to establish

the next generation as described above. This means that adults were between 5–10

days old at the time of egg collection, and we may therefore have inadvertently selected

for early life fecundity.

At the time of the experiment described below, generations of selection were as

follows: 61 for the 18°C standard diet, 51 for the 18°C diluted and low P:C diets, 89 for

the 25°C standard diet, 75 for the 25°C diluted and low P:C diets, 100 for the 28°C

standard diet, 81 for the 28°C diluted diet, and 84 for the 28°C low P:C diet. Due to

logistical limitations, three replicate lines were randomly selected from the five set up

for each selection regime to perform the experiment described below. Prior to the

experiment, all lines were placed in common garden conditions at 25°C on a standard

diet for at least two generations to minimise parental effects (De Villemereuil *et al*.

2016).

Plastic and evolved shifts in larval thermotolerance

Experimental set up

After two generations of common garden rearing, adults were transferred to laying

plates containing egg-laying medium and a surface layer of autoclaved yeast to induce

oviposition. Laying plates were placed at 25°C for 14 hours to allow egg-laying. Eggs

were collected from laying plates and transferred to vials containing 7.2 mL of standard

diet, at a density of twenty eggs per vial (Agnew *et al*. 2002). Five replicate vials were set

up for each replicate line per selection regime. First instar larvae (hatched 24 hours

after egg transfer) were exposed to the hardening and acute stress treatments. Vials

were maintained at 25°C under 12:12 hour light:dark cycles throughout the experiment,

except when exposing larvae to the hardening and acute stress treatments described

below. This common garden experimental design enabled us to investigate the effects

of genetic adaptation to combined thermal-nutritional stress on larval thermotolerance

and thermal plasticity.

Assessing induced (plastic) and basal thermotolerance

Larval cold plasticity was assessed by subjecting 24-hour-old first instar larvae, reared from embryos at 25°C on a standard diet, to a two-hour hardening treatment in a 0°C water bath, followed by a 24-hour recovery period at 25°C. After recovery, these 48-hour-old larvae were exposed to a four-hour acute cold shock at 0°C in a water bath and then returned to 25°C under 12:12 hour light: dark cycles to complete development. Basal cold tolerance was assessed by exposing 48-hour-old larvae, reared from embryos at 25°C on a standard diet, to a four-hour acute cold shock at 0°C in a water bath, without prior hardening treatment. The larvae were then allowed to complete development at 25°C under 12:12 hour light:dark cycles. 48-hour-old larvae were exposed to the acute cold shock treatment to equalise the stage at which larvae were exposed to the acute stress across the hardened and basal treatments.

Larval heat plasticity was assessed by subjecting 24-hour-old first instar larvae, again reared from embryos at 25°C on a standard diet, to a one-hour hardening treatment in a 35°C water bath, followed by a 24-hour recovery period at 25°C. After recovery, these 48-hour-old larvae were exposed to an acute heat shock of 39°C for 30 minutes. Those larvae were then allowed to develop at 25°C under 12:12 hour light:dark cycles until adult eclosion. Basal heat tolerance was assessed by exposing 48-hour-old larvae, reared from embryos on the standard diet at 25°C, to an acute heat shock treatment of 39°C for 30 minutes without any hardening treatment. They were then allowed to complete development at 25°C under 12:12 hour light:dark cycles.

A control treatment was also included, where vials contained larvae that were not

exposed to any heat or cold stress treatments and were maintained on control food at

25°C under 12:12 hour light:dark cycles until adult eclosion.

The temperatures and durations for both heat and cold shock hardening and acute

treatments were determined by pilot experiments (details in supplementary materials).

Throughout these treatments, all vials were temporarily sealed with plastic caps and

parafilm to prevent water leakage. After adult eclosion, larval-adult viability was

subsequently measured as the ratio of the number of emerged adult flies to the initial

egg number deposited in the vials.

Statistical analyses

All data analyses and visualisations were performed in R (R Core Team 2024). The

larval-adult viability data from all treatments were fitted with generalised linear mixed-

effects models with a binomial distribution using the *lme4* package (Bates *et al*. 2015).

All model weights were set to the number of eggs deposited per vial (20 eggs).

To test for the effects of long-term selection under combined thermal-nutritional stress

on larval-adult viability under control (25°C standard diet) conditions, selection

treatments (combinations of selection temperature and diet) were included as a fixed

effect, while block (the day that eggs were picked to set up the experiment) and

replicate line were included as non-nested (crossed) random effects to account for

variation between blocks or variation among replicate lines; these random effects were

included to improve model accuracy but were not tested for statistical significance

(Bates *et al*. 2015).

To test for the effects of selection on larval basal thermotolerance and thermal

plasticity, selection temperature, selection diet, hardening treatment (hardened and

non-hardened), and their three-way interaction were included as fixed effects. Block

and replicate line were treated as non-nested (crossed) random effects to account for

any effect of block or variation among replicate lines; these random effects were not

tested for statistical significance (Bates *et al*. 2015). Interactions between the random

and fixed effects were also not tested because they were not core to our question.

Significance of the fixed effects was assessed using ANOVA using the *car* package (Fox

et al. 2001), which uses the Wald Chi-square test to generate p-values for the fixed

effects (Bolker *et al*. 2009; Rebolledo *et al*. 2023). Post hoc multiple comparisons were

performed using the *emmean* package (Lenth 2023) to test for significant differences in

larval-adult viability across levels of each factor. To quantify the amount of variation in

the data explained by our models, we estimated conditional R^2 values which take both

fixed and random effects in the model into account (Nakagawa *et al*. 2017) using the

MuMIn package (Nakagawa and Schielzeth 2013), since traditional measures of effect

size are not applicable to generalised linear mixed models (Nakagawa *et al*. 2017).

Results

Our aim was to understand the effects of long-term adaptation to combined thermal-

nutritional stress on larval-adult viability, thermotolerance, and thermal plasticity. To

address this, we first asked whether adaptation to combinations of three temperatures

(18°C, 25°C, and 28°C) and three diets (standard, diluted and low P:C) resulted in

evolved shifts in viability under control (25°C, standard diet) conditions. We then

assessed basal and induced (plasticity in) cold and heat tolerance of *D. melanogaster*

larvae from the evolved lines.

Larval-adult viability under control conditions

We first explored whether there was evidence for any response to selection in larval-

adult viability under control (non-stress) conditions. We compared larval-adult viability

from all selection lines to our control 25°C standard diet line, and found that selection

treatments had no effect on larval-adult viability (conditional R^2 = 0.14, Table 1, Figure

1). This implies that adaptation to combinations of temperature and diet did not incur

costs in the form of reductions in viability under control conditions.

Basal cold tolerance

We then explored whether adaptation to combined thermal-nutritional stress altered

basal cold tolerance. We found a significant selection diet effect (conditional R^2 = 0.33,

Table 2) and a significant two-way interaction between selection temperature and

selection diet on basal cold tolerance (Table 2, Figure 2), meaning that any response to

selection diet further changes with the selection temperature. This interaction is

illustrated by the fact that lines selected on both diluted and low P:C diets had higher

basal cold tolerance than those selected on standard diets but only when they had

evolved at 18°C and 28°C, and not 25°C (Figure 2, Table S2). Because the diluted and

low P:C diets had the same concentration of protein, this suggests that the interaction

between protein concentration and temperature is driving these effects irrespective of

carbohydrate levels.

Plasticity in cold tolerance (cold hardening capacity)

Larval cold tolerance was significantly affected by selection diet, cold hardening

treatment, and a three-way interaction between selection temperature, selection diet,

and cold hardening treatment (conditional R^2 = 0.26, Table 3, Figure 2). This means that

the larvae showed a plastic shift in cold tolerance (a significant hardening response)

but that this plastic shift was dependent on the selection diet and selection

temperature. Hardening induced an increase in cold tolerance in larvae from most

selection treatments, but not for those selected on the low P:C diet at 18°C and 28°C

and those selected on the standard diet at 25°C (Figure 2, Table S3). This means that

selection on diluted diet increased plasticity in cold tolerance in general, while

selection on low P:C diet increased plasticity only at the optimal temperature, and

selection on standard diet (higher protein) increased plasticity in cold tolerance only at

non-optimal temperatures.

Basal heat tolerance

We then explored the effects of long-term adaptation to combined thermal-nutritional

stress on basal heat tolerance. Basal heat tolerance was significantly affected by the

selection temperature, but not selection diet (conditional R^2 = 0.12, Table 4, Figure 3).

The 25°C selection lines had higher basal heat tolerance than the 28°C, but no

difference was found in basal heat tolerance between the 18°C and 25°C or between

the 18°C and 28°C selection lines (Figure S1, Table S4). This suggests that long-term

selection at higher temperatures does not necessarily lead to an increase in basal heat

tolerance, contrary to previous studies (Cavicchi *et al.* 1995).

Plasticity in heat tolerance (heat hardening capacity)

Lastly, we explored the effects of long-term adaptation to combined thermal-nutritional

stress on plasticity in heat tolerance. We found a three-way interaction between

selection temperature, selection diet, and hardening treatment (conditional R^2 = 0.21,

Table 5, Table S5). This means that plastic shifts in heat tolerance depended on both

the selected diet and selected temperature.

In contrast to the positive cold hardening response described above, where hardening

increased cold resistance in most selection lines (Figure 2, Table S3), heat hardening

resulted in no change or a reduction in heat tolerance (Figure 3, Table S5), and these

effects varied across selection diet and selection temperature. Lines selected at 25°C

on low P:C diets and at 28°C on standard diets, which showed no significant response

to hardening (Figure 3, Table S5); however, heat hardening resulted in a significant

reduction in heat tolerance in all other lines (Figure 3, Table S5).

Discussion

Temperature and nutrition are amongst the most common stressors experienced by animals in nature, and this will become increasingly so with ongoing climate change (Rosenblatt and Schmitz 2016). However, assessments of climate risk have largely focused on the ability to tolerate thermal extremes (Hoffmann *et al.* 2003; Bowler 2005; Loeschcke and Sørensen 2005; Teets *et al.* 2020; Ma *et al.* 2021), rather than sensitivity to combinations of stressors that organisms will increasingly experience in nature. Genetic adaptation will play a key role in the long-term population persistence (Potvin and Tousignant 1996; Gienapp *et al.* 2008; Lande 2009). However, we know very little about the extent to which populations harbour the genetic variation needed to adapt to combined thermal-nutritional stress, and whether such adaptation may be constrained by trade-offs. We also have little insight into the consequences of adaptation to both temperature and nutrition on the ability to shift phenotypes via plasticity, and whether shifts in responses can be predicted by adaptation to a single stressor alone. We addressed these outstanding questions using lines of *D. melanogaster* whose larvae evolved under combinations of three temperatures (18°C, 25°C, and 28°C) and three diets (standard, diluted, and low P:C) (Alton *et al.* 2024), and assessed the extent to which adaptation to these stressors resulted in reduced viability or shifts in larval thermotolerance and thermal plasticity.

Temperature and nutrition interact to affect evolutionary shifts in pre-adult viability of D. melanogaster under control conditions

Previous studies have shown that populations harbour the genetic variation needed to adapt to diet (Kolss *et al.* 2009; Kristensen *et al.* 2011) and temperature (Sørensen *et al.*

2001; Kimura 2004; Sgrò *et al.* 2010; Freda *et al.* 2019) individually and in combination

(Bochdanovits and De Jong 2003), which is consistent with the current study. However,

these previous studies provide mixed and context-dependent evidence for trade-offs

associated with adaptation. For example, adaptation to changes in nutrition alone can

result in reduced (Kristensen *et al.* 2011) or increased (Leftwich *et al.* 2017) pre-adult

viability regardless of the test diet, or an increase in viability only when tested on the

evolved diet (Kolss *et al.* 2009). Similarly, thermal evolution can result in increased pre-

adult survival only when tested at the temperature at which populations evolved

(Partridge *et al.* 1994; Santos *et al.* 2006).

Such complexity is also evident when larvae are forced to adapt to both temperature

and diet simultaneously. Specifically, Bochdanovits and De Jong (2003) found that

adaptation to poor larval food in *D. melanogaster* resulted in increased viability when

lines were selected at 27.5°C but decreased viability when lines were selected at

17.5°C. In contrast, we found that adaptation to low-protein diets (the diluted and low

P:C diets) reduced larval viability compared to larvae evolved on standard food when

selected at 28°C, whereas this pattern was largely reversed in lines evolving at 25°C.

Lines evolved on different diets did not differ in their larval viability when evolved at

18°C. Our results suggest that simultaneous adaptation to nutritional stress and

elevated temperature may negatively impact population growth, at least when reared

under control (non-stress) conditions. Studies testing evolved shifts in viability and

trade-offs under a range of conditions, such as using a reciprocal design, are needed to

better understand the consequences for population growth.

Adaptation to temperature and nutrition in combination affects larval thermotolerance

and thermal plasticity

Previous studies on adult thermal adaptation (Huey *et al.* 1991; Cavicchi *et al.* 1995;

Bubliy and Loeschcke 2005) and dietary plasticity (Hazel 1995; Kostal and Simek 1998;

Andersen *et al.* 2010; Colinet *et al.* 2013) suggest that adaptation to lower temperatures

or a high-carbohydrate diet increases basal cold tolerance. We predicted that lines

exposed to 18°C and the low P:C diet (high carbohydrate) would evolve higher basal

cold tolerance compared to lines from other selection temperatures and diets,

although this assumption ignores complex interactions between temperature and diet,

and is based on studies focussed largely on adult traits. Surprisingly, lines selected on

poor diets, both diluted and low P:C, evolved higher basal cold tolerance compared to

lines selected on a standard diet, but only when they were selected at the non-optimal

temperatures of 18°C and 28°C. This contradicts previous single nutritional stressor

studies (Hazel 1995; Kostal and Simek 1998; Andersen *et al.* 2010; Colinet *et al.* 2013)

that suggest a high-carbohydrate diet increases cold tolerance, except that Kolss et al.

(2009) were consistent with our findings and suggested that adaptation to poor (25%

diluted) food at 25°C had no effect on basal cold tolerance. We found that cold

tolerance may be enhanced when populations are forced to adapt to combinations of

non-optimal thermal and low-protein nutritional conditions, as the diluted and low P:C

diets were protein-matched. This further emphasises that evolutionary responses to

combined stressors cannot be predicted by single-stressor studies alone. More studies

are needed to better understand the effects of a broader range of temperature and

nutritional contexts on basal cold tolerance and the underlying mechanisms.

While some studies suggest trade-offs between basal cold tolerance and cold plasticity

(Bubliy and Loeschcke 2005; Nyamukondiwa *et al*. 2011; Noh *et al*. 2017), we found that

most experimental evolution lines showed a positive cold hardening response—cold

tolerance increased with hardening. This suggests that basal cold tolerance and

plasticity in cold tolerance may have evolved together, depending on the environmental

conditions to which populations are exposed. Further studies using different rearing

conditions are recommended to further assess the trade-offs between basal cold

tolerance and cold plasticity.

Adaptation to elevated temperatures (Cavicchi *et al*. 1995; Bettencourt *et al*. 1999) and

plastic responses to high-protein diets (Andersen *et al*. 2010) have been shown to

increase adult heat tolerance. We therefore predicted that selection at 28°C on the

standard (high-protein) diet would increase larval heat tolerance. However, we found

that larvae selected at 25°C had the highest average basal heat tolerance. This

discrepancy between our results (a significant effect of selection temperature but not

diet) and previous studies (Andersen *et al*. 2010, which reported a significant effect of

selection diet) may be due to the fact that previous studies focused on adult, not larval,

thermotolerance, and insect life stages can differ in their thermal sensitivity (Kingsolver

and Buckley 2020). For instance, thermal stress tolerance in *D. buzzatii* was either

independent or weakly correlated across life stages (Sarup *et al*. 2006). Similarly,

different life stages of *D. melanogaster* also vary in their basal heat tolerance (Freda *et*

al. 2019; Moghadam *et al*. 2019), although the precise mechanisms controlling this

variation remain unclear (Bowler and Terblanche 2008).

We also found that plasticity in heat tolerance became maladaptive after adaptation to

most of the temperature-diet selection treatments. Specifically, the hardening

treatment seemed to impose further heat stress on larvae and reduced, rather than

increased, their survival and heat tolerance. Our results are partly consistent with

previous work by Cavicchi *et al*. (1995), who found increased heat tolerance after

hardening in lines adapted to 18°C and 25°C, while lines adapted to 28°C lost the ability

to mount a hardening response. Additionally, the 28°C lines showed increased

sensitivity to higher hardening temperatures (43°C), leading to a decrease in their

hardened heat tolerance (Cavicchi *et al*. 1995). In contrast, we found that none of the

evolved lines showed a significant positive hardening response; all significant

hardening responses reduced larval heat tolerance.

Our results could be explained by plasticity in heat tolerance reaching a physiological

limit. We found very weak evidence for evolved shifts in basal heat tolerance,

suggesting that the starting population may have had limited genetic variation for larval

heat tolerance (Loeschcke and Krebs 1996; Lansing *et al*. 2000) and its ability to evolve

plasticity. Limited responses to selection on larval heat tolerance also suggest that a

trade-off between basal and hardened plasticity (Gunderson and Stillman 2015; van

Heerwaarden *et al*. 2016; van Heerwaarden and Kellermann 2020) is unlikely to explain

our findings.

Another explanation could be that the threshold for inducing a heat hardening response

might have shifted down as a consequence of long-term adaptation to combined

thermal-nutritional stress in the present study (Cavicchi *et al*. 1995; van Heerwaarden

et al. 2024). Cavicchi *et al*. (1995) found that lines adapted to 18°C had a lower

threshold for inducing a hardening response compared to those adapted to 25°C. In

contrast, they found that lines adapted to 28°C appeared to lose the capacity to mount

a hardening response at lower hardening temperatures and exhibited a negative

hardening response at the highest temperature. This is in part consistent with our

results.

Plasticity of larval heat tolerance has been shown to vary across assay temperature and

duration of exposure, and is species-specific (Chen *et al*. 1991; Moghadam *et al*. 2019;

van Heerwaarden *et al*. 2024). Chen *et al*. (1991) demonstrated that the duration of

hardening treatment significantly affected adult emergence of *Sarcophaga crassipalpis*

after a heat shock; adult emergence was maximised with 2 hours of hardening at 40°C,

but only achieved 50% with a 1-hour hardening treatment. Moghadam et al., (2019)

further showed that heat hardening capacity depends on the temperature of acute heat

stress. Hardening treatments had a negative effect on subsequent heat tolerance in 3[rd]

instar larvae of *D. melanogaster* when the acute stress temperature exceeded 40°C,

but not when it was below 40°C. Our choice of temperature and duration for hardening

and acute heat stress treatments might have been too stressful for the selected larvae.

In our pilot study (see supplementary materials), larvae selected on the standard diet

across different selection temperatures showed positive hardening responses for heat

tolerance (Figure S3, albeit the effect was small). This means that positive hardening

responses may be observed if a lower acute stress temperature was used. Future

studies assessing evolutionary responses of thermotolerance and thermal plasticity

should explore a broader range of temperatures, nutritional contexts, hardening

treatments/durations under a reciprocal design, and across different life stages to

better understand the complexity of adaptation to environmental change.

Finally, we acknowledge that the mass-bred population was maintained for only two

generations before the selection experiment was initiated, meaning it was likely not

fully adapted to the laboratory conditions, and therefore not at equilibrium (Kellermann

et al. 2015). This raises the possibility that linkage disequilibrium persisted in our lines

during the first 4–5 generations of selection, which may have influenced the results.

Nonetheless, we believe this study profvides important insight into the complexities of

adaptation to combinations of stressors.

In conclusion, we found that long-term adaptation to low-protein diets and elevated

temperature simultaneously resulted in reduced larval viability, at least under control

conditions. Further, we observed unpredictable shifts in larval basal thermotolerance

and its plasticity as a result of adaptation to combined thermal-nutritional stress, which

could not be predicted from studies of thermal or nutritional adaptation individually.

We found that basal cold tolerance and plasticity in cold tolerance may co-evolve.

However, the evolved lines seemed to lose the ability to increase heat tolerance via

positive heat hardening responses, and largely displayed reductions in heat tolerance

following hardening. Taken together, our results suggest that being forced to adapt to

shifts in both the thermal and nutritional environments simultaneously may increase

the vulnerability of ectotherms to ongoing climate change.

Results – table and graphs

Table 1. Results of multiple comparisons on the effects of selection treatment on larval-adult viability under control (no stress) conditions. Significance is defined as a p-value less than 0.05.

Contrast: Selection treatment	Estimate	SE	Z ratio	P-value
18°C standard diet – 25°C standard diet	-0.030	0.142	-0.211	1.000
18°C diluted diet – 25°C standard diet	0.041	0.143	0.285	1.000
18°C low P:C diet – 25°C standard diet	0.230	0.144	1.585	0.813
25°C standard diet – 25°C diluted diet	-0.357	0.148	-2.414	0.276
25°C standard diet – 25°C low P:C diet	-0.194	0.145	-1.342	0.919
25°C standard diet – 28°C standard diet	-0.337	0.148	-2.280	0.354
25°C standard diet – 28°C diluted diet	-0.056	0.143	-0.390	1.000
25°C standard diet – 28°C low P:C diet	0.048	0.141	0.337	1.000

Table 2. Results of analysis of variance testing for effects of selection temperature, selection diet, and their interaction on larval basal (non-hardened) cold tolerance. Significance is defined as a p-value less than 0.05.

Response: Larval-adult viability	Chi-square	Df	P-value
selection temperature	1.125	2	0.121
selection diet	12.367	2	0.002
selection temperature: selection diet	11.922	4	0.018

Table 3. Results of analysis of variance testing for effects of selection temperature, selection diet, cold hardening treatment, and their interaction on larval cold tolerance. Significance is defined as a p-value less than 0.05.

Response: Larval-adult viability	Chi-square	Df	P-value

	Chi-square	Df	P-value
selection temperature	4.677	2	0.096
selection diet	15.226	2	<0.001
hardening treatment	70.606	1	<0.001
selection temperature: selection diet	6.969	4	0.138
selection temperature: hardening treatment	1.948	2	0.378
selection diet: hardening treatment	1.498	2	0.473
selection temperature: selection diet: hardening treatment	16.389	4	0.003

578 **Table 4**. Results of analysis of variance testing for effects of selection temperature,

579 selection diet, and their interaction on larval basal (non-hardened) heat tolerance.

580 Significance is defined as a p-value less than 0.05.

Response: Larval-adult viability	Chi-square	Df	P-value
selection temperature	6.142	2	0.046
selection diet	2.038	2	0.361
selection temperature: selection diet	1.079	4	0.898

582 **Table 5**. Results of analysis of variance testing for effects of selection temperature,

583 selection diet, heat hardening treatment, and their interaction on larval heat tolerance.

584 Significance is defined as a p-value less than 0.05.

Response: Larval-adult viability	Chi-square	Df	P-value
selection temperature	7.072	2	0.029
selection diet	2.557	2	0.278
hardening treatment	25.277	1	<0.001

selection temperature: selection diet	31.516	4	<0.001
selection temperature: hardening treatment	1.824	2	0.402
selection diet: hardening treatment	9.289	2	0.010
selection temperature: selection diet: hardening treatment	21.281	4	<0.001

Figure 1. Mean larval-adult viability of experimental evolution lines tested under control treatments. Control treatment involved larvae developing on standard (control) food at 25°C under 12:12 light:dark cycles until adult eclosion. Error bars indicate ± 1 standard error.

Figure 2. Mean larval-adult viability of basal (non-hardened) and hardened (2 h at 0°C)

experimental evolution lines after exposure to an acute cold shock treatment of 4 h at

0°C. Error bars indicate ± 1 standard error.

Figure 3. Mean larval-adult viability of basal (non-hardened) and hardened (1 h at 35°C)

experimental evolution lines after exposure to an acute heat shock treatment of 0.5 h at

39°C. Error bars indicate ± 1 standard error.

References

Agnew P, Hide M, Sidobre C, Michalakis Y (2002) 'A minimalist approach to the effects

of density-dependent competition on insect life-history traits' *Ecological*

Entomology **27**, 396–402. doi:10.1046/j.1365-2311.2002.00430.x

Almeida de Carvalho MJ, Mirth CK (2017) 'Food intake and food choice are altered by

the developmental transition at critical weight in Drosophila melanogaster' *Animal*

Behaviour **126**, 195–208. doi:10.1016/j.anbehav.2017.02.005

Alton LA, Kutz T, Bywater CL, Lombardi E, Cockerell FE, Layh S, Winwood-Smith H,

Arnold PA, Beaman JE, Walter GM, Monro K, Mirth CK, Sgrò CM, White CR (2024)

'Temperature and nutrition do not interact to shape the evolution of metabolic

rate' *Philosophical Transactions of the Royal Society B: Biological Sciences* **379**,.

doi:10.1098/rstb.2022.0484

Andersen LH, Kristensen TN, Loeschcke V, Toft S, Mayntz D (2010) 'Protein and

carbohydrate composition of larval food affects tolerance to thermal stress and

desiccation in adult Drosophila melanogaster' *Journal of Insect Physiology* **56**,

336–340. doi:10.1016/j.jinsphys.2009.11.006

Austin CJ, Moehring AJ (2019) 'Local thermal adaptation detected during multiple life

stages across populations of Drosophila melanogaster' *Journal of Evolutionary*

Biology **32**, 1342–1351. doi:10.1111/jeb.13530

Australian Academy of Science (2021) 'The risks to Australia of a 3°C warmer world.'

Available at https://www.science.org.au/supporting-science/science-policy-and-

analysis/reports-and-publications/risks-australia-three-degrees-c-warmer-world

Bates D, Mächler M, Bolker B, Walker S (2015) 'Fitting linear mixed-effects models using

lme4' *Journal of Statistical Software* **67**, 1–54. doi:10.18637/jss.v067.i01

Bettencourt BR, Feder ME, Cavicchi S (1999) 'Experimental evolution of Hsp70

expression and thermotolerance in Drosophila melanogaster' *Evolution* **53**, 484–

492. doi:10.1111/j.1558-5646.1999.tb03783.x

Bochdanovits Z, De Jong G (2003) 'Experimental evolution in Drosophila melanogaster:

Interaction of temperature and food quality selection regimes' *Evolution* **57**, 1829–

1836. doi:10.1111/j.0014-3820.2003.tb00590.x

Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, White JSS

(2009) 'Generalized linear mixed models: a practical guide for ecology and

evolution' *Trends in Ecology and Evolution* **24**, 127–135.

doi:10.1016/j.tree.2008.10.008

Bowes G (1993) 'Facing the inevitable: Plants and increasing atmospheric CO_2' *Annual*

Review of Plant Physiology and Plant Molecular Biology **44**, 309–332.

doi:10.1146/annurev.pp.44.060193.001521

Bowler K (2005) 'Acclimation, heat shock and hardening' *Journal of Thermal Biology* **30**,

125–130. doi:10.1016/j.jtherbio.2004.09.001

Bowler K, Terblanche JS (2008) 'Insect thermal tolerance: What is the role of ontogeny,

ageing and senescence?' *Biological Reviews* **83**, 339–355. doi:10.1111/j.1469-

185X.2008.00046.x

Bubliy OA, Kristensen TN, Kellermann V, Loeschcke V (2012) 'Plastic responses to four

environmental stresses and cross-resistance in a laboratory population of

Drosophila melanogaster' *Functional Ecology* **26**, 245–253. doi:10.1111/j.1365-2435.2011.01928.x

Bubliy OA, Loeschcke V (2005) 'Correlated responses to selection for stress resistance and longevity in a laboratory population of Drosophila melanogaster' *Journal of Evolutionary Biology* **18**, 789–803. doi:10.1111/j.1420-9101.2005.00928.x

Cavicchi S, Guerra D, Torre V La, Huey RB (1995) 'Chromosomal analysis of heat-shock tolerance in Drosophila melanogaster evolving at different temperatures in the laboratory' *Evolution* **49**, 676. doi:10.2307/2410321

Chakraborty A, Sgrò CM, Mirth CK (2020) 'Does local adaptation along a latitudinal cline shape plastic responses to combined thermal and nutritional stress?' *Evolution* **74**, 2073–2087. doi:10.1111/evo.14065

Chen C, Lee RE, Denlinger DL (1991) 'Cold shock and heat shock: a comparison of the protection generated by brief pretreatment at less severe temperatures' *Physiological Entomology* **16**, 19–26. doi:10.1111/j.1365-3032.1991.tb00539.x

Chevin L-M, Lande R, Mace GM (2010) 'Adaptation, plasticity, and extinction in a changing environment: Towards a predictive theory' *PLoS Biology* **8**, e1000357. doi:10.1371/journal.pbio.1000357

Colinet H, Larvor V, Bical R, Renault D (2013) 'Dietary sugars affect cold tolerance of Drosophila melanogaster' *Metabolomics* **9**, 608–622. doi:10.1007/s11306-012-0471-z

Czajka MC, Lee RE (1990) 'A rapid cold-hardening response protecting against cold shock injury in Drosophila melanogaster' *Journal of Experimental Biology* **148**, 245–

254. doi:10.1242/jeb.148.1.245

DaMatta FM, Grandis A, Arenque BC, Buckeridge MS (2010) 'Impacts of climate

changes on crop physiology and food quality' *Food Research International* **43**,

1814–1823. doi:10.1016/j.foodres.2009.11.001

Diffenbaugh NS, Field CB (2013) 'Changes in ecologically critical terrestrial climate

conditions' *Science* **341**, 486–492. doi:10.1126/science.1237123

Dwyer SA, Ghannoum O, Nicotra A, Von Caemmerer S (2007) 'High temperature

acclimation of C4 photosynthesis is linked to changes in photosynthetic

biochemistry' *Plant, Cell and Environment* **30**, 53–66. doi:10.1111/j.1365-

3040.2006.01605.x

Fox J, Weisberg S, Price B (2001) 'car: Companion to Applied Regression' *CRAN Contrib.*

Packag. doi:10.32614/CRAN.package.car

Frankino WA, Bakota E, Dworkin I, Wilkinson GS, Wolf JB, Shingleton AW (2019)

'Individual cryptic scaling relationships and the evolution of animal form'

Integrative and Comparative Biology **59**, 1411–1428. doi:10.1093/icb/icz135

Freda PJ, Ali ZM, Heter N, Ragland GJ, Morgan TJ (2019) 'Stage-specific genotype-by-

environment interactions for cold and heat hardiness in Drosophila melanogaster'

Heredity **123**, 479–491. doi:10.1038/s41437-019-0236-9

Gerken AR, Eller OC, Hahn DA, Morgan TJ (2015) 'Constraints, independence, and

evolution of thermal plasticity: Probing genetic architecture of long-and short-term

thermal acclimation' *Proceedings of the National Academy of Sciences of the*

United States of America **112**, 4399–4404. doi:10.1073/pnas.1503456112

Ghalambor CK, McKay JK, Carroll SP, Reznick DN (2007) 'Adaptive versus non-adaptive

phenotypic plasticity and the potential for contemporary adaptation in new

environments' *Functional Ecology* **21**, 394–407. doi:10.1111/j.1365-

2435.2007.01283.x

Gienapp P, Teplitsky C, Alho JS, Mills JA, Merilä J (2008) 'Climate change and evolution:

Disentangling environmental and genetic responses' *Molecular Ecology* **17**, 167–

178. doi:10.1111/j.1365-294X.2007.03413.x

Gowik U, Westhoff P (2011) 'The Path from C3 to C4 photosynthesis' *Plant Physiology*

155, 56–63. doi:10.1104/pp.110.165308

Gunderson AR, Stillman JH (2015) 'Plasticity in thermal tolerance has limited potential

to buffer ectotherms from global warming' *Proceedings of the Royal Society B:*

Biological Sciences **282**, 20150401. doi:10.1098/rspb.2015.0401

Harshman LG, Hoffmann AA (2000) 'Laboratory selection experiments using

Drosophila: What do they really tell us?' *Trends in Ecology and Evolution* **15**, 32–36.

doi:10.1016/S0169-5347(99)01756-5

Hawlena D, Schmitz OJ (2010) 'Herbivore physiological response to predation risk and

implications for ecosystem nutrient dynamics' *Proceedings of the National*

Academy of Sciences of the United States of America **107**, 15503–15507.

doi:10.1073/pnas.1009300107

Hazel JR (1995) 'Thermal adaptation in biological membranes: Is homeoviscous

adaptation the explanation?' *Annual Review of Physiology* **57**, 19–42.

doi:10.1146/annurev.ph.57.030195.000315

van Heerwaarden B, Kellermann V (2020) 'Does plasticity trade off with basal heat

tolerance?' *Trends in Ecology and Evolution* **35**, 874–885.

doi:10.1016/j.tree.2020.05.006

van Heerwaarden B, Kellermann V, Sgrò CM (2016) 'Limited scope for plasticity to

increase upper thermal limits' *Functional Ecology* **30**, 1947–1956.

doi:10.1111/1365-2435.12687

van Heerwaarden B, Sgrò C, Kellermann VM (2024) 'Threshold shifts and developmental

temperature impact trade-offs between tolerance and plasticity' *Proceedings of*

the Royal Society B: Biological Sciences **291**, 20232700.

doi:10.1098/rspb.2023.2700

Hoffmann AA, Chown SL, Clusella-Trullas S (2013) 'Upper thermal limits in terrestrial

ectotherms: How constrained are they?' *Functional Ecology* **27**, 934–949.

doi:10.1111/j.1365-2435.2012.02036.x

Hoffmann AA, Hallas R, Sinclair C, Partridge L (2001) 'Rapid loss of stress resistance in

drosophila melanogaster under adaptation to laboratory culture' *Evolution* **55**,

436–438. doi:10.1111/j.0014-3820.2001.tb01305.x

Hoffmann AA, Harshman LG (1999) 'Desiccation and starvation resistance in

Drosophila: Patterns of variation at the species, population and intrapopulation

levels' *Heredity* **83**, 637–643. doi:10.1046/j.1365-2540.1999.00649.x

Hoffmann AA, Sørensen JG, Loeschcke V (2003) 'Adaptation of Drosophila to

temperature extremes: Bringing together quantitative and molecular approaches'

Journal of Thermal Biology **28**, 175–216. doi:10.1016/S0306-4565(02)00057-8

Holleley CE, Hocking AD, Schubert TL, Whitehead MR (2008) 'Control of Penicillium

roqueforti (Thom) infection in cultures of Drosophila melanogaster (Meigen)

(Diptera: Drosophilidae)' *Australian Journal of Entomology* **47**, 149–152.

doi:10.1111/j.1440-6055.2007.00630.x

Huey RB, Carlson M, Crozier L, Frazier M, Hamilton H, Harley C, Hoang A, Kingsolver JG

(2002) 'Plants versus animals: Do they deal with stress in different ways?'

Integrative and Comparative Biology **42**, 415–423. doi:10.1093/icb/42.3.415

Huey RB, Patridge L, Fowler K (1991) 'Thermal sensitivity of Drosophila melanogaster

responds rapidly to laboratory natural selection' *Evolution* **45**, 751–756.

doi:10.2307/2409925

Jakobs R, Gariepy TD, Sinclair BJ (2015) 'Adult plasticity of cold tolerance in a

continental-temperate population of Drosophila suzukii' *Journal of Insect*

Physiology **79**, 1–9. doi:10.1016/j.jinsphys.2015.05.003

Jensen D, Overgaard J, Sørensen JG (2007) 'The influence of developmental stage on

cold shock resistance and ability to cold-harden in Drosophila melanogaster'

Journal of Insect Physiology **53**, 179–186. doi:10.1016/j.jinsphys.2006.11.008

Jerbi-Elayed M, Foray V, Tougeron K, Grissa-Lebdi K, Hance T (2021) 'Developmental

temperature affects life-history traits and heat tolerance in the aphid parasitoid

aphidius colemani' *Insects* **12**, 1–12. doi:10.3390/insects12100852

Jin J, Armstrong R, Tang C (2019) 'Impact of elevated CO_2 on grain nutrient

concentration varies with crops and soils – A long-term FACE study' *Science of the*

Total Environment **651**, 2641–2647. doi:10.1016/j.scitotenv.2018.10.170

753 Kalra B, Tamang AM, Parkash R (2017) 'Cross-tolerance effects due to adult heat

754  hardening, desiccation and starvation acclimation of tropical drosophilid-

755  Zaprionus indianus' *Comparative Biochemistry and Physiology -Part A : Molecular*

756  *and Integrative Physiology* **209**, 65–73. doi:10.1016/j.cbpa.2017.04.014

757 Karve SM, Bhave D, Nevgi D, Dey S (2016) 'Escherichia coli populations adapt to

758  complex, unpredictable fluctuations by minimizing trade-offs across

759  environments' *Journal of Evolutionary Biology* **29**, 2545–2555.

760  doi:10.1111/jeb.12972

761 Kellermann V, van Heerwaarden B (2019) 'Terrestrial insects and climate change:

762  adaptive responses in key traits' *Physiological Entomology* **44**, 99–115.

763  doi:10.1111/phen.12282

764 Kellermann V, van Heerwaarden B, Sgrò CM (2017) 'How important is thermal history?

765  Evidence for lasting effects of developmental temperature on upper thermal limits

766  in Drosophila melanogaster' *Proceedings of the Royal Society B: Biological*

767  *Sciences* **284**, 14–16. doi:10.1098/rspb.2017.0447

768 Kellermann V, Hoffmann AA, Kristensen TN, Moghadam NN, Loeschcke V (2015)

769  'Experimental evolution under fluctuating thermal conditions does not reproduce

770  patterns of adaptive clinal differentiation in Drosophila melanogaster' *American*

771  *Naturalist* **186**, 582–593. doi:10.1086/683252

772 Kellermann V, Sgrò CM (2018) 'Evidence for lower plasticity in CTMAX at warmer

773  developmental temperatures' *Journal of Evolutionary Biology* **31**, 1300–1312.

774  doi:10.1111/jeb.13303

Kelty JD, Lee RE (2001) 'Rapid cold-hardening of Drosophila melanogaster (Diptera:

Drosophilidae) during ecologically based thermoperiodic cycles' *Journal of*

Experimental Biology **204**, 1659–1666. doi:10.1242/jeb.204.9.1659

Kimura MT (2004) 'Cold and heat tolerance of drosophilid flies with reference to their

latitudinal distributions' *Oecologia* **140**, 442–449. doi:10.1007/s00442-004-1605-4

Kingsolver JG, Buckley LB (2020) 'Ontogenetic variation in thermal sensitivity shapes

insect ecological responses to climate change' *Current Opinion in Insect Science*

41, 17–24. doi:10.1016/j.cois.2020.05.005

Kingsolver JG, MacLean HJ, Goddin SB, Augustine KE (2016) 'Plasticity of upper thermal

limits to acute and chronic temperature variation in Manduca sexta larvae' *Journal*

of Experimental Biology **219**, 1290–1294. doi:10.1242/jeb.138321

Kolss M, Vijendravarma RK, Schwaller G, Kawecki TJ (2009) 'Life-history consequences

of adaptation to larval nutritional stress in drosophila' *Evolution* **63**, 2389–2401.

doi:10.1111/j.1558-5646.2009.00718.x

Kostal V, Simek P (1998) 'Changes in fatty acid composition of phospholipids and

triacylglycerols after cold-acclimation of an aestivating insect prepupa' *Journal of*

Comparative Physiology - B Biochemical, Systemic, and Environmental Physiology

168, 453–460. doi:10.1007/s003600050165

Kreuzwieser J, Gessler A (2010) 'Global climate change and tree nutrition: Influence of

water availability' *Tree Physiology* **30**, 1221–1234. doi:10.1093/treephys/tpq055

Kristensen TN, Overgaard J, Loeschcke V, Mayntz D (2011) 'Dietary protein content

affects evolution for body size, body fat and viability in Drosophila melanogaster'

Biology Letters **7**, 269–272. doi:10.1098/rsbl.2010.0872

Kutz TC, Sgrò CM, Mirth CK (2019) 'Interacting with change: Diet mediates how larvae

respond to their thermal environment' (C Williams, Ed.) *Functional Ecology* **33**,

1940–1951. doi:10.1111/1365-2435.13414

Lande R (2009) 'Adaptation to an extraordinary environment by evolution of phenotypic

plasticity and genetic assimilation' *Journal of Evolutionary Biology* **22**, 1435–1446.

doi:10.1111/j.1420-9101.2009.01754.x

Lansing E, Justesen J, Loeschcke V (2000) 'Variation in the expression of Hsp70, the

major heat-shock protein, and thermotolerance in larval and adult selection lines

of Drosophila melanogaster' *Journal of Thermal Biology* **25**, 443–450.

doi:10.1016/S0306-4565(00)00008-5

Lee, RE (1989) 'Insect cold-hardiness: To freeze or not to freeze' *BioScience* **39**, 308–

313. doi:10.2307/1311113

Leftwich PT, Nash WJ, Friend LA, Chapman T (2017) 'Adaptation to divergent larval diets

in the medfly, Ceratitis capitata' *Evolution* **71**, 289–303. doi:10.1111/evo.13113

Lenth R V. (2023) 'emmeans: Estimated Maarginal Means, aka Least-Square Means. R

package version 1.8.8.' Available at https://cran.r-project.org/package=emmeans

Leonard AM, Lancaster LT (2020) 'Maladaptive plasticity facilitates evolution of thermal

tolerance during an experimental range shift' *BMC Evolutionary Biology* **20**, 1–11.

doi:10.1186/s12862-020-1589-7

Loeschcke V, Krebs RA (1996) 'Selection for heat-shock resistance in larval and in adult

Drosophila buzzatii: Comparing direct and indirect responses' *Evolution* **50**, 2354–

2359. doi:10.1111/j.1558-5646.1996.tb03623.x

Loeschcke V, Sørensen JG (2005) 'Acclimation, heat shock and hardening - A response

from evolutionary biology' *Journal of Thermal Biology* **30**, 255–257.

doi:10.1016/j.jtherbio.2004.12.005

Lukac M, Calfapietra C, Lagomarsino A, Loreto F (2010) 'Global climate change and tree

nutrition: Effects of elevated CO2 and temperature' *Tree Physiology* **30**, 1209–

1220. doi:10.1093/treephys/tpq040

Ma C-S, Ma G, Pincebourde S (2021) 'Survive a warming climate: Insect responses to

extreme high temperatures' *Annual Review of Entomology* **66**, 163–184.

doi:10.1146/annurev-ento-041520-074454

Manenti T, Loeschcke V, Moghadam NN, Sørensen JG (2015) 'Phenotypic plasticity is

not affected by experimental evolution in constant, predictable or unpredictable

fluctuating thermal environments' *Journal of Evolutionary Biology* **28**, 2078–2087.

doi:10.1111/jeb.12735

Markow TA, Raphael B, Dobberfuhl D, Breitmeyer CM, Elser JJ, Pfeiler E (1999)

'Elemental stoichiometry of Drosophila and their hosts' *Functional Ecology* **13**, 78–

84. doi:10.1046/j.1365-2435.1999.00285.x

Masson-Delmotte V, Zhai P, Pörtner H-O, Roberts D, Skea J, Shukla PR, Pirani A,

Moufouma-Okia W, Péan C, Pidcock R, Connors S, Matthews JBR, Chen Y, Zhou X,

Gomis MI, Lonnoy E, Maycock T, Tignor M, Waterfield T (2018) 'Intergovernmental

panel on climate change-global warming of 1.5 degrees Celcius.' Available at

https://report.ipcc.ch/sr15/pdf/sr15_spm_final.pdf%0Ahttp://www.ipcc.ch/report

/sr15/

Matavelli C, Carvalho MJA, Martins NE, Mirth CK (2015) 'Differences in larval nutritional

requirements and female oviposition preference reflect the order of fruit

colonization of Zaprionus indianus and Drosophila simulans' *Journal of Insect*

Physiology **82**, 66–74. doi:10.1016/j.jinsphys.2015.09.003

Min KT, Benzer S (1997) 'Wolbachia, normally a symbiont of Drosophila, can be virulent,

causing degeneration and early death' *Proceedings of the National Academy of*

Sciences of the United States of America **94**, 10792–10796.

doi:10.1073/pnas.94.20.10792

Mitchell KA, Boardman L, Clusella-Trullas S, Terblanche JS (2017) 'Effects of nutrient

and water restriction on thermal tolerance: A test of mechanisms and hypotheses'

Comparative Biochemistry and Physiology -Part A : Molecular and Integrative

Physiology **212**, 15–23. doi:10.1016/j.cbpa.2017.06.019

Moghadam NN, Ketola T, Pertoldi C, Bahrndorff S, Kristensen TN (2019) 'Heat hardening

capacity in Drosophila melanogaster is life stage-specific and juveniles show the

highest plasticity' *Biology Letters* **15**, 20180628. doi:10.1098/rsbl.2018.0628

Nakagawa S, Johnson PCD, Schielzeth H (2017) 'The coefficient of determination R2

and intra-class correlation coefficient from generalized linear mixed-effects

models revisited and expanded' *Journal of the Royal Society Interface* **14**,.

doi:10.1098/rsif.2017.0213

Nakagawa S, Schielzeth H (2013) 'A general and simple method for obtaining R2 from

generalized linear mixed-effects models' *Methods in Ecology and Evolution* **4**, 133–

863 142. doi:10.1111/j.2041-210x.2012.00261.x

864 Noble DWA, Radersma R, Uller T (2019) 'Plastic responses to novel environments are

865  biased towards phenotype dimensions with high additive genetic variation'

866  *Proceedings of the National Academy of Sciences of the United States of America*

867  **116**, 13452–13461. doi:10.1073/pnas.1821066116

868 Noh S, Everman ER, Berger CM, Morgan TJ (2017) 'Seasonal variation in basal and

869  plastic cold tolerance: Adaptation is influenced by both long- and short-term

870  phenotypic plasticity' *Ecology and Evolution* **7**, 5248–5257. doi:10.1002/ece3.3112

871 Nyamukondiwa C, Terblanche JS, Marshall KE, Sinclair BJ (2011) 'Basal cold but not

872  heat tolerance constrains plasticity among Drosophila species (Diptera:

873  Drosophilidae)' *Journal of Evolutionary Biology* **24**, 1927–1938. doi:10.1111/j.1420-

874  9101.2011.02324.x

875 Padilla DK, Savedo MM (2013) 'A systematic review of phenotypic plasticity in marine

876  invertebrate and plant systems' *Advances in Marine Biology* **65**, 67–94.

877  doi:10.1016/B978-0-12-410498-3.00002-1

878 Partridge L, Barrie B, Fowler K, French V (1994) 'Thermal evolution of pre-adult life

879  history traits in Drosophila melanogaster' *Journal of Evolutionary Biology* **7**, 645–

880  663. doi:10.1046/j.1420-9101.1994.7060645.x

881 Pecl GT, Araújo MB, Bell JD, Blanchard J, Bonebrake TC, Chen IC, Clark TD, Colwell RK,

882  Danielsen F, Evengård B, Falconi L, Ferrier S, Frusher S, Garcia RA, Griffis RB,

883  Hobday AJ, Janion-Scheepers C, Jarzyna MA, Jennings S, Lenoir J, Linnetved HI,

884  Martin VY, McCormack PC, McDonald J, Mitchell NJ, Mustonen T, Pandolfi JM,

Pettorelli N, Popova E, Robinson SA, Scheffers BR, Shaw JD, Sorte CJB, Strugnell

JM, Sunday JM, Tuanmu MN, Vergés A, Villanueva C, Wernberg T, Wapstra E,

Williams SE (2017) 'Biodiversity redistribution under climate change: Impacts on

ecosystems and human well-being' *Science* **355**, 1–9.

doi:10.1126/science.aai9214

Polak M, Kroeger DE, Cartwright IL, Ponce DeLeon C (2004) 'Genotype-specific

responses of fluctuating asymmetry and of preadult survival to the effects of lead

and temperature stress in Drosophila melanogaster' *Environmental Pollution* **127**,

145–155. doi:10.1016/S0269-7491(03)00238-0

Potvin C, Tousignant D (1996) 'Evolutionary consequences of simulated global change:

genetic adaptation or adaptive phenotypic plasticity' *Oecologia* **108**, 683–693.

doi:10.1007/BF00329043

Pumhan N, Tian M, Xu L-L, Jiang J, Liu T-X, Zhang S-Z (2020) 'Effects of heat stress and

exposure duration on survival characteristics of different developmental stages of

Propylaea japonica, a dominant aphidophagous ladybeetle in China' *Crop*

Protection **130**, 105054. doi:10.1016/j.cropro.2019.105054

Van Der Putten WH, Macel M, Visser ME (2010) 'Predicting species distribution and

abundance responses to climate change: Why it is essential to include biotic

interactions across trophic levels' *Philosophical Transactions of the Royal Society*

B: Biological Sciences **365**, 2025–2034. doi:10.1098/rstb.2010.0037

R Core Team (2024) 'R: A language and environment for statistical computing. R

Foundation for Statistical Computing' Available at https://www.r-project.org/

Rako L, Hoffmann AA (2006) 'Complexity of the cold acclimation response in

Drosophila melanogaster' *Journal of Insect Physiology* **52**, 94–104.

doi:10.1016/j.jinsphys.2005.09.007

Rebolledo AP, Sgrò CM, Monro K (2023) 'When is warmer better? Disentangling within-

and between-generation effects of thermal history on early survival' *Functional*

Ecology **37**, 2488–2499. doi:10.1111/1365-2435.14398

Rho MS, Lee KP (2017) 'Temperature-driven plasticity in nutrient use and preference in

an ectotherm' *Oecologia* **185**, 401–413. doi:10.1007/s00442-017-3959-4

Richardson SJ, Press MC, Parsons AN, Hartley SE (2002) 'How do nutrients and

warming impact on plant communities and their insect herbivores? A 9-year study

from a sub-Arctic heath' *Journal of Ecology* **90**, 544–556. doi:10.1046/j.1365-

2745.2002.00681.x

Rodrigues YK, Beldade P (2020) 'Thermal Plasticity in Insects' Response to Climate

Change and to Multifactorial Environments' *Frontiers in Ecology and Evolution* **8**,

1–12. doi:10.3389/fevo.2020.00271

Rose MR, Passananti HB, Chippindale AK, Phelan JP, Matos M, Teotónio H, Mueller LD

(2005) 'The effects of evolution are local: Evidence from experimental evolution in

Drosophila' *Integrative and Comparative Biology* **45**, 486–491.

doi:10.1093/icb/45.3.486

Rosenblatt AE, Schmitz OJ (2016) 'Climate change, nutrition, and bottom-up and top-

down food web processes' *Trends in Ecology & Evolution* **31**, 965–975.

doi:10.1016/j.tree.2016.09.009

Santos M, Brites D, Laayouni H (2006) 'Thermal evolution of pre-adult life history traits,

geometric size and shape, and developmental stability in Drosophila subobscura'

Journal of Evolutionary Biology **19**, 2006–2021. doi:10.1111/j.1420-

9101.2006.01139.x

Sarup P, Sørensen JG, Dimitrov K, Barker JSF, Loeschcke V (2006) 'Climatic adaptation

of Drosophila buzzatii populations in southeast Australia' *Heredity* **96**, 479–486.

doi:10.1038/sj.hdy.6800828

Scharf I, Galkin N, Halle S (2015) 'Disentangling the consequences of growth

temperature and adult acclimation temperature on starvation and thermal

tolerance in the red flour beetle' *Evolutionary Biology* **42**, 54–62.

doi:10.1007/s11692-014-9298-z

Seebacher F, White CR, Franklin CE (2015) 'Physiological plasticity increases resilience

of ectothermic animals to climate change' *Nature Climate Change* **5**, 61–66.

doi:10.1038/nclimate2457

Sejerkilde M, Sørensen JG, Loeschcke V (2003) 'Effects of cold- and heat hardening on

thermal resistance in Drosophila melanogaster' *Journal of Insect Physiology* **49**,

719–726. doi:10.1016/S0022-1910(03)00095-7

Sgrò CM, Blows MW (2004) 'The genetic covariance among clinal environments after

adaptation to an environmental gradient in Drosophila serrata' *Genetics* **167**,

1281–1291. doi:10.1534/genetics.103.026120

Sgrò CM, Overgaard J, Kristensen TN, Mitchell KA, Cockerell FE, Hoffmann AA (2010) 'A

comprehensive assessment of geographic variation in heat tolerance and

hardening capacity in populations of Drosophila melanogaster from Eastern

Australia' *Journal of Evolutionary Biology* **23**, 2484–2493. doi:10.1111/j.1420-

9101.2010.02110.x

Sgrò CM, Terblanche JS, Hoffmann AA (2016) 'What can plasticity contribute to Insect

responses to climate change?' *Annual Review of Entomology* **61**, 433–451.

doi:10.1146/annurev-ento-010715-023859

Silva-Soares NF, Nogueira-Alves A, Beldade P, Mirth CK (2017) 'Adaptation to new

nutritional environments: larval performance, foraging decisions, and adult

oviposition choices in Drosophila suzukii' *BMC Ecology* **17**, 21.

doi:10.1186/s12898-017-0131-2

Sørensen JG, Dahlgaard J, Loeschcke V (2001) 'Genetic variation in thermal tolerance

among natural populations of Drosophila buzzatii: Down regulation of Hsp70

expression and variation in heat stress resistance traits' *Functional Ecology* **15**,

289–296. doi:10.1046/j.1365-2435.2001.00525.x

Stillwell RC, Wallin WG, Hitchcock LJ, Fox CW (2007) 'Phenotypic plasticity in a

complex world: Interactive effects of food and temperature on fitness components

of a seed beetle' *Oecologia* **153**, 309–321. doi:10.1007/s00442-007-0748-5

Teets NM, Gantz JD, Kawarasaki Y (2020) 'Rapid cold hardening: ecological relevance,

physiological mechanisms and new perspectives' *Journal of Experimental Biology*

223, 1–12. doi:10.1242/JEB.203448

Thomas CD, Cameron A, Green RE, Bakkenes M, Beaumont LJ, Collingham YC,

Erasmus BFN, de Siqueira MF, Grainger A, Hannah L, Hughes L, Huntley B, van

Jaarsveld AS, Midgley GF, Miles L, Ortega-Huerta MA, Townsend Peterson A,

Phillips OL, Williams SE (2004) 'Extinction risk from climate change' *Nature* **427**,

145–148. doi:10.1038/nature02121

Urban MC, Richardson JL, Freidenfelds NA (2014) 'Plasticity and genetic adaptation

mediate amphibian and reptile responses to climate change' *Evolutionary*

Applications **7**, 88–103. doi:10.1111/eva.12114

De Villemereuil P, Gaggiotti OE, Mouterde M, Till-Bottraud I (2016) 'Common garden

experiments in the genomic era: New perspectives and opportunities' *Heredity*

116, 249–254. doi:10.1038/hdy.2015.93

Walters RJ, Blanckenhorn WU, Berger D (2012) 'Forecasting extinction risk of

ectotherms under climate warming: An evolutionary perspective' *Functional*

Ecology **26**, 1324–1338. doi:10.1111/j.1365-2435.2012.02045.x

Wiens JJ (2016) 'Climate-related local extinctions are already widespread among plant

and animal species' *PLoS Biology* **14**, 1–18. doi:10.1371/journal.pbio.2001104

Willot Q, Loos B, Terblanche JS (2021) 'Interactions between developmental and adult

acclimation have distinct consequences for heat tolerance and heat stress

recovery' *Journal of Experimental Biology* **224**, 1–8. doi:10.1242/jeb.242479