Inferring metabolite states from spatial transcriptomes using multiple graph neural

network

Jiaxu Luo[1], Daosheng Ai[2,3]*, Wenzhi Sun[1,3]*

[1] School of Basic Medical Sciences, Capital Medical University, Beijing 100069, China

[2] Academy for Advanced Interdisciplinary Studies, Peking University, Beijing 100871, China

[3] Chinese Institute for Brain Research, Beijing 102206, China

* To whom the correspondence should be addressed:

Wenzhi Sun, Ph.D (Lead contact)

Email: sunwenzhi@cibr.ac.cn

Daosheng Ai, Ph.D

Email: aidaosheng@cibr.ac.cn

Keywords: spatial transcriptome, single cell transcriptome, graph neural network,

metabolic flux estimation

Abstract

Metabolism serves as the pivotal interface connecting genotype and phenotype in various

contexts, such as cancer reprogramming and immune metabolic reprogramming. Compared to the

transcriptome, the development of the single-cell metabolome faces significant challenges. While

various methods exist for predicting metabolite levels from transcriptome, their efficacy remains

limited. We developed an efficient and adaptable algorithm known as Multiple Graph-based Flux

Estimation Analysis (MGFEA). MGFEA enables rapid inference from million-level single-cell

transcriptome datasets and achieves accuracy comparable to that of scFEA. Additionally, MGFEA

can detect metabolite biomarkers in different cancer bulk RNA-seq datasets. As an attempt to

integrate multi-omics dataset, MGFEA can further improve the accuracy of these inferences by

leveraging additional metabolome.

Introduction

In the intricate realm of cellular biology, common cells diligently uphold metabolic homeostasis within their internal milieu, a crucial state that ensures the proper functioning of distinct functional proteins. Notably, the diverse nature of cells entails varied modes of sustaining metabolic equilibrium. For instance, neoplastic cells exhibit metabolic reprogramming, altering their metabolic profiles to adapt to their environment [1–5]. Similarly immune cells dynamically adjust to effectively adapt and respond to their microenvironment [6–8].

Over the last decade, there has been significant advancement in single-cell transcriptomics technology, leading to the accumulation of a substantial number of precise single-cell databases [9–12]. However, the advancement of single-cell metabolomics has lagged behind that of single-cell transcriptomics largely due to inherent technical bottlenecks [13,14]. Though progress has been slow, contemporary computational tools are now capable of characterizing metabolism through transcription. Enrichment-based methods have demonstrated significant impact in the field of functional genomics research, yet they are predominantly utilized for qualitative analysis [15–18]. Constraint-based models have shown significant promise by their ability to deduce the rate of metabolic reactions without the prerequisite detection of numerous kinetic parameters [18,19]. Consequently, novel computational models have emerged for predicting flux state at the single cell resolution. These include, but are not limited to, scFBA [20], scFEA [21], Compass [22], and METAFlux [23]. Despite their innovative nature, the efficiency of these models still falls short of optimal levels.

Building upon the foundations established by scFEA and a series of constraint-based models, we introduce a novel modeling framework for inferring metabolic flux based on both metabolic

network guided gene interaction graph and spatial information graph (Fig. 1a). This framework

aims to estimate the divergences in metabolic reactions among cells by utilizing the known

gene-reaction relationships in the Genome Scale Metabolism model (GSMM) alongside

single-cell transcriptomics dataset. The code implementation and relevant dataset are available on

GitHub (https://github.com/Sunwenzhilab/MGFEA).

Fig. 1: An algorithmic framework to estimate metabolic states based on spatial and single cell

transcriptomic dataset, and computational performance measurement.

a, MGFEA algorithmic framework. VAE: variational autoencoder VGAE: variational graph

autoencoder scRNAseq: single cell transcriptome dataset spRNAseq: spatial transcriptome dataset

MALDI: Marix-Assisted Laser Desorption Ionization SMA: spatial multimodal analysis dataset. A

paired spRNAseq and MALDI dataset.

b, left: Comparison of peak memory usage between scFEA and MGFEA for different sized

datasets three repeats each point. right: Comparison of time cost between scFEA and MGFEA for

different sized datasets three repeats each point. scFEA reported out of memory error in the 4 large

datasets, so there is no data point of scFEA in the figure.

c, left: Comparison of peak memory usage between scFEA and MGFEA for different formats

of stereo-seq dataset. Right: Comparison of computation time between scFEA and MGFEA for

different formats of stereo-seq dataset. scFEA reported out of memory error in the 4 large datasets,

so there is no data point for scFEA in the figure.

Result

Overview of MGFEA framework

MGFEA pipeline combined the framework of representation learning and the metabolic flux constraint framework such as scFEA and adopts a suitable module for sparse matrix. MGFEA is a flexible framework which achieved the efficient and accurate metabolic inference for various types of datasets, such as single cell RNA seq dataset, spatial RNA seq dataset. MGFEA extracted cell embeddings from expression matrix and integrated gene co-expression information from expression matrix, spatial information from spatial RNAseq and expert knowledge from GSMM model into the gene embeddings. MGFEA used the dot product to combine the two types of embedding and used the dot product as weights to enhance the cell expression embedding. At the last layer, full connected layer transformed the enhanced cell expression embedding into the metabolic flux under the constraint of the metabolic flux loss. With the result of metabolic flux, MGFEA could infer all of the metabolites' relative stock level in the single cell.

Compared with scFEA, MGFEA demonstrated three innovations. First, our framework exhibits remarkable flexibility, featuring multiple modules that can accept inputs in various data formats, including single-cell transcriptome datasets, spatial transcriptome datasets, and spatial multimodal analysis (SMA) datasets [24]. MGFEA employed anndata format and sparse matrix module to achieve compatibility for large datasets. Second, MGFEA utilized representation learning to extract and integrate gene interaction information from expression matrix, spatial information from spatial RNAseq and expert knowledge from GSMM model. By leveraging Variational Graph Autoencoder (VGAE) or Variational Autoencoder (VAE), our framework extracts cellular embeddings from the transcriptome dataset. Additionally, another VGAE is

employed to extract inter-gene metabolic information derived from the knowledge-guided graph

of GSMM, thereby generating gene embeddings. We then computed the dot product of these two

distinct embedding types and applied a softmax transformation along the cell axis, to derive a

weight matrix. This matrix was subsequently applied to enhance the original transcriptome matrix.

Third, based on the premise which non-metabolic genes also affect metabolic flux, MGFEA used

the same stoichiometry matrix from scFEA and other GSMM models to guided fully connected

multiple layer perceptron (MLP) to transform the extracted cell embeddings into metabolic flux in

the cell.

To promote the development of metabolic prediction methods based on metabolic graph, we

also provided three types of metabolic graphs for each species: the first from scFEA, the second

from flux-estimator [42], and the third from the GSMM model Recon3D and IMM1865 [26,27].

Although flux-estimator [42] only subgraphs for user access, we integrated the majority of these

into a single comprehensive graph to ensure a fair comparison. Graphs serve as an important role

in the constraint based methods. In this article, when referencing the small graph, we will

abbreviate the names of two models. For the large graph, we will append an "-L" suffix to the

different models. Similarly, IMM1865 will be referenced to as "IMM." Subsequently, the

improved matrix serves as input, enabling the neural network to autonomously learn without fixed

gene-reaction relationships. Through unsupervised metabolic flux inference, our framework

derives metabolic flux, facilitating the characterization of metabolite imbalance levels across

different cells analogous to metabolomics analysis. Furthermore, owing to the flexibility of flux

estimation analysis method, we also provided an additional reference module to leverage the

metabolic information present in the SMA dataset thereby further optimizing the algorithm's

performance. Based on the existing contributions of scFEA [21], flux-estimator [42], Recon3D [26]

and IMM1865 [27], we also offered three types of graph in h5ad format and the reference module

can also be applied on the other constraint based methods. The above two items represent our

modest contribution to the field of constraint-based inferences methods about metabolism.

MGFEA outperformed in memory usage and time cost

Based on the data preprocess pipeline which is suitable for the sparse matrix, MGFA showed

high computational performance. Compared with scFEA, MGFEA demonstrated significant

advantages in computational performance, particularly in its adaptability to the growing scale of

single-cell datasets and its efficiency regarding memory usage and computational speed (Fig. 1b).

In the same device, scFEA pipeline reported out of memory error in the large datasets, but

MGFEA showed measurable performance. Considering the scalability of stereo-seq datasets [43],

we selected five thresholds, resulting in the generation of five different datasets sizes, each with

varying sizes and library depth of spots. According to stereo-seq datasets, MGFEA demonstrated

better performance than scFEA in different resolution of stereo-seq datasets (Fig. 1c). According

to the benchmark of the daily largest single cell RNAseq datasets, MGFEA exhibited improved

performance in terms of computational resource utilization.

Depmap dataset benchmark confirms framework efficiency and discovery of potential

metabolites biomarker

Metabolic reprogramming is the classical features of tumor. Tumor metabolic reprogramming

is the important format of tumor rearranging tumor microenvironment and resisting the

chemotherapeutic agents [44]. We validated the performance of MGFEA in the public datasets

which owns transcriptome and metabolome information of human cancer cell lines. To assess the

effect of dataset size to model accuracy, we utilized a public dataset paired with metabolomics

dataset from human cancer cell lines provided by the Cancer Dependency Map (DepMap) Project

[36,37]. We selected all overlapped metabolites among the different metabolic graphs and the

metabolomics dataset for comparable accuracy assessment across different metabolic graphs. Then

we calculated the relative mean square error between the predictions and metabolomics result. We

presented the results of the prediction from scFEA, scFEA-L, MGFEA, MGFEA-L and

MGFEA-IMM. As the dataset size increased, the performance of two flux estimation analysis

algorithms improved. The usage of flux estimation analysis algorithm on larger datasets showed

higher inference accuracy and the performance of predicted results from MGFEA consistently

outperformed that of scFEA (Fig. 2a).

As a supplemental experiments, we cultured three types of cancer cell lines: Hela, U87 and

SH-SY5Y for our in house paired transcriptome and metabolome dataset. Based on the targeted

metabolomics detection, we confirmed that for the ten detected metabolites, the predictions from

MGFEA-L and scFEA-L matched with the true concentration distribution in fumarate and Uridine

Monophosphate across three cancer cell lines targeted metabolomics detection (Fig. 2b). But both

MGFEA and scFEA show incorrect trends in other two significant differential metabolites:

beta-alanine and deoxyadenosine. Two models reported the same predictions but not matched with

the validation completely on succinate (Fig. S1c), we will discuss the phenomenon in the

discussion section.

MGFEA could be used in the discovery of tumor potential metabolic biomarkers. In an

application utilizing single-cell transcriptome dataset from gliomas sampled across different

regions [45], we observed a significant difference in lactate levels between cells in the tumor core

and those in the peripheral regions as predicted by our model (Fig. S1a). This finding is similar

with the functional magnetic resonance imaging results [46]. Leveraging the abundant

transcriptome datasets from TCGA [20], we tried to find the potential biomarker with the

prediction from MGFEA. MGFEA classified of patients as two groups based on the median

relative predicted metabolites level from MGFEA. For example, MGFEA identified four

metabolites as the potential biomarkers that could distinguish patients with poor survival outcomes

(Fig 2c). Citrulline is predicted as a biomarker associated with patients' better prognosis (Fig 2c).

Citrulline has been found as the products of nitric oxide(NO) synthase, which catalyzed the

reaction which produce NO. Similar research supported our predictions that oral administration of

L-arginine or hydroxyurea significantly increased brain tumor barrier permeability when

compared with the nontreated control rat [47]. Thereby patients whose tumor owns higher

citrulline level could be more sensitive to the chemotherapeutic agents. MGFEA predicted that

fumarate is associated with patients' bad prognosis. Fumarate has been validated that correlated

with the inhibitory function of CD8 positive T cell [48]. Similar measured results about Glutamate

can be found in a published metabolomics dataset [49] (Fig. S1b). Glutamate is found in higher

concentrations in IDH wild-type gliomas but is lower in IDH-mutant gliomas. Our predictions are

consistent with metabolomic findings, as IDH-mutant patients, who generally have better

prognosis, show lower glutamate levels. Research has indicated that glioma release glutamate to

improve their growth by utilizing its neurotoxicity [50,51]. MGFEA also predicted succinate

correlated with the patients' bad prognosis. Glioblastoma cells improved the succinate level to

accommodate the hypoxic environment which is similar with our predictions [52]. MGFEA

showed comparable accuracy with scFEA in DepMap datasets and could be used for the

identification of potential metabolites' biomarkers in TCGA and more human datasets.

Fig. 2: Comparison of prediction results from scFEA and MGFEA in DepMap dataset and

predicted biomarker in TCGA glioma dataset.

a, The relative mean square error comparison of metabolic predictions from scFEA, MGFEA,

scFEA-L, MGFEA-L and MGFEA-IMM on DepMap datasets.

b, The prediction result from metabolomics, scFEA, scFEA-L, MGFEA and MGFEA-L at

Fumarate and UMP.

c, The prognosis of TCGA glioma patients is correlated with several predicted metabolites:

citrulline, fumarate, succinate and glutamate. Red lines mean the statistics from patients with

higher predicted metabolites. Blue lines mean the statistics from patients with lower predicted

metabolites.

Fig. S1: Comparison of MGFEA and scFEA in terms of cancer metabolites imbalance level.

a, Predicted relative lactate levels of lactate in different glioma regions by MGFEA and

198 scFEA.

199  b, Metabolomics result about glutamate in IDH wild type and IDH mutant type glioma

200 samples from Wang et al.

201  c, Inhouse datasets metabolomics results and prediction results from scFEA and MGFEA

202 about succinate content in Hela, U87 and SHSY5Y cell lines.

205 **MGFEA integrated non metabolic genes and metabolomics into the whole framework**

206  With the spatial multimodal analysis (SMA) paired spatial RNAseq and Matrix-assisted laser

207 desorption/ionization (MALDI) datasets, we could validate the performance of MGFEA and

208 validate the true metabolic information integration module of MGFEA. For the SMA dataset, we

209 selected one of the slices to compare the prediction results of different models against MALDI

210 results (Fig. 3a-d). The finding indicated that MGFEA captured spatial similarity of different spots

211 and showed significant better performance when using the raw matrix as input. In both models of

212 MGFEA, imputation did not yield significant improvements in performance (Fig. 3a, c).

213 Consistent with scFEA reports, MAGIC [31] improved its performance (Fig. 3b, d). STAGATE

214 [32] which integrates spatial information demonstrated no improvement in scFEA and small

215 improvement in scFEA-L. We compared three methods which select variable genes for MGFEA

216 metabolic gene interaction graph construction: common highly variable genes (HVG) defined by

217 normalized dispersion, spatial autocorrelation metrics Moran's I (SRR) and spatial differential

218 expressed genes (SDE). By leveraging HVG, MGFEA demonstrated better performance than the

219 other methods without imputation. Although HVG methods did not demonstrate improvement

compared to the other two methods, they also did not result in decrement in performance.

Therefore, we used HVG as the default parameter of MGFEA. With respect to the relative mean

square error, MGFEA outperformed scFEA with different imputation preprocess (Fig. 3b, d).

Given the paired nature of the SMA dataset, leveraging the metabolomics results as a

reference for MGFEA predictions can improve performance (Fig. 3e). We observed improvements

in accuracy and reductions in variance when using the MALDI reference, although statistical

significance was not achieved (Fig. 3f).

Fig. 3: Comparison of prediction results from scFEA and MGFEA in SMA dataset and the

reference module which integrates metabolomics and gene expression.

a, c Relative mean square error between the prediction by MGFEA and MGFEA-L with

different variable genes selection methods, highly variable genes, spatial correlated genes and

spatial differential genes.

b, d Relative mean square error between the prediction by scFEA, MGFEA, scFEA-L,

MGFEA-L in SMA dataset with the three different imputation methods, no imputation, MAGIC

and STAGATE.

e, Comparison of dopamine prediction basic MGFEA and referenced MGFEA with raw

matrix and imputated matrix as input.

f, Statistics of reference module contribution in MGFEA relative mean square error.

Unpaired brain transcriptome and metabolome datasets confirmed efficiency of MGFEA

Leveraging the unpaired metabolome and spatial transcriptome datasets, MGFEA showed

efficient performance on differential metabolites identification. Based on the metabolomics dataset

of different brain regions from Shao et al [53] and visium sagittal spatial RNAseq dataset, we

present the results of the metabolome, the prediction from scFEA, MGFEA, MGFEA-IMM,

scFEA-L and MGFEA-L (Fig. 4a). Compared to scFEA, MGFEA showed lower relative mean

square error across all metabolic graphs. We provided several examples of prediction results (Fig.

4b). MGFEA exhibited comparable prediction accuracy to scFEA in two types of metabolic graphs

(Fig. 4c, d). Along with the significant differential metabolites identified through metabolomics,

MGFEA demonstrated greater accuracy in classifying relative differences across various brain

region (Fig. 4c). When considering the top-ranked predicted differential metabolites from each

model, MGFEA exhibited outperformed scFEA (Fig. 4d).

Fig. 4: Comparison of prediction results from scFEA and MGFEA in unpaired visium sagittal

dataset and Shao metabolomics dataset.

a, Results of relative mean square error between Shao datasets and the prediction of scFEA,

MGFEA, scFEA-L, MGFEA-L and MGFEA-IMM for the relative metabolite level of mouse

brain.

b, Detection of log fold change of residual metabolites from different brain region pairs using

metabolome, scFEA and MGFEA. COR: cortex, HYT: hypothalamus, BST: brain stem, CBL:

cerebellum, OB: olfactory bulb.

c, The ROC-AUC area of models in relative difference classification within significant

differential metabolites from metabolome.

d, The ROC-AUC area of models in relative difference classification within top differential

predicted metabolites from models.

Discussion

MGFEA is designed for fast inference on large datasets and is particularly adept at inferring metabolic states in tumor samples, leveraging the rich transcriptomic public data resource. Our pipeline improved computational performance significantly and could be applied in the analysis of large datasets even million level datasets [54,55] and the future application on the insilico perturbation gene functional analysis which needs thousands repeats.

Tumor heterogeneity is associated with the bad prognosis of patients [56]. In contrast to the metabolome, transcriptome enabled researchers to acquire single cell transcript information at an affordable price [57]. Thereby we used MGFEA in tumor samples for the discovery of biomarkers. To validate the efficiency of models, we prepared the in-housed dataset of three cancer cell lines. In the in-housed dataset, both MGFEA and scFEA predicted the correct distribution of fumarate and uridine monophosphate. Interestingly, scFEA and MGFEA reported the same prediction about succinate based on the transcriptome, but different from the targeted metabolomics detections. There are several possible reasons about the phenomena, for example, the correlation between metabolites and transcript is weak [24]. Enzyme catalyzed metabolites transformation, enzyme is translated from transcript, but the correlation between protein and transcript is even weak [58]. Although transcriptome and proteome can't work as the proxy of each other [59], the question of which more accurately represents the actual functions performed by cells, the transcriptome or the proteome, should be rigorously assessed through experimental validation from multiple aspects. Metabolome quantified the metabolites' relative level in the time points of samples collection. Although the algorithms computed the relative level of metabolites based on the key enzyme expression, the difference between the inferred and actual measurements becomes more noticeable

in the non-steady state scenario of culture media. Two models' prediction approved the succinate

of SH-SY5Y is higher than other two cell lines, the inconsistency of prediction and measurement

could also bring new assumption: succinate is very important for the proliferation of SH-SY5Y or

the related TCA cycle genes are reprogramed in SH-SY5Y. The deeper understanding may be

proposed between the different result from theoretical model and experimental observation. The

essence of the phenomenon is worthy to explore further for our metabolites-transcript consistency

understanding.

In an attempt to validate the potential of MGFEA on the further exploitation of public

transcriptome datasets, we used MGFEA to discover novel validated metabolites from TCGA

datasets and demonstrated the potential of our flexible and efficient framework. Of the four

metabolites shown in figure 2c, most of them have been found to engage the progression of tumor

[47,48,50,52].

For instance, we performed metabolite inference validation using the SMA dataset, which

seamlessly integrates histologic data from various modalities within the MGFEA framework.

Although our reference module demonstrated subtle improvement, but our attempt demonstrated

constraint-based methods or flux estimation models such as scFEA [21], compass [22],

METAFlux [23] and MGFEA which is compatible with high throughput single cell transcriptome

datasets are also suitable for the integration of multi-omics datasets consists of MALDI, spatial

RNAseq and spatial proteomics. Although the correlation between different modalities is weak,

the integration of multiple modalities is also promising to produce novel knowledge and even

novel research field in the future.

With the development of spatial metabolome technique [60–62], or the metabolites aptamer

technique [63], it may be easier for the acquirement of the metabolic and transcript information of

our interested samples, with the novel inference algorithm based on the genotype information, the

understanding of interaction of genotype and phenotype could further develop and help with the

human health.

In summary, MGFEA demonstrates the ability to make fast and accurate inferences about the

metabolic state of a sample based on its transcriptome. It provides an algorithmic framework that

can easily integrate both transcriptional and metabolic modalities from the same samples, making

it a valuable tool for multimodal data integration. The further development of MGFEA can

provide inspiration for the emergence of a mature integration framework across multi-omics fields,

such as transcriptomics, proteomics, and metabolomics.

Methods

MGFEA framework

MGFEA framework (Fig. 1a) consists of data preprocess, metabolic graph integration, cell and gene embedding extraction, embedding augmentation layer and flux transformation layer.

X, x_k : expression matrix, expression vector of kth cell

X_r : matrix of spatial metabolomics matrix from spatial multimodal analysis (SMA) datasets [24]

T^g, T^c : the extracted Eigen vector matrix in the gene and cell dimension of the expression matrix using Principal Component Analysis

u, v : cell embedding, gene embedding

G^m, G^s, A^m, A^s : metabolic network guided gene interaction graph, spatial information graph , metabolic network guided gene interaction graph adjacent matrix, spatial information graph adjacent matrix

φ_k, θ_k : cell variational graph autoencoder [25] VGAE encoder parameter, decoder parameter

φ_G, θ_G : gene VGAE encoder parameter, decoder parameter

F : Flux matrix of all cells in dataset

f : flux vector of single cell

S : stoichiometry matrix of GSMM model

GSMM model preprocess

We employed two published GSMM model, Recon3D [26] and IMM1865 [27], for MGFEA prediction of relative metabolites level. The original models have large numbers of metabolites consists of the same metabolites located in different organelles. In our modified models, we

removed duplicated metabolites and used function find_blocked_reaction from python package

cobrapy [28] to remove most blocked reactions.

MGFEA metabolic interaction graph preprocess

According the metabolic stoichiometry matrix and gene co-expression relationship from

expression matrix, we transformed all the information into a gene interaction graph. The graph

incorporated metabolic relationships from the GSMM model, along with the intrinsic gene

co-expression information in the expression matrix, as the edges between genes. For metabolic

edges, we construct edges between genes connected by reaction or metabolite. For gene

co-expression edges, we calculated the expression correlation of metabolic genes with highly

variable genes (HVG). Edge connections are established between the top k highly variable genes

and the metabolic genes with the highest correlation in their expression. To ensure the information

density of HVG is comparable to that of metabolic genes, we compute the normalized dispersion

of all metabolic genes. Then, the sum of normalized dispersion is used as the threshold value to

select the top k HVG genes whose corresponding statistics equals to that of the metabolic genes.

According to the above method, we construct a gene adjacency matrix A^m.

360
MGFEA embedding

MGFEA takes a preprocessed expression matrix X and a preprocessed GSMM model as input.

Inspired by the GLUE [29] framework, we employed two separate VGAEs to learn cell

embeddings in the spatial transcriptomic dataset and gene embeddings in metabolic networks

separately. For the cell embedding u, the input consists of the expression matrix X and spatial

coordinate information of spots A^s. The obtained latent layer embedding u serves as the

representation of different cells.

For single cell RNAseq dataset input, the training of VGAE satisfies the following loss function:

369
$$L_{cell}(\varphi_k, \theta_k) = E_{x_k \sim p_{data}(x_k)} \left[E_{u \sim q(u|x_k; \varphi_k)} log p(x_k \mid u; \theta_k) - KL\left(q(u \mid x_k; \varphi_k) \parallel p(u) \right) \right]$$

370 For spatial RNAseq dataset input, the training of VGAE satisfies the following loss function:

371
$$L_{cell}(\varphi_k, \theta_k) = E_{i,j \sim A^s} \left[E_{u \sim q(u|x_k, A^s; \varphi_k)} log p(A^s \mid u; \theta_k) - KL\left(q(u \mid x_k, A^s; \varphi_k) \parallel p(u) \right) \right]$$

372 For gene embedding v, Using the metabolic gene interaction graph as input, the VGAE learns the

373 intrinsic relationships of genes and acquires gene embeddings to represent different genes.

374 The loss function of VGAE which exported gene embeddings satisfies:

375
$$L_{gene}(\varphi_G, \theta_G) = E_{i,j \sim A^m} \left[log p(A^m \mid v, \theta_{G_m}) - KL[q(v \mid A^m, T^g, \varphi_{G_m}) \parallel p(v)] \right]$$

376 The loss of two VGAE is the latent loss of MGFEA.

377
$$Ll(\varphi_{G_s}, \theta_{G_s}, \varphi_{G_m}, \theta_{G_m}) = \frac{1}{K} \Sigma_{k=1}^{K} L_{cell} + L_{gene}$$

The two VGAEs learned the representations of the transcriptional cell states u and gene

representations v defined jointly by the metabolic network and co-expression information. The

former represents a conventional cellular state and incorporates both the expression matrix and

spatial coordinate information of spots of spatial transcriptomics. The latter means the gene

representations defined by gene interaction graph. The gene representations mean the genes'

location in metabolic space. The dot product of two representations shares the same matrix form

as the original expression matrix. The form is used for embedding enhancement.

Taking into account the inherent projection nature of the dot product, we interpret the dot product

of the two as a projection of the cellular state representation vector onto the metabolic space. For

scRNAseq, by utilizing the difference between this projection and the original transcriptional

expression matrix as a loss function, we enable VAE to rationalize the cell representations it learns.

For spRNAseq, VGAE learns to reconstruct the spatial coordinate graph and gene interaction

graph to rationalize the obtained cell embeddings and gene embeddings.

MGFEA embedding augmentation

With the above framework, we are able to generate matrix containing both metabolic gene

relationships and cell transcriptional states. Then we conducted softmax transformation between

different genes within a cell based on the generated matrix. This transformation yields a matrix

which contains genes' weights in cells. Subsequently, we performed element-wise multiplication

(Hadamard product) between weights matrix and the original expression matrix X. Through this

process, we enhance the cell specific metabolic features in the original matrix X to preserve the

transcriptional states and the augmented matrix X_a improved the process of MGFEA

flux-estimation.

$$X_a = Softmax_i(v^T u) \otimes X$$
$$F = MLP(X_a)$$

MGFEA-Flux estimation

The balance of metabolites in reaction network is influenced by both influx and efflux.

Considering that the efficiency of enzymes in metabolic networks is regulated by the regulatory

genes, the transcriptional state of the cell plays a crucial role in influencing metabolic balance [30].

Building upon this premise, we utilized the transcriptional expression matrices of metabolic genes,

along with a restricted set of highly variable genes, as input. We finally employed a Multilayer

perceptron to estimate the fluxes of all metabolic modules (reactions).

$$L_{flux} = \sum_{i}^{reactions} \sum_{j}^{metabolites} [|S \cdot f| + |f| - f + |sgn(f)| - sgn(f)]$$

$$L_{total} = \lambda_f L_{flux} + (1 - \lambda_f) L_l \left(\varphi_{G_g}, \theta_{G_g}, \varphi_{G_m}, \theta_{G_m} \right)$$

Reference based framework

We use transcriptome and metabolome paired dataset [24] to offer a reference for

spatialGraphFEA learning. We modify flux loss and add a reference loss for our model. We add

reciprocal of flux to prohibit maintained decrease of flux loss. We use metabolism quantitative

information as a reference and use a mean square error formula to forced predicted metabolism

quantification into reference result.

$$L_{reference} = \frac{1}{K} \, mse(S \cdot f - X_r)$$

$$L_{total} = \lambda_f L_{flux} + \lambda_l L_l \left(\varphi_{G_g}, \theta_{G_g}, \varphi_{G_m}, \theta_{G_m} \right) + (1 - \lambda_f - \lambda_l) L_{reference}$$

Parameters

The weights of flux loss should be confirmed based on the epoch size. We usually used 0.5 as a

default weight of flux loss. As a semi-supervised framework, MGFEA's best parameter should be

confirmed manually. When the epoch size is too big, model appears overfitting and the results

even appeared as an opposite direction to the truth, we used the weight of flux loss to make flux

loss converge as the training ends. We can also increase the relative weights of flux direction.

Dataset preprocess

Expression matrix X is loaded in h5ad format and is normalized and log transformed. We used

MAGIC [31] and STAGATE [32] for imputation. Reactions without expression in the expression

matrix are removed, and the modified reaction network was used for MGFEA reaction prediction.

We used highly variable genes detected by scanpy [33], spatial differential genes from spatialDE

[34] and spatial correlated genes by Moran's I from squidpy [35] for the gene interaction graph

construction.

Relative mean square error

In all instances where the relative mean square error was utilized, we first filtered out all

nonoverlapping metabolites between the predicted results and the truth, we then scaled the

metabolomics dataset and the metabolites' stock level output from the model, calculating the

relative mean squared error between the predicted results and the truth. For cases involving Shao

metabolomics dataset, datasets from Dependency Map (DepMap) Project [36,37] and our inhouse

dataset, since the vectors are not paired, so we first computed the mean metabolites level in

different regions before calculating the relative mean square error for different metabolites.

Receiver operating characteristic curve (ROC)-area under curve (AUC)

We transformed the correct direction between the different brain region pairs into the binary label.

Thereby we can employed ROC-AUC metrics to assess the capability on classifying correct

relative level between different brain regions of the different models. We used the true log

transformed fold change between pairs of brain region as true label and the models' predicted

mean of log transformed fold change as predicted value for the ROC-AUC calculation.

Memory usage and time consumption

We utilized python package memory-profiler(https://github.com/pythonprofilers/memory_profiler)

to measure memory usage and total time consumption of different models.

Experimental validation of Cancer Cell lines

We ordered cell lines from the vector center at Chinese institute brain research, obtaining

U-87-MG (EK-Bioscience Cat.No: CC-Y1528) and HeLa (EK-Bioscience Cat.No: CC-Y1211).

We acquired SH-SY5Y cell line (YC-D014) from Ubigene. We cultured cell in 90% DMEM and

10% FBS. We passaged cells every two days. The cell lines were cultured with 10cm plates. We

amplified cell lines to 3-4 plates. For each cell line, once the cells reached confluence, we first

removed the culture medium. We digested the cells with 0.25% trypsin for 3-5 minutes and

neutralized the trypsin with 90% DMEM and 10% FBS. We pipetted to detach the cells and

collected all cell mix in one 15ml centrifuge tube. After centrifuging to collect the cells, we

resuspended them in PBS. Following repeated washes, we counted the cells with Countstar(Alit

Biotech) and diluted them to 10^6 cells/ml. Then we separated 1ml cell suspension into a

centrifuge tube and centrifuged the cells. The supernatant was removed and the pellets are stored

at -80 degrees.

RNA sequencing

The FastPure Cell/Tissue Total RNA Isolation Kit V2 (Vazyme RC112) was used to isolate total

RNA from cell lines pellets. VAHTS Universal V6 RNA-seq Library Prep Kit (Vazyme NR604)

was employed to generate sequencing libraries from the isolated total RNA. MGI2000 was used

for sequencing the libraries. Samples are multiplexed in each lane, which yielded targeted number

of paired-end, 100bp reads for each sample.

Bulk RNA-seq data preprocess

We remove low quality reads with Trimmomatic [38], mapped reads with STAR [39] and generate

counts matrix with featurecounts [40]. We used combat to remove batch effect between our

in-house dataset and DepMap dataset [36,37]. The preprocessed dataset was used for subsequent

flux estimation analysis.

Metabolomics detection

We used targeted metabolomic analysis, Metabolites from the cells were extracted with 80%

Acetonitrile by vigorous vortex and centrifugation at 22 000g for 20 min at 4 °C. The supernatants

were used for analysis. The mix is vortexed and centrifuged. We used suspension for analysis.

Chromatographic separation was performed on a I Class ultra-high-performance liquid

chromatography system (Waters, Milford, Massachusetts, USA), with an InfinityLab Poroshell

120 HILIC-Z column (2.1 mm ×100 mm, 2.7 μm, agilent) and the following gradient: 0min,

100%B; 4min 84%B; 11min 40%B; 12min 40%B; 13min 100%B; 17min 10%B. Mobile phase A

was 10 mM ammonium acetate in water. Mobile phase B was 10 mM ammonium acetate in 90%

acetonitrile. The flow rate was 0.4 mL/min. The column temperature was kept at 35 °C and the

autosampler was kept at 8 °C. The injection volume was 5 μL. Mass data acquisition of the

metabolites was performed using a Triple QuadTM 7500 mass spectrometer (SCIEX, Framingham,

MA) equipped with an electrospray ion source in multiple reaction monitoring (MRM) mode. The

parameters of the electrospray ion source were:

neg: A：10mM ammonium acetate，pH=8.5　B：10% 10mM ammonium acetate，pH=8.5，90%

Acetonitrile

pos: A：10mM ammonium formate，pH=3　B：10% 10mM ammonium formate，pH=3，90%

Acetonitrile

The MRM transitions of all of the derivatized metabolites were shown in followed Table:

	metabolites	Q1	Q3
Group 1	BETA-ALANINE	90.1	44.2
Group 2	CITRULLINE	176	159
Group 3	DEOXYADENOSINE	252	136
Group 4	HYPOXANTHINE	137	110
Group 5	NICOTINAMIDE	123.1	80
Group 6	PROLINE	116.1	70.1
Group 7	PUTRESCINE	89	72
Group 8	URIDINE MONOPHOSPHATE	325	97
Group 9	URACIL	113	70
Group 10	FUMARATE	115	71
Group 11	PYRUVATE	87	43
Group 12	SUCCINATE	117	73

Metabolomic data analysis

SCIEX was used to process and integrate the components' peaks and provide integrated extracted

ion chromatograms for each metabolite component in all cell line samples and internal standard

samples. The generated results are normalized to the internal standard samples and the normalized

results are used for absolute quantification with the aid of the calibration curve.

TCGA survival analysis

We used easyTCGA (https://github.com/ayueme/easyTCGA) to download TCGA clinical dataset

and the expression matrix. We scaled the expression transcript per million (TPM) matrix using

log2 transformation and utilized combat [41] to remove batch effects between glioma and GBM.

After prediction of MGFEA, we divided all samples into two groups based on predicted

metabolite levels and conducted Kastle–Meyer test to identify which predicted metabolite serves

as a biomarker.

Data availability

All datasets used in this study have been published and can be obtained in h5ad format from

https://cellxgene.cziscience.com/datasets. The raw sequencing data of cancer cell line datasets

have been deposited at CNGBdb under the accession number CNP0007635.

We used Recon3D and IMM1865 as human and mouse GSMM model. Our raw file and

preprocess code can be obtained from our github site (https://github.com/Sunwenzhilab/MGFEA).

Detailed message and URLs of datasets is recorded in Table 1

528   Table 1

datasets	Source	website
4M brain dataset	Cao et al. [54]	https://cellxgene.cziscience.com/collections/c114c2 0f-1ef4-49a5-9c2e-d965787fb90c
1M GBM dataset	Ruiz Moreno et al. [55]	https://cellxgene.cziscience.com/collections/999f2a1 5-3d7e-440b-96ae-2c806799c08c
200k brain dataset	Siletti et al. [11]	https://cellxgene.cziscience.com/collections/283d65 eb-dd53-496d-adb7-7570c7caa443
50k brain dataset	Smith et al. [64]	https://cellxgene.cziscience.com/collections/e02201 d7-f49f-401f-baf0-1eb1406546c0
10k brain dataset	Siletti et al. [11]	https://cellxgene.cziscience.com/collections/283d65 eb-dd53-496d-adb7-7570c7caa443
Yu dataset	Yu et al. [45]	https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc =GSE117891
Stereo-seq dataset	Stereo-seq	http://116.6.21.110:8090/share/dd965cba-7c1f-40b2 -a275-0150890e005f
SMA dataset	Vicari et al. [24]	https://data.mendeley.com/datasets/w7nw4km7xd/1
Visium sagittal brain section	10X Genomics	https://www.10xgenomics.com/datasets/preservation -method-comparison-on-visium-cytassist-fixed-froz en-mouse-brain-sagittal-11-mm-capture-area-2-stan dard
Shao dataset	Shao et al. [53]	https://www.nature.com/articles/s41392-023-01552- y
DepMap transcriptome	Ghandi et al. [36]	https://depmap.org/portal/download/custom/
DepMap metabolome	Li et al. [37]	https://depmap.org/portal/download/custom/

Code availability

The code and related dataset can be accessible from the following GitHub respiratory

(https://github.com/Sunwenzhilab/MGFEA).

Author Contributions

D.A and J.L. conceived the project. W.S. supervised the whole project. J.L. implemented all model

codes. J.L. and D.A. performed all data analysis. All authors reviewed and edited the manuscript.

Acknowledgement

We thank all the member of Sun Laboratory and other colleagues from CIBR for their feedback on

this work.

Funding

This project is financially supported by STI2030-Major Projects (2022ZD0204700) and CIBR

Internal Fund to W.S..

References

1. Finley LWS. What is cancer metabolism? Cell 2023; 0:

2. Martínez-Reyes I, Chandel NS. Cancer metabolism: looking forward. Nat. Rev.
Cancer 2021; 21:669–680

3. Melone MAB, Valentino A, Margarucci S, et al. The carnitine system and cancer
metabolic plasticity. Cell Death Dis. 2018; 9:1–12

4. Bunse L, Pusch S, Bunse T, et al. Suppression of antitumor T cell immunity by the
oncometabolite (R)-2-hydroxyglutarate. Nat. Med. 2018; 24:1192–1203

5. Notarangelo G, Spinelli JB, Perez EM, et al. Oncometabolite d-2HG alters T cell
metabolism to impair CD8+ T cell function. Science 2022; 377:1519–1529

6. Panieri E, Santoro MM. ROS signaling and redox biology in endothelial cells. Cell.
Mol. Life Sci. 2015; 72:3281–3303

7. Panieri E, Santoro MM. ROS homeostasis and metabolism: a dangerous liason in
cancer cells. Cell Death Dis. 2016; 7:e2253–e2253

8. O'Neill LAJ, Kishton RJ, Rathmell J. A guide to immunometabolism for
immunologists. Nat. Rev. Immunol. 2016; 16:553–565

9. La Manno G, Siletti K, Furlan A, et al. Molecular architecture of the developing
mouse brain. Nature 2021; 596:92–96

10. Yao Z, van Velthoven CTJ, Kunst M, et al. A high-resolution transcriptomic and
spatial atlas of cell types in the whole mouse brain. Nature 2023; 624:317–332

11. Siletti K, Hodge R, Mossi Albiach A, et al. Transcriptomic diversity of cell types
across the adult human brain. Science 2023; 382:eadd7046

12. Chen X, Huang Y, Huang L, et al. A brain cell atlas integrating single-cell
transcriptomes across human brain regions. Nat. Med. 2024; 30:2679–2691

13. Lanekoff I, Sharma VV, Marques C. Single-cell metabolomics: where are we and
where are we going? Curr. Opin. Biotechnol. 2022; 75:102693

14. Guo S, Zhang C, Le A. The limitless applications of single-cell metabolomics.
Curr. Opin. Biotechnol. 2021; 71:115–122

15. Yu G, Wang L-G, Han Y, et al. clusterProfiler: an R package for comparing
biological themes among gene clusters. Omics J. Integr. Biol. 2012; 16:284–287

16. Xu S, Hu E, Cai Y, et al. Using clusterProfiler to characterize multiomics data.
Nat. Protoc. 2024; 19:3292–3320

17. Zhao K, Rhee SY. Interpreting omics data with pathway enrichment analysis.
Trends Genet. TIG 2023; 39:308–319

18. Hrovatin K, Fischer DS, Theis FJ. Toward modeling metabolic state from
single-cell transcriptomics. Mol. Metab. 2022; 57:101396

19. Orth JD, Thiele I, Palsson BØ. What is flux balance analysis? Nat. Biotechnol.
2010; 28:245–248

20. Damiani C, Maspero D, Filippo MD, et al. Integration of single-cell RNA-seq data
into population models to characterize cancer metabolism. PLOS Comput. Biol. 2019;
15:e1006733

21. Alghamdi N, Chang W, Dang P, et al. A graph neural network model to estimate
cell-wise metabolic flux using single-cell RNA-seq data. Genome Res. 2021;

gr.271205.120

22. Wagner A, Wang C, Fessler J, et al. Metabolic modeling of single Th17 cells
reveals regulators of autoimmunity. Cell 2021; 184:4168-4185.e21

23. Huang Y, Mohanty V, Dede M, et al. Characterizing cancer metabolism from bulk
and single-cell RNA-seq data using METAFlux. Nat. Commun. 2023; 14:4883

24. Vicari M, Mirzazadeh R, Nilsson A, et al. Spatial multimodal analysis of
transcriptomes and metabolomes in tissues. Nat. Biotechnol. 2023; 1–5

25. Kipf TN, Welling M. Variational Graph Auto-Encoders. 2016;

26. Brunk E, Sahoo S, Zielinski DC, et al. Recon3D enables a three-dimensional view
of gene variation in human metabolism. Nat. Biotechnol. 2018; 36:272–281

27. Khodaee S, Asgari Y, Totonchi M, et al. iMM1865: A New Reconstruction of
Mouse Genome-Scale Metabolic Model. Sci. Rep. 2020; 10:6177

28. Ebrahim A, Lerman JA, Palsson BO, et al. COBRApy: COnstraints-Based
Reconstruction and Analysis for Python. BMC Syst. Biol. 2013; 7:74

29. Cao Z-J, Gao G. Multi-omics single-cell data integration and regulatory inference
with graph-linked embedding. Nat. Biotechnol. 2022; 40:1458–1466

30. Kierans SJ, Taylor CT. Regulation of glycolysis by the hypoxia-inducible factor
(HIF): implications for cellular physiology. J. Physiol. 2021; 599:23–37

31. D van D, R S, J N, et al. Recovering Gene Interactions from Single-Cell Data
Using Data Diffusion. Cell 2018; 174:

32. K D, S Z. Deciphering spatial domains from spatially resolved transcriptomics
with an adaptive graph attention auto-encoder. Nat. Commun. 2022; 13:

33. Wolf FA, Angerer P, Theis FJ. SCANPY: large-scale single-cell gene expression
data analysis. Genome Biol. 2018; 19:15

34. Svensson V, Teichmann SA, Stegle O. SpatialDE: identification of spatially
variable genes. Nat. Methods 2018; 15:343–346

35. Palla G, Spitzer H, Klein M, et al. Squidpy: a scalable framework for spatial
omics analysis. Nat. Methods 2022; 19:171–178

36. Ghandi M, Huang FW, Jané-Valbuena J, et al. Next-generation characterization of
the Cancer Cell Line Encyclopedia. Nature 2019; 569:503–508

37. Li H, Ning S, Ghandi M, et al. The landscape of cancer cell line metabolism. Nat.
Med. 2019; 25:850–860

38. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 2014; 30:2114–2120

39. Dobin A, Davis CA, Schlesinger F, et al. STAR: ultrafast universal RNA-seq
aligner. Bioinformatics 2013; 29:15–21

40. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. Bioinformatics 2014; 30:923–930

41. Leek JT, Johnson WE, Parker HS, et al. The sva package for removing batch
effects and other unwanted variation in high-throughput experiments. Bioinformatics
2012; 28:882–883

42. Zhang Z, Zhu H, Dang P, et al. FLUXestimator: a webserver for predicting
metabolic flux and variations using transcriptomics data. Nucleic Acids Res. 2023;
51:W180–W190

43. Chen A, Liao S, Cheng M, et al. Spatiotemporal transcriptomic atlas of mouse
organogenesis using DNA nanoball-patterned arrays. Cell 2022; 185:1777-1792.e21
44. Zhao Q, Zhang T, Yang H. ScRNA-seq identified the metabolic reprogramming
of human colonic immune cells in different locations and disease states. Biochem.
Biophys. Res. Commun. 2022; 604:96–103
45. Yu K, Hu Y, Wu F, et al. Surveying brain tumor heterogeneity by single-cell
RNA-sequencing of multi-sector biopsies. Natl. Sci. Rev. 2020; 7:1306–1318
46. Kubelt C, Peters S, Ahmeti H, et al. Intratumoral Distribution of Lactate and the
Monocarboxylate Transporters 1 and 4 in Human Glioblastoma Multiforme and Their
Relationships to Tumor Progression-Associated Markers. Int. J. Mol. Sci. 2020;
21:6254
47. Yin D, Wang X, Konda BM, et al. Increase in Brain Tumor Permeability in
Glioma-Bearing Rats with Nitric Oxide Donors. Clin. Cancer Res. 2008;
14:4002–4009
48. Cheng J, Yan J, Liu Y, et al. Cancer-cell-derived fumarate suppresses the
anti-tumor capacity of CD8+ T cells in the tumor microenvironment. Cell Metab.
2023; 35:961-978.e10
49. Wang L-B, Karpova A, Gritsenko MA, et al. Proteogenomic and metabolomic
characterization of human glioblastoma. Cancer Cell 2021; 39:509-528.e20
50. Takano T, Lin JH-C, Arcuino G, et al. Glutamate release promotes growth of
malignant gliomas. Nat. Med. 2001; 7:1010–1015
51. de Groot J, Sontheimer H. Glutamate and the Biology of Gliomas. Glia 2011;
59:1181–1189
52. Yang S, Zhan Q, Su D, et al. HIF1α/ATF3 partake in PGK1 K191/K192
succinylation by modulating P4HA1/succinate signaling in glioblastoma.
Neuro-Oncol. 2024; 26:1405–1420
53. Shao Y, Fu Z, Wang Y, et al. A metabolome atlas of mouse brain on the global
metabolic signature dynamics following short-term fasting. Signal Transduct. Target.
Ther. 2023; 8:1–12
54. Cao J, O'Day DR, Pliner HA, et al. A human cell atlas of fetal gene expression.
Science 2020; 370:eaba7721
55. Ruiz-Moreno C, Salas SM, Samuelsson E, et al. Harmonized single-cell landscape,
intercellular crosstalk and tumor architecture of glioblastoma. 2022;
2022.08.27.505439
56. Dagogo-Jack I, Shaw AT. Tumour heterogeneity and resistance to cancer
therapies. Nat. Rev. Clin. Oncol. 2018; 15:81–94
57. Svensson V, Vento-Tormo R, Teichmann SA. Exponential scaling of single-cell
RNA-seq in the past decade. Nat. Protoc. 2018; 13:599–604
58. Fulcher JM, Markillie LM, Mitchell HD, et al. Parallel measurement of
transcriptomes and proteomes from same single cells using nanodroplet splitting. Nat.
Commun. 2024; 15:1–13
59. Wang D, Eraslan B, Wieland T, et al. A deep proteome and transcriptome
abundance atlas of 29 healthy human tissues. Mol. Syst. Biol. 2019; 15:e8503
60. Weigt D, Sammour DA, Ulrich T, et al. Automated analysis of lipid drug-response

markers by combined fast and high-resolution whole cell MALDI mass spectrometry biotyping. Sci. Rep. 2018; 8:11260

61. Wang Y, Li S, Qian K. Nanoparticle-based applications by atmospheric pressure matrix assisted desorption/ionization mass spectrometry. Nanoscale Adv. 2023; 5:6804–6818

62. Sun Z, Wang F, Liu Y, et al. Recent strategies for improving MALDI mass spectrometry imaging performance towards low molecular weight compounds. TrAC Trends Anal. Chem. 2024; 175:117727

63. Tan JH, Fraser AG. Quantifying metabolites using structure-switching aptamers coupled to DNA sequencing. Nat. Biotechnol. 2025; 1–4

64. Smith RS, Florio M, Akula SK, et al. Early role for a Na+,K+-ATPase (ATP1A3) in brain development. Proc. Natl. Acad. Sci. 2021; 118:e2023333118