Description of new micro-colonial fungi species *Neophaeococcomyces mojavensis*, *Coniosporium tulheliwenetii, and Taxawa tesnikishii* cultured from biological soil crusts

Tania Kurbessoian[1], Sarah A. Ahmed[2], Yu Quan[2], Sybren de Hoog[2], Jason E. Stajich [1],*

[1]*Department of Microbiology and Plant Pathology and Institute of Integrative Genome Biology, University of California, 92521, Riverside, CA, USA*

[2]*Radboudumc/CWZ Ceter of Expertise for Mycology, Nijmegen, 6525GA Nijmegen, The Netherlands*

*Correspondence: jason.stajich@ucr.edu; Tel.: +1 951.827.2363 (J.E.S.)

Abstract

Black yeasts and relatives comprise Micro-Colonial Fungi (MCFs) which are slow-growing stress-tolerant micro-eukaryotes that specialize in extreme environments. MCFs are paraphyletic and found in the Orders *Chaetothyriales* (*Eurotiomycetes*) and *Dothideales* (*Dothidiomycetes*). We have isolated and described three new MCFs species from desert biological soil crusts (BSCs) collected from two arid land regions: Joshua Tree National Park (Mojave Desert) and UC Natural Reserve at Boyd Deep Canyon (confluence of Mojave and Sonoran Deserts). BSCs are composite assemblages of cyanobacteria, eukaryotic algae, fungi, lichens, and bryophytes embedded into the surface of desert soils, providing a protective buffer against the harsh desert environment. Our work focused on one type of desert BSC, the cyanolichen crust dominated by *Collema sp.* Using culture-dependent protocols, three MCFs were axenically isolated from their respective samples along with the extracted DNA. Their genomes were sequenced using Illumina and Nanopore, and finally assembled and annotated using hybrid assembly approaches and established bioinformatics pipelines to conduct final taxonomic phylogenetic analysis and placement. The three species described here are unique specimen from desert BSCs, here we

introduce, *Neophaeococcomyces mojavensis* (*Chaetothyriales)*, *Cladosporium tulheliwenetii*

(*Dothideales*), and *Taxawa tesnikishii* (*Dothideales*).

Introduction

Biological Soil Crusts

Dryland ecosystems constitute some of the largest terrestrial biomes, collectively covering

47.2% of Earth's land surface (6.15 billion hectares) (Lal 2004). Drylands are expanding due to

climate change and population pressures, while also supporting over 38% of the global human

population (Lal 2004; Maestre et al. 2012). Biological Soil Crusts (BSCs) cover up to 70% of

dryland ecosystems (Rozenstein and Karnieli 2015). They are an ancient feature of dryland

ecosystems, with the earliest known fossil records indicating nearly 2.6 billion years ago

(Beraldi-Campesi and Retallack 2016). The initial phases of BSC formation in drylands are the

stabilization of the soil surface by filamentous cyanobacteria, followed by the colonization of

lichens and bryophytes (Weber et al. 2016). Within dryland ecosystems, the carbon and nitrogen

dynamics are often described as the "pulse-reserve model" where precipitation events happen

sporadically, stimulating intermittent biological activity which generates biomass and organic

matter. After extended periods of heat and dryness, the biological activity goes into reserve and

becomes dormant again (Pietrasiak et al. 2013). BSCs also increase soil fertility by increasing

soil stability and adding fixed atmospheric carbon and nitrogen to the soil underneath, allowing

for greater complex microbial and micro-faunal populations (Housman et al. 2007).

Within BSCs is a consortium of cyanobacteria, algae, diazotrophic bacteria, lichen, fungi, and

bryophytes that colonize and stabilize the soil surfaces against natural forces like wind and water

(Warren et al. 2019). These microorganisms recruit partners to create a mutualistic symbiotic

relationship which helps maintain a self-sustaining micro-ecosystem. Phototrophic organisms fix

carbon that is later used by surrounding micro-organisms as an energy source (Green et al.

2016). The presence of chemoheterotrophic bacteria and cyanobacteria indicates that BSCs are

capable of fixing atmospheric nitrogen to increase levels of fixed nitrogen for nearby living

vascular plants (Belnap and Harper 1995; Evans and Ehleringer 1993). BSCs have been shown

to reduce surface erodibility as filaments of cyanobacterial sheath material lipopolysaccharides

(LPS) entangle surface particles and create a crust that is more resistant to entrainment than the

layers below (Lal 2004).

60
Different ecological sites promote different crust taxa. BSCs are classified into types based on

the dominant photosynthetic microorganism in the crust which includes: cyanobacteria, algae,

lichens, or mosses to produce Cyano-Lichenized Crust, Light Algal Crusts, Dark Algal Crusts,

Lichenized Crust, Rough Moss Crust and Smooth Moss Crust (Pietrasiak 2005). The community

types are classified based on dominant taxa and their distinct morphologies (Pietrasiak et al.

2013). BSC's main roles include: 1) mediating almost all inputs and outputs (gasses, nutrients,

water) to and from the strata above and below the surface, 2) being the zone of high nutrient

deposition, transformation, and availability, 3) structuring temporal, spatial, and compositional

aspects of the surrounding vascular plant community, and 4) facilitating the direct delivery of

carbon, nutrients, and water by BSCs from the soil interspace to nearby vascular plants.

Micro-colonial Fungi

'Black yeasts and relatives' comprise a large amalgamate of melanized fungi. Among these are

micro-colonial fungi (MCF) or rock-inhabiting fungi (RIF), prevalent names used to describe a

special group of melanized fungi. Micro-colonial fungi are ubiquitous and found in the most

extreme environments. They are present in widely divergent habitats, such as on the indoor

rubbers of dishwashers, in oil deposits, on the surfaces of Mediterranean monuments, on the

radioactive walls of the Chernobyl Station, in mutilating skin infections in third-world countries,

in human superficial and invasive infections, or colonizing lungs of cystic fibrosis patients (de

Hoog et al. 2005; Hoog et al. 2000; Vicente et al. 2008; Gümral et al. 2014; Saunte et al. 2012;

Thanh and Hien 2019; Mehdiabadi and Schultz ; Little and Currie 2008; de Hoog and

Hermanides-Nijhof 1977). MCFs are compelling not only because of where we can discover

them and their unique physiological features, but also because of how little we know about them.

85
MCFs have three main characteristics: meristematic growth, presence of 1,8-DHN melanin, and

membrane-associated carotenoids and intracellular mycosporine-like amino acids (Staley et al.

1982; Gorbushina et al. 2003). These features distinguish them from other fungi, yet

phylogenetically they fall into two paraphyletic Ascomycota clades, the *Dothideomycetes* and

Chaetothyriomycetes. Melanin is a highly complex secondary metabolite that is produced by

many organisms through different evolutionary lineages (Siletti et al. 2017; Wheeler 1983). Most

importantly in MCFs, it has been used as a virulence factor as a non-specific armor against host

immune system to infect its host (Steenbergen and Casadevall 2003; Casadevall et al. 2003;

Casadevall et al. 2000; Nosanchuk et al. 2015; Taborda et al. 2008; Gómez and Nosanchuk

2003). Available research indicates melanin has other functional roles, such as metal chelation,

photoprotection, protection from heat and cold stress, binding to antifungals to reduce efficacy,

radioprotection (physical shielding and quenching of cytotoxic free radicals), protection from

desiccation, interacting with a range of electromagnetic radiation frequencies and mechanical-

chemical cellular strength (Dadachova et al. 2008; Casadevall et al. 2003; Gorbushina et al.

2003). Understanding poorly studied fungal function in one extreme environment could yield

insight into MCF adaptation in other extreme environments.

102
The objective of this study was to collect MCF from biological soil crust samples from desert

drylands. During our collection, we isolated 3 MCF that had not been described previously. Our

efforts to converse with indigenous tribes allowed us to provide a new naming scheme for our

strains. We also used a new methodology described herein using BUSCO nucleotide and amino

acid sequences to build a multi-locus phylogenetic tree, which proved to be similar to a more

conventional tree of MCF which used four highly conserved markers.

109
Materials and Methods

Sampling

Biological soil crusts were collected from two locations in Southern California. The first crust

location at the northern edge of Joshua Tree National Park. The second crust location was from

Boyd Deep Canyon, near the campgrounds in the UC Natural Reserve. Metadata and coordinates

can be found in **Table 1**. All materials were collected under permits from the BLM or permission

of the UC National Reserves.

Culturing and isolation

Media for isolation, cultivation, and collection used were Malt Extract Yeast Extract (MEYE)

media for nutrient-rich complex media and nutrient-poor, oligotrophic media Glucose

Asparagine Agar (GAA). The recipe for both media can be found in our protocols.io and the

step-by-step protocol for culturing MCFs (Kurbessoian 2019)). The mixing of the sample with

the media provides an environment with different layers of dissolved oxygen. Fast-growing fungi

require full oxygenation and grow on the surface of the media, while slow-growing MCFs

propagate on the bottom of the culture plate, requiring low levels of oxygen. This also allowed a

much easier way to culture and select the strains. Once grown, axenic MCFs were isolated by

using a flamed loop to pick and place onto clean media.

128
Morphological analysis

Strains were grown on Malt Extract Agar (MEA; Oxoid, England), Oatmeal Agar (OA; Oxoid)

and Potato dextrose Agar (PDA; Sigma, Germany) incubated at 28°C for 14 days under UV

light. Colonies were photographed with a Nikon C-DSD230 dissecting microscope. Microscopic

mounts were made in lactic acid and examined using a light microscope (Nikon Eclipse 80i).

Pictures were taken with a digital camera (Nikon Digital Sight, DS-5M) and adjusted in Adobe

Photoshop CS.

136
DNA extraction

Each isolate was grown on MEYE for approximately 1.5 weeks at 22°C. Genomic DNA was

extracted from hyphae collected from MEYE plates using the CTAB protocol described in our

protocols.io document (Carter-House et al. 2020). Melanin was removed using multiple

iterations of phenol:chloroform and chloroform washes of our sample. Genomic DNA was

measured by Nanodrop and diluted to ~28ng/µl. DNA extractions were sent to the UCR

Genomics core (Riverside, CA) for 2x150bp sequencing on an Illumina NovoSeq 6000. DNA

from all three isolates was also extracted and sequenced on Oxford Nanopore (ONT) platform

with library preparation and sequencing following the manufacturer's directions (Oxford

Nanopore, Oxford United Kingdom). Flow cell versions FAK68780, FAQ05493, FAL40904,

and FAL35715 were used along with base-calling on the cluster using guppy (*v.3.4.4.+a296cb*)

(Wick et al. 2019).

149
ITS sequencing

To identify the isolates, extracted DNA was used as a template for PCR amplification using

fungal primers ITS1-F (CTTGGTCATTTAGAGGAAGTAA) and ITS4

(TCCTCCGCTTATTGATATGC) along with the PCR setup. PCR was performed in 25 μL

reaction volumes using 12.5 μL of Taq 2X MasterMix (NEB), 9.5μL water, 1μL of 20 picomol

ITS1-F, 1μL of 20 picomol ITS4 and 1μL of template DNA. Using Bio-Rad thermal cycler, the

samples were passed through denaturation and enzyme activation at 95°C for 5 minutes,

followed by amplification for 35 cycles at the following conditions: 30 seconds at 95°C, 1

minute at 72°C, and 30 seconds at 65°C. A final 5-minute extension at 72°C completes the

protocol. PCR results typically yield between 50-60 ng of amplified DNA.

Genome assembly and annotation

Initial genome assemblies were constructed for the three MCF isolates using Illumina

sequencing. Each MCF was also re-sequenced using Oxford Nanopore technology. All genomes

were *de novo* assembled with the AAFTF pipeline (*v.0.2.3*) (Palmer and Stajich 2022) which

performs read QC and filtering with BBTools bbduk (*v.38.86)* (Bushnell 2014) followed by

SPAdes (*v.3.15.2*) (Bankevich et al. 2012) assembly using default parameters, followed by

screening to remove short contigs < 200 bp and contamination using NCBI's VecScreen. The

BUSCO ascomycota_odb10 database (Manni et al. 2021) was used to determine how complete

the assembly 3 isolates were. A hybrid assembly of each was generated using MaSUrCa *(v.3.3.4)*

(Zimin et al. 2013) as the assembler using both Nanopore and Illumina sequencing reads.

General default parameters were used except: CA_PARAMETERS=cgwErrorRate=0.15,

NUM_THREADS=16, and JF_SIZE=200000000.

We predicted genes in each near-complete genome assembly with Funannotate (*v1.8.1*) (Palmer

and Stajich 2020). A masked genome was created by generating a library of sequence repeats

with the RepeatModeler pipeline (Smit and Hubley 2008). These species-specific predicted

repeats were combined with fungal repeats in the RepBase (Bao et al. 2015) to identify and mask

repetitive regions in the genome assembly with RepeatMasker (*v.4-1-1*) (SMIT 2004). To predict

genes, *ab initio* gene predictors SNAP (*v.2013_11_29*) (Korf 2004) and AUGUSTUS (*v.3.3.3*)

(Stanke et al. 2006)were used along with additional gene models by GeneMark.HMM-ES

(*v.4.62_lic*) (Brůna et al. 2020), and GlimmerHMM (*v.3.0.4*) (Majoros et al. 2004) utilize a self-

training procedure to optimize *ab initio* predictions. Additional exon evidence to provide hints to

gene predictors was generated by DIAMOND BLASTX alignment of SwissprotDB proteins and

polished by Exonerate (*v.2.4.0*) (Slater and Birney 2005). Finally, EvidenceModeler (*v.1.1.1*)

(Haas et al. 2008) generated consensus gene models in Funannotate that were constructed using

default evidence weights. Non-protein-coding tRNA genes were predicted by tRNAscan-SE

(*v.2.0.9*) (Lowe and Chan 2016).

187
The annotated genomes were processed with antiSMASH (*v.5.1.1*) (Blin et al. 2021) to predict

secondary metabolite biosynthesis gene clusters. These annotations were also incorporated into

the functional annotation using Funannotate. Putative protein functions were assigned to genes

based on sequence similarity to InterProScan5 (*v.5.51-85.0*) (Jones et al. 2014), Pfam (*v.35.0*)

(Finn et al. 2014), Eggnog (*v.2.1.6-d35afda*) (Huerta-Cepas et al. 2019), dbCAN2 (*v.9.0*) (Zhang

et al. 2018) and MEROPS (*v.12.0*) (Rawlings et al. 2018) databases relying on NCBI BLAST

(*v.2.9.0+*) (Sofi et al. 2022) and HMMer (*v.3.3.2*) (Potter et al. 2018). Gene Ontology terms were

assigned to protein products based on the inferred homology based on these sequence similarity

analyses. The final annotation produced by Funannotate was deposited in NCBI as a genome

assembly with gene model annotation.

198
199   Sequence Read Archive (SRA) files and genome assembly and annotation files can be found

200   under BioProject PRJNA631111.

201
202   *Mating type determination*

203   The Mating Type (MAT) locus was identified by searching for homologous MAT genes. The

204   identified homologous regions were examined for their conserved synteny of the MAT locus

205   using clinker (Gilchrist and Chooi 2021) and a custom Biopython script (Cock *et al.* 2009)

206   (Kurbessoian 2022) to extract the annotated region of the genome which contained the locus.

207
Telomeric repeat determination

Identification of telomeric repeat sequences was performed using the FindTelomeres.py script

(https://github.com/JanaSperschneider/FindTelomeres). Briefly, this searches for chromosomal

assembly with a regular expression pattern for telomeric sequences at each scaffold's 5' and 3'

end. Telomere repeat sequences were also predicted using the Telomere Identification toolkit

(tidk) (*v.0.1.5*) "explore" option (https://github.com/tolkit/telomeric-identifier).

214
215   *Inferring phylogenetic relationships of species*

216   Inference of the species relationship was made through phylogenetic trees containing the three

217   new MCF strains described here and available named species with sequence data.

218
A multi-gene tree was generated using collected BT2 (Beta-Tubulin), CAM (Calmodulin), ITS

(Inter-Transcribed Spacer), and LSU (Large Subunit) genes from a total of 32 species. These

samples were chosen to compare to the three MCF species with closely related species to

confirm identification as a novel lineage. The final selection of 32 species was based on

iterations of sequence searching and tree building data analyses to determine close relatives. To

include genes from genomes, genome assemblies were searched by TBLASTN tool of NCBI-

BLAST (*v.2.9.0+*) using *Zymoseptoria tritici* genes (AAS55060.1 for Beta tubulin) and

(AEI69650.1 for Calmodulin). The collection of ITS and LSU genes were obtained from NCBI-

Genbank based on BLASTN searches. Each gene file contained FASTA formatted sequences

from each species or isolate, the sequences were aligned with MUSCLE (*v.5.1*) and trimmed for

low quality alignment regions using Clipkit (*v.1.3.0*). Nucleotide substitution models were

determined for each alignment with IQTREE2. The alignments were combined into a single

'supermatrix' alignment with combine_multiseq_aln.py script provided in PHYling. The script

also produced a partition file listing the sub-locations of each marker gene alignment in the

super-alignment. IQTree2 was run to find a substitution model fit for and partition merging. To

reduce the computational burden that occurs with including a large number of organism

ribosomal sequences we used the relaxed hierarchical clustering algorithm (Lanfear et al. 2014)

using the tag -m MF+MERGE and running 1000 bootstrapping on the alignment.

237
Choosing to work with indigenous tribes

Our choice for naming of the three MCFs isolated from BSCs is rooted in the traditional

indigenous land where these were found. *Neophaeococcomyces mojaviensis* was isolated from

BSCs collected from the south-east region of Joshua Tree National Park, territories ancestral to

Yuhaaviatam/Maarenga'yam (Serrano) and Newe Segobia (Western Shoshone) Tribes (Brown

and Boyd 1922; Kroeber 1925). The isolates of *Coniosporium tulheliwenetii* and *Taxawa*

tesnikishii were isolated from BSCs collected from UC Reserve Boyd Deep Canyon (Reserve

doi:10.21973/N3V66D), land of the traditional ancestral Cahuilla Tribe (White 2001; Hooper

1920). The word *tulheliweneti* in Cahuilla means, black-spread-be-noun, while *Taxaw* means

body, and *tesnikish* means yellow. In Cahuilla traditions, Taxaw tesnikish is the deity that

resided in Boyd Deep valley. To recognize the origins, we consulted with Tribal linguists and

speakers to derive their names from Cahuilla and Serrano terms that describe either geographic

region, oral tradition, or features that help distinguish them.

Results

252
Biological soil crust collection

Biological soil crusts were collected from the two regions described in **Table 1**. The method of

sterile collection is through the use of sterile empty Petri dishes, stamped on top, with a sterile

spatula used to grab the soil from underneath, flip the sample into the dish, place the lid on top

and seal it with parafilm.

Culturing results

Samples material was used to culture fungi using a collection of techniques in our laboratory at

UC Riverside. More than 50+ fungal culture specimens were collected from biological soil

crusts, 18 were MCF cultures. The pour plate protocol used allows for fast-growing fungi to

appear on the top, while MCF grows within the bottom layer of media in Petri dishes. Samples

were excised from deep in the media and allowed to grow on fresh MEYE. Each sample had the

DNA extracted and each strain ITS was Sanger sequenced to determine the identity. JES_112

was initially identified as a *Knufia* species. JES_115 and JES_119 were identified as *Dothideales*

species, but there remained ambiguity on the exact species identification as a near perfect

sequence match was not found for ITS1 to assign a name.

Table 1. Three species were collected from biological soil crusts collected from two desert locations
in Southern California. Strain JES_112 was isolated from the south east region of Joshua Tree National
Park in the Mojave Desert, while JES_115 and JES_119 were both collected from biological soil crusts
collected from the campground site in the UC Natural Reserve of Boyd Deep Canyon.

Species	Date	Location	Coordinates
JES_112	November 2017	Joshua Tree National Park, CA	34.1026 °N, -115.4542°W
JES_115	July 2019	Boyd Deep Canyon Reserve, CA	34.0182 °N, -116.0262 °W

JES_119	July 2019	Boyd Deep Canyon Reserve, CA	34.0182 °N, -116.0262 °W

Sequencing and assembly of MCF isolates

To gain information on the genome content for the recovered MCF isolates, we sequenced and assembled the genomes of the three isolates (**Table 2**). Each isolate was sequenced with Illumina MiSeq and Oxford Nanopore (ONT) technology and constructed into a hybrid assembly that combined both read types (**Table 2**). The depth of coverage ranged from 25-43x Illumina coverage across the three specimens, while Nanopore ranged from 0.83-194.58x. Assessment of the genome completeness was performed with BUSCO (Benchmarking Universal Single-Copy Orthologs) and results indicated the genomes had ranges of 73-92.9% complete gene copies found, with 73-92.8% of the markers appearing as single copies in the assembly, a low fraction of duplication (0.1-0.2). Contig counts ranged from 38-480, while the average genome assembly size is about 30 Mbps for all isolates. The L50 statistic, a measure of the count of the smallest number of contigs whose sum makes up half of the genome size, ranged from 9-88, and the N50 statistic, a measure of the smallest contig length for covering half of the genome size, ranging from 117kbp - 1.232 Mbp.

To further assess completeness, we tested for the presence of telomeric repeat units "TTTAGGG/CCCTAA". These were identified as repeat arrays at both ends of contigs in JES_115 and JES_119, 61 total telomeres and 55 total telomeres respectively, though the results were inconclusive for JES_112. No telomeric repeat-containing contigs were detected in

JES_112 which could be due to the fragmented nature of the genome assembly. The fragmented

assembly also is reflected in the lower BUSCO score for this genome (**Table 2**).

The number of predicted tRNA genes ranged from 76-197, the total number of protein-coding

genes predicted ranged from 9,029-10,920, and secondary metabolite cluster counts ranged from

15-27, 27 resulting from JES_112.

Table 2. Genome assembly and annotation summary statistics. All strains were sequenced with
Illumina and Nanopore sequencing. With the added sequencing reads, we were able to glean telomere
information along with genome completeness. It is clear that JES_112 initially was very contaminated as
the BUSCO score is lower, while JES_115 and JES_119 are much closer to a more complete genome.

	JES_112	JES_115	JES_119
Scaffold Count	478	64	38
Total Length	32,593,754 bp	32,607,911 bp	29,383,553 bp
Minimum Length	561	4823	1783
Maximum Length	427,480	1,739,523	2,828,030
Mean Scaffold Length	68,187.77	509,498.61	773,251.39
Scaffold L50	87	14	9
Scaffold N50	117,971	966,560	1,232,081
Scaffold L90	282	31	24
Scaffold N90	32,261	507,175	558,687
BUSCO Complete % Illumina	84.7	90.2	97.4

BUSCO Complete % Nanopore	73.4	92.6	92.9
BUSCO Duplicate % Nanopore	0.2	0.1	0.1
BUSCO Single % Nanopore	73.2	92.5	92.8
GC%	48.25	52.55	51.98
Illumina coverage	34.58x	43.45x	25.38x
Nanopore coverage	0.83x	20.44x	194.58x
Telomere Count	0, 0F, 0R	61, 29 F, 32 R	55, 29 F, 26 R
Number of Genes	10,996	9,304	9,386
mRNA genes	10,924	9,226	9,189
tRNA genes	72	78	197
GO Terms	6,001	5,089	5,321
InterProScan	8,523	7,079	7,291
EGGNOG	9,934	8,487	8,511
PFAM	7,520	5,982	6,219
CAZYme	309	257	346
MEROPS	344	284	291
Secretion	674	532	631

Mating type loci determination

The mating types were determined using prior work on members of Chaetothyriales and

Dothideomycetes including *Exophiala dermatitidis*, and *Aureobasidium pullulans* (Metin et al.

2019; Teixeira et al. 2017). Using collected nucleotide and protein FASTA sequences, BLAST

was used to identify the genes found in these loci. The genes in this locus are generally flanked

by SLA2 and APN2 and may sometimes have alternative genes within. **Figure 2** was generated

using JES_112, *Exophiala aquamarina* CBS 119918, *E. dermatitidis* NIH/UT8656, *E.*

dermatitidis CBS 115663, and *Capronia coronata* CBS 617.96. The MAT locus is flanked by

two genes, SLA2 (purple) and APN2 (orange). The MAT 1-1 gene, MAT 1-1-4 (green), and

MAT 1-1-1 (teal) are observed in *E. dermatitidis* CBS 115663 and *Capronia coronata*, while *E.*

dermatitidis NIH/UT8656 contains MAT 1-2 (pink) as does *Capronia coronata*. Our organism in

question, *Neophaeococcomyces mojaviensis* contains the MAT 1-2 gene along with the MAT 1-

1-1 gene. This is similarly seen with *N. aloes* FJII-L3-CM-P2. This is confusing as the MAT 1-1-

1 gene usually pairs with MAT 1-1-4. The gene arrow in blue is indicated as an L-type calcium

channel domain, while the gene in yellow is an unidentified functional gene or domain.

322
Figure 3 was generated using JES_115 and JES_119 and two other species including,

Aureobasidium pullulans EXF-150 and *Coniosporium apollinis* CBS 100218. All species

described here only have the MAT 1-2 gene. The MAT locus is flanked by two genes, SLA2

(purple) and APN2 (orange). *Dothideomycetes* mating type loci look very different from the

Chaetothyriales mating type loci. Other genes included in this locus concern only a single MAT

locus, MAT 1-2 (pink), and rRNA adenine demethylase (yellow). Certain genes like the AP

endonuclease (teal) and a hypothetical protein (blue) are not seen in all four genomes.

MCF morphology

Strains showed limited and slow growth with black colonies on all media tested. Strain JES_119

produced abundant unilateral, hyaline budding cells which soon inflated, developed cruciate

septation, and became melanized. Strain JES_115 was hyphal with lateral chains of swollen cells

which occasionally detached. Strain JES_112 lacked sporulation, the thallus consisting of

melanized densely septate hyphae with series of somewhat inflated cells

338
Figure 1. Microscopic morphology of isolated MCF. (1) *Taxawa tesnikishii* JES_119, colonies on
MEA (A), OA (B), and PDA (C) agars, meristematic cells (D-F), and budding cells (G-M); (2)
Coniosporium tulheliwenetii JES_115, colonies on MEA (A), OA (B), and PDA (C) agars, hyphae with
lateral branches of swollen cells reluctantly leading to arthroconidia (D-J); (3) *Neophaeococcomyces*
mojaviensis JES_112, colonies on MEA (A), OA (B), and PDA (C) agars, hyphae producing intercalary
swollen cells without conidiation (D-J). Microscopic mounts were made in lactic acid and examined using
a light microscope. The scale depicted here is 25µm.

Figure 2. Mating type loci determined for Chaetothyriales JES_112 here called *Neophaeococcomyces mojaviensis.* Other MCFs added to this analysis include *Neophaeococcomyces aloes* FJII-L3-CM-P2*, Exophiala dermatitidis* NIH/UT8656, *E. dermatitidis* CBS 115663, and *Capronia coronata* CBS 617.96. The MAT locus is flanked by two genes, SLA2 (purple) and APN2 (orange). The MAT 1-1 gene, MAT 1-1-4 (green), and MAT 1-1-1 (teal) are observed in *E. dermatitidis* CBS 115663 and *C. coronata*, while *N. aloes* and *E. dermatitidis* NIH/UT8656 contain MAT 1-2 (pink) as does *C. coronata*. Our organism in question, *N. mojaviensis* contains the MAT 1-2 and MAT 1-1-1 gene which is confusing as it is not common to lose the MAT 1-1-4. The gene arrow in blue is indicated as an L-type calcium channel domain, while the gene in yellow is an unidentified functional gene or domain.

Figure 3. Mating type loci determined for Dothideomycetes JES_115 and JES_119, here called *Coniosporium tulheliwenetii* **and** *Taxawa tesnikishi***.** Other dothideaceous black yeast-like fungi in this analysis include *Aureobasidium pullulans* EXF-150, and *Coniosporium apollinis CBS* 100218. All species described here only have the MAT 1-2 gene. The MAT locus is flanked by two genes, SLA2 (purple) and APN2 (orange). *Dothideomycetes* mating type loci look very different from the *Chaetothyriales* mating type loci. Other genes included in this locus include only a single MAT locus, MAT 1-2 (pink). Another gene included in this locus is rRNA adenine demethylase (yellow). Certain genes like the AP endonuclease (teal) and a hypothetical protein (blue) are not seen in all four genomes.

Secondary metabolite gene cluster predictions

During the annotation step, we processed the genomes through antiSMASH (**Table 3**), to detect

Biosynthetic Gene Clusters (BGCs) of secondary metabolites from the annotated genome

sequence. Once predicted metabolite of note, beta-lactone was found in all three MCF. A further

literature search indicates the importance of beta-lactone in industry as there have been multiple

products derived from these chemicals including, antifungals and anti-cancer agents (Robinson et

al. 2019). A BLASTP search of the JES_112 beta-lactone gene identified homologs to Acetyl-

CoA synthetase + 2-isopropylmalate synthases + bile acid dehydrogenase, zinc-binding alcohol

dehydrogenase + acetoacetate-CoA + NAD-dependent alcohol dehydrogenase, and AMP-

binding protein + 2-isopropylmalate synthases.

378
379   A second predicted metabolite of note was the siderophore gene cluster. These genes are

380   involved in iron transport, is a more well-known metabolite that could provide the organism

381   with virulence and protection by sequestering iron molecules from its surrounding environment

382   (Johnson 2008; Sass et al. 2019) and was primarily detected in JES_112.

383
Additionally, the BSC search revealed fungal-RiPPs (Ribosomal and post-transcriptionally

modified proteins) in JES_119. The homolog was best classified as a gamma-glutamyltransferase

+ cytochrome protein. Fungal-RiPPs have been used as an alternative source of bioactive

metabolites and could be of industrial interest (Vogt and Künzler 2019). Additional BGCs were

found including Type-1 Polyketide synthases (T1PKS), Type-3 Polyketide synthases, non-

ribosomal peptide synthases, non-ribosomal peptide synthase-like, and terpenes.

390
391

Table 3. Predicted secondary metabolite gene clusters. The BGCs predicted include multiple copies of PKS and NRPS genes, but siderophores were only recovered in JES_112. There was a higher number of BGCs predicted in JES_112 (27) our *Chaetothyriales* species, than JES_115 and JES_119 (15) our *Dothideomycetes*. This table lists the numbers of each. JES_112 has a much higher number (27) of secondary metabolites than the other two (15).

	T1PKS	Terpene	T3PKS	NRPS	NRPS-like	Siderophore	Arylpolyene	Betalactone	Fungal RiPP
JES_112	6	6	2	4	6	1	1	1	0
JES_115	5	4	0	2	3	0	0	1	0
JES_119	1	4	0	1	7	0	0	1	1

Four-locus gene trees

The multi-gene internal-transcribed space (ITS1), large ribosomal subunit (LSU) beta-tubulin, calmodulin tree (**Figure 4** and **Figure 5)** places the new isolates in context of named species in the known genera (**Table 3**).

Forty-three isolates were used, and the alignment of the sequences were trimmed, and concatenated to run the IQTree program and generate a concise multi-loci gene tree for *Neophaeococcomyces mojaviensis* JES112. Here this strain is placed in the genus *Neophaeococcomyces*, along with *Exophiala dermatitidis* NIH/UT8656, *E. radicis* P2854, *E. oligospoerma* CBS 265.49, *E. oligospoerma* CBS 127587, and *E. xenobiotica* CBS 11817, *Fonsecaea pedrosoi* CBS 271.37, *Capronia coronata* CBS 617.96, *Cladophialophora immunda* CBS 834.96, *C. carrionii* CBS 260.83, *C. yegresii* CBS 114405, *Neophaeococcomyces aloes* FIJI-L3-CM-P2*, N. catenatus* CBS650.76*, N. oklahomaensis* EMSL 3313*, Knufia petricola* MA5789*, K. separata* CGMCC3.17337*, K. mediterranea* CCFEE 6211*, K. mediterranea*

CCFEE6205, *Knufia* sp. MV-2018-MLT-8, *K. walvisbayicola* CBS 146989, *K. tsunedae*

FMR10621, *Bradymyces yunnanensis* CGmCC3.17350, *B. alpinus* CCFEE 5493, *Strelitziana*

africana ICMP 21758 and ICMP 21760, and *Tuber melanosporum* Mel8 as the outgroup.

The third phylogenetic tree, **Figure 5**, uses the same method described above as in **Figure 4**. 25

strains of Dothideomycetes MCF were included to place JES_115 and JES_119 into their proper

clades. MCF strains included a collection of *Coniosporium apollinis* CBS 100218, CBS 352.97,

Aureobasidium mangrovei IBRCM30265, *Aureobasidium* sp. AWRI4619-CO-2020, *Zalaria*

obscura FKI-L7-BK-DRAB2, *Polyplosphaeria fusca* CBS 125425, *Leptosphaeria maculans*

JN3, *Leptosphaeria biglobosa* CA1, *Phaeosphaeria* sp. MPI-PUGE-AT-0046c, *Alternaria solani*

NL03003, *Cenococcum geophilum* 1.58, *Venturia pyrina* ICMP11032, *Coniosporium uncinatum*

CBS 100219, and CCFEE 6149, *Scorias spongiosa* HLSS1, *Polychaeton citri* CBS 116435,

Zymoseptoria tritici CBS 100329, *Neohortaea acidophila* CBS 113389, *Salinomyces thailandica*

CCFEE 6315, *Hortaea werneckii* EXF-2000, *Baudoinia panamericana* UAMH 10762,

Teratosphaeria guachensis CMW 17545, *Friedmanniomyces simplex* CCFEE 5184,

Friedmanniomyces endolithicus CCFEE 5311, and *Tuber melanosporum* Mel8 as the outgroup.

JES_115 groups near *Coniosporium* spp., *w*hile JES_119 groups near *Aureobasidium* spp.

430
Taxonomy

***Neophaococcomyces mojaviensis sp. nov.*, Kurbessoian, Ahmed, de Hoog and Stajich -**

Mycobank (848161). Description: Colonies on MEA, OA, PDA after two weeks of incubation

at 25°C attain diameters of 1.6, 2, 2.2, and 1.6 mm respectively. Hyphae 2.9-5.4 μm diameter,

dark brown or black, smooth or occasionally rough walled, cells eventually inflating up to 8 μm

in diameter. The proliferation of inflated cells might result in the formation of multicellular

bodies terminal or intercalary in the hyphae measuring 13.8-31.8 μm, multiseptate with

transverse and longitudinal septa; bodies round or irregular, dark brown to black and thick-

walled. No conidia were observed. Sexual state unknown.

Holotype: ATCC (TSD-444), preserved at the herbarium collection of the Westerdijk Fungal

Biodiversity Institute, Utrecht, The Netherlands. Living culture ex-type ATCC (TSD-444).

Diagnosis: Strain JES_112 clusters in the order *Chaetothyriales*, relatively close to the numerous

rock-inhabiting species of the genus *Knufia*, and amidst the species of the small genus

Neophaeococcomyces. Both genera are assigned to the family *Trichomeriaceae*, containing

'sooty molds' with oligotrophic lifestyles colonizing inert surfaces (Chomnunti et al. 2012). The

cluster *Neophaeococcomyces,* based on *N. catenatus* as generic type species, as yet contains only

four species, which mostly were isolated from low nutrient habitats, such as the bark of *Aloe*,

stone, or air. Several species of the genus show budding with melanized cells, and frequently

cells cohere in short chains. Often lateral branches of short inflated cells are inserted on hyphae;

the cells may be released as arthroconidia which then swell or appear yeast-like, or remain

coherent.

Coniosporium tulheliwenetii sp. nov.**, Kurbessoian, Ahmed, de Hoog and Stajich -**

MycoBank (848162).

Description: Colonies on MEA, OA, PDA, after 2 weeks of incubation at 25°C attain diameter

restricted, measuring 1.1 cm, 0.72 cm, and 0.75cm, respectively. Hyphae 4.6-8.4 μm diameter

showing lateral chains of swollen cells 7.7 - 10.8 μm. The cells however remain intact, and no

conidia was observed. Teleomorph unknown.

Holotype: CBS (150075), and herbarium CBS H-25288, preserved at the herbarium collection

of the Westerdijk Fungal Biodiversity Institute, Utrecht, The Netherlands. Living culture ex-type

CBS (150075).

Diagnosis: Strain JES_115 was morphologically similar to *Knufia* species in the presence of

lateral chains of swollen cells inserted on main hyphae, the cells eventually being released as

conidia, but was phylogenetically remote. It clustered with some species of rock-inhabiting

described in *Coniosporium,* in the order *Pleosporales*. Interestingly, the original descriptions,

which were made before sequence data were available, classified them all into a single genus.

The genus *Coniosporium* is one of the classical genera of meristematic fungi. However, most

species were described in the 19th century, including the generic type species *C. olivaceum*, and

voucher specimens are lost. Nevertheless, Haridas et al. (2020) introduced the family

Coniosporiaceae and order *Coniosporales* using the few recently described rock-inhabiting

fungi, *C. apollinis* and *C. uncinatum* as reference (Haridas et al. 2020). It should be noted that

the family *Coniosporiaceae* was already used by Nannfeldt for the genus *Coniospora*.

Taxawa **gen. nov. Kurbessoian, Ahmed, de Hoog and Stajich - MycoBank (848163).**

Type species: *Taxawa tesnikishii*

Taxawa tesnikishii sp. nov. Kurbessoian, Ahmed, de Hoog and Stajich - MycoBank (848164).

Description: Colonies moist, black, regular and flat on OA and heaped irregular on MEA and

PDA, measuring 1.4-1.8 cm in diameter growing on MEA, OA, and PDA at 25°C . Hyphae

smooth, thick-walled, measuring 6.1-10.3 μm wide. Yeast cells abundant produce unilateral

budding 2.7-4.8 μm, subsequently inflating and becoming crucially septate and melanized with

age. Clusters of black meristematic bodies (9.0-56.8 μm) were observed. Sexual state unknown.

Holotype: CBS (150074), and herbarium CBS H-25287, preserved at the herbarium collection

of the Westerdijk Fungal Biodiversity Institute, Utrecht, The Netherlands. Living culture ex-type

CBS (150074).

Diagnosis: Strain JES_119 showed similar morphology to the cultural states of e.g. *Sydowia* and

Comminutispora, and has been described in the asexual genus *Phaeotheca*. Thambugala et al.

(2014) erected the family *Aureobasidiaceae* for fungi in this relationship, but also this name

existed already, *Aureobasidiaceae* Ciferri (1958) (Thambugala et al. 2014), based on the same

species, *Aureobasidium pullulans* (Ciferri 1958). Only a small number of extant species has been

sequenced, and the genome trees show an even larger taxon sampling effect, but given our

current inability to assign strain JES_119 to any genus, we introduce the genus *Taxawa*.

492
493   **Discussion**
494
BSCs are an untapped resource of novel microorganisms, using non-traditional methods of

culturing may be a method to uncover more unavailable species for sequencing. Using a pour

plate method for culturing allows the microorganism to avoid complete oxygen saturation (or

oxygen toxicity), and instead can grow in low to micro-aerophilic concentrations of oxygen

(Stevens 1995). Collecting MCF using this method has proved to be efficient, as fast-growing

fungi overtake the surface of plates, while MCF can tolerate lower oxygen concentrations, thus

localizing to the bottom of plates. Preparing axenic cultures of MCF is then relatively easy and

can also increase the rate of isolation as MCF tends to take two weeks to grow. A hypothesis

about the facultative aerobic or microaerophilic nature of MCF could also be seen in the

phenotype of these species. Morphological growth on surface of petri plates shows unusual

furrowing and clumping growth (JES_119, JES_115), some strains (JES_112) grow underneath

the surface of the plate which seem to be MCF adaptations to the high concentration of oxygen

present on petri plates.

The high melanin content of MCF complicates efforts to extract high molecular weight (HMW)

DNA for sequencing, however adjustments to standard fungal CTAB can dramatically improve

the gDNA purity. In the case for these BSC MCF, an increase in phenol:chloroform and

chloroform washes allow for melanin to be removed completely from the DNA samples. Another

consideration to help concentrate more DNA is to incubate the DNA sample in -20°C overnight

during the DNA precipitation step. Using long read sequencing technology (Oxford ONT) it is

best to have a very high concentration of clean DNA to process and thus could be the limiting

factor for these sequencing procedures. Using hybrid methods of sequencing (Illumina and

Nanopore) provides scientists the proper backbone to develop *de-novo* assemblies (Stevens

1995; Khezri et al. 2021; Chen et al. 2020; Saud et al. 2021). It is vital to have as near complete

assemblies of species as possible, this provides the scientific community with the best tools for

proper genomic assessments. For future research endeavors, a second nanopore flow cell would

be necessary to help build a better assembly for strain JES_112 (*Neophaeococcomyces*

mojaviensis) as the telomere analysis failed on this first round of assembly. The other two strains

had enough depth of coverage that allowed for different programs to optimize the assembly and

have more closed telomere contigs.

Scientific literature describing novel MCF always note the clonal or asexual nature, describing

"missing" sexual states and thus always being in a haploid state. The MAT locus is one of the

pivotal regions in the fungal genome that provides more accurate ploidy and sexual states of

fungi, and using this data to describe the three new MCF assisted in understanding the sexual

states. JES_115 and JES_119 as the two *Dothideomycetes* representatives indicate heterothallic

sexual states having only the MAT1-2 gene, *vs.* homothallic species having representatives of

both MAT1-1 and MAT1-2 genes. Better genome assembly would help better describe the

JES_112 MAT locus as there is the presence of MAT1-2 but also a portion of the MAT1-1 gene

(as it is composed of two genes, MAT1-1-1 and MAT1-1-4). The high melanin content of MCF

could prove to be difficult when testing for ploidy in the scientific "gold standard" method (flow

cytometry). But future research would most benefit from having this method optimized to

understand not only the new MCF described in this work but also for future MCF considerations.

Possibly the sexual states have not yet been described or seen for MCFs and may require a

particular environmental stimulus to produce the proper structures to reproduce. Possibly, other

forms of propagation are found in other habitats. A prototypical example is *Auerobasidium*

pullulans which shows meristematic growth on glass or rock but is a rapidly expanding yeast-

like fungus in culture. There has also been an increase in literature describing clonal MCF having

high hybridization rates that create pseudo but stable diploids without the process of

recombination (Gostinčar et al. 2022). This was not detected in these three species described in

this paper.

MCF morphology indicates usual slow growth (about two weeks) seen in most MCF described

previously (Selbmann et al. 2014; Chowdhary et al. 2014; Hölker et al. 2004; Zhao et al. 2010;

Kejžar et al. 2013). JES_119, as it develops over time, increases its melanin production and

develops from a bright orange-yellow color to a more deeply pigmented black. The budding cells

are unilateral and are uncommon in other described MCF, and it was evaluated as a previously

undescribed genus and species in Class *Dothideomycetes* as *Taxawa tesnikishii.* Strain JES_115

has a phenotype that is commonly seen in MCF, but due to its complicated taxonomy history, the

genus *Coniosporium* remains as a genus of doubtful identity. The species JES_112 is one of the

few described *Neophaeococcomyces* available anywhere and thus an important contribution to

the scientific community.

Finding MCF from extreme environments such as BSCs may point to harbored secondary

metabolites either absorbed from surrounding species through horizontal gene transfer (HGT) or

adapted from time. MCF are known to use secondary metabolites to help survive such extremes

and are thus an untapped resource for the identification of novel metabolites. An example to be

described by Dr. Erin Carr, *Neophaeococcomyces* sp, has been shown to have an obligate

bacterial hitchhiker (Carr 2022). JES_115 and JES_119 both contain about 15 secondary

metabolites detected by antiSMASH, which is a considerable number when compared to other

fungi (Rateb and Ebel 2011). The roles of secondary metabolites in modern human industry

applications are vast (Demain 2014) and thus would be important to test and compare across all

MCF.

568
Figure 4. Chaetothyriales multi-locus gene tree. Multi-loci phylogenetic tree using 4 highly-conserved
genes including, beta-tubulin, calmodulin, internal-transcribed sequence (ITS), and the large subunit of
the Ribosome (LSU). Seven representative strains with loci were used, which were aligned, trimmed, and
concatenated to run the IQTree program and generate a concise multi-loci gene tree for
Neophaeococcomyces mojaviensis JES112. Here this strain is placed in the *Neophaeococcomyces* genera.
574

576
Figure 5. Dothideomycetes multi-locus gene tree. Multi-loci phylogenetic tree using 4 highly-
conserved genes including, beta-tubulin, calmodulin, internal-transcribed sequence (ITS), and the large
subunit of the Ribosome (LSU). Twenty-five representative strains with loci were used, which were
aligned, trimmed, and concatenated to run the IQTree program and generate a concise multi-loci gene tree
for JES_115 and JES_119 or *Coniosporium tulheliwenetii* and *Taxawa tesnikishii* respectively. Here
strain JES_115 is grouping with other *Coniosporium* spp., while JES_119 is grouping with but distinctly
apart from *Aureobasidium* spp.

584
585
Table 4. Gene Tree Nucleotide Accession numbers. Collected accession numbers of alternative strains
to build Figures 4 and 5. The remaining empty spaces indicate the need to pull the sequences using
BLAST. Once collected each gene populated a gene FASTA file, aligned, clipped and IQTREE2, a tree-
building program, was run to determine the gene models for each gene. Once determined, the gene
alignment lengths were calculated and partitioning scripts were described and used to build a larger tree,
again using IQTREE2.

Species	Beta-Tubulin	Calmodulin	Internal-Transcribed Spacer	Large Subunit (Ribosomal)
*Neophaeococcomyces mojaviensis*JES112	-	-	-	-
Taxawa tesnikishii JES119	-	-	-	-
Alternaria solani NL03003	MK388240.1	MH243808.1	NR_172408.1	ON416546.1
Anthracina ramosa CGMCC3.16372	KP226556.1	-	NR_172152.1	KP174923.1
Anthracina saxicola CGMCC3.17315	KP226554.1	-	NR_172151.1	KP174921.1
Arthrocladium caudatum CBS457.67	LT558710.1	-	NR_145398.1	MH870732.1
Arthrocladium tardum CBS 127021	-	-	NR_154706.1	NG_057089.1
Arthrocladium tropicale CBS 134926	KX822406.1	-	NR_154724.1	NG_057119.1
Aureobasidium mangrovei IBRCM30265	XM_029904852.1	XM_041025809.1	NR_174637.1	NG_078639.1
*Aureobasidium sp.*AWRI4619-CO-2020	XM_029904852.1	XM_041025809.1	MK256457.1	MK968773.1
Baudoinia panamericana UAMH10762	XM_007673885.1	XM_007675264.1	NR_153615.1	KT186497.1
Bradymyces alpinus CCFEE 5493	LN589970.1	-	NR_132844.1	NG_058641.1
Bradymyces yunnanensis CGMCC 3.17350	KP226545.1	-	KP174865.1	KP174926.1
Capronia coronata CBS 617.96	XM_007728982.1	XM_007721556.1	NR_154745.1	NG_067831.1

Cenococcum geophilum 1.58	-	-	**KC967410.1**	**MN817694.1**
Cladophialophora carrionii CBS 260.83	**KF928582.1**	**XM_008724000.1**	**MH861582.1**	**MH873312.1**
Cladophialophora_yegresii CBS 114405	**EU137209.1**	-	**NR_111284.1**	**KX822323.1**
Cladophialophora immunda CBS 834.96	**EU137203.1**	-	**NR_111283.1**	**MH874242.1**
Coniosporium apollinis CBS 100218	-	**XM_007785495.1**	**NR_159787.1**	**GU250898.1**
Coniosporium uncinatum CBS 100219	-	-	**NR_145343.1**	**NG_058806.1**
Coniosporium uncinatum CCFEE 6149	-	-	**MZ573424.1**	**MZ573424.1**
Cyphellophora aestiva CBS 239.91	**MW297547.1**	-	**MH861291.1**	**MH873056.1**
Cyphellophora laciniata CBS 239.91	**KF928604.1**	-	**NR_121335.1**	**MH873933.1**
Cyphellophora sessilis KNU-002	**KX431943.1**	-	**KX431941.1**	**KX431942.1**
Cyphellophora vermispora CBS 228.86	**KC455227.1**	-	**KC455244.1**	**KC455257.1**
Exophiala dermatitidis NIH.UT8656	-	**XM_009160450.1**	**JH226131.1**	**ON543213.1**
Exophiala oligosperma CBS 265.49	**EF551507.1**	**XM_016404560.1**	**MH856519.1**	**MH868049.1**
Exophiala oligosperma CBS 127587	**KF928570.1**	-	**MH864631.1**	**MH876068.1**
Exophiala radicis P2854	**KT723463.1**	-	**NR_158397.1**	**NG_069319.1**

Exophiala xenobiotica CBS 118157	NW_01356249 0.1	XM_01345465 6.1	OP132917.1	MH875251.1
Fonsecaea pedrosoi CBS 271.37	XM_0134308 09.1	XM_01342686 4.1	MT023597.1	KF155185.1
Friedmanniomyces endolithicus CCFEE 5311	-	-	NR_144917.1	GU250367.1
Friedmanniomyces simplex CCFEE 5184	-	-	NR_155079.1	GU250368.1
Hortaea werneckii EXF-2000	-	-	NR_145338.1	NG_057773.1
Knufia petricola MA5789	-	-	NR_111871.1	MT126669.1
Knufia mediterranea CCFEE 6211	-	-	KP791793.1	KR781080.1
Knufia mediterranea CCFEE 6205	-	-	KP791794.1	KR781081.1
Knufia separata CGMCC 3.17337	KP226564.1	-	NR_172233.1	KP174931.1
Knufia tsunedae FMR10621	-	-	NR_132842.1	HG003672.1
Knufia walvisbayicola CBS 146989	MZ078266.1	-	NR_173046.1	NG_076737.1
Knufia sp. MV-2018 MLT-8	-	-	MT636977.1	MT636937.1
Leptosphaeria biglobosa CA1	AY749005.1	-	NR_111620.1	JX681093.1
Leptosphaeria maculans JN3	AY749012.1	XM_00384071 9.1	M96384.1	JF740306.1
Lithohypha aloicola CPC35996	MN556837.1	-	NR_166313.1	NG_068316.1
Lithohypha guttulata FMR 20100	OX346320.1	-	OX346376.1	OX346377.1

Metulocladosporiella musicola CBS 110960	-	**MG934537.1**	**MH862870.1**	**NG_067447.1**
Metulocladosporiella samutensis CBS 143921	-	**MG934540.1**	**NR_161056.1**	**MG934453.1**
Neohortaea acidophila CBS 113389	**XM_03373332 87.1**	**XM_03373393 0.1**	**NR_169648.1**	**OL739260.1**
Neophaeococcomyces aloes FJII-L3-CM-P2	-	-	**NR_132069.1**	**KF777234.1**
Neophaeococcomyces catenatus CBS 650.76	-	-	**AF050277.1**	**MH872793.1**
Neophaeococcomyces oklahomaensis EMSL3313	-	-	**MW798187.1**	**MW798182.1**
*Phaeosphaeria sp.*MPI-PUGE-AT-0046c	-	-	**NR_155675.1**	**NG_068582.1**
Polychaeton citri CBS 116435	-	-	**GU214649.1**	**OM238161.1**
Polyplosphaeria fusca CBS 125425	-	-	**NR_119397.1**	**MH875147.1**
Salinomyces thailandica CCFEE 6315	-	-	**NR_171710.1**	**MH875167.1**
Scorias spongiosa HLSS1	-	-	**NR_154422.1**	**MH866910.1**
Strelitziana africana ICMP 21758	-	-	**MK210538.1**	**MK210499.1**
Strelitziana africana ICMP 21760	-	-	**MK210540.1**	**MK210501.1**
Teratosphaeria gauchensis CMW 17545	-	**KF902726.1**	**NR_144976.1**	**EU019295.2**
Trichomerium cicatricatum CGMCC3.17307	**KP226540.1**	-	**NR_172232.1**	**KP174947.1**

Trichomerium deniqulatum MFLUCC10-0884	-	-	**NR_132965.1**	**NG_059479.1**
Trichomerium gleosproum MFLUCC10-0087	-	-	**JX313656.1**	**JX313662.1**
Trichomerium sp. XDY-2021a-IFRDCC3302	-	-	**MN901488.1**	**MZ822131.1**
Tuber melanosporum	**MZ458420.1**	**XM_002841184.1**	**GQ917052.1**	**JF807486.1**
Venturia pyrina ICMP11032	**EU853839.1**	-	**NR_169905.1**	**AY849967.1**
Zalaria obscura FKI-L7-BK-DRAB2	-	-	**NR_153466.1**	**MT103097.1**
Zymoseptoria tritici CBS 100329	**XM_003856727.1**	**KF254299.1**	**KP852517.1**	**MK788267.1**

Conclusions

Biological soil crusts are an untapped resource for novel microorganisms as evidenced by our

work. The MCF we collected are incredibly slow growing and could be easily missed during

culturing attempts. The method described here (and detailed in **Appendix 1**) improved the

chances for isolation of these species. MCFs have been isolated from different types of extreme

environments including BSC's.

An interesting consideration for species JES_112 or *Neophaeococcomyces mojaviensis* is

through personal communications with another MCF scientist, Erin Carr, who also isolated a

species of *Neophaococcomyces*, named "Crusty" from a biological soil crust. Their species was

associated with a pink *Methylobacterium* symbiont, which she calls "Light Pinky"and "Dark

Pinky". Her assessment of this symbiosis suggests the role of the symbiont is involved in aerobic

anoxygenic photosynthesis and auxin production (Carr 2022). Through our multiple culturing

attempts, there was no indication of a pink symbiont, but the possible bacteria signal within our

secondary metabolite assessment could be attributed to a contamination from this bacterium.

JES_112 will be the second known genome of the genus *Neophaeococcomyces* found in

GenBank as JAPDRQ000000000.

Species JES_119 or *Taxawa tesnikishi* is the first isolated of this phenotype. This species was

closer to *Aureobasidium*, yet phenotypically is very different from identified *Aureobasidium*

species. Using fungal phenotype and phylogenomics, we were able to determine the

microorganism is different from known *Aureobasidium* species and may be of a new genus,

hence the naming *Taxawa* as its genus name. *Taxawa tesnikishi* is the first known genus and

species, now available in Genbank as JAPDRO000000000.

Species JES_115 or *Coniosporium tulheliwenetii* was also collected from the BSC's from Boyd

Deep Canyon and was thus honored with a Cahuilla term used to describe BSC phenotypic

features seen on the soil. The genus *Coniosporium* has historically been confusing, as the earliest

strains were collected in the late 1800s. Initial collections of *Coniosporium* were placed in

Chaetothyriales, while later similar MCF phenotypes were also grouped with *Coniosporium*, to

be later considered as Dothideomycetes instead. A careful reorganization and assessment of the

genus is desperately needed. Genome assembly is now available as JAPDRP000000000.

Acknowledgments

This work was performed in part at the University of California Natural Reserve System (Boyd

Deep Canyon Desert Research Center) Reserve DOI: (doi:10.21973/N3V66D) (UCNRS

Information Manager 1965) and at Joshua Tree National Park under permits generated through

the help and guidance of Nuttapon Pombubpa, permit numbers: JOTR-2015-SCI-0033, JOTR-

2016-SCI-0036, JOTR-2017-SCI-0044, JOTR-2018-SCI-0035, JOTR-2019-SCI-0031. Many

thanks to Nicole Pietrasiak for her extensive knowledge in biological soil crusts. Many thanks

also to Joshua Tree Reserve members who helped facilitate the admission of our Joshua Tree

sample, specifically Anna R Tegarden, Ann Hitchcock, and Jay Goodwin.

638
639 **Funding**

640
641 T.K was partially supported by NIH grant R01 AI127548 (support to J.E.S. via Deborah Hogan).

642 Computational analyses were performed at the University of California – Riverside HPCC,

643 supported by grants from the National Science Foundation (DBI – 1429826 and 2215705) and

644 NIH (S10OD016290). JES is a CIFAR Fellow in the Fungal Kingdom: Threats and

645 Opportunities program.

646
647 **References:**

649 Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI,
650 Pham S, Prjibelski AD, et al. 2012. SPAdes: a new genome assembly algorithm and its applications to
651 single-cell sequencing. J Comput Biol. 19(5):455–477.

652
653 Bell AA, Wheeler MH. 1986. Biosynthesis and Functions of Fungal Melanins. Annu Rev Phytopathol.
654 24(1):411–451.

655
656 Belnap J, Harper KT. 1995. Influence of cryptobiotic soil crusts on elemental content of tissue of two
657 desert seed plants. Arid Soil Research and Rehabilitation. 9(2):107–115.

658
659 Beraldi-Campesi H, Retallack GJ. 2016b. Terrestrial Ecosystems in the Precambrian. In: Weber B, Büdel
660 B, Belnap J, editors. Biological Soil Crusts: An Organizing Principle in Drylands. Cham: Springer
661 International Publishing. p. 37–54.

662
663 Blin K, Shaw S, Kloosterman AM, Charlop-Powers Z, van Wezel GP, Medema MH, Weber T. 2021.
664 antiSMASH 6.0: improving cluster detection and comparison capabilities. Nucleic Acids Res.
665 49(W1):W29–W35.
666

667 Brown J, Boyd J. 1922. History of San Bernardino and Riverside Counties: With Selected Biography of
668 Actors and Witnesses of the Period of Growth and Achievement. Western Historical Association.
669
670 Brůna T, Lomsadze A, Borodovsky M. 2020. GeneMark-EP : eukaryotic gene prediction with self-
671 training in the space of genes and proteins. NAR Genomics and Bioinformatics. 2(2).
672 doi:10.1093/nargab/lqaa026. http://dx.doi.org/10.1093/nargab/lqaa026.
673
674 Bushnell B. 2014. BBMap: A fast, accurate, splice-aware aligner. Lawrence Berkeley National Lab.
675 (LBNL), Berkeley, CA (United States) Report No.: LBNL-7065E. https://www.osti.gov/biblio/1241166.
676
677 Carr EC. 2022. Discovering Novel Polyextremotolerant Fungi, and Determining Their Ecological Role
678 Within the Biological Soil Crust Consortium.
679
680 Casadevall A, Rosas AL, Nosanchuk JD. 2000. Melanin and virulence in *Cryptococcus neoformans*.
681 Current Opinion in Microbiology. 3(4):354–358. doi:10.1016/s1369-5274(00)00103-x.
682
683 Casadevall A, Steenbergen JN, Nosanchuk JD. 2003. "Ready made" virulence and "dual use" virulence
684 factors in pathogenic environmental fungi — the *Cryptococcus neoformans* paradigm. Current Opinion in
685 Microbiology. 6(4):332–337. doi:10.1016/s1369-5274(03)00082-1.
686
687 Chomnunti P, Bhat DJ, Jones EBG, Chukeatirote E, Bahkali AH, Hyde KD. 2012. Trichomeriaceae, a
688 new sooty mould family of Chaetothyriales. Fungal Divers. 56(1):63–76.
689
690 Chowdhary A, Perfect J, de Hoog GS. 2014. Black Molds and Melanized Yeasts Pathogenic to Humans.
691 Cold Spring Harb Perspect Med. 5(8):a019570.
692
693 Ciferri R. 1958. Affinity between fungi of the verrucous dermatitis and sooty-molds. Mycopathol Mycol
694 Appl. 9(3):197–200.
695
696 Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, Friedberg I, Hamelryck T, Kauff F,
697 Wilczynski B, et al. 2009. Biopython: freely available Python tools for computational molecular biology
698 and bioinformatics. Bioinformatics. 25(11):1422–1423.
699
700 Dadachova E, Bryan RA, Howell RC, Schweitzer AD, Aisen P, Nosanchuk JD, Casadevall A. 2008. The
701 radioprotective properties of fungal melanin are a function of its chemical composition, stable radical
702 presence and spatial arrangement. Pigment Cell Melanoma Res. 21(2):192–199.
703
704 Dadachova E, Casadevall A. 2008. Ionizing radiation: how fungi cope, adapt, and exploit with the help of
705 melanin. Curr Opin Microbiol. 11(6):525–531.
706
707 Demain AL. 2014. Valuable Secondary Metabolites from Fungi. In: Martín J-F, García-Estrada C,
708 Zeilinger S, editors. Biosynthesis and Molecular Genetics of Fungal Secondary Metabolites. New York,
709 NY: Springer New York. p. 1–15.
710

711 Evans RD, Ehleringer JR. 1993. A break in the nitrogen cycle in aridlands? Evidence from δp15N of
712 soils. Oecologia. 94(3):314–317.
713
714 Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, Heger A, Hetherington K, Holm L,
715 Mistry J, et al. 2014. Pfam: the protein families database. Nucleic Acids Res. 42(Database issue):D222–
716 30.
717
718 Gilchrist CLM, Chooi Y-H. 2021 Jan 18. Clinker & clustermap.js: Automatic generation of gene cluster
719 comparison figures. Bioinformatics. doi:10.1093/bioinformatics/btab007.
720
721 Gómez BL, Nosanchuk JD. 2003. Melanin and fungi. Curr Opin Infect Dis. 16(2):91–96.
722
723 Gorbushina AA, Whitehead K, Dornieden T, Niesse A, Schulte A, Hedges JI. 2003. Black fungal
724 colonies as units of survival: hyphal mycosporines synthesized by rock-dwelling microcolonial fungi. Can
725 J Bot. 81(2):131–138.
726
727 Gostinčar C, Sun X, Černoša A, Fang C, Gunde-Cimerman N, Song Z. 2022. Clonality, inbreeding, and
728 hybridization in two extremotolerant black yeasts. Gigascience. 11. doi:10.1093/gigascience/giac095.
729
730 Green TGA, Allan Green TG, Proctor MCF. 2016. Physiology of Photosynthetic Organisms Within
731 Biological Soil Crusts: Their Adaptation, Flexibility, and Plasticity. Biological Soil Crusts: An
732 Organizing Principle in Drylands.:347–381. doi:10.1007/978-3-319-30214-0_18.
733
734 Green TGA, Broady PA. 2001. Biological soil crusts of Antarctica. In: Ecological Studies. Berlin,
735 Heidelberg: Springer Berlin Heidelberg. (Ecological studies: analysis and synthesis. Berlin, Heidelberg,
736 New York NY). p. 133–139.
737
738 Gümral R, Tümgör A, Saraçlı MA, Yıldıran ŞT, Ilkit M, de Hoog GS. 2014. Black yeast diversity on
739 creosoted railway sleepers changes with ambient climatic conditions. Microb Ecol. 68(4):699–707.
740
741 Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, White O, Buell CR, Wortman JR. 2008.
742 Automated eukaryotic gene structure annotation using EVidenceModeler and the Program to Assemble
743 Spliced Alignments. Genome Biol. 9(1):R7.
744
745 Haridas S, Albert R, Binder M, Bloem J, LaButti K, Salamov A, Andreopoulos B, Baker SE, Barry K,
746 Bills G, et al. 2020. 101 Dothideomycetes genomes: A test case for predicting lifestyles and emergence of
747 pathogens. Stud Mycol. 96:141–153.
748
749 Hölker U, Bend J, Pracht R, Tetsch L, Müller T, Höfer M, de Hoog GS. 2004. *Hortaea acidophila*, a new
750 acid-tolerant black yeast from lignite. Antonie Van Leeuwenhoek. 86(4):287–294.
751
752 de Hoog GS, Hermanides-Nijhof EJ. 1977. The Black Yeasts and Allied Hyphomycetes, Studies in
753 Mycology. Baarn, The Netherlands: Centraalbureau voor Schimmelcultures.
754

de Hoog GS, Matos T, Sudhadham M, Luijsterburg KF, Haase G. 2005. Intestinal prevalence of the neurotropic black yeast *Exophiala (Wangiella) dermatitidis* in healthy and impaired individuals. Mycoses. 48(2):142–145.

Hoog GS de, Queiroz-Telles F, Haase G, Fernandez-Zeppenfeldt G, Angelis DA, H. G. Gerrits van den Ende A, Matos T, Peltroche-Llacsahuanga H, Pizzirani-Kleiner AA, Rainer J, et al. 2000a. Black fungi: clinical and pathogenic approaches. Med Mycol. 38(sup1):243–250.

Hooper L. 1920. The Cahuilla Indians. University of California Press.

Housman DC, Yeager CM, Darby BJ, Sanford RL, Kuske CR, Neher DA, Belnap J. 2007. Heterogeneity of soil nutrients and subsurface biota in a dryland ecosystem. Soil Biol Biochem. 39(8):2138–2149.

Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, Mende DR, Letunic I, Rattei T, Jensen LJ, et al. 2019. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. Nucleic Acids Res. 47(D1):D309–D314.

James TY, Stajich JE, Hittinger CT, Rokas A. 2020. Toward a Fully Resolved Fungal Tree of Life. Annu Rev Microbiol. 74:291–313.

Johnson L. 2008. Iron and siderophores in fungal–host interactions. Mycol Res. 112(2):170–183.

Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A, Nuka G, et al. 2014. InterProScan 5: genome-scale protein function classification. Bioinformatics. 30(9):1236–1240.

Kejžar A, Gobec S, Plemenitaš A, Lenassi M. 2013. Melanin is crucial for growth of the black yeast *Hortaea werneckii* in its natural hypersaline environment. Fungal Biology. 117(5):368–379. doi:10.1016/j.funbio.2013.03.006. http://dx.doi.org/10.1016/j.funbio.2013.03.006.

Korf I. 2004. Gene finding in novel genomes. BMC Bioinformatics. 5:59.

Kroeber AL. 1925. Handbook of the Indians of California. U.S. Government Printing Office.

Kurbessoian T. 2022. tania-k/Chapter2.3MCF: V1. https://zenodo.org/record/7259500.

Lal R. 2004b. Carbon sequestration in dryland ecosystems. Environ Manage. 33(4):528–544.

Lanfear R, Calcott B, Kainer D, Mayer C, Stamatakis A. 2014. Selecting optimal partitioning schemes for phylogenomic datasets. BMC Evol Biol. 14:82.

Little AEF, Currie CR. 2008. Black yeast symbionts compromise the efficiency of antibiotic defenses in fungus-growing ants. Ecology. 89(5):1216–1222.

Lowe TM, Chan PP. 2016. tRNAscan-SE On-line: integrating search and context for analysis of transfer
RNA genes. Nucleic Acids Res. 44(W1):W54–7.

Maestre FT, Soliveres S, Gotelli NJ, Quero JL, Berdugo M. 2012. Response to Comment on "Plant
Species Richness and Ecosystem Multifunctionality in Global Drylands." Science. 337(6091):155–155.
doi:10.1126/science.1220620. http://dx.doi.org/10.1126/science.1220620.

Majoros WH, Pertea M, Salzberg SL. 2004. TigrScan and GlimmerHMM: two open source ab initio
eukaryotic gene-finders. Bioinformatics. 20(16):2878–2879.

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update: Novel and
Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of
Eukaryotic, Prokaryotic, and Viral Genomes. Mol Biol Evol. 38(10):4647–4654.

Mehdiabadi NJ, Schultz TR. Natural history and phylogeny of the fungus-farming ants (Hymenoptera:
Formicidae: Myrmicinae: Attini). [accessed 2022 Mar 29].

Metin B, Döğen A, Yıldırım E, de Hoog GS, Heitman J, Ilkit M. 2019. Mating type (MAT) locus and
possible sexuality of the opportunistic pathogen *Exophiala dermatitidis*. Fungal Genetics and Biology.
124:29–38. doi:10.1016/j.fgb.2018.12.011.

Nosanchuk JD, Stark RE, Casadevall A. 2015. Fungal melanin: What do we know about structure? Front
Microbiol. 6:1463.

Palmer JM, Stajich J. 2020. Funannotate v1.8.1: Eukaryotic genome annotation.
https://zenodo.org/record/4054262.

Pietrasiak N. 2005. Field guide to classify biological soil crusts for ecological site evaluation. NRCS.

Pietrasiak N, Regus JU, Johansen JR, Lam D, Sachs JL, Santiago LS. 2013. Biological soil crust
community types differ in key ecological functions. Soil Biol Biochem. 65:168–171.

Potter SC, Luciani A, Eddy SR, Park Y, Lopez R, Finn RD. 2018. HMMER web server: 2018 update.
Nucleic Acids Res. 46(W1):W200–W204.

Rateb ME, Ebel R. 2011. Secondary metabolites of fungi from marine habitats. Nat Prod Rep. 28(2):290–
344.

Rawlings ND, Barrett AJ, Thomas PD, Huang X, Bateman A, Finn RD. 2018. The MEROPS database of
proteolytic enzymes, their substrates and inhibitors in 2017 and a comparison with peptidases in the
PANTHER database. Nucleic Acids Res. 46(D1):D624–D632.

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other things).
Methods Ecol Evol. 3(2):217–223.

Robinson SL, Christenson JK, Wackett LP. 2019. Biosynthesis and chemical diversity of β-lactone
natural products. Nat Prod Rep. 36(3):458–475.

Rozenstein O, Karnieli A. 2015. Identification and characterization of Biological Soil Crusts in a sand
dune desert environment across Israel–Egypt border using LWIR emittance spectroscopy. J Arid Environ.
112:75–86.

Sass G, Ansari SR, Dietl A-M, Déziel E, Haas H, Stevens DA. 2019. Intermicrobial interaction:
*Aspergillus fumigatus*siderophores protect against competition by *Pseudomonas aeruginosa*. PLoS One.
14(5):e0216085.

Saunte DM, Tarazooie B, Arendrup MC, de Hoog GS. 2012a. Black yeast-like fungi in skin and nail: it
probably matters. Mycoses. 55(2):161–167.

Siletti CE, Zeiner CA, Bhatnagar JM. 2017. Distributions of fungal melanin across species and soils. Soil
Biol Biochem. 113:285–293.

Slater GSC, Birney E. 2005. Automated generation of heuristics for biological sequence comparison.
BMC Bioinformatics. 6:31.

SMIT A. F. A. 2004. Repeat-Masker Open-3.0. http://www.repeatmasker.org.

Sofi MY, Shafi A, Masoodi KZ. 2022. NCBI BLAST. Bioinformatics for Everyone.:95–102.
doi:10.1016/b978-0-323-91128-3.00021-5.

Staley JT, Palmer F, Adams JB. 1982. Microcolonial fungi: common inhabitants on desert rocks?
Science. 215(4536):1093–1095.

Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B. 2006. AUGUSTUS: ab initio
prediction of alternative transcripts. Nucleic Acids Res. 34(Web Server issue):W435–9.

Steenbergen JN, Casadevall A. 2003. The origin and maintenance of virulence for the human pathogenic
fungus Cryptococcus neoformans. Microbes Infect. 5(7):667–675.

Stevens TO. 1995. Optimization of media for enumeration and isolation of aerobic heterotrophic bacteria
from the deep terrestrial subsurface. J Microbiol Methods. 21(3):293–303.

Taborda CP, da Silva MB, Nosanchuk JD, Travassos LR. 2008b. Melanin as a virulence factor of
Paracoccidioides brasiliensis and other dimorphic pathogenic fungi: a minireview. Mycopathologia.
165(4-5):331–339.

Teixeira MM, Moreno LF, Stielow BJ, Muszewska A, Hainaut M, Gonzaga L, Abouelleil A, Patané JSL,
Priest M, Souza R, et al. 2017. Exploring the genomic diversity of black yeasts and relatives
(Chaetothyriales, Ascomycota). Stud Mycol. 86:1–28.
889
890 Thambugala KM, Ariyawansa HA, Li Y-M, Boonmee S, Hongsanan S, Tian Q, Singtripop C, Bhat DJ,
891 Camporesi E, Jayawardena R, et al. 2014. Dothideales. Fungal Divers. 68(1):105–158.
892
893 Thanh VN, Hien DD. 2019. *Moniliella floricola* sp. nov., a species of black yeast isolated from flowers.
894 International Journal of Systematic and Evolutionary Microbiology. 69(1):87–92.
895 doi:10.1099/ijsem.0.003099. http://dx.doi.org/10.1099/ijsem.0.003099.
896
897 UCNRS Information Manager. 1965. Boyd deep canyon desert research center. doi:10.21973/N3V66D.
898
899 Vicente VA, Attili-Angelis D, Pie MR, Queiroz-Telles F, Cruz LM, Najafzadeh MJ, de Hoog GS, Zhao J,
900 Pizzirani-Kleiner A. 2008. Environmental isolation of black yeast-like fungi involved in human infection.
901 Stud Mycol. 61:137–144.
902
903 Vogt E, Künzler M. 2019. Discovery of novel fungal RiPP biosynthetic pathways and their application
904 for the development of peptide therapeutics. Appl Microbiol Biotechnol. 103(14):5567–5581.
905
906 Wang L-G, Lam TT-Y, Xu S, Dai Z, Zhou L, Feng T, Guo P, Dunn CW, Jones BR, Bradley T, et al.
907 2020. Treeio: An R Package for Phylogenetic Tree Input and Output with Richly Annotated and
908 Associated Data. Mol Biol Evol. 37(2):599–603.
909
910 Warren SD, Clair LL St, Stark LR, Lewis LA, Pombubpa N, Kurbessoian T, Stajich JE, Aanderud ZT.
911 2019. Reproduction and dispersal of biological soil crust organisms. Front Ecol Evol. 7.
912 doi:10.3389/fevo.2019.00344.
913
914 Weber B, Bowker M, Zhang Y, Belnap J. 2016. Natural Recovery of Biological Soil Crusts After
915 Disturbance. In: Weber B, Büdel B, Belnap J, editors. Biological Soil Crusts: An Organizing Principle in
916 Drylands. Cham: Springer International Publishing. p. 479–498.
917
918 Wheeler MH. 1983. Comparisons of fungal melanin biosynthesis in ascomycetous, imperfect and
919 basidiomycetous fungi. Trans Br Mycol Soc. 81(1):29–36.
920
921 White PM. 2001. California Indians and Their Reservations: An Online Dictionary. San Diego State
922 University Library.
923
924 Wick RR, Judd LM, Holt KE. 2019. Performance of neural network basecalling tools for Oxford
925 Nanopore sequencing. Genome Biol. 20(1):129.
926
Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. 2017. Ggtree : An r package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. Methods Ecol Evol.
8(1):28–36.

Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, Busk PK, Xu Y, Yin Y. 2018. dbCAN2: a meta

server for automated carbohydrate-active enzyme annotation. Nucleic Acids Res. 46(W1):W95–W101.

Zhao J, Zeng J, de Hoog GS, Attili-Angelis D, Prenafeta-Boldú FX. 2010. Isolation and identification of

black yeasts by enrichment on atmospheres of monoaromatic hydrocarbons. Microb Ecol. 60(1):149–156.

Zimin AV, Marçais G, Puiu D, Roberts M, Salzberg SL, Yorke JA. 2013. The MaSuRCA genome

assembler. Bioinformatics. 29(21):2669–2677. doi:10.1093/bioinformatics/btt476.

http://dx.doi.org/10.1093/bioinformatics/btt476.

941