Description of new micro-colonial fungi species *Neophaeococcomyces mojavensis,*
***Coniosporium tulheliwenetii, and Taxawa tesnikishii* cultured from biological soil**
crusts

Tania Kurbessoian[1], Sarah A. Ahmed[2], Yu Quan[2], Sybren de Hoog[2], Jason E. Stajich [1,*]
[1]Department of Microbiology and Plant Pathology and Institute of Integrative Genome Biology,
University of California, 92521, Riverside, CA, USA
[2]Radboudumc/CWZ Ceter of Expertise for Mycology, Nijmegen, 6525GA Nijmegen, The Netherlands

*Correspondence: jason.stajich@ucr.edu; Tel.: +1 951.827.2363 (J.E.S.)

10
11

Abstract

Black yeasts and relatives comprise Micro-Colonial Fungi (MCFs) which are slow-growing stress-

tolerant micro-eukaryotes that specialize in extreme environments. MCFs are paraphyletic and

found in the Orders *Chaetothyriales* (*Eurotiomycetes*) and *Dothideales* (*Dothidiomycetes*). We

have isolated and described three new MCFs species from desert biological soil crusts (BSCs)

collected from two arid land regions: Joshua Tree National Park (Mojave Desert) and UC Natural

Reserve at Boyd Deep Canyon (confluence of Mojave and Sonoran Deserts). BSCs are composite

assemblages of cyanobacteria, eukaryotic algae, fungi, lichens, and bryophytes embedded into the

surface of desert soils, providing a protective buffer against the harsh desert environment. Our

work focused on one type of desert BSC, the cyanolichen crust dominated by *Collema sp.* Using

culture-dependent protocols, three MCFs were axenically isolated from their respective samples

along with the extracted DNA. Their genomes were sequenced using Illumina and Nanopore, and

finally assembled and annotated using hybrid assembly approaches and established bioinformatics

pipelines to conduct final taxonomic phylogenetic analysis and placement. The three species

described here are unique specimen from desert BSCs, here we introduce, *Neophaeococcomyces*

mojavensis (Chaetothyriales), *Cladosporium tulheliwenetii (Dothideales)*, and *Taxawa tesnikishii*

(*Dothideales*).

Introduction

Biological Soil Crusts

Dryland ecosystems constitute some of the largest terrestrial biomes, collectively covering 47.2%

of Earth's land surface (6.15 billion hectares) (Lal 2004). Drylands are expanding due to climate

change and population pressures, while also supporting over 38% of the global human population

(Lal 2004; Maestre et al. 2012). Biological Soil Crusts (BSCs) cover up to 70% of dryland

ecosystems (Rozenstein and Karnieli 2015). They are an ancient feature of dryland ecosystems,

with the earliest known fossil records indicating nearly 2.6 billion years ago (Beraldi-Campesi and

Retallack 2016). The initial phases of BSC formation in drylands are the stabilization of the soil

surface by filamentous cyanobacteria, followed by the colonization of lichens and bryophytes

(Weber et al. 2016). Within dryland ecosystems, the carbon and nitrogen dynamics are often

described as the "pulse-reserve model" where precipitation events happen sporadically,

stimulating intermittent biological activity which generates biomass and organic matter. After

extended periods of heat and dryness, the biological activity goes into reserve and becomes

dormant again (Pietrasiak et al. 2013). BSCs also increase soil fertility by increasing soil stability

and adding fixed atmospheric carbon and nitrogen to the soil underneath, allowing for greater

complex microbial and micro-faunal populations (Housman et al. 2007).

Within BSCs is a consortium of cyanobacteria, algae, diazotrophic bacteria, lichen, fungi, and

bryophytes that colonize and stabilize the soil surfaces against natural forces like wind and water

(Warren et al. 2019). These microorganisms recruit partners to create a mutualistic symbiotic

relationship which helps maintain a self-sustaining micro-ecosystem. Phototrophic organisms fix

carbon that is later used by surrounding micro-organisms as an energy source (Green et al. 2016).

The presence of chemoheterotrophic bacteria and cyanobacteria indicates that BSCs are capable

of fixing atmospheric nitrogen to increase levels of fixed nitrogen for nearby living vascular plants

(Belnap and Harper 1995; Evans and Ehleringer 1993). BSCs have been shown to reduce surface

erodibility as filaments of cyanobacterial sheath material lipopolysaccharides (LPS) entangle

surface particles and create a crust that is more resistant to entrainment than the layers below (Lal

2004).

60
Different ecological sites promote different crust taxa. BSCs are classified into types based on the

dominant photosynthetic microorganism in the crust which includes: cyanobacteria, algae, lichens,

or mosses to produce Cyano-Lichenized Crust, Light Algal Crusts, Dark Algal Crusts, Lichenized

Crust, Rough Moss Crust and Smooth Moss Crust (Pietrasiak 2005). The community types are

classified based on dominant taxa and their distinct morphologies (Pietrasiak et al. 2013). BSC's

main roles include: 1) mediating almost all inputs and outputs (gasses, nutrients, water) to and

from the strata above and below the surface, 2) being the zone of high nutrient deposition,

transformation, and availability, 3) structuring temporal, spatial, and compositional aspects of the

surrounding vascular plant community, and 4) facilitating the direct delivery of carbon, nutrients,

and water by BSCs from the soil interspace to nearby vascular plants.

Micro-colonial Fungi

'Black yeasts and relatives' comprise a large amalgamate of melanized fungi. Among these are

micro-colonial fungi (MCF) or rock-inhabiting fungi (RIF), prevalent names used to describe a

special group of melanized fungi. Micro-colonial fungi are ubiquitous and found in the most

extreme environments. They are present in widely divergent habitats, such as on the indoor rubbers

of dishwashers, in oil deposits, on the surfaces of Mediterranean monuments, on the radioactive

walls of the Chernobyl Station, in mutilating skin infections in third-world countries, in human

superficial and invasive infections, or colonizing lungs of cystic fibrosis patients (de Hoog et al.

2005; Hoog et al. 2000; Vicente et al. 2008; Gümral et al. 2014; Saunte et al. 2012; Thanh and

Hien 2019; Mehdiabadi and Schultz ; Little and Currie 2008; de Hoog and Hermanides-Nijhof

1977). MCFs are compelling not only because of where we can discover them and their unique

physiological features, but also because of how little we know about them.

85
MCFs have three main characteristics: meristematic growth, presence of 1,8-DHN melanin, and

membrane-associated carotenoids and intracellular mycosporine-like amino acids (Staley et al.

1982; Gorbushina et al. 2003). These features distinguish them from other fungi, yet

phylogenetically they fall into two paraphyletic Ascomycota clades, the *Dothideomycetes* and

Chaetothyriomycetes. Melanin is a highly complex secondary metabolite that is produced by many

organisms through different evolutionary lineages (Siletti et al. 2017; Wheeler 1983). Most

importantly in MCFs, it has been used as a virulence factor as a non-specific armor against host

immune system to infect its host (Steenbergen and Casadevall 2003; Casadevall et al. 2003;

Casadevall et al. 2000; Nosanchuk et al. 2015; Taborda et al. 2008; Gómez and Nosanchuk 2003).

Available research indicates melanin has other functional roles, such as metal chelation,

photoprotection, protection from heat and cold stress, binding to antifungals to reduce efficacy,

radioprotection (physical shielding and quenching of cytotoxic free radicals), protection from

desiccation, interacting with a range of electromagnetic radiation frequencies and mechanical-

chemical cellular strength (Dadachova et al. 2008; Casadevall et al. 2003; Gorbushina et al. 2003).

Understanding poorly studied fungal function in one extreme environment could yield insight into

MCF adaptation in other extreme environments.

102
The objective of this study was to collect MCF from biological soil crust samples from desert

drylands. During our collection, we isolated 3 MCF that had not been described previously. Our

efforts to converse with indigenous tribes allowed us to provide a new naming scheme for our

strains. We also used a new methodology described herein using BUSCO nucleotide and amino

acid sequences to build a multi-locus phylogenetic tree, which proved to be similar to a more

conventional tree of MCF which used four highly conserved markers.

109
Materials and Methods

Sampling

Biological soil crusts were collected from two locations in Southern California. The first crust

location at the northern edge of Joshua Tree National Park. The second crust location was from

Boyd Deep Canyon, near the campgrounds in the UC Natural Reserve. Metadata and coordinates

can be found in **Table 1**. All materials were collected under permits from the BLM or permission

of the UC National Reserves.

Culturing and isolation

Media for isolation, cultivation, and collection used were Malt Extract Yeast Extract (MEYE) media for nutrient-rich complex media and nutrient-poor, oligotrophic media Glucose Asparagine Agar (GAA). The recipe for both media can be found in our protocols.io and the step-by-step protocol for culturing MCFs (Kurbessoian 2019)). The mixing of the sample with the media provides an environment with different layers of dissolved oxygen. Fast-growing fungi require full oxygenation and grow on the surface of the media, while slow-growing MCFs propagate on the bottom of the culture plate, requiring low levels of oxygen. This also allowed a much easier way to culture and select the strains. Once grown, axenic MCFs were isolated by using a flamed loop to pick and place onto clean media.

Morphological analysis

Strains were grown on Malt Extract Agar (MEA; Oxoid, England), Oatmeal Agar (OA; Oxoid) and Potato dextrose Agar (PDA; Sigma, Germany) incubated at 28°C for 14 days under UV light. Colonies were photographed with a Nikon C-DSD230 dissecting microscope. Microscopic mounts were made in lactic acid and examined using a light microscope (Nikon Eclipse 80i). Pictures were taken with a digital camera (Nikon Digital Sight, DS-5M) and adjusted in Adobe Photoshop CS.

DNA extraction

Each isolate was grown on MEYE for approximately 1.5 weeks at 22°C. Genomic DNA was extracted from hyphae collected from MEYE plates using the CTAB protocol described in our protocols.io document (Carter-House et al. 2020). Melanin was removed using multiple iterations of phenol:chloroform and chloroform washes of our sample. Genomic DNA was measured by

Nanodrop and diluted to ~28ng/µl. DNA extractions were sent to the UCR Genomics core

(Riverside, CA) for 2x150bp sequencing on an Illumina NovoSeq 6000. DNA from all three

isolates was also extracted and sequenced on Oxford Nanopore (ONT) platform with library

preparation and sequencing following the manufacturer's directions (Oxford Nanopore, Oxford

United Kingdom). Flow cell versions FAK68780, FAQ05493, FAL40904, and FAL35715 were

used along with base-calling on the cluster using guppy (*v.3.4.4.+a296cb*) (Wick et al. 2019).

147
ITS sequencing

To identify the isolates, extracted DNA was used as a template for PCR amplification using fungal

primers ITS1-F (CTTGGTCATTTAGAGGAAGTAA) and ITS4

(TCCTCCGCTTATTGATATGC) along with the PCR setup. PCR was performed in 25 µL

reaction volumes using 12.5 µL of Taq 2X MasterMix (NEB), 9.5µL water, 1µL of 20 picomol

ITS1-F, 1µL of 20 picomol ITS4 and 1µL of template DNA. Using Bio-Rad thermal cycler, the

samples were passed through denaturation and enzyme activation at 95°C for 5 minutes, followed

by amplification for 35 cycles at the following conditions: 30 seconds at 95°C, 1 minute at 72°C,

and 30 seconds at 65°C. A final 5-minute extension at 72°C completes the protocol. PCR results

typically yield between 50-60 ng of amplified DNA.

Genome assembly and annotation

Initial genome assemblies were constructed for the three MCF isolates using Illumina sequencing.

Each MCF was also re-sequenced using Oxford Nanopore technology. All genomes were *de novo*

assembled with the AAFTF pipeline (*v.0.2.3*) (Palmer and Stajich 2022) which performs read QC

and filtering with BBTools bbduk (*v.38.86*) (Bushnell 2014) followed by SPAdes (*v.3.15.2*)

(Bankevich et al. 2012) assembly using default parameters, followed by screening to remove short

contigs < 200 bp and contamination using NCBI's VecScreen. The BUSCO ascomycota_odb10

database (Manni et al. 2021) was used to determine how complete the assembly 3 isolates were. A

hybrid assembly of each was generated using MaSUrCa *(v.3.3.4)* (Zimin et al. 2013) as the

assembler using both Nanopore and Illumina sequencing reads. General default parameters were

used except: CA_PARAMETERS=cgwErrorRate=0.15, NUM_THREADS=16, and

JF_SIZE=200000000.

170
We predicted genes in each near-complete genome assembly with Funannotate (*v1.8.1*) (Palmer

and Stajich 2020). A masked genome was created by generating a library of sequence repeats with

the RepeatModeler pipeline (Smit and Hubley 2008). These species-specific predicted repeats

were combined with fungal repeats in the RepBase (Bao et al. 2015) to identify and mask repetitive

regions in the genome assembly with RepeatMasker (*v.4-1-1*) (SMIT 2004). To predict genes, *ab*

initio gene predictors SNAP (*v.2013_11_29*) (Korf 2004) and AUGUSTUS (*v.3.3.3*) (Stanke et al.

2006)were used along with additional gene models by GeneMark.HMM-ES (*v.4.62_lic*) (Brůna et

al. 2020), and GlimmerHMM (*v.3.0.4*) (Majoros et al. 2004) utilize a self-training procedure to

optimize *ab initio* predictions. Additional exon evidence to provide hints to gene predictors was

generated by DIAMOND BLASTX alignment of SwissprotDB proteins and polished by Exonerate

(*v.2.4.0*) (Slater and Birney 2005). Finally, EvidenceModeler (*v.1.1.1*) (Haas et al. 2008) generated

consensus gene models in Funannotate that were constructed using default evidence weights. Non-

protein-coding tRNA genes were predicted by tRNAscan-SE (*v.2.0.9*) (Lowe and Chan 2016).

The annotated genomes were processed with antiSMASH (*v.5.1.1*) [(Blin et al. 2021)](#) to predict

secondary metabolite biosynthesis gene clusters. These annotations were also incorporated into

the functional annotation using Funannotate. Putative protein functions were assigned to genes

based on sequence similarity to InterProScan5 (*v.5.51-85.0*) [(Jones et al. 2014)](#), Pfam (*v.35.0*)

[(Finn et al. 2014)](#), Eggnog (*v.2.1.6-d35afda*) [(Huerta-Cepas et al. 2019)](#), dbCAN2 (*v.9.0*) [(Zhang](#)

[et al. 2018)](#) and MEROPS (*v.12.0*) [(Rawlings et al. 2018)](#) databases relying on NCBI BLAST

(*v.2.9.0+)* [(Sofi et al. 2022)](#) and HMMer (*v.3.3.2*) [(Potter et al. 2018)](#). Gene Ontology terms were

assigned to protein products based on the inferred homology based on these sequence similarity

analyses. The final annotation produced by Funannotate was deposited in NCBI as a genome

assembly with gene model annotation.

195
196 Sequence Read Archive (SRA) files and genome assembly and annotation files can be found under

197 BioProject [PRJNA631111](#).

198
199 *Mating type determination*

200 The Mating Type (MAT) locus was identified by searching for homologous MAT genes. The

201 identified homologous regions were examined for their conserved synteny of the MAT locus using

202 clinker [(Gilchrist and Chooi 2021)](#) and a custom Biopython script [(Cock *et al.* 2009)](#) [(Kurbessoian](#)

203 [2022)](#) to extract the annotated region of the genome which contained the locus.

204
Telomeric repeat determination

Identification of telomeric repeat sequences was performed using the FindTelomeres.py script

(https://github.com/JanaSperschneider/FindTelomeres). Briefly, this searches for chromosomal

assembly with a regular expression pattern for telomeric sequences at each scaffold's 5' and 3'

end. Telomere repeat sequences were also predicted using the Telomere Identification toolkit (tidk)

(*v.0.1.5*) "explore" option (https://github.com/tolkit/telomeric-identifier).

211
212 *Inferring phylogenetic relationships of species*

213 Inference of the species relationship was made through phylogenetic trees containing the three

214 new MCF strains described here and available named species with sequence data.

215
A multi-gene tree was generated using collected BT2 (Beta-Tubulin), CAM (Calmodulin), ITS

(Inter-Transcribed Spacer), and LSU (Large Subunit) genes from a total of 32 species. These

samples were chosen to compare to the three MCF species with closely related species to confirm

identification as a novel lineage. The final selection of 32 species was based on iterations of

sequence searching and tree building data analyses to determine close relatives. To include genes

from genomes, genome assemblies were searched by TBLASTN tool of NCBI-BLAST (*v.2.9.0+*)

using *Zymoseptoria tritici* genes (AAS55060.1 for Beta tubulin) and (AEI69650.1 for

Calmodulin). The collection of ITS and LSU genes were obtained from NCBI-Genbank based on

BLASTN searches. Each gene file contained FASTA formatted sequences from each species or

isolate, the sequences were aligned with MUSCLE (*v.5.1*) and trimmed for low quality alignment

regions using Clipkit (*v.1.3.0*). Nucleotide substitution models were determined for each alignment

with IQTREE2. The alignments were combined into a single 'supermatrix' alignment with

combine_multiseq_aln.py script provided in PHYling. The script also produced a partition file

listing the sub-locations of each marker gene alignment in the super-alignment. IQTree2 was run

to find a substitution model fit for and partition merging. To reduce the computational burden that

occurs with including a large number of organism ribosomal sequences we used the relaxed

hierarchical clustering algorithm (Lanfear et al. 2014) using the tag -m MF+MERGE and running

1000 bootstrapping on the alignment.

234
Choosing to work with indigenous tribes

Our choice for naming of the three MCFs isolated from BSCs is rooted in the traditional indigenous

land where these were found. *Neophaeococcomyces mojaviensis* was isolated from BSCs collected

from the south-east region of Joshua Tree National Park, territories ancestral to

Yuhaaviatam/Maarenga'yam (Serrano) and Newe Segobia (Western Shoshone) Tribes (Brown

and Boyd 1922; Kroeber 1925). The isolates of *Coniosporium tulheliwenetii* and *Taxawa*

tesnikishii were isolated from BSCs collected from UC Reserve Boyd Deep Canyon (Reserve

doi:10.21973/N3V66D), land of the traditional ancestral Cahuilla Tribe (White 2001; Hooper

1920). The word *tulheliweneti* in Cahuilla means, black-spread-be-noun, while *Taxaw* means

body, and *tesnikish* means yellow. In Cahuilla traditions, Taxaw tesnikish is the deity that resided

in Boyd Deep valley. To recognize the origins, we consulted with Tribal linguists and speakers to

derive their names from Cahuilla and Serrano terms that describe either geographic region, oral

tradition, or features that help distinguish them.

Results

249
Biological soil crust collection

Biological soil crusts were collected from the two regions described in **Table 1**. The method of

sterile collection is through the use of sterile empty Petri dishes, stamped on top, with a sterile

spatula used to grab the soil from underneath, flip the sample into the dish, place the lid on top and

seal it with parafilm.

Culturing results

Samples material was used to culture fungi using a collection of techniques in our laboratory at

UC Riverside. More than 50+ fungal culture specimens were collected from biological soil crusts,

18 were MCF cultures. The pour plate protocol used allows for fast-growing fungi to appear on

the top, while MCF grows within the bottom layer of media in Petri dishes. Samples were excised

from deep in the media and allowed to grow on fresh MEYE. Each sample had the DNA extracted

and each strain ITS was Sanger sequenced to determine the identity. JES_112 was initially

identified as a *Knufia* species. JES_115 and JES_119 were identified as *Dothideales* species, but

there remained ambiguity on the exact species identification as a near perfect sequence match was

not found for ITS1 to assign a name.

Sequencing and assembly of MCF isolates

To gain information on the genome content for the recovered MCF isolates, we sequenced and

assembled the genomes of the three isolates (**Table 2**). Each isolate was sequenced with Illumina

MiSeq and Oxford Nanopore (ONT) technology and constructed into a hybrid assembly that

combined both read types (**Table 2**). The depth of coverage ranged from 25-43x Illumina coverage

across the three specimens, while Nanopore ranged from 0.83-194.58x. Assessment of the genome

completeness was performed with BUSCO (Benchmarking Universal Single-Copy Orthologs) and

results indicated the genomes had ranges of 73-92.9% complete gene copies found, with 73-92.8%

of the markers appearing as single copies in the assembly, a low fraction of duplication (0.1-0.2).

Contig counts ranged from 38-480, while the average genome assembly size is about 30 Mbps for

all isolates. The L50 statistic, a measure of the count of the smallest number of contigs whose sum

makes up half of the genome size, ranged from 9-88, and the N50 statistic, a measure of the

smallest contig length for covering half of the genome size, ranging from 117kbp - 1.232 Mbp.

To further assess completeness, we tested for the presence of telomeric repeat units

"TTTAGGG/CCCTAA". These were identified as repeat arrays at both ends of contigs in JES_115

and JES_119, 61 total telomeres and 55 total telomeres respectively, though the results were

inconclusive for JES_112. No telomeric repeat-containing contigs were detected in JES_112

which could be due to the fragmented nature of the genome assembly. The fragmented assembly

also is reflected in the lower BUSCO score for this genome (**Table 2**).

284
285   The number of predicted tRNA genes ranged from 76-197, the total number of protein-coding

286   genes predicted ranged from 9,029-10,920, and secondary metabolite cluster counts ranged from

287   15-27, 27 resulting from JES_112.

288
289

Mating type loci determination

The mating types were determined using prior work on members of Chaetothyriales and

Dothideomycetes including *Exophiala dermatitidis*, and *Aureobasidium pullulans* (Metin et al.

2019; Teixeira et al. 2017). Using collected nucleotide and protein FASTA sequences, BLAST

was used to identify the genes found in these loci. The genes in this locus are generally flanked by

SLA2 and APN2 and may sometimes have alternative genes within. **Figure 2** was generated using

JES_112, *Exophiala aquamarina* CBS 119918, *E. dermatitidis* NIH/UT8656, *E. dermatitidis* CBS

115663*, and *Capronia coronata* CBS 617.96. The MAT locus is flanked by two genes, SLA2

(purple) and APN2 (orange). The MAT 1-1 gene, MAT 1-1-4 (green), and MAT 1-1-1 (teal) are

observed in *E. dermatitidis* CBS 115663 and *Capronia coronata*, while *E. dermatitidis*

NIH/UT8656 contains MAT 1-2 (pink) as does *Capronia coronata*. Our organism in question,

Neophaeococcomyces mojaviensis contains the MAT 1-2 gene along with the MAT 1-1-1 gene.

This is similarly seen with *N. aloes* FJII-L3-CM-P2. This is confusing as the MAT 1-1-1 gene

usually pairs with MAT 1-1-4. The gene arrow in blue is indicated as an L-type calcium channel

domain, while the gene in yellow is an unidentified functional gene or domain.

306
Figure 3 was generated using JES_115 and JES_119 and two other species including,

Aureobasidium pullulans EXF-150 and *Coniosporium apollinis* CBS 100218. All species

described here only have the MAT 1-2 gene. The MAT locus is flanked by two genes, SLA2

(purple) and APN2 (orange). *Dothideomycetes* mating type loci look very different from the

Chaetothyriales mating type loci. Other genes included in this locus concern only a single MAT

locus, MAT 1-2 (pink), and rRNA adenine demethylase (yellow). Certain genes like the AP

endonuclease (teal) and a hypothetical protein (blue) are not seen in all four genomes.

314
MCF morphology

Strains showed limited and slow growth with black colonies on all media tested. Strain JES_119

produced abundant unilateral, hyaline budding cells which soon inflated, developed cruciate

septation, and became melanized. Strain JES_115 was hyphal with lateral chains of swollen cells

which occasionally detached. Strain JES_112 lacked sporulation, the thallus consisting of

melanized densely septate hyphae with series of somewhat inflated cells

During the annotation step, we processed the genomes through antiSMASH (**Table 3**), to detect

Biosynthetic Gene Clusters (BGCs) of secondary metabolites from the annotated genome

sequence. Once predicted metabolite of note, beta-lactone was found in all three MCF. A further

literature search indicates the importance of beta-lactone in industry as there have been multiple

products derived from these chemicals including, antifungals and anti-cancer agents (Robinson et

al. 2019). A BLASTP search of the JES_112 beta-lactone gene identified homologs to Acetyl-

CoA synthetase + 2-isopropylmalate synthases + bile acid dehydrogenase, zinc-binding alcohol

dehydrogenase + acetoacetate-CoA + NAD-dependent alcohol dehydrogenase, and AMP-binding

protein + 2-isopropylmalate synthases.

331
332   A second predicted metabolite of note was the siderophore gene cluster. These genes are involved

333   in iron transport, is a more well-known metabolite that could provide the organism with virulence

334   and protection by sequestering iron molecules from its surrounding environment (Johnson 2008;

335   Sass et al. 2019) and was primarily detected in JES_112.

336
Additionally, the BSC search revealed fungal-RiPPs (Ribosomal and post-transcriptionally

modified proteins) in JES_119. The homolog was best classified as a gamma-glutamyltransferase

+ cytochrome protein. Fungal-RiPPs have been used as an alternative source of bioactive

metabolites and could be of industrial interest (Vogt and Künzler 2019). Additional BGCs were

found including Type-1 Polyketide synthases (T1PKS), Type-3 Polyketide synthases, non-

ribosomal peptide synthases, non-ribosomal peptide synthase-like, and terpenes.

343
344

Four-locus gene trees

The multi-gene internal-transcribed space (ITS1), large ribosomal subunit (LSU) beta-tubulin,

calmodulin tree (**Figure 4** and **Figure 5)** places the new isolates in context of named species in

the known genera (**Table 3**).

Forty-three isolates were used, and the alignment of the sequences were trimmed, and concatenated

to run the IQTree program and generate a concise multi-loci gene tree for *Neophaeococcomyces*

mojaviensis JES112. Here this strain is placed in the genus *Neophaeococcomyces*, along with

Exophiala dermatitidis NIH/UT8656, *E. radicis* P2854, *E. oligospoerma* CBS 265.49, *E.*

oligospoerma CBS 127587, and *E. xenobiotica* CBS 11817, *Fonsecaea pedrosoi* CBS 271.37,

Capronia coronata CBS 617.96, *Cladophialophora immunda* CBS 834.96, *C. carrionii* CBS

260.83, *C. yegresii* CBS 114405, *Neophaeococcomyces aloes* FIJI-L3-CM-P2*, N. catenatus*

CBS650.76*, N. oklahomaensis* EMSL 3313*, Knufia petricola* MA5789*, K. separata*

CGMCC3.17337*, K. mediterranea* CCFEE 6211*, K. mediterranea* CCFEE6205*, Knufia* sp. MV-

2018-MLT-8*, K. walvisbayicola* CBS 146989*, K. tsunedae* FMR10621*, Bradymyces yunnanensis*

CGmCC3.17350*, B. alpinus* CCFEE 5493*, Strelitziana africana* ICMP 21758 and ICMP 21760*,*

and *Tuber melanosporum* Mel8 as the outgroup.

361
The third phylogenetic tree, **Figure 5**, uses the same method described above as in **Figure 4**. 25

strains of Dothideomycetes MCF were included to place JES_115 and JES_119 into their proper

clades. MCF strains included a collection of *Coniosporium apollinis* CBS 100218*,* CBS 352.97*,*

Aureobasidium mangrovei IBRCM30265*, Aureobasidium* sp. AWRI4619-CO-2020*, Zalaria*

obscura FKI-L7-BK-DRAB2*, Polyplosphaeria fusca* CBS 125425*, Leptosphaeria maculans* JN3*,*

Leptosphaeria biglobosa CA1*, Phaeosphaeria* sp. MPI-PUGE-AT-0046c*, Alternaria solani*

NL03003*, Cenococcum geophilum* 1.58*, Venturia pyrina* ICMP11032*, Coniosporium uncinatum*

CBS 100219*,* and CCFEE 6149*, Scorias spongiosa* HLSS1*, Polychaeton citri* CBS 116435*,*

Zymoseptoria tritici CBS 100329*, Neohortaea acidophila* CBS 113389*, Salinomyces thailandica*

CCFEE 6315*, Hortaea werneckii* EXF-2000*, Baudoinia panamericana* UAMH 10762*,*

Teratosphaeria guachensis CMW 17545*, Friedmanniomyces simplex* CCFEE 5184*,*

Friedmanniomyces endolithicus CCFEE 5311*,* and *Tuber melanosporum* Mel8 as the outgroup.

JES_115 groups near *Coniosporium* spp.*, w*hile JES_119 groups near *Aureobasidium* spp*.*

Taxonomy

Neophaococcomyces mojaviensis sp. nov.*, Kurbessoian, Ahmed, de Hoog and Stajich -

Mycobank (848161). Description: Colonies on MEA, OA, PDA after two weeks of incubation at

25°C attain diameters of 1.6, 2, 2.2, and 1.6 mm respectively. Hyphae 2.9-5.4 μm diameter, dark

brown or black, smooth or occasionally rough walled, cells eventually inflating up to 8 μm in

diameter. The proliferation of inflated cells might result in the formation of multicellular bodies

terminal or intercalary in the hyphae measuring 13.8-31.8 μm, multiseptate with transverse and

longitudinal septa; bodies round or irregular, dark brown to black and thick-walled. No conidia

were observed. Sexual state unknown.

Holotype: ATCC (TSD-444), preserved at the herbarium collection of the Westerdijk Fungal

Biodiversity Institute, Utrecht, The Netherlands. Living culture ex-type ATCC (TSD-444).

Diagnosis: Strain JES_112 clusters in the order *Chaetothyriales*, relatively close to the numerous

rock-inhabiting species of the genus *Knufia*, and amidst the species of the small genus

Neophaeococcomyces. Both genera are assigned to the family *Trichomeriaceae*, containing 'sooty

molds' with oligotrophic lifestyles colonizing inert surfaces (Chomnunti et al. 2012). The cluster

Neophaeococcomyces, based on *N. catenatus* as generic type species, as yet contains only four

species, which mostly were isolated from low nutrient habitats, such as the bark of *Aloe*, stone, or

air. Several species of the genus show budding with melanized cells, and frequently cells cohere

in short chains. Often lateral branches of short inflated cells are inserted on hyphae; the cells may

be released as arthroconidia which then swell or appear yeast-like, or remain coherent.

***Coniosporium tulheliwenetii sp. nov.*, Kurbessoian, Ahmed, de Hoog and Stajich - MycoBank**

(848162).

Description: Colonies on MEA, OA, PDA, after 2 weeks of incubation at 25°C attain diameter

restricted, measuring 1.1 cm, 0.72 cm, and 0.75cm, respectively. Hyphae 4.6-8.4 µm diameter

showing lateral chains of swollen cells 7.7 - 10.8 µm. The cells however remain intact, and no

conidia was observed. Teleomorph unknown.

Holotype: CBS (150075), and herbarium CBS H-25288, preserved at the herbarium collection

of the Westerdijk Fungal Biodiversity Institute, Utrecht, The Netherlands. Living culture ex-type

CBS (150075).

Diagnosis: Strain JES_115 was morphologically similar to *Knufia* species in the presence of lateral

chains of swollen cells inserted on main hyphae, the cells eventually being released as conidia, but

was phylogenetically remote. It clustered with some species of rock-inhabiting described in

Coniosporium, in the order *Pleosporales*. Interestingly, the original descriptions, which were made

before sequence data were available, classified them all into a single genus. The genus

Coniosporium is one of the classical genera of meristematic fungi. However, most species were

described in the 19th century, including the generic type species *C. olivaceum*, and voucher

specimens are lost. Nevertheless, Haridas et al. (2020) introduced the family *Coniosporiaceae* and

order *Coniosporales* using the few recently described rock-inhabiting fungi, *C. apollinis* and *C.*

uncinatum as reference (Haridas et al. 2020). It should be noted that the family *Coniosporiaceae*

was already used by Nannfeldt for the genus *Coniospora*.

417
***Taxawa* gen. nov. Kurbessoian, Ahmed, de Hoog and Stajich - MycoBank (848163).**

Type species: *Taxawa tesnikishii*

Taxawa tesnikishii sp. nov. Kurbessoian, Ahmed, de Hoog and Stajich - MycoBank (848164).

Description: Colonies moist, black, regular and flat on OA and heaped irregular on MEA and

PDA, measuring 1.4-1.8 cm in diameter growing on MEA, OA, and PDA at 25°C . Hyphae

smooth, thick-walled, measuring 6.1-10.3 μm wide. Yeast cells abundant produce unilateral

budding 2.7-4.8 μm, subsequently inflating and becoming crucially septate and melanized with

age. Clusters of black meristematic bodies (9.0-56.8 μm) were observed. Sexual state unknown.

Holotype: CBS (150074), and herbarium CBS H-25287, preserved at the herbarium collection

of the Westerdijk Fungal Biodiversity Institute, Utrecht, The Netherlands. Living culture ex-type

CBS (150074).

Diagnosis: Strain JES_119 showed similar morphology to the cultural states of e.g. *Sydowia* and

Comminutispora, and has been described in the asexual genus *Phaeotheca*. Thambugala et al.

(2014) erected the family *Aureobasidiaceae* for fungi in this relationship, but also this name

existed already, *Aureobasidiaceae* Ciferri (1958) (Thambugala et al. 2014), based on the same

species, *Aureobasidium pullulans* (Ciferri 1958). Only a small number of extant species has been

sequenced, and the genome trees show an even larger taxon sampling effect, but given our current

inability to assign strain JES_119 to any genus, we introduce the genus *Taxawa*.

Discussion

BSCs are an untapped resource of novel microorganisms, using non-traditional methods of

culturing may be a method to uncover more unavailable species for sequencing. Using a pour plate

method for culturing allows the microorganism to avoid complete oxygen saturation (or oxygen

toxicity), and instead can grow in low to micro-aerophilic concentrations of oxygen (Stevens

1995). Collecting MCF using this method has proved to be efficient, as fast-growing fungi

overtake the surface of plates, while MCF can tolerate lower oxygen concentrations, thus

localizing to the bottom of plates. Preparing axenic cultures of MCF is then relatively easy and

can also increase the rate of isolation as MCF tends to take two weeks to grow. A hypothesis about

the facultative aerobic or microaerophilic nature of MCF could also be seen in the phenotype of

these species. Morphological growth on surface of petri plates shows unusual furrowing and

clumping growth (JES_119, JES_115), some strains (JES_112) grow underneath the surface of the

plate which seem to be MCF adaptations to the high concentration of oxygen present on petri

plates.

The high melanin content of MCF complicates efforts to extract high molecular weight (HMW)

DNA for sequencing, however adjustments to standard fungal CTAB can dramatically improve

the gDNA purity. In the case for these BSC MCF, an increase in phenol:chloroform and

chloroform washes allow for melanin to be removed completely from the DNA samples. Another

consideration to help concentrate more DNA is to incubate the DNA sample in -20°C overnight

during the DNA precipitation step. Using long read sequencing technology (Oxford ONT) it is

best to have a very high concentration of clean DNA to process and thus could be the limiting

factor for these sequencing procedures. Using hybrid methods of sequencing (Illumina and

Nanopore) provides scientists the proper backbone to develop *de-novo* assemblies (Stevens 1995;

Khezri et al. 2021; Chen et al. 2020; Saud et al. 2021). It is vital to have as near complete

assemblies of species as possible, this provides the scientific community with the best tools for

proper genomic assessments. For future research endeavors, a second nanopore flow cell would

be necessary to help build a better assembly for strain JES_112 (*Neophaeococcomyces*

mojaviensis) as the telomere analysis failed on this first round of assembly. The other two strains

had enough depth of coverage that allowed for different programs to optimize the assembly and

have more closed telomere contigs.

Scientific literature describing novel MCF always note the clonal or asexual nature, describing

"missing" sexual states and thus always being in a haploid state. The MAT locus is one of the

pivotal regions in the fungal genome that provides more accurate ploidy and sexual states of

fungi, and using this data to describe the three new MCF assisted in understanding the sexual

states. JES_115 and JES_119 as the two *Dothideomycetes* representatives indicate heterothallic

sexual states having only the MAT1-2 gene, *vs.* homothallic species having representatives of

both MAT1-1 and MAT1-2 genes. Better genome assembly would help better describe the

JES_112 MAT locus as there is the presence of MAT1-2 but also a portion of the MAT1-1 gene

(as it is composed of two genes, MAT1-1-1 and MAT1-1-4). The high melanin content of MCF

could prove to be difficult when testing for ploidy in the scientific "gold standard" method (flow

cytometry). But future research would most benefit from having this method optimized to

understand not only the new MCF described in this work but also for future MCF considerations.

Possibly the sexual states have not yet been described or seen for MCFs and may require a

particular environmental stimulus to produce the proper structures to reproduce. Possibly, other

forms of propagation are found in other habitats. A prototypical example is *Auerobasidium*

pullulans which shows meristematic growth on glass or rock but is a rapidly expanding yeast-

like fungus in culture. There has also been an increase in literature describing clonal MCF having

high hybridization rates that create pseudo but stable diploids without the process of

recombination (Gostinčar et al. 2022). This was not detected in these three species described in

this paper.

MCF morphology indicates usual slow growth (about two weeks) seen in most MCF described

previously (Selbmann et al. 2014; Chowdhary et al. 2014; Hölker et al. 2004; Zhao et al. 2010;

Kejžar et al. 2013). JES_119, as it develops over time, increases its melanin production and

develops from a bright orange-yellow color to a more deeply pigmented black. The budding cells

are unilateral and are uncommon in other described MCF, and it was evaluated as a previously

undescribed genus and species in Class *Dothideomycetes* as *Taxawa tesnikishii.* Strain JES_115

has a phenotype that is commonly seen in MCF, but due to its complicated taxonomy history, the

genus *Coniosporium* remains as a genus of doubtful identity. The species JES_112 is one of the

few described *Neophaeococcomyces* available anywhere and thus an important contribution to the

scientific community.

Finding MCF from extreme environments such as BSCs may point to harbored secondary

metabolites either absorbed from surrounding species through horizontal gene transfer (HGT) or

adapted from time. MCF are known to use secondary metabolites to help survive such extremes

and are thus an untapped resource for the identification of novel metabolites. An example to be

described by Dr. Erin Carr, *Neophaeococcomyces* sp, has been shown to have an obligate bacterial

hitchhiker (Carr 2022). JES_115 and JES_119 both contain about 15 secondary metabolites

detected by antiSMASH, which is a considerable number when compared to other fungi (Rateb

and Ebel 2011). The roles of secondary metabolites in modern human industry applications are

vast (Demain 2014) and thus would be important to test and compare across all MCF.

511
512   **Conclusions**
513
514   Biological soil crusts are an untapped resource for novel microorganisms as evidenced by our

515   work. The MCF we collected are incredibly slow growing and could be easily missed during

516   culturing attempts. The method described here (and detailed in **Appendix 1**) improved the chances

517   for isolation of these species. MCFs have been isolated from different types of extreme

518   environments including BSC's.

519
An interesting consideration for species JES_112 or *Neophaeococcomyces mojaviensis* is through

personal communications with another MCF scientist, Erin Carr, who also isolated a species of

Neophaococcomyces, named "Crusty" from a biological soil crust. Their species was associated

with a pink *Methylobacterium* symbiont, which she calls "Light Pinky"and "Dark Pinky". Her

assessment of this symbiosis suggests the role of the symbiont is involved in aerobic anoxygenic

photosynthesis and auxin production (Carr 2022). Through our multiple culturing attempts, there

was no indication of a pink symbiont, but the possible bacteria signal within our secondary

metabolite assessment could be attributed to a contamination from this bacterium. JES_112 will

be the second known genome of the genus *Neophaeococcomyces* found in GenBank as

JAPDRQ000000000.

Species JES_119 or *Taxawa tesnikishi* is the first isolated of this phenotype. This species was

closer to *Aureobasidium*, yet phenotypically is very different from identified *Aureobasidium*

species. Using fungal phenotype and phylogenomics, we were able to determine the

microorganism is different from known *Aureobasidium* species and may be of a new genus, hence

the naming *Taxawa* as its genus name. *Taxawa tesnikishi* is the first known genus and species,

now available in Genbank as JAPDRO000000000.

Species JES_115 or *Coniosporium tulheliwenetii* was also collected from the BSC's from Boyd

Deep Canyon and was thus honored with a Cahuilla term used to describe BSC phenotypic features

seen on the soil. The genus *Coniosporium* has historically been confusing, as the earliest strains

were collected in the late 1800s. Initial collections of *Coniosporium* were placed in

Chaetothyriales, while later similar MCF phenotypes were also grouped with *Coniosporium*, to be

later considered as Dothideomycetes instead. A careful reorganization and assessment of the genus

is desperately needed. Genome assembly is now available as JAPDRP000000000.

Acknowledgments

This work was performed in part at the University of California Natural Reserve System (Boyd

Deep Canyon Desert Research Center) Reserve DOI: (doi:10.21973/N3V66D) (UCNRS

Information Manager 1965) and at Joshua Tree National Park under permits generated through

the help and guidance of Nuttapon Pombubpa, permit numbers: JOTR-2015-SCI-0033, JOTR-

2016-SCI-0036, JOTR-2017-SCI-0044, JOTR-2018-SCI-0035, JOTR-2019-SCI-0031. Many

thanks to Nicole Pietrasiak for her extensive knowledge in biological soil crusts. Many thanks

also to Joshua Tree Reserve members who helped facilitate the admission of our Joshua Tree

sample, specifically Anna R Tegarden, Ann Hitchcock, and Jay Goodwin.

556
557   **Funding**
558

T.K was partially supported by NIH grant R01 AI127548 (support to J.E.S. via Deborah Hogan).

Computational analyses were performed at the University of California – Riverside HPCC,

supported by grants from the National Science Foundation (DBI – 1429826 and 2215705) and

NIH (S10OD016290). JES is a CIFAR Fellow in the Fungal Kingdom: Threats and

Opportunities program.

Figure Legends:

Figure 1. Microscopic morphology of isolated MCF. (1) *Taxawa tesnikishii* JES_119, colonies on MEA
(A), OA (B), and PDA (C) agars, meristematic cells (D-F), and budding cells (G-M); (2) *Coniosporium*
tulheliwenetii JES_115, colonies on MEA (A), OA (B), and PDA (C) agars, hyphae with lateral branches
of swollen cells reluctantly leading to arthroconidia (D-J); (3) *Neophaeococcomyces mojaviensis* JES_112,
colonies on MEA (A), OA (B), and PDA (C) agars, hyphae producing intercalary swollen cells without
conidiation (D-J). Microscopic mounts were made in lactic acid and examined using a light microscope.
The scale depicted here is 25µm.

Figure 2. Mating type loci determined for Chaetothyriales JES_112 here called *Neophaeococcomyces
mojaviensis. Other MCFs added to this analysis include *Neophaeococcomyces aloes* FJII-L3-CM-P2*,*
Exophiala dermatitidis NIH/UT8656, *E. dermatitidis* CBS 115663, and *Capronia coronata* CBS 617.96.
The MAT locus is flanked by two genes, SLA2 (purple) and APN2 (orange). The MAT 1-1 gene, MAT 1-
1-4 (green), and MAT 1-1-1 (teal) are observed in *E. dermatitidis* CBS 115663 and *C. coronata*, while *N.*
aloes and *E. dermatitidis* NIH/UT8656 contain MAT 1-2 (pink) as does *C. coronata*. Our organism in
question, *N. mojaviensis* contains the MAT 1-2 and MAT 1-1-1 gene which is confusing as it is not common
to lose the MAT 1-1-4. The gene arrow in blue is indicated as an L-type calcium channel domain, while the
gene in yellow is an unidentified functional gene or domain.

Figure 3. Mating type loci determined for Dothideomycetes JES_115 and JES_119, here called
Coniosporium tulheliwenetii **and *Taxawa tesnikishi*.** Other dothideaceous black yeast-like fungi in this

analysis include *Aureobasidium pullulans* EXF-150, and *Coniosporium apollinis CBS* 100218. All species
described here only have the MAT 1-2 gene. The MAT locus is flanked by two genes, SLA2 (purple) and
APN2 (orange). *Dothideomycetes* mating type loci look very different from the *Chaetothyriales* mating
type loci. Other genes included in this locus include only a single MAT locus, MAT 1-2 (pink). Another
gene included in this locus is rRNA adenine demethylase (yellow). Certain genes like the AP endonuclease
(teal) and a hypothetical protein (blue) are not seen in all four genomes.
592
593   **Figure 4. Chaetothyriales multi-locus gene tree.** Multi-loci phylogenetic tree using 4 highly conserved
594   genes including, beta-tubulin, calmodulin, internal-transcribed sequence (ITS), and the large subunit of
595   the Ribosome (LSU). Seven representative strains with loci were used, which were aligned, trimmed, and
596   concatenated to run the IQTree program and generate a concise multi-loci gene tree for
597   *Neophaeococcomyces mojaviensis* JES112. Here this strain is placed in the *Neophaeococcomyces* genera.
598
Figure 5. Dothideomycetes multi-locus gene tree. Multi-loci phylogenetic tree using 4 highly
conserved genes including, beta-tubulin, calmodulin, internal-transcribed sequence (ITS), and the large
subunit of the Ribosome (LSU). Twenty-five representative strains with loci were used, which were
aligned, trimmed, and concatenated to run the IQTree program and generate a concise multi-loci gene tree
for JES_115 and JES_119 or *Coniosporium tulheliwenetii* and *Taxawa tesnikishii* respectively. Here
strain JES_115 is grouping with other *Coniosporium* spp., while JES_119 is grouping with but distinctly
apart from *Aureobasidium* spp.
606
607   **Table 1. Three species were collected from biological soil crusts collected from two desert locations**
608   **in Southern California.** Strain JES_112 was isolated from the south east region of Joshua Tree National
609   Park in the Mojave Desert, while JES_115 and JES_119 were both collected from biological soil crusts
610   collected from the campground site in the UC Natural Reserve of Boyd Deep Canyon.
611
612   **Table 2. Genome assembly and annotation summary statistics.** All strains were sequenced with
613   Illumina and Nanopore sequencing. With the added sequencing reads, we were able to glean telomere
614   information along with genome completeness. It is clear that JES_112 initially is contaminated as the
615   BUSCO score is lower, while JES_115 and JES_119 are much closer to a more complete genome.
616
617   **Table 3. Predicted secondary metabolite gene clusters.** The BGCs predicted include multiple copies of
618   PKS and NRPS genes, but siderophores were only recovered in JES_112. There was a higher number of
619   BGCs predicted in JES_112 (27) our *Chaetothyriales* species, than JES_115 and JES_119 (15) our
620   *Dothideomycetes*. This table lists the numbers of each. JES_112 has a much higher number (27) of
621   secondary metabolites than the other two (15).
622
623   **Table 4. Gene Tree Nucleotide Accession numbers.** Collected accession numbers of alternative strains
624   to build Figures 4 and 5. The remaining empty spaces indicate the need to pull the sequences using BLAST.
625   Once collected each gene populated a gene FASTA file, aligned, clipped and IQTREE2, a tree-building
626   program, was run to determine the gene models for each gene. Once determined, the gene alignment lengths
627   were calculated and partitioning scripts were described and used to build a larger tree, using IQTREE2.
628
629

References:

Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI,
Pham S, Prjibelski AD, et al. 2012. SPAdes: a new genome assembly algorithm and its applications to
single-cell sequencing. J Comput Biol. 19(5):455–477.

Bell AA, Wheeler MH. 1986. Biosynthesis and Functions of Fungal Melanins. Annu Rev Phytopathol.
24(1):411–451.

Belnap J, Harper KT. 1995. Influence of cryptobiotic soil crusts on elemental content of tissue of two
desert seed plants. Arid Soil Research and Rehabilitation. 9(2):107–115.

Beraldi-Campesi H, Retallack GJ. 2016b. Terrestrial Ecosystems in the Precambrian. In: Weber B, Büdel
B, Belnap J, editors. Biological Soil Crusts: An Organizing Principle in Drylands. Cham: Springer
International Publishing. p. 37–54.

Blin K, Shaw S, Kloosterman AM, Charlop-Powers Z, van Wezel GP, Medema MH, Weber T. 2021.
antiSMASH 6.0: improving cluster detection and comparison capabilities. Nucleic Acids Res.
49(W1):W29–W35.

Brown J, Boyd J. 1922. History of San Bernardino and Riverside Counties: With Selected Biography of
Actors and Witnesses of the Period of Growth and Achievement. Western Historical Association.

Brůna T, Lomsadze A, Borodovsky M. 2020. GeneMark-EP : eukaryotic gene prediction with self-
training in the space of genes and proteins. NAR Genomics and Bioinformatics. 2(2).
doi:10.1093/nargab/lqaa026. http://dx.doi.org/10.1093/nargab/lqaa026.

Bushnell B. 2014. BBMap: A fast, accurate, splice-aware aligner. Lawrence Berkeley National Lab.
(LBNL), Berkeley, CA (United States) Report No.: LBNL-7065E. https://www.osti.gov/biblio/1241166.

Carr EC. 2022. Discovering Novel Polyextremotolerant Fungi, and Determining Their Ecological Role
Within the Biological Soil Crust Consortium.

Casadevall A, Rosas AL, Nosanchuk JD. 2000. Melanin and virulence in *Cryptococcus neoformans*.
Current Opinion in Microbiology. 3(4):354–358. doi:10.1016/s1369-5274(00)00103-x.

Casadevall A, Steenbergen JN, Nosanchuk JD. 2003. "Ready made" virulence and "dual use" virulence
factors in pathogenic environmental fungi — the *Cryptococcus neoformans* paradigm. Current Opinion in
Microbiology. 6(4):332–337. doi:10.1016/s1369-5274(03)00082-1.

Chomnunti P, Bhat DJ, Jones EBG, Chukeatirote E, Bahkali AH, Hyde KD. 2012. Trichomeriaceae, a
new sooty mould family of Chaetothyriales. Fungal Divers. 56(1):63–76.

Chowdhary A, Perfect J, de Hoog GS. 2014. Black Molds and Melanized Yeasts Pathogenic to Humans.
Cold Spring Harb Perspect Med. 5(8):a019570.
675
676 Ciferri R. 1958. Affinity between fungi of the verrucous dermatitis and sooty-molds. Mycopathol Mycol
677 Appl. 9(3):197–200.
678
679 Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, Friedberg I, Hamelryck T, Kauff F,
680 Wilczynski B, et al. 2009. Biopython: freely available Python tools for computational molecular biology
681 and bioinformatics. Bioinformatics. 25(11):1422–1423.
682
683 Dadachova E, Bryan RA, Howell RC, Schweitzer AD, Aisen P, Nosanchuk JD, Casadevall A. 2008. The
684 radioprotective properties of fungal melanin are a function of its chemical composition, stable radical
685 presence and spatial arrangement. Pigment Cell Melanoma Res. 21(2):192–199.
686
687 Dadachova E, Casadevall A. 2008. Ionizing radiation: how fungi cope, adapt, and exploit with the help of
688 melanin. Curr Opin Microbiol. 11(6):525–531.
689
690 Demain AL. 2014. Valuable Secondary Metabolites from Fungi. In: Martín J-F, García-Estrada C,
691 Zeilinger S, editors. Biosynthesis and Molecular Genetics of Fungal Secondary Metabolites. New York,
692 NY: Springer New York. p. 1–15.
693
694 Evans RD, Ehleringer JR. 1993. A break in the nitrogen cycle in aridlands? Evidence from δp15N of
695 soils. Oecologia. 94(3):314–317.
696
697 Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, Heger A, Hetherington K, Holm L,
698 Mistry J, et al. 2014. Pfam: the protein families database. Nucleic Acids Res. 42(Database issue):D222–
699 30.
700
701 Gilchrist CLM, Chooi Y-H. 2021 Jan 18. Clinker & clustermap.js: Automatic generation of gene cluster
702 comparison figures. Bioinformatics. doi:10.1093/bioinformatics/btab007.
703
704 Gómez BL, Nosanchuk JD. 2003. Melanin and fungi. Curr Opin Infect Dis. 16(2):91–96.
705
706 Gorbushina AA, Whitehead K, Dornieden T, Niesse A, Schulte A, Hedges JI. 2003. Black fungal
707 colonies as units of survival: hyphal mycosporines synthesized by rock-dwelling microcolonial fungi. Can
708 J Bot. 81(2):131–138.
709
710 Gostinčar C, Sun X, Černoša A, Fang C, Gunde-Cimerman N, Song Z. 2022. Clonality, inbreeding, and
711 hybridization in two extremotolerant black yeasts. Gigascience. 11. doi:10.1093/gigascience/giac095.
712
713 Green TGA, Allan Green TG, Proctor MCF. 2016. Physiology of Photosynthetic Organisms Within
714 Biological Soil Crusts: Their Adaptation, Flexibility, and Plasticity. Biological Soil Crusts: An
715 Organizing Principle in Drylands.:347–381. doi:10.1007/978-3-319-30214-0_18.
716

717 Green TGA, Broady PA. 2001. Biological soil crusts of Antarctica. In: Ecological Studies. Berlin,
718 Heidelberg: Springer Berlin Heidelberg. (Ecological studies: analysis and synthesis. Berlin, Heidelberg,
719 New York NY). p. 133–139.
720
721 Gümral R, Tümgör A, Saraçlı MA, Yıldıran ŞT, Ilkit M, de Hoog GS. 2014. Black yeast diversity on
722 creosoted railway sleepers changes with ambient climatic conditions. Microb Ecol. 68(4):699–707.
723
724 Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, White O, Buell CR, Wortman JR. 2008.
725 Automated eukaryotic gene structure annotation using EVidenceModeler and the Program to Assemble
726 Spliced Alignments. Genome Biol. 9(1):R7.
727
728 Haridas S, Albert R, Binder M, Bloem J, LaButti K, Salamov A, Andreopoulos B, Baker SE, Barry K,
729 Bills G, et al. 2020. 101 Dothideomycetes genomes: A test case for predicting lifestyles and emergence of
730 pathogens. Stud Mycol. 96:141–153.
731
732 Hölker U, Bend J, Pracht R, Tetsch L, Müller T, Höfer M, de Hoog GS. 2004. *Hortaea acidophila*, a new
733 acid-tolerant black yeast from lignite. Antonie Van Leeuwenhoek. 86(4):287–294.
734
735 de Hoog GS, Hermanides-Nijhof EJ. 1977. The Black Yeasts and Allied Hyphomycetes, Studies in
736 Mycology. Baarn, The Netherlands: Centraalbureau voor Schimmelcultures.
737
738 de Hoog GS, Matos T, Sudhadham M, Luijsterburg KF, Haase G. 2005. Intestinal prevalence of the
739 neurotropic black yeast *Exophiala (Wangiella) dermatitidis* in healthy and impaired individuals. Mycoses.
740 48(2):142–145.
741
742 Hoog GS de, Queiroz-Telles F, Haase G, Fernandez-Zeppenfeldt G, Angelis DA, H. G. Gerrits van den
743 Ende A, Matos T, Peltroche-Llacsahuanga H, Pizzirani-Kleiner AA, Rainer J, et al. 2000a. Black fungi:
744 clinical and pathogenic approaches. Med Mycol. 38(sup1):243–250.
745
746 Hooper L. 1920. The Cahuilla Indians. University of California Press.
747
748 Housman DC, Yeager CM, Darby BJ, Sanford RL, Kuske CR, Neher DA, Belnap J. 2007. Heterogeneity
749 of soil nutrients and subsurface biota in a dryland ecosystem. Soil Biol Biochem. 39(8):2138–2149.
750
751 Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, Mende DR, Letunic
752 I, Rattei T, Jensen LJ, et al. 2019. eggNOG 5.0: a hierarchical, functionally and phylogenetically
753 annotated orthology resource based on 5090 organisms and 2502 viruses. Nucleic Acids Res.
754 47(D1):D309–D314.
755
756 James TY, Stajich JE, Hittinger CT, Rokas A. 2020. Toward a Fully Resolved Fungal Tree of Life. Annu
757 Rev Microbiol. 74:291–313.
758
759 Johnson L. 2008. Iron and siderophores in fungal–host interactions. Mycol Res. 112(2):170–183.
760

761   Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A, Nuka
762   G, et al. 2014. InterProScan 5: genome-scale protein function classification. Bioinformatics. 30(9):1236–
763   1240.
764
765   Kejžar A, Gobec S, Plemenitaš A, Lenassi M. 2013. Melanin is crucial for growth of the black yeast
766   *Hortaea werneckii* in its natural hypersaline environment. Fungal Biology. 117(5):368–379.
767   doi:10.1016/j.funbio.2013.03.006. http://dx.doi.org/10.1016/j.funbio.2013.03.006.
768
769   Korf I. 2004. Gene finding in novel genomes. BMC Bioinformatics. 5:59.
770
771   Kroeber AL. 1925. Handbook of the Indians of California. U.S. Government Printing Office.
772
773   Kurbessoian T. 2022. tania-k/Chapter2.3MCF: V1. https://zenodo.org/record/7259500.
774
775   Lal R. 2004b. Carbon sequestration in dryland ecosystems. Environ Manage. 33(4):528–544.
776
777   Lanfear R, Calcott B, Kainer D, Mayer C, Stamatakis A. 2014. Selecting optimal partitioning schemes for
778   phylogenomic datasets. BMC Evol Biol. 14:82.
779
780   Little AEF, Currie CR. 2008. Black yeast symbionts compromise the efficiency of antibiotic defenses in
781   fungus-growing ants. Ecology. 89(5):1216–1222.
782
783   Lowe TM, Chan PP. 2016. tRNAscan-SE On-line: integrating search and context for analysis of transfer
784   RNA genes. Nucleic Acids Res. 44(W1):W54–7.
785
786   Maestre FT, Soliveres S, Gotelli NJ, Quero JL, Berdugo M. 2012. Response to Comment on "Plant
787   Species Richness and Ecosystem Multifunctionality in Global Drylands." Science. 337(6091):155–155.
788   doi:10.1126/science.1220620. http://dx.doi.org/10.1126/science.1220620.
789
790   Majoros WH, Pertea M, Salzberg SL. 2004. TigrScan and GlimmerHMM: two open source ab initio
791   eukaryotic gene-finders. Bioinformatics. 20(16):2878–2879.
792
793   Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update: Novel and
794   Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of
795   Eukaryotic, Prokaryotic, and Viral Genomes. Mol Biol Evol. 38(10):4647–4654.
796
797   Mehdiabadi NJ, Schultz TR. Natural history and phylogeny of the fungus-farming ants (Hymenoptera:
798   Formicidae: Myrmicinae: Attini). [accessed 2022 Mar 29].
799
800   Metin B, Döğen A, Yıldırım E, de Hoog GS, Heitman J, Ilkit M. 2019. Mating type (MAT) locus and
801   possible sexuality of the opportunistic pathogen *Exophiala dermatitidis*. Fungal Genetics and Biology.
802   124:29–38. doi:10.1016/j.fgb.2018.12.011.
803

Nosanchuk JD, Stark RE, Casadevall A. 2015. Fungal melanin: What do we know about structure? Front
Microbiol. 6:1463.

Palmer JM, Stajich J. 2020. Funannotate v1.8.1: Eukaryotic genome annotation.
https://zenodo.org/record/4054262.

Pietrasiak N. 2005. Field guide to classify biological soil crusts for ecological site evaluation. NRCS.

Pietrasiak N, Regus JU, Johansen JR, Lam D, Sachs JL, Santiago LS. 2013. Biological soil crust
community types differ in key ecological functions. Soil Biol Biochem. 65:168–171.

Potter SC, Luciani A, Eddy SR, Park Y, Lopez R, Finn RD. 2018. HMMER web server: 2018 update.
Nucleic Acids Res. 46(W1):W200–W204.

Rateb ME, Ebel R. 2011. Secondary metabolites of fungi from marine habitats. Nat Prod Rep. 28(2):290–
344.

Rawlings ND, Barrett AJ, Thomas PD, Huang X, Bateman A, Finn RD. 2018. The MEROPS database of
proteolytic enzymes, their substrates and inhibitors in 2017 and a comparison with peptidases in the
PANTHER database. Nucleic Acids Res. 46(D1):D624–D632.

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other things).
Methods Ecol Evol. 3(2):217–223.

Robinson SL, Christenson JK, Wackett LP. 2019. Biosynthesis and chemical diversity of β-lactone
natural products. Nat Prod Rep. 36(3):458–475.

Rozenstein O, Karnieli A. 2015. Identification and characterization of Biological Soil Crusts in a sand
dune desert environment across Israel–Egypt border using LWIR emittance spectroscopy. J Arid Environ.
112:75–86.

Sass G, Ansari SR, Dietl A-M, Déziel E, Haas H, Stevens DA. 2019. Intermicrobial interaction:
*Aspergillus fumigatus*siderophores protect against competition by *Pseudomonas aeruginosa*. PLoS One.
14(5):e0216085.

Saunte DM, Tarazooie B, Arendrup MC, de Hoog GS. 2012a. Black yeast-like fungi in skin and nail: it
probably matters. Mycoses. 55(2):161–167.

Siletti CE, Zeiner CA, Bhatnagar JM. 2017. Distributions of fungal melanin across species and soils. Soil
Biol Biochem. 113:285–293.

Slater GSC, Birney E. 2005. Automated generation of heuristics for biological sequence comparison.
BMC Bioinformatics. 6:31.

SMIT A. F. A. 2004. Repeat-Masker Open-3.0. http://www.repeatmasker.org.

Sofi MY, Shafi A, Masoodi KZ. 2022. NCBI BLAST. Bioinformatics for Everyone.:95–102.
doi:10.1016/b978-0-323-91128-3.00021-5.

Staley JT, Palmer F, Adams JB. 1982. Microcolonial fungi: common inhabitants on desert rocks?
Science. 215(4536):1093–1095.

Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B. 2006. AUGUSTUS: ab initio
prediction of alternative transcripts. Nucleic Acids Res. 34(Web Server issue):W435–9.

Steenbergen JN, Casadevall A. 2003. The origin and maintenance of virulence for the human pathogenic
fungus Cryptococcus neoformans. Microbes Infect. 5(7):667–675.

Stevens TO. 1995. Optimization of media for enumeration and isolation of aerobic heterotrophic bacteria
from the deep terrestrial subsurface. J Microbiol Methods. 21(3):293–303.

Taborda CP, da Silva MB, Nosanchuk JD, Travassos LR. 2008b. Melanin as a virulence factor of
Paracoccidioides brasiliensis and other dimorphic pathogenic fungi: a minireview. Mycopathologia.
165(4-5):331–339.

Teixeira MM, Moreno LF, Stielow BJ, Muszewska A, Hainaut M, Gonzaga L, Abouelleil A, Patané JSL,
Priest M, Souza R, et al. 2017. Exploring the genomic diversity of black yeasts and relatives
(Chaetothyriales, Ascomycota). Stud Mycol. 86:1–28.

Thambugala KM, Ariyawansa HA, Li Y-M, Boonmee S, Hongsanan S, Tian Q, Singtripop C, Bhat DJ,
Camporesi E, Jayawardena R, et al. 2014. Dothideales. Fungal Divers. 68(1):105–158.

Thanh VN, Hien DD. 2019. *Moniliella floricola* sp. nov., a species of black yeast isolated from flowers.
International Journal of Systematic and Evolutionary Microbiology. 69(1):87–92.
doi:10.1099/ijsem.0.003099. http://dx.doi.org/10.1099/ijsem.0.003099.

UCNRS Information Manager. 1965. Boyd deep canyon desert research center. doi:10.21973/N3V66D.

Vicente VA, Attili-Angelis D, Pie MR, Queiroz-Telles F, Cruz LM, Najafzadeh MJ, de Hoog GS, Zhao J,
Pizzirani-Kleiner A. 2008. Environmental isolation of black yeast-like fungi involved in human infection.
Stud Mycol. 61:137–144.

Vogt E, Künzler M. 2019. Discovery of novel fungal RiPP biosynthetic pathways and their application
for the development of peptide therapeutics. Appl Microbiol Biotechnol. 103(14):5567–5581.

Wang L-G, Lam TT-Y, Xu S, Dai Z, Zhou L, Feng T, Guo P, Dunn CW, Jones BR, Bradley T, et al.
2020. Treeio: An R Package for Phylogenetic Tree Input and Output with Richly Annotated and
Associated Data. Mol Biol Evol. 37(2):599–603.

Warren SD, Clair LL St, Stark LR, Lewis LA, Pombubpa N, Kurbessoian T, Stajich JE, Aanderud ZT.
2019. Reproduction and dispersal of biological soil crust organisms. Front Ecol Evol. 7.
doi:10.3389/fevo.2019.00344.

Weber B, Bowker M, Zhang Y, Belnap J. 2016. Natural Recovery of Biological Soil Crusts After
Disturbance. In: Weber B, Büdel B, Belnap J, editors. Biological Soil Crusts: An Organizing Principle in
Drylands. Cham: Springer International Publishing. p. 479–498.

Wheeler MH. 1983. Comparisons of fungal melanin biosynthesis in ascomycetous, imperfect and
basidiomycetous fungi. Trans Br Mycol Soc. 81(1):29–36.

White PM. 2001. California Indians and Their Reservations: An Online Dictionary. San Diego State
University Library.

Wick RR, Judd LM, Holt KE. 2019. Performance of neural network basecalling tools for Oxford
Nanopore sequencing. Genome Biol. 20(1):129.

Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. 2017. Ggtree : An r package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. Methods Ecol Evol.
8(1):28–36.

Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, Busk PK, Xu Y, Yin Y. 2018. dbCAN2: a meta
server for automated carbohydrate-active enzyme annotation. Nucleic Acids Res. 46(W1):W95–W101.

Zhao J, Zeng J, de Hoog GS, Attili-Angelis D, Prenafeta-Boldú FX. 2010. Isolation and identification of
black yeasts by enrichment on atmospheres of monoaromatic hydrocarbons. Microb Ecol. 60(1):149–156.

Zimin AV, Marçais G, Puiu D, Roberts M, Salzberg SL, Yorke JA. 2013. The MaSuRCA genome
assembler. Bioinformatics. 29(21):2669–2677. doi:10.1093/bioinformatics/btt476.
http://dx.doi.org/10.1093/bioinformatics/btt476.

Aureobasidium_pullulans_EXF-150_MAT

XP_029765312.1 XP_029765311.1 XP_029766327.1 XP_029765566.1 XP_029765964.1 XP_029765963.1 XP_029765962.1 XP_029765961.1 XP_029765960.1

Taxawa_tesnikishii_JES119_MAT

H2203_009196 H2203_009195 H2203_009187 H2203_009186 H2203_009185 H2203_009184 H2203_007159 H2203_007158 H2203_007157 H2203_007156

Coniosporium_tulheliwenetii_JES115_MAT

H2199_005258 H2199_005257 H2199_005371 H2199_005370 H2199_005369 H2199_005368 H2199_000674 H2199_000673 H2199_000672 H2199_000671

Coniosporium_apollinis_CBS100218_MAT

XP_007781593.1 XP_007781592.1 XP_007781591.1 XP_007782976.1 XP_007782975.1 XP_007782974.1 XP_007779098.1 XP_007779097.1

2.5kb

rRNA adenine demethylase
APN2
MAT 1-2
SLA2
Ribonucleoprotein
AP endonuclease
Hypothetical protein

0.3

Table 1. Three species were collected from biological soil crusts collected from two desert locations in Southern California. Strain JES_112 was isolated from the south east region of Joshua Tree National Park in the Mojave Desert, while JES_115 and JES_119 were both collected from biological soil crusts collected from the campground site in the UC Natural Reserve of Boyd Deep Canyon.

Species	Date	Location	Coordinates
JES_112	November 2017	Joshua Tree National Park, CA	34.1026 °N, -115.4542°W
JES_115	July 2019	Boyd Deep Canyon Reserve, CA	34.0182 °N, -116.0262 °W
JES_119	July 2019	Boyd Deep Canyon Reserve, CA	34.0182 °N, -116.0262 °W

Table 2. Genome assembly and annotation summary statistics. All strains were sequenced with Illumina and Nanopore sequencing. With the added sequencing reads, we were able to glean telomere information along with genome completeness. It is clear that JES_112 is contaminated as the BUSCO score is lower, while JES_115 and JES_119 are much closer to a more complete genome.

	JES_112	JES_115	JES_119
Scaffold Count	478	64	38
Total Length	32,593,754 bp	32,607,911 bp	29,383,553 bp
Minimum Length	561	4823	1783
Maximum Length	427,480	1,739,523	2,828,030
Mean Scaffold Length	68,187.77	509,498.61	773,251.39
Scaffold L50	87	14	9
Scaffold N50	117,971	966,560	1,232,081
Scaffold L90	282	31	24
Scaffold N90	32,261	507,175	558,687
BUSCO Complete % Illumina	84.7	90.2	97.4
BUSCO Complete % Nanopore	73.4	92.6	92.9
BUSCO Duplicate % Nanopore	0.2	0.1	0.1
BUSCO Single % Nanopore	73.2	92.5	92.8
GC%	48.25	52.55	51.98

Illumina coverage	34.58x	43.45x	25.38x
Nanopore coverage	0.83x	20.44x	194.58x
Telomere Count	0, 0F, 0R	61, 29 F, 32 R	55, 29 F, 26 R
Number of Genes	10,996	9,304	9,386
mRNA genes	10,924	9,226	9,189
tRNA genes	72	78	197
GO Terms	6,001	5,089	5,321
InterProScan	8,523	7,079	7,291
EGGNOG	9,934	8,487	8,511
PFAM	7,520	5,982	6,219
CAZYme	309	257	346
MEROPS	344	284	291
Secretion	674	532	631

Table 3. Predicted secondary metabolite gene clusters. The BGCs predicted include multiple copies of PKS and NRPS genes, but siderophores were only recovered in JES_112. There was a higher number of BGCs predicted in JES_112 (27) our *Chaetothyriales* species, than JES_115 and JES_119 (15) our *Dothideomycetes*. This table lists the numbers of each. JES_112 has a much higher number (27) of secondary metabolites than the other two (15).

	T1PKS	Terpene	T3PKS	NRPS	NRPS-like	Siderophore	Arylpolyene	Betalactone	Fungal RiPP
JES 112	6	6	2	4	6	1	1	1	0
JES 115	5	4	0	2	3	0	0	1	0
JES 119	1	4	0	1	7	0	0	1	1

Table 4. Gene Tree Nucleotide Accession numbers. Collected accession numbers of alternative strains to build Figures 4 and 5. The remaining empty spaces indicate the need to pull the sequences using BLAST. Once collected each gene populated a gene FASTA file, aligned, clipped and IQTREE2, a tree-building program, was run to determine the gene models for each gene. Once determined, the gene alignment lengths were calculated and partitioning scripts were described and used to build a larger tree, using IQTREE2.

Species	Beta-Tubulin	Calmodulin	Internal-Transcribed Spacer	Large Subunit (Ribosomal)
*Neophaeococcomyces mojaviensis*JES112	-	-	-	-
Taxawa tesnikishii JES119	-	-	-	-
Alternaria solani NL03003	**MK388240.1**	**MH243808.1**	**NR_172408.1**	**ON416546.1**
Anthracina ramosa CGMCC3.16372	**KP226556.1**	-	**NR_172152.1**	**KP174923.1**
Anthracina saxicola CGMCC3.17315	**KP226554.1**	-	**NR_172151.1**	**KP174921.1**
Arthrocladium caudatum CBS457.67	**LT558710.1**	-	**NR_145398.1**	**MH870732.1**
Arthrocladium tardum CBS 127021	-	-	**NR_154706.1**	**NG_057089.1**
Arthrocladium tropicale CBS 134926	**KX822406.1**	-	**NR_154724.1**	**NG_057119.1**
Aureobasidium mangrovei IBRCM30265	**XM_02990485 2.1**	**XM_04102580 9.1**	**NR_174637.1**	**NG_078639.1**
*Aureobasidium sp.*AWRI4619-CO-2020	**XM_02990485 2.1**	**XM_04102580 9.1**	**MK256457.1**	**MK968773.1**
Baudoinia panamericana UAMH10762	**XM_0076738 85.1**	**XM_00767526 4.1**	**NR_153615.1**	**KT186497.1**
Bradymyces alpinus CCFEE 5493	**LN589970.1**	-	**NR_132844.1**	**NG_058641.1**

Bradymyces yunnanensis CGMCC 3.17350	**KP226545.1**	-	**KP174865.1**	**KP174926.1**
Capronia coronata CBS 617.96	**XM_0077289 82.1**	**XM_00772155 6.1**	**NR_154745.1**	**NG_067831.1**
Cenococcum geophilum 1.58	-	-	**KC967410.1**	**MN817694.1**
Cladophialophora carrionii CBS 260.83	**KF928582.1**	**XM_00872400 0.1**	**MH861582.1**	**MH873312.1**
Cladophialophora_yegre sii CBS 114405	**EU137209.1**	-	**NR_111284.1**	**KX822323.1**
Cladophialophora immunda CBS 834.96	**EU137203.1**	-	**NR_111283.1**	**MH874242.1**
Coniosporium apollinis CBS 100218	-	**XM_00778549 5.1**	**NR_159787.1**	**GU250898.1**
Coniosporium uncinatum CBS 100219	-	-	**NR_145343.1**	**NG_058806.1**
Coniosporium uncinatum CCFEE 6149	-	-	**MZ573424.1**	**MZ573424.1**
Cyphellophora aestiva CBS 239.91	**MW297547.1**	-	**MH861291.1**	**MH873056.1**
Cyphellophora laciniata CBS 239.91	**KF928604.1**	-	**NR_121335.1**	**MH873933.1**
Cyphellophora sessilis KNU-002	**KX431943.1**	-	**KX431941.1**	**KX431942.1**
Cyphellophora vermispora CBS 228.86	**KC455227.1**	-	**KC455244.1**	**KC455257.1**
Exophiala dermatitidis NIH.UT8656	-	**XM_00916045 0.1**	**JH226131.1**	**ON543213.1**
Exophiala oligosperma CBS 265.49	**EF551507.1**	**XM_01640456 0.1**	**MH856519.1**	**MH868049.1**

Exophiala oligosperma CBS 127587	**KF928570.1**	-	**MH864631.1**	**MH876068.1**
Exophiala radicis P2854	**KT723463.1**	-	**NR_158397.1**	**NG_069319.1**
Exophiala xenobiotica CBS 118157	**NW_013562490.1**	**XM_013454656.1**	**OP132917.1**	**MH875251.1**
Fonsecaea pedrosoi CBS 271.37	**XM_013430809.1**	**XM_013426864.1**	**MT023597.1**	**KF155185.1**
Friedmanniomyces endolithicus CCFEE 5311	-	-	**NR_144917.1**	**GU250367.1**
Friedmanniomyces simplex CCFEE 5184	-	-	**NR_155079.1**	**GU250368.1**
Hortaea werneckii EXF-2000	-	-	**NR_145338.1**	**NG_057773.1**
Knufia petricola MA5789	-	-	**NR_111871.1**	**MT126669.1**
Knufia mediterranea CCFEE 6211	-	-	**KP791793.1**	**KR781080.1**
Knufia mediterranea CCFEE 6205	-	-	**KP791794.1**	**KR781081.1**
Knufia separata CGMCC 3.17337	**KP226564.1**	-	**NR_172233.1**	**KP174931.1**
Knufia tsunedae FMR10621	-	-	**NR_132842.1**	**HG003672.1**
Knufia walvisbayicola CBS 146989	**MZ078266.1**	-	**NR_173046.1**	**NG_076737.1**
Knufia sp. MV-2018 MLT-8	-	-	**MT636977.1**	**MT636937.1**
Leptosphaeria biglobosa CA1	**AY749005.1**	-	**NR_111620.1**	**JX681093.1**
Leptosphaeria maculans JN3	**AY749012.1**	**XM_003840719.1**	**M96384.1**	**JF740306.1**

Lithohypha aloicola CPC35996	**MN556837.1**	-	**NR_166313.1**	**NG_068316.1**
Lithohypha guttulata FMR 20100	**OX346320.1**	-	**OX346376.1**	**OX346377.1**
Metulocladosporiella musicola CBS 110960	-	**MG934537.1**	**MH862870.1**	**NG_067447.1**
Metulocladosporiella samutensis CBS 143921	-	**MG934540.1**	**NR_161056.1**	**MG934453.1**
Neohortaea acidophila CBS 113389	**XM_033733287.1**	**XM_033733930.1**	**NR_169648.1**	**OL739260.1**
Neophaeococcomyces aloes FJII-L3-CM-P2	-	-	**NR_132069.1**	**KF777234.1**
Neophaeococcomyces catenatus CBS 650.76	-	-	**AF050277.1**	**MH872793.1**
Neophaeococcomyces oklahomaensis EMSL3313	-	-	**MW798187.1**	**MW798182.1**
*Phaeosphaeria sp.*MPI-PUGE-AT-0046c	-	-	**NR_155675.1**	**NG_068582.1**
Polychaeton citri CBS 116435	-	-	**GU214649.1**	**OM238161.1**
Polyplosphaeria fusca CBS 125425	-	-	**NR_119397.1**	**MH875147.1**
Salinomyces thailandica CCFEE 6315	-	-	**NR_171710.1**	**MH875167.1**
Scorias spongiosa HLSS1	-	-	**NR_154422.1**	**MH866910.1**
Strelitziana africana ICMP 21758	-	-	**MK210538.1**	**MK210499.1**
Strelitziana africana ICMP 21760	-	-	**MK210540.1**	**MK210501.1**

Teratosphaeria gauchensis CMW 17545	-	KF902726.1	NR_144976.1	EU019295.2
Trichomerium cicatricatum CGMCC3.17307	KP226540.1	-	NR_172232.1	KP174947.1
Trichomerium deniqulatum MFLUCC10-0884	-	-	NR_132965.1	NG_059479.1
Trichomerium gleosproum MFLUCC10-0087	-	-	JX313656.1	JX313662.1
Trichomerium sp. XDY-2021a-IFRDCC3302	-	-	MN901488.1	MZ822131.1
Tuber melanosporum	MZ458420.1	XM_002841184.1	GQ917052.1	JF807486.1
Venturia pyrina ICMP11032	EU853839.1	-	NR_169905.1	AY849967.1
Zalaria obscura FKI-L7-BK-DRAB2	-	-	NR_153466.1	MT103097.1
Zymoseptoria tritici CBS 100329	XM_003856727.1	KF254299.1	KP852517.1	MK788267.1