Prosapip1 in the dorsal hippocampus mediates synaptic protein composition, long-term

potentiation, and spatial memory

Zachary W. Hoisington[1], Himanshu Gangal[2], Khanhky Phamluong[1], Chhavi Shukla[1], Yann

Ehinger[1], Jeffrey J. Moffat[1], Gregg E. Homanics[3], Jun Wang[2], and Dorit Ron[1]

[1]Alcohol and Addiction Research Group, Department of Neurology, University of California San

Francisco; [2]Department of Neuroscience and Experimental Therapeutics, School of Medicine,

Texas A&M University Health Science Center; [3]Department of Anesthesiology and Perioperative

Medicine, University of Pittsburgh

*Correspondence: D. Ron, Department of Neurology, University of California, San Francisco, 675

Nelson Rising Lane, BOX 0663, San Francisco, CA, USA. dorit.ron@ucsf.edu

Abstract

Prosapip1 is a brain-specific protein localized to the postsynaptic density, where it promotes dendritic spine maturation in primary hippocampal neurons. However, nothing is known about the role of Prosapip1 *in vivo*. To examine this, we utilized the Cre-loxP system to develop a Prosapip1 neuronal knockout mouse. We found that Prosapip1 controls the synaptic localization of its binding partner SPAR, along with PSD-95 and the GluN2B subunit of the NMDA receptor (NMDAR) in the dorsal hippocampus (dHP). We next sought to identify the potential contribution of Prosapip1 to the activity and function of the NMDAR and found that Prosapip1 plays an important role in NMDAR-mediated transmission and long-term potentiation (LTP) in the CA1 region of the dHP. As LTP is the cellular hallmark of learning and memory, we examined the consequences of neuronal knockout of Prosapip1 on dHP-dependent memory. We found that global or dHP-specific neuronal knockout of Prosapip1 caused a deficit in learning and memory whereas developmental, locomotor, and anxiety phenotypes were normal. Taken together, Prosapip1 in the dHP promotes the proper localization of synaptic proteins which, in turn, facilitates LTP driving recognition, social, and spatial learning and memory.

Introduction

ProSAP-interacting protein 1 (Prosapip1), encoded by the gene *LZTS3*, is a brain-specific protein that shows particularly high expression in the cortex, cerebellum, and hippocampus (Wendholt et al., 2006; Laguesse et al., 2017). Prosapip1 is highly enriched in the PSD of excitatory synapses and plays an important role in dendritic spine maturation in primary hippocampal neurons (Wendholt et al., 2006; Dolnik et al., 2016). The PSD is a complex protein network located at the postsynaptic membrane of neurons, namely excitatory neurons (Boeckers et al., 2002; Kaizuka and Takumi, 2018). The PSD is crucial for synaptic transmission and plasticity (Boeckers et al., 2002; Kaizuka and Takumi, 2018), and abnormalities in the PSD network are linked to various neuropsychiatric and neurodegenerative disorders (Kaizuka and Takumi, 2018), such as schizophrenia, autism spectrum disorder (ASD), depression, and Alzheimer's disease (Kaizuka and Takumi, 2018). The PSD contains scaffolding proteins such as PSD-95, Shank, and Homer (Kennedy, 2018). The major scaffold in the PSD is actin (Qualmann et al., 2004), which recycles between globular (G-actin) and filamentous (F-actin), and is crucial in reorganizing and stabilizing the PSD (Qualmann et al., 2004; Kuriu et al., 2006).

Prosapip1 has a C-terminal Fez1 domain that it shares with the other members of the Fezzin family (LAPSER1, PSD-Zip70, N4BP3) (Wendholt et al., 2006). Fezzins interact with the PDZ (PSD-95/Dlg1/ZO-1) domain of SH3 and multiple ankyrin repeat domains (Shank) family proteins. Shanks are also known as proline-rich synapse-associated proteins (ProSAPs) and were originally identified as proteins primarily localizing in the PSD of excitatory synapses (Boeckers et al., 1999; Sheng and Kim, 2000; Boeckers et al., 2002). Prosapip1 specifically interacts with the PSD protein, Shank3 (Wendholt et al., 2006). Prosapip1 also interreacts with the spine-associated Rap GTPase-activating protein (SPAR) (Wendholt et al., 2006; Reim et al., 2016). SPAR catalyzes the conversion of the small G-protein Rap from its active, GTP-bound form to its inactive GDP-bound form, thereby inhibiting Rap activity. SPAR is an essential modulator of spine morphology through its interaction with actin (Pak et al., 2001; Richter et al.,

2007). In hippocampal neurons, it is localized to dendritic spines and leads to spine head

enlargement (Pak et al., 2001).   SPAR also associates with PSD-95/NMDAR-complex

components (Matsuura et al., 2022). Wendholt et al reported that in primary hippocampal neurons

Prosapip1 regulates the synaptic levels of SPAR by scaffolding it to Shank3 (Wendholt et al.,

2006; Dolnik et al., 2016; Reim et al., 2016).

We previously found that *Prosapip1* mRNA to protein translation is controlled by the

mechanistic target of rapamycin complex 1 (mTORC1) in the nucleus accumbens (NAc) of mice

consuming large quantities of alcohol (Laguesse et al., 2017). We further showed that Prosapip1

interacts with SPAR in the NAc and leads to alcohol-dependent synaptic and structural plasticity

(Laguesse et al., 2017). Specifically, we found that knockdown of Prosapip1 in the NAc led to a

reduction in F-actin, while overexpression of Prosapip1 led to an increase in F-actin (Laguesse et

al., 2017). Knockdown of Prosapip1 led to a reduction of mature, mushroom-type spines and an

increase in immature, thin spines (Laguesse et al., 2017). Finally, we found that Prosapip1 in the

NAc contributes to alcohol self-administration and reward (Laguesse et al., 2017). However,

nothing is known about the normal *in vivo* role of Prosapip1 in the central nervous system (CNS).

Here, we report that Prosapip1 in the dorsal hippocampus (dHP) controls the localization of

synaptic proteins at the PSD and contributes to long-term potentiation (LTP) as well as

recognition, social, and spatial learning and memory.

Materials and Methods

Animals

Prosapip1(flx/flx) mice were generated as described below. Syn1-Cre(+) mice were

purchased from Jackson Laboratory (Stock #003966, Bar Harbor, Maine). Prosapip1(flx/flx) and

Prosapip1(flx/flx);Syn1-Cre mice were bred and group housed in a 12-hour light-dark cycle room

that was temperature- and humidity-controlled. Unrestricted amounts of food and water were

provided. Animal procedures were approved by the UCSF Institutional Animal Care and Use

Committee (IACUC).

Generation of Prosapip1 Floxed Mice

Guide RNA binding sites in intron 2 and in the 3'UTR located in Exon 5 of Prosapip1 (a.k.a,

Lzts3) (**Figure 1A**) were identified using the CRISPRator tool (Labuhn et al., 2018) within the

CCTop (Stemmer et al., 2015) online platform (http://crispr.cos.uni-heidelberg.de/). The gRNA

binding sites are located ~3.3kb from each other. These gRNA target sites were used to produce

two Alt-R CRISPR-Cas9 crRNAs (IDT DNA, Coralville, IA) which were individually hybridized to

a universal 67-mer Alt-R CRISPR-Cas9 tracrRNA (IDT DNA) to produce two gRNAs. Two 140 nt

long PAGE purified Ultramer single stranded DNA Oligos (IDT DNA) with three phosphorotioate

modifications on each end (Renaud et al., 2016) that were homologous to the target loci in intron

2 and Exon 5 (**Figure 1B**) were used as a repair templates.

We introduced each loxP site into 1-cell and 2-cell stages of embryonic development (Horii

et al., 2017). Single cell C57BL/6J embryos were electroporated with gRNA#6 (200ng/µl), IDT Alt-

R® HiFi Cas9 Nuclease V3 protein (75ng/µl), and Exon 5 repair template (200ng/µl) using a

BioRad Gene-Pulser Xcell electroporator in a 1 mm-gap slide electrode (Protech International,

#501P1-10) using square-wave pulses (five repeats of 3 msec 25V pulses with 100 ms interpulse

intervals). Following this first electroporation, embryos were cultured overnight. Surviving two-cell

embryos were electroporated with gRNA#7 (200ng/µl), IDT Alt-R® HiFi Cas9 Nuclease V3 protein

(75ng/µl), and intron 2 repair template (200ng/µl) under conditions described above. Surviving

embryos were transferred to the oviducts of CD1 (Charles River) pseudopregnant recipient

females. Offspring were genotyped for the intron 2 lox insertion using PCR (forward primer 5'

AGAGAAGTCTACGCTGTAGTCAG 3' and reverse primer 5' AAGCGGGAAGGTAGAGAGGT 3';

wild type product = 449bp, floxed product = 489bp) followed by Sanger sequencing. Offspring

were genotyped for the Exon 5 lox insertion using PCR (forward primer 5'

TGCACAACCTTCTGACACGT 3' and reverse primer 5' AGGGCACAGACAGTAGCACT 3'; wild

type product = 408bp, floxed product = 448bp) followed by Sanger sequencing. Two founder

animals were found that harbored loxP insertions at both the intron 2 and Exon 5 sites. When

mated to C57BL/6J females, the loxP sites segregated indicating that they were on different

chromosomes. Therefore, F1 offspring that harbored only the loxP insertion at the intron 2 site

were used as 1-cell embryo donors for insertion of the Exon 5 loxP site using electroporation

conditions described above. One male offspring was produced that harbored both the intron 2

and Exon 5 loxP insertions on the same chromosome. This male was mated to C57BL/6J females

to establish the floxed mouse line used here.

Guide RNA off target sites were predicted and ranked using CRISPOR (Concordet and

Haeussler, 2018). The top 11 and 10 sites for gRNA#6 and gRNA#7, respectively based on CFD

score were amplified from this founder mouse DNA and Sanger sequenced. All predicted off

target sites analyzed were wild type (data not shown).

Reagents

Antibodies: Rabbit anti-SPAR (SIPA1L1) (1:500) (25086-1-AP) antibodies were purchased from

ProteinTech. Mouse anti-Shank3 antibodies (1:500) (#ab93607) were purchased from Abcam.

Rabbit anti-GluN2B (1:1000) (#4212), Rabbit anti-GluA1 (1:1000) (#13185S), and Rabbit anti-

CREB (1:500) (#9197) antibodies were purchased from Cell Signaling. Goat anti-GluN2A

antibodies (1:500) (#SC-1468) were purchased from Santa Cruz. Mouse anti-GAPDH antibodies

(1:10,000) (#G8795) were purchased from Sigma. Mouse anti-PSD-95 antibodies (1:100,000)

(#05-494) were purchased from Millipore (Upstate). Donkey anti-rabbit horseradish peroxidase

(HRP), donkey anti-goat HRP, and donkey anti-mouse HRP conjugated secondary antibodies

(1:5000) were purchased from Jackson ImmunoResearch (West Grove, PA). Enhance

Chemiluminescence reagent (ECL) was purchased from Millipore (Burlington, MA). EDTA-free

complete mini–Protease Inhibitor Cocktails was from Roche (Indianapolis, IN). Pierce

bicinchoninic acid (BCA) protein assay kit was purchased from Thermo Scientific (Rockford, IL).

NuPAGE Bis-Tris precast gels were purchased from Life Technologies (Carlsbad, CA).

Viruses: AAV8-Ef1a-mCherry-IRES-Cre (1×10^{13} vg/ml #55632) and AAV2-CMV-EGFP (1×10^{13}

vg/ml, #105530) were purchased from Addgene.

Crude synaptosomal fractionation

Tissue was first homogenized at 4°C in 500 µl of Krebs buffer including 125 mM NaCl, 1.2 mM

KCl, 1.2 mM $MgSO_4$, 1.2 mM $CaCl_2$, 22 mM Na_2CO_3, 1 mM NaH_2PO_4, 10 mM Glucose, 0.32 M

sucrose and protease/phosphatase inhibitors. A portion of the homogenate (100 µl) was saved

as total homogenate (H), while the remaining homogenate was diluted by adding 500 µl of Krebs

buffer to the remaining 400 µl. The total homogenate was transferred to a 1.5 ml Eppendorf tube.

The glass homogenizer was washed with 500 µl of Krebs buffer, and the wash solution was added

to the remaining homogenate. The sample was then centrifuged at 1,000g at 4°C for 10 minutes.

The supernatant (S1) was collected. The process was repeated, and then the S1 supernatant

was centrifuged at 16,000g at 4°C for 20 min. The supernatant (S2) was saved, and the pellet

was kept on ice. The resulting pellet contained the synaptosomal fraction. The pellet was then

resuspended in 500 ul of Krebs buffer and centrifuged at 16,000g 4°C for 20 minutes and

resuspended in radio immunoprecipitation assay (RIPA) buffer (containing 50 mM Tris-HCL, 5

mM EDTA, 120 mM NaCl, 1%NP-40, 0.1% deoxycholate, 0.5% SDS, and protease and

phosphatase inhibitors) for analysis.

Western blot analysis

Tissue collected from mice was homogenized in ice-cold RIPA buffer using a sonic dismembrator.

Protein concentration was determined using the BCA™ protein assay kit. 30 μg of each tissue

lysate was loaded for separation by SDS-PAGE (4-12%), followed by transfer onto a nitrocellulose

membrane at 300mA for 2 hours. The membranes were then blocked with 5% milk-PBS

containing 0.1% Tween 20 at room temperature for 30 minutes before being probed with the

appropriate primary antibodies overnight at 4ºC. Following washing, the membranes were

incubated with HRP-conjugated secondary antibodies for one hour at room temperature and then

visualized using ECL. Band intensities were quantified using ImageJ software (NIH,

https://imagej.net/ij/).

Electrophysiology

Preparation of slices is outlined in (Gangal et al., 2023). In brief, coronal sections of the

hippocampus (250 μm) were cut in an ice-cold solution containing the following: 40 mM NaCl,

148.5 mM sucrose, 4.5 mM KCl, 1.25 mM NaH_2PO_4, 25 mM $NaHCO_3$, 0.5 mM $CaCl_2$, 7 mM

$MgSO_4$, 10 mM dextrose, 1 mM sodium ascorbate, 3 mM myo-inositol, 3 mM sodium pyruvate

and saturated with a 95% O_2 and 5% CO_2. After cutting, the slices were incubated for 45 minutes

in an external solution containing the following: 125 mM NaCl, 4.5 mM KCl, 2.5 mM $CaCl_2$, 1.3

mM $MgSO_4$, 1.25 mM NaH_2PO_4, 25 mM $NaHCO_3$, 15 mM sucrose, 15 mM glucose and saturated

with 95% O_2 and 5% CO_2.

Field potential recordings were conducted as detailed in previous studies (Wang et al.,

2012)**.** Specifically, the procedure involved the use of stimulation electrodes filled with artificial

cerebrospinal fluid (aCSF) and placed in the CA1 region of the dHP. The recording electrode,

filled with 1M NaCl, was positioned in the CA1 region, approximately 100-150 μm away from the

stimulating electrodes. The optimal location for the recording electrode was determined based on

the site where a stable fiber volley and field potential response could be consistently observed

following the delivery of a single electrical stimulation pulse (2 ms). The stimulation intensity was

carefully adjusted to elicit 50% of the maximum possible response. Baseline field potentials were

meticulously recorded over 10 mins at 10-second intervals. The HFS protocol for LTP included a

regimen of 100 Hz, with 100 pulses every 20 seconds, repeated four times. Following the HFS

protocol, field potential recordings were extended for 30 min. LTP was quantified based on the

averaged fEPSP as a percentage of the baseline, comparing the measurements at 20-30 minutes

post-HFS between the Prosapip1(flx/flx);Syn1-Cre(-) and Prosapip1(flx/flx);Syn1-Cre(+) groups.

To inhibit GABAergic transmission, picrotoxin (100 μM) was applied to the bath. The

measurement of fEPSP was performed using a MultiClamp 700B amplifier integrated with

Clampex 10.4 software provided by Molecular Devices.

Whole-cell recordings were executed following the protocols previously established

(Wang et al., 2012; Ma et al., 2018). For these recordings, a cesium-based intracellular solution

was employed. This solution comprised of the following components 119 mM $CsMeSO_4$, 8 mM

Tetraethylammonium chloride (TEA-Cl), 15 mM HEPES, 0.6 mM ethylene glycol tetraacetic acid

(EGTA), 0.3 mM Na_3GTP, 4 mM MgATP, 5 mM QX-314-Cl, and 7 mM phosphocreatine. The pH

of this solution was adjusted to 7.3 using CsOH. The temperature of the recording bath was

maintained at a constant 32°C, and the perfusion speed was set between 2-3 ml/min. For the

recordings, CA1 neurons were voltage-clamped at a holding potential of -70 mV. In order to

measure NMDAR-mediated transmission, a low external concentration of Mg^{2+} (0.05 mM) was

used. This was coupled with NBQX (10 μM) and picrotoxin (100 μM) to block AMPA receptor

(AMPAR)-mediated synaptic transmission and inhibitory synaptic currents. NMDA-induced

currents were measured by bath-applying NMDA (20 μM) for 30 seconds, with holding currents

recorded every 5 seconds.

Input-output curves for NMDAR-mediated EPSC were created by electrical stimulation

of varying intensities which were delivered through electrodes strategically placed within the CA1

region. The paired-pulse ratio (PPR) was calculated by dividing the amplitude of the second

electrically-evoked EPSC by that of the first, with an interval of 100 ms between the two pulses.

Additionally, gap-free recordings of spontaneous excitatory postsynaptic currents were conducted

over a duration of 2 minutes.

Behavior

All behavioral paradigms were completed between the hours of 9:00 and 18:00. Mice were

tested between 8 and 20 weeks of age and were age-matched based on experiment. The mice

were moved from their housing room, currently in the "light cycle," to the dimly lit behavior room

(10-15 lux) 30 minutes before the experiment and allowed to acclimate. The light/dark box,

elevated plus maze, and Barnes maze experiments were done in full room light. During

experiments, white noise was played through a speaker in the room at 50dB to reduce influence

from noise outside of the room. The researcher remained in the room during the trial, but a wall

was placed between the arena and the experimenter to prohibit mice from seeing the researcher.

Mice were handled for 3 days for 1-2 minutes per day before experimentation began to reduce

handling anxiety. All behavioral analyses were recorded and analyzed using Noldus Ethovision

XT software.

Locomotion

Mice were placed in a 43cm x 43cm open field and allowed to explore for 30 minutes. The

center point of the mouse was tracked, and the primarily dependent variable was total locomotion

during the trial. Other tracked variables were average and maximum velocity, center entrances,

time spent in center, and locomotion binned into 1-minute periods to examine evolution of

exploration of the space.

Novel Object Recognition

Mice were placed in a 43cm x 43cm chamber with two similar objects and allowed 20 seconds of total familiarization time (nose point within 2cm radius of object), or 5 total minutes in the chamber, whichever came first, before being returned to home cage. After 24 hours, mice were allowed to explore the test space containing one object from the familiarization trial (familiar object) and one novel object until total object interaction time of 20 seconds was reached. If 20 seconds of total interaction time was not reached before the 5-minute time limit, the trial was excluded (3 trials). Mice that were climbing on objects were gently but quickly moved back to the starting position.

Novelty T-Maze

The protocol was adapted from (Sanderson et al., 2009). Mice were placed in the "start" arm of a T-shaped maze. The dimensions of each transparent arm were 30cm×10cm×20cm (L×W×H). During training, the entrance to one arm ("novel arm", assignment counterbalanced across groups) was blocked with clear plastic. Training consisted of five, 2-minute trials in which each subject was allowed to explore the "start" and "familiar" arms. The training trials were separated by a 1-minute inter-trial interval. After the training sessions, the plastic blocking the "novel" arm was removed and mice were allowed to explore all arms of the maze. The test session began when the center point of the mouse left the start arm and ended when 2 minutes of total time was spent in the novel or familiar arm. Mice that showed anxiety-related behavior and an aversion to either arm during the test (no arm entrances, <250cm total movement) were excluded (2 mice).

3-Chamber Social Interaction

Mice were placed in the center chamber of a transparent 3-chamber arena and allowed 3 minutes to habituate to the space. The mouse was then placed into the closed center chamber

and the trial was started by removing the doors and allowing 15 minutes to explore the "social" or

"empty" chamber (Part I). The social chamber had a sex-matched, naïve mouse (4-5 weeks of

age) in a cylindrical interaction cage, while the empty chamber had only the interaction cage. The

chamber time and interaction time was recorded, with interaction time being defined as

experimental mouse nose point being within interaction zone (5cm from interaction cage).

Immediately after Part I, a novel mouse, also sex-matched and naïve, was introduced in the empty

cage and mice were allowed 15 minutes to explore "Novel" or "Familiar" chamber (Part II). The

familiar mouse is the social mouse from Part I. The locations of the novel and familiar mice were

switched to reduce side preference for this interaction partner.

Barnes Maze

The protocol was adapted from (Pitts, 2018). The Barnes maze apparatus is a 122cm-

diameter, white-acrylic circle with 40 evenly spaced, 5cm holes drilled 2.5cm from the edge of the

circle. This experiment was done in light room conditions to increase the motivation to escape.

The volume of the white noise was also increased (90dB). Four visual cues were placed on each

side of the platform, which consisted of brightly colored shapes (purple diamond, yellow star,

green cube, red triangle).

The mouse was first habituated to the escape tunnel, a small box filled with Alphadry

bedding, in its home cage for 1 minute. It was then habituated to the experimental conditions. The

mouse was placed in the center of the apparatus and allowed to explore until it entered the escape

tunnel or 5 minutes elapsed. If the mouse did not organically reach the escape tunnel within the

time limit, it was led to the exit. There was at least 1 hour between habituation and the onset of

acquisition training. The escape tunnel was moved between habituation and training trials.

There were 4 training trials per day for 4 consecutive days. There was an inter-trial interval

of 30 minutes. During training trials, the position of the escape tunnel remained at a fixed location

relative to the spatial cues. The mouse was placed in the center of the platform and tracked until

it fully entered the escape tunnel or 5 minutes elapsed. If the mouse failed to reach the escape

tunnel in the allotted time, it was led to the escape by the researcher. Following each trial, the

platform was cleaned with 70% alcohol and the bedding inside the escape chamber was replaced.

For each trial, several variables were tracked to assess performance. These include

distance traveled, latency to exit, incorrect hole visits (primary errors), and incorrect hole revisits

(secondary errors). Errors are defined as nose point entering a hole that does not contain the exit.

Finally, based on these errors, the search strategy was characterized. Mice searched the maze

serially, spatially, or randomly. Serial searching mice spent most of the time on the periphery

performing a systematic search of the holes in a clockwise or counterclockwise manner, with 2 or

less direction changes. Spatial search strategy was defined as <10 primary errors and a

significantly reduced path and latency to exit. These mice use the surroundings to determine the

shortest path to the exit and often are within 1-2 holes of the destination. All other results were

defined as random search strategy. Random search strategy was usually characterized by

multiple direction changes and skipping between holes, with many primary and secondary errors.

24 hours after the last training trial, mice were tested with a probe trial. The escape tunnel

was removed before the mouse was placed on the platform for 5 minutes. Total time spent in

each quartile was recorded, along with visits to former correct and incorrect escapes.

Light/Dark Box

Mice were placed in the light side of a Light/Dark box apparatus and movement within the

apparatus was recorded from above for 10 minutes. The "light" side of the box had translucent

walls and no ceiling, illuminated by the overhead room light. The "dark" side had opaque walls

and a visible-light-filtering ceiling, allowing the mouse to experience a dark environment but the

infrared camera to continue recording. We recorded the latency to enter the dark, transitions

between zones, and time spent in zones.

Elevated Plus Maze

The elevated plus maze apparatus is a white "+"-shaped platform elevated 50cm above the floor with oppositely positioned "open" arms and "closed" arms. The closed arms are enclosed by 30cm high opaque walls, while the open arms have no railing. Each mouse was tested for 5 min after being placed onto the center platform facing an open arm. The number of open and closed arm entries and the time spent on the various sections of the EPM was recorded. The amount of exploration into the open arm is calculated to measure anxiety-like behavior.

Stereotaxic Surgery

Mice underwent stereotaxic surgery as described in (Ehinger et al., 2021). Specifically, mice were anesthetized by vaporized isoflurane and were then headfixed in a stereotaxic frame (David Kopf Instruments). The experimental virus, or relevant control, was infused into the dorsal hippocampus (anteroposterior (AP) -2.3, mediolateral (ML) ±1.7, dorsoventral (DV) -1.7 mm measured from bregma) using stainless steel injectors (33 gauge; Small Parts Incorporated) connected to Hamilton syringes (10 µl, 1701). Animals received 1µl of virus bilaterally at a rate of 0.1µl/min controlled by an automatic pump (Harvard Apparatus). After the infusion was complete, the injectors remained at the site for 10 minutes to allow diffusion of the virus. Mice were allowed to recover in their home cages for at least 3 weeks before testing to allow for maximal viral expression.

Confirmation of Viral Expression

At the end of the experimental timeline, animals were euthanized via cervical dislocation and the brains were removed. The brains were placed on ice and dissected into 1-mm coronal sections. The fluorescent protein expressed by the virus (either GFP or mCherry) was visualized using an EVOS FL tabletop fluorescent microscope (ThermoFisher Scientific). Images were taken

for future reference. Animals that failed to exhibit fluorescence associated with viral

overexpression were excluded from the study (4 mice).

Statistical Analysis

Biochemical analysis Parametric tests were performed on data deemed by the D'Agostino-

Pearson omnibus (K2) test to be derived from a normally distributed population. Data were

analyzed using a two-tailed t-test with Welch's correction for normal populations. Mann-Whitney

tests were performed on data derived from non-normal populations. The results were determined

to be statistically significant if the p-value was less than 0.05.

Electrophysiology analysis Parametric tests were performed on data deemed to be

derived from a normally distributed population. Unpaired t-tests were performed on the average

measurement over the experimental period. When multiple intensities were used, a two-way

repeated measures ANOVA was performed. *Post hoc* pairwise comparisons (Tukey's) were

performed after a significant result in the ANOVA. The results were determined to be statistically

significant if the p-value was less than 0.05.

Behavioral analysis Parametric tests were performed on data deemed to be derived from

a normally distributed population. Behavioral experiments were first assessed using a three-way

ANOVA with primary variables being genotype, sex, and experimental variable where appropriate.

Where there was no significant difference between sexes, the data was consolidated by genotype.

A two-way ANOVA was then performed on the consolidated data (genotype x experimental

variable). *Post hoc* Šidák's multiple comparison's test was performed to measure experimental

differences directly. When there was only one experimental variable, an unpaired t-test was

performed. If the variance was unequal, Welch's correction was applied. The results were

determined to be statistically significant if the p-value was less than 0.05.

Results

Generation and characterization of Prosapip1(flx/flx) mice

To elucidate the role of Prosapip1 in the CNS, we generated a mouse line with flox sites flanking the Prosapip1 gene. Guide RNA binding sites were identified in intron 2 and in the 3' UTR of exon 5 of Prosapip1 (*Lzts3*) (**Figure 1A**). PAGE purified Ultramer single stranded DNA oligos that were homologous to the target loci in intron 2 and exon 5 (**Figure 1B**) were used as repair templates.

To construct a conditional knockout of Prosapip1 in neurons, we crossed the homozygous Prosapip1 floxed (Prosapip1(flx/flx)) mouse with a mouse line expressing Cre recombinase under the control of a synapsin promoter (Syn1-Cre(+)). Male Prosapip1(flx/flx);Syn1-Cre(-/-) mice were mated with female Prosapip1(flx/flx);Syn1-Cre(+/-) mice (**Figure 1C**), which generated Prosapip1(flx/flx);Syn1-Cre(+) and Prosapip1(flx/flx);Syn1-Cre(-) mice, which were used as controls. The neuronal knockout was confirmed using western blot analysis. As shown in **Figure 1D**, Prosapip1(flx/flx);Syn1-Cre(-) mice have Prosapip1 protein levels comparable to a C57BL/6 mouse, while Prosapip1(flx/flx);Syn1-Cre(+) show a complete knockout of Prosapip1 protein in the dHP.

We examined the breeding history of the newly developed line to identify any abnormalities. The litter sizes of Prosapip1(flx/flx);Syn1-Cre matings followed a normal distribution with an average litter size between 5 and 6 (**Figure 1E, Table 1**). The proportion of male and female pups were near expectations, with 52% of pups being male and 48% female (**Figure 1F**). Inheritance of Syn1-Cre(+) was also split evenly, with 53% of pups from the line being born with the transgene (**Figure 1G**). Body weight of male and female mice was monitored every 2 weeks from birth to assess potential developmental effects. While males were significantly heavier than females throughout development, there was no significant effect of the genotype on weight over time (**Figure 1H, Table 1**).

We next assessed baseline locomotor and exploratory behavior of these mice. Upon

reaching adulthood, mice were placed in an open field and allowed to explore for 20 minutes, and

total distance traveled and time spent in the center of the open field were measured. Deletion of

neuronal Prosapip1 did not alter locomotor activity (**Figure 1I, Table 1**). Furthermore, there were

no differences between genotypes when assessing the time spent in the center of the open field

(**Figure 1J, Table 1**).

Prosapip1 is required for synaptic localization of PSD proteins

Prosapip1 is a scaffolding protein that recruits SPAR to the PSD and interconnects it with

Shank3 in primary hippocampal neurons (Wendholt et al., 2006; Reim et al., 2016). First, we

examined if the synaptic localization of SPAR was affected by Prosapip1 neuronal knockout. To

do this, we isolated the synaptic fraction from the total homogenate. We confirmed the efficacy of

the crude synaptosomal fraction isolation by measuring enrichment of synapsin, a protein found

at the synapse, and lack of cAMP response element-binding protein (CREB), a transcription factor

located in the nucleus (**Figure 2A**). We analyzed the total (**Figure 2B**) and synaptic (**Figure 2C**)

levels of SPAR in the dHP of Prosapip1(flx/flx);Syn1-Cre(-) and Prosapip1(flx/flx);Syn1-Cre(+)

male and female mice. Total levels of SPAR were similar between the groups (**Figure 2B, Table**

1). However, we detected a significant reduction in synaptic SPAR in Prosapip1(flx/flx);Syn1-

Cre(+) mice as compared to the Prosapip1(flx/flx);Syn1-Cre(-) group (**Figure 2C, Table 1**).

Prosapip1 interconnects SPAR to Shank3, and SPAR forms a complex with PSD-95

(Wendholt et al., 2006; Reim et al., 2016; Matsuura et al., 2022). Therefore, we next examined

the total and synaptic levels of Shank3 and PSD-95. Prosapip1 deletion did not affect the total or

synaptic levels of Shank3 (**Figure 2D-G, Table 1**). We also found that Prosapip1(flx/flx);Syn1-

Cre(+) mice have similar levels of total PSD-95 (**Figure 2D, 2H, Table 1**) compared to

Prosapip1(flx/flx);Syn1-Cre(-) controls. However, when examining synaptic PSD-95,

Prosapip1(flx/flx);Syn1-Cre(+) mice displayed drastically reduced levels compared to controls

(**Figure 2E, 2I, Table 1**). Collectively, these results imply that Prosapip1 in the dHP is necessary

for synaptic localization of SPAR and PSD-95, but not Shank3.

SPAR and Shank3 are both known to interact with the NMDAR and AMPAR, and PSD-95

stabilizes NMDAR surface localization (Naisbitt et al., 1999; Tu et al., 1999; Pak et al., 2001;

Boeckers et al., 2002; Arons et al., 2012; Won et al., 2016; Coley and Gao, 2019; Matsuura et al.,

2022). As the synaptic localization of SPAR and PSD-95 is disrupted in the absence of Prosapip1,

we next tested the synaptic levels of NMDAR subunits GluN2A and GluN2B, along with AMPAR

subunit GluA1. There was no change in the overall levels of the NMDAR subunits GluN2A and

GluN2B or the AMPAR subunit GluA1 (**Figure 2J, 2L, 2N, 2P, Table 1**). The synaptic levels of

GluN2A and GluA1 were also unaltered in the dHP of Prosapip1(flx/flx);Syn1-Cre(+) mice

compared to Prosapip1(flx/flx);Syn1-Cre(-) controls (**Figure 2K, 2M, 2Q, Table 1**). In contrast, the

synaptic levels of NMDAR subunit GluN2B were significantly reduced in Prosapip1(flx/flx);Syn1-

Cre(+) mice (**Figure 2K, 2Q, Table 1**). Together, Prosapip1 controls the synaptic localization of

NMDAR subunit GluN2B, but not GluN2A or GluA1, in the dHP.

Prosapip1 in the dorsal hippocampus plays a role in NMDA receptor-mediated

transmission and long-term potentiation

The GluN2B subunit of the NMDAR is critically involved in learning and memory, a

process reliant on LTP (Nachtigall et al., 2024) Therefore, we examined the contribution of

Prosapip1 to GluN2B-mediated LTP. To examine this, hippocampal slices from male and female

Prosapip1(flx/flx);Syn1-Cre(-) and Prosapip1(flx/flx);Syn1-Cre(+) mice were prepared for

electrophysiology recordings. Bipolar stimulating electrodes were positioned in the CA1 region of

the dHP. Simultaneously, the recording electrode was placed approximately 100-150 μm away

from the stimulating electrode (**Figure 3A**). The administration of a single electrical pulse (2 ms)

elicited a fiber volley, followed by a field excitatory postsynaptic potential response (fEPSPs)

(**Figure 3B**). Before inducing LTP, a stable baseline of fEPSPs was established for 10 minutes

(**Figure 3C**). Upon establishing a baseline of fEPSPs, high-frequency stimulation (HFS) was

applied at 100 Hz for 1 second, repeated four times at 20-second intervals, to trigger LTP. In the

control Prosapip1(flx/flx);Syn1-Cre(-) mice, there was a significant increase in the fEPSP

amplitude following HFS, indicating a successful LTP induction (**Figure 3C**), which persisted for

at least 30 minutes (**Figure 3C**). However, LTP was absent in the Prosapip1(flx/flx);Syn1-Cre(+)

mice (**Figure 3C**). A comparative analysis of the normalized fEPSPs, conducted 20-30 minutes

post-HFS, revealed a significantly reduced fEPSP amplitude in the Prosapip1(flx/flx);Syn1-Cre(+)

group compared to the Prosapip1(flx/flx);Syn1-Cre(-) controls (**Figure 3D, Table 1**). Together

these findings strongly suggest Prosapip1 plays an important role in dHP LTP.

LTP induction in the hippocampus is known to rely on the functioning of postsynaptic

NMDARs. Therefore, we aimed to discern if there were any differences between

Prosapip1(flx/flx);Syn1-Cre(-) and Prosapip1(flx/flx);Syn1-Cre(+) mice in terms of NMDAR-

mediated excitatory postsynaptic currents (EPSCs) in dHP CA1 pyramidal neurons. A whole-cell

voltage-clamp recording technique was employed for this purpose. First, a low concentration of

external Mg^{2+} (0.05 mM) was used to facilitate the removal of the magnesium block from the

NMDAR channel. Concurrently, AMPARs and $GABA_A$ receptors ($GABA_ARs$) were

pharmacologically inhibited. NMDAR-EPSCs were recorded in response to an escalating series

of stimulus intensities. The amplitude of NMDAR-EPSCs was significantly reduced in the

Prosapip1(flx/flx);Syn1-Cre(+) mice compared to the controls (**Figure 3E, Table 1**). Next, we

investigated whether LTP inhibition because of Prosapip1 deletion is due to post and/or

presynaptic changes. First, we investigated postsynaptic changes by measuring NMDA-induced

currents. NMDA (20 µM) was bath applied for 30 seconds and peak current was measured. We

observed that the peak current in the Prosapip1(flx/flx);Syn1-Cre(+) mice was markedly lower

than that in their control counterparts (**Figure 3F, Table 1**). These data suggest that postsynaptic

NMDAR responsiveness was decreased the in the dHP of Prosapip1(flx/flx);Syn1-Cre(+) mice as

compared to Prosapip1(flx/flx);Syn1-Cre(-) mice. Finally, to examine the presence of any

presynaptic alterations in glutamatergic transmission in CA1 neurons upon Prosapip1 deletion,

we measured the paired-pulse ratio (PPR, 100 millisecond interval) of electrically evoked EPSCs

in CA1 dHP neurons. Surprisingly we found that PPR was significantly higher in the

Prosapip1(flx/flx);Syn1-Cre(+) group as compared to Prosapip1(flx/flx);Syn1-Cre(-) mice (**Figure**

3G, Table 1), indicating a probable reduction in presynaptic glutamate release onto CA1 neurons

in the Prosapip1(flx/flx);Syn1-Cre(+) mice (Zucker and Regehr, 2002). Together, the data imply

that the Prosapip1 is crucial for the induction of hippocampal LTP, which is likely attributed to the

downregulation of NMDAR function.

Prosapip1 contributes to spatial memory

Prosapip1 was originally identified in hippocampal neurons (Wendholt et al., 2006; Reim

et al., 2016), a brain region associated with memory (Broadbent et al., 2004). Given the altered

PSD composition and subsequent loss of LTP in this region after Prosapip1 knockout, we

hypothesized that Prosapip1 contributes to memory-dependent behavior. First, we tested whether

Prosapip1 contributes to long-term recognition memory in male and female mice using the novel

object recognition (NOR) test with an inter-trial interval of 24 hours (Leger et al., 2013; Lueptow,

2017). Prosapip1(flx/flx);Syn1-Cre(-) animals showed a significant preference for the novel object,

indicating long-term memory of the familiar object and its spatial location (**Figure 4A, Table 1**).

Contrarily, male and female Prosapip1(flx/flx);Syn1-Cre(+) mice spent a similar amount of time

exploring both objects, performing at chance levels (**Figure 4A, Table 1**). The lack of novel object

recognition in Prosapip1 knockout mice implies a loss of object characteristic acquisition or long-

term recognition memory.

To determine whether Prosapip1 contributes to spatial learning or working memory, male

and female mice performed a novelty T-maze test (d'Isa et al., 2021). Mice were allowed to

explore two arms of a "T"-shaped maze during training trials while the third arm was blocked. All

three arms were available during the test. The experiment utilized a 1-minute inter-trial interval,

performance of which is reliant on hippocampal LTP (Sanderson et al., 2009). During the test trial,

Prosapip1(flx/flx);Syn1-Cre(-) mice primarily explored the novel arm of the T-maze, indicated by

a positive difference score (time spent in novel arm – time spent in familiar arm) (**Figure 4B,**

Table 1). Prosapip1(flx/flx);Syn1-Cre(+) mice explored the arms at comparable levels, performing

significantly worse than control animals, and indicating a lack of spatial learning for the familiar

arm of the maze (**Figure 4B, Table 1**). This difference can be seen visually in the heatmap

(**Figure 4C**). Taken together, these data suggest that Prosapip1 is required for spatial learning

and working memory.

We next utilized the 3-chamber social interaction (3CSI) test. The 3CSI test has two

stages. The first part measures the preference for social interaction with a juvenile interaction

partner.   We   found   that   male   and   female   Prosapip1(flx/flx);Syn1-Cre(-)   and

Prosapip1(flx/flx);Syn1-Cre(+) mice behave similarly, with both significantly preferring the

interaction partner over the empty chamber (**Figure 4D, Table 1**). In the second stage of the

3CSI, the mouse has the choice between interacting with a familiar partner (partner from part 1)

or a novel interaction partner. This assesses the social novelty recognition of the subject.

Prosapip1(flx/flx);Syn1-Cre(-) mice displayed a preference for the novel interaction partner,

exhibited by the increase in time spent in close proximity to the novel partner (**Figure 4E, Table**

1). Prosapip1(flx/flx);Syn1-Cre(+) mice spent a similar amount of time with the familiar and novel

interaction partners (**Figure 4E, Table 1**). The lack of discrimination from Prosapip1(flx/flx);Syn1-

Cre(+) mice implies a loss of recognition of social novelty or loss of social memory.

While the Prosapip1(flx/flx);Syn1-Cre(+) mice show impaired performance in the NOR and

novelty T-maze tests, these results may be attributed to the learning/acquisition process or

storage of memory. To confirm that Prosapip1 is important for processes underlying spatial

learning and memory, we employed the Barnes maze (Pitts, 2018). The Barnes maze is a white

plastic circle with holes evenly drilled around the perimeter. Underneath one of these holes is an

exit tunnel. The protocol has a training and testing phase, where spatial learning and memory,

respectively, are separately examined. First, mice were habituated to the platform. There were no

differences in baseline exploratory profile between Prosapip1(flx/flx);Syn1-Cre(-) and

Prosapip1(flx/flx);Syn1-Cre(+) mice (**Extended Figure 4A, Table 1**). The primary variable

examined during training trials was the path to escape, which tracked the distance traveled from

the starting point to the exit compartment. A shorter path to escape is more efficient and indicates

learning of the paradigm objective and exit location. Prosapip1(flx/flx);Syn1-Cre(-) mice

immediately improve performance, with a significant decrease in path to escape observed as

soon as the third trial (**Figure 4F, Table 1**). Prosapip1(flx/flx);Syn1-Cre(+) mice improved

performance over time but more gradually and performed significantly worse than control mice

over the entirety of training (**Figure 4F, Table 1**). Prosapip1(flx/flx);Syn1-Cre(-) mice showed a

significantly reduced path to escape by the second trial (**Figure 4F, Table 1**). In addition,

Prosapip1(flx/flx);Syn1-Cre(+) mice committed significantly more primary (incorrect hole visit) and

secondary (incorrect hole revisit) errors throughout training (**Figure 4G, Table 1**).

Differences in training performance can be explained by examining the search strategy of

each mouse. There are three defined search strategies: random, serial, and spatial, which

correspond to an increase in task efficiency (**Figure 4H**) (Rosenfeld and Ferguson, 2014). Mice

searching serially methodically inspected holes with minimal directional changes (Rosenfeld and

Ferguson, 2014). Spatial searchers used environmental cues to navigate efficiently to the exit

with less than 10 primary errors (Rosenfeld and Ferguson, 2014). In contrast, random search

strategies involved frequent direction changes and errors, with mice often skipping between holes

(Rosenfeld and Ferguson, 2014). Prosapip1(flx/flx);Syn1-Cre(-) mice mainly searched randomly

during the first trial, but 66% of mice utilized a serial search strategy by the second trial (**Figure**

4I). Spatial searching appeared as early as trial 3, and its ratio increased until it had become a

majority on the last day of training, with 77% of mice spatially searching on the final trial (**Figure**

4I). In the Prosapip1(flx/flx);Syn1-Cre(+) mice, randomly searching was a prevalent strategy

throughout training, with 39% of all trials being classified as random (**Figure 4J, Table 1**). More

than 50% of Prosapip1(flx/flx);Syn1-Cre(+) mice switched to serial search strategy by the third

day of training, which also explains the gradual decrease in path to escape (**Figure 4J, Table 1**).

Interestingly, less than 2% of all trials completed by Prosapip1(flx/flx);Syn1-Cre(+) mice utilized

the spatial search strategy (**Figure 4J, Table 1**). Both groups remember the context of the Barnes

maze and the objective to escape, but Prosapip1(flx/flx);Syn1-Cre(+) mice fail to remember the

spatial location of the exit. These results confirm the conclusion that Prosapip1 is required for

spatial learning.

The probe trial of the Barnes maze specifically assesses the spatial memory of the mouse.

During the probe trial, the exit was removed, and the mouse was allowed to explore the platform

for 5 minutes. If the mouse retained spatial memory of the exit, it spent significantly more time

exploring the quartile associated with the exit compartment compared to chance (25% of the

time). Control mice performed above chance value, spending an average of 116.84 seconds in

the exit quartile (**Figure 4K, Table 1**). Prosapip1(flx/flx);Syn1-Cre(+) mice, conversely, performed

around chance, spending an average of 68.52 seconds in the exit quartile (**Figure 4K, Table 1**).

This was significantly less than control mice and can be observed in the probe trial heat map

(**Figure 4L**). Thus, Prosapip1 is required for spatial memory in the Barnes maze.

To exclude the possibility that Prosapip1(flx/flx);Syn1-Cre(+) mice preferred familiar

situations compared to novel situations due to an increase in anxiety, we used the light/dark box

and elevated plus maze paradigms to assess the baseline anxiety of these mice. The light/dark

box consists of two chambers where the subject may choose to explore the well-lit ("light") or

visible-light-blocked ("dark") side of a chamber (Bourin and Hascoet, 2003). Spending less time

in the light side of the chamber implies an increase in anxiety (Bourin and Hascoet, 2003). Both

Prosapip1(flx/flx); Syn1-Cre(-) and Prosapip1(flx/flx);Syn1-Cre(+) mice spent significantly more

time in the dark side compared to the light side, but there was no significant difference between

the genotypes (**Extended Figure 4B, Table 1**). Mice also underwent the elevated plus maze test.

The elevated plus maze is elevated above the floor and consists of two "closed" arms with opaque

walls and two "open" arms with no railing. Time exploring the open arm is used to assess anxiety

(Walf and Frye, 2007). Prosapip1(flx/flx);Syn1-Cre(-) and Prosapip1(flx/flx);Syn1-Cre(+) mice

spent a similar amount of time exploring the open arm (**Extended Figure 4C, Table 1**). Taken

together, our data suggest that Prosapip1 is not involved in anxiety-related behaviors.

Prosapip1 in the dorsal hippocampus contributes to recognition, social, and spatial

memory

Memory is often associated with the dHP (Pilly and Grossberg, 2012). As our data suggest

that Prosapip1 contributes to recognition, social, and spatial memory, we next determined

whether the dHP is the loci of Prosapip1's contribution to learning and memory processes. To do

so, the dHP of Prosapip1(flx/flx) mice was infected with an adeno-associated virus (AAV)

expressing Cre to knockout Prospaip1 specifically in this region. An AAV expressing solely GFP

was used as a control (**Figure 5A**). Western blot analysis of the AAV-Cre-infected dHP shows

efficient deletion of Prosapip1 protein (**Figure 5B**).

We repeated the experimental battery performed on the genetically manipulated mice on

the mice infused with AAV-GFP or AAV-Cre in the dHP. In the open field, AAV-GFP- and AAV-

Cre-infected mice exhibited similar levels of locomotion (**Figure 5C, Table 1**) and spent a similar

amount of time in the center of the field (**Figure 5D, Table 1**). The region-specific knockout of

Prosapip1 did not affect exploratory behavior. We also directly assessed anxiety-like behavior

using the light/dark box. Mice infected with AAV-GFP or AAV-Cre both spent comparable amounts

of time in the light and dark chamber, preferring the dark (**Figure 5E, Table 1**). Therefore,

Prosapip1 in the dHP does not control locomotion or anxiety-like behavior.

Next, we examined the role of Prosapip1 in the dHP on memory. We first assessed

recognition memory using the novel object recognition test. The AAV-GFP-infected mice showed

a significant preference for the novel object, indicating functioning long-term recognition memory

(**Figure 5F, Table 1**). Mice infected with AAV-Cre, however, showed no significant preference for

either object (**Figure 5F, Table 1**). Similarly, in the novelty T-maze test, AAV-GFP-infected mice

spent much of the test exploring the novel arm of the maze, indicated by a positive difference

score (**Figure 5G, Table 1**). AAV-Cre-infected mice spent significantly less time exploring the

novel arm of the maze, performing at chance levels (**Figure 5G, Table 1**). Together, these results

suggest that Prosapip1 in the dHP specifically contributes to recognition and spatial working

memory.

We also assessed the effects of region-specific Prosapip1 knockout on sociability and

social memory using the 3-chamber social interaction test. Both AAV-GFP- and AAV-Cre-infected

mice showed a significant preference for interacting with a social partner over the empty cage

(**Figure 5H, Table 1**). When a novel interaction partner was introduced, AAV-GFP mice

significantly preferred interacting with the novel partner (**Figure 5I, Table 1**). AAV-Cre-infected

mice did not show a significant preference (**Figure 5I, Table 1**).

Finally, AAV-GFP- and AAV-Cre-infected mice underwent the Barnes maze procedure to

separately assess spatial learning and memory. During training trials, AAV-Cre-infected mice

performed significantly worse when examining path to escape (**Figure 5J, Table 1**). While AAV-

GFP-infected mice quickly improved exit strategy, AAV-Cre-infected mice were more variable and

took a longer path, on average (**Figure 5J, Table 1**). After training, during the probe trial, that

AAV-GFP-infected mice spent significantly more time in the exit quartile compared to AAV-Cre-

infected mice (**Figure 5K, Table 1**). Together, our data suggests that Prosapip1 in the dHP is

specifically involved in the formation of spatial memory.

Discussion

Our findings suggest that Prosapip1 in the dHP is responsible for the synaptic localization of SPAR and PSD-95, and consequently, for the membranal localization of GluN2B. Furthermore, our data suggest that Prosapip1 in the dHP plays an important role in LTP as well as learning and memory.

Prosapip1 is a key protein in the PSD

Prosapip1 is highly enriched in the PSD of primary hippocampal neurons (Wendholt et al., 2006; Reim et al., 2016), and our results suggest that Prosapip1 could be crucial to the formation and stability of PSD complexes. Prosapip1 scaffolds SPAR to Shank3 in the PSD (Wendholt et al., 2006; Dolnik et al., 2016; Reim et al., 2016). We found that knockout of Prosapip1 reduced the synaptic localization of SPAR but not Shank3. Reim et al. reported that Prosapip1 is directly regulating SPAR levels in the PSD in primary hippocampal neurons (Reim et al., 2016). Our data supports this study, as Prosapip1 knockout *in vivo* led to a reduction of synaptic SPAR. SPAR plays a role in the formation of F-actin, which is important for dendritic spine maturation (Pak et al., 2001). We previously showed Prosapip1 interacts with SPAR in the NAc of mice (Laguesse et al., 2017), and Prosapip1 knockdown reduces F-actin content while overexpression increases F-actin content in the NAc (Laguesse et al., 2017). Furthermore, we showed that Prosapip1 plays a role in spine maturation in the NAc (Laguesse et al., 2017). Our results suggest that the recruitment of SPAR to the PSD via Prosapip1 in the dHP will promote F-actin formation and lead to dendritic spine maturation, however these possibilities merit further examined.

Both SPAR and Shank3 interact with PSD-95 (Naisbitt et al., 1999; Tu et al., 1999; Pak et al., 2001). SPAR binds PSD-95 directly (Pak et al., 2001), and Shank3 associates with Guanylate kinase-associated protein (GKAP), which binds PSD-95 (Naisbitt et al., 1999; Tu et al., 1999).

PSD-95 binds to NMDAR subunits GluN2A and GluN2B and stabilizes NMDAR surface

localization (Won et al., 2016; Coley and Gao, 2019). Studies have shown that PSD-95 deficiency

causes an imbalance of NMDAR and AMPAR synaptic presence, altering glutamatergic

transmission (Lin et al., 2004; Elias et al., 2006; Zhang et al., 2013; Chen et al., 2015).

Surprisingly, we discovered that loss of Prosapip1 resulted in a marked reduction of synaptic

PSD-95, and a correlated reduction in the synaptic compartmentalization of the NMDAR subunit

GluN2B, raising the possibility that the interaction between Prosapip1 and PSD-95 is required for

synaptic membranal localization of GluN2B. However, we did not observe a change in the

synaptic level of GluN2A. In addition, we did not find that the localization of the AMPAR subunit

GluA1 was altered as a result of Prosapip1 knockout. In line with our findings, Pak et al. showed

that SPAR binds PSD-95 and GluN2B, but not GluA1 (Pak et al., 2001). Based on our results and

these studies, the complex controlling AMPAR synaptic localization is different, and likely

independent of Prosapip1 in the dHP. Together, our data suggest that Prosapip1 is necessary for

the formation and/or maintenance of the NMDAR-associated PSD network.

We previously discovered that Prosapip1's translation depends on mTORC1 in the NAc

of mice consuming excessive levels of alcohol, which in turn promotes F-actin formation, dendritic

spine maturation and alcohol-reward memory and the reinforcement of alcohol consumption

(Laguesse et al., 2017). mTORC1 activation by alcohol also induces the translation of the

scaffolding protein Homer and PSD-95 in the NAc of mice (Beckley et al., 2016; Liu et al., 2017).

mTORC1 plays an essential role in the translation of a subset of mRNAs to proteins. In the CNS,

mTORC1 promotes the translation of synaptic proteins at dendrites (Buffington et al., 2014), and

plays an important LTP, learning, and memory (Hoeffer and Klann, 2010; Graber et al., 2013).

Therefore, it would be of great interest to test the hypothesis that learning-dependent activation

of mTORC1 in the dHP activates the translational machinery at synapses and therefore increases

the translation of Prosapip1 which in turn recruits proteins such as PSD-95 to stabilize the PSD-

95 complex.

Prosapip1 is required for LTP in the dHP

In our study, we discovered that mice lacking Prosapip1 exhibited markedly diminished

NMDAR-mediated synaptic transmission, which is in line with the fact that there is reduced

synaptic GluN2B. We also found that Prosapip1 knockout led to a significant reduction in the

formation of LTP. LTP is a persistent increase in synaptic strength between neurons that can last

for months *in vivo* (Abraham et al., 2002; Cooke and Bliss, 2006) and is the cellular hallmark of

learning and memory. It is often induced by HFS, which causes extensive glutamate release and

significant AMPAR activity, resulting in large membrane depolarization (Ma et al., 2018). This

depolarization facilitates the removal of Mg^{2+} blockage in NMDARs, thereby activating them and

leading to calcium influx (Ma et al., 2018). This influx triggers the calmodulin/CaMKII pathway,

initiating a series of cascades that enhance AMPAR phosphorylation and trafficking, culminating

in sustained synaptic strengthening (Huganir and Nicoll, 2013; Herring and Nicoll, 2016).

Therefore, the induction of hippocampal LTP heavily relies on NMDAR activation, with alterations

in NMDAR function significantly affecting the establishment of hippocampal LTP and its

associated learning and memory processes. In Prosapip1 knockout mice, the reduction in

NMDAR functionality likely results in decreased calcium influx during high-frequency stimulation

and diminished activation of the calmodulin/CaMKII pathway, leading to weaker synaptic

strengthening.

Prosapip1 is required for learning and memory

As detailed above, we found that Prosapip1 neuronal knockout mice have deficits in LTP in the dHP, and LTP is the hallmark of learning and memory (Bliss and Collingridge, 1993; Frey and Morris, 1998; Collingridge et al., 2004). We also found that global neuronal knockout and dHP-specific knockout of Prosapip1 results in deficits in spatial learning and memory. Specifically, we showed that Prosapip1 in the dHP is required for normal performance in the novel object recognition, novelty T-maze, 3-chamber social interaction, and Barnes maze tests, which all require a unique form of learning and memory (Davis et al., 1992; Broadbent et al., 2010; Barker and Warburton, 2011; Rosenfeld and Ferguson, 2014; Lueptow, 2017; Pitts, 2018; Sanchez-Rodriguez et al., 2022). The novel object recognition test is primarily examining recognition memory, and with an inter-trial interval of 24 hours, specifically long-term memory (Broadbent et al., 2010; Barker and Warburton, 2011; Antunes and Biala, 2012; Lueptow, 2017; Cinalli et al., 2020). The novelty T-maze focuses on both spatial learning and memory, and with its short, 1-minute inter-trial interval, this procedure is examining spatial working memory (Sharma et al., 2010; d'Isa et al., 2021). The 3-chamber social interaction test is assessing baseline sociability and also social recognition memory (Meira et al., 2018; Tzakis and Holahan, 2019; Wang and Zhan, 2022; Cope et al., 2023; Wei et al., 2024). Finally, the Barnes maze tests both spatial learning and working memory in intra-day trials, while simultaneously testing long-term contextual and spatial memory in inter-day trials and the probe test (Bach et al., 1995; Sharma et al., 2010; Rosenfeld and Ferguson, 2014; Pitts, 2018). Interestingly, loss of hippocampal LTP has been shown to impair spatial, but not contextual memory in the Barnes maze (Bach et al., 1995). As presented here, Prosapip1 knockout mice significantly reduced distance traveled to exit but did not switch to spatial searching. In this example, Prosapip1 knockout mice are retaining the contextual understanding of escaping the platform, but do not recall the spatial location of the exit. Additionally, the link between NMDAR function and learning and memory is well established

(Newcomer et al., 2000; Li and Tsien, 2009). For example, blockage of hippocampal NMDA

receptors impairs spatial learning in rats (Davis et al., 1992; Bye and McDonald, 2019). Prosapip1

is likely controlling the reinforcement of learning and memory by PSD scaffolding, stabilization,

and GluN2B synaptic localization, leading to LTP.

We observed that Prosapip1 knockout specifically in the dHP replicated the recognition,

social, and spatial learning and memory deficits exhibited by the global neuronal knockout mice,

suggesting that Prosapip1 is controlling these learning and memory processes specifically in the

dHP. The dHP primarily controls memory formation and recall (Eichenbaum, 1997; Broadbent et

al., 2004; Squire et al., 2004; Pilly and Grossberg, 2012). We found that LTP in the CA1 subregion

of the dHP was reliant on Prosapip1. The CA1 subregion is critically involved in contextual

memory, object recognition memory, and spatial memory (Tsien et al., 1996; Lee and Kesner,

2004; Daumas et al., 2005; Sanderson et al., 2009; Sharma et al., 2010; Stevenson et al., 2018;

Bye and McDonald, 2019; Cinalli et al., 2020; Jeong and Singer, 2022). Loss of Prosapip1 in CA1

is likely leading to decreased performance in the novel object recognition, novelty T-maze, and

Barnes maze tests. The lack of social recognition displayed by Prosapip1(flx/flx);Syn1-Cre(+)

mice and AAV-Cre-infected mice is likely attributed to the loss of Prosapip1 in the CA2 subregion

of the dHP, which is the primary subregion controlling social recognition memory (Meira et al.,

2018; Tzakis and Holahan, 2019; Wang and Zhan, 2022; Cope et al., 2023; Wei et al., 2024).

Specifically, silencing the CA2 subregion of the dHP impairs social memory formation and

consolidation (Meira et al., 2018). However, the CA3 and DG subregions of the dHP are also

involved in spatial and contextual memory (Broadbent et al., 2004; Lee and Kesner, 2004;

Daumas et al., 2005). As our conditional knockout strategy resulted in Prosapip1 deletion from

the whole dHP, further studies are required to dissect the subregion specificity of the contribution

of Prosapip1 to recognition, social, and spatial learning and memory processes.

Memory consists of three primary processes: encoding, consolidation, and retrieval

(Straube, 2012). In this study, the defect in memory function is likely due to a failure to encode

new information (Bye and McDonald, 2019) or consolidate this "short-term" into "long-term"

memory (Yang et al., 2022). The spatial T-maze experiment utilized a short inter-trial interval of

one minute which requires working spatial memory (Sharma et al., 2010), and Prosapip1 knockout

mice exhibited a failure to encode new information. Similarly, the Barnes maze training trials were

separated by an inter-trial interval of 30 minutes, but Prosapip1 knockout mice did not acquire

spatial memory between training trials, nor during the longer consolidation periods between days,

again implying a failure to encode spatial information or consolidate this information. The lack of

synaptic localization of GluN2B is likely underlying the loss of memory encoding or consolidation

(Nachtigall et al., 2024). It is unlikely that NMDAR dysfunction is affecting the retrieval of memory,

as studies have exhibited rats' ability to use previously acquired spatial information during

NMDAR blockage (Bast et al., 2005; Mackes and Willner, 2006; Bye and McDonald, 2019).

Prosapip1 belongs to the Fezzin family of proteins (Wendholt et al., 2006). It is important

to note that other Fezzins do not compensate for the loss of Prosapip1 in the dHP. Knockout of

other Fezzins, like PSD-Zip70, also lead to cognitive deficits (Mayanagi et al., 2015). However,

these deficits were attributed to the action of PSD-Zip70 in the PFC. Therefore, one could

hypothesize that proteins in this family enact their function in specific brain subregions.

In summary, Prosapip1 in the dorsal hippocampus is integral to the synaptic localization

of SPAR, PSD-95, and GluN2B, which are required for the formation of LTP and subsequent

spatial learning and memory behavior. Abnormalities with PSD proteins are associated with

neuropsychiatric disorders (Kaizuka and Takumi, 2018), and further unraveling of the

physiological role of Prosapip1 may unlock insights into normal and abnormal mechanisms of

learning and memory.

Funding and Disclosure: Supported by NIH grant AA020889 (G.H.). ZWH was partially funded

by GM007175. None of the authors have a conflict of interest.

Acknowledgements: Some figures were created in part with BioRender.com. We thank Carolyn

Ferguson for expert technical assistance.

Author Contributions: DR conceived the project. GH developed the mouse line. ZWH, JW,

and DR designed the experiments. ZWH, HG, KP, and CS conducted the experiments and

analyzed the data. ZWH, HG, YE, JW, and DR wrote the manuscript.

References

Abraham WC, Logan B, Greenwood JM, Dragunow M (2002) Induction and experience-
dependent consolidation of stable long-term potentiation lasting months in the
hippocampus. J Neurosci 22:9626-9634.

Antunes M, Biala G (2012) The novel object recognition memory: neurobiology, test procedure,
and its modifications. Cogn Process 13:93-110.

Arons MH, Thynne CJ, Grabrucker AM, Li D, Schoen M, Cheyne JE, Boeckers TM, Montgomery
JM, Garner CC (2012) Autism-associated mutations in ProSAP2/Shank3 impair synaptic
transmission and neurexin-neuroligin-mediated transsynaptic signaling. J Neurosci
32:14966-14978.

Bach ME, Hawkins RD, Osman M, Kandel ER, Mayford M (1995) Impairment of spatial but not
contextual memory in CaMKII mutant mice with a selective loss of hippocampal LTP in the
range of the theta frequency. Cell 81:905-915.

Barker GR, Warburton EC (2011) When is the hippocampus involved in recognition memory? J
Neurosci 31:10721-10731.

Bast T, da Silva BM, Morris RG (2005) Distinct contributions of hippocampal NMDA and AMPA
receptors to encoding and retrieval of one-trial place memory. J Neurosci 25:5845-5856.

Beckley JT, Laguesse S, Phamluong K, Morisot N, Wegner SA, Ron D (2016) The First Alcohol
Drink Triggers mTORC1-Dependent Synaptic Plasticity in Nucleus Accumbens Dopamine
D1 Receptor Neurons. J Neurosci 36:701-713.

Bliss TV, Collingridge GL (1993) A synaptic model of memory: long-term potentiation in the
hippocampus. Nature 361:31-39.

Boeckers TM, Bockmann J, Kreutz MR, Gundelfinger ED (2002) ProSAP/Shank proteins - a
family of higher order organizing molecules of the postsynaptic density with an emerging
role in human neurological disease. J Neurochem 81:903-910.

Boeckers TM, Winter C, Smalla KH, Kreutz MR, Bockmann J, Seidenbecher C, Garner CC,

Gundelfinger ED (1999) Proline-rich synapse-associated proteins ProSAP1 and ProSAP2

interact with synaptic proteins of the SAPAP/GKAP family. Biochem Biophys Res

Commun 264:247-252.

Bourin M, Hascoet M (2003) The mouse light/dark box test. Eur J Pharmacol 463:55-65.

Broadbent NJ, Squire LR, Clark RE (2004) Spatial memory, recognition memory, and the

hippocampus. Proc Natl Acad Sci U S A 101:14515-14520.

Broadbent NJ, Gaskin S, Squire LR, Clark RE (2010) Object recognition memory and the rodent

hippocampus. Learn Mem 17:5-11.

Buffington SA, Huang W, Costa-Mattioli M (2014) Translational control in synaptic plasticity and

cognitive dysfunction. Annu Rev Neurosci 37:17-38.

Bye CM, McDonald RJ (2019) A Specific Role of Hippocampal NMDA Receptors and Arc Protein

in Rapid Encoding of Novel Environmental Representations and a More General Long-

Term Consolidation Function. Front Behav Neurosci 13:8.

Chen X, Levy JM, Hou A, Winters C, Azzam R, Sousa AA, Leapman RD, Nicoll RA, Reese TS

(2015) PSD-95 family MAGUKs are essential for anchoring AMPA and NMDA receptor

complexes at the postsynaptic density. Proc Natl Acad Sci U S A 112:E6983-6992.

Cinalli DA, Jr., Cohen SJ, Guthrie K, Stackman RW, Jr. (2020) Object Recognition Memory:

Distinct Yet Complementary Roles of the Mouse CA1 and Perirhinal Cortex. Front Mol

Neurosci 13:527543.

Coley AA, Gao WJ (2019) PSD-95 deficiency disrupts PFC-associated function and behavior

during neurodevelopment. Sci Rep 9:9486.

Collingridge GL, Isaac JT, Wang YT (2004) Receptor trafficking and synaptic plasticity. Nat Rev

Neurosci 5:952-962.

Concordet JP, Haeussler M (2018) CRISPOR: intuitive guide selection for CRISPR/Cas9 genome

editing experiments and screens. Nucleic Acids Res 46:W242-W245.

Cooke SF, Bliss TV (2006) Plasticity in the human central nervous system. Brain 129:1659-1673.

Cope EC, Wang SH, Waters RC, Gore IR, Vasquez B, Laham BJ, Gould E (2023) Activation of
816   the CA2-ventral CA1 pathway reverses social discrimination dysfunction in Shank3B
817   knockout mice. Nat Commun 14:1750.

818   d'Isa R, Comi G, Leocani L (2021) Apparatus design and behavioural testing protocol for the
819   evaluation of spatial working memory in mice through the spontaneous alternation T-
820   maze. Sci Rep 11:21177.

821   Daumas S, Halley H, Frances B, Lassalle JM (2005) Encoding, consolidation, and retrieval of
822   contextual memory: differential involvement of dorsal CA3 and CA1 hippocampal
823   subregions. Learn Mem 12:375-382.

824   Davis S, Butcher SP, Morris RG (1992) The NMDA receptor antagonist D-2-amino-5-
825   phosphonopentanoate (D-AP5) impairs spatial learning and LTP in vivo at intracerebral
826   concentrations comparable to those that block LTP in vitro. J Neurosci 12:21-34.

827   Dolnik A, Kanwal N, Mackert S, Halbedl S, Proepper C, Bockmann J, Schoen M, Boeckers TM,
828   Kuhl SJ, Schmeisser MJ (2016) Sipa1l3/SPAR3 is targeted to postsynaptic specializations
829   and interacts with the Fezzin ProSAPiP1/Lzts3. J Neurochem 136:28-35.

830   Ehinger Y, Morisot N, Phamluong K, Sakhai SA, Soneja D, Adrover MF, Alvarez VA, Ron D (2021)
831   cAMP-Fyn signaling in the dorsomedial striatum direct pathway drives excessive alcohol
832   use. Neuropsychopharmacology 46:334-342.

833   Eichenbaum H (1997) Declarative memory: insights from cognitive neurobiology. Annu Rev
834   Psychol 48:547-572.

835   Elias GM, Funke L, Stein V, Grant SG, Bredt DS, Nicoll RA (2006) Synapse-specific and
836   developmentally regulated targeting of AMPA receptors by a family of MAGUK scaffolding
837   proteins. Neuron 52:307-320.

838   Frey U, Morris RG (1998) Synaptic tagging: implications for late maintenance of hippocampal
839   long-term potentiation. Trends Neurosci 21:181-188.

Gangal H, Xie X, Huang Z, Cheng Y, Wang X, Lu J, Zhuang X, Essoh A, Huang Y, Chen R, Smith
LN, Smith RJ, Wang J (2023) Drug reinforcement impairs cognitive flexibility by inhibiting
striatal cholinergic neurons. Nat Commun 14:3886.

Graber TE, McCamphill PK, Sossin WS (2013) A recollection of mTOR signaling in learning and
memory. Learn Mem 20:518-530.

Herring BE, Nicoll RA (2016) Long-Term Potentiation: From CaMKII to AMPA Receptor
Trafficking. Annu Rev Physiol 78:351-365.

Hoeffer CA, Klann E (2010) mTOR signaling: at the crossroads of plasticity, memory and disease.
Trends Neurosci 33:67-75.

Horii T, Morita S, Kimura M, Terawaki N, Shibutani M, Hatada I (2017) Efficient generation of
conditional knockout mice via sequential introduction of lox sites. Sci Rep 7:7891.

Huganir RL, Nicoll RA (2013) AMPARs and synaptic plasticity: the last 25 years. Neuron 80:704-
717.

Jeong N, Singer AC (2022) Learning from inhibition: Functional roles of hippocampal CA1
inhibition in spatial learning and memory. Curr Opin Neurobiol 76:102604.

Kaizuka T, Takumi T (2018) Postsynaptic density proteins and their involvement in
neurodevelopmental disorders. J Biochem 163:447-455.

Kennedy MB (2018) The Protein Biochemistry of the Postsynaptic Density in Glutamatergic
Synapses Mediates Learning in Neural Networks. Biochemistry 57:4005-4009.

Kuriu T, Inoue A, Bito H, Sobue K, Okabe S (2006) Differential control of postsynaptic density
scaffolds via actin-dependent and -independent mechanisms. J Neurosci 26:7693-7706.

Labuhn M, Adams FF, Ng M, Knoess S, Schambach A, Charpentier EM, Schwarzer A, Mateo JL,
Klusmann JH, Heckl D (2018) Refined sgRNA efficacy prediction improves large- and
small-scale CRISPR-Cas9 applications. Nucleic Acids Res 46:1375-1385.

Laguesse S, Morisot N, Shin JH, Liu F, Adrover MF, Sakhai SA, Lopez MF, Phamluong K, Griffin
WC, 3rd, Becker HC, Bender KJ, Alvarez VA, Ron D (2017) Prosapip1-Dependent

866 Synaptic Adaptations in the Nucleus Accumbens Drive Alcohol Intake, Seeking, and

867 Reward. Neuron 96:145-159 e148.

868 Lee I, Kesner RP (2004) Differential contributions of dorsal hippocampal subregions to memory

869 acquisition and retrieval in contextual fear-conditioning. Hippocampus 14:301-310.

870 Leger M, Quiedeville A, Bouet V, Haelewyn B, Boulouard M, Schumann-Bard P, Freret T (2013)

871 Object recognition test in mice. Nat Protoc 8:2531-2537.

872 Li F, Tsien JZ (2009) Memory and the NMDA receptors. N Engl J Med 361:302-303.

873 Lin Y, Skeberdis VA, Francesconi A, Bennett MV, Zukin RS (2004) Postsynaptic density protein-

874 95 regulates NMDA channel gating and surface expression. J Neurosci 24:10138-10148.

875 Liu F, Laguesse S, Legastelois R, Morisot N, Ben Hamida S, Ron D (2017) mTORC1-dependent

876 translation of collapsin response mediator protein-2 drives neuroadaptations underlying

877 excessive alcohol-drinking behaviors. Mol Psychiatry 22:89-101.

878 Lueptow LM (2017) Novel Object Recognition Test for the Investigation of Learning and Memory

879 in Mice. J Vis Exp:55718.

880 Ma T, Cheng Y, Roltsch Hellard E, Wang X, Lu J, Gao X, Huang CCY, Wei XY, Ji JY, Wang J

881 (2018) Bidirectional and long-lasting control of alcohol-seeking behavior by corticostriatal

882 LTP and LTD. Nat Neurosci 21:373-383.

883 Mackes JL, Willner J (2006) NMDA antagonist MK-801 impairs acquisition of place strategies, but

884 not their use. Behav Brain Res 175:112-118.

885 Matsuura K, Kobayashi S, Konno K, Yamasaki M, Horiuchi T, Senda T, Hayashi T, Satoh K,

886 Arima-Yoshida F, Iwasaki K, Negishi L, Yasui-Shimizu N, Kohu K, Kawahara S, Kirino Y,

887 Nakamura T, Watanabe M, Yamamoto T, Manabe T, Akiyama T (2022) SIPA1L1/SPAR1

888 Interacts with the Neurabin Family of Proteins and is Involved in GPCR Signaling. J

889 Neurosci 42:2448-2473.

Mayanagi T, Yasuda H, Sobue K (2015) PSD-Zip70 Deficiency Causes Prefrontal Hypofunction

Associated with Glutamatergic Synapse Maturation Defects by Dysregulation of Rap2

Activity. J Neurosci 35:14327-14340.

Meira T, Leroy F, Buss EW, Oliva A, Park J, Siegelbaum SA (2018) A hippocampal circuit linking

dorsal CA2 to ventral CA1 critical for social memory dynamics. Nat Commun 9:4163.

Nachtigall EG, de CMJ, Izquierdo I, Furini CRG (2024) Cellular mechanisms of contextual fear

memory reconsolidation: Role of hippocampal SFKs, TrkB receptors and GluN2B-

containing NMDA receptors. Psychopharmacology (Berl) 241:61-73.

Naisbitt S, Kim E, Tu JC, Xiao B, Sala C, Valtschanoff J, Weinberg RJ, Worley PF, Sheng M

(1999) Shank, a novel family of postsynaptic density proteins that binds to the NMDA

receptor/PSD-95/GKAP complex and cortactin. Neuron 23:569-582.

Newcomer JW, Farber NB, Olney JW (2000) NMDA receptor function, memory, and brain aging.

Dialogues Clin Neurosci 2:219-232.

Pak DT, Yang S, Rudolph-Correia S, Kim E, Sheng M (2001) Regulation of dendritic spine

morphology by SPAR, a PSD-95-associated RapGAP. Neuron 31:289-303.

Pilly PK, Grossberg S (2012) How do spatial learning and memory occur in the brain? Coordinated

learning of entorhinal grid cells and hippocampal place cells. J Cogn Neurosci 24:1031-

1054.

Pitts MW (2018) Barnes Maze Procedure for Spatial Learning and Memory in Mice. Bio Protoc 8.

Qualmann B, Boeckers TM, Jeromin M, Gundelfinger ED, Kessels MM (2004) Linkage of the actin

cytoskeleton to the postsynaptic density via direct interactions of Abp1 with the

ProSAP/Shank family. J Neurosci 24:2481-2495.

Reim D, Weis TM, Halbedl S, Delling JP, Grabrucker AM, Boeckers TM, Schmeisser MJ (2016)

The Shank3 Interaction Partner ProSAPiP1 Regulates Postsynaptic SPAR Levels and the

Maturation of Dendritic Spines in Hippocampal Neurons. Front Synaptic Neurosci 8:13.

Renaud JB, Boix C, Charpentier M, De Cian A, Cochennec J, Duvernois-Berthet E, Perrouault L,
Tesson L, Edouard J, Thinard R, Cherifi Y, Menoret S, Fontaniere S, de Croze N,
Fraichard A, Sohm F, Anegon I, Concordet JP, Giovannangeli C (2016) Improved Genome
Editing Efficiency and Flexibility Using Modified Oligonucleotides with TALEN and
CRISPR-Cas9 Nucleases. Cell Rep 14:2263-2272.

Richter M, Murai KK, Bourgin C, Pak DT, Pasquale EB (2007) The EphA4 receptor regulates
neuronal morphology through SPAR-mediated inactivation of Rap GTPases. J Neurosci
27:14205-14215.

Rosenfeld CS, Ferguson SA (2014) Barnes maze testing strategies with small and large rodent
models. J Vis Exp:e51194.

Sanchez-Rodriguez I, Temprano-Carazo S, Jeremic D, Delgado-Garcia JM, Gruart A, Navarro-
Lopez JD, Jimenez-Diaz L (2022) Recognition Memory Induces Natural LTP-like
Hippocampal Synaptic Excitation and Inhibition. Int J Mol Sci 23:10806.

Sanderson DJ, Good MA, Skelton K, Sprengel R, Seeburg PH, Rawlins JN, Bannerman DM
(2009) Enhanced long-term and impaired short-term spatial memory in GluA1 AMPA
receptor subunit knockout mice: evidence for a dual-process memory model. Learn Mem
16:379-386.

Sharma S, Rakoczy S, Brown-Borg H (2010) Assessment of spatial memory in mice. Life Sci
87:521-536.

Sheng M, Kim E (2000) The Shank family of scaffold proteins. J Cell Sci 113 (Pt 11):1851-1856.

Squire LR, Stark CE, Clark RE (2004) The medial temporal lobe. Annu Rev Neurosci 27:279-306.

Stemmer M, Thumberger T, Del Sol Keyer M, Wittbrodt J, Mateo JL (2015) CCTop: An Intuitive,
Flexible and Reliable CRISPR/Cas9 Target Prediction Tool. PLoS One 10:e0124633.

Stevenson RF, Zheng J, Mnatsakanyan L, Vadera S, Knight RT, Lin JJ, Yassa MA (2018)
Hippocampal CA1 gamma power predicts the precision of spatial memory judgments.
Proc Natl Acad Sci U S A 115:10148-10153.

Straube B (2012) An overview of the neuro-cognitive processes involved in the encoding,
consolidation, and retrieval of true and false memories. Behav Brain Funct 8:35.

Tsien JZ, Huerta PT, Tonegawa S (1996) The essential role of hippocampal CA1 NMDA receptor-
dependent synaptic plasticity in spatial memory. Cell 87:1327-1338.

Tu JC, Xiao B, Naisbitt S, Yuan JP, Petralia RS, Brakeman P, Doan A, Aakalu VK, Lanahan AA,
Sheng M, Worley PF (1999) Coupling of mGluR/Homer and PSD-95 complexes by the
Shank family of postsynaptic density proteins. Neuron 23:583-592.

Tzakis N, Holahan MR (2019) Social Memory and the Role of the Hippocampal CA2 Region.
Front Behav Neurosci 13:233.

Walf AA, Frye CA (2007) The use of the elevated plus maze as an assay of anxiety-related
behavior in rodents. Nat Protoc 2:322-328.

Wang J, Ben Hamida S, Darcq E, Zhu W, Gibb SL, Lanfranco MF, Carnicella S, Ron D (2012)
Ethanol-mediated facilitation of AMPA receptor function in the dorsomedial striatum:
implications for alcohol drinking behavior. J Neurosci 32:15124-15132.

Wang X, Zhan Y (2022) Regulation of Social Recognition Memory in the Hippocampal Circuits.
Front Neural Circuits 16:839931.

Wei X, Wang J, Yang E, Zhang Y, Qian Q, Li X, Huang F, Sun B (2024) Efr3b is essential for
social recognition by modulating the excitability of CA2 pyramidal neurons. Proc Natl Acad
Sci U S A 121:e2314557121.

Wendholt D, Spilker C, Schmitt A, Dolnik A, Smalla KH, Proepper C, Bockmann J, Sobue K,
Gundelfinger ED, Kreutz MR, Boeckers TM (2006) ProSAP-interacting protein 1
(ProSAPiP1), a novel protein of the postsynaptic density that links the spine-associated
Rap-Gap (SPAR) to the scaffolding protein ProSAP2/Shank3. J Biol Chem 281:13805-
13816.

Won S, Incontro S, Nicoll RA, Roche KW (2016) PSD-95 stabilizes NMDA receptors by inducing
the degradation of STEP61. Proc Natl Acad Sci U S A 113:E4736-4744.

Yang X, Gong R, Qin L, Bao Y, Fu Y, Gao S, Yang H, Ni J, Yuan TF, Lu W (2022) Trafficking of

NMDA receptors is essential for hippocampal synaptic plasticity and memory

consolidation. Cell Rep 40:111217.

Zhang H, Etherington LA, Hafner AS, Belelli D, Coussen F, Delagrange P, Chaouloff F, Spedding

M, Lambert JJ, Choquet D, Groc L (2013) Regulation of AMPA receptor surface trafficking

and synaptic plasticity by a cognitive enhancer and antidepressant molecule. Mol

Psychiatry 18:471-484.

Zucker RS, Regehr WG (2002) Short-term synaptic plasticity. Annu Rev Physiol 64:355-405.

Figure 1. Generation and characterization of Prosapip1(flx/flx) mice

(A) Identification of guide RNA binding sites in intron 2 and the 3' UTR of exon 5 of Prosapip1.

(B) PAGE purified Ultramer single stranded DNA oligos that were homologous to the target loci

in intron 2 and exon 5 were used as repair templates. **(C)** Genetic crossing scheme of the

Prosapip1(flx/flx);Syn1-Cre mice. Male Prosapip1(flx/flx);Syn1-Cre(-/-) mice were mated with

female Prosapip1(flx/flx);Syn1-Cre(+/-) mice, leading to litters of Prosapip1(flx/flx);Syn1-Cre(-/-)

or Prosapip1(flx/flx);Syn1-Cre(+/-) mice. **(D)** The dHP of C57BL/6, Prosapip1(flx/flx);Syn1-Cre(-)

and Prosapip1(flx/flx);Syn1-Cre(+) mice was dissected. Prosapip1 was detected using anti-

Prosapip1 antibodies. GAPDH was used as a loading control. **(E)** Histogram of litter size from

Prosapip1 mating pairs. X axis depicts number of pups per litter, while Y axis depicts numbers of

litters at that size. **(F)** Proportion of male and female offspring from Prosapip1 matings. **(G)**

Proportion of Cre(-) and Cre(+) offspring from Prosapip1 matings. **(H)** Body weight of male and

female Prosapip1(flx/flx);Syn1-Cre(-) and Prosapip1(flx/flx);Syn1-Cre(+) mice was measured

biweekly from birth to assess overt developmental deficits. Data are represented as mean±SEM

and analyzed using three-way ANOVA (**Table 1**). n = 10 (Prosapip1(flx/flx);Syn1-Cre(-) male), 5

(Prosapip1(flx/flx);Syn1-Cre(-) female), 11 (Prosapip1(flx/flx);Syn1-Cre(+) male), 11

(Prosapip1(flx/flx);Syn1-Cre(+) female). **(I-J)** Mice were placed in an open field and locomotion

was recorded for 20 minutes. Total distance traveled during the open field test **(I)** and time spent

in the center of the open field **(J)**. Data represented as mean±SEM and analyzed using an

unpaired t-test (**Table 1**). ns, non-significant. n = 22 (Prosapip1(flx/flx);Syn1-Cre(-)), 21

(Prosapip1(flx/flx);Syn1-Cre(+)).

1000

1001

Figure 2. Prosapip1 is required for synaptic localization of PSD proteins

(A) The levels of Prosapip1 (top), Synapsin (middle), and CREB (bottom) in the dorsal hippocampus of Prosapip1(flx/flx);Syn1-Cre(-) mice were measured in the total and crude synaptosomal fraction. **(B-I)** Total levels of SPAR **(B),** Shank3 **(D, F)**, and PSD-95 **(D, H)** alongside the synaptic levels of SPAR **(C)**, Shank3 **(E, G)**, and PSD-95 **(E, I)** were measured in the dHP of Prosapip1(flx/flx);Syn1-Cre(-) and Prosapip1(flx/flx);Syn1-Cre(+) mice using western blot analysis. Protein levels were normalized to GAPDH and presented as a percentage of the average of the Prosapip1(flx/flx);Syn1-Cre(-) values. Data are represented as mean±SEM and analyzed using unpaired two-tailed t-test with Welch's correction (**Table 1**). **$p<0.01$, ****$p<0.0001$; ns, non-significant. n = 5 per group (B-G), 9 (Prosapip1(flx/flx);Syn1-Cre(-)) and 10 (Prosapip1(flx/flx);Syn1-Cre(+)) (H-I). **(J-Q)** The total levels of GluN2A **(J, L)**, GluN2B **(J, N)**, and GluA1 **(J, P)** and the synaptic levels of GluN2A **(K, M)**, GluN2B **(K, O)**, and GluA1 **(K, Q)** were measured dHP of Prosapip1(flx/flx);Syn1-Cre(-) and Prosapip1(flx/flx);Syn1-Cre(+) mice via western blot analysis. Protein levels were normalized to GAPDH and presented as a percentage of the average of the Prosapip1(flx/flx);Syn1-Cre(-) values. Data are represented as mean±SEM and analyzed using unpaired two-tailed t-test with Welch's correction (**Table 1**). *$p<0.05$, **$p<0.01$, ***$p<0.001$; ns, non-significant. n = 4 (Prosapip1(flx/flx);Syn1-Cre(-)), 5 (Prosapip1(flx/flx);Syn1-Cre(+)).

Figure 3. Prosapip1 in the dorsal hippocampus plays a role in NMDA receptor-mediated transmission and long-term potentiation

(A) Location of stimulating and recording electrodes within the hippocampal CA1 region. **(B)** Sample field excitatory postsynaptic potential (fEPSP) traces recorded before (pre) and after (post) administering high-frequency stimulation (HFS) (100Hz, 100 pulses every 20 seconds) in both Prosapip1(flx/flx);Syn1-Cre(-) (Cre(-)) and Prosapip1(flx/flx);Syn1-Cre(+) (Cre(+)) groups. **(C)** A stable baseline of fEPSPs was established for 10 minutes before application of HFS and fEPSPs were recorded for 30 minutes after HFS. Time course of fEPSPs before and after HFS. **(D)** Quantification of average of fEPSP amplitudes measured between 30-40 minutes. Data are represented as mean±SEM and analyzed using unpaired two-tailed t-test (**Table 1**). **p<0.01. n = 10 slices from 6 mice (10/6) (Prosapip1(flx/flx);Syn1-Cre(-)) and (9/5) (Prosapip1(flx/flx);Syn1-Cre(+)). **(E)** Cells were clamped at -65 mV and the bath contained both DNQX and PTX to block AMPA and GABA mediated responses. Voltage clamp whole cell recordings and representative electrically evoked NMDA currents in Prosapip1(flx/flx);Syn1-Cre(-) (Cre(-)) and Prosapip1(flx/flx);Syn1-Cre(+) (Cre(+)) mice at four stimulation intensities (left). Summarized responses for Prosapip1(flx/flx);Syn1-Cre(-) (Cre(-)) and Prosapip1(flx/flx);Syn1-Cre(+) (Cre(+)) CA1 neurons quantified by average at each stimulating intensity; #p < 0.05 by two-way repeated measures ANOVA followed by post-hoc Tukey test (Prosapip1(flx/flx);Syn1-Cre(-) vs. Prosapip1(flx/flx);Syn1-Cre(+)) at the same stimulating intensities, *p<0.05. n = 14/4 (Prosapip1(flx/flx);Syn1-Cre(-)) and 11/4 (Prosapip1(flx/flx);Syn1-Cre(+)). **(F)** Representative currents evoked in the CA1 neurons after NMDA bath application (20 uM, 30s) in Prosapip1(flx/flx);Syn1-Cre(-) (Cre(-)) and Prosapip1(flx/flx);Syn1-Cre(+) (Cre(+)) mice (left). Average of the peak current elicited by each mouse (right). Data are represented as mean±SEM and analyzed using unpaired two-tailed t-test (**Table 1**). **p<0.01. n = 13/4 (Prosapip1(flx/flx);Syn1-Cre(-)) and 11/4 (Prosapip1(flx/flx);Syn1-Cre(+)). **(G)** In voltage clamp

recordings, 2 electrical stimulations (100 ms separation) were provided to elicit 2 responses in

the CA1 neurons. PPR was calculated as the amplitude of peak 2/amplitude of peak 1.

Representative paired-pulse ratio in Prosapip1(flx/flx);Syn1-Cre(-) (Cre(-)) and

Prosapip1(flx/flx);Syn1-Cre(+) (Cre(+)) mice (left). Average PPR for Prosapip1(flx/flx);

Syn1-Cre(-) (Cre(-)) and Prosapip1(flx/flx);Syn1-Cre(+) (Cre(+)) (right). Data are represented as

mean±SEM and analyzed using unpaired two-tailed t-test (**Table 1**). *$p < 0.05$. n = 11/3

(Prosapip1(flx/flx);Syn1-Cre(-)) and 13/4 (Prosapip1(flx/flx);Syn1-Cre(+)).

1052

Figure 4. Prosapip1 contributes to recognition, social, and spatial memory

(A) Mice underwent the novel object recognition test, where they were first allowed to explore two similar objects. After 24 hours, one of the familiar objects was replaced by a novel object, and mice were again allowed to explore and interact with the objects. Time spent interacting with the familiar and novel object. n = 22 (Prosapip1(flx/flx);Syn1-Cre(-)), 20 (Prosapip1(flx/flx);Syn1-Cre(+)). **(B-C)** In the novelty T-maze test, mice were allowed to explore two arms of a three-armed, T-shaped maze. There were 5 training trials separated by a 1-minute inter-trial interval. During testing, the third "novel" arm was unblocked and allowed to be explored. **(B)** Difference score (time exploring novel arm – time exploring familiar arm) performance during the novelty T-maze test. A positive difference score indicates preference for the novel arm of the maze. n = 22 (Prosapip1(flx/flx);Syn1-Cre(-)), 19 (Prosapip1(flx/flx);Syn1-Cre(+)). **(C)** Heat map of group average time spent in each arm of the T-maze during the test trial. **(D-E)** Mice performed the 3-chamber social interaction test. Specifically, they were placed in the center chamber of a 3-chamber apparatus and allowed to freely explore for 15 minutes for two trials. During the first trial, one chamber was paired with a juvenile interaction partner (social), while the other chamber contained only the empty interaction cage (empty). During the second trial, one chamber was paired with the familiar mouse from the first trial (familiar), and the other chamber contained a novel juvenile interaction partner (novel). **(D)** Time spent in the empty and social-paired chamber respectively in the first portion of the 3-chamber social interaction test. n = 14 (Prosapip1(flx/flx);Syn1-Cre(-)), 17 (Prosapip1(flx/flx);Syn1-Cre(+)). **(E)** Time spent in the familiar and novel chamber during the second portion of the 3-chamber social interaction test. n = 14 (Prosapip1(flx/flx);Syn1-Cre(-)), 17 (Prosapip1(flx/flx);Syn1-Cre(+)). **(F-K)** Mice performed the Barnes maze test, where they were placed in the center of a white plastic platform with 40 uniformly distributed holes around the perimeter, one of which had an exit compartment placed underneath. The goal of the trial was to escape into the exit compartment. There were 4 training

trials a day over the course of 4 days, separated by an inter-trial interval of 30 minutes. 24 hours

after the last training trial mice were placed back onto the platform but with the exit compartment

removed (probe trial) and allowed to explore for 5 minutes. **(F)** Average distance traveled from

start point to exit during the Barnes maze training trials. n = 9 (Prosapip1(flx/flx);Syn1-Cre(-)), 12

(Prosapip1(flx/flx);Syn1-Cre(+)). **(G)** Primary (filled circles) and secondary (hollow circles) errors

committed during Barnes maze training. Primary errors are an incorrect hole visit and secondary

errors are an incorrect hole revisit. n = 9 (Prosapip1(flx/flx);Syn1-Cre(-)), 12

(Prosapip1(flx/flx);Syn1-Cre(+)). **(H)** The method of searching utilized by each mouse for each

training trial was qualified. Example path to exit from mice exhibiting random, serial, and spatial

search strategies. **(I-J)** Ratio of search strategy utilization by Prosapip1(flx/flx);Syn1-Cre(-) **(I)**

and Prosapip1(flx/flx);Syn1-Cre(+) **(J)** mice during Barnes maze training. **(K-L)** Time spent in

exit-associated quartile during the probe trial and associated heatmaps **(L)**. n = 9

(Prosapip1(flx/flx);Syn1-Cre(-)), 12 (Prosapip1(flx/flx);Syn1-Cre(+)). Data are represented as

mean±SEM and analyzed using two-way ANOVA (A,C,D,E), Welch's t-test (B), three-way ANOVA

(F), or Mann-Whitney test (J) (**Table 1**). *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

1093

Figure 5. Prosapip1 in the dorsal hippocampus contributes to recognition, social, and spatial memory

(A) Images of AAV-GFP and AAV-Cre overexpression in the dHP of Prosapip1(flx/flx) mice. **(B)** Western blot analysis of Prosapip1 protein level in the dHP in non-infected mice compared to mice infected with AAV-Cre. **(C-D)** Mice infected with AAV-GFP or AAV-Cre in the dHP were placed in an open field, and their behavior was recorded for 20 minutes. The total distance traveled **(C)** and the time spent in the center of the field **(D)** were measured during the test. n = 19 (AAV-GFP), 15 (AAV-Cre). **(E)** Mice infected with AAV-GFP or AAV-Cre in the dHP were placed on the light side of a light/dark box apparatus and allowed to explore for 10 minutes. Time spent in the dark and light chamber during the light/dark box test. n = 19 (AAV-GFP), 15 (AAV-Cre). **(F)** Mice infected with AAV-GFP or AAV-Cre in the dHP underwent the novel object recognition test. Briefly, they were familiarized with two similar objects before one was switched for a novel object after 24 hours. Cumulative time spent exploring the familiar and novel object during the novel object recognition test. n = 16 (AAV-GFP), 17 (AAV-Cre). **(G)** Mice infected with AAV-GFP or AAV-Cre in the dHP underwent the novelty T-maze test, where they were first allowed to explore two arms of a three-armed, T-shaped maze. During the testing phase, the third "novel" arm was unblocked and made available for exploration. Difference score (time exploring novel arm – time exploring familiar arm) of time spent exploring the novel arm of the T-maze. A heatmap of each average group performance during the test is also presented. n = 18 (AAV-GFP), 15 (AAV-Cre). **(H-I)** Mice infected with AAV-GFP or AAV-Cre in the dHP performed the 3-chamber social interaction test. Briefly, they were placed in the center chamber and allowed to freely explore for 15 minutes during two trials. In the first trial, one chamber had a juvenile interaction partner, and the other was empty. In the second trial, one chamber had the familiar mouse from the first trial, and the other had a new juvenile interaction partner. **(H)** Cumulative time spent in the empty and social chamber of the 3-chamber social interaction test. n = 18 (AAV-

GFP), 15 (AAV-Cre). **(I)** Cumulative time spent in the familiar and novel chambers in the 3-

chamber social interaction test. n = 18 (AAV-GFP), 15 (AAV-Cre). **(J-K)** Mice infected with AAV-

GFP or AAV-Cre in the dHP underwent the Barnes maze experiment. They were placed in the

center of a platform with 40 evenly spaced holes around the perimeter, one of which led to an exit

compartment. Mice underwent 4 training trials per day for 4 days, with 30-minute intervals

between trials. Twenty-four hours after the last training trial, they were placed back on the platform

without the exit compartment (probe trial) and allowed to explore for 5 minutes. **(J)** Average

distance traveled from start point to exit during the Barnes maze training trials. n = 8 (AAV-GFP),

6 (AAV-Cre). **(K)** Time spent in the exit quartile during the probe trial. n = 8 (AAV-GFP), 6 (AAV-

Cre). Data are represented as mean±SEM and analyzed using two-way ANOVA (E, F, H, I, J),

Welch's t-test (C, D, G), or Mann-Whitney test (K) (**Table 1**). *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$,

****$p < 0.0001$.

1131

Extended Figure 4. Prosapip1 knockout did not affect Barnes maze exploratory behavior or baseline anxiety

(A) Before the Barnes maze training trials began, mice were placed on the platform to assess baseline exploratory behavior and locomotion in the environment. The exit compartment was in a different location than training trials. Distance traveled during the Barnes maze habituation trial. Data represented as mean±SEM and analyzed using unpaired t-test (**Table 1**). ns, non-significant. n = 6 (Prosapip1(flx/flx);Syn1-Cre(-)), 12 Prosapip1(flx/flx);Syn1-Cre(+). **(B)** To assess anxiety, mice were placed on the light side of a light/dark box apparatus and allowed to explore for 10 minutes. Time spent in the light and dark chamber during the test. Data represented as mean±SEM and analyzed using a two-way ANOVA (**Table 1**). ****$p<0.0001$. n = 15 (Prosapip1(flx/flx);Syn1-Cre(-)), 16 (Prosapip1(flx/flx);Syn1-Cre(+)). **(C)** Mice were placed in the center of a plus-shaped maze and allowed to explore two closed and two open arms for 5 minutes to assess anxiety and exploratory behavior. Time spent in the open arm of the elevated plus maze. Open-arm time was measured and analyzed using unpaired two-tailed t-test (**Table 1**). ns, non-significant. n = 7 (Prosapip1(flx/flx);Syn1-Cre(-)), 17 (Prosapip1(flx/flx);Syn1-Cre(+)).

Figure 1

Generation and characterization of Prosapip1(flx/flx) mice

A

B

Intron 2

WT   AAGCACTGCTTTTCTCAACCAGTCCAATCTCCCCGGCCCATGAAACTGAA

Flox   AAGCACTGCTTTTCTCAACCAGTCCAATCTCCCCGGCCCATGAAACTGAA

GAATTC ATAACTTCGTATAGCATACATTATACGAAGTTAT

EcoRI   loxP

Exon 5 3'UTR

WT   GCGCATGGAGCAAGGCTCCCTTTCTGGGGAAGCCTCGTGTTGGAGATAG

Flox   GCGCATGGAGCAAGGCTCCCTTTCTGGGGAAGCCTCGTGTTGGAGATAG

GATATC ATAACTTCGTATAGCATACATTATACGAAGTTAT

EcoRV   loxP

C

Prosapip1(flx/flx);Syn1-Cre(–/–)   Prosapip1(flx/flx);Syn1-Cre(+/–)

♂   X   ♀

Prosapip1(flx/flx);Syn1-Cre(–/–) or Prosapip1(flx/flx);Syn1-Cre(+/–)

D

E Litter Size — Mean: 5.52

F Sex — 48% Female, 52% Male, Total = 358

G Genotype — 53% / 47%, Total = 319

H Body Weight

I Total Distance — ns

J Total Spent in Center — ns

Figure 2

Prosapip1 is required for synaptic localization of PSD proteins

Figure 3

**Prosapip1 in the dorsal hippocampus plays a role in NMDA
receptor-mediated transmission and long-term potentiation**

Figure 4

Prosapip1 contributes to recognition, social, and spatial memory

Figure 5

Prosapip1 in the dorsal hippocampus contributes to recognition, social, and spatial memory

**Prosapip1 knockout did not affect Barnes maze exploratory
behavior or baseline anxiety**

Table 1. Statistics

Figure	n	Sex Difference	Statistical Test	Effect	Result	p-value	Post-hoc	Post-hoc comparison	p-value
1E	77		D'Agostino & Pearson	Normality	K2 = 5.480	0.0646			
1H	10 (Cre- Male), 5 (Cre- Female), 11 (Cre+ Male), 11 (Cre+ Female)	Yes	Three-way ANOVA	Genotype	$F_{(1,33)}$ = 0.7842	0.3823			
				Sex	$F_{(1,33)}$ = 23.94	< 0.0001			
				Time	$F_{(5,158)}$ = 588.4	< 0.0001			
				Genotype x Sex	$F_{(1,33)}$ = 1.345	0.2545			
				Genotype x Time	$F_{(5,158)}$ = 0.1476	0.9805			
				Sex x Time	$F_{(5,158)}$ = 2.345	0.0437			
				Genotype x Sex x Time	$F_{(5,158)}$ = 0.3764	0.8643			
1I	22 (Cre-), 21 (Cre+)	No	Unpaired t-test	Genotype	$t_{(41)}$ = 1.460	0.1518			
1J	22 (Cre-), 21 (Cre+)	No	Mann-Whitney Test	Genotype	U = 220	0.8007			
2B	5 (Cre-), 5 (Cre+)		Unpaired t-test	Genotype	$t_{(6.963)}$ = 1.427	0.1968			
2C	5 (Cre-), 5 (Cre+)		Unpaired t-test	Genotype	$t_{(5.880)}$ = 5.183	0.0022			
2F	5 (Cre-), 5 (Cre+)		Unpaired t-test	Genotype	$t_{(6.932)}$ = 1.345	0.221			
2G	5 (Cre-), 5 (Cre+)		Unpaired t-test	Genotype	$t_{(6.720)}$ = 0.1306	0.8999			
2H	9 (Cre-), 10 (Cre+)		Unpaired t-test	Genotype	$t_{(16.38)}$ = 0.5588	0.5839			
2I	9 (Cre-), 10 (Cre+)		Unpaired t-test	Genotype	$t_{(16.44)}$ = 6.095	< 0.0001			
2L	4 (Cre-), 5 (Cre+)		Unpaired t-test	Genotype	$t_{(4.429)}$ = 0.1050	0.921			
2M	4 (Cre-), 5 (Cre+)		Unpaired t-test	Genotype	$t_{(6.689)}$ = 0.1758	0.8656			
2N	4 (Cre-), 5 (Cre+)		Unpaired t-test	Genotype	$t_{(6.700)}$ = 1.675	0.1398			
2O	4 (Cre-), 5 (Cre+)		Unpaired t-test	Genotype	$t_{(6.332)}$ = 5.524	0.0012			
2P	4 (Cre-), 5 (Cre+)		Unpaired t-test	Genotype	$t_{(6.956)}$ = 0.1831	0.8599			
2Q	4 (Cre-), 5 (Cre+)		Unpaired t-test	Genotype	$t_{(6.060)}$ = 0.8548	0.4252			
3D	10/6 (Cre-), 9/5 (Cre+)		Unpaired t-test	Genotype	$t_{(17)}$ = 3.933	0.0011			
3E	14/4 (Cre-), 11/4 (Cre+)		Two-way ANOVA	Genotype	$F_{(1,3)}$ = 9.441	0.044	Tukey	0.2 (Cre- vs. Cre+)	0.345
				Intensity	$F_{(1,3)}$ = 9.441	< 0.001		0.4 (Cre- vs. Cre+)	0.025
				Genotype x Intensity	$F_{(1,3)}$ = 1.376	0.258		0.6 (Cre- vs. Cre+)	0.034
								0.9 (Cre- vs. Cre+)	0.046
3F	3/4 (Cre-), 11/4 (Cre+)		Unpaired t-test	Genotype	$t_{(22)}$ = 3.094	0.00529			
3G	11/3 (Cre-), 13/4 (Cre+)		Unpaired t-test	Genotype	$t_{(22)}$ = 2.099	0.0475			
4A	22 (Cre-), 20 (Cre+)	No	Two-way ANOVA	Genotype	$F_{(1,40)}$ = 0.3079	= 0.5821	Šidák	Familiar vs. Novel (Cre-)	<0.0001
				Object	$F_{(1,40)}$ = 34.32	< 0.0001		Familiar vs. Novel (Cre+)	0.4871
				Genotype x Object	$F_{(1,40)}$ = 18.37	0.0001			
4B	22 (Cre-), 19 (Cre+)	No	Unpaired t-test	Genotype	$t_{(26.08)}$ = 5.434	< 0.0001			
4D	14 (Cre-), 17 (Cre+)	No	Two-way ANOVA	Genotype	$F_{(1,29)}$ = 5.419	0.0271	Šidák	Empty vs. Social (Cre-)	0.0006
				Social	$F_{(1,29)}$ = 34.30	< 0.0001		Empty vs. Social (Cre+)	0.0005
				Genotype x Social	$F_{(1,29)}$ = 0.05655	0.8137			
4E	14 (Cre-), 17 (Cre+)	No	Two-way ANOVA	Genotype	$F_{(1,29)}$ = 1.463	0.2362	Šidák	Familiar vs. Novel (Cre-)	0.0008
				Social Novelty	$F_{(1,29)}$ = 17.60	0.0002		Familiar vs. Novel (Cre+)	0.1451
				Genotype x Social Novelty	$F_{(1,29)}$ = 2.947	0.0967			
Ext. 4A	6 (Cre-), 12 (Cre+)	No	Unpaired t-test	Genotype	$t_{(16)}$ = 0.2635	0.7955			
4F	9 (Cre-), 12 (Cre+)	No	Two-way ANOVA	Genotype	$F_{(1,19)}$ = 102.0	< 0.0001	Šidák	Trial 2 (Cre- vs. Cre+)	<0.0001
				Trial	$F_{(15,285)}$ = 7.899	< 0.0001		Trial 3 (Cre- vs. Cre+)	0.0067
				Genotype x Trial	$F_{(15,285)}$ = 1.401	0.1454		Trial 4 (Cre- vs. Cre+)	0.0018
								Trial 5 (Cre- vs. Cre+)	0.0002
								Trial 6 (Cre- vs. Cre+)	0.002
								Trial 7 (Cre- vs. Cre+)	0.0002
								Trial 8 (Cre- vs. Cre+)	0.0016
								Trial 9 (Cre- vs. Cre+)	<0.0001
								Trial 10 (Cre- vs. Cre+)	0.0006
								Trial 11 (Cre- vs. Cre+)	0.0049
								Trial 12 (Cre- vs. Cre+)	0.0169
								Trial 13 (Cre- vs. Cre+)	0.0098
4G	9 (Cre-), 12 (Cre+)	No	Three-way ANOVA	Genotype	$F_{(1,19)}$ = 100.7	< 0.0001			
				Trial	$F_{(15,289)}$ = 7.289	< 0.0001			
				Error Type	$F_{(0.4468, 8.489)}$ = 4.55	0.0772			
				Genotype x Trial	$F_{(15,289)}$ =1.529	0.094			
				Genotype x Error Type	$F_{(1,19)}$ = 15.09	0.001			
				Trial x Error Type	$F_{(6.157, 117)}$ = 2.865	0.0115			
				Genotype x Trial x Error Type	$F_{(15,285)}$ = 1.831	0.0303			
4K	9 (Cre-), 12 (Cre+)	No	Mann-Whitney Test	Genotype	U = 8	0.0004			
Ext. 4B	15 (Cre-), 16 (Cre+)	No	Two-way ANOVA	Chamber	$F_{(1,29)}$ = 92.69	< 0.0001	Šidák	Light vs. Dark (Cre-)	<0.0001
				Genotype	$F_{(1,29)}$ = 0.9504	0.3377		Light vs. Dark (Cre+)	<0.0001
				Genotype x Chamber	$F_{(1,29)}$ = 0.09645	0.7584			
Ext. 4C	7 (Cre-), 17 (Cre+)	No	Mann-Whitney Test	Genotype	U = 42	0.2878			
5C	19 (AAV-GFP), 15 (AAV-Cre)	No	Unpaired t-test	Treatment	$t_{(29.59)}$ = 1.961	0.0594			
5D	19 (AAV-GFP), 15 (AAV-Cre)	No	Unpaired t-test	Treatment	$t_{(29.72)}$ = 1.066	0.2951			
5E	19 (AAV-GFP), 15 (AAV-Cre)	No	Two-way ANOVA	Treatment	$F_{(1,32)}$ = 2.361	0.1342	Šidák	Light vs. Dark (AAV-GFP)	<0.0001
				Chamber	$F_{(1,32)}$ = 46.74	< 0.0001		Light vs. Dark (AAV-Cre)	0.0003
				Treatment x Chamber	$F_{(1,32)}$ = 0.1588	0.6929			
5F	16 (AAV-GFP), 17 (AAV-Cre)	No	Two-way ANOVA	Treatment	$F_{(1,31)}$ = 3.069	0.0897	Šidák	Familiar vs. Novel (AAV-GFP)	< 0.0001
				Object	$F_{(1,31)}$ = 36.30	< 0.0001		Familiar vs. Novel (AAV-Cre)	0.9965
				Treatment x Object	$F_{(1,31)}$ = 35.05	< 0.0001			
5G	18 (AAV-GFP), 15 (AAV-Cre)	No	Unpaired t-test	Treatment	$t_{(26.04)}$ = 3.777	0.0008			
5H	18 (AAV-GFP), 15 (AAV-Cre)	No	Two-way ANOVA	Treatment	$F_{(1,31)}$ = 0.0003957	0.9843	Šidák	Empty vs. Social (AAV-GFP)	<0.0001
				Social	$F_{(1,31)}$ = 31.60	< 0.0001		Empty vs. Social (AAV-Cre)	0.0062
				Treatment x Social	$F_{(1,31)}$ = 0.7781	0.3845			
5I	18 (AAV-GFP), 15 (AAV-Cre)	No	Two-way ANOVA	Treatment	$F_{(1,31)}$ = 0.2045	0.6543	Šidák	Familiar vs. Novel (AAV-GFP)	0.0303
				Social Novelty	$F_{(1,31)}$ = 9.777	0.0038		Familiar vs. Novel (AAV-Cre)	0.1319
				Treatment x Social Novelty	$F_{(1,31)}$ = 0.1131	0.7389			
5J	8 (AAV-GFP), 6 (AAV-Cre)	No	Two-way ANOVA	Treatment	$F_{(1,12)}$ = 22.95	0.0004	Šidák	Trial 5 (AAV-GFP vs. AAV-Cre)	<0.0001
				Trial	$F_{(15,180)}$ = 6.246	< 0.0001		Trial 9 (AAV-GFP vs. AAV-Cre)	0.0026
				Treatment x Trial	$F_{(15,180)}$ = 2.033	0.0153		Trial 10 (AAV-GFP vs. AAV-Cre)	<0.0001
								Trial 14 (AAV-GFP vs. AAV-Cre)	0.0475
5K	8 (AAV-GFP), 6 (AAV-Cre)	No	Mann-Whitney Test	Treatment	U = 3	0.0047			