HIV-1 infection promotes neuroinflammation and neuron pathogenesis in novel microglia-containing cerebral organoids

Srinivas D. Narasipura[#1], Janet P. Zayas[#1], Michelle K. Ash[1], Anjelica Reyes[1], Tanner Shull[1], Stephanie Gambut[1], Jeffrey R. Schneider[1], Ramon Lorenzo-Redondo[2,3], Lena Al-Harthi[1]* and João I. Mamede[1]*

1- Department of Microbial Pathogens and Immunity, Rush University Medical Center, Chicago, Illinois, USA.

2- Department of Medicine, Division of Infectious Diseases, Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA.

3- Center for Pathogen Genomics and Microbial Evolution, Institute for Global Health, Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA

[#]Equal contributors

*Co-corresponding authors:

joao_mamede@rush.edu

lena_al-harthi@rush.edu

Abstract

Cerebral organoids (COs) are a valuable tool to study the intricate interplay between glial cells and neurons in brain development and disease, including HIV‑associated neuroinflammation. We developed a novel approach to generate microglia containing COs (CO-iMs) by co-culturing hematopoietic progenitors and induced pluripotent stem cells. This approach allowed for the differentiation of microglia within the organoids concomitantly to the neuronal progenitors. CO-iMs exhibited higher efficiency in generation of CD45$^+$/CD11b$^+$/Iba-1$^+$ microglia cells compared to conventional COs with physiologically relevant proportion of microglia (~7%). CO-iMs exhibited substantially higher expression of microglial homeostatic and sensome markers as well as markers for the complement cascade. CO-iMs showed susceptibility to HIV infection resulting in a significant increase in several pro-inflammatory cytokines/chemokines and compromised neuronal function, which were abrogated by addition of antiretrovirals. Thus, CO-iM is a robust model to decipher neuropathogenesis, neurological disorders, and viral infections of brain cells in a 3D culture system.

Key words: Cerebral organoids, microglia, HIV, neuropathogenesis, neuroinflammation, iPSC, HPC.

Introduction:

Neurodegenerative diseases are debilitating disorders characterized by deterioration of brain function and affect millions of people worldwide. Currently, the most-utilized models to study brain diseases involve *postmortem* human brain tissue[1,2], non-human primates (NHP)[3], small animal models[4–7], *in vitro* 2-dimentional (2D) cultures of primary cells (mostly fetal in origin)[8] or cell lines[9-11]. While relevant in the context of their specific studies, many of these models are limited by availability, cost, and the differences between species[,5,9]. Cerebral organoids have emerged as a tool to study the human brain in health and disease [10,11]. COs provide a three-dimensional (3D) architecture of resident brain cells (neurons, astrocytes, oligodendrocytes, and microglia), albeit the lack vascularization of the tissue environment. They are developed from embryonic stem cells (ESCs) or induced pluripotent stem cells (iPSCs) to form 3D tissue organization and spatial arrangement of diverse CNS cell types that recapitulate the intricate pattern and functionality of the brain tissue[12–15]. Organoids of various parts of the brain, such as forebrain, midbrain, and hippocampus, are a powerful tool to study neurodegenerative diseases, neurodevelopmental disorders, and the interplay between brain cells under homeostatic and diseased conditions[16]. As the genetic manipulation of iPSCs is very well established, COs allow for the study of specific molecular mechanisms related to brain biology and neurovirology. Furthermore, COs express genes associated with human cortical development, and the developing neurons display active neural networks[17,18].

Despite their progress as a powerful model platform to study neuronal diseases at an organ level, COs lack proper abundance of cell types that are of non-ectodermal origin, such as microglia[19]. To circumvent this, short term co-cultures of pre-differentiated microglia are engrafted to organoids; however, the extent to which these incorporated microglia survive and function in these structures is unknown[20–23].

Human immunodeficiency virus (HIV) invades the brain as early as 8 days post-infection[24–26], initiating an inflammatory cascade leading to neuronal injury clinically manifesting as HIV associated neurocognitive disorders (HAND). Close to 50% of people living with HIV (PLWH) suffer from HAND[27,28], a spectrum of neurocognitive impairments that range from mild cognitive and motor impairments to severe dementia. HAND can significantly affect the quality of life and daily functioning of those impacted[29]. While mechanisms driving HAND are multifaceted, low level of HIV in the brain under antiretroviral therapy (ART) contributes to

persistent neuroinflammation[29,30]. Evidence of pockets of replication in peripheral tissues despite

suppressive ART exist, where the presence of low levels of viral RNA, DNA, and proteins, is

thought to be the primary driver of neuroinflammation[31].

There are numerous models to study HIV neuropathogenesis, each with its own

limitations. While human samples are ideal for studying HIV in the brain, post-mortem samples

of the brain are often difficult to obtain, and these cohorts are usually heterogeneous in terms of

their treatment regimens and comorbidities, as well as the method of preservation utilized during

brain harvest. It is also difficult to conduct longitudinal assessment of brain tissue from PLWH.

Non-human primates have served as an important model of HIV/SIV neuropathogenesis, yet

SIVmac and SHIVs are not genetically identical to HIV-1 and there are profound differences in

species-specific responses including differential molecular aspects to viral replication, disease

progression and responses[32,33]. Humanized mouse models while having human immune cells,

contain vast majority of host cells in the brain that are resistant to HIV infection[34]. These studies

are also conducted in the backdrop of graft versus host disease (GVHD)[35], which develops in all

humanized models and has to be controlled for[34,36]. These caveats to existing models to study HIV

in the brain underscore the need to develop models of brain cells in 3D culture systems that

recapitulate the human condition.

Microglia, the resident myeloid cells of the brain, are the major cell type infected by HIV

in the CNS[37,38]. Astrocytes, the most abundant glial cells in the brain, are infected to a lesser extent

and have mechanisms to suppress HIV transcription[39–42]. Macrophages and infiltrating T cells are

also known to harbor HIV in the brain[43–45]. Development of appropriate models to dissect the role

of microglia in HIV infection becomes paramount not only to understand HIV associated

neuroinflammation, but also to study the brain as a reservoir for HIV. This brain reservoir

contributes to the overall reservoir of HIV in the body, as HIV can egress from the brain to seed

other lymphoid tissues[46]. Recently, Tang *et al*., have shown that microglia isolated from adult

human brain are susceptible to HIV infection and lack significant viral genetic diversity, indicating

the microglial reservoir is seeded early in infection[47]. However, these studies exploit monolayer

cell culture of microglia in the absence of other cell types. Using innately developed microglia in

a CO model, Ormel *et al*. demonstrated HIV-1 infection of microglia[48]. In this model, the

limitations include presence of lower numbers of microglia (≤1%) compared to the human brain

and hence requirement of a high amounts of virus (100ng/CO) to achieve a successful infection.

dos Reis *et al*., engrafted infected microglia (adult primary or a cell line) into COs[49]. While this

method is adequate for incorporating microglia into COs, it has several drawbacks, as the microglia

engrafted into COs are previously infected rather than infected after incorporation into the tissue

environment, and the study uses cell cultures that are difficult to obtain for large-scale experiments.

Recently, in an elegant study, Schafer *et al*., xenotransplanted erythromyeloid progenitors

incorporated COs into adult NOD/SCID mouse brain and showed generation of functionally

mature human microglia that can operate within a physiologically relevant vascularized brain

environment[50]. Although this model can provide excellent insights into the biology and pathology

of human microglia, the laborious surgery methods make it impractical to conduct large scale

studies. Further, complications to study HIV pathogenesis may arise due to resistance of mouse

immune system to HIV infection and fast occurrence of GVDH to humanization.

Hematopoietic progenitors (HPCs) also known as erythromyeloid progenitors (EMPs)

derived from iPSC differentiate into microglia when properly cued[51]. Utilizing this principle, we

developed an approach to generate COs containing microglia by combining HPCs with iPSCs

during the first stage of embryoid body formation. We demonstrate successful proliferation,

differentiation, and maturation of incorporated HPCs into microglia in mature COs. These cerebral

organoids containing microglia (CO-iMs) express high levels of microglia specific homeostatic

and sensome markers. They were successfully infected with HIV and exhibited augmented

neuroinflammatory characteristics with infection. Further, by incorporating constitutively

expressing red fluorescent protein (RFP) into HPCs, we demonstrate that microglia in CO-iMs are

specifically amenable to genetic modifications.

Materials and Methods:

Culture and maintenance of iPSCs

Human iPSC lines were obtained from Rutgers University Cell and DNA Repository (RUCDR)

Infinite Biologics (ID# NN0003920, source; male, fibroblast) and Coriell Institute for Medical

Research (Camden, NJ; ID#AG27602, Source; male, fibroblast) maintained in complete mTeSR+

medium (Stem Cell Technologies; Vancouver, Canada) containing 0.1% penicillin/streptomycin

and propagated as small aggregates using accutase or ReLeSR (Stem Cell Technologies). All iPSC

colonies were cultured at 37°C in a 5% CO_2 incubator on 6-well tissue culture plates coated with

matrigel (Corning, NY). Coating of plates was performed by thoroughly mixing matrigel thawed

on ice with ice cold DMEM/F-12 medium at 1:100 dilution followed by incubating 1ml/well for

1h at room temperature (RT) or 37°C in 5% CO_2. iPSCs were routinely validated for markers,

OCT4, TRA-1-60 and Ki-67 using flow cytometry.

Labelling of iPSC with turboRFP and generation of HPCs

TurboRFP expressing lentiviral particles under the control of EF1a promoter (EF1A-RFP-LV) was

purchased from Cellomics (Halethorpe, MD, # PLV-10072). iPSC from RUCDR (termed iPSC-

Ru) were dissociated into single cells with accutase and plated at $0.3x10^6$ cells per well of a 6-well

plate in mTeSR+. Next day, cells were replaced with fresh mTeSR+ and infected with EF1A-RFP-

LV viral particles at 0.1 to 1.0 transduction units/cell. Three days later, wells were replaced with

fresh mTeSR+ media supplemented with puromycin (0.75 μg/ml). Media was changed every other

day and healthy colonies formed under lowest viral concentrations were expanded for at least two

more passages under puromycin (6-8 days per passage) before freezing the cells for stocking or

differentiating into HPCs.

Both wild type (WT) and RFP labeled iPSCs were differentiated to HPCs using STEMdiff

hematopoietic kit (Stem Cell Technologies). Briefly, iPSCs were dissociated into small colony

clumps via ReLeSR treatment (incubating as per STEMCELL protocol, followed by minimal

gentle pipetting) and plated at different dilutions of colonies per well of a 6-well plate in mTeSR+.

On the next 24 to 48h after confirming for optimal size and number of colonies (5-10 small

colonies/cm^2), mTSR+ media was replaced with fresh media A and the protocol was followed for

the next 12-14 days as per kit instructions (Stem Cell Technologies). On day 12 and day 14, round

floating HPCs were harvested and cultured for up to three more days in X-vivo-15 basal media

(Lonza, Switzerland), supplemented with 100 ng/mL M-CSF (Invitrogen, Massachusetts), 25

ng/mL IL-3 (R&D), 2 mM glutamax (Invitrogen), 0.055 mM β-mercaptoethanol (Invitrogen), 100

U/mL penicillin, and 100 μg/mL streptomycin. HPSCs phenotype was confirmed by flow

cytometry before the cells were used for experiments or frozen for later usage. HPCs were

routinely validated for expression of markers; CD45, CD43, and CD34 via flow cytometry.

Images of iPSC colonies and HPCs were captured under bright field or fluorescence using

a Keyence BZ-810 fluorescence microscope (Keyence, IL, USA) at 20x magnification. Cells were

frozen in Cryo-SFM freezing media (Promo Cell, Heidelberg, Germany).

Generation of cerebral organoids

Cerebral organoids were generated using STEMdiff cerebral organoid kit (Stem Cell

Technologies) with minor modifications[15,45]. On day 0, iPSC colonies were suspended into single

cells using accutase. To generate a regular CO, 10,000 iPSCs were incubated in organoid formation

media; alternatively, to generate a CO-iM, 10,000 iPSCs were co-cultured with 3,000 HPCs and

incubated in organoid formation media. Ultra-low binding, round bottom 96-well plates (Corning)

were used for embryoid body (EB) formation. Heat stable recombinant human bFGF (Gibco) at

5U/ml was supplemented to organoid formation media only on day 0. In the case of CO-iMs,

macrophage colony stimulating factor (MCSF, 50 ng/ml) and interleukin 3 (IL-3, 20 U/ml) were

used throughout organoid development to support survival, proliferation and differentiation of

HPCs to microglia. In order to prevent the fusion of multiple organoids during the maturation stage

(around day 20), each organoid was separated into single wells in low binding 24-well plates with

approximately 700 µl maturation media per organoid and incubated at 37°C, 5% CO_2 on an orbital

shaker at 90 rpm/min. Anti-adherent solution (Stem Cell Technologies) was used to create low

binding plates.

Single cell dissociation of organoids and flow cytometry

Organoids were gently washed with sterile HBSS, cut into small pieces, and incubated with papain

(30 units/ml, Stem Cell Technologies) along with human recombinant DNAseI (20 U/ml, Roche)

for 20-30 min at 37°C, 5% CO_2 on an orbital shaker at 90 rpm/min. Preparation and activation of

papain solution was performed as per the instructions (Stem Cell Technologies). One ml of papain

solution per organoid per well in a 12-well plate was used. Approximately, every 10 minutes,

organoids were dissociated by vigorous trituration (10~15 times) using a 1 mL pipette tip. By 30

min, most of the organoids dissociated into single cells and the remaining small tissue and cell

aggregates were gently pressed using the back of a 1ml syringe, and triturated. The whole cell

suspension was passed through a cell strainer (40 µM) and washed with 2-3ml of sterile HBSS.

The flow through containing single cells were centrifuged at 1300rpm for 8 min, washed again

with sterile HBSS, resuspended in 1ml HBSS, counted using a cell counter and processed for flow
cytometry. This protocol consistently resulted in >80% of cell viability as measured by tryphan
blue or calcein violet-AM (BioLegend).

iPSC colonies cultured on matrigel coated plates were detached with accutase and washed
twice with sterile PBS. Floating HPCs were also washed twice with sterile PBS. Single cells from
iPSC, HPCs or organoids were resuspended in 100 µl of PBS, incubated with Fc block (Cat
#564220, BD Biosciences, NJ) for 10 min at RT as per instructions and subjected to either cell
surface staining for surface markers using staining buffer (BD Biosciences), or intracellular
staining using perm/fix reagents (BD Biosciences). For intracellular staining, cells were incubated
with antibodies at 4°C for 60 min in the dark, washed three times thoroughly with perm/wash
buffer, resuspended in perm/wash buffer and run on a Fortessa flow cytometer (BD Biosciences).
Primary conjugated antibodies used for flow are listed in supplementary Table 1. Data was
analyzed using FlowJo software (v10.0.0) and normalized to the respective isotype IgGs or
fluorescent minus one (FMOs) controls. For compensation, 2 drops of anti-mouse Ig, k beads (BD
Biosciences) were incubated with primary antibodies for 20 min at 4°C, washed once with PBS,
then a drop of negative control BD comp beads (BD Biosciences) was added at the end and fixed
with 2% paraformaldehyde. FlowJo was used to set compensation parameters.

RNA and DNA extraction from organoids

Organoids were stored in RNAlater (500 µl/organoid) at -20°C until processing. RNA extraction
was performed using the RNeasy mini kit (Qiagen, Hilden, Germany). Briefly, organoid was
quickly brought to room temperature, RNA later was carefully removed and replaced with 500 µl
of RLT lysis buffer in sterile tubes containing lysing matrix D beads (Cat #6913100, MP
Biomedicals). Tubes were placed in bead beater (BeadBlaster 24, Bechmark Scientific, NJ) at a
speed of 4 M/S, for 30 sec with 15 sec intervals for 5 cycles to completely disintegrate and dissolve
organoids in RLT buffer. Further steps were followed according to kit protocol. Column DNAse
digestion was performed, and elution was done using only 40 µl of elution buffer. RNA was
quantified using a nanodrop and samples that had 260/280 ratio of >1.8 and 260/230 ratios between
2.0-2.2 were selected for further processing. Complementary DNA (cDNA) was synthesized using
a Qscript kit (QuantaBio, MA) as per kit instructions in a T-100 (Bio-Rad) thermocycler.

For DNA extraction, organoids were removed from RNAlater, cut into small pieces and

suspended in RLT buffer (700 µl/organoid) containing proteinase K. Samples were incubated at

56°C for 30 min-2h with intermittent vortexing until all the pieces were dissolved. Next steps were

followed according to DNAeasy Kit (Qiagen). DNA was eluted in 40 µl elution buffer and

quantified using a nanodrop. Samples with 260/280 ratio of >1.8 and 260/230 ratios between 2.0-

2.2 were considered for further processing.

Quantitative real time PCR

Real time quantitative RT-PCR (qRT-PCR) was performed on cDNA samples (at least 1/20

volume) using SYBR green master mix. Samples were loaded on a 96-well fast block and run on

a QuantStudio 5 Flex (Thermo Fisher) instrument. Protocol settings were as follows: hold stage

(95 °C for 10 minutes), 45 cycles of PCR stage (95 °C for 20 seconds and 60 °C for 1 minute) and

melt curve stage (95°C for 15 seconds, 60 °C for 1 minute and 95 °C for 15 seconds). Primers used

are listed in supplementary Table 2. Cycle threshold (Ct) values were subtracted with Ct values of

housekeeping genes (GAPDH and 18S), relative expressions of genes were computed using $2^{-\Delta\Delta Ct}$

method and are expressed as fold changes compared to respective control groups.

Similarly, real time quantitative PCR (qPCR) was performed on DNA samples using

Taqman based primers and probes for HIV DNA and human DNA essentially as previously

described[52]. Ct values of HIV amplifications were normalized to human DNA and are expressed

as fold changes compared to respective control groups.

HIV infection of cerebral organoids

Infectious particles of HIV-1$_{Ba-L}$ (Cat# ARP-510, BEI Resources) were produced in peripheral

blood mononuclear cells (PBMCs) isolated from a healthy seronegative donor using Ficol-

Hypaque density gradient centrifugation. Approximately 100 million cells were cultured in

complete RPMI 1640 medium which included 10% heat-inactivated fetal bovine serum (Gemini

Bio Products, Calabasas, CA), 2 mM L-glutamine (GlutaMAX supplement; ThermoFisher), 20

U/mL IL-2 (AIDS Reagent Program, Germantown, MD) and 1% penicillin/streptomycin (Sigma-

Aldrich, St. Louis, MO), Cells were activated for 3 days with 1 µg/mL/million cells of purified

NA/LE mouse anti-human CD3 and CD28 (BD Biosciences) and infected with HIV$_{BaL}$ at 0.01

ng/10^6 cells. PBMCs were monitored to express at least 6–10% intracellular p24$^+$ in T lymphocytes

before harvesting the supernatant. Supernatant was spun at 5000 rpm for 10 min to remove cell

debris and concentrated via Lenti-X as per the instructions (Takara Bio). HIV-Gag-iGFP_JRFL

(the following reagent was obtained through the NIH HIV Reagent Program, Division of AIDS,

NIAID, NIH: Human Immunodeficiency Virus (HIV) Gag-iGFP_JRFL, ARP-12456, contributed

by Dr. Benjamin Chen), containing an eGFP detailed in fused to GAG was produced and purified

as previously [53,54]. Purified viruses were quantified using a HIV1 p24 ELISA kit (#ab218268,

Abcam, Waltham, MA). Approximately, 60 to70-day old organoids were infected with 10ng/ml

concentration of HIV in a low-binding 24 well plates at one organoid/well/700 µl of maturation

media containing MCSF and IL-3. Three days later, the wells were gently washed and replaced

with fresh maturation media. cART treatment consisted of plasma levels of nevirapine (NVP, 10

µM) and dolutegravir (DTG, 3.90 µg/mL) that were added 24 h prior to infection, at the time of

infection, and after day 3 wash.

Immunofluorescence of organoids

Organoids were flash frozen fresh in cryomolds containing Tissue-Tek OCT (optimal cutting

temperature) compound. Organoid specimens were washed twice in PBS and acclimated to OCT

for few minutes in a small weight boat. Then, organoids were transferred into a labeled cryomold

where they were oriented and fully immersed in OCT while avoiding bubble formation. The

cryomolds containing the OCT immersed organoids were immediately transferred and submerged

for about 10 seconds into a cooling bath consisting of a beaker containing 2-propanol inside a dry

ice bucket until the OCT compound solidified or was visibly frozen. OCT blocks were kept on dry

ice until stored at -80°C. or sectioned (10-20µm) into cover slips using a microtome cryostat and

stored at -80°C.

Sections were subject to multiplex immunofluorescence (mIF) staining for target cells and

imaged by widefield deconvolution or spinning disk confocal microscopy.[55] mIF consisted of a

series of consecutive rounds of staining and bleaching after initial fixing and blocking of tissues.

Briefly, sections in cover slips were mounted and glued into home-made polyurethane resin

staining chambers and fixed for 5 minutes at room temperature in the dark using a PIPES/PFA

fixing solution (PIPES buffer pH 6.8, and 10% paraformaldehyde). After removing the fixing

solution, tissues were washed three times with PBS and blocked for 1 hour at room temperature

using donkey serum blocking buffer (PBS, donkey serum, 10% sodium azide, and 10% triton X-100). All staining cocktails were prepared in blocking buffer and sections were stained for 1 hour at room temperature or overnight at 4°C. Additionally, bleaching was accomplished by exposing the tissues to light for 1 hour at room temperature in bleaching buffer (7 volumes of distilled water, 2 volumes of 1M NaHCO₃, and 1 volume of 30% H₂O₂). After removing the bleaching solution, tissues were washed three times with PBS and kept on PBS at 4°C, or immediately used for a new round of staining. Antibodies used to stain COs are indicated and listed in supplementary Table 3. Ashlar (v1.18.0 https://github.com/labsyspharm/ashlar) was used to align max intensity projection of the Z-stacks (0.5um spacing) for the multiple mIF rounds with max pixel deviation of 200 and filter sigma of 2.

Images were acquired with a Nikon wide-field Ti-e2 equipped with an LED light source (Spectra X) and a PCO edge 4.2BI camera using standard Chroma DAPI/GFP/TRITC/Cy5 polychroic and emission filters for DAPI/GFP/TRITC with oil 40x magnification or with a Nikon Ti-e2 spinning disk confocal (Crest X-Light V3) equipped with a celesta laser light source. Widefield images obtained with the widefield Nikon TIe-2 microscope were deconvolved with FlowDec using the Lucy-Richardson algorithm with the support of Pims relying on python 3.x using calculated PSFs from FlowDec as previously reported [56,57].

Multiplex ELISA

Luminex assay for simultaneous quantification of multiple cytokines and chemokines (custom built, premixed kit) in the organoid supernatants were performed in a 96-well microplate format as per kit instructions (Luminex performance assay, R & D Systems). Data capture was done using a Luminex Flexmap 3D analyzer (ThermoFisher) and data analysis used a five-parameter logistic (5-PL) curve fit of a standard curve per analyte.

Bulk RNA sequencing (RNAseq), Single cell RNA sequencing (scRNAseq), and bioinformatics analysis.

Total RNA was extracted from organoids using the RNeasy kit (Qiagen) and checked for concentration, RIN/RQN and 28S/18S ratios using Agilent 4150 TapeStation system (Agilent Technologies). Transcriptome sequencing was performed by DNBSEQ™ sequencing technology platform (BGI/Ionomics) which includes stranded library preparation, 100 bp paired-end

sequencing and ≥30million reads. Sequencing data was demultiplexed and trimmed using

Trimmomatic v0.36 to remove adapters and low-quality reads. Trimmed reads were aligned to the

Homo sapiens reference genome GRCh38 and transcripts quantified using the Hisat2-StringTie

pipeline[58]. Differential gene expression analysis of the quantified gene transcripts was performed

with DESeq2 v.1.42.0 R package using R v.4.3.2. After retaining genes with nonzero total read

count, we identified differentially expressed genes (DEGs) either between CO-iMs and COs; HIV-

infected and uninfected CO-iMs; or between ART-treated and untreated HIV-infected CO-iMs

using as cut-offs an absolute log2 fold change > 1 and a false discovery rate (FDR) < 0.05 using

the Benjamin-Hochberg procedure. Gene enrichment analyses for each comparison were

subsequently performed using gene set enrichment analysis (GSEA) to identify specific Gene

Ontologies (GO), KEGG, and REACTOME pathways associated with CO-iMs and/or HIV

infection and ART treatment. We performed GSEA using clusterProfiler v.4.10.0 in R with all

lists of genes ranked by the corresponding log2 fold change. For these analyses all genes whose

gene symbols could be mapped to ENTREZ Ids using the org.Hs.eg.db v.3.18.0 Bioconductor

annotation package were included.

For scRNAseq four CO-iMs were individually processed for isolation of single cells using

papain and DNAseI treatment aspreviously described. The single cell suspensions from the

organoids were resuspended in Parse cell buffer containing BSA, manually counted using a

hemocytometer and assessed for cell viability by calcein violet-AM. The samples were centrifuged

at 1300 rpm for 10 minutes and then fixed/permeabilized using the Evercode fixation v2 kit (Parse

Biosciences, ECF2101) and stored at -80°C according to the instructions. The bar-coding and

library preparation for single cells were performed using the spit-pool based approach (Evercode™

WT Mini v2, Parse Biosciences). In brief, ∼27,000 fixed cells from each organoid were pooled

and underwent three rounds of barcoding. The final pooled cells were divided into 2 equal libraries

for cDNA synthesis and library construction to recover 10,000 cells. Multiplexed libraries were

sequenced using an Illumina NovaSeq X Plus system, with a 10B Flowcell and PE150 sequencing.

Fastq files were generated from bcl files and processed using the 'Parse Biosciences analysis

pipeline' using the GRCh38 reference genome to generate a gene expression matrix. After quality

filtering, 6,669 and 6,331 estimated cells for each library were analyzed with 92,691 and 91,357

reads per cell, respectively. 40,666 and 40,588 genes were analyzed with a median of 4,387 and

4,279 genes per cell for each library. Seurat v.5.0.3 R package was used for analysis. We filtered

cell barcodes with reads from less than 200 genes or more than 7,500, as well as barcodes with

more than 10% of mitochondrial genes that resulted in 11,440 remaining barcodes for analysis.

We used the harmony method with 5000 variable features and 30 principal components to integrate

the dataset from the two libraries and the Leiden method for clustering with 0.3 resolution using

10 principal components. DEGs that were upregulated in each cluster and expressed by a minimum

of 25% of the cells in that cluster compared to the rest of the cells were identified using Seurat's

implementation of the MAST algorithm Finally, ScType method R function using the curated

ScType database filtered for brain tissue excluding cancer cellswas used to annotate the clusters

obtained.

Statistics

Statistical analyses were performed using Prism software (GraphPad Prism, San Diego, CA).

Comparisons between groups were done using an unpaired, non-parametric, Mann-Whitney test.

Experiments were conducted independently at least three times, and the data represented as

standard error of mean (SEM). p value ≤ 0.05 is considered statistically significant. * Represents

$p \leq 0.05$, ** represents $p \leq 0.01$ and *** represents $p \leq 0.001$.

Results:

Microglia are efficiently differentiated in organoids produced by combining iPSC with HPCs

during EB formation.

Given that HPCs (or erythromyeloid progenitor cells, EMPs), when provided with proper cues can

differentiate into microglia *in vitro*[58], we hypothesized that by mixing HPCs with iPSCs during

EB formation stage and providing appropriate conditions (MCSF and IL-3) allows HPCs to

proliferate and differentiate into microglia in line with organoid maturation. We first confirmed

iPSC lines to be of high quality by morphology (colonies with tight borders with less than 5%

differentiation, **Suppl. Fig. 1A**) and by phenotype (expression of classical markers, OCT-4 and

TRA-1-60 pluripotency markers, and Ki-67 proliferation marker, **Suppl. Fig. 1B**). We then

differentiated unlabeled, or RFP labeled, iPSCs into HPCs and confirmed for its small spherical

floating cells morphology (**Suppl. Fig. 1C**), and expression of key progenitor marker, CD43

(>80% of cells) along with stemness marker, CD34 (~60% of cells) and myeloid marker, CD45

(**Suppl. Figs. 1D and E**). In addition, efficient labeling of iPSCs and HPCs with constitutively

expressing turboRFP was also confirmed (**Suppl. Figs. 1F-I**). Next, we generated unguided

cerebral organoids (COs), as described[12,48] and generated CO-iMs by mixing iPSCs with HPCs at

a 10:3 ratio on day 0 and incubated with MCSF (50ng/ml) and IL-3 (20U/ml) throughout the

formation, expansion and maturation stages. The schematic in **Figure 1A** outlines both

approaches. We did not detect noticeable changes in the morphological features of organoids at

any stage by adding HPCs in CO-iMs from the conventional COs developed only from iPSCs (**Fig.**

1B). Further, incorporation of HPCs did not affect the size of the COs at any measured

developmental stage (**Fig. 1C**). Cyclic multiplex immunofluorescence (mIF) imaging of

approximately 60 to 70-day old CO-iMs revealed robust presence of $TMEM119^+$ as well as $Iba1^+$

microglia cells (**Figs. 1D-F and Suppl. Figs. 2A-D**). Furthermore, mIF confirmed the presence of

similar numbers of astrocytes ($GFAP^+/S100\beta^+$) and neurons ($Tuj1^{+/}NeuN^+$) in CO-iMs and COs.

Additional lineage specific markers for glial and neuronal cells were used as well, which revealed

the presence of differential marker expression on these cell types, for example not all $GFAP^+$ are

$S100\beta^+$. Similarly, not all $NeuN^+$ are $Tuj1^+$ or $MAP2^+$ (**Figs. 1D-F and Suppl. Figs. 2A-D**). We

quantified astrocytic ($GFAP^+$ and $S100\beta^+$), neuronal ($Tuj1^+$) or NPC ($Nestin^+$) populations in these

organoids via flow cytometric analysis (**Suppl. Figs. 2E-H**). Incorporation of HPCs did not affect

the relative proportion of NPC (~30-37% $Nestin^+$), astrocytes (~38-40% $S100b^+$ and ~45-55%

$GFAP^+$), or neuronal cells (~45-50% $Tuj1^+$) between COs and CO-iMs (**Suppl. Figs. 2I and J**).

However, there was a significant increase (~4.6 fold) in myeloid ($CD45^+$) cells in CO-iMs

compared to COs (**Figs. 2A and B).** In line with these findings, we observed a significant increase

(~3.5-4 folds) in CD11b/Iba1 double positive cells (**Figs. 2C**). As expected, CD11b/Iba1 double

positive cells were only found in $CD45^+$ population and they were devoid in $CD45^-$ population

(**Fig. 2A**, bottom dot plots). Interestingly, within the $CD45^+$ myeloid population, there were similar

proportions of CD11b/Iba1 single positive or double positive cells, both in COs and CO-iMs (**Fig.**

2D). In parallel, in organoids generated using iPSCs-CRL, there was a significantly higher (~2

folds) proportion of CD11b/Iba1 double positive cells; however, there was no difference in $CD45^+$

population in CO-iMs when compared to COs (**Figs. 2E and F**). In addition, in CO-iMs generated

with incorporated HPCs stably expressing RFP, most of RFP^+ cells were $CD45^+$ (>60%, **Figs. 2G**

and H). As previously documented[44], we found $CD45^+/Iba1^+$ microglia population in COs, albeit

to lower levels (**Figs. 2B, C, F and H**). Taken together, these results demonstrate that HPCs

incorporated at the beginning of EB formation will successfully proliferate and differentiate into

mature microglia at the same timeframe with the overall growth and maturation of the neurons and

astrocytes in the brain organoids.

Transcription analysis exhibit induced expression of key microglia specific genes in CO-iMs

To further characterize the myeloid cells in CO-iMs for proliferation, abundance, differentiation

and maturation, we quantified transcriptional profiles of important homeostatic microglia markers

such as *CD11B*, *IBA1* and *TMEM119*[59] along with *CD45*, a myeloid marker expressed at lower

levels compared to monocytes/macrophages in steady state microglia[60]. We found that expression

levels of all these specific genes were significantly higher in CO-iMs when compared to

conventional COs. The trend was similar when normalized to either *GAPDH* (**Fig. 3A**) or *18S*

endogenous controls (**Suppl. Fig. 3A**). Interestingly, in CO-iMs, the level of *CD45* induction was

much lower compared to *CD11B* or *IBA1* (~8-10 fold vs 60-100 fold; **Fig. 3A**), suggesting that

low expression of *CD45* in combination with high *CD11B* perhaps indicates a resting steady-state

feature of microglia in CO-iMs[60,61]. Expression levels of astrocyte markers (*S100β, ACSBG1* and

EAAT1), neuronal progenitor/immature neuron markers (*Vimentin* and *Nestin*) and mature

neuronal markers (*MAP2* and *NEUN*) were similar between CO-iMs and COs (**Figs. 3B-D** and

Suppl. Fig. 3B). Further, *APOE*, a lipid transporter, mainly expressed in astrocytes was also found

to be unchanged (**Fig. 3B** and **Suppl. Fig. 3B).**

Similar observations with respect to gene expressions were noted in organoids generated

using iPSC-CRL line. Wherein, expression of *CD11B* and *IBA1* were significantly higher in CO-

iMs in comparison to COs (**Suppl. Figs. 3C and D**). *S100b* and *Nestin* levels were unchanged

between CO and CO-iMs (**Suppl. Figs. 3E and F**). The expression levels of *CD45* mRNA and

TMEM119 mRNAs were not significantly different between COs and CO-iMs with this iPSC line

(**Suppl. Figs. 3C and D**) as it was found with CD45 protein expression (**Fig. 2E**). These findings

reflect inherent intra- and inter-group variations as observed among batches and to the different

iPSCs cell lines[14,18,62].

To assess the differences between COs and CO-iMs, we characterized the global

transcriptome profile and compared COs and CO-iMs by RNAseq analysis. Gene enrichment

analysis using differentially expressed genes demonstrated striking differences in genes involved

in immunity such as phagocytosis, leukocyte migration, chemotaxis, cell migration, and others

(**Fig. 4A**) and pathways such as complement and coagulation, interactions of cytokine receptors

with cytokines and foreign antigens, TLR signaling, etc. (**Fig. 4B**). Differential gene expression

(DGE) analysis identified 1412 (5.6% of the total number of genes tested) genes that showed

significant upregulation in CO-iMs compared to COs (FDR<0.05 and log2 fold-change>1) and

1486 (5.9% of the total tested) downregulated genes, indicating significant changes associated to

the inclusion of microglia in the COs (**Fig. 4C and Suppl. File 1**). Subsequently, we performed a

Gene Set Enrichment Analysis (GSEA) with the entire list of quantified genes ranked by log2 fold

change of the CO-iM vs CO comparison. We observed a significant enrichment of genes related

to complement, cytokine expression, and innate immunity (**Figs. 4A** and **B**) amongst genes

upregulated in CO-iMs when compared with COs (**Fig. 4C**). While enrichment for genes related

to neuronal and astrocytic homeostatic function was detected amongst the downregulated genes.

Interestingly, we find genes with important brain functions to be statistically significantly

upregulated in CO-iMs, such as *CDA* and *PKD2L1* that have BBB and oligodendrocyte

myelination functions, respectively. Conversely, *LINGO4*, a gene anticipated to operate either

upstream or within the positive regulation of synapse assembly, is downregulated, which might

point to important microglia molecular function in synapse regulation as reviewed[64].

Microglia sense changes in their surroundings for immune regulation and host defense

functions. Therefore, we quantified the expression of microglia-specific sensome genes listed from

a well-established cluster[65,66]. Heatmap analysis indicated significant enrichment for the majority

of these genes in CO-iMs when compared to COs (**Fig. 4D**). Validation by qRT-PCR analysis for

some of these genes including *CX3CR1*, *TLR4*, *SLCO2B1*, *P2RY12, CSF1-R* and *TMEM119*

exhibited significant induction of all these genes in CO-iMs compared to COs, except for *P2RY12*

(**Figs. 3A, 4E** and **Suppl. Figs. 3A and 4A**). MCSF and IL34 are required for the survival,

proliferation, and maturation of microglia[61,62]. CSF1-R is the common receptor for MCSF and IL34.

We provided MCSF along with IL-3 throughout the culture period. Substantial induction of *CSF1-*

R mRNA in CO-iMs (**Fig. 4E** and **Supp. Fig. 4A**) along with similar yet noticeable levels of *IL34*

mRNA in both types of organoids (**Fig. 4F** and **Supp. Fig. 4B**) indicated a self-sufficient milieu,

not only for initial proliferation and differentiation of HPCs into microglia, but also for survival,

proliferation, and maturation of microglia in our 60+ day old, cultured organoids. Microglia plays

an important role in synapse pruning (removal of excess and less active synapses), via classical

complement pathway, in a developing brain. Synapse pruning is pivotal in maintenance and

refinement of synapses[67]. We assessed the expression levels of *C1Q* and *C3*, two early key players

of complement pathways via qPCR. These two mRNAs were significantly elevated in CO-iMs

(**Fig. 4G** and **Suppl. Fig. 4C**). Similar observation was made in CO-iMs generated from iPSC-

CRL, except for *C1Q*, as it did not change between two types of organoids (**Suppl. Figs. 4D**). In

addition to *C3*, heatmap analysis indicated induction of additional members of complement

pathways including *C7*, *C3AR1* and *MASP1* in CO-iMs (**Fig. 4H**). While *C6* was unchanged

between CO and CO-iMs, *C5* and *C4B* were significantly downregulated in CO-iMs (**Fig. 4H**),

suggesting a homeostatic maintenance of complement cascade. Collectively, these results

demonstrate that microglia generated in this manner in CO-iMs exhibit adult-like features

including expression of homeostatic and immune surveillance genes.

468   While targeted approaches and bulk RNAseq capture the overall trends of the CO-iMs, and

mIF can capture the different expression profiles and different spatial cellular distributions, they

are less powerful in detecting unknown or less abundant cell types and subtypes. Thus, to capture

the molecular identity of different cell types more accurately, we isolated single cells from four

CO-iMs (approximately 4-month-old), pooled into 2 groups and performed scRNAseq on a total

of 13,000 cells. Graph-based cell clustering based on transcriptional patterns detected 13 distinct

clusters that we could assign to 11 main CNS cell types (**Fig. 4I**), as indicated by the bigger bubbles

within each cluster in Fig. 4I, including various types of neurons (mature, serotonergic and others),

microglia (**Suppl. Fig. 3F**), astrocytes (**Suppl. Fig. 3H**), oligodendrocyte precursor cells, and

others. A heat map with the most highly expressed genes in each cluster is shown (**Suppl. Fig.**

3G). Taken together, scRNAseq confirmed the emergence of mature microglia in CO-iMs,

assuring that CO-iMs presented here are a good representation of different brain cells fortifying

the importance of these cell culture 3D models to neural and neurovirology research.

Microglia are robustly infected with HIV in CO-iMs

As CO-iMs contain sufficient and physiologically relevant proportions of microglia cells, we used

CO-iMs as a model to study HIV infection. Brain organoids have been previously used to model

HIV, CMV and ZIKV infections[48,49,63,64]. However, for HIV infections, mature microglia were

infected first and then allowed to traffic into COs[48,49]. Doing so will not allow us to study the

complete establishment, spread, and consequences of HIV infection at its different stages. To

address the questions, on the levels of microglia susceptibility to HIV and the neuropathology

within an organoid to closely mimic natural mode of infection in the CNS, CO-iMs were infected

with two different infectious HIV isolates, HIV-1$_{Ba-L}$ (R5 tropic) and HIV-Gag-iGFP_JRFL

(neurotropic envelope where the viral particles are green due to fusion of a fluid phase GFP within

Gag)[65]. HIV infection did not affect the morphology of the organoids up to the measured 30 days

post-infection (**Figs. 5A and B**), as reported for other viral infections[64,66,67]. Extracellular p24

quantification by ELISA indicated a significantly higher amount of virus released from CO-iMs

when compared to COs. This release was inhibited in the presence of cART, which was also used

as a control for viral input in the supernatant (**Fig. 5C**). Similarly, qPCR on these infected

organoids showed a 6 to 8-fold increase in HIV DNA when compared to COs (**Fig. 5D**). A

representative amplification plot for viral and human DNA is shown in **Figure 5E**. These data

demonstrate a robust productive infection in CO-iMs in comparison to COs. Cyclic mIF analysis

demonstrated co-localization of HIV p24$^+$ (mAb staining or iGFP-Gag+) in Iba$^+$ microglia cells

(**Figs. 5F-I**). Furthermore, we observed viral particles in regions surrounding microglia infected

cells and regions with heavy concentration of viral particles surrounding microglia that are not

infected (arrows with dotted lines **Fig. 5H**). Flow cytometry-based quantification analysis

indicated that HIV-iGFP+ (or p24$^+$) cells were positive for CD45$^+$, while the CD45$^-$ population

was negative (**Figs. 5J and K**). We also report significantly higher CD45$^+$/p24$^+$ cells in CO-iMs

compared to COs (**Fig. 5L**). Further, by flow cytometry we found that the majority of HIV$^+$

(iGFP+) cells were CD11b$^+$ which was almost undetectable after ART treatment (**Suppl. Fig. 5A**).

Our findings demonstrate robust HIV infection in microglia, which are also the major source of

productive infection in CO-iMs fully modeling results reported from post-mortem brains and

humanized mice studies.

HIV infection of CO-iMs results in heightened inflammatory condition.

To determine the immunological and cellular responses from HIV infection, we examined the

expression of inflammatory cytokines at the RNA and protein levels by bulk RNA sequencing,

qRT-PCR, and multiplex ELISA. Multiplex ELISA indicated significantly higher secretion of

several inflammatory cytokines such as IL-6, IP-10, TNFα, IL-8, MCP1, INFα, IL1β and INFβ in

the supernatant of CO-iMs infected with HIV compared to non-infected CO-iMs (**Fig. 5M**). We

then evaluated the mRNA expression of these cytokines under similar conditions via qRT-PCR.

We found that many of these cytokine transcripts were induced in response to HIV infection,

including *IP10* and *TNFα*, which were statistically significant (**Fig. 5N and Suppl. Fig. 5B**). In

parallel, heatmap analysis via RNAseq, showed a strong induction of many of the cytokines and

chemokines in CO-iMs infected with HIV when compared to non-infected controls, including IL6,

IP10 and TNFα (**Fig. 5O**), which was abrogated by cART treatment (**Fig. 5N and Suppl. Figs.**

5B, C). While *IL1β, INFβ, and IL1α* mRNAs showed no relative increase at the measured

timeframes post HIV infection (**Fig. 5N** and **Suppl. Fig. 5D**); their measured protein levels

significantly increased (**Fig. 5M**), which may reflect a longstanding permanence of these cytokines

in this system after transcriptional shutdown.

Transcriptome analysis between HIV infected and non-infected CO-iMs showed strong

upregulation and activation of immune responses (IFNs signaling), *IL-10* (log2FC = 5.65), *IL-6* (

log2FC = 3.58), *IL-33* (log2FC = 2.834) and antigen processing and presentation (**Figs. 6A and**

B). Conversely, HIV-1 infection resulted in suppression of pathways of glucuronidation, neuron

fate commitment, neuropeptide functions, axon and forebrain development, and lipoprotein

function (**Figs. 6A and B**). Employing DEG analysis, we identified a total of 1826 upregulated

genes (7% of the total number of genes tested) (FDR<0.05 and log2 fold-change>1) and 2158

(8.3% of the total tested) downregulated genes with HIV infection, indicating significant responses

to infection in CO-iMs (**Fig. 6C and Suppl. File 2**). Of note, we find that caspase-5, a protein that

is associated with pyroptosis, cell death, and the activation of the inflammasome as a response to

neuronal trauma resulting in cell death[68–70] is highly upregulated with infection (log2FC = 9.26).

Further, DTHD1, a widely uncharacterized protein found to be involved in apoptotic events in the

brain and found to be active and highly expressed in the brains of Alzhiemer's and Multiple

Sclerosis lesions[71,72] is highly upregulated (log2FC = 23.83). In addition, other transcripts that are

important for neuronal homeostasis and function, such as psynaptophysin (log2FC = -1.7); *MAP2*

(log2FC = -1.77), and *EAAT2* (log2FC = -1.46) were downregulated approximately between 60-

70% with infection. Strikingly, *NOS1* (log2FC = 22.34), a gene that is involved in nitric oxide

synthesis from L-arginine, which has many neurotransmitter properties and has been shown to

relate to neurotoxicity and neurodegenerative diseases was also highly upregulated with HIV

infection (**Fig. 6C and Suppl. File 2**). Interestingly, cART treatment with and without infection

only mildly reduced the expression of DEGs (<1%, **Suppl. Fig. 5E and Suppl. File 3**).

Taken together, these data demonstrate robust infection and replication of HIV in CO-iMs,
driven primarily by HIV infection of microglia in this system. The infection drives inflammatory
responses in these organoids that lead to poor neuronal health. Furthermore, we find that HIV-1
infection and replication of CO-iMs have an impact in neuronal function and development,
modeling one of the hallmarks of HIV-1 infection, where there is a big biological impact in the
brain.

Discussion

We developed a microglia containing CO model by co-culturing iPSCs and HPCs from the same
line, at the beginning stage of EB formation. In this 3D culture model, HPCs proliferate and mature
into microglia, as indicated by robust expression of microglia specific homeostatic markers, as
well as sensome markers in 60+ day old organoids. In the same system and in parallel, iPSCs
differentiate into NPCs, which further differentiate into mature neurons and astrocytes. Notably,
by introducing HPCs at the formation stage along the 3D formation stage (expansion to maturation
stage), rather than the distinct engraftment of fully differentiated microglia we gain unique
advantages such as: (i) mimicking the brain cell composition and cellular interactions from the
formation of CO-iMs; (ii) use of less HPCs/microglia per organoid as this method does not depend
on the efficiency of microglia post-engraftment; and, (iii) the ability to regulate the number of
HPCs and in turn microglia, per organoid. Further, by harnessing the self-organization capability
of iPSCs instead of NPCs we generated brain organoids (unguided type), rather than neurospheres,
which can mimic the cytoarchitecture and developmental trajectories found. Importantly, we show
that microglia in our CO-iM model can be successfully and preferentially infected with HIV at a
particularly high percentage which led to heightened inflammatory responses.

Microglia are known to account for around 5–10% of total cells in the CNS[73] with
astrocytes and neurons being the major cell types at similar proportions, depending on brain region
and size of neurons, but typically within 20-40%[74]. In the CO-iM model, on average, around 7.3%
of total cells were microglia (double positive for CD11b and Iba1) compared to less than 1% in
conventional COs. In addition, astrocytes (around 35-50%) and neurons (around 44-51%) were
abundant, and their defining transcripts and antigens were well expressed in CO-iM. Thus, this
novel method can generate organoids with physiologically relevant contents of microglia,

astrocytes, neurons, and other brain cells providing a platform to study interactions between CNS
cell types in a 3D conformation.

CSF1 and IL34 are important proteins for growth and maturation of microglia.
Significantly induced expression of *CSF-1R*, a common receptor for both CSF1 and IL34 in
combination with strong expression of *IL34* at transcript levels suggests a self-sufficient milieu for
survival and maturation of microglia in our CO-iM organoids. Further, significant induction of
multiple sensome markers and the general immunity related transcripts by RNAseq in CO-iMs is
a compelling indication that microglia are functional in our organoid model. In addition, elevated
levels of *C1q* and *C3*, the two main players of initial activation of complement cascade, along with
C3AR1, C7 and *MASP1* suggests that microglia may be playing an important role in synapse
pruning and refinement in our CO-iM model. At the same time, heightened complement activation
could be detrimental and is associated with cognitive impairment because of excessive pruning[75–
77]. CO-iMs exhibited downregulation of some of its components including *C4B* and *C5* suggesting
a balanced regulation of pathway to maintain a homeostatic condition. Keeping in mind the high
interest of late to develop targeted therapeutics against this pathway to mitigate neurodegeration[78–
81], we believe that CO-iMs could provide a good platform toward high throughput screening of
small molecules and drugs that can specifically regulate complement activation. Taken together,
we can conclude that microglia in CO-iMs are functional for their known immunity and particular
susceptibility to HIV infection.

HIV enters the brain within two weeks of acute infection and within the brain, microglia
and macrophages are the major targets of HIV infection. In the post cART era, it is imperative to
understand and dissect the role of microglia in HIV neuropathogenesis to gain insights into two
major unsolved issues, HAND and HIV latency. Several recent studies have shown that microglia
are successfully infected with HIV resulting in abrogated immune responses. Limitations of these
studies include either the use of monolayer cultures of microglia[47], incorporation of externally
infected microglia cells into maturing organoids[82], or use of high amounts of virus particles to
infect low abundant microglia that are developed innately in COs[48]. As microglia do not present
the well-characterized SAMHD1 restriction to HIV-1 in myeloid cells (macrophages and dendritic
cells)[83], we found them to be infected at a particularly high ratio soon after viral inoculation (within
5 days) where the viruses were added into the culture media and crossed the organoid tissue layers

to initiate replication deep into the organoids. We find that when comparing CO-iMs with and without HIV infection, there is a high upregulation of immune markers and their release with infection. We also compared inoculated CO-iMs with and without antiretrovirals that prevent seeding of proviruses into the host genome and found that the transcriptome related to immunity is dependent on viral replication. These allowed us to exclude the effect of extracellular/non-cytoplasmic immune responses such as TLR activation due to the presence of viruses in the inoculum. Interestingly, we find that not all microglia become infected despite the presence of high virus amounts in regions seeded with microglia, denoting a diversity of cells and a possible protective early triggering of innate responses and cytokine milieu from the initial rounds of replication that warrants further investigation. Infection of the CO-iMs result in strong neuroinflammation profiles, both at a transcriptomic (**Figs. 5N and O**) and cytokine level (**Fig. 5M**), and infection results in a transcriptomic profile associated with suppression of normal neural function and development (**Fig. 6**). Despite controlling viral spread and systemic replication, modern antiretroviral therapies do not prevent any of the late-stages of the HIV life cycle as they focus on reverse transcriptase and integrase inhibitors, allowing for the production of nucleic acids, viral protein, and viruses capable of maturation and fusion to other cells[84–87]. Thus, our model provides a unique platform to test specific molecular interactions between HIV particles and the cellular innate immune machinery that detects HIV and is known to mount cascades of immune responses. For example, innate sensing of foreign nucleic acids by cGAS/STING in blood myeloid cells occurs within the first hours of infection followed by a severe down regulation of their activation[88–90].

The CO-iM model offers a suitable platform where microglia with or without infection can crosstalk with other glial and neuronal cell types in a human-brain-like environment. As the cellular origin of CO-iMs are iPSCs, this is a suitable platform to detail mechanistic functions related to specific genes and to specific cellular lineages (HPCs vs NPCs), as iPSCs are easily genetically manipulated. In the context of HIV, this platform provides a unique opportunity to model the interaction of human microglia with their neuronal environment and renders it suitable for unmasking the role of microglia and its specific genes in viral-related inflammation, its role in HIV persistence under cART, and the biological consequences of infection to astrocytic and neuronal health.

Author contributions

JIM, SDN, and LA conceptualized and designed the study. SDN, JPZ, JIM performed the study. All interpreted the data. MKA contributed to the generation of organoids, and manuscript editing. AR, TS and SG helped with organoid slicing, qRT-PCR and p24 ELISA, respectively. RLR performed RNAseq and scRNAseq analysis. SDN and JIM wrote the manuscript with input from all authors.

Declaration of interests

Authors declare no competing interests.

Acknowledgments

This work was in part supported by grants from the NIH R01DA055497 and R01NS108796 to LA, Walder Foundation innovation Top Up award to JRS, R01MH125778, P01AI169600, and the P30AI117943 Third Coast Center for AIDS Research (TC-CFAR) to RLR, and R21MH129205 and R61DA058348 to JIM. Library preparation and sequencing were performed at Northwestern University NUSeq core. Cytokine panels were performed by the Rush University Biomarker Development core.

Figure Legends:

Figure 1: Generation and characterization of CO-iMs. (A) Schematic flow diagram indicating the timeline as well as all developmental stages for generation of CO and CO-iMs. **(B)** Representative bright field images of a CO and a CO-iM through different stages of their development. Scale bars, 100 μm. **(C)** Diameter sizes of CO and CO-iMs measured at key time points during their differentiation (n=5 per time point). **(D)** Left, representative multiplex IF image of a CO-iM stained for nucleus (Hoechst, white), microglia (TMEM119, red), neurons (green, Tuj1) and astrocytes (GFAP, blue) Scale bar, 150 μm. Panel on the right showing individual staining for indicated markers and a merged image from all. Scale bars, 100 μm. **(E and F)** Zoomed insets of i), and ii) regions in (D). Scale bars, 100 μm.

Figure 2: Phenotypic characterization reflecting enrichment of microglia in CO-iMs. CO and CO-iMs were processed and dissociated into single cell suspension, stained, and analyzed for cell surface and intracellular proteins by flow cytometry. **(A)** Representative dot plots showing $CD45^+$ and $CD45^-$ populations from total cells (top), and $CD11b^+$ and $Iba1^+$ populations (bottom). Arrows indicate populations analyzed within $CD45^+$ or $CD45^-$ gates. **(B and C)** Cumulative data indicating percent $CD45^+$ and CD11b /Iba1 double positive cells in total population of organoids generated using iPSC-Ru line (n=7 for COs and CO-iMs, respectively) **(D)** Cumulative data showing percent CD11b/Iba1 single positive as well as double positive cells within $CD45^+$ population in organoids generated using iPSC-Ru line (n=3 for COs, and n=5 for CO-iMs). **(E and F)** Cumulative data showing percent CD11b/Iba1 single as well as double positive cells within $CD45^+$ population in organoids generated using iPSC-CRL line (n=4 for COs, and n=6 for CO-iMs). **(G)** Representative dot plots showing that majority of RFP^+ cells are also $CD45^+$ cells. **(H)** Cumulative data indicating percent $RFP^+ CD45^+$ population (n=3 for COs, and n=5 for CO-iMs).

Each symbol represents an individual organoid. Data are presented as mean ± SD using non-parametric Mann-Whitney test. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.

Figure 3: Genotypic characterization reflecting enrichment of microglia markers in CO-iMs. qRT-PCR analysis of COs (n=3-5) and CO-iMs (n=5-7). mRNA levels were normalized to the endogenous reference GAPDH and expressed as fold change relative to CO using $2^{-(\Delta\Delta Ct)}$. Fold change for each sample in COs was assessed by subtracting its delta Ct from average delta Ct, followed by $2^{-(\Delta\Delta Ct)}$. **(A)** mRNA levels of homeostatic microglia markers: *CD45*, *CD11B*, *IBA1* and *TMEM119*. **(B)** mRNA levels of astrocyte markers: *S100b*, *ACSBG1*, *EAAT1* and *APOE.* **(C and D)** mRNA levels of neuronal progenitor and mature neuronal markers: *Vimentin, Nestin*, *MAP2* and *NeuN*. Each symbol represents an individual organoid. Data are presented as mean ± SD using non-parametric Mann-Whitney test. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.

Figure 4: Transcriptional Profile of CO and CO-iMs. RNA-seq analysis of ~65-day old iPSC-Ru line derived COs (n = 3) and CO-iMs (n = 3). **(A)** Dot plot showing the enriched GO terms of biological processes and molecular functions amongst activated and suppressed genes identified using absolute log2 fold change > 1 and a false discovery rate (FDR) < 0.05 using the Benjamin-

Hochberg procedure in CO-iMs when compared to COs. **(B)** Dot plot showing the enriched

REACTOME pathways (FDR <0.1) in activated as well as suppressed genes in CO-iMs when

compared to COs. For **(A)** and **(B)** the x-axis indicates the ratio of genes identified by our analysis

and the total number of genes that constitute the pathway, the size of the dot is based on gene count

in the pathway, and the color of the dot shows the pathway enrichment significance. **(C)** Volcano

plot comparison of CO-iMs vs COs indicating average log2 (fold change) versus log10 (FDR) for

all genes. Genes upregulated and downregulated by 2-fold change and FDR < 0.05 are labeled

with red dots. **(D)** A heatmap showing expression of microglia sensome markers in individual

replicates of CO-iMs and COs, (n = 3 for CO-iMs and n = 3 for COs). Color indicates the

expression level scaled for each gene by centering and scaling. **(E)** qRT-PCR quantification to

determine mRNA levels of microglia immune sensome markers: *CX3CR1*, *SLCO2B1*, *TLR4*,

CSF1-R and *P2RY12* **(F)** qRT-PCR quantification to determine mRNA levels of *IL34*, an essential

cytokine for microglia maturation and survival. **(G)** qRT-PCR quantification to determine mRNA

levels of *C3* and *C1Q*, components of complement pathway. **(H)** Heatmap showing expression of

microglia complement pathway genes in individual replicates of CO-iMs and COs, (n = 3 for CO-

iMs and n = 3 for COs). Color indicates the expression level scaled for each gene. **(I)** Uniform

Manifold Approximation and Projection (UMAP) plot for dimension reduction of 11,440 barcoded

single cells derived from 4 individual CO-iMs (~120 day old) subjected to scRNA seq analysis

exhibiting 12 distinct clusters as identified by graph-based clustering of cell-specific gene

expression. Subsequent cell annotation allowed us to define the most likely cell type corresponding

to each cluster. The bubble plot shows all the cell types that were considered by ScType for cluster

annotation that were assigned to at least 100 cells with the exception of cluster 10 where no

assignment reached that cut-off and the only assignment obtained is shown. The outer bubbles

correspond to each cluster with size reflecting the number of cells in the cluster, while the inner

bubbles correspond to considered cell types for each cluster, with the biggest bubble corresponding

to assigned cell type, except for cluster 3 and 10 where we did not obtain high quality assignments.

Figure 5: Robust response of microglia to HIV-1 infection in CO-iMs. **(A)** Representative

bright field images showing the morphology of CO and CO-iMs (~60 day old) after 15 or 30 days

post infection (DPI) with HIV-1 or mock. Scale bars, 500 μm. **(B)** Diameter sizes of CO and CO-

iMs measured 15 DPI or 30 DPI with HIV or mock (n=2 for COs, and n=3 for CO-iMs). **(C)**

Organoids were infected with HIV_{BaL} (at 10 ng/ml) on day 0. At 3 DPI, organoids were washed

once with PBS and replaced with fresh media. Then, at 5 DPI, the culture supernatant was collected

and used to quantify p24 by ELISA. n=3 for each condition with organoids generated from two

different hiPSC cell lines (RUCDR and CRL, respectively). **(D)** Organoids were infected with

HIV_{BaL} or HIV-iGFP-JRFL as explained in (C), genomic DNA was extracted and quantified for

HIV DNA using Taqman qPCR and represented as fold change to COs infected with HIV_{BaL} or

HIV-iGFP_JRFL (n=3 for COs, n=3 for CO-iM + HIV_{BaL}, and n=4 for CO-iMs+HIV-iGFP_JRFL).

(E) A representative Taqman qPCR amplification plot showing amplification of human DNA as

well as HIV-GAG for all samples shown in (D). **(F)** Representative mFI images of a CO-iM

showing IBA1$^+$/HIV$^+$ microglia after 5 DPI HIV infection. Scale bars, 100 μm. **(G)** Zoomed in

insets of regions i) and ii) from (G) showing p24$^+$ (green), Iba1$^+$ (red) cells. Scale bars, 25 μm. **(H**

and I) mFI images of infected CO-iMs exhibiting p24$^+$ (green) and Iba1$^+$ (red) co-expressing cells

(arrows with solid lines) as well as free virus (arrows with dotted lines) nearby Iba1$^+$ cells. Scale

bars, 50 and 10 μm, respectively. **(J and K)** Organoids were infected with HIV_{BaL} for 5 days as

explained, dissociated into single cell suspension and stained for cell surface (CD45) and

intracellular (p24) markers for flow cytometry analysis. Representative dot plots for CD45 and

p24 populations in COs (J, left) and in CO-iMs (K, left). **(L)** Cumulative data indicating percentage

of p24$^+$ cells in CD45$^+$ and CD45$^-$ populations, respectively. n=3 for COs, and n=4 for CO-iMs.

(M) Multiplex ELISA to quantify various pro-inflammatory cytokines/chemokines in the

supernatants of CO-iMs infected with HIV_{BaL} for 5 days (n=6), or mock (n=6) for 5 days. **(N)** qPCR

analysis for mRNA expression of various pro-inflammatory cytokines/chemokines (as indicated

in the figures) in CO-iMs either infected with HIV_{BaL} for 5 days, or mock infected, or infected and

treated with cART. n=3 in each category. (O) Heatmap showing relative expression of a list of

inflammatory cytokines/chemokines in infected vs uninfected CO-iMs (n = 3 for each group).

Color indicates the expression level scaled for each gene.

Each symbol represents an individual organoid. Data are presented as mean ± SD using non-

parametric Mann-Whitney test. *p < 0.05, **p < 0.01, ***p < 0.001.

Figure 6: Transcriptional Profile of CO and CO-iMs infected with HIV. RNA-seq analysis

of 65-day old iPSC-Ru line derived CO-iMs infected with HIV_{BaL} (n = 3) for 5 days or mock

infected (n = 3) or infected+cART treated (n = 3) for 5 days. **(A)** REACTOME dot plot showing

the pathways (FDR <0.1) enriched in activated and suppressed gene in infected CO-iMs when

compared with their uninfected counterparts. **(B)** Dot plot showing the GO terms (FDR <0.05) of

biological processes enriched in activated and suppressed genes in infected CO-iMs when

compared with their uninfected counterparts. For (A) and (B) the x-axis indicates the ratio of

genes identified by our analysis and the total number of genes that constitute the pathway, the

size of the dot is based on gene count in the pathway, and the color of the dot shows the pathway

enrichment significance. **(C)** Volcano plot comparison of infected vs uninfected CO-iMs

indicating average log2 (fold change) versus log10 (FDR) for all genes. Genes upregulated and

downregulated by 2-fold change and FDR < 0.05 are labeled with red dots.

References

1. Mizee, M.R., Miedema, S.S.M., van der Poel, M., Adelia, Schuurman, K.G., van Strien, M.E., Melief, J., Smolders, J., Hendrickx, D.A., Heutinck, K.M., et al. (2017). Isolation of primary microglia from the human post-mortem brain: effects of ante- and post-mortem variables. Acta Neuropathol. Commun. *5*, 16. https://doi.org/10.1186/s40478-017-0418-8.

2. Cochrane, CatherineR., Angelovich, ThomasA., Byrnes, SarahJ., Waring, E., Guanizo, AleksC., Trollope, GemmaS., Zhou, J., Vue, J., Senior, L., Wanicek, E., et al. (2022). Intact HIV Proviruses Persist in the Brain Despite Viral Suppression with ART. Ann. Neurol. *92*, 532–544. https://doi.org/10.1002/ana.26456.

3. Feng, G., Jensen, FrancesE., Greely, HenryT., Okano, H., Treue, S., Roberts, AngelaC., Fox, JamesG., Caddick, S., Poo, M., Newsome, WilliamT., et al. (2020). Opportunities and limitations of genetically modified nonhuman primate models for neuroscience research. Proc. Natl. Acad. Sci. PNAS *117*, 24022–24031. https://doi.org/10.1073/pnas.2006515117.

4. Jennings, C.G., Landman, R., Zhou, Y., Sharma, J., Hyman, J., Movshon, J.A., Qiu, Z., Roberts, A.C., Roe, A.W., Wang, X., et al. (2016). Opportunities and challenges in modeling human brain disorders in transgenic primates. Nat. Neurosci. *19*, 1123–1130. https://doi.org/10.1038/nn.4362.

5. Agarwal, Y., Beatty, C., Biradar, S., Castronova, I., Ho, S., Melody, K., and Bility, M. (2020). Moving beyond the mousetrap: current and emerging humanized mouse and rat models for investigating prevention and cure strategies against HIV infection and associated pathologies. Retrovirology *17*, 8. https://doi.org/10.1186/s12977-020-00515-3.

6. Urbach, Y.K., Bode, F.J., Nguyen, H.P., Riess, O., and Von Hörsten, S. (2010). Neurobehavioral Tests in Rat Models of Degenerative Brain Diseases. In Rat Genomics, I. Anegon, ed. (Humana Press), pp. 333–356. https://doi.org/10.1007/978-1-60327-389-3_24.

7. Pauletti, A., Terrone, G., Shekh-Ahmad, T., Salamone, A., Ravizza, T., Rizzi, M., Pastore, A., Pascente, R., Liang, L.-P., Villa, B.R., et al. (2019). Targeting oxidative stress improves disease outcomes in a rat model of acquired epilepsy. Brain *142*, e39–e39. https://doi.org/10.1093/brain/awz130.

8. Gage, F.H., and Temple, S. (2013). Neural Stem Cells: Generating and Regenerating the Brain. Neuron *80*, 588–601. https://doi.org/10.1016/j.neuron.2013.10.037.

9. Dezonne, R.S., Sartore, R.C., Nascimento, J.M., Saia-Cereda, V.M., Romão, L.F., Alves-Leon, S.V., de Souza, J.M., Martins-de-Souza, D., Rehen, S.K., and Gomes, F.C.A. (2017). Derivation of Functional Human Astrocytes from Cerebral Organoids. Sci. Rep. *7*, 45091. https://doi.org/10.1038/srep45091.

10. Sun, N., Meng, X., Liu, Y., Song, D., Jiang, C., and Cai, J. (2021). Applications of brain organoids in neurodevelopment and neurological diseases. J. Biomed. Sci. *28*, 30. https://doi.org/10.1186/s12929-021-00728-4.

11. Eichmüller, O.L., and Knoblich, J.A. (2022). Human cerebral organoids — a new tool for clinical neurology research. Nat. Rev. Neurol. *18*, 661–680. https://doi.org/10.1038/s41582-022-00723-9.

12. Lancaster, M.A., Renner, M., Martin, C.-A., Wenzel, D., Bicknell, L.S., Hurles, M.E., Homfray, T.,
Penninger, J.M., Jackson, A.P., and Knoblich, J.A. (2013). Cerebral organoids model human brain
development and microcephaly. Nature *501*, 373–379. https://doi.org/10.1038/nature12517.

13. Subramanian, L., Bershteyn, M., Paredes, M.F., and Kriegstein, A.R. (2017). Dynamic behaviour of
human neuroepithelial cells in the developing forebrain. Nat. Commun. *8*, 14167.
https://doi.org/10.1038/ncomms14167.

14. Velasco, S., Kedaigle, A.J., Simmons, S.K., Nash, A., Rocha, M., Quadrato, G., Paulsen, B., Nguyen, L.,
Adiconis, X., Regev, A., et al. (2019). Individual brain organoids reproducibly form cell diversity of
the human cerebral cortex. Nature *570*, 523–527. https://doi.org/10.1038/s41586-019-1289-x.

15. Di Lullo, E., and Kriegstein, A.R. (2017). The use of brain organoids to investigate neural
development and disease. Nat. Rev. Neurosci. *18*, 573–584. https://doi.org/10.1038/nrn.2017.107.

16. Hopkins, H.K., Traverse, E.M., and Barr, K.L. (2021). Methodologies for Generating Brain Organoids
to Model Viral Pathogenesis in the CNS. Pathogens *10*, 1510.
https://doi.org/10.3390/pathogens10111510.

17. Camp, J.G., Badsha, F., Florio, M., Kanton, S., Gerber, T., Wilsch-Bräuninger, M., Lewitus, E., Sykes,
A., Hevers, W., Lancaster, M., et al. (2015). Human cerebral organoids recapitulate gene expression
programs of fetal neocortex development. Proc. Natl. Acad. Sci. *112*, 15672–15677.
https://doi.org/10.1073/pnas.1520760112.

18. Quadrato, G., Nguyen, T., Macosko, E.Z., Sherwood, J.L., Min Yang, S., Berger, D.R., Maria, N.,
Scholvin, J., Goldman, M., Kinney, J.P., et al. (2017). Cell diversity and network dynamics in
photosensitive human brain organoids. Nature *545*, 48–53. https://doi.org/10.1038/nature22047.

19. Mansour, A.A., Schafer, S.T., and Gage, F.H. (2021). Cellular complexity in brain organoids: Current
progress and unsolved issues. Semin. Cell Dev. Biol. *111*, 32–39.
https://doi.org/10.1016/j.semcdb.2020.05.013.

20. Qian, X., Nguyen, H.N., Song, M.M., Hadiono, C., Ogden, S.C., Hammack, C., Yao, B., Hamersky, G.R.,
Jacob, F., Zhong, C., et al. (2016). Brain-Region-Specific Organoids Using Mini-bioreactors for
Modeling ZIKV Exposure. Cell *165*, 1238–1254. https://doi.org/10.1016/j.cell.2016.04.032.

21. Muffat, J., Li, Y., Yuan, B., Mitalipova, M., Omer, A., Corcoran, S., Bakiasi, G., Tsai, L.-H., Aubourg, P.,
Ransohoff, R.M., et al. (2016). Efficient derivation of microglia-like cells from human pluripotent
stem cells. Nat. Med. *22*, 1358–1367. https://doi.org/10.1038/nm.4189.

22. Abreu, C.M., Gama, L., Krasemann, S., Chesnut, M., Odwin-Dacosta, S., Hogberg, H.T., Hartung, T.,
and Pamies, D. (2018). Microglia Increase Inflammatory Responses in iPSC-Derived Human
BrainSpheres. Front. Microbiol. *9*.

23. Lin, Y.-T., Seo, J., Gao, F., Feldman, H.M., Wen, H.-L., Penney, J., Cam, H.P., Gjoneska, E., Raja, W.K.,
Cheng, J., et al. (2018). APOE4 Causes Widespread Molecular and Cellular Alterations Associated
with Alzheimer's Disease Phenotypes in Human iPSC-Derived Brain Cell Types. Neuron *98*, 1141-
1154.e7. https://doi.org/10.1016/j.neuron.2018.05.008.

24. Valcour, V., Chalermchai, T., Sailasuta, N., Marovich, M., Lerdlum, S., Suttichom, D., Suwanwela,
N.C., Jagodzinski, L., Michael, N., Spudich, S., et al. (2012). Central Nervous System Viral Invasion
and Inflammation During Acute HIV Infection. J. Infect. Dis. *206*, 275–282.
https://doi.org/10.1093/infdis/jis326.

25. Liu, J., Ghneim, K., Sok, D., Bosche, WilliamJ., Li, Y., Chipriano, E., Berkemeier, B., Oswald, K.,
Borducchi, E., Cabral, C., et al. (2016). Antibody-mediated protection against SHIV challenge includes
systemic clearance of distal virus. Sci. AAAS *353*, 1045–1049.
https://doi.org/10.1126/science.aag0491.

26. Deeks, S.G., Overbaugh, J., Phillips, A., and Buchbinder, S. (2015). HIV infection. Nat. Rev. Dis. Primer
1, 15035. https://doi.org/10.1038/nrdp.2015.35.

27. Wang, Y., Liu, M., Lu, Q., Farrell, M., Lappin, J.M., Shi, J., Lu, L., and Bao, Y. (2020). Global prevalence
and burden of HIV-associated neurocognitive disorder: A meta-analysis. Neurology *95*, e2610–
e2621. https://doi.org/10.1212/WNL.0000000000010752.

28. Zhou, L., and Saksena, N.K. (2013). HIV Associated Neurocognitive Disorders. Infect. Dis. Rep. *5*, e8.
https://doi.org/10.4081/idr.2013.s1.e8.

29. Spudich, S., Robertson, K.R., Bosch, R.J., Gandhi, R.T., Cyktor, J.C., Mar, H., Macatangay, B.J., Lalama,
C.M., Rinaldo, C., Collier, A.C., et al. (2019). Persistent HIV-infected cells in cerebrospinal fluid are
associated with poorer neurocognitive performance. J. Clin. Invest. *129*, 3339–3346.
https://doi.org/10.1172/JCI127413.

30. Gisolf, E.H., van Praag, R.M.E., Jurriaans, S., Portegies, P., Goudsmit, J., Danner, S.A., Lange, J.M.A.,
and Prins, J.M. (2000). Increasing Cerebrospinal Fluid Chemokine Concentrations Despite
Undetectable Cerebrospinal Fluid HIV RNA in HIV-1-Infected Patients Receiving Antiretroviral
Therapy. JAIDS J. Acquir. Immune Defic. Syndr. *25*, 426.

31. Lorenzo-Redondo, R., Fryer, HelenR., Bedford, T., Kim, E.-Y., Archer, J., Kosakovsky Pond, SergeiL.,
Chung, Y.-S., Penugonda, S., Chipman, JeffreyG., Fletcher, CourtneyV., et al. (2016). Persistent HIV-1
replication maintains the tissue reservoir during therapy. Nature *530*, 51–56.
https://doi.org/10.1038/nature16933.

32. Fauci, A., and Desrosiers, R. (1997). Pathogenesis of HIV and SI.

33. Post, K., Olson, E.D., Naufer, M.N., Gorelick, R.J., Rouzina, I., Williams, M.C., Musier-Forsyth, K., and
Levin, J.G. (2016). Mechanistic differences between HIV-1 and SIV nucleocapsid proteins and cross-
species HIV-1 genomic RNA recognition. Retrovirology *13*, 89. https://doi.org/10.1186/s12977-016-
0322-5.

34. Zhang, J., Diehl, G.E., and Littman, D.R. (2008). Relief of Preintegration Inhibition and
Characterization of Additional Blocks for HIV Replication in Primary Mouse T Cells. PLoS ONE *3*,
e2035. https://doi.org/10.1371/journal.pone.0002035.

35. Greenblatt, M.B., Vbranac, V., Tivey, T., Tsang, K., Tager, A.M., and Aliprantis, A.O. (2012). Graft
versus Host Disease in the Bone Marrow, Liver and Thymus Humanized Mouse Model. PLOS ONE *7*,
e44664. https://doi.org/10.1371/journal.pone.0044664.

36. Shultz, L.D., Ishikawa, F., and Greiner, D.L. (2007). Humanized mice in translational biomedical
research. Nat. Rev. Immunol. *7*, 118–130. https://doi.org/10.1038/nri2017.

37. Wallet, C., De Rovere, M., Van Assche, J., Daouad, F., De Wit, S., Gautier, V., Mallon, P.W.G.,
Marcello, A., Van Lint, C., Rohr, O., et al. (2019). Microglial Cells: The Main HIV-1 Reservoir in the
Brain. Front. Cell. Infect. Microbiol. *9*.

38. Ash, M.K., Al-Harthi, L., and Schneider, J.R. (2021). HIV in the Brain: Identifying Viral Reservoirs and
Addressing the Challenges of an HIV Cure. Vaccines *9*, 867.
https://doi.org/10.3390/vaccines9080867.

39. Churchill, M.J., Gorry, P.R., Cowley, D., Lal, L., Sonza, S., Purcell, D.F.J., Thompson, K.A., Gabuzda, D.,
McArthur, J.C., Pardo, C.A., et al. (2006). Use of laser capture microdissection to detect integrated
HIV-1 DNA in macrophages and astrocytes from autopsy brain tissues. J. Neurovirol. *12*, 146–152.
https://doi.org/10.1080/13550280600748946.

40. Trillo-Pazos, G., Diamanturos, A., Rislove, L., Menza, T., Chao, W., Belem, P., Sadiq, S., Morgello, S.,
Sharer, L., and Volsky, D.J. (2003). Detection of HIV-1 DNA in microglia/macrophages, astrocytes and
neurons isolated from brain tissue with HIV-1 encephalitis by laser capture microdissection. Brain
Pathol. Zurich Switz. *13*, 144–154. https://doi.org/10.1111/j.1750-3639.2003.tb00014.x.

41. Narasipura, S.D., Kim, S., and Al-Harthi, L. (2014). Epigenetic Regulation of HIV-1 Latency in
Astrocytes. J. Virol. *88*, 3031–3038. https://doi.org/10.1128/jvi.03333-13.

42. Narasipura, S.D., Henderson, L.J., Fu, S.W., Chen, L., Kashanchi, F., and Al-Harthi, L. (2012). Role of β-
Catenin and TCF/LEF Family Members in Transcriptional Activity of HIV in Astrocytes. J. Virol. *86*,
1911–1921. https://doi.org/10.1128/JVI.06266-11.

43. Lamers, S.L., Rose, R., Maidji, E., Agsalda-Garcia, M., Nolan, D.J., Fogel, G.B., Salemi, M., Garcia, D.L.,
Bracci, P., Yong, W., et al. (2016). HIV DNA Is Frequently Present within Pathologic Tissues Evaluated
at Autopsy from Combined Antiretroviral Therapy-Treated Patients with Undetectable Viral Loads. J.
Virol. *90*, 8968–8983. https://doi.org/10.1128/JVI.00674-16.

44. Fischer-Smith, T., Tedaldi, E.M., and Rappaport, J. (2008). CD163/CD16 Coexpression by Circulating
Monocytes/Macrophages in HIV: Potential Biomarkers for HIV Infection and AIDS Progression. AIDS
Res. Hum. Retroviruses *24*, 417–421. https://doi.org/10.1089/aid.2007.0193.

45. Koppensteiner, H., Brack-Werner, R., and Schindler, M. (2012). Macrophages and their relevance in
Human Immunodeficiency Virus Type I infection. Retrovirology *9*, 82. https://doi.org/10.1186/1742-
4690-9-82.

46. Lutgen, V., Narasipura, S.D., Barbian, H.J., Richards, M., Wallace, J., Razmpour, R., Buzhdygan, T.,
Ramirez, S.H., Prevedel, L., Eugenin, E.A., et al. (2020). HIV infects astrocytes in vivo and egresses
from the brain to the periphery. PLoS Pathog. *16*, e1008381.
https://doi.org/10.1371/journal.ppat.1008381.

47. Tang, Y., Chaillon, A., Gianella, S., Wong, L.M., Li, D., Simermeyer, T.L., Porrachia, M., Ignacio, C.,
Woodworth, B., Zhong, D., et al. Brain microglia serve as a persistent HIV reservoir despite durable
antiretroviral therapy. J. Clin. Invest. *133*, e167417. https://doi.org/10.1172/JCI167417.

48. Ormel, P.R., Vieira de Sá, R., van Bodegraven, E.J., Karst, H., Harschnitz, O., Sneeboer, M.A.M.,
Johansen, L.E., van Dijk, R.E., Scheefhals, N., Berdenis van Berlekom, A., et al. (2018). Microglia
innately develop within cerebral organoids. Nat. Commun. *9*, 4167. https://doi.org/10.1038/s41467-
018-06684-2.

49. dos Reis, R.S., Sant, S., Keeney, H., Wagner, M.C.E., and Ayyavoo, V. (2020). Modeling HIV-1
neuropathogenesis using three-dimensional human brain organoids (hBORGs) with HIV-1 infected
microglia. Sci. Rep. *10*, 15209. https://doi.org/10.1038/s41598-020-72214-0.

50. Schafer, S.T., Mansour, A.A., Schlachetzki, J.C.M., Pena, M., Ghassemzadeh, S., Mitchell, L., Mar, A.,
Quang, D., Stumpf, S., Ortiz, I.S., et al. (2023). An in vivo neuroimmune organoid model to study
human microglia phenotypes. Cell *186*, 2111-2126.e20. https://doi.org/10.1016/j.cell.2023.04.022.

51. McQuade, A., Coburn, M., Tu, C.H., Hasselmann, J., Davtyan, H., and Blurton-Jones, M. (2018).
Development and validation of a simplified method to generate human microglia from pluripotent
stem cells. Mol. Neurodegener. *13*, 67. https://doi.org/10.1186/s13024-018-0297-x.

52. Albalawi, Y.A., Narasipura, S.D., Olivares, L.J., and Al-Harthi, L. (2022). CD4dim CD8bright T Cells
Home to the Brain and Mediate HIV Neuroinvasion. J. Virol. *96*, e00804-22.
https://doi.org/10.1128/jvi.00804-22.

53. Mamede, J.I., and Hope, T.J. (2016). Detection and Tracking of Dual-Labeled HIV Particles Using
Wide-Field Live Cell Imaging to Follow Viral Core Integrity. In HIV Protocols Methods in Molecular
Biology. (Humana Press, New York, NY), pp. 49–59. https://doi.org/10.1007/978-1-4939-3046-3_4.

54. Mamede, J.I., Cianci, G.C., Anderson, M.R., and Hope, T.J. (2017). Early cytoplasmic uncoating is
associated with infectivity of HIV-1. Proc. Natl. Acad. Sci. U. S. A. *114*, E7169–E7178.
https://doi.org/10.1073/pnas.1706245114.

55. Tan, W.C.C., Nerurkar, S.N., Cai, H.Y., Ng, H.H.M., Wu, D., Wee, Y.T.F., Lim, J.C.T., Yeong, J., and Lim,
T.K.H. (2020). Overview of multiplex immunohistochemistry/immunofluorescence techniques in the
era of cancer immunotherapy. Cancer Commun. *40*, 135–153. https://doi.org/10.1002/cac2.12023.

56. Carter, S.D., Mamede, J.I., Hope, T.J., and Jensen, G.J. (2020). Correlated cryogenic fluorescence
microscopy and electron cryo-tomography shows that exogenous TRIM5α can form hexagonal
lattices or autophagy aggregates in vivo. Proc. Natl. Acad. Sci. *117*, 29702–29711.
https://doi.org/10.1073/pnas.1920323117.

57. Yoh, S.M., Mamede, J.I., Lau, D., Ahn, N., Sánchez-Aparicio, M.T., Temple, J., Tuckwell, A., Fuchs,
N.V., Cianci, G.C., Riva, L., et al. (2022). Recognition of HIV-1 capsid by PQBP1 licenses an innate
immune sensing of nascent HIV-1 DNA. Mol. Cell, S1097-2765(22)00555-X.
https://doi.org/10.1016/j.molcel.2022.06.010.

58. Abud, E.M., Ramirez, R.N., Martinez, E.S., Healy, L.M., Nguyen, C.H.H., Newman, S.A., Yeromin, A.V.,
Scarfone, V.M., Marsh, S.E., Fimbres, C., et al. (2017). iPSC-Derived Human Microglia-like Cells to
Study Neurological Diseases. Neuron *94*, 278-293.e9.
https://doi.org/10.1016/j.neuron.2017.03.042.

59. Jurga, A.M., Paleczna, M., and Kuter, K.Z. (2020). Overview of General and Discriminating Markers of
Differential Microglia Phenotypes. Front. Cell. Neurosci. *14*.

60. Greter, M., and Merad, M. (2013). Regulation of microglia development and homeostasis. Glia *61*,
121–127. https://doi.org/10.1002/glia.22408.

61. Ford, A.L., Goodsall, A.L., Hickey, W.F., and Sedgwick, J.D. (1995). Normal adult ramified microglia
separated from other central nervous system macrophages by flow cytometric sorting. Phenotypic
differences defined and direct ex vivo antigen presentation to myelin basic protein-reactive CD4+ T
cells compared. J. Immunol. Baltim. Md 1950 *154*, 4309–4321.

62. Hernández, D., Rooney, L.A., Daniszewski, M., Gulluyan, L., Liang, H.H., Cook, A.L., Hewitt, A.W., and
Pébay, A. (2022). Culture Variabilities of Human iPSC-Derived Cerebral Organoids Are a Major Issue
for the Modelling of Phenotypes Observed in Alzheimer's Disease. Stem Cell Rev. Rep. *18*, 718–731.
https://doi.org/10.1007/s12015-021-10147-5.

63. Xu, R., Boreland, A.J., Li, X., Erickson, C., Jin, M., Atkins, C., Pang, Z.P., Daniels, B.P., and Jiang, P.
(2021). Developing human pluripotent stem cell-based cerebral organoids with a controllable
microglia ratio for modeling brain development and pathology. Stem Cell Rep. *16*, 1923–1937.
https://doi.org/10.1016/j.stemcr.2021.06.011.

64. Brown, R.M., Rana, P.S.J.B., Jaeger, H.K., O'Dowd, J.M., Balemba, O.B., and Fortunato, E.A. (2019).
Human Cytomegalovirus Compromises Development of Cerebral Organoids. J. Virol. *93*, e00957-19.
https://doi.org/10.1128/JVI.00957-19.

65. Hübner, W., Chen, P., Portillo, A.D., Liu, Y., Gordon, R.E., and Chen, B.K. (2007). Sequence of Human
Immunodeficiency Virus Type 1 (HIV-1) Gag Localization and Oligomerization Monitored with Live
Confocal Imaging of a Replication-Competent, Fluorescently Tagged HIV-1. J. Virol. *81*, 12596–
12607. https://doi.org/10.1128/JVI.01088-07.

66. Dang, J., Tiwari, S.K., Lichinchi, G., Qin, Y., Patil, V.S., Eroshkin, A.M., and Rana, T.M. (2016). Zika
Virus Depletes Neural Progenitors in Human Cerebral Organoids through Activation of the Innate
Immune Receptor TLR3. Cell Stem Cell *19*, 258–265. https://doi.org/10.1016/j.stem.2016.04.014.

67. Watanabe, M., Buth, J.E., Vishlaghi, N., Torre-Ubieta, L. de la, Taxidis, J., Khakh, B.S., Coppola, G.,
Pearson, C.A., Yamauchi, K., Gong, D., et al. (2017). Self-Organized Cerebral Organoids with Human-
Specific Features Predict Effective Drugs to Combat Zika Virus Infection. Cell Rep. *21*, 517–532.
https://doi.org/10.1016/j.celrep.2017.09.047.

68. De Rivero Vaccari, J.P., Lotocki, G., Marcillo, A.E., Dietrich, W.D., and Keane, R.W. (2008). A
Molecular Platform in Neurons Regulates Inflammation after Spinal Cord Injury. J. Neurosci. *28*,
3404–3414. https://doi.org/10.1523/JNEUROSCI.0157-08.2008.

69. Espinosa-Oliva, A.M., García-Revilla, J., Alonso-Bellido, I.M., and Burguillos, M.A. (2019). Brainiac
Caspases: Beyond the Wall of Apoptosis. Front. Cell. Neurosci. *13*, 500.
https://doi.org/10.3389/fncel.2019.00500.

70. Lin, W.-P., Xiong, G.-P., Lin, Q., Chen, X.-W., Zhang, L.-Q., Shi, J.-X., Ke, Q.-F., and Lin, J.-H. (2016).
Heme oxygenase-1 promotes neuron survival through down-regulation of neuronal NLRP1

expression after spinal cord injury. J. Neuroinflammation *13*, 52. https://doi.org/10.1186/s12974-
016-0521-y.

71. McInnis, J.J., Sood, D., Guo, L., Dufault, M.R., Garcia, M., Passaro, R., Gao, G., Zhang, B., and Dodge,
J.C. (2023). Unravelling neuronal and glial differences in ceramide composition, synthesis, and
sensitivity to toxicity. Preprint, https://doi.org/10.1101/2023.12.06.569570
https://doi.org/10.1101/2023.12.06.569570.

72. Elkjaer, M.L., Hartebrodt, A., Oubounyt, M., Weber, A., Vitved, L., Reynolds, R., Thomassen, M.,
Rottger, R., Baumbach, J., and Illes, Z. (2024). Single-Cell Multi-Omics Map of Cell Type–Specific
Mechanistic Drivers of Multiple Sclerosis Lesions. Neurol. Neuroimmunol. Neuroinflammation *11*,
e200213. https://doi.org/10.1212/NXI.0000000000200213.

73. Frost, J.L., and Schafer, D.P. (2016). Microglia: Architects of the Developing Nervous System. Trends
Cell Biol. *26*, 587–597. https://doi.org/10.1016/j.tcb.2016.02.006.

74. Herculano-Houzel, S. (2014). The glia/neuron ratio: How it varies uniformly across brain structures
and species and what that means for brain physiology and evolution. Glia *62*, 1377–1391.
https://doi.org/10.1002/glia.22683.

75. Hong, S., Beja-Glasser, V.F., Nfonoyim, B.M., Frouin, A., Li, S., Ramakrishnan, S., Merry, K.M., Shi, Q.,
Rosenthal, A., Barres, B.A., et al. (2016). Complement and microglia mediate early synapse loss in
Alzheimer mouse models. Science *352*, 712–716. https://doi.org/10.1126/science.aad8373.

76. Markarian, M., Krattli, R.P., Baddour, J.D., Alikhani, L., Giedzinski, E., Usmani, M.T., Agrawal, A.,
Baulch, J.E., Tenner, A.J., and Acharya, M.M. (2021). Glia-Selective Deletion of Complement C1q
Prevents Radiation-Induced Cognitive Deficits and Neuroinflammation. Cancer Res. *81*, 1732–1744.
https://doi.org/10.1158/0008-5472.CAN-20-2565.

77. Vasek, M.J., Garber, C., Dorsey, D., Durrant, D.M., Bollman, B., Soung, A., Yu, J., Perez-Torres, C.,
Frouin, A., Wilton, D.K., et al. (2016). A complement-microglial axis drives synapse loss during virus-
induced memory impairment. Nature *534*, 538–543. https://doi.org/10.1038/nature18283.

78. Gavriilaki, M., Kimiskidis, V.K., and Gavriilaki, E. (2020). Precision Medicine in Neurology: The
Inspirational Paradigm of Complement Therapeutics. Pharmaceuticals *13*, 341.
https://doi.org/10.3390/ph13110341.

79. Garred, P., Tenner, A.J., and Mollnes, T.E. (2021). Therapeutic Targeting of the Complement System:
From Rare Diseases to Pandemics. Pharmacol. Rev. *73*, 792–827.
https://doi.org/10.1124/pharmrev.120.000072.

80. Mastellos, D.C., Ricklin, D., and Lambris, J.D. (2019). Clinical promise of next-generation
complement therapeutics. Nat. Rev. Drug Discov. *18*, 707–729. https://doi.org/10.1038/s41573-
019-0031-6.

81. Zelek, W.M., Xie, L., Morgan, B.P., and Harris, C.L. (2019). Compendium of current complement
therapeutics. Mol. Immunol. *114*, 341–352. https://doi.org/10.1016/j.molimm.2019.07.030.

82. dos Reis, R.S., Sant, S., Keeney, H., Wagner, M.C.E., and Ayyavoo, V. (2020). Modeling HIV-1
neuropathogenesis using three-dimensional human brain organoids (hBORGs) with HIV-1 infected
microglia. Sci. Rep. *10*, 15209. https://doi.org/10.1038/s41598-020-72214-0.

83. Cenker, J.J., Stultz, R.D., and McDonald, D. (2017). Brain Microglial Cells Are Highly Susceptible to
HIV-1 Infection and Spread. AIDS Res. Hum. Retroviruses *33*, 1155–1165.
https://doi.org/10.1089/aid.2017.0004.

84. Pollack, R.A., Jones, R.B., Pertea, M., Bruner, K.M., Martin, A.R., Thomas, A.S., Capoferri, A.A., Beg,
S.A., Huang, S.-H., Karandish, S., et al. (2017). Defective HIV-1 proviruses are expressed and can be
recognized by cytotoxic T lymphocytes which shapes the proviral landscape. Cell Host Microbe *21*,
494-506.e4. https://doi.org/10.1016/j.chom.2017.03.008.

85. Klatt, N.R., Chomont, N., Douek, D.C., and Deeks, S.G. (2013). Immune activation and HIV
persistence: Implications for curative approaches to HIV infection. Immunol. Rev. *254*, 326–342.
https://doi.org/10.1111/imr.12065.

86. Imamichi, H., Dewar, R.L., Adelsberger, J.W., Rehm, C.A., O'Doherty, U., Paxinos, E.E., Fauci, A.S.,
and Lane, H.C. (2016). Defective HIV-1 proviruses produce novel protein-coding RNA species in HIV-
infected patients on combination antiretroviral therapy. Proc. Natl. Acad. Sci. U. S. A. *113*, 8783–
8788. https://doi.org/10.1073/pnas.1609057113.

87. Mohammadzadeh, N., Roda, W., Branton, W.G., Clain, J., Rabezanahary, H., Zghidi-Abouzid, O.,
Gelman, B.B., Angel, J.B., Cohen, E.A., Gill, M.J., et al. (2021). Lentiviral Infections Persist in Brain
despite Effective Antiretroviral Therapy and Neuroimmune Activation. mBio *12*, e0278421.
https://doi.org/10.1128/mBio.02784-21.

88. Yoh, S.M., Mamede, J.I., Lau, D., Ahn, N., Sánchez-Aparicio, M.T., Temple, J., Tuckwell, A., Fuchs,
N.V., Cianci, G.C., Riva, L., et al. (2022). Recognition of HIV-1 capsid by PQBP1 licenses an innate
immune sensing of nascent HIV-1 DNA. Mol. Cell *82*, 2871-2884.e6.
https://doi.org/10.1016/j.molcel.2022.06.010.

89. Gao, D., Wu, J., Wu, Y.-T., Du, F., Aroh, C., Yan, N., Sun, L., and Chen, Z.J. (2013). Cyclic GMP-AMP
Synthase Is an Innate Immune Sensor of HIV and Other Retroviruses. Science *341*, 903–906.
https://doi.org/10.1126/science.1240933.

90. Wang, P.-H., Fung, S.-Y., Gao, W.-W., Deng, J.-J., Cheng, Y., Chaudhary, V., Yuen, K.-S., Ho, T.-H.,
Chan, C.-P., Zhang, Y., et al. (2018). A novel transcript isoform of STING that sequesters cGAMP and
dominantly inhibits innate nucleic acid sensing. Nucleic Acids Res. *46*, 4054–4071.
https://doi.org/10.1093/nar/gky186.

1062

A)

iPSC & HPC culture

CO generation iPSC single cells

iPSC+HPC single cells CO-iM generation

0d EB formation — 5d Induction stage — 7d Expansion stage — 10d ---- Maturation ---- 50+d Assay → Days

B)

	Day 2	Day 5	Day 8	Day 60
iPSC single cells → CO				
iPSC +HPC single cells → CO-iM				

C)

- CO
- CO-iM

Diameter (µm) vs Time (days): 1, 2, 5, 7, 15, 60

D)

Hoechst
Neurons (Tuj1)
Astrocytes (GFAP)
Microglia (TMEM119)

100µm

Hoechst | TMEM119-AF647 | GFAP-AF488 | Iba1-AF555 | Tuj1-AF647
MAP2-PE | NeuN-AF647 | ALD1-Cy7 | S100β-FITC | Mix all

E)

i
Hoechst
Neurons (Tuj1)
Astrocytes (GFAP)
Microglia (TMEM119)
100µm

F)

ii
Hoechst
Neurons (Tuj1)
Astrocytes (GFAP)
Microglia (TMEM119)
100µm