Multi-dimensionality of tree communities structure host-parasitoid networks and their

phylogenetic composition

Ming-Qiang Wang[1,2#], Shi-Kun Guo[1,3#], Peng-Fei Guo[4#], Juan-Juan Yang[1,3], Guo-Ai Chen[1,3],

Douglas Chesters[1,5], Michael Orr[1,6], Ze-Qing Niu[1], Michael Staab[7], Jing-Ting Chen[1,3], Yi Li[10],

Qing-Song Zhou[1,5], Felix Fornoff[8], Xiaoyu Shi[1], Shan Li[10], Massimo Martini[8], Alexandra-Maria

Klein[8], Andreas Schuldt[9], Xiaojuan Liu[10], Keping Ma[5,10], Helge Bruelheide[11], Arong Luo[1,5,13]*,

Chao-Dong Zhu[1,3,5,12,13]*

1 CAS Key Laboratory of Zoological Systematics and Evolution, Institute of Zoology, Chinese

Academy of Sciences, 1 Beichen West Road, Chaoyang District, Beijing, 100101, China

2 CAS Key Laboratory of Mountain Ecological Restoration and Bioresource Utilization &

Biodiversity Conservation Key Laboratory of Sichuan Province, Chengdu Institute of Biology,

Chinese Academy of Sciences, 4 Renmin South Road, Wuhou District, Chengdu, 610041, China

3 College of Biological Sciences, University of Chinese Academy of Sciences, No.19A Yuquan

Road, Shijingshan District, Beijing, 100049, China

4 College of Pharmacy, Guizhou University of Traditional Chinese Medicine, Guiyang 550025,

China

5 International College, University of Chinese Academy of Sciences, No.19A Yuquan Road,

Shijingshan District, Beijing, 100049, China

6 Entomologie, Staatliches Museum für Naturkunde Stuttgart, Stuttgart, Germany.

7 Ecological Networks, Technical University Darmstadt, Schnittspahnstraße 3, 64287 Darmstadt,

Germany

8 Department of Nature Conservation and Landscape Ecology, Albert-Ludwigs-University

Freiburg, Tennenbacher Straße 4, 79106 Freiburg, Germany

9 Forest Nature Conservation, University of Göttingen, Büsgenweg 3, 37077, Göttingen,

Germany

10 State Key Laboratory of Vegetation and Environmental Change, Institute of Botany,

Chinese Academy of Sciences, 20 Nanxincun, Xiangshan, Haidian District, Beijing 100093,

China

11 Institute of Biology/Geobotany and Botanical Garden, Martin Luther University

Halle-Wittenberg, Große Steinstraße 79/80, 06108 Halle (Saale), Germany

12 State Key Laboratory of Integrated Pest Management, Institute of Zoology, Chinese Academy

of Sciences, Beijing, 100101, China

13 Key Laboratory of Animal Biodiversity Conservation and Integrated Pest Management

(Chinese Academy of Sciences), Beijing, 100101, China

[#]Ming-Qiang Wang, Shi-Kun Guo and Peng-Fei Guo equally contributed to this manuscript.

*Correspondence author: luoar@ioz.ac.cn; zhucd@ioz.ac.cn, Phone: +86-10-64807085, Fax:

+86-10-64807099

Data availability

Data is available on Science Data Bank at https://www.scidb.cn/en/s/Ujq22u, and will be available

on the BEF-China project database at https://data.botanik.uni-halle.de/bef-china/datasets.

Conflict of interest statement

The authors have no conflict of interest.

Author contributions

C.D.Z., A.L. and M.Q.W. conceived the idea for the manuscript; C.D.Z., A.M.K., S.K.G., P.F.G.

and M.Q.W. designed the study. S.K.G., P.F.G., J.J.Y., G.A.C., Z.Q.N., M.S., J.T.C., Y.L., Q.S.Z.,

F.F., X.S., S.L., M.M., A.M.K., A.S., X.L., K.M. and H.B. collected and/or contributed data; D.C.,

M.Q.W. and S.K.G. conducted the phylogenetic analysis; M.Q.W. S.KG. and P.F.G. conducted

the statistical analyses and wrote the manuscript, with input from F.F., A.S., M.S. and M.M and all

coauthors.

Abstract

Environmental factors can influence ecological networks, but these effects are poorly understood

in the realm of the phylogeny of host-parasitoid interactions. Especially, we lack a comprehensive

understanding of the ways that biotic factors, including plant species richness, overall community

phylogenetic and functional composition of consumers, and abiotic factors such as microclimate,

determining host–parasitoid network structure and host–parasitoid community dynamics. To

address this, we leveraged a five-year dataset of trap-nesting bees and wasps and their parasitoids

collected in a highly-controlled, large-scale subtropical tree biodiversity experiment. We tested for

effects of tree species richness, tree phylogenetic and functional diversity, and species and

phylogenetic composition on species and phylogenetic diversity of both host and parasitoid

communities and the composition of their interaction networks. We show that multiple

components of tree diversity and canopy cover impacted both, species and phylogenetic

composition of hosts and parasitoids. Generally, phylogenetic associations between hosts and

parasitoids reflected non-randomly structured interactions between phylogenetic trees of hosts and

parasitoids. Further, host-parasitoid network structure was influenced by tree species richness, tree

phylogenetic diversity, and canopy cover. Our study indicates that the composition of higher

trophic levels and corresponding interaction networks are determined by plant diversity and

canopy cover especially via trophic links in species-rich ecosystems.

Introduction

Understanding the ecological consequences of biodiversity loss is an increasingly important

task in ecology, given the ongoing biodiversity crisis (Isbell et al. 2022). Representing the

interdependencies among organisms, ecological networks reflect whether and how species

interact with each other across trophic levels, playing an indispensable role in assessing

ecosystem stability and integrity (De Ruiter et al. 1995, Harvey et al. 2017). Changes in

network structure of higher trophic levels usually coincide with variations in their diversity

and community composition, which could be in turn affected by the changes in producers via

trophic cascades (Barnes et al. 2018, Gonzalez et al. 2020). However, we still lack a

generalizable framework for how these networks and especially their phylogenetic

interdependencies respond to biodiversity loss in ecosystems, such as changes in the tree

diversity of forests (Tylianakis et al. 2008, Grossman et al. 2018) . To better understand

species coexistence and its role for biodiversity conservation, we must further study the

mechanisms on dynamics of networks from multiple perspectives (Tittensor et al. 2014,

Brondizio et al. 2019). Interactions can be viewed from a top-down or bottom-up perspective.

Previous studies have shown there might be asymmetric effects across trophic levels (Vidal

and Murphy 2018), with both factors shaping multitrophic communities together(Hunter et al.

1992). Moreover, the diversity and community of higher trophic levels could be also driven

by microclimate, which could be determined by plant structuring, such as canopy cover

(Fornoff et al. 2021, Perlík et al. 2023). Understanding how canopy cover responds to

changing biodiversity is therefore imperative to predict how ongoing environmental changes

will impact the functioning and stability of ecosystems (Hines et al. 2019).

A network that unites bottom-up and top-down processes in many ecosystems and is

prone to strong alterations due to environmental change is the interaction network between

parasitoids and their hosts (Tylianakis et al. 2006, Jeffs and Lewis 2013). Insect parasitoids

attack and feed on and eventually kill their insect hosts (Godfray and Godfray 1994).

Parasitoids are thought to be particularly sensitive to environmental changes, because species

in higher trophic levels usually have smaller population sizes and their host fluctuations may

cascade up to impact the parasitoids' presence or absence. Therefore, insect host-parasitoid

systems are ideal for studying the relationships between community-level changes and species

interactions (Jeffs and Lewis 2013). Previous studies mainly focused on the influence of

abiotic factors on host-parasitoid interactions, such as elevation and habitat structure (e.g.

Valladares et al. 2012, Maunsell et al. 2015, Grass et al. 2018), or sometimes interaction

structure (e.g. Cagnolo et al. 2011). However, the role of multiple components of plant

diversity (i.e. taxonomic, functional and phylogenetic diversity) in modifying host-parasitoid

interaction networks, as key biotic determinants of overall ecosystem structure, remains

relatively little explored (but see Staab et al. 2016). Recent studies mainly focused on basic

diversity associations between host and parasitoids (Ebeling et al. 2012, Schuldt et al. 2019,

Guo et al. 2021). These studies have demonstrated both direct and indirect effects (i.e. one

pathway and more pathways via other variables) of plant diversity on both hosts and

parasitoids diversity, possibly via increased niche space and resource availability (Guo et al.

2021). Nevertheless, how these patterns propagate to their interaction networks is still unclear.

Moreover, the effects of changing plant diversity are not always obvious when looking at

plant species richness. The other diversity components (e.g. plant phylogenetic diversity)

have been shown to better predict diversity-dependent bottom-up effects on host-parasitoid

networks (e.g. Staab et al. 2016). It is especially important to take phylogenetic dependences

(e.g. phylogenetic diversity or phylogenetic congruence) into account when it comes to the

phylogenetic structure within and across trophic levels (Webb et al. 2002, Emerson and

Gillespie 2008). This makes it vital to account for multiple dimensions of biodiversity (e.g.

taxonomic, phylogenetic, functional) and relevant trophic interactions (Peralta et al. 2015,

Volf et al. 2017, Wang et al. 2020). These components might jointly affect host-parasitoid

networks in a system with high species diversity. Forests have garnered special attention

lately, because they represent complex and large ecosystems susceptible to global change (De

Frenne et al. 2021, Popkin 2021). Understanding how multiple dimensions of biodiversity

modulate the effects of tree diversity loss on the structure and interaction strength of

host-parasitoid networks clearly requires further study (Staab et al. 2016, Fornoff et al. 2019).

Here, we leverage standardized trap nests for solitary cavity-nesting bees, wasps, and

their parasitoids in a large-scale subtropical forest biodiversity experiment to test how

multiple dimensions of tree diversity and community composition influence host-parasitoid

network structure. A multi-faceted approach is particularly important when considering that

associations between trophic levels might be non-random and phylogenetically structured

(Volf et al. 2018, Wang et al. 2020). We aimed to discern the primary components of the

diversity and composition of tree communities that affect higher trophic level interactions via

quantifying the strength and complexity of associations between hosts and parasitoid. We

expected that (a) multiple tree community metrics, such as species, phylogenetic and

functional diversity and species community composition can structure host and parasitoid

community compositions, especially via phylogenetic processes (e.g. lineages of trophic

levels diverge and evolve over time), as species interactions often show phylogenetic

conservatism (e.g. Pellissier et al. 2013, Peralta et al. 2015). Further, we hypothesized that (b)

host-parasitoid networks will be more complex and stable with increasing tree species

richness due to potential links from higher richness of hosts and parasitoids promoted by

more tree species, and (c) both, community and interaction network changes can also be

related to abiotic factors, such as canopy cover, which might play a role in structuring

hymenopteran communities (Haddad et al. 2011, Fornoff et al. 2021). By better understanding

these dynamics, we can begin building a generalized framework for understanding

host-parasitoid interactions in forest ecosystems.

Results

Overall, 34,398 brood cells were collected from 13,267 tubes across five years of sampling

(2015, 2016, 2018, 2019, and 2020). Six families of hosts and seventeen families of

parasitoids were identified. Among them, we found 56 host species (12 bees and 44 wasps,

mean abundance and richness are 400 and 45, respectively, for each plot) and 50 parasitoid

species (38 Hymenoptera and 12 Diptera, mean abundance and richness are 14 and 9,

respectively, for each plot). The full species list and their abundances are given in Table S1.

Overall, our sampling was adequate for analysis (especially for hosts), as confirmed by the

sampling completness evaluation (Fig. S1).

Community composition of hosts and parasitoids

Host species composition was significantly related to the species and phylogenetic

composition of the trees and parasitoid communities (NMDS axis scores; i.e. preserving the

rank order of pairwise dissimilarities between samples), as well as to canopy cover, tree mean

phylogenetic distance (MPD), elevation, and eastness (sine-transformed radian values of

aspect) (Fig. 1a, Table 1, Table S2). Parasitoid species composition was significantly

associated with host phylogenetic diversity, tree functional diversity, tree MPD, eastness, and

elevation, and was significantly related to tree species composition, host species composition,

and canopy cover (Fig. 1b, Table 1, Table S3). Host phylogenetic composition was affected

by tree species composition, tree MPD, tree functional diversity, canopy cover, eastness,

elevation and was especially affected by parasitoid species and phylogenetic composition (Fig.

1c, Table 1, Table S4). For parasitoid phylogenetic composition, significant relationships

were found with tree species and phylogenetic composition, host species composition, tree

functional diversity, canopy cover, elevation, and eastness (Fig. 1d, Table 1, Table S5). The

PERMANOVA analysis also highlighted the important role of canopy cover for host and

parasitoid community (Table S6-9). The Mantel test revealed a consistent pattern with the

NMDS analysis, highlighting a pronounced relationship between the species composition of

hosts and parasitoids (Table S10). However, the correlation between the phylogenetic

composition of hosts and parasitoids was not significant.

An effect of host composition on the composition of the parasitoid communities was

further indicated by a significant parafit test (p = 0.032) for testing the hypothesis of

coevolution between a clade of hosts and a clade of parasites, suggesting nonrandom

associations in the phylogenetic structure of parasitoid and host communities (Fig. 2, Fig. S4).

Host-parasitoid network associations

The linear regression model results showed that host vulnerability, linkage density, and

robustness of parasitoids were significantly negatively related to tree species richness, while

remaining unaffected by other environmental covariables (Table 2, Fig. 3; except for

elevation, which was marginally significantly related to robustness). Interaction evenness was

significantly negatively associated with canopy cover, and interaction evenness was also

negatively related to eastness (Fig. 4c; Table 2). Parasitoid generality was only marginally

associated with canopy cover, and was not related to tree species richness or the other

environmental variables. In the alternative models (tree species richness replaced by tree

MPD), host vulnerability and linkage density were significantly positively related to tree

MPD (Fig. 4a, 4b; Table S11), while robustness of parasitoids was negatively related to tree

MPD (Fig. S5, Table S11). The results of other network metrics (parasitoid generality and

interaction evenness) were consistent with those of the primary models. Tree mean nearest

taxon distance (MNTD) was unrelated to any network indices.

The results of the null model analysis suggested that our metrics calculated by the

observed network were significantly different from a random distribution (72, 71, and 77 out

of 85 values for parasitoid generality, host vulnerability, and linkage density, respectively; all

values for robustness, and interaction evenness), strongly demonstrating that interactions

between species were not driven by random processes.

Discussion

Our study demonstrates that tree species richness and phylogenetic diversity play key roles in

modulating interacting communities of hosts and parasitoids. These interactions are further

structured by the phylogenetic associations between hosts and parasitoids. Moreover, canopy

cover partly determined host-parasitoid network indices patterns, including host vulnerability,

linkage density, and interaction evenness. This result supports a recent finding that the

structure of host-parasitoid networks is also mediated by changes in microclimate, which is

directly related to canopy cover, to some extent (Fornoff et al. 2021). These patterns were

highly associated with multiple tree diversity metrics (tree species, phylogenetic and

functional diversity), and compositional changes which are key to understand how

host-parasitoid interactions may be impacted by biodiversity loss of lower trophic levels in

food webs through trait- and phylogeny-based processes.

Community composition associations

Host species composition was influenced by several factors, including the species and

phylogenetic composition of trees and parasitoids, tree diversity (species richness and MPD),

and other biotic factors such as elevation and slope (Fig. 1a; Table 1). The effects of tree

diversity and composition on host species composition agree with previous studies where

solitary bee and wasp species composition were related to plant community structure (e.g.

Loyola and Martins 2008). It seems likely that these results are based on bee linkages to

pollen resources and predatory wasp linkages to the diverse of food sources, which may

themselves be closely linked to resource heterogeneity increasing with tree species richness

(Reitalu et al. 2019, Staab and Schuldt 2020).

We also found that tree MPD, FD, and species composition affect parasitoid species

composition (Fig. 1b; Table 1), as many studies have found significant relationships between

plant and parasitoid diversity in forests, including tree phylogenetic diversity (Staab et al.

2016), functional diversity (Rodriguez et al. 2019), and structural diversity (Schuldt et al.

2019). Similar to predators, parasitoids might also be more active or efficient with increasing

tree community dissimilarity due to higher prey resources or lower intraguild parasitism

caused by more diverse habitats (Finke and Denno 2002). On the other hand, our results also

show that host species composition and parasitoid species composition relate to each other

and their phylogenetic compositions, which are structured by tree communities to some extent.

This pattern could propagate to the adjacent two higher trophic interactions through both

top-down and bottom-up control.

Both host and parasitoid phylogenetic composition were related to tree species composition

(Fig. 1c, d; Table 1). This pattern has important implications for cascading effects among

trophic levels, in that producer communities could structure a higher trophic level community

via an intermediate trophic level. However, while both host and parasitoid phylogenetic

composition was related to tree MPD, only parasitoids responded to tree phylogenetic

composition. This may be because there are many caterpillar-hunting wasps in our host

communities, and the community composition of caterpillars were usually correlated with tree

phylogenetic communities (Wang et al. 2019). Therefore, the prey organisms highly

associated with tree phylogenetic composition (e.g. caterpillars) might indirectly determine

predatory wasp (host) phylogenetic composition, as recently found for interactions between

plants, caterpillars and spiders (Chen et al. 2023). This could be further tested by analyzing

the food directly used by the wasps (caterpillars). For parasitoids, tree phylogenetic

composition might drive the process of community assembly through trophic cascades (e.g.

from plants to parasitoids via herbivores and host wasps) (Webb et al. 2002, Cavender-Bares

et al. 2009). Additionally, parasitoid phylogenetic composition can be indirectly influenced

by tree structural diversity (e.g. host availability in plots with higher heterogeneity; Schuldt et

al. 2019), which can be determined by conserved traits across phylogenies (Webb et al. 2002).

The phylogenetic associations between hosts and parasitoids exhibited a nonrandom structure

(significant *parafit* correlation; Fig. 2) between the phylogenetic trees of the host and their

parasitoids (see also Peralta et al. 2015). Such pattern could be further confirmed by the

significant association between host phylogenetic composition and parasitoid phylogenetic

composition (Fig. 1c), which suggested that their interactions are phylogenetically structured

to some extent. However, this significant pattern was observed only in the NMDS analysis

and not in the Mantel test, suggesting that the non-random interactions between hosts and

parasitoids could not be simply predicted by their community similarity and associations

between the phylogenetic composition of hosts and parasitoids are more complex and require

further investigation in the future.

Moreover, the species and phylogenetic composition of hosts and parasitoids was also

related to abiotic factors, especially to canopy cover, which has been considered especially

important to microclimate (Sobek et al. 2009, Fornoff et al. 2021). In future studies, it will be

useful to incorporate other, more direct metrics of microclimate, such as local temperature

and humidity, to determine the proximal drivers of these microclimatic effects (Ma et al. 2010,

Fornoff et al. 2021).

Community-level host-parasitoid networks

Tree species richness did not significantly influence the diversity of hosts targeted by

parasitoids (parasitoid generality), but caused a significant increase in the diversity of

parasitoids per host species (host vulnerability) (Fig. 3a; Table 2). This is likely because niche

differentiation often influences network specialization via potential higher resource diversity

in plots with higher tree diversity (Lopez-Carretero et al. 2014). Such positive relationship

between host vulnerability and tree species richness suggested that host-parasitoid

interactions could be driven through bottom-up effects via benefit from tree diversity. For

example, parasitoid species increases more than host diversity with increasing tree species

richness (Guo et al. 2021), resulting in an increase of host vulnerability at community level.

According to the enemies hypothesis (Root 1973), which posits a positive effects of plant

richness on natural enemies, the higher trophic levels in our study (e.g. predators and

parasitoids) would benefit from tree diversity and regulate herbivores thereby (Staab and

Schuldt 2020). Indeed, previous studies at the same site found that bee parasitoid richness and

abundance were positively related to tree species richness, but not their bee hosts (Fornoff et

al. 2021, Guo et al. 2021). Because our dataset considered all hosts and reflects an overall

pattern of host-parasitoid interactions, the effects of tree species richness on parasitoid

generality might be more complex and difficult to predict, as we found that neither tree

species richness nor tree MPD were related to parasitoid generality.

Linkage density was positively related to tree species richness (Fig. 3b; Table 2), supporting

the food web theory, which predicts in our case that network complexity (linkage density)

depends on the number of plants (Blüthgen and Staab 2024). Although trees were not directly

included as a trophic level in our networks, potential network complexity increased with tree

species richness, likely enabling higher network stability/resistance (Ebeling et al. 2011,

Staab et al. 2015). For example, a network might be more sensitive to extinctions because of

key species loss due to lower linkage density and lower redundancy (Blüthgen and Staab

2024). However, parasitoid robustness was unexpectedly negatively related to tree species

richness (Fig. 3c; Table 2). It was expected that higher trophic levels would be more robust,

less influenced by perturbations from lower trophic levels, when plant diversity is higher, as

more potential interactions at lower trophic levels should theoretically increase redundancy

and resilience of connected higher levels (Blüthgen and Klein 2011, Fornoff et al. 2019).

Dilution effects may explain this, as plots with higher richness held fewer individuals of a

given tree species. If there are strong prey item (caterpillars, grasshoppers, etc.) preferences

for one species, there may be fewer individuals per area or they may be more densely

aggregated and less likely to be encountered by parasitoids. This increased stochasticity in

parasitoid wasps could benefit hosts by reducing parasitism pressure overall, weakening

top-down controls.

Similar to tree species richness, tree MPD was also positively correlated with host

vulnerability and linkage density (Fig. 4a, b), meaning that the mean number of parasitoids

per host species and number of links within the host-parasitoid system can also be promoted

by tree MPD. This is in agreement with several recent studies showing that plant phylogenetic

diversity not only affects herbivores but also higher tropic levels (Pellissier et al. 2013, Staab

et al. 2020, Wang et al. 2020). Our results suggest that the specialization and complexity of

higher trophic levels can also be affected by plant phylogenetic diversity. This pattern can be

traced to the effects of habitat heterogeneity caused by tree species richness and MPD on

higher trophic levels via bottom-up control. The effects of tree MPD were consistent with

effects of tree species richness on robustness of parasitoids to host loss. This result suggests

that higher trophic levels are sensitive to changes in both plant phylogenetic relatedness and

general species dissimilarity via trophic interactions, even the hosts are not all directly

interacting with plants, bees excluded. Therefore, it may be that stronger linkages would be

found when exclusively exploring such plant-herbivore-parasitoid systems.

Interaction evenness was significantly negatively related to canopy cover(Fig. 4c),

further reinforcing the important role of canopy-cover modulated microclimate (likely

temperature and humidity) for trophic interactions (Sobek et al. 2009, Fornoff et al. 2021).

Our results agree with a previous study on ants, where plant-insect interactions were more

even with more open canopies (Dáttilo and Dyer 2014). In our case, canopy cover might

change hymenopteran species evenness and then further influence interaction evenness.

Certain host species tended to nest in plots with higher canopy cover, which might decrease

the interaction evenness by favoring parasitoids of fewer, more dominant hosts. This pattern

would become more significant when more host and parasitoid species are in a plot, given the

positive relationship between higher trophic level diversity and canopy cover.

Future prospects

Overall, our study enables new insights into the dynamics of host-parasitoid interactions

under varying environmental conditions, an important step toward building a synthetic model

for such biodiversity. A key finding was that although parasitoids and hosts respond to tree

species richness, top-down control seems predominant in parasite-host interaction, though

whether this holds for others antagonistic interactions requires further investigation. Different

trophic levels and functional groups of species responded differently to experimental changes

in plant communities (Fornoff et al. 2021, Guo et al. 2021). This highlights the complexities

of building multi-trophic networks and calls for more studies across habitat types and taxa, to

test the generality of our findings. Future studies should also consider the role of

host/parasitoid functional traits, because they might play a critical role in modifying network

structures and ecosystem functioning.

Materials and methods

Study sites design

This study was conducted in the BEF-China biodiversity experiment, which is the largest tree

diversity experiment worldwide. The experiment is located in a subtropical forest near

Xingangshan, Jiangxi province, south-east China (29° 08$'$ −29° 11$'$ N, 117° 90$'$ −117°

93$'$ E). The mean annual temperature is 16.7° C and mean annual precipitation 1821 mm

(Yang et al. 2013). The experiment includes two study sites (Site A and Site B), 4 km apart

from each other, that were established in 2009 (Site A) and 2010 (Site B) respectively. A total

of 566 plots (25.8 $\times$ 25.8 m) were designed on the two sites, and per plot 400 trees were

initially planted in 20 rows and 20 columns with a planting distance of 1.29 m. A tree species

richness gradient (1, 2, 4, 8, 16 and 24 species) was established at each site, based on a

species pool of 40 local, broadleaved tree species (Bruelheide et al. 2014). The tree species

pools of the two plots are nonoverlapping (16 species for each site). The composition of tree

species within the study plots is based on a "broken-stick" design (see Bruelheide et al. 2014).

For our study, at both sites (site A and site B) eight plots of each tree species richness

level (1, 2, 4, 8) were randomly selected, as well as six and two plots of 16 and 24 mixtures.

In addition, at site B eight additional monocultures were sampled (Fornoff et al. 2021),

resulting in 48 plots in Site B (including 16 monocultures, eight plots for each 2, 4, 8 mixtures

and six and two plots of 16 and 24 mixtures. In total, 88 study plots were used (40 plots on

Site A and 48 plots on Site B, see Fig. S2).

Sampling

We collected trap nests monthly to sample solitary bees and wasps (Staab et al. 2018) in the

88 plots from September to November in 2015 and April to November in 2016, 2018, 2019

and 2020. For each plot, we installed two poles with trap nests (11 m apart from each other

and 9 m away from the nearest adjacent plots) along a SW–NE diagonal (following the design

of Ebeling et al. 2012). Each pole stood 1.5 m above ground, and each trap nest consisted of

two PVC tubes (length: 22 cm $\times$ diameter: 12.5 cm) filled with 75 $\pm$ 9 (SD) reed

internodes of 20 cm length and diameters varying between 0.1 and 2.0 cm (Staab et al. 2014,

Fornoff et al. 2021). Every month, we sampled the reeds with nesting hymenopterans and

replaced them with internodes of the same diameter. All the samples were reared in glass test

tubes under ambient conditions until specimens hatched. We identified hatched hosts and

parasitoids to species or morphospecies (Supplementary Table S1) based on reference

specimens (vouchered at the Institute of Zoology, CAS, Beijing). We were interested in the

general patterns of host-parasitoid interactions at the community level, so for the analysis we

did not distinguish between the two life-history strategies of parasitoids (true parasitoids and

kleptoparasitoids, including hymenopteran and dipteran parasitoids) because they both have

the same ecological result, death of host brood cells. We evaluated our sampling

completeness with r package "*iNEXT*" (Hsieh et al. 2016).

DNA extraction and amplification

All specimens were sequenced for a region of the mitochondrial cytochrome c oxidase

subunit I (COI) gene (Hebert et al. 2003). We extracted whole-genomic DNA of hosts and

parasitoids using DNeasy Blood & Tissue Kits (QIAGEN GmbH, Hilden, Germany),

following the manufacturer's protocols. COI sequences of samples were amplified using

universal primer pairs, LCO1490 (GGTCAACAAATCATAAAGATATTGG) as the forward

primer and HCO2198 (TAAACTTCAGGGTGACCAAAAAATCA) or HCOout

(CCAGGTAAAATTAAAATATAAACTTC) as the reverse primer. We carried out

polymerase chain reactions (PCR) in 96-well plates with 30 μl reactions containing 10 μl

ddH2O, 15 μl Premix PrimeSTAR HS (TaKaRa), 1 ul of each primer at 10 μM, and 3 μl

template genomic DNA using a thermo cycling profile. The PCR procedure as follows: 94℃

for 1 min; 94℃ for 1 min, 45℃ for 1.5 min and 72℃ for 1.5 min, cycle for 5 times; 94℃

for 2 min, 58℃ for 1.5 min and 72℃ for 1 min, cycle for 36 times; 72℃ for 5 min. We

performed all PCRs on an Eppendorf Mastercycler gradient, which were then visualized on a

1% agarose gel. Samples with clean single bands were sequenced after PCR purification

using BigDye v3.1 on an ABI 3730xl DNA Analyser (Applied Biosystems).

Sequence alignment and phylogenetic analysis

We applied MAFFT (Misawa, Katoh, Kuma, & Miyata, 2002) to align all sequences, then

translated the nucleotides into amino acids via MEGA v7.0 (Kumar, Stecher, & Tamura, 2016)

to check for the presence of stop codons with manual adjustments. Host and parasitoid

sequences were then aligned against the references using a Perl-based DNA barcode aligner

(Chesters, 2019).

We employed two strategies to improve the phylogenetic structure of a DNA barcode

phylogeny, which demonstrably improve resulting phylogeny-based diversity indices

(Macías-Hernández et al. 2020). These include the integration of 1) molecular sequences of

the plot data and 2) phylogenetic relationships from other molecular datasets. Integration was

achieved following Wang et al. (2020) and Chesters (2020): reference DNA barcodes of

Hymenoptera and Diptera were downloaded from the BOLD API

(www.boldsystems.org/index.php/API_Public), which were variously processed (e.g. to retain

only fully taxonomically labelled barcodes, to remove low quality or mislabeled entries, and

to dereplicate to a single exemplar per species), and then aligned (Chesters 2019). A single

outgroup was included for which we selected the most appropriate insect order sister to

Diptera and Hymenoptera (Misof et al. 2014), a representative of the order Psocoptera

(Psocidae, *Psocus leidyi*). We then constructed a phylogeny of the references and subjects,

with references constrained according to the method described earlier (Chesters 2020). A

number of backbone topologies were integrated for setting hard and soft constraints, including

a transcriptomics-derived topology (Chesters 2020), a mitogenome tree of insects (Chesters

2017), Diptera-specific trees (Wiegmann et al. 2011, Cranston et al. 2012, Ament 2017) and

Hymenoptera-specific trees (Peters et al. 2011, Branstetter et al. 2017, Cardinal 2018). The

constrained inference was conducted with RaxML version 8 (Stamatakis 2014) under the

standard GTRGAMMA DNA model with 24 rate categories. According to the backbone trees

used, most taxa present were monophyletic with a notable exception of Crabronidae, for

which there is emerging phylogenomic evidence of its polyphyly (Sann et al. 2018).

Tree phylogenetic diversity, functional diversity, and environmental covariates

The phylogenetic diversity of the tree communities was quantified by calculating wood

volume-weighted phylogenetic Mean Pairwise Distance (MPD) (Tucker et al. 2017). Tree

wood volume was estimated from data on basal area and tree height (Bongers et al. 2021)

measured in the center of each plot. Moreover, to represent variations towards the tips of the

phylogeny beyond MPD, we additionally calculated Mean Nearest Taxon Distance (MNTD),

which is a measure that quantifies the distance between each species and its nearest neighbor

on the phylogenetic tree (Webb 2000). Phylogenetic metrics of trees (tree MPD and MNTD)

were calculated based on a maximum likelihood phylogenetic tree available for the tree

species in our study area (Purschke et al. 2017). Considering that predatory wasps mainly

feed on herbivorous caterpillars, we calculated tree functional diversity to test the indirect

effects on hymenopteran communities and relevant network indices. Specifically, seven leaf

traits were used for calculation of tree functional diversity, which was calculated as the mean

pairwise distance in trait values among tree species, weighted by tree wood volume, and

expressed as Rao's Q (Ricotta and Moretti 2011), including specific leaf area, leaf toughness,

leaf dry matter content, leaf carbon content, ratio of leaf carbon to nitrogen, leaf magnesium

content, and leaf calcium content. These functional traits were commonly related to higher

trophic levels in our study area, such as herbivores and predators (Wang et al. 2020, Chen et

al. 2023), which are the main food resources of our predatory wasps. All of the traits were

measured on pooled samples of sun-exposed leaves of a minimum of five tree individuals per

species following standard protocols (Pérez-Harguindeguy et al. 2003).

As our analyses mainly compare community patterns among study plots, we additionally

considered potential effects of environmental variation by using plot means of slope,

elevation, "eastness" (sine-transformed radian values of aspect), and "northness"

(cosine-transformed radian values of aspect) as environmental covariates that characterize the

heterogeneity of the study plots. We also accounted for the potential effects of canopy cover

at plot level for host-parasitoid interactions, as it can structure hymenopteran communities

(Perlík et al. 2023). Canopy cover was calculated as in Fornoff et al. (2021) based on

hemispherical photographs.

Statistical analysis

All analyses were conducted in R 4.1.2 with the packages *ape*, *vegan*, *picante*, *bipartite*, and

caper (http://www.R-project.org). Prior to analysis, samples from the five years (2015, 2016,

2018, 2019, and 2020) were pooled at the plot level to discern overall and generalizable

effects permeating this system. We excluded three plots with no living trees because of high

mortality, resulting in 85 plots in the final analysis.

Composition of trees, hosts and parasitoids

The species and phylogenetic composition of trees, hosts, and parasitoids were quantified

with nonmetric multidimensional scaling (NMDS) analysis based on Morisita-Horn distances.

The minimum number of required dimensions in the NMDS based on the reduction in stress

value was determined in the analysis (k = 2 in our case). We centered the results to acquire

maximum variance on the first dimension, and used the principal components rotation in the

analysis. The phylogenetic composition was calculated by mean pairwise distance among the

host or parasitoid communities per plot with the R package "*picante*" applying the *'mpd'*

function. To test the influence of study plot heterogeneity on these relationships, we fitted

their standardized values (see vectors in Table S2) to the ordination on the basis of a

regression with the NMDS axis scores (Quinn and Keough 2002). NMDS was widely used to

summarize the variation in species composition across plots. The two axes extracted from the

NMDS represent gradients in community composition, where each axis reflects a subset of

the compositional variation. These axes should not be interpreted in isolation, as the overall

species composition is co-determined by their combined variation. For clarity, results were

interpreted based on the relationships of variables with the compositional gradients captured

by both axes together. For the analysis, we considered tree species richness, tree functional

and phylogenetic diversity, canopy cover, and environmental covariates (elevation, eastness,

northness, and slope) as plot characteristics. We assessed the significance of correlations with

permutation tests (permutation: $n= 999$). To strengthen the robustness of our findings based

on NMDS, we further validated the composition results using Mantel test and PERMANOVA

(with 'adonis2') for correlation between communities and relationships between communities

and environmental variables.

Phylogenetic match of hosts and parasitoids

In addition, we used a parafit test (9,999 permutations) with the R package "*ape*" to test

whether the associations were non-random between hosts and parasitoids. This is widely used

to assess host-parasite co-phylogeny by analyzing the congruence between host and parasite

phylogenies using a distance-based matrix approach. The species that were not attacked by

parasitoids or failed to generate sequences were excluded from the analyses. For species

abundance and composition, see Table S1.

Host-parasitoid interactions

We constructed quantitative host-parasitoid networks at community level with the R package

"*bipartite*" for each plot of the two sites. Bees and wasps were considered together as hosts

because there were too few abundant bee species to analyze separate interaction networks for

bees and wasps. We calculated five indices to quantitatively characterize the structure of the

interaction networks (Blüthgen and Staab 2024): weighted parasitoid generality (effective

number of host species per parasitoid species), weighted host vulnerability (effective number

of parasitoid species attacking a host species), robustness (degree of network stability),

linkage density (degree of network specialization), and interaction evenness (degree of

network evenness). Parasitoid generality was defined as the weighted mean number of host

species per parasitoid species, $G_{qw} = \sum_{j=1}^{J} \frac{A_j}{m} 2^{H_j}$, with A_j being the number of interactions of

parasitoid species j, m the total number of interactions of all species, and H_j the Shannon

diversity of interactions of species j. Host vulnerability was the weighted mean number of

parasitoid species per host species, $\text{Vulnerability} = \sum_{i=1}^{I} \frac{A_i}{m} 2^{H_i}$ (Bersier et al. 2002).

Robustness was defined as the area under the extinction curve, reflecting the degree of

decreases of one trophic level with the random elimination species of the other trophic levels,

here using the robustness index for higher trophic levels (i.e. parasitoids). The robustness

calculation was also weighted based on interaction abundance. For linkage density, $L_q =$

$0.5 \left(\sum_{j=1}^{J} \frac{A_j}{m} 2^{H_j} + \sum_{i=1}^{I} \frac{A_i}{m} 2^{H_i} \right)$, we used the realized proportion of possible links between the

two trophic levels as the mean number of interactions per species across the entire network

(Tylianakis et al. 2007). Interaction evenness was defined as $E_S = -\sum_i \sum_j p_{ij} \ln p_{ij} / \ln IJ$,

which is used to describe Shannon's evenness of network interactions (Dormann et al. 2009).

To check whether all network indices significantly differ from chance across all study plots,

we used Patefield null models (Dormann et al. 2009) to compare observed indices with

simulated values (10,000 times).

Linear models

To test the effects of tree species richness, tree phylogenetic, and functional diversity, as well

as canopy cover and the other environmental covariates (including slope, elevation, eastness,

and northness) on the six network indices, we used linear mixed-effect models. For our

analyses, the study sites were considered as random effects, and the other variables were

treated as fixed effects (see above). Given the strong correlation between tree species richness

and tree MPD (Pearson's r = 0.74, p<0.01), we excluded tree MPD in the models where tree

species richness was a predictor. To evaluate the potential effects caused by tree MPD, we also ran alternative models where tree species richness was replaced with tree MPD. We simplified all models by gradually removing non-significant factors to obtain the most parsimonious model with the lowest AICc. To ensure that the analyses were not strongly affected by multicollinearity, the correlations among all predictors were tested (Fig. S3), and variance inflation factors (VIF) of our statistical models were checked.

Acknowledgements

We thank the BEF-China consortium for support (especially, Bo Yang). We thank Mr. Yinquan Qi for his help with the collection. This work was supported by the National Key Research Development Program of China (2022YFF0802300), the National Science Fund for Excellent Young Scholars (32122016), the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB310304), the National Natural Science Foundation, China (32100343, 32070465), and the National Science Fund for Distinguished Young Scholars (31625024). MQW was supported by the Alexander von Humboldt research fellowships. CDZ's lab is funded by the Key Program of the National Natural Science Foundation of China (No. 32330013) and also has been continuously supported by grants from the Key Laboratory of the Zoological Systematics and Evolution of the Chinese Academy of Sciences (grant number 2008DP173354).

541   Table 1 Environmental correlates of dissimilarity matrixes with predictors (NMDS on

542   Morisita-Horn dissimilarity) across the study plots. Significant P-values are indicated in bold.

543   See Table S2-S5 for the complete information.

	Host species community	Parasitoid species community	Host phylogenetic community	Parasitoid phylogenetic community
Tree phylogeny NMDS1	0.225	0.422	0.386	0.274
Tree phylogeny NMDS2	**0.003**	0.12	0.128	**0.024**
Tree composition NMDS1	**0.001**	**0.001**	**0.001**	**0.001**
Tree composition NMDS2	0.604	0.418	0.433	**0.031**
Canopy cover	**0.001**	**0.001**	**0.001**	**0.004**
Tree species richness	**0.035**	0.122	0.100	0.094
Elevation	**0.005**	**0.007**	**0.001**	**0.001**
Eastness	0.079	0.045	**0.04**	**0.001**
Northness	0.49	0.837	0.821	0.340
Slope	**0.031**	0.507	0.507	0.959
Tree FD (Rao's Q)	0.094	**0.019**	**0.021**	**0.031**
Tree MPD	**0.005**	**0.021**	**0.013**	0.223
Host phylogeny NMDS1	-	**0.016**	-	0.584
Host phylogeny NMDS2	-	**0.027**	-	0.914
Host composition NMDS1	-	**0.001**	-	**0.008**
Host composition NMDS2	-	0.169	-	0.138
Parasitoid phylogeny NMDS1	**0.001**	-	**0.001**	-
Parasitoid phylogeny NMDS2	0.462	-	0.058	-
Parasitoid composition NMDS1	**0.001**	-	**0.001**	-
Parasitoid composition NMDS2	**0.014**	-	**0.001**	-

546 Table 2. Summary results of linear mix-effects models for parasitoid generality, host

547 vulnerability, robustness, linkage density, and interaction evenness of host-parasitoid network

548 indices at the community level across the tree species richness gradient. Standardized

549 parameter estimates (with standard errors, t and P values) are shown for the variables retained

550 in the minimal models.

		Est.	SE	t	P
Parasitoid generality	Intercept	0.176	0.016	10.96	<0.001
	Canopy cover	0.033	0.016	2.03	0.046
Host vulnerability	Intercept	2.873	0.08	35.89	<0.001
	Elevation	-0.140	0.08	-1.66	0.101
	Tree species richness: Site A	0.150	0.13	1.20	0.234
	Tree species richness: Site B	0.230	0.11	2.12	0.037
Robustness of parasitoids	Intercept	0.630	0.01	84.43	<0.001
	Tree species richness: Site A	-0.022	0.01	-1.99	0.049
	Tree species richness: Site B	-0.019	0.01	-1.90	0.061
Linkage density	Intercept	2.038	0.04	50.99	<0.001
	Elevation	-0.078	0.04	-1.85	0.069
	Tree species richness: Site A	0.106	0.06	1.70	0.094
	Tree species richness: Site B	0.106	0.04	2.68	0.009
Interaction evenness	Intercept	0.511	0.009	59.12	0.025
	Canopy cover	-0.037	0.007	-5.06	<0.001
	Eastness	-0.018	0.007	-2.50	0.015

Fig. 1. Ordination plot of the non-metric multidimensional scaling (NMDS) analysis
of (a) host species composition, (b) parasitoid species composition, (c) host
phylogenetic composition, and (d) parasitoid phylogenetic composition across the
study plots (filled circles) in the BEF-China experiment. Stress = 0.23, 0.23, 0.24 and
0.20, respectively. Arrows indicate significant (at $p < 0.05$) correlations of
environmental variables with NMDS axis scores. Lengths of arrows are proportional
to the strength of the correlations. Red crosses refer to the host or parasitoid species in
each community. See Table S2-S5 in the Supplementary Material for abbreviations
and statistical values.

Fig. 2. Dendrogram of phylogenetic congruence for the host species (below) and associated parasitoid species (above) recorded in the study. Each rectangle represents a different superfamily (for host species) or family (for parasitoid species). H1: Pompilidae H2: Apoidea H3: Vespidae; P1: Sarcophagidae P2: Phoridae P3: Bombyliidae P4: Trigonalyidae P5: Mutillidae P6: Megachilidae P7: Chrysididae P8: Ichneumonidae P9: Chalcidoidea. The trophic network of hosts and parasitoids was non-randomly structured (parafit test: P = 0.032). Host and parasitoid species names are given in Fig. S4.

575   Fig. 3. Community-level relationships of network between tree species richness and (a)

576   vulnerability, (b) linkage density, and (c) robustness of parasitoids. Values were adjusted for

577   covariates of the final regression model. Regression lines (with 95% confidence bands) show

578   significant ($p < 0.05$) relationships. Note that axes are on a log scale for tree species richness.

Fig. 4. Community-level relationships of network between tree mean pairwise phylogenetic distance and (a) vulnerability and (b) linkage density and community-level relationships of network between canopy cover and (c) interaction evenness. Values were adjusted for covariates of the final regression model. Regression lines (with 95% confidence bands) show significant ($p < 0.05$) relationships.

References

Ament, D. C. 2017. Phylogeny of Phorinae sensu latu (Diptera: Phoridae) inferred from a
morphological analysis with comprehensive taxon sampling and an uncommon
method of character coding. Zoological Journal of the Linnean Society **181**:151-188.
Barnes, A. D., M. Jochum, J. S. Lefcheck, N. Eisenhauer, C. Scherber, M. I. O'Connor, P. de
Ruiter, and U. Brose. 2018. Energy Flux: The Link between Multitrophic Biodiversity
and Ecosystem Functioning. Trends in Ecology & Evolution **33**:186-197.
Bersier, L.-F., C. Banašek-Richter, and M.-F. Cattin. 2002. Quantitative Descriptors of
Food-Web Matrices. Ecology **83**:2394-2407.
Blüthgen, N., and A.-M. Klein. 2011. Functional complementarity and specialisation: The role
of biodiversity in plant–pollinator interactions. Basic and Applied Ecology
12:282-291.
Blüthgen, N., and M. Staab. 2024. A critical evaluation of network approaches for studying
species interactions. Annual Review of Ecology, Evolution, and Systematics **55**.
Bongers, F. J., B. Schmid, H. Bruelheide, F. Bongers, S. Li, G. von Oheimb, Y. Li, A. Cheng, K.
Ma, and X. Liu. 2021. Functional diversity effects on productivity increase with age in
a forest biodiversity experiment. Nature Ecology & Evolution.
Branstetter, M. G., J. T. Longino, P. S. Ward, and B. C. Faircloth. 2017. Enriching the ant tree
of life: Enhanced UCE bait set for genome‐scale phylogenetics of ants and other
Hymenoptera. Methods in Ecology and Evolution **8**:768-776.
Brondizio, E. S., J. Settele, S. Díaz, and H. T. Ngo. 2019. Global assessment report on
biodiversity and ecosystem services of the Intergovernmental Science-Policy
Platform on Biodiversity and Ecosystem Services.
Bruelheide, H., K. Nadrowski, T. Assmann, J. Bauhus, S. Both, F. Buscot, X. Y. Chen, B. Y. Ding,
W. Durka, A. Erfmeier, J. L. M. Gutknecht, D. L. Guo, L. D. Guo, W. Hardtle, J. S. He, A.
M. Klein, P. Kuhn, Y. Liang, X. J. Liu, S. Michalski, P. A. Niklaus, K. Q. Pei, M.
Scherer-Lorenzen, T. Scholten, A. Schuldt, G. Seidler, S. Trogisch, G. von Oheimb, E.
Welk, C. Wirth, T. Wubet, X. F. Yang, M. J. Yu, S. R. Zhang, H. Z. Zhou, M. Fischer, K. P.
Ma, and B. Schmid. 2014. Designing forest biodiversity experiments: general
considerations illustrated by a new large experiment in subtropical China. Methods
in Ecology and Evolution **5**:74-89.
Cagnolo, L., A. Salvo, and G. Valladares. 2011. Network topology: patterns and mechanisms
in plant‐herbivore and host‐parasitoid food webs. Journal of Animal Ecology
80:342-351.
Cardinal, S. 2018. Bee (Hymenoptera: Apoidea: Anthophila) diversity through time. Insect
biodiversity: science and society **2**:851-867.
Cavender-Bares, J., K. H. Kozak, P. V. Fine, and S. W. Kembel. 2009. The merging of
community ecology and phylogenetic biology. Ecology Letters **12**:693-715.
Chen, J.-T., M.-Q. Wang, Y. Li, D. Chesters, A. Luo, W. Zhang, P.-F. Guo, S.-K. Guo, Q.-S. Zhou,
K.-P. Ma, H. Bruelheide, A. Schuldt, and C.-D. Zhu. 2023. Functional and phylogenetic
relationships link predators to plant diversity via trophic and non-trophic pathways.
Proceedings of the Royal Society B **290**:20221658.

Chesters, D. 2017. Construction of a Species-Level Tree of Life for the Insects and Utility in
Taxonomic Profiling. Syst Biol **66**:426-439.
Chesters, D. 2019. aligner. pl. Computer software available at
https://sourceforge.net/projects/dna- barcode-aligner/.

Chesters, D. 2020. The phylogeny of insects in the data-driven era. Systematic Entomology
45:540-551.
Cranston, P. S., N. B. Hardy, and G. E. Morse. 2012. A dated molecular phylogeny for the
Chironomidae (Diptera). Systematic Entomology **37**:172-188.
Dáttilo, W., and L. Dyer. 2014. Canopy openness enhances diversity of ant–plant interactions
in the Brazilian Amazon rain forest. Biotropica **46**:712-719.
De Frenne, P., J. Lenoir, M. Luoto, B. R. Scheffers, F. Zellweger, J. Aalto, M. B. Ashcroft, D. M.
Christiansen, G. Decocq, and K. De Pauw. 2021. Forest microclimates and climate
change: Importance, drivers and future research agenda. Global change biology
27:2279-2297.
De Ruiter, P. C., A.-M. Neutel, and J. C. Moore. 1995. Energetics, patterns of interaction
strengths, and stability in real ecosystems. Science **269**:1257-1260.
Dormann, C. F., J. Fründ, N. Blüthgen, and B. Gruber. 2009. Indices, graphs and null models:
analyzing bipartite ecological networks. The Open Ecology Journal **2**:7-24.
Ebeling, A., A.-M. Klein, and T. Tscharntke. 2011. Plant–flower visitor interaction webs:
Temporal stability and pollinator specialization increases along an experimental
plant diversity gradient. Basic and Applied Ecology **12**:300-309.
Ebeling, A., A.-M. Klein, W. W. Weisser, and T. Tscharntke. 2012. Multitrophic effects of
experimental changes in plant diversity on cavity-nesting bees, wasps, and their
parasitoids. Oecologia **169**:453-465.
Emerson, B. C., and R. G. Gillespie. 2008. Phylogenetic analysis of community assembly and
structure over space and time. Trends in Ecology & Evolution **23**:619-630.
Finke, D. L., and R. F. Denno. 2002. Intraguild predation diminished in complex‐structured
vegetation: implications for prey suppression. Ecology **83**:643-652.
Fornoff, F., A. M. Klein, N. Blüthgen, and M. Staab. 2019. Tree diversity increases robustness
of multi-trophic interactions. Proceedings: Biological sciences, The Royal Society
286:20182399.
Fornoff, F., M. Staab, C.-D. Zhu, and A.-M. Klein. 2021. Multi-trophic communities
re-establish with canopy cover and microclimate in a subtropical forest biodiversity
experiment. Oecologia **196**:289-301.
Godfray, H. C. J., and H. Godfray. 1994. Parasitoids: behavioral and evolutionary ecology.
Princeton University Press.
Gonzalez, A., R. M. Germain, D. S. Srivastava, E. Filotas, L. E. Dee, D. Gravel, P. L. Thompson,
F. Isbell, S. Wang, and S. Kéfi. 2020. Scaling‐up biodiversity‐ecosystem
functioning research. Ecology Letters **23**:757-776.
Grass, I., B. Jauker, I. Steffan-Dewenter, T. Tscharntke, and F. Jauker. 2018. Past and
potential future effects of habitat fragmentation on structure and stability of
plant–pollinator and host–parasitoid networks. Nature Ecology & Evolution
2:1408-1417.

674 Grossman, J. J., M. Vanhellemont, N. Barsoum, J. Bauhus, H. Bruelheide, B. Castagneyrol, J.
675   Cavender-Bares, N. Eisenhauer, O. Ferlian, D. Gravel, A. Hector, H. Jactel, H. Kreft, S.
676   Mereu, C. Messier, B. Muys, C. Nock, A. Paquette, J. Parker, M. P. Perring, Q. Ponette,
677   P. B. Reich, A. Schuldt, M. Staab, M. Weih, D. C. Zemp, M. Scherer-Lorenzen, and K.
678   Verheyen. 2018. Synthesis and future research directions linking tree diversity to
679   growth, survival, and damage in a global network of tree diversity experiments.
680   Environmental and Experimental Botany **152**:68-89.
681 Guo, P.-F., M.-Q. Wang, M. Orr, Y. Li, J.-T. Chen, Q.-S. Zhou, M. Staab, F. Fornoff, G.-H. Chen,
682   N.-L. Zhang, A.-M. Klein, and C.-D. Zhu. 2021. Tree diversity promotes predatory
683   wasps and parasitoids but not pollinator bees in a subtropical experimental forest.
684   Basic and Applied Ecology **53**:134-142.
685 Haddad, N. M., G. M. Crutsinger, K. Gross, J. Haarstad, and D. Tilman. 2011. Plant diversity
686   and the stability of foodwebs. Ecology Letters **14**:42-46.
687 Harvey, E., I. Gounand, C. L. Ward, and F. Altermatt. 2017. Bridging ecology and conservation:
688   from ecological networks to ecosystem function. Journal of Applied Ecology
689   **54**:371-379.
690 Hebert, P. D., S. Ratnasingham, and J. R. deWaard. 2003. Barcoding animal life: cytochrome c
691   oxidase subunit 1 divergences among closely related species. Proceedings: Biological
692   sciences, The Royal Society **270 Suppl 1**:S96-99.
693 Hines, J., A. Ebeling, A. D. Barnes, U. Brose, C. Scherber, S. Scheu, T. Tscharntke, W. W.
694   Weisser, D. P. Giling, A. M. Klein, and N. Eisenhauer. 2019. Chapter Nine - Mapping
695   change in biodiversity and ecosystem function research: food webs foster
696   integration of experiments and science policy. Pages 297-322 *in* N. Eisenhauer, D. A.
697   Bohan, and A. J. Dumbrell, editors. Advances in Ecological Research. Academic Press.
698 Hsieh, T. C., K. H. Ma, A. Chao, and G. McInerny. 2016. iNEXT: an R package for rarefaction
699   and extrapolation of species diversity (Hill numbers). Methods in Ecology and
700   Evolution **7**:1451-1456.

701 Hunter, M. D., M. D. Hunter, T. Ohgushi, and P. W. Price. 1992. Interactions within herbivore
702   communities mediated by the host plant: the keystone herbivore concept.
703 Isbell, F., P. Balvanera, A. S. Mori, J. S. He, J. M. Bullock, G. R. Regmi, E. W. Seabloom, S.
704   Ferrier, O. E. Sala, and N. R. Guerrero‐Ramírez. 2022. Expert perspectives on
705   global biodiversity loss and its drivers and impacts on people. Frontiers in Ecology
706   and the Environment.
707 Jeffs, C. T., and O. T. Lewis. 2013. Effects of climate warming on host–parasitoid interactions.
708   Ecological Entomology **38**:209-218.
709 Lopez-Carretero, A., C. Diaz-Castelazo, K. Boege, and V. Rico-Gray. 2014. Evaluating the
710   spatio-temporal factors that structure network parameters of plant-herbivore
711   interactions. PLoS ONE **9**:e110430.
712 Loyola, R. D., and R. P. Martins. 2008. Habitat structure components are effective predictors
713   of trap-nesting Hymenoptera diversity. Basic and Applied Ecology **9**:735-742.
714 Ma, S., A. Concilio, B. Oakley, M. North, and J. Chen. 2010. Spatial variability in microclimate
715   in a mixed-conifer forest before and after thinning and burning treatments. Forest
716   Ecology and Management **259**:904-915.

Macías-Hernández, N., M. Domènech, P. Cardoso, B. C. Emerson, P. A. V. Borges, J.
Lozano-Fernandez, O. S. Paulo, A. Vieira, A. Enguídanos, F. Rigal, I. R. Amorim, and M.
A. Arnedo. 2020. Building a Robust, Densely-Sampled Spider Tree of Life for
Ecosystem Research. Diversity **12**:288.
Maunsell, S. C., R. L. Kitching, C. J. Burwell, and R. J. Morris. 2015. Changes in host-parasitoid
food web structure with elevation. Journal of Animal Ecology **84**:353-363.
Misof, B., S. Liu, K. Meusemann, R. S. Peters, A. Donath, C. Mayer, P. B. Frandsen, J. Ware, T.
Flouri, and R. G. Beutel. 2014. Phylogenomics resolves the timing and pattern of
insect evolution. Science **346**:763-767.
Pellissier, L., C. Ndiribe, A. Dubuis, J. N. Pradervand, N. Salamin, A. Guisan, and S. Rasmann.
2013. Turnover of plant lineages shapes herbivore phylogenetic beta diversity along
ecological gradients. Ecology Letters **16**:600-608.
Peralta, G., C. M. Frost, R. K. Didham, A. Varsani, and J. M. Tylianakis. 2015. Phylogenetic
diversity and co-evolutionary signals among trophic levels change across a habitat
edge. Journal of Animal Ecology **84**:364-372.
Pérez-Harguindeguy, N., S. Díaz, F. Vendramini, J. H. C. Cornelissen, D. E. Gurvich, and M.
Cabido. 2003. Leaf traits and herbivore selection in the field and in cafeteria
experiments. Austral Ecology **28**:642-650.
Perlík, M., D. Kraus, H. Bußler, L. Neudam, S. Pietsch, U. Mergner, D. Seidel, P. Sebek, and S.
Thorn. 2023. Canopy openness as the main driver of aculeate Hymenoptera and
saproxylic beetle diversity following natural disturbances and salvage logging. Forest
Ecology and Management **540**:121033.
Peters, R. S., B. Meyer, L. Krogmann, J. Borner, K. Meusemann, K. Schütte, O. Niehuis, and B.
Misof. 2011. The taming of an impossible child: a standardized all-in approach to the
phylogeny of Hymenoptera using public database sequences. BMC Biology **9**:1-14.
Popkin, G. 2021. Forest fight. Science **374**:1184-1189.
Purschke, O., S. G. Michalski, H. Bruelheide, and W. Durka. 2017. Phylogenetic turnover
during subtropical forest succession across environmental and phylogenetic scales.
Ecology and Evolution **7**:11079-11091.
Quinn, G. P., and M. J. Keough. 2002. Experimental Design and Data Analysis for Biologists.
Cambridge University Press, Cambridge.
Reitalu, T., A. E. Bjune, A. Blaus, T. Giesecke, A. Helm, I. Matthias, S. M. Peglar, J. S. Salonen,
H. Seppä, and V. Väli. 2019. Patterns of modern pollen and plant richness across
northern Europe. Journal of Ecology **107**:1662-1677.
Ricotta, C., and M. Moretti. 2011. CWM and Rao's quadratic diversity: a unified framework
for functional ecology. Oecologia **167**:181-188.
Rodriguez, A., J. L. Pohjoismäki, and J. Kouki. 2019. Diversity of forest management
promotes parasitoid functional diversity in boreal forests. Biological Conservation
238:108205.
Root, R. B. 1973. Organization of a Plant‐Arthropod Association in Simple and Diverse
Habitats: The Fauna of Collards (Brassica Oleracea). Ecological Monographs
43:95-124.

Sann, M., O. Niehuis, R. S. Peters, C. Mayer, A. Kozlov, L. Podsiadlowski, S. Bank, K.
Meusemann, B. Misof, and C. Bleidorn. 2018. Phylogenomic analysis of Apoidea
sheds new light on the sister group of bees. Bmc Evolutionary Biology **18**:1-15.
Schuldt, A., A. Ebeling, M. Kunz, M. Staab, C. Guimarães-Steinicke, D. Bachmann, N.
Buchmann, W. Durka, A. Fichtner, F. Fornoff, W. Härdtle, L. R. Hertzog, A.-M. Klein, C.
Roscher, J. Schaller, G. von Oheimb, A. Weigelt, W. Weisser, C. Wirth, J. Zhang, H.
Bruelheide, and N. Eisenhauer. 2019. Multiple plant diversity components drive
consumer communities across ecosystems. Nature Communications **10**:1460.
Sobek, S., T. Tscharntke, C. Scherber, S. Schiele, and I. Steffan-Dewenter. 2009. Canopy vs.
understory: does tree diversity affect bee and wasp communities and their natural
enemies across forest strata? Forest Ecology and Management **258**:609-615.
Staab, M., N. Blüthgen, and A.-M. Klein. 2015. Tree diversity alters the structure of a
tri-trophic network in a biodiversity experiment. Oikos **124**:827-834.
Staab, M., H. Bruelheide, W. Durka, S. Michalski, O. Purschke, C. D. Zhu, and A. M. Klein.
2016. Tree phylogenetic diversity promotes host-parasitoid interactions.
Proceedings of the Royal Society B: Biological Sciences **283**.
Staab, M., X. Liu, T. Assmann, H. Bruelheide, F. Buscot, W. Durka, A. Erfmeier, A. M. Klein, K.
Ma, S. Michalski, T. Wubet, B. Schmid, A. Schuldt, and J. Koricheva. 2020. Tree
phylogenetic diversity structures multitrophic communities. Functional Ecology
35:521-534.
Staab, M., M. Ohl, C.-D. Zhu, and A.-M. Klein. 2014. A unique nest-protection strategy in a
new species of spider wasp. PLoS ONE **9**:e101592.
Staab, M., G. Pufal, T. Tscharntke, A. M. Klein, and G. Iossa. 2018. Trap nests for bees and
wasps to analyse trophic interactions in changing environments—A systematic
overview and user guide. Methods in Ecology and Evolution **9**:2226-2239.
Staab, M., and A. Schuldt. 2020. The influence of tree diversity on natural enemies—a review
of the "enemies" hypothesis in forests. Current Forestry Reports **6**:243-259.
Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics **30**:1312-1313.
Tittensor, D. P., M. Walpole, S. L. Hill, D. G. Boyce, G. L. Britten, N. D. Burgess, S. H. Butchart,
P. W. Leadley, E. C. Regan, R. Alkemade, R. Baumung, C. Bellard, L. Bouwman, N. J.
Bowles-Newark, A. M. Chenery, W. W. Cheung, V. Christensen, H. D. Cooper, A. R.
Crowther, M. J. Dixon, A. Galli, V. Gaveau, R. D. Gregory, N. L. Gutierrez, T. L. Hirsch,
R. Hoft, S. R. Januchowski-Hartley, M. Karmann, C. B. Krug, F. J. Leverington, J. Loh, R.
K. Lojenga, K. Malsch, A. Marques, D. H. Morgan, P. J. Mumby, T. Newbold, K.
Noonan-Mooney, S. N. Pagad, B. C. Parks, H. M. Pereira, T. Robertson, C. Rondinini, L.
Santini, J. P. Scharlemann, S. Schindler, U. R. Sumaila, L. S. Teh, J. van Kolck, P.
Visconti, and Y. Ye. 2014. A mid-term analysis of progress toward international
biodiversity targets. Science **346**:241-244.
Tucker, C. M., M. W. Cadotte, S. B. Carvalho, T. J. Davies, S. Ferrier, S. A. Fritz, R. Grenyer, M.
R. Helmus, L. S. Jin, A. O. Mooers, S. Pavoine, O. Purschke, D. W. Redding, D. F.
Rosauer, M. Winter, and F. Mazel. 2017. A guide to phylogenetic metrics for

conservation, community ecology and macroecology. Biological Reviews of the
Cambridge Philosophical Society **92**:698-715.
Tylianakis, J. M., R. K. Didham, J. Bascompte, and D. A. Wardle. 2008. Global change and
species interactions in terrestrial ecosystems. Ecology Letters **11**:1351-1363.
Tylianakis, J. M., T. Tscharntke, and A.-M. Klein. 2006. Diversity, ecosystem function, and
stability of parasitoid–host interactions across a tropical habitat gradient. Ecology
87:3047-3057.
Tylianakis, J. M., T. Tscharntke, and O. T. Lewis. 2007. Habitat modification alters the
structure of tropical host–parasitoid food webs. Nature **445**:202.
Valladares, G., L. Cagnolo, and A. Salvo. 2012. Forest fragmentation leads to food web
contraction. Oikos **121**:299-305.
Vidal, M. C., and S. M. Murphy. 2018. Bottom-up vs. top-down effects on terrestrial insect
herbivores: a meta-analysis. Ecology Letters **21**:138-150.
Volf, M., P. Pyszko, T. Abe, M. Libra, N. Kotaskova, M. Sigut, R. Kumar, O. Kaman, P. T.
Butterill, J. Sipos, H. Abe, H. Fukushima, P. Drozd, N. Kamata, M. Murakami, and V.
Novotny. 2017. Phylogenetic composition of host plant communities drives
plant-herbivore food web structure. Journal of Animal Ecology **86**:556-565.
Volf, M., S. T. Segar, S. E. Miller, B. Isua, M. Sisol, G. Aubona, P. Simek, M. Moos, J. Laitila, J.
Kim, J. Zima, Jr., J. Rota, G. D. Weiblen, S. Wossa, J. P. Salminen, Y. Basset, and V.
Novotny. 2018. Community structure of insect herbivores is driven by conservatism,
escalation and divergence of defensive traits in Ficus. Ecology Letters **21**:83-92.
Wang, M.-Q., Y. Li, D. Chesters, P. Anttonen, H. Bruelheide, J.-T. Chen, W. Durka, P.-F. Guo,
W. Haerdtle, K. Ma, S. G. Michalski, B. Schmid, G. von Oheimb, C.-S. Wu, N.-L. Zhang,
Q.-S. Zhou, A. Schuldt, and C.-D. Zhu. 2019. Multiple components of plant diversity
loss determine herbivore phylogenetic diversity in a subtropical forest experiment.
Journal of Ecology **107**:2697-2712.
Wang, M. Q., Y. Li, D. Chesters, H. Bruelheide, K. Ma, P. F. Guo, Q. S. Zhou, M. Staab, C. D.
Zhu, and A. Schuldt. 2020. Host functional and phylogenetic composition rather than
host diversity structure plant-herbivore networks. Molecular Ecology **29**:2747-2762.
Webb, C. O. 2000. Exploring the Phylogenetic Structure of Ecological Communities: An
Example for Rain Forest Trees. Am Nat **156**:145-155.
Webb, C. O., D. D. Ackerly, M. A. McPeek, and M. J. Donoghue. 2002. Phylogenies and
Community Ecology. Annual Review of Ecology and Systematics **33**:475-505.
Wiegmann, B. M., M. D. Trautwein, I. S. Winkler, N. B. Barr, J.-W. Kim, C. Lambkin, M. A.
Bertone, B. K. Cassel, K. M. Bayless, and A. M. Heimberg. 2011. Episodic radiations in
the fly tree of life. Proceedings of the National Academy of Sciences **108**:5690-5695.