Multi-dimensionality of tree communities structure host-parasitoid networks and their phylogenetic composition

Ming-Qiang Wang[1,2#], Shi-Kun Guo[1,3#], Peng-Fei Guo[4#], Juan-Juan Yang[1,3], Guo-Ai Chen[1,3], Douglas Chesters[1,5], Michael Orr[1,6], Ze-Qing Niu[1], Michael Staab[7], Jing-Ting Chen[1,3], Yi Li[10], Qing-Song Zhou[1,5], Felix Fornoff[8], Xiaoyu Shi[1], Shan Li[10], Massimo Martini[8], Alexandra-Maria Klein[8], Andreas Schuldt[9], Xiaojuan Liu[10], Keping Ma[5,10], Helge Bruelheide[11], Arong Luo[1,5,13]*, Chao-Dong Zhu[1,3,5,12,13]*

CAS Key Laboratory of Zoological Systematics and Evolution, Institute of Zoology, Chinese Academy of Sciences, 1 Beichen West Road, Chaoyang District, Beijing, 100101, China

CAS Key Laboratory of Mountain Ecological Restoration and Bioresource Utilization & Biodiversity Conservation Key Laboratory of Sichuan Province, Chengdu Institute of Biology, Chinese Academy of Sciences, 4 Renmin South Road, Wuhou District, Chengdu, 610041, China

College of Biological Sciences, University of Chinese Academy of Sciences, No.19A Yuquan Road, Shijingshan District, Beijing, 100049, China

College of Pharmacy, Guizhou University of Traditional Chinese Medicine, Guiyang 550025, China

International College, University of Chinese Academy of Sciences, No.19A Yuquan Road, Shijingshan District, Beijing, 100049, China

Entomologie, Staatliches Museum für Naturkunde Stuttgart, Stuttgart, Germany.

Ecological Networks, Technical University Darmstadt, Schnittspahnstraße 3, 64287 Darmstadt, Germany

Department of Nature Conservation and Landscape Ecology, Albert-Ludwigs-University Freiburg, Tennenbacher Straße 4, 79106 Freiburg, Germany

Forest Nature Conservation, University of Göttingen, Büsgenweg 3, 37077, Göttingen, Germany

State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, 20 Nanxincun, Xiangshan, Haidian District, Beijing 100093, China

Institute of Biology/Geobotany and Botanical Garden, Martin Luther University Halle-Wittenberg, Große Steinstraße 79/80, 06108 Halle (Saale), Germany

State Key Laboratory of Integrated Pest Management, Institute of Zoology, Chinese Academy of Sciences, Beijing, 100101, China

Key Laboratory of Animal Biodiversity Conservation and Integrated Pest Management (Chinese Academy of Sciences), Beijing, 100101, China

[#]Ming-Qiang Wang, Shi-Kun Guo and Peng-Fei Guo equally contributed to this manuscript.

*Correspondence author: luoar@ioz.ac.cn; zhucd@ioz.ac.cn, Phone: +86-10-64807085, Fax: +86-10-64807099

Data availability

Data is available on Science Data Bank at https://www.scidb.cn/en/s/Ujq22u, and will be available on the BEF-China project database at https://data.botanik.uni-halle.de/bef-china/datasets.

Conflict of interest statement

The authors have no conflict of interest.

Author contributions

C.D.Z., A.L. and M.Q.W. conceived the idea for the manuscript; C.D.Z., A.M.K., S.K.G., P.F.G. and M.Q.W. designed the study. S.K.G., P.F.G., J.J.Y., G.A.C., Z.Q.N., M.S., J.T.C., Y.L., Q.S.Z., F.F., X.S., S.L., M.M., A.M.K., A.S., X.L., K.M. and H.B. collected and/or contributed data; D.C., M.Q.W. and S.K.G. conducted the phylogenetic analysis; M.Q.W. S.KG. and P.F.G. conducted the statistical analyses and wrote the manuscript, with input from F.F., A.S., M.S. and M.M and all coauthors.

Abstract

Environmental factors can influence ecological networks, but these effects are poorly understood

in the realm of the phylogeny of host-parasitoid interactions. Especially, we lack a comprehensive

understanding of the ways that biotic factors, including plant diversity, tree identity, genetic

diversity, overall community composition of higher trophic levels, and abiotic factors such as

microclimate, interact to determine host–parasitoid network structure and host–parasitoid

community dynamics. To address this, we leveraged a five-year dataset of trap-nesting bees and

wasps and their parasitoids collected in a highly-controlled, large-scale subtropical tree

biodiversity experiment. We tested for effects of tree species richness, tree phylogenetic and

functional diversity, and taxonomic and phylogenetic composition on taxonomic, phylogenetic,

and network composition of both host and parasitoid communities. We show that multiple

components of tree diversity, tree composition, and canopy cover impacted both, taxonomic and

phylogenetic composition of hosts and parasitoids. Generally, top-down control was stronger than

bottom-up control via phylogenetic association between hosts and parasitoids, reflecting

non-randomly structured interactions between phylogenetic trees of hosts and parasitoids. Further,

host-parasitoid network structure was influenced by tree species richness, tree phylogenetic

diversity, and canopy cover. Our study indicates that the composition of higher trophic levels and

corresponding interaction networks are determined by habitat structure and heterogeneity, which

is maintained by trees and especially via phylogenetic links in species-rich ecosystems.

Introduction

Understanding the ecological consequences of biodiversity loss is an increasingly important task

in ecology, given the ongoing biodiversity crisis (Isbell et al. 2022). Representing the

interdependencies among organisms, ecological networks reflect whether and how species interact

with each other across trophic levels, playing an indispensable role in assessing ecosystem

stability and integrity (De Ruiter et al. 1995, Harvey et al. 2017). Changes in network structure

usually coincide with variations in diversity and community of trophic levels, which could be in

turn affected by the changes in producers via trophic cascades (Barnes et al. 2018, Gonzalez et al.

2020). However, we still lack a generalizable framework for how these networks and especially

their phylogenetic interdependencies respond to basal changes of biodiversity loss in ecosystems,

such as changes in the tree diversity of forests (Tylianakis et al. 2008, Grossman et al. 2018) . To

better understand species codependence and its role for biodiversity conservation, we must further

study the complex dynamics of networks from multiple angles (Tittensor et al. 2014, Brondizio et

al. 2019). Interactions can be viewed from a top-down or bottom-up perspective. Higher trophic

levels can impact lower trophic levels through antagonistic species interactions, while lower

trophic levels can provide resources to support higher trophic levels, Previous studies have shown

there might be asymmetric effects between top-down and bottom-up control by different

mediators (Vidal and Murphy 2018), with both factors shaping multitrophic communities

together(Hunter et al. 1992). Moreover, the diversity and community of higher trophic levels

could be driven by environmental microclimate, which could be determined by plant structuring,

such as   canopy cover (Fornoff et al. 2021, Perlík et al. 2023). Understanding how these forces

respond to changing biodiversity is therefore imperative to predict how ongoing environmental

changes will impact the functioning and stability of ecosystems (Hines et al. 2019).

A crucial network that unites bottom-up and top-down processes in many ecosystems and is

prone to strong alterations due to environmental change is the interaction network between

parasitoids and their hosts (Tylianakis et al. 2006, Jeffs and Lewis 2013). Insect parasitoids attack

and feed on and eventually kill their insect hosts (Godfray and Godfray 1994). Parasitoids are

thought to be particularly sensitive to environmental changes, because species in higher trophic

levels usually have smaller population sizes and the potential specialization of their hosts may

cascade up to impact the parasitoids' presence or absence. Therefore, insect host-parasitoid

systems are ideal for studying the relationships between habitat-level changes and species

interactions (Jeffs and Lewis 2013). Previous studies on host-parasitoid interactions mainly

focused on the influence of abiotic factors, such as elevation and habitat structure (e.g. Valladares

et al. 2012, Maunsell et al. 2015, Grass et al. 2018), or sometimes interaction structure (e.g.

Cagnolo et al. 2011). However, the role of multiple components of plant diversity in modifying

host-parasitoid interaction networks, as key biotic determinants of overall ecosystem structure,

remains relatively little explored. Recent studies mainly focused on basic diversity associations (Ebeling et al. 2012, Schuldt et al. 2019, Guo et al. 2021). These studies have demonstrated both direct and indirect effects of plant diversity on both hosts and parasitoids, possibly via increased niche space and resource availability (Guo et al. 2021). Nevertheless, how these patterns propagate to their interaction networks is still unclear. Moreover, the effects of changing plant diversity are not always obvious when looking at plant species richness, and other diversity components such as plant phylogenetic diversity have been shown to better predict diversity-dependent bottom-up effects on host-parasitoid networks (e.g. Staab et al. 2016). It is especially important to take phylogenetic dependences into account when it comes to the phylogenetic structure within and across trophic levels (Webb et al. 2002, Emerson and Gillespie 2008). This makes it vital to account for multiple dimensions of biodiversity and relevant trophic interactions (e.g. taxonomic, phylogenetic, functional; Peralta et al. 2015, Volf et al. 2017, Wang et al. 2020). These components might jointly affect host-parasitoid networks in a system with high species diversity. Forests have garnered special attention lately, because they represent complex and large ecosystems susceptible to global change (add citation here). Understanding how such relationships modulate the effects of tree diversity loss on the structure and interaction strength of host-parasitoid networks clearly requires further study (Staab et al. 2016, Fornoff et al. 2019).

Here, we leverage standardized trap nests for solitary cavity-nesting bees, wasps, and their parasitoids in a large-scale subtropical forest biodiversity experiment to test how multiple dimensions of tree diversity and community composition influence host-parasitoid network structure. A multi-faceted approach is particularly important when considering that associations between trophic levels might be non-random and phylogenetically structured (Volf et al. 2018, Wang et al. 2020). We aimed to quantify the strength and stability of associations between hosts and parasitoids to both discern the primary components of the diversity and composition of tree communities that affect higher trophic level interactions We expected that (a) multiple tree community metrics, such as taxonomic, phylogenetic and functional diversity and phylogenetic and species community composition can structure host and parasitoid community compositions, especially via phylogenetic processes, as species interactions often show phylogenetic

conservatism (e.g. Pellissier et al. 2013, Peralta et al. 2015). Further, we hypothesized that (b)

host-parasitoid networks will be more complex and stable with increasing tree species richness

due to potential links from higher richness of hosts and parasitoids promoted by more tree species,

and (c) both, community and interaction network changes can also be related to abiotic factors,

such as canopy cover, which might play a role in structuring hymenopteran communities (Haddad

et al. 2011, Fornoff et al. 2021). By better understanding these dynamics, we can begin building a

generalized framework for understanding host-parasitoid interactions in forest ecosystems.

Results

Overall, 34,398 brood cells were collected from 13,267 tubes across five years of sampling

(2015, 2016, 2018, 2019, and 2020). Six families of hosts and seventeen families of

parasitoids were identified. Among them, we found 56 host species (12 bees and 44 wasps)

and 50 parasitoid species (38 Hymenoptera and 12 Diptera). The full species list and their

abundances are given in Table S1.

Community composition of hosts and parasitoids

Host species composition was significantly related to the species and phylogenetic

composition of the tree and parasitoid communities (NMDS scores), as well as to canopy

cover, tree MPD, elevation, and eastness (Fig. 1a, Table 1, Table S2). Parasitoid species

composition was significantly associated with host phylogenetic diversity, tree functional

diversity, tree MPD, eastness, and elevation, and was significantly related to tree species

composition, host species composition, and canopy cover (Fig. 1b, Table 1, Table S3). Host

phylogenetic composition was affected by tree species composition, tree MPD, tree functional

diversity, canopy cover, eastness, elevation and was especially obviously affected by

parasitoid species and phylogenetic composition (Fig. 1c, Table 1, Table S4). For parasitoid

phylogenetic composition, significant relationships were found with tree species and

phylogenetic composition, host species composition, tree functional diversity, canopy cover,

elevation, and eastness (Fig. 1d, Table 1, Table S5).

An effect of host composition on the composition of the parasitoid communities was

further indicated by a significant parafit test (p = 0.032), suggesting nonrandom associations

in the phylogenetic structure of parasitoid and host communities (Fig. 2, Fig. S3).

Host-parasitoid network associations

The linear regression model results showed that host vulnerability, linkage density, and

robustness of parasitoids were significantly negatively related to tree species richness, while

remaining unaffected by other environmental covariables (Table 2, Fig. 3; except for

elevation, which was marginally significantly related to robustness). Interaction evenness was

significantly negatively associated with canopy cover, and interaction evenness was also

negatively related to eastness (Fig. 4c; Table 2).. Generality was only marginally associated

with canopy cover, and was not related to tree species richness or the other environmental

factors. In the alternative models (tree species richness replaced by tree MPD), vulnerability

and linkage density were significantly positively related to tree MPD (Fig. 4a, 4b; Table S6),

while parasitoid robustness was negatively related to tree MPD (Fig. S4, Table S6). The

results of other network metrics (generality and interaction evenness) were consistent with

those of the primary models

The results of the null model analysis suggested that our metrics calculated by the

observed network were significantly different from a random distribution (72, 71, and 77 out

of 85 values for generality, vulnerability, and linkage density, respectively; all values for

robustness, and interaction evenness), strongly demonstrating that interactions between

species were not driven by random processes.

Discussion

Our study demonstrates that tree species richness and phylogenetic diversity play key roles in

modulating communities of hosts and parasitoids and their interactions with each other. These

interactions are further structured by the phylogenetic associations between hosts and

parasitoids. Moreover, canopy cover partly determined host-parasitoid networks, supporting a

recent finding that the structure of host-parasitoid networks is also mediated by changes in

microclimate to some extent (Fornoff et al. 2021). These patterns were highly associated with

multiple tree diversity metrics (tree taxonomic, phylogenetic and functional diversity), and

compositional changes which are key to understanding how host-parasitoid interactions may

be impacted by biodiversity loss through trait- and phylogeny-based processes.

Community composition associations

Host species composition was influenced by several factors, including the taxonomic and

phylogenetic composition of trees and parasitoids, tree diversity (species richness and MPD),

and other biotic factors. The effects of tree diversity and composition on host species

composition agree with previous studies where solitary bee and wasp species composition

were related to plant community structure (e.g. Loyola and Martins 2008). It seems likely that

these results are based on bee linkages to pollen resources and predatory wasp linkages to the

diverse of food sources, which may themselves be closely linked to tree species richness

(Reitalu et al. 2019, Staab and Schuldt 2020).

We also found that tree MPD, FD, and species composition affect parasitoid species

composition, as many studies have found significant relationships between plant and

parasitoid diversity in forests, including tree phylogenetic diversity (Staab et al. 2016),

functional diversity (Rodriguez et al. 2019), and structural diversity (Schuldt et al. 2019).

Similar to predators, parasitoids might also be more active or efficient with increasing tree

community dissimilarity due to higher prey resources or lower intraguild parasitism caused by

more diverse habitats (Finke and Denno 2002). On the other hand, our results also show that

host species composition and parasitoid species composition relate to each other and their

phylogenetic compositions, which are structured by tree communities to some extent. This

pattern could propagate to the adjacent two higher trophic interactions through both top-down

and bottom-up control.

Both host and parasitoid phylogenetic composition were related to tree species

composition. This pattern has important implications for cascading effects among trophic

levels, in that producer communities could structure a higher trophic level community via an

intermediate trophic level. As previous studies usually found weak effects of plants on higher

trophic levels (e.g. Cappelli et al. 2022).   However, while both host and parasitoid

phylogenetic composition was related to tree MPD, only parasitoids responded to tree

phylogenetic composition. This may be because there are many caterpillar-hunting wasps in

our host communities, and the community composition of caterpillars were usually correlated

with tree phylogenetic communities(Wang et al. 2019). Therefore, the prey highly associated

with tree phylogenetic composition (e.g. caterpillars) might indirectly determine predatory

wasp (host) phylogenetic composition, as recently found for plants-caterpillars-spiders (Chen

et al. 2023). This could be further tested by collecting the food directly used by the wasps

(caterpillars). For parasitoids, tree phylogenetic composition might drive the process of

community assembly through trophic cascades (e.g. from plants to parasitoids via herbivores

and host wasps) (Webb et al. 2002, Cavender-Bares et al. 2009). Additionally, parasitoid

phylogenetic composition can be influenced by tree structural diversity (e.g. host availability

in plots with higher heterogeneity; Schuldt et al. 2019), which can be determined by

conserved traits across phylogenies (Webb et al. 2002). The phylogenetic associations

between hosts and parasitoids exhibited a nonrandom structure (significant *parafit* correlation)

between the phylogenetic trees of the host and their parasitoids (see also Peralta et al. 2015).

Interestingly, we found that only host phylogenetic composition was affected by parasitoid

phylogenetic composition (and not vice-versa). This asymmetry suggests that top-down

control (parasitoids to hosts) was stronger than bottom-up control (hosts to parasitoids),

supporting prior hypotheses, demonstrating the strong control of parasitoids (Vidal and

Murphy 2018). This pattern might reflect disproportionate interactions between hosts and

parasitoids, with one trophic level more strongly influencing the other in some instances, but

further study is needed.

Moreover, the species and phylogenetic composition of hosts and parasitoids was also

related to abiotic factors, especially to canopy cover, which has been considered especially

important (Sobek et al. 2009, Fornoff et al. 2021). In future studies, it will be useful to

incorporate other, more direct metrics of microclimate, such as local temperature and

humidity, to determine the proximal drivers of these microclimatic effects (Ma et al. 2010,

Fornoff et al. 2021).

Community-level host-parasitoid networks

Tree community species richness did not significantly influence the diversity of hosts targeted

by parasitoids (generality), but caused a significant increase in   the diversity of parasitoids

per host species (vulnerability). This is likely because niche differentiation often influences

network specialization via potential higher resource diversity in plots with higher tree

diversity (Lopez-Carretero et al. 2014). For the significant relationship between vulnerability

and tree species richness, a potential explanation is that host-parasitoid interactions could be

driven through bottom-up effects. Moreover, higher trophic levels will particularly benefit

from tree diversity, inducing parasitoid species increases more than host diversity with

increasing tree species richness (Guo et al. 2021). These will further increase vulnerability at

community level. According to the enemies hypothesis (Root 1973), which posits a positive

effects of plant richness on natural enemies, the higher trophic levels in our study (e.g.

predators and parasitoids) would benefit from tree diversity and regulate herbivores thereby

(Staab and Schuldt 2020). Indeed, previous studies at the same site found that bee parasitoid

richness and abundance were positively related to tree species richness, but not their bee hosts

(Fornoff et al. 2021, Guo et al. 2021). Because our dataset considered all hosts and reflects an

overall pattern of host-parasitoid interactions, the effects of tree species richness on generality

might be more complex and difficult to predict, as we found that neither tree species richness

nor tree MPD were related to generality. Thus, our results again indicate that top-down

control imposed by higher trophic levels is more significant than bottom-up control with an

increasing number of plant species.

Linkage density was positively related to tree species richness, supporting the food web

theory, which predicts in our case that network complexity (linkage density) depends on the

number of plants (Melián and Bascompte 2002). Although trees were not directly included as

a trophic level in our networks, potential network complexity increased with tree species

richness, likely enabling higher network stability/resistance (Ebeling et al. 2011). For

example, a network might be more sensitive to extinctions because of key species loss due to

lower linkage density and lower redundancy (e.g. Naeem and Li 1997). However, parasitoid

robustness was unexpectedly negatively related to tree species richness. It was expected that

higher trophic levels would be more robust, less influenced by perturbations from lower

trophic levels, when plant diversity is higher, as more potential interactions at lower trophic

levels should theoretically increase redundancy and resilience of connected higher levels

(Blüthgen and Klein 2011, Fornoff et al. 2019). Dilution effects may explain this, as plots

with higher richness held fewer individuals of a given tree species. If there are strong prey

item (caterpillars, grasshoppers, etc.) preferences for one species, there may be fewer or they

may be more densely aggregated and less likely to be encountered by parasitoids. This

increased stochasticity in parasitoid wasps could benefit hosts by reducing parasitism pressure

overall, weakening top-down controls.

Similar to tree species richness, tree MPD was also positively correlated with

vulnerability and linkage density, meaning that the mean number of parasitoids per host

species and number of links within the host-parasitoid system can also be promoted by tree

MPD, in agreement with several recent studies (Pellissier et al. 2013, Staab et al. 2020, Wang

et al. 2020). Our results suggest that the specialization and complexity of higher trophic levels

can also be affected by plant phylogenetic diversity. This pattern can be traced to the effects

of habitat heterogeneity caused by tree species richness and MPD on higher trophic levels via

bottom-up control. The effects of tree MPD were consistent with effects of tree species

richness on robustness of parasitoids to host loss. This result suggests that higher trophic

levels are sensitive to changes in both plant phylogenetic relatedness and general species

dissimilarity via trophic interactions, even the hosts are not all directly interacting with plants,

bees excluded. Therefore, it may be that stronger linkages would be found when exclusively

exploring such plant-herbivore-parasitoid systems.

Interaction evenness was significantly negatively related to canopy cover, further

reinforcing an important role of microclimate (likely temperature and humidity; (Sobek et al.

2009, Fornoff et al. 2021). Our results agree with a previous study on ants, where plant-insect

interactions were more even with more open canopies (Dáttilo and Dyer 2014). In our case,

canopy cover might change hymenopteran species evenness and then further influence

interaction evenness. certain host species tended to nest in plots with higher canopy cover,

which might decrease the interaction evenness by favoring parasitoids of fewer, more

dominant hosts. This pattern would become more significant when more host and parasitoid

species are in a plot, given the positive relationship between higher trophic level diversity and

canopy cover.

Future prospects

Overall, our study enables new insights into the dynamics of host-parasitoid interactions

under varying environmental conditions, an important step toward building a synthetic model

for such biodiversity. A key finding was that although parasitoids and hosts respond to tree

species richness, top-down control seems predominant in parasite-host interaction, though

whether this holds for others antagonistic interactions requires further investigation.

Different trophic levels and functional groups of species responded differently to lower level

changes (Fornoff et al. 2021, Guo et al. 2021). This highlights the complexities of building

such models and calls for more studies across habitat types and taxa, to test the generality of

our findings. Future studies should also consider the role of host/parasitoid functional traits,

because they might play a critical role in modifying network structures and ecosystem

functioning.

Materials and methods

Study sites design

This study was conducted in the BEF-China biodiversity experiment, which is the largest tree diversity experiment worldwide. The experiment is located in a subtropical forest near Xingangshan, Jiangxi province, south-east China (29°08′–29°11′N, 117°90′–117°93′E). The mean annual temperature is 16.7°C and mean annual precipitation 1821 mm (Yang et al. 2013). The experiment includes two study sites (Site A and Site B), 4 km apart from each other, that were established in 2009 (Site A) and 2010 (Site B) respectively. A total of 566 plots (25.8×25.8 m) were designed on the two sites, and per plot 400 trees were initially planted in 20 rows and 20 columns with a planting distance of 1.29 m. A tree species richness gradient (1, 2, 4, 8, 16 and 24 species) was established at each site, based on a species pool of 40 local, broadleaved tree species (Bruelheide et al. 2014).

For our study, at both sites (site A and site B) eight plots of each tree species richness level (1, 2, 4, 8) were randomly selected, as well as six and two plots of 16 and 24 mixtures. In addition, at site B eight additional monocultures were sampled (Fornoff et al. 2021), resulting 48 plots in Site B (including 16 monocultures, eight plots for each 2, 4, 8 mixtures and six and two plots of 16 and 24 mixtures. In total, 88 study plots were used (40 plots on Site A and 48 plots on Site B, see Fig. S1).

Sampling

We collected trap nests monthly to sample solitary bees and wasps (Staab et al. 2018) in the 88 plots from September to November in 2015 and April to November in 2016, 2018, 2019 and 2020. For each plot, we installed two poles with trap nests (11 m apart from each other and 9 m away from the nearest adjacent plots) along a SW–NE diagonal centrally per plot (following the design of Ebeling et al. 2012). Each pole stood 1.5 m above ground, and each trap nest consisted of two PVC tubes (length: 22 cm × diameter: 12.5 cm) filled with 75 ± 9 (SD) reed internodes of 20 cm length and diameters varying between 0.1 and 2.0 cm (Staab et al. 2014, Fornoff et al. 2021).

Every month, we sampled the reeds with nesting hymenopterans and replaced them with

internodes of the same diameter. All the samples were reared in glass test tubes under ambient

conditions until specimens hatched. We identified hatched hosts and parasitoids to species or

morphospecies (Supplementary Table S1) based on reference specimens (vouchered at the

Institute of Zoology, CAS, Beijing). We were interested in the general patterns of host-parasitoid

interactions at the community level, so for the analysis we did not distinguish between the two

life-history strategies of parasitoids (true parasitoids and kleptoparasitoids, including

hymenopteran and dipteran parasitoids) because they both have the same ecological result, death

of host brood cells.

DNA extraction and amplification

All specimens were sequenced for a region of the mitochondrial cytochrome c oxidase subunit I

(COI) gene (Hebert et al. 2003). We extracted whole-genomic DNA of hosts and parasitoids using

DNeasy Blood & Tissue Kits (QIAGEN GmbH, Hilden, Germany), following the manufacturer's

protocols. COI sequences of samples were amplified using universal primer pairs, LCO1490

(GGTCAACAAATCATAAAGATATTGG) as the forward primer and HCO2198

(TAAACTTCAGGGTGACCAAAAAATCA) or HCOout

(CCAGGTAAAATTAAAATATAAACTTC) as the reverse primer. We carried out polymerase

chain reactions (PCR) in 96-well plates with 30 μl reactions containing 10 μl ddH2O, 15 μl

Premix PrimeSTAR HS (TaKaRa), 1 ul of each primer at 10 μM, and 3 μl template genomic DNA

using a thermo cycling profile. The PCR procedure as follows: 94℃ for 1 min; 94℃ for 1 min, 45℃

for 1.5 min and 72℃ for 1.5 min, cycle for 5 times; 94℃ for 2 min, 58℃ for 1.5 min and 72℃ for

1 min, cycle for 36 times; 72℃ for 5 min. We performed all PCRs on an Eppendorf Mastercycler

gradient, which were then visualized on a 1% agarose gel. Samples with clean single bands were

sequenced after PCR purification using BigDye v3.1 on an ABI 3730xl DNA Analyser (Applied

Biosystems).

Sequence alignment and phylogenetic analysis

We applied MAFFT (Misawa, Katoh, Kuma, & Miyata, 2002) to align all sequences, then

translated the nucleotides into amino acids via MEGA v7.0 (Kumar, Stecher, & Tamura, 2016) to

check for the presence of stop codons with manual adjustments. Host and parasitoid sequences

were then aligned against the references using a Perl-based DNA barcode aligner (Chesters,

2019).

We employed two strategies to improve the phylogenetic structure of a DNA barcode

phylogeny, which demonstrably improve resulting phylogeny-based diversity indices

(Macías-Hernández et al. 2020). These include the integration of 1) molecular sequences of the

plot data and 2) phylogenetic relationships from other molecular datasets. Integration was

achieved following Wang et al. (2020) and Chesters (2020): reference DNA barcodes of

Hymenoptera and Diptera were downloaded from the BOLD API

(www.boldsystems.org/index.php/API_Public), which were variously processed (e.g. to retain

only fully taxonomically labelled barcodes, to remove low quality or mislabeled entries, and to

dereplicate to a single exemplar per species), and then aligned (Chesters 2019). A single outgroup

was included for which we selected the most appropriate insect order sister to Diptera and

Hymenoptera (Misof et al. 2014), a representative of the order Psocoptera (Psocidae, *Psocus*

leidyi). We then constructed a phylogeny of the references and subjects, with references

constrained according to the method described earlier (Chesters 2020). A number of backbone

topologies were integrated for setting hard and soft constraints, including a

transcriptomics-derived topology (Chesters 2020), a mitogenome tree of insects (Chesters 2017),

Diptera-specific trees (Wiegmann et al. 2011, Cranston et al. 2012, Ament 2017) and

Hymenoptera-specific trees (Peters et al. 2011, Branstetter et al. 2017, Cardinal 2018). The

constrained inference was conducted with RaxML version 8 (Stamatakis 2014) under the standard

GTRGAMMA DNA model with 24 rate categories. According to the backbone trees used, most

taxa present were monophyletic with a notable exception of Crabronidae, for which there is

emerging phylogenomic evidence of its polyphyly (Sann et al. 2018).

Tree phylogenetic diversity, functional diversity, and environmental covariates

The phylogenetic diversity of the tree communities was quantified by calculating wood

volume-weighted phylogenetic Mean Pairwise Distance (MPD) (Tucker et al. 2017). Tree

wood volume was estimated from data on basal area and tree height (Bongers et al. 2021)

measured in the center of each plot. Moreover, to represent variations towards the tips of the

phylogeny beyond MPD, we additionally calculated Mean Nearest Taxon Distance (MNTD),

which is a measure that quantifies the distance between each species and its nearest neighbor

on the phylogenetic tree (Webb 2000). Phylogenetic metrics of trees were calculated based on

a maximum likelihood phylogenetic tree available for the tree species in our study area

(Purschke et al. 2017). Considering that predatory wasps mainly feed on herbivorous

caterpillars, we calculated tree functional diversity to test the indirect effects on

hymenopteran communities and relevant network indices. Specifically,   seven leaf traits

were expressed as Rao's Q (Ricotta and Moretti 2011), including specific leaf area, leaf

toughness, leaf dry matter content, leaf carbon content, ratio of leaf carbon to nitrogen, leaf

magnesium content, and leaf calcium content. These functional traits were commonly related

to higher trophic levels in our study area (Wang et al. 2020, Chen et al. 2023), which are the

main food resources of our predatory wasps. All of the traits were measured on pooled

samples of sun-exposed leaves of a minimum of five tree individuals per species following

standard protocols (Pérez-Harguindeguy et al. 2003).

As our analyses mainly compare community patterns among study plots, we additionally

considered potential effects of environmental variation by using plot means of slope,

elevation, "eastness" (sine-transformed radian values of aspect), and "northness"

(cosine-transformed radian values of aspect) as environmental covariates that characterize the

heterogeneity of the study plots. We also accounted for the potential effects of canopy cover

at plot level for host-parasitoid interactions, as it can structure hymenopteran communities

(Perlík et al. 2023). Canopy cover was calculated as in Fornoff et al. (2021) based on

above-nest images.

Statistical analysis

All analyses were conducted in R 4.1.2 with the packages *ape*, *vegan*, *picante*, *bipartite*, and *caper*

(http://www.R-project.org). Prior to analysis, samples from the five years (2015, 2016, 2018, 2019,

and 2020) were pooled at the plot level to discern overall and generalizable effects permeating this

system. We excluded three plots with no living trees because of high mortality, resulting in 85

plots in the final analysis.

Composition of trees, hosts and parasitoids

The species and phylogenetic composition of trees, hosts, and parasitoids were quantified at

each plot with nonmetric multidimensional scaling (NMDS) analysis based on Morisita-Horn

distances. The phylogenetic composition was calculated by mean pairwise distance among the

host or parasitoid communities per plot with the R package "*picante.*" To test the influence of

study plot heterogeneity on these relationships, we fitted their standardized values to the

ordination on the basis of a regression with the NMDS axis scores (Quinn and Keough 2002).

For the analysis, we considered tree species richness, tree functional and phylogenetic

diversity, canopy cover, and environmental covariates (elevation, eastness, northness, and

slope) as plot characteristics. We assessed the significance of correlations with permutation

tests (permutation: *n*= 999).

Phylogenetic match of hosts and parasitoids

In addition, we used a parafit test (9,999 permutations) with the R package "*ape*" to test

whether the associations were non-random between hosts and parasitoids. The species that

were not attacked by parasitoids or failed to generate sequences were excluded from the

analyses. For species abundance and composition, see Table S1.

Host-parasitoid interactions

We constructed quantitative host-parasitoid networks with the R package "*bipartite*" for the

level of host and parasitoid communities across for each plot of the two sites. All hosts were

grouped because there were too few abundant bee species to separate them out from other

hosts. We calculated six indices to quantitatively characterize the structure of the interaction

networks: weighted generality (effective number of host species per parasitoid species),

weighted vulnerability (Effective number of parasitoid species attacking a host species),

robustness (degree of network stability), linkage density (degree of network specialization),

and interaction evenness (degree of network evenness). Generality was defined as the

weighted mean number of host species per parasitoid species, $G_{qw} = \sum_{j=1}^{J} \frac{A_j}{m} 2^{H_j}$, with A_j

being the number of interactions of parasitoid species j, m the total number of interactions of

all species, and H_j the Shannon diversity of interactions of species j. Vulnerability was the

weighted mean number of parasitoid species per host species, $\text{Vulnerability} = \sum_{i=1}^{I} \frac{A_i}{m} 2^{H_i}$

(Bersier et al. 2002). Robustness was defined as the area under the extinction curve, reflecting

the degree of decreases of one trophic level with the elimination of the other trophic levels,

here using the robustness index for higher trophic levels (i.e. parasitoids). For linkage density,

$L_q = 0.5 \left(\sum_{j=1}^{J} \frac{A_j}{m} 2^{H_j} + \sum_{i=1}^{I} \frac{A_i}{m} 2^{H_i} \right)$, we used the realized proportion of possible links

between the two trophic levels as the mean number of interactions per species across the

entire network (Tylianakis et al. 2007). Interaction evenness was defined as $E_S =$

$-\sum_i \sum_j p_{ij} \ln p_{ij} / \ln IJ$, which is used to describe Shannon's evenness of network interactions

(Dormann et al. 2009). To check whether all network indices significantly differ from chance

across all study plots, we used Patefield null models (Dormann et al. 2009) to compare

observed indices with simulated values (10,000 times).

Linear mix-effects models To test the effects of tree species richness, tree phylogenetic, and

functional diversity, as well as canopy cover and the other environmental covariates

(including slope, elevation, eastness, and northness) on the six network indices, we used

linear mixed-effect models. For our analyses, the study sites were considered as random

effects, and the others were treated as fixed effects (see above). Given the strong correlation

between tree species richness and tree MPD (Pearson's r = 0.74, p<0.01), we excluded tree

MPD in the models where tree species richness was a predictor. To evaluate the potential

effects caused by tree MPD, we also ran alternative models where tree species richness was

replaced with tree MPD. We simplified all models by gradually removing non-significant

factors to obtain the most parsimonious model with the lowest AICc.   To ensure that the

analyses were not strongly affected by multicollinearity, the correlations among all predictors

were tested (Fig. S2), and variance inflation factors (VIF) of our statistical models were checked.

Acknowledgements

We thank the BEF-China consortium for support (especially, Bo Yang). We thank Mr.

Yinquan Qi for his help with the collection. This work was supported by the National Key

Research Development Program of China (2022YFF0802300), the National Science Fund for

Excellent Young Scholars (32122016), the Strategic Priority Research Program of the

Chinese Academy of Sciences (XDB310304), the National Natural Science Foundation,

China (32100343, 32070465), and the National Science Fund for Distinguished Young

Scholars (31625024). MQW was supported by the Alexander von Humboldt research

fellowships. CDZ's lab is funded by the Key Program of the National Natural Science

Foundation of China (No. 32330013) and also has been continuously supported by grants

from the Key Laboratory of the Zoological Systematics and Evolution of the Chinese

Academy of Sciences (grant number 2008DP173354).

Table 1 Environmental correlates of dissimilarity matrixes with predictors (NMDS on

Morisita-Horn dissimilarity) across the study plots. Significant P-values are indicated in bold.

See Table S2-S5 for the complete information.

	Host species community	Parasitoid species community	Host phylogenetic community	Parasitoid phylogenetic community
Tree phylogeny NMDS1	0.225	0.422	0.386	0.274
Tree phylogeny NMDS2	**0.003**	0.12	0.128	**0.024**
Tree composition NMDS1	**0.001**	**0.001**	**0.001**	**0.001**
Tree composition NMDS2	0.604	0.418	0.433	**0.031**
Canopy cover	**0.001**	**0.001**	**0.001**	**0.004**
Tree species richness	**0.035**	0.122	0.100	0.094
Elevation	**0.005**	**0.007**	**0.001**	**0.001**
Eastness	0.079	0.045	**0.04**	**0.001**
Northness	0.49	0.837	0.821	0.340
Slope	**0.031**	0.507	0.507	0.959
Tree FD (Rao's Q)	0.094	**0.019**	**0.021**	**0.031**
Tree MPD	**0.005**	**0.021**	**0.013**	0.223
Host phylogeny NMDS1	-	**0.016**	-	0.584
Host phylogeny NMDS2	-	**0.027**	-	0.914
Host composition NMDS1	-	**0.001**	-	**0.008**
Host composition NMDS2	-	0.169	-	0.138
Parasitoid phylogeny NMDS1	**0.001**	-	**0.001**	-
Parasitoid phylogeny NMDS2	0.462	-	0.058	-
Parasitoid composition NMDS1	**0.001**	-	**0.001**	-
Parasitoid composition NMDS2	**0.014**	-	**0.001**	-

501 Table 2. Summary results of linear mix-effects models for generality, vulnerability,

502 robustness, linkage density, and interaction evenness of host-parasitoid network indices at the

503 community level across the tree species richness gradient. Standardized parameter estimates

504 (with standard errors, t and P values) are shown for the variables retained in the minimal

505 models.

		Est.	SE	t	P
Generality	Intercept	0.176	0.016	10.96	<0.001
	Canopy cover	0.033	0.016	2.03	0.046
Vulnerability	Intercept	2.860	0.197	14.49	0.044
	Tree species richness	0.180	0.079	2.29	0.025
Robustness of parasitoids	Intercept	0.630	0.007	85.15	<0.001
	Tree species richness	-0.021	0.007	-2.77	0.007
Linkage density	Intercept	2.033	0.093	21.83	0.029
	Tree species richness	0.106	0.040	2.68	0.009
Interaction evenness	Intercept	0.511	0.009	59.12	0.025
	Canopy cover	-0.037	0.007	-5.06	<0.001
	Eastness	-0.018	0.007	-2.50	0.015

Fig. 1. Ordination plot of the non-metric multidimensional scaling (NMDS) analysis

of (a) host taxonomic composition, (b) parasitoid taxonomic composition, (c) host

phylogenetic composition, and (d) parasitoid phylogenetic composition across the

study plots (filled circles) in the BEF-China experiment. Stress = 0.23, 0.23, 0.24 and

0.20, respectively. Arrows indicate significant (at $p < 0.05$) correlations of

environmental variables with NMDS axis scores. Lengths of arrows are proportional

to the strength of the correlations. Red crosses refer to the host or parasitoid species in

each community. See Table S2-S5 in the Supplementary Material for abbreviations

and statistical values.

Fig. 2. Dendrogram of phylogenetic congruence for the host species (below) and associated parasitoid species (above) recorded in the study. Each rectangle represents a different superfamily (for host species) or family (for parasitoid species). H1: Pompilidae H2: Apoidea H3: Vespidae; P1: Sarcophagidae P2: Phoridae P3: Bombyliidae P4: Trigonalyidae P5: Mutillidae P6: Megachilidae P7: Chrysididae P8: Ichneumonidae P9: Chalcidoidea. The trophic network of hosts and parasitoids was non-randomly structured (parafit test: P = 0.032). Host and parasitoid species names are given in Fig. S3.

Fig. 3. Community-level relationships of network between tree species richness and (a) vulnerability, (b) linkage density, and (c) robustness of parasitoids. Values were adjusted for covariates of the final regression model. Regression lines (with 95% confidence bands) show significant (p < 0.05) relationships. Note that axes are on a log scale for tree species richness.

Fig. 4. Community-level relationships of network between tree mean pairwise phylogenetic
distance and (a) vulnerability and (b) linkage density and community-level relationships of
network between canopy cover and (c) interaction evenness. Values were adjusted for
covariates of the final regression model. Regression lines (with 95% confidence bands) show
significant ($p < 0.05$) relationships.

References

Ament, D. C. 2017. Phylogeny of Phorinae sensu latu (Diptera: Phoridae) inferred from a
morphological analysis with comprehensive taxon sampling and an uncommon method of
character coding. Zoological Journal of the Linnean Society **181**:151-188.
Barnes, A. D., M. Jochum, J. S. Lefcheck, N. Eisenhauer, C. Scherber, M. I. O'Connor, P. de Ruiter, and
U. Brose. 2018. Energy Flux: The Link between Multitrophic Biodiversity and Ecosystem
Functioning. Trends in Ecology & Evolution **33**:186-197.
Bersier, L.-F., C. Banašek-Richter, and M.-F. Cattin. 2002. Quantitative Descriptors of Food-Web
Matrices. Ecology **83**:2394-2407.
Blüthgen, N., and A.-M. Klein. 2011. Functional complementarity and specialisation: The role of
biodiversity in plant–pollinator interactions. Basic and Applied Ecology **12**:282-291.
Bongers, F. J., B. Schmid, H. Bruelheide, F. Bongers, S. Li, G. von Oheimb, Y. Li, A. Cheng, K. Ma, and X.
Liu. 2021. Functional diversity effects on productivity increase with age in a forest
biodiversity experiment. Nature Ecology & Evolution.
Branstetter, M. G., J. T. Longino, P. S. Ward, and B. C. Faircloth. 2017. Enriching the ant tree of life:
Enhanced UCE bait set for genome-scale phylogenetics of ants and other Hymenoptera.
Methods in Ecology and Evolution **8**:768-776.
Brondizio, E. S., J. Settele, S. Díaz, and H. T. Ngo. 2019. Global assessment report on biodiversity and
ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and
Ecosystem Services.
Bruelheide, H., K. Nadrowski, T. Assmann, J. Bauhus, S. Both, F. Buscot, X. Y. Chen, B. Y. Ding, W.
Durka, A. Erfmeier, J. L. M. Gutknecht, D. L. Guo, L. D. Guo, W. Hardtle, J. S. He, A. M. Klein, P.
Kuhn, Y. Liang, X. J. Liu, S. Michalski, P. A. Niklaus, K. Q. Pei, M. Scherer-Lorenzen, T. Scholten,
A. Schuldt, G. Seidler, S. Trogisch, G. von Oheimb, E. Welk, C. Wirth, T. Wubet, X. F. Yang, M. J.
Yu, S. R. Zhang, H. Z. Zhou, M. Fischer, K. P. Ma, and B. Schmid. 2014. Designing forest
biodiversity experiments: general considerations illustrated by a new large experiment in
subtropical China. Methods in Ecology and Evolution **5**:74-89.
Cagnolo, L., A. Salvo, and G. Valladares. 2011. Network topology: patterns and mechanisms in plant-
herbivore and host-parasitoid food webs. Journal of Animal Ecology **80**:342-351.
Cappelli, S. L., N. A. Pichon, T. Mannall, and E. Allan. 2022. Partitioning the effects of plant diversity on
ecosystem functions at different trophic levels. Ecological Monographs **92**.
Cardinal, S. 2018. Bee (Hymenoptera: Apoidea: Anthophila) diversity through time. Insect biodiversity:
science and society **2**:851-867.
Cavender-Bares, J., K. H. Kozak, P. V. Fine, and S. W. Kembel. 2009. The merging of community
ecology and phylogenetic biology. Ecology Letters **12**:693-715.
Chen, J.-T., M.-Q. Wang, Y. Li, D. Chesters, A. Luo, W. Zhang, P.-F. Guo, S.-K. Guo, Q.-S. Zhou, K.-P. Ma,
H. Bruelheide, A. Schuldt, and C.-D. Zhu. 2023. Functional and phylogenetic relationships link
predators to plant diversity via trophic and non-trophic pathways. Proceedings of the Royal
Society B **290**:20221658.
Chesters, D. 2017. Construction of a Species-Level Tree of Life for the Insects and Utility in Taxonomic
Profiling. Syst Biol **66**:426-439.
Chesters, D. 2019. aligner. pl. Computer software available at https://sourceforge.net/projects/dna-
barcode-aligner/.

Chesters, D. 2020. The phylogeny of insects in the data-driven era. Systematic Entomology **45**:540-551.

Cranston, P. S., N. B. Hardy, and G. E. Morse. 2012. A dated molecular phylogeny for the Chironomidae (Diptera). Systematic Entomology **37**:172-188.

Dáttilo, W., and L. Dyer. 2014. Canopy openness enhances diversity of ant–plant interactions in the Brazilian Amazon rain forest. Biotropica **46**:712-719.

De Ruiter, P. C., A.-M. Neutel, and J. C. Moore. 1995. Energetics, patterns of interaction strengths, and stability in real ecosystems. Science **269**:1257-1260.

Dormann, C. F., J. Fründ, N. Blüthgen, and B. Gruber. 2009. Indices, graphs and null models: analyzing bipartite ecological networks. The Open Ecology Journal **2**:7-24.

Ebeling, A., A.-M. Klein, and T. Tscharntke. 2011. Plant–flower visitor interaction webs: Temporal stability and pollinator specialization increases along an experimental plant diversity gradient. Basic and Applied Ecology **12**:300-309.

Ebeling, A., A.-M. Klein, W. W. Weisser, and T. Tscharntke. 2012. Multitrophic effects of experimental changes in plant diversity on cavity-nesting bees, wasps, and their parasitoids. Oecologia **169**:453-465.

Emerson, B. C., and R. G. Gillespie. 2008. Phylogenetic analysis of community assembly and structure over space and time. Trends in Ecology & Evolution **23**:619-630.

Finke, D. L., and R. F. Denno. 2002. Intraguild predation diminished in complex-structured vegetation: implications for prey suppression. Ecology **83**:643-652.

Fornoff, F., A. M. Klein, N. Blüthgen, and M. Staab. 2019. Tree diversity increases robustness of multi-trophic interactions. Proceedings: Biological sciences, The Royal Society **286**:20182399.

Fornoff, F., M. Staab, C.-D. Zhu, and A.-M. Klein. 2021. Multi-trophic communities re-establish with canopy cover and microclimate in a subtropical forest biodiversity experiment. Oecologia **196**:289-301.

Godfray, H. C. J., and H. Godfray. 1994. Parasitoids: behavioral and evolutionary ecology. Princeton University Press.

Gonzalez, A., R. M. Germain, D. S. Srivastava, E. Filotas, L. E. Dee, D. Gravel, P. L. Thompson, F. Isbell, S. Wang, and S. Kéfi. 2020. Scaling-up biodiversity-ecosystem functioning research. Ecology Letters **23**:757-776.

Grass, I., B. Jauker, I. Steffan-Dewenter, T. Tscharntke, and F. Jauker. 2018. Past and potential future effects of habitat fragmentation on structure and stability of plant–pollinator and host–parasitoid networks. Nature Ecology & Evolution **2**:1408-1417.

Grossman, J. J., M. Vanhellemont, N. Barsoum, J. Bauhus, H. Bruelheide, B. Castagneyrol, J. Cavender-Bares, N. Eisenhauer, O. Ferlian, D. Gravel, A. Hector, H. Jactel, H. Kreft, S. Mereu, C. Messier, B. Muys, C. Nock, A. Paquette, J. Parker, M. P. Perring, Q. Ponette, P. B. Reich, A. Schuldt, M. Staab, M. Weih, D. C. Zemp, M. Scherer-Lorenzen, and K. Verheyen. 2018. Synthesis and future research directions linking tree diversity to growth, survival, and damage in a global network of tree diversity experiments. Environmental and Experimental Botany **152**:68-89.

Guo, P.-F., M.-Q. Wang, M. Orr, Y. Li, J.-T. Chen, Q.-S. Zhou, M. Staab, F. Fornoff, G.-H. Chen, N.-L. Zhang, A.-M. Klein, and C.-D. Zhu. 2021. Tree diversity promotes predatory wasps and parasitoids but not pollinator bees in a subtropical experimental forest. Basic and Applied Ecology **53**:134-142.

631   Haddad, N. M., G. M. Crutsinger, K. Gross, J. Haarstad, and D. Tilman. 2011. Plant diversity and the
632   stability of foodwebs. Ecology Letters **14**:42-46.

633   Harvey, E., I. Gounand, C. L. Ward, and F. Altermatt. 2017. Bridging ecology and conservation: from
634   ecological networks to ecosystem function. Journal of Applied Ecology **54**:371-379.

635   Hebert, P. D., S. Ratnasingham, and J. R. deWaard. 2003. Barcoding animal life: cytochrome c oxidase
636   subunit 1 divergences among closely related species. Proceedings: Biological sciences, The
637   Royal Society **270 Suppl 1**:S96-99.

638   Hines, J., A. Ebeling, A. D. Barnes, U. Brose, C. Scherber, S. Scheu, T. Tscharntke, W. W. Weisser, D. P.
639   Giling, A. M. Klein, and N. Eisenhauer. 2019. Chapter Nine - Mapping change in biodiversity
640   and ecosystem function research: food webs foster integration of experiments and science
641   policy. Pages 297-322 *in* N. Eisenhauer, D. A. Bohan, and A. J. Dumbrell, editors. Advances in
642   Ecological Research. Academic Press.

643   Hunter, M. D., M. D. Hunter, T. Ohgushi, and P. W. Price. 1992. Interactions within herbivore
644   communities mediated by the host plant: the keystone herbivore concept.

645   Isbell, F., P. Balvanera, A. S. Mori, J. S. He, J. M. Bullock, G. R. Regmi, E. W. Seabloom, S. Ferrier, O. E.
646   Sala, and N. R. Guerrero-Ramírez. 2022. Expert perspectives on global biodiversity loss and
647   its drivers and impacts on people. Frontiers in Ecology and the Environment.

648   Jeffs, C. T., and O. T. Lewis. 2013. Effects of climate warming on host–parasitoid interactions.
649   Ecological Entomology **38**:209-218.

650   Lopez-Carretero, A., C. Diaz-Castelazo, K. Boege, and V. Rico-Gray. 2014. Evaluating the
651   spatio-temporal factors that structure network parameters of plant-herbivore interactions.
652   PLoS ONE **9**:e110430.

653   Loyola, R. D., and R. P. Martins. 2008. Habitat structure components are effective predictors of
654   trap-nesting Hymenoptera diversity. Basic and Applied Ecology **9**:735-742.

655   Ma, S., A. Concilio, B. Oakley, M. North, and J. Chen. 2010. Spatial variability in microclimate in a
656   mixed-conifer forest before and after thinning and burning treatments. Forest Ecology and
657   Management **259**:904-915.

658   Macías-Hernández, N., M. Domènech, P. Cardoso, B. C. Emerson, P. A. V. Borges, J. Lozano-Fernandez,
659   O. S. Paulo, A. Vieira, A. Enguídanos, F. Rigal, I. R. Amorim, and M. A. Arnedo. 2020. Building a
660   Robust, Densely-Sampled Spider Tree of Life for Ecosystem Research. Diversity **12**:288.

661   Maunsell, S. C., R. L. Kitching, C. J. Burwell, and R. J. Morris. 2015. Changes in host-parasitoid food
662   web structure with elevation. Journal of Animal Ecology **84**:353-363.

663   Melián, C. J., and J. Bascompte. 2002. Food web structure and habitat loss. Ecology Letters **5**:37-46.

664   Misof, B., S. Liu, K. Meusemann, R. S. Peters, A. Donath, C. Mayer, P. B. Frandsen, J. Ware, T. Flouri,
665   and R. G. Beutel. 2014. Phylogenomics resolves the timing and pattern of insect evolution.
666   Science **346**:763-767.

667   Naeem, S., and S. Li. 1997. Biodiversity enhances ecosystem reliability. Nature **390**:507-509.

668   Pellissier, L., C. Ndiribe, A. Dubuis, J. N. Pradervand, N. Salamin, A. Guisan, and S. Rasmann. 2013.
669   Turnover of plant lineages shapes herbivore phylogenetic beta diversity along ecological
670   gradients. Ecology Letters **16**:600-608.

671   Peralta, G., C. M. Frost, R. K. Didham, A. Varsani, and J. M. Tylianakis. 2015. Phylogenetic diversity and
672   co-evolutionary signals among trophic levels change across a habitat edge. Journal of Animal
673   Ecology **84**:364-372.

Pérez-Harguindeguy, N., S. Díaz, F. Vendramini, J. H. C. Cornelissen, D. E. Gurvich, and M. Cabido. 2003.
Leaf traits and herbivore selection in the field and in cafeteria experiments. Austral Ecology
28:642-650.
Perlík, M., D. Kraus, H. Bußler, L. Neudam, S. Pietsch, U. Mergner, D. Seidel, P. Sebek, and S. Thorn.
2023. Canopy openness as the main driver of aculeate Hymenoptera and saproxylic beetle
diversity following natural disturbances and salvage logging. Forest Ecology and
Management **540**:121033.
Peters, R. S., B. Meyer, L. Krogmann, J. Borner, K. Meusemann, K. Schütte, O. Niehuis, and B. Misof.
2011. The taming of an impossible child: a standardized all-in approach to the phylogeny of
Hymenoptera using public database sequences. BMC Biology **9**:1-14.
Purschke, O., S. G. Michalski, H. Bruelheide, and W. Durka. 2017. Phylogenetic turnover during
subtropical forest succession across environmental and phylogenetic scales. Ecology and
Evolution **7**:11079-11091.
Quinn, G. P., and M. J. Keough. 2002. Experimental Design and Data Analysis for Biologists. Cambridge
University Press, Cambridge.
Reitalu, T., A. E. Bjune, A. Blaus, T. Giesecke, A. Helm, I. Matthias, S. M. Peglar, J. S. Salonen, H. Seppä,
and V. Väli. 2019. Patterns of modern pollen and plant richness across northern Europe.
Journal of Ecology **107**:1662-1677.
Ricotta, C., and M. Moretti. 2011. CWM and Rao's quadratic diversity: a unified framework for
functional ecology. Oecologia **167**:181-188.
Rodriguez, A., J. L. Pohjoismäki, and J. Kouki. 2019. Diversity of forest management promotes
parasitoid functional diversity in boreal forests. Biological Conservation **238**:108205.
Root, R. B. 1973. Organization of a Plant-Arthropod Association in Simple and Diverse Habitats: The
Fauna of Collards (Brassica Oleracea). Ecological Monographs **43**:95-124.
Sann, M., O. Niehuis, R. S. Peters, C. Mayer, A. Kozlov, L. Podsiadlowski, S. Bank, K. Meusemann, B.
Misof, and C. Bleidorn. 2018. Phylogenomic analysis of Apoidea sheds new light on the sister
group of bees. Bmc Evolutionary Biology **18**:1-15.
Schuldt, A., A. Ebeling, M. Kunz, M. Staab, C. Guimarães-Steinicke, D. Bachmann, N. Buchmann, W.
Durka, A. Fichtner, F. Fornoff, W. Härdtle, L. R. Hertzog, A.-M. Klein, C. Roscher, J. Schaller, G.
von Oheimb, A. Weigelt, W. Weisser, C. Wirth, J. Zhang, H. Bruelheide, and N. Eisenhauer.
2019. Multiple plant diversity components drive consumer communities across ecosystems.
Nature Communications **10**:1460.
Sobek, S., T. Tscharntke, C. Scherber, S. Schiele, and I. Steffan-Dewenter. 2009. Canopy vs. understory:
does tree diversity affect bee and wasp communities and their natural enemies across forest
strata? Forest Ecology and Management **258**:609-615.
Staab, M., H. Bruelheide, W. Durka, S. Michalski, O. Purschke, C. D. Zhu, and A. M. Klein. 2016. Tree
phylogenetic diversity promotes host-parasitoid interactions. Proceedings of the Royal
Society B: Biological Sciences **283**.
Staab, M., X. Liu, T. Assmann, H. Bruelheide, F. Buscot, W. Durka, A. Erfmeier, A. M. Klein, K. Ma, S.
Michalski, T. Wubet, B. Schmid, A. Schuldt, and J. Koricheva. 2020. Tree phylogenetic
diversity structures multitrophic communities. Functional Ecology **35**:521-534.
Staab, M., M. Ohl, C.-D. Zhu, and A.-M. Klein. 2014. A unique nest-protection strategy in a new species
of spider wasp. PLoS ONE **9**:e101592.

Staab, M., G. Pufal, T. Tscharntke, A. M. Klein, and G. Iossa. 2018. Trap nests for bees and wasps to
analyse trophic interactions in changing environments—A systematic overview and user
guide. Methods in Ecology and Evolution **9**:2226-2239.
Staab, M., and A. Schuldt. 2020. The influence of tree diversity on natural enemies—a review of the
"enemies" hypothesis in forests. Current Forestry Reports **6**:243-259.
Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics **30**:1312-1313.
Tittensor, D. P., M. Walpole, S. L. Hill, D. G. Boyce, G. L. Britten, N. D. Burgess, S. H. Butchart, P. W.
Leadley, E. C. Regan, R. Alkemade, R. Baumung, C. Bellard, L. Bouwman, N. J. Bowles-Newark,
A. M. Chenery, W. W. Cheung, V. Christensen, H. D. Cooper, A. R. Crowther, M. J. Dixon, A.
Galli, V. Gaveau, R. D. Gregory, N. L. Gutierrez, T. L. Hirsch, R. Hoft, S. R. Januchowski-Hartley,
M. Karmann, C. B. Krug, F. J. Leverington, J. Loh, R. K. Lojenga, K. Malsch, A. Marques, D. H.
Morgan, P. J. Mumby, T. Newbold, K. Noonan-Mooney, S. N. Pagad, B. C. Parks, H. M. Pereira,
T. Robertson, C. Rondinini, L. Santini, J. P. Scharlemann, S. Schindler, U. R. Sumaila, L. S. Teh, J.
van Kolck, P. Visconti, and Y. Ye. 2014. A mid-term analysis of progress toward international
biodiversity targets. Science **346**:241-244.
Tucker, C. M., M. W. Cadotte, S. B. Carvalho, T. J. Davies, S. Ferrier, S. A. Fritz, R. Grenyer, M. R.
Helmus, L. S. Jin, A. O. Mooers, S. Pavoine, O. Purschke, D. W. Redding, D. F. Rosauer, M.
Winter, and F. Mazel. 2017. A guide to phylogenetic metrics for conservation, community
ecology and macroecology. Biological Reviews of the Cambridge Philosophical Society
92:698-715.
Tylianakis, J. M., R. K. Didham, J. Bascompte, and D. A. Wardle. 2008. Global change and species
interactions in terrestrial ecosystems. Ecology Letters **11**:1351-1363.
Tylianakis, J. M., T. Tscharntke, and A.-M. Klein. 2006. Diversity, ecosystem function, and stability of
parasitoid–host interactions across a tropical habitat gradient. Ecology **87**:3047-3057.
Tylianakis, J. M., T. Tscharntke, and O. T. Lewis. 2007. Habitat modification alters the structure of
tropical host–parasitoid food webs. Nature **445**:202.
Valladares, G., L. Cagnolo, and A. Salvo. 2012. Forest fragmentation leads to food web contraction.
Oikos **121**:299-305.
Vidal, M. C., and S. M. Murphy. 2018. Bottom-up vs. top-down effects on terrestrial insect herbivores:
a meta-analysis. Ecology Letters **21**:138-150.
Volf, M., P. Pyszko, T. Abe, M. Libra, N. Kotaskova, M. Sigut, R. Kumar, O. Kaman, P. T. Butterill, J.
Sipos, H. Abe, H. Fukushima, P. Drozd, N. Kamata, M. Murakami, and V. Novotny. 2017.
Phylogenetic composition of host plant communities drives plant-herbivore food web
structure. Journal of Animal Ecology **86**:556-565.
Volf, M., S. T. Segar, S. E. Miller, B. Isua, M. Sisol, G. Aubona, P. Simek, M. Moos, J. Laitila, J. Kim, J.
Zima, Jr., J. Rota, G. D. Weiblen, S. Wossa, J. P. Salminen, Y. Basset, and V. Novotny. 2018.
Community structure of insect herbivores is driven by conservatism, escalation and
divergence of defensive traits in Ficus. Ecology Letters **21**:83-92.
Wang, M.-Q., Y. Li, D. Chesters, P. Anttonen, H. Bruelheide, J.-T. Chen, W. Durka, P.-F. Guo, W.
Haerdtle, K. Ma, S. G. Michalski, B. Schmid, G. von Oheimb, C.-S. Wu, N.-L. Zhang, Q.-S. Zhou,
A. Schuldt, and C.-D. Zhu. 2019. Multiple components of plant diversity loss determine
herbivore phylogenetic diversity in a subtropical forest experiment. Journal of Ecology
107:2697-2712.

Wang, M. Q., Y. Li, D. Chesters, H. Bruelheide, K. Ma, P. F. Guo, Q. S. Zhou, M. Staab, C. D. Zhu, and A.
Schuldt. 2020. Host functional and phylogenetic composition rather than host diversity
structure plant-herbivore networks. Molecular Ecology **29**:2747-2762.
Webb, C. O. 2000. Exploring the Phylogenetic Structure of Ecological Communities: An Example for
Rain Forest Trees. Am Nat **156**:145-155.
Webb, C. O., D. D. Ackerly, M. A. McPeek, and M. J. Donoghue. 2002. Phylogenies and Community
Ecology. Annual Review of Ecology and Systematics **33**:475-505.
Wiegmann, B. M., M. D. Trautwein, I. S. Winkler, N. B. Barr, J.-W. Kim, C. Lambkin, M. A. Bertone, B. K.
Cassel, K. M. Bayless, and A. M. Heimberg. 2011. Episodic radiations in the fly tree of life.
Proceedings of the National Academy of Sciences **108**:5690-5695.