Brochosomes as an antireflective camouflage coating for leafhoppers

Wei Wu[1*], Qianzhuo Mao[1], Zhuangxin Ye[1], Zhenfeng Liao[2], Hong-Wei Shan[1,] Jun-Min Li[1],

Chuan-Xi Zhang[1], Jian-Ping Chen[1*]

[1]State Key Laboratory for Managing Biotic and Chemical Threats to the Quality and Safety of

Agro-products, Key Laboratory of Biotechnology in Plant Protection of Ministry of Agriculture

and Zhejiang Province, Institute of Plant Virology, Ningbo University, Ningbo 315211, China.

[2]State Key Laboratory for Sustainable Control of Pest and Disease, Institute of Virology and

Biotechnology, Zhejiang Academy of Agricultural Sciences, Hangzhou, 310021, China.

*Corresponding author. Email: jianpingchen@nbu.edu.cn (J.-P.C.);

wuwei_19861115@163.com (W.W.)

Abstract:

In nature, insects face immense predation pressure, where visual cues play a vital role in predators locating them. To counter this threat, insects employ a variety of nano- and microstructures on their cuticular layer to manipulate and interact with light, enhancing anti-reflective properties and providing camouflage or reducing detectability by predators. Leafhoppers have a unique extra-cuticular coating called brochosome, yet its anti-reflective functions and protein composition remain unclear. Our study demonstrates strong anti-reflective properties of brochosomes, effectively reducing reflectance on the cuticle surface, especially in the ultraviolet spectrum, to improve evasion from visual predators. Furthermore, we identify four novel structural proteins of the brochosome (BSM) for the first time. Inhibiting their synthesis by RNAi alters brochosome morphology, impacting the optical properties of the cuticle surface. Evolutionary origin analysis of BSM suggests that brochosomes likely originated from a process involving duplication–divergence. Our study reveals that leafhoppers employ a unique camouflage strategy by secreting brochosomes as anti-reflection nano-coatings, enabling them to evade natural predators and contributing to their evolutionary success.

Keywords:

Brochosome, Leafhopper, Camouflage, Antireflective, Predation, BSM

Introduction:

Predation exerts significant selective pressure on the evolution of various species, driving the development of anti-predator strategies across the animal kingdom(Galloway et al., 2020; Stevens & Ruxton, 2018). Camouflage is a widely employed strategy, allowing animals to blend into their surroundings by adjusting their coloration and patterns to reduce detection by visual predators(Hughes et al., 2019; Skelhorn & Rowe, 2016). Nevertheless, the presence of predators with multispectral vision, capable of perceiving non-visible light spectra like infrared and ultraviolet, can compromise traditional color and pattern-based camouflage(Galloway et al., 2020; Jones et al., 2007; Stevens, 2007; Thery & Gomez, 2010), paralleling the challenges of military camouflage under non-visible light(Xi et al., 2023). Consequently, achieving alignment between an animal's surface optical characteristics and its surroundings becomes a crucial requirement in visual camouflage strategies.

Insects, one of the most diverse animal groups on earth, contend with intense predation pressure(Cinel et al., 2020; Sheikh et al., 2017). Many of their predators, including birds, reptiles, and predatory arthropods, rely on visual cues, with some possessing highly developed ultraviolet (UV) vision, where UV light plays a critical role in locating insect prey(Lim & Ben-Yakir, 2020; Stobbe et al., 2008). Furthermore, the natural backgrounds where insects reside, such as leaves, tree bark, and soil, exhibit minimal light reflection(Endler, 1997). Consequently, light reflections originating from an insect's body significantly elevate its risk of exposure to predators(Tovee, 1995). To counter this challenge, the various nano- and microstructural features of the insect cuticle have excellent antireflective properties that affect insect body coloration, reduce surface light reflection and aid camouflage(Vukusic & Sambles, 2003; Watson et al., 2017). The first instances of anti-reflective structures were observed in the compound eyes of moths, the corneas of which are covered by hexagonally arranged protrusions that are approximately 200 nanometers apart (Bernhard, 1965; Blagodatski et al., 2015; Clapham & Hutley, 1973). Subsequent research has revealed that anti-reflective structures are widespread on insect body surfaces, primarily found on compound eyes and wings(Blagodatski et al., 2015; Chotard et al., 2022; Ho et al., 2016; Monteiro et al., 2019; Stavenga et al., 2005). These nano- and micro-structures on the cuticular layer are believed to possess anti-reflective properties, effectively minimizing light reflections from their bodies and

thus preventing detection by predators(Blagodatski et al., 2015; Chotard et al., 2022; Ho et al.,

2016; Monteiro et al., 2019; Stavenga et al., 2005).

Leafhoppers (Cicadellidae), one of the largest insect families with over 22,000 species,

possess unique extra-cuticular coating known as brochosome. Brochosomes, typically hollow,

honeycomb-like spheres with a diameter ranging from 0.2 to 0.6 micrometers, are synthesized

in the Malpighian tubules, secreted through the hindgut, and applied as a coating on the fresh

cuticle following molting(Rakitov, 2000; Rakitov, 2009). Brochosomes primarily comprise lipids

and proteins, with protein content ranging from 45% to 75%, but the exact proteins responsible

for brochosome formation remain unknown(Li et al., 2022; Rakitov et al., 2018; Yuan et al.,

2023). Some have proposed that brochosomes may serve as a protective layer on the

leafhopper's cuticle surface, with a potential role in enhancing the hydrophobicity of their cuticle

surface, providing defense against pathogens and predators(Rakitov & Gorb, 2013a; Rakitov,

2009). Apart from these known function of hydrophobicity and the shedding of brochosomes in

helping leafhopper escape spider webs, there is currently no definitive biological evidence

supporting other hypothesized functions of brochosomes(Lin et al., 2021; Rakitov & Gorb,

2013a, 2013b). Furthermore, it has been suggested by certain researchers that the

brochosomes present on the surface of leafhoppers may play a role in their camouflage

abilities(Swain, 1936). In recent years, studies investigating bio-inspired materials that mimic

the structure of brochosomes have demonstrated remarkable antireflective properties(Lei et al.,

2020; Yang et al., 2017), and further elucidated the antireflective mechanisms based on the

brochosome geometry(Wang et al., 2024). These findings offer valuable insights, suggesting

that leafhoppers may employ a distinctive strategy, distinct from other insects, to achieve anti-

reflection properties on their surfaces. Consequently, we propose the hypothesis that

brochosomes may function as a camouflage coating on the leafhopper's cuticle surface, aiding

in the reduction of surface optical reflections and enabling the leafhopper to remain concealed,

thereby evading detection by visual predators and ultimately escaping predation.

In this study, we combined spectrophotometry, electron microscopy, transcriptome analysis,

proteome analysis, gene function validation, and leafhopper and jumping spider bioassays to

characterize the protein composition and the function of brochosome in leafhopper camouflage.

We focused our research on the rice green leafhopper, *Nephotettix cincticeps*, a well-known

agricultural pest with an extensive research history(Hibino, 1996; Wei et al., 2018; Yan et al.,

2021). Our results revealed the pronounced effectiveness of brochosome coverage in reducing

light reflection on leafhoppers *N. cincticeps*, especially within the scope of the ultraviolet

spectrum. By conducting RNAi experiments, we successfully identified four brochosome

structural protein for the first time. The suppression of their synthesis induced changes in

brochosome morphology, influencing the optical characteristics of the leafhopper's cuticle

surface. Furthermore, our analysis of the evolutionary origin of brochosomes indicates that they

likely originated through a process involving duplication–divergence. Our findings support

brochosomes as a unique camouflage coating that enables leafhoppers to evade visual

predators.

Results

Brochosomes are a distinctive coating on the cuticle surface of leafhopper *N. cincticeps.*

The cuticle surface of rice green leafhopper *N. cincticeps* is coated with brochosomes

(Figure 1A and B), which are approximately 350nm in diameter and featuring a hollow sphere

with a honeycomb-like structure (Figure 1C and D). The leafhopper *N. cincticeps* possesses

two pairs of Malpighian tubules, each consisting of a proximal, distal, and terminal segment

(Figure 1E). Brochosomes are synthesized in the distal segment of the Malpighian tubule, which

exhibits a swollen and rod-shaped appearance (Figure 1E). The epithelial cells in this segment

contain large spherical nuclei, an extensive rough endoplasmic reticulum, and multiple Golgi

regions (Figure 1E and F). Within the Golgi-derived vacuole, the brochosomes undergo

progressive development, transforming their initially spherical granules into a honeycomb-like

surface with closely spaced invaginations (Figure 1G-I). Mature brochosomes are stored in

secretory vacuoles at the boundaries of the cells and are subsequently secreted into the lumen

of the Malpighian tubules (Figure 1F).

The optical properties of the leafhopper's cuticle surface are intricately connected to the

brochosome coating.

To investigate the interplay between brochosome coverage and the optical features of *N.*

cincticeps cuticle surface, we observed the relationship for male and female adults at 5, 10, 15,

20, and 25 days post-eclosion. *N. cincticeps* exhibited a gradual shift in body color with

increasing eclosion time, manifesting clear distinctions between males and females (Figure 2A-

figure supplement 1A and B). Males transformed from light green to dark green, while females

progressively transitioned from light green to translucent, displaying an iridescence (Figure 2A-

figure supplement 1A and B). Furthermore, the brightness of the leafhopper cuticle surface

under UV light intensified with the extension of post-eclosion time, and the UV reflectance of

female cuticle surface was significantly higher than that of males after 15 days post-eclosion

(Figure 2B-figure supplement 1A and B).

To validate the correlation between brochosome coverage, body color, and UV light

reflectance, we examined the distribution of brochosomes on the cuticle surface of male and

female *N. cincticeps* at various post-eclosion intervals using scanning electron microscopy

(SEM). SEM results unveiled a gradual reduction in brochosome coverage on the cuticle

surface of both sexes as post-eclosion time progressed. Specifically, male coverage decreased

from 90% to approximately 40%, whereas female coverage descended from 90% to around 10%

(Figure 2C-figure supplement 1A-C). Microscopic analyses of the Malpighian tubule revealed

that, with prolonged post-eclosion time, the distal segment of the tubule underwent gradual

atrophy, particularly pronounced at 15 days post-eclosion. Notably, females exhibited a more

marked atrophy in the distal segment compared to males. By 25 days post-eclosion, the distal

segment of the Malpighian tubule in females became indistinguishable from other segments

(Figure 2D-figure supplement 1A and B). This denotes a progressive reduction in post-eclosion

brochosome synthesis, resulting in a corresponding decrease in brochosome coverage on the

cuticle surface. Furthermore, the distribution of brochosomes on the leafhopper cuticle surface

may be correlated with its optical characteristics. To substantiate this hypothesis, we analyzed

the light reflection characteristics of leafhopper forewings at various post-eclosion time points.

The findings revealed a positive correlation between brochosome coverage on the forewings

and light reflection values, underscoring the remarkable anti-reflective attributes of the

brochosome coating on leafhopper cuticle surface (Figure 2E-figure supplement 1D and E).

Brochosomal coverage was found to significantly decrease the reflectance of ultraviolet light

on the leafhopper's surface, reducing it from approximately 30% to 20%. Additionally, the

reflectance of visible and infrared light was also notably diminished, dropping from around 20%

to 10% (Figure 2E-figure supplement 1D and E). In summary, the presence of brochosomal

coverage resulted in an overall reduction of surface light reflection by approximately 30%,

highlighting its substantial antireflective properties.

Preferential capture of leafhoppers with fewer brochosomes by jumping spiders

Given that UV light serves as a crucial visual cue for arthropod predators, especially

common visual hunters like jumping spiders(Cronin & Bok, 2016; Li & Lim, 2005; Morehouse

et al., 2017; Silberglied, 1979; Zou et al., 2011), the function of the brochosome coating on

leafhoppers' cuticle surface could be to avoid predation by reduction of UV light reflectance.

This prompted us to investigate the effect of brochosome coatings on reducing predation risk

in leafhoppers. Predation experiments were conducted using *Plexippus paykulli* (Figure 2F-

figure supplement 2), a common jumping spider in rice fields known to prey on

leafhoppers(Yang et al., 2018). We observed the spiders' feeding behavior, noting the time of

the first attack and the targeted leafhopper, using these metrics to evaluate predation efficiency

(Taylor et al., 2014; Walker & Rypstra, 2002). In the experimental group comprising females

and males at 5 days post-eclosion, *P. paykulli* did not exhibit a clear predation preference

(Figure 2G). However, in the experimental group involving females and males at 25 days post-

eclosion, there was a higher probability of predation on female *N. cincticeps* compared to male

ones (Figure 2H). For male *N. cincticeps* in both the 5-day and 25-day post-eclosion

experimental groups, *P. paykulli* showed a stronger inclination towards preying on individuals

that were eclosed for 25 days, resulting in significantly shorter predation times compared to

those at 5 days post-eclosion (Figure 2I). These patterns were consistently observed for female

leafhoppers in both the 5-days and 25-days post-eclosion (Figure 2J). The cumulative findings

strongly suggest a distinct preference of the jumping spider *P. paykulli* for individuals with lower

brochosome coverage on the cuticle surface. Based on these experiments, we proposed the

hypothesis that the brochosome coating on the leafhopper cuticle surface may mitigate

predation risk by reducing surface light reflection, particularly in the UV spectrum.

Identifying the major structural proteins of brochosomes.

Despite the initial discovery of brochosomes in the early 1950s and the confirmation in the

1960s that they are protein-lipid particles(Gouranton & Maillet, 1967), the specific protein

composition of brochosomes remains unknown. Recent studies suggested that they are

primarily composed of brochosomins (BSM) and brochosome-associated proteins (BSAP)(Li et

al., 2022; Rakitov et al., 2018). BSM, a novel class of secreted proteins with molecular weights

ranging from 21 to 40 kDa, is considered as the major structural component of

brochosomes(Rakitov et al., 2018). Proteomics studies have identified the proteins composing

Brochosomes, but the specific proteins involved in their formation remain unclear.

To further elucidate the essential protein components of brochosomes, transcriptomic and

proteomic data derived from brochosome of *N. cincticeps* were comprehensively analyzed.

Based on our prior integrated analysis of the brochosome transcriptome and proteome(Wu et

al., 2023), in conjunction with literature search results, we selected 50 candidate genes for

functional analysis using RNAi-mediated gene silencing. Through RNAi experiments, we

successfully identified four genes encoding brochosome structural proteins that led to

morphological changes in brochosomes (Figure 3A). According to the coding sequence (CDS)

length of these genes, the four proteins were named as BSM-1 to BSM-4 (GenBank accession

number PP273097, PP273098, PP273099, PP273100) (Figure 3B). Homology analysis

suggested that BSM-coding genes might be paralogous (Figure supplement 3A). Although

BSM2 and BSM3 exhibited low sequence homology, their protein structures were highly

conserved (Figure supplement 3B-C). To validate the functions of these four BSM-coding genes,

we synthesized dsRNA from two non-overlapping regions of each gene to minimize off-target

effects (Figure supplement 4A). RT-qPCR revealed that both individual and mixed dsRNA

injections effectively suppressed the expression of the BSM-coding genes, with mixed

injections achieving higher efficiency (Figure 3C-F-figure supplement 4C-J). SEM observations

showed that both injection methods induced significant morphological changes in brochosomes,

characterized by increased diameters and deformed honeycomb-like structures (Figure 3A-

figure supplement 4B). Statistical analysis of SEM data from mixed dsRNA injections indicated

a 60-70% reduction in brochosome distribution area and a 20% incidence of morphologically

abnormal brochosomes compared to dsGFP controls (Figure supplement 5A-B). Additionally,

temporal and spatial expression analyses demonstrated that the BSM-coding genes were

specifically expressed in the Malpighian tubules (Figure 3G-J) and exhibited relatively stable

expression during the early post-eclosion period, followed by a gradual decline after 10 days

(Figure 3K-N). By 25 days post-eclosion, the expression of BSM-coding genes declined to

around 10% in females and 30-40% in males, respectively (Figure 3K-N). This phenomenon is

consistent with our initial microscopic observations, suggesting that the gradual reduction in

brochosome synthesis contributes to the decrease of brochosome coverage on the cuticle

surface of *N. cincticeps* after adulthood.

Brochosome coating diminishes light reflection and facilitates predator avoidance

To further investigate the function of brochosomes in leafhopper cuticle surface associated

with light reflection and predator avoidance, we implemented RNAi to simultaneously suppress

the expression of all four BSM-coding genes with ds*BSM* mixture injection. RT-qPCR results

demonstrated a substantial downregulation in the expression of these genes compared to the

ds*GFP* control (Figure 4F). Notably, under ultraviolet light, both male and female leafhoppers

exhibited a significant increase in UV reflection on their cuticle surface following ds*BSM*

treatment (Figure 4A). Spectral data corroborated these findings, showing elevated emission

values in the ultraviolet spectrum on the leafhopper's cuticle surface (Figure 4G). Scanning

electron microscopy showed that dsBSM treatment significantly altered brochosome

morphology and distribution (Figure 4B and C-figure supplement 6A and B). The distribution

area decreased by 80%, and deformed brochosomes accumulated, contrasting with the

uniform distribution in the dsGFP treatment (Figure supplement 5A). Additionally, nearly 30%

of brochosomes exhibited significant morphological changes following dsBSM treatment

(Figure supplement 5B). Transmission electron microscopy of the Malpighian tubules indicated

numerous Golgi-derived vacuoles without brochosome distribution in epithelial cells after

ds*BSM* treatment (Figure 4D and E-figure supplement 6C and D). Predation experiments with

jumping spiders revealed an increased predation preference for leafhoppers treated with

ds*BSM*. For male leafhoppers, the proportion attacked of the predator was 70%, with a

significantly shorter predation time (71.3 s) compared to the ds*GFP* control (100.4 s) (Figure

4H). Similarly, for females, there was a higher predation rate (68.3%) and a significantly shorter

predation time (79.6 s) for ds*BSM* treated groups compared to the ds*GFP* ones (30% and 109.9

s, respectively) (Figure 4I).

Normal structure of the brochosome correlates with its antireflective properties

To further elucidate the correlation between brochosome morphology and its optical

performance, we collected brochosomes from the wings of leafhoppers treated with ds*GFP* and

ds*BSM*, respectively. Subsequently, these brochosomes were applied to the wings of brown

planthoppers and quartz slides. SEM observations revealed that the coverage of brochosomes

derived from both ds*GFP* and ds*BSM* treatments on the wings of brown planthoppers and

quartz slides were nearly identical (Figure supplement 7). Spectral measurements indicated

that the application of brochosomes from both ds*GFP* and ds*BSM* treatments effectively

reduced the reflectance of the brown planthopper wings and quartz slides. It should be noted

that brochosomes from the ds*GFP* treatment exhibited significantly higher anti-reflective

performance compared to those from the ds*BSM* treatment (Figure 4J and K). Additionally,

purified BSM proteins applied to quartz glass did not show improved anti-reflective performance

over purified GST protein (Figure supplement 8), indicating a strong correlation between

brochosome geometry and optical performance.

Brochosomes as a camouflage coating for the leafhoppers in the family Cicadellidae

Based on the forementioned findings, brochosomes of *N. cincticeps* can be considered as

an anti-reflective stealth coating against visual recognition by the predator *P. paykulli*. However,

it remains unknown whether brochosomes also serve as a stealth coating in other leafhopper

species. Therefore, we extended our investigation to include additional leafhopper species in

the family Cicadellidae, including *Recilia dorsalis*, *Empoasca onukii*, and *Psammotettix alienus*.

We carefully removed the brochosomes from the forewings of these leafhopper species using

acetone, and found a significant enhancement in UV reflectance on the forewings of the

leafhoppers (Figure 5A and B). In addition to *N. cincticeps*, we observed a considerably higher

brightness of acetone-treated forewings compared to the untreated ones under UV light for *R.*

dorsalis, *E. onukii*, and *P. alienus* (Figure 5A). This result suggests that brochosomes as an

anti-reflective stealth coating may be commonly existed in various species of the family

Cicadellidae.

Previous research indicated that the brochosome is a unique secretion in Cicadellidae

(Rakitov & Gorb, 2013b; Rakitov et al., 2018). To investigate the conservation of genes linked

to brochosome synthesis in Cicadellidae, we systematically screened these four BSM coding

genes across 116 other hemipteran insect species (Supplementary Table 2). Results of

homology analysis reveal that BSM coding genes are orphan genes restricted to the clade of

Membracoidea. However, there are variations in the species distribution of different BSM

coding genes. In Membracoidea species, homologous genes for BSM3 and 4 are observed,

while BSM2 is identified in Cicadellidae insects, and BSM1 in Deltocephalinae insects (Figure

5C). An intriguing observation is that in the majority of Cicadellidae species with confirmed

brochosome distribution, homologous gene distribution for BSM2 is detectable. In Cicadellidae

species without brochosome distribution and in the majority of Membracidae, BSM2 genes are

largely absent (Figure 5C). Since Membracidae evolved from Cicadellidae, we hypothesize that

BSM homologous gene duplication/loss may be a key factor in brochosome formation/loss. In

conclusion, our data suggest that in Cicadellidae, brochosome is synthesized by a conserved

group of BSM coding genes, making it a widely distributed anti-reflective camouflage coating

in Cicadellidae.

Discussion

Leafhoppers exhibit a distinctive grooming behavior, resulting in the deposition of a unique

extra-cuticular coating known as brochosome(Rakitov, 2009). However, the precise

characteristics and exact adaptive significance of this specialized coating remain largely

unknown. Although they are widely believed to confer hydrophobicity, protecting leafhoppers

from water and their own excreta(Rakitov & Gorb, 2013a), similar functionalities are attributed

to cuticular waxes on the external surfaces of various insects, including leafhoppers(Andersen,

1979; Bello et al., 2022). Consequently, the functionalities of brochosomes may extend beyond

this conventional understanding. In this study, we demonstrated that brochosomes exhibit

robust anti-reflective properties, decreasing the reflectance of the leafhopper cuticle surface,

particularly in the ultraviolet spectral range. This reduction may reduce visibility to visual

predators, contributing to the evasion and predation avoidance(Vukusic & Sambles, 2003;

Watson et al., 2017). Thus, brochosomes function as a natural camouflage coating on the

leafhopper cuticle surface, playing a pivotal role in reducing leafhoppers' visibility to visual

predators and offering a distinct advantage for their survival in predation scenarios as illustrated

in Fig. 6.

The present work has elucidated the function of brochosome as a natural anti-reflective

camouflage coating on leafhoppers' extra-cuticular, with its coverage directly influencing the

anti-reflective properties of the cuticle surface (Figure 2A-E and Figure 4A-G). Previous studies

suggested that arthropods employ various nanostructures formed by proteins, lipids, wax, or

adhesive substances on their cuticle surfaces to achieve anti-reflectivity (Silberglied, 1979).

These nanostructures, initially identified in the compound eyes of moths, have been

subsequently found to be widely distributed on the cuticle surfaces of insects (Bernhard, 1965;

Blagodatski et al., 2015; Chotard et al., 2022; Clapham & Hutley, 1973; Ho et al., 2016; Monteiro

et al., 2019; Stavenga et al., 2005). The nano- and microstructures on insect cuticle surfaces

generate a gradual change in refractive index near the surface, leading to antireflectivity (Raut

et al., 2011). The degree of reflection reduction depends on the shape of these nanostructures,

studies in *Drosophila melanogaster* have confirmed that modifying the morphology of

nanostructures in compound eyes directly impacts their optical performance (Kryuchkov et al.,

2020). In this study, a thorough examination of the structural protein composition of

brochosome, coupled with RNAi-mediated suppression of BSM coding gene expression,

yielded brochosomes with regular structure (ds*GFP* treatment) and structurally abnormal

brochosomes (ds*BSM* treatment) (Figure 4B-E). Subsequent in vitro experiments provided

further evidence of the correlation between the normal structure of brochosome and its anti-

reflective performance (Figure 4J and K). This correlation is likely attributed to the fact that the

diameter of the hollow pits in brochosome structure (approximately 100nm) is considerably

smaller than most spectral wavelengths(Burks et al., 2022; Rakitov, 2009). As light passes

through these small pits, it undergoes diffraction, while light passing through the ridges of

brochosome induces scattering. The interference of diffracted and scattered light between

different pits and ridges contributes to the observed extinction features in brochosome(Raut et

al., 2011). Biomimetic material and mathematical modeling analyses have clarified the

relationship between the extinction features of brochosome and the spacing between hollow

pits(Banerjee et al., 2023; Lei et al., 2020; Wang et al., 2024; Yang et al., 2017). Therefore, the

regular morphology of brochosome is a robust guarantee for its excellent anti-reflective

performance.

Moreover, we noticed a distinct contrast in the anti-reflective capabilities when applying

brochosomes with regular and irregular structures onto different substrates (quartz glass and

brown planthopper wings), especially in the UV region (Figure 4J and K). This indicates that

the reduction in reflected light is not only related to the structure of brochosome but also to

other factors. SEM observations revealed that some brochosome is embedded in the grooves

of the leafhopper cuticle surface or brown planthopper wings, which is not observed on the

smooth surface of quartz glass (Figure 4B-figure supplement 6). When the wavelength of light

is higher than the size of the structure and the surface has a gradient refractive index, light

interacts entirely with the rough surface, causing the light to gradually bend(Raut et al., 2011).

Combined with the complex structure formed by brochosomes and nanostructures on the

cuticle surface, this result in multiple internal reflections of light within these structures,

increasing light absorption and significantly enhancing anti-reflective performance. Furthermore,

the aromatic ring structures, conjugated double bonds, and peptide bonds in the proteins of

both the cuticle and brochosomes may contribute to the absorption of ultraviolet light, providing

another potential reason for their anti-reflective capabilities in the ultraviolet spectral region.

Therefore, the anti-reflective performance of brochosome on the insect cuticle surface might

be a result of the combined effect of brochosome and cuticular nanostructures.

UV light serves as a crucial visual cue for various insect predators, enhancing foraging,

navigation, mating behavior, and prey identification (Cronin & Bok, 2016; Morehouse et al.,

2017; Silberglied, 1979). Predators such as birds, reptiles, and predatory arthropods often rely

on UV vision to detect prey (Church et al., 1998; Li & Lim, 2005; Zou et al., 2011). However,

UV reflectance from insect cuticles can disrupt camouflage, increasing the risk of detection and

predation, as natural backgrounds like leaves, bark, and soil typically reflect minimal UV light

(Endler, 1997; Li & Lim, 2005; Tovee, 1995). To mitigate this risk, insects often possess anti-

reflective cuticular structures that reduce UV and broad-spectrum light reflectance. This

strategy is widespread among insects, including cicadas, dragonflies, and butterflies, and has

been shown to decrease predator detection rates (Hooper et al., 2006; Siddique et al., 2015;

Zhang et al., 2006). For example, the compound eyes of moths feature hexagonal

protuberances that reduce UV reflectance, aiding nocturnal concealment (Blagodatski et al.,

2015; Stavenga et al., 2005). In butterflies, UV reflectance from eyespots on wings can attract

predators, but reducing UV reflectance or eyespot size can lower predation risk and enhance

camouflage (Chan et al., 2019; Lyytinen et al., 2004). Hence, the reflection of ultraviolet light

from the insect cuticle surface increases the risk of predation by disrupting camouflage (Tovee,

1995) . In this study, we utilized the jumping spider *P. paykulli* to explore the impact of the

brochosome coating on the leafhopper's camouflage against predation. Hunting spiders, such

as jumping spiders and wolf spiders, are primary predators of leafhoppers(Liu et al., 2015;

Oraze & Grigarick, 1989) and the wavelength of UV light plays a crucial role as a visual cue for

jumping spiders in identifying and locating their prey compare to the other wavelengths(Li &

Lim, 2005; Zou et al., 2011). Our results revealed that *P. paykulli* displayed a preference for

preying on *N. cincticeps* with a lower brochosome coverage on their cuticle surface (Figure 2G-

J and Figure 4H-I). Thus, we hypothesize that the combination of brochosome and leafhopper

cuticular microstructures exhibits superior anti-reflective performance in the UV region,

effectively reducing UV reflection from leafhopper bodies and consequently diminishing the

predator's ability to identify them. Additionally, spectral measurement data indicated that the

wing reflection spectrum of the brown planthopper (Figure 4J) closely resembled the body

spectrum of 25-day-old female *N. cincticeps* (Figure 2E). It is possible that, in the same living

environment, other Hemipteran insects would be more conspicuous to visual predators than

leafhoppers. Brochosome coating could provide an advantage for leafhoppers to evade visual

predators, thereby enhancing their survival rates in interspecific competition. This adaptation

could potentially favor the survival of Cicadellidae in environments with high predation pressure,

allowing them to thrive in a wider range of environments. Additionally, within the leafhopper

population, the brochosome coverage on the cuticle surface of older individuals was notably

lower than that of younger individuals, making elderly leafhoppers more susceptible to be

detected and captured by predators (Figure 2F-I-figure supplement 1). This may help to

eliminate older individuals who have lost their reproductive capacity in the population, thereby

ensuring the vitality and reproductive power of the population. Therefore, we consider

brochosome, as a form of anti-reflective camouflage coating on the leafhopper's cuticle surface,

represents a more advanced evolutionary strategy against visual predation.

We successfully identified four proteins essential for the structure of brochosomes for the

first time (Figure 3). These BSM-encoded genes exhibit typical orphan gene characteristics,

consistent with previous findings(Li et al., 2022; Rakitov et al., 2018; Yuan et al., 2023). Orphan

genes originate from processes like de novo evolution, horizontal gene transfer (HGT), and

duplication–divergence(Light et al., 2014; Tautz & Domazet-Lošo, 2011). To elucidate the

evolutionary origin of the four identified BSM genes, we aligned them with all prokaryotic

genomes in NCBI and approximately 1000 arthropod genomes. Interestingly, the homologous

genes of these BSM genes were exclusively found in leafhopper genomes in the family

Cicadellidae, suggesting that the origin of these BSM genes might not be due to de novo

evolution or horizontal gene transfer. Combining the species distribution of the four identified

BSM genes (as well as their homology) (Figure 5C), we hypothesize that the BSMs were

potentially originated from a process involving duplication–divergence. In this scenario, a new

gene would be created through gene duplication or transposition, undergoing fast adaptive

evolution and losing similarity to the original gene(Light et al., 2014; Tautz & Domazet-Lošo,

2011). For instance, BSM4 is widely present in Membracoidea species and also has

homologous genes in *Clastoptera arizonana* of Cereopoidea, indicating that BSM4 may have

a common ancestral gene in Cicadomorpha. The specialization of the distal segment of the

Malpighian tubule is a characteristic shared by the three major lineages of the infraorder

Cicadomorpha: Cercopoidea(spittle-bugs), Cicadoidea(cicadas), and Membracoidea

(leafhoppers and treehoppers) (Rakitov, 2002). Cicadas coat the surface of their burrows with

secretions from the Malpighian tubules, spittlebug nymphs have their integuments covered with

foam synthesized by the Malpighian tubules, and leafhoppers coat their integuments with

brochosomes synthesized by the Malpighian tubules (Chang et al., 2019; Rakitov, 2002). The

different demands for secretions from the Malpighian tubules among these insects may be the

main reason for the rapid adaptive evolution of BSM genes.

Among Auchenorrhyncha insects, most Cicadellidae (leafhoppers) secrete brochosomes,

while many Membracidae (treehoppers) exhibit grooming behavior but lack specialized setae

(Rakitov, 1996). Considering that Membracidae are derived from Cicadellidae (Dietrich et al.,

2001), the formation of the symbiotic relationship between treehoppers and ants might due to the loss of brochosome synthesis capability. Ants create a protective environment around treehoppers, deterring threats and reducing predation pressure, providing a relatively secure living environment, leading to lower predation pressure compared to leafhoppers (Del-Claro & Oliveira, 2000; Nelson & Mooney, 2022). Consequently, the need for the protective disguise represented by brochosomes is likely diminished. We hypothesize that environmental changes drive brochosome evolution, supported by the distribution of BSM2 and BSM3 paralogs in Auchenorrhyncha. BSM3 homologs are widespread in Auchenorrhyncha, while BSM2 is restricted to brochosome-secreting Cicadellidae and absent in most Membracidae.The high structural homology between BSM2 and BSM3, despite low sequence similarity, suggests that BSM2 arose from BSM3 duplication. We propose that gene duplication/loss events are the primary factors underlying differences in brochosome synthesis between Cicadellidae and Membracidae. Gene duplication/loss is a common mechanism for functional gain/loss in eukaryotes, promoting protein diversity and redundancy (Wong & Belov, 2012). Thus, gene duplication may be a key driver of brochosome formation and functional diversity in leafhoppers.

In summary, our research has underscored the indispensable role of brochosomes as an essential anti-reflective camouflage coating for leafhoppers, conferring a significant advantage in evading visually-oriented predators. We have identified four key BSM encoding genes, a groundbreaking discovery in brochosome research, which suggests that gene duplication may be involved in the formation of these adaptive structures. Our study not only deepens our understanding of leafhoppers' anti-predator strategies but also sheds light on the distinct evolutionary trajectories employed by these insects within their ecological niche. This contributes to the development of biomimicry and the advancement of state-of-the-art camouflage technologies. Furthermore, while our study has provided valuable insights into the structure and function of brochosomes, there are still gaps in our knowledge regarding the evolutionary origins of BSMs. Future research should aim to fill these gaps, as a more comprehensive understanding of BSMs and their role in brochosome formation will enhance our appreciation of the complex evolutionary processes that have shaped these unique structures. Additionally, the analysis of BSMs in a broader range of leafhopper species will help elucidate the relationship between BSM composition and the morphological and functional

diversity of brochosomes, further informing our understanding of the ecological and

evolutionary dynamics of this insect group.

Materials and Methods

Insect

The leafhopper *N. cincticeps* and *R. dorsalis* adults were collected from a rice field in Jiaxing,

Zhejiang Province, China in September 2020. *P. alienus* adults were collected from a wheat

field in Shenyang, Liaoning Province, China in May 2020. *E. onukii* adult were collected from a

tea-garden in Fuzhou, Fijian province, China in May 2020. The identity of the leafhopper

species was confirmed using stereomicroscopy and mitochondrial cytochrome oxidase subunit

1 (CO1) sequence analysis (Supplementary Table 1). Collected leafhopper *N. cincticeps* and

R. dorsalis are maintained in insect-proof greenhouses at 26 ± 1°C under a 16:8 h light:dark

photoperiod and 50 ± 5% relative humidity on rice variety TaiChung Native 1(TN1).

The jumping spider *P. paykulli* were collected from a rice field in Ningbo, Zhejiang Province,

China. Species identification was confirmed by assessing its morphological characteristics

using stereomicroscopy and CO1 sequence analysis (Supplementary Table 1). Jumping

spiders were fed about their own mass of leafhoppers *N. cincticeps* three times each week and

examined 14-52 days following collection, as previously reported (Taylor et al., 2014).

Electron microscopy

The Malpighian tubules of *N. cincticeps* were dissected using an astereomicroscope, serially

fixed overnight at 4°C with 2.5% glutaraldehyde in 0.1 mol/L PBS, and post-fixed for 2 hours at

room temperature with 1% OsO4. Following that, samples were dehydrated in a series of

ethanol solutions (50%, 70%, 80%, 90%, and 95%), then permeabilized with 100% ethanol and

100% acetone, followed by a series of epoxies in acetone (50%, 75%, and 100%). The pierced

tissue was then immersed in epoxy resin and baked for more than 24 hours at 70°C. Ultrathin

slices were cut and stained with uranyl acetate for 15 minutes and lead citrate for 5 minutes,

and examined by Hitachi H7800 microscope (Hitachi, Japan).

Scanning electron microscopy was used to observe the morphology of leafhopper

brochosomes. Samples were prepared as previously described(Rakitov & Gorb, 2013a).

Leafhopper forewings were obtained by dissection under the stereomicroscope and glued onto

SEM metal stubs with a double-sided carbon tape. The dried samples were coated with gold

for 1 min in a MC1000 Ion Sputter Coater (Hitachi, Japan) and viewed by a HITACHI Regulus

8100 SEM (Hitachi, Japan).

Spider predation experiment

Jumping spiders were starved for three days prior to the predation experiment to ensure

that participating spiders were hungry enough to attack leafhoppers in our predation preference

tests, but not so hungry that they would attack the first prey they encountered (without being

choosy)(Taylor et al., 2014). For the jumping spiders to prey on, we placed two leafhoppers in

a 9 cm diameter test arena (the bottom of the dish was covered with filter paper). Before starting

the test, the spiders were placed in the arena for 15 minutes to adapt, and then they were

released and permitted to prey on the leafhoppers. We directly observed the feeding process

of the jumping spiders and recorded when the spiders attacked the first leafhopper and which

leafhopper they preyed on. The predation finishes when the first leafhopper is attacked or when

the jumping spider does not attack any leafhopper within 5 minutes. Each pair of jumping spider

predation tests were independently replicated more than 60 times.

The distribution of cuticle surface brochosomes and the optical properties of male and

female insects at various periods post-eclosion

To investigate the distribution of brochosomes on the cuticle surface of adult *N. cincticeps*

at various post-eclosion periods, we first collected female and male adult leafhoppers at 5-, 10-,

15-, 20-, and 25-days post-eclosion and dissected the forewings under a Nikon SMZ25

microscope (Nikon, Japan). The forewings were gold-sprayed and examined using HITACHI

Regulus 8100 SEM (Hitachi, Japan). At each time point, 30 samples were randomly selected

for SEM imaging. A 10 μm × 10 μm area within the SEM images was used to assess

brochosome coverage, and ImageJ was employed for analysis(Schneider et al., 2012). Since

brochosomes are synthesized at the distal section of the Malpighian tubule and the morphology

of the Malpighian tubule in leafhoppers can also reflect brochosomes synthesis, we used Nikon

SMZ25 microscope (Nikon, Japan) to examine the morphology of the Malpighian tubule in

female and male adult leafhoppers at 5-, 10-, 15-, 20-, and 25-days post-eclosion.

To examine the optical properties of male and female insects at various periods post-

eclosion, we first collected female and male adult leafhoppers at 5-, 10-, 15-, 20-, and 25-days

post-eclosion, freeze-killed them and placed them under white or UV light for observation

through a Nikon SMZ25 microscope (Nikon, Japan). Following that, the forewings of the

leafhoppers were meticulously dissected, and the specular reflectance spectra of the forewings

(25 × 25 μm) were precisely measured utilizing a 20/30 PV UV-Vis-NIR

microspectrophotometer (CRAIC Technologies Inc., USA). UV-Vis-NIR reflectance spectra

(250 to 1000 nm) were obtained using a 10× UV-absorbing glass objective lens, with a minimum

of five replicates per sample.

Identification of brochosome structural protein-coding genes

Candidate brochosome structural protein-coding genes were identified through a combined

analysis of transcriptomic and proteomic data. To knock down the expression of these

candidates, RNAi was conducted by microinjecting double-stranded RNA (dsRNA) into the

abdomens of fifth instar nymphs. The dsRNAs targeting 500-1000 bp regions of Malpighian

tubule-specific expression genes or GFP were synthesized in vitro using the T7 RiboMAX

Express RNAi System (Promega, USA). The 50 fifth instar nymphs of *N. cincticeps* were

microinjected with 30 nl dsRNA (0.5 μg μl^{-1}) using the Nanoject II Auto-Nanoliter Injector

(Drummond, USA). Thereafter, they were transferred to healthy rice seedlings for recovery.

Forewings of treated leafhoppers were collected 7 days after microinjection (about 4-5 days

post-eclosion), sprayed with gold, and analyzed by HITACHI Regulus 8100 SEM (Hitachi,

Japan). Screening identified four brochosome structural protein-coding genes, and inhibiting

their expression resulted in significant changes in brochosome morphology. The full-length

sequence of these genes was obtained using by 5'-rapid amplification of cDNA ends (RACE)

and 3'-RACE using SMARTer RACE 5'/3' Kit (Clontech, USA).

To analyze the temporal expression profiles of four brochosome structural protein-coding

genes (GenBank accession number PP273097, PP273098, PP273099, PP273100), the whole

body, salivary gland, midgut, ovary, and testis were dissected from 200 adult insects, total RNA

in different tissue were extracted using TRIzol Reagent (Invitrogen, USA). The relative

expression of these genes in different tissues of *N. cincticeps* adults was detected by RT-qPCR

assay using a QuantStudio™ 5 Real-Time PCR system (Thermo Fisher Scientific, USA). The

detected transcript levels were normalized to the transcript level of the housekeeping gene

elongation factor 1 alpha (EF1α) (GenBank accession number AB836665) and estimated by

the $2^{-\triangle\triangle Ct}$ (cycle threshold) method. Using the same procedure, the expression levels of

these brochosome structural protein-coding genes were evaluated in female and male adult

leafhoppers at 5-, 10-, 15-, 20-, and 25-days post-eclosion.

To investigate the effects of four BSM-encoding genes on brochosome morphology and

distribution, dsRNAs were designed to target two non-overlapping regions within the sequences

of these genes. These dsRNAs were microinjected into the abdomens of leafhoppers both

individually and in combination, with dsRNA targeting GFP serving as a negative control.

Following a seven-day incubation period post-injection, changes in BSM gene expression were

assessed in both control and treated groups via RT-qPCR. The morphology and distribution of

brochosomes on the leafhopper cuticle were examined and documented using SEM.

Additionally, ImageJ software was employed to quantify the proportion of morphologically

abnormal brochosomes and the distribution area of brochosomes on the cuticle surface after

injection of the dsRNA mixtures targeting the two non-overlapping regions of each BSM gene,

as well as after dsGFP treatment.

Knocking down in vivo expression of BSM-coding genes in *N. cincticeps*

Following the prior procedure, dsRNA produced from four BSM-coding genes was injected

into the abdomen of fifth instar nymphs of *N. cincticeps* by microinjection, and a control group

injected with ds*GFP* was established. After a seven-day incubation post-injection, alterations

in the expression of the four BSM genes were monitored in the ds*BSM* and ds*GFP* treated

leafhoppers using RT-qPCR. The synthesis of brochosomes within the Malpighian tubules was

investigated through TEM analysis. The morphology and distribution of brochosomes on the

leafhopper cuticle were observed and imaged using SEM. To examine the optical properties,

the ds*BSM* and ds*GFP* treated leafhoppers, freeze-killed them and placed them under white or

UV light for observation through a Nikon SMZ25 microscope (Nikon, Japan). Subsequently, the

forewings of the leafhoppers were meticulously dissected, and the specular reflectance spectra

of the forewings (25 × 25 μm) were precisely measured utilizing a 20/30 PV UV-Vis-NIR

microspectrophotometer (CRAIC Technologies Inc., USA). UV-Vis-NIR reflectance spectra

(250 to 1000 nm) were obtained using a 10× UV-absorbing glass objective lens, with a minimum

of five replicates per sample. Finally, the predation preference of the jumping spiders on the

ds*BSM* and ds*GFP* treated leafhoppers was examined by a predation experiment.

In vitro measurements were conducted to assess the relationship between brochosome

morphology and their optical performance.

To collect approximately 500 *N. cincticeps* treated with ds*GFP* or ds*BSM*, dissect the

forewings of the leafhoppers and immerse them in acetone in a 50 mL centrifuge tube. Place

the centrifuge tubes on a room temperature orbital shaker at 50 rpm for 12 hours to separate

the forewings from the leafhoppers due to friction. Following this, centrifuge at 1000g for 10

minutes to separate the brochosomes. Resuspend the particles in fresh acetone, briefly

sonicate, then centrifuge at 1000g for 10 minutes. Repeat this process three times. Resuspend

the collected brochosomes from ds*GFP* and ds*BSM*-treated leafhopper forewings in acetone

and adjust the two suspension solutions to the same OD280 reading. Next, using a pipette,

carefully drop the brochosome solution onto quartz slides and brown planthopper wings. Allow

the acetone to completely evaporate at room temperature. At the same time, we also set up

quartz slides and brown planthopper wings that were treated with acetone alone as negative

controls. Prepared quartz slides and brown planthopper wings can be observed under SEM to

examine the distribution of brochosomes. Additionally, UV-Vis-NIR reflectance spectra (250 to

1000 nm) were obtained using a 20/30 PV UV-Vis-NIR microspectrophotometer (CRAIC

Technologies Inc., USA), with a minimum of five replicates per sample.

To investigate the correlation between BSM proteins and the optical characteristics of

brochosomes, the four genes encoding BSM proteins were separately cloned into pET-28a or

pGEX-4T2 vectors. Following induction,the expressed proteins were purified from cell lysates

via batch affinity chromatography using either glutathione (GSH) agarose or nickel-

nitrilotriacetic acid (Ni-NTA) agarose,depending on the fusion tag. GST protein was produced

following the identical protocol from cells harboring the empty pGEX-4T2 vector. The purified

BSM fusion proteins and GST protein were normalized to an equivalent OD280 value. These

protein solutions were then meticulously spotted onto quartz slides and permitted to dry

completely at ambient temperature. In parallel, quartz slides treated exclusively with PBS

served as negative controls. The quartz slides were subsequently analyzed to acquire UV-Vis-

NIR reflectance spectra (ranging from 250 to 1000 nm) using a 20/30 PV UV-Vis-NIR

microspectrophotometer (CRAIC Technologies Inc.,USA), with a minimum of five technical

replicates conducted per sample.

Removal of leafhopper forewing brochosome by acetone

Previous research has shown that acetone may efficiently remove brochosomes from the

cuticle surface of leafhoppers(Rakitov et al., 2018). We manually removed the brochosomes

from the forewings of leafhoppers using acetone and examined the imaging under white or UV

light for observation through a Nikon SMZ25 microscope (Nikon, Japan). UV-Vis-NIR

reflectance spectra (250 to 1000 nm) were obtained using a 20/30 PV UV-Vis-NIR

microspectrophotometer (CRAIC Technologies Inc., USA), with a minimum of five replicates

per sample.

Bioinformatics analysis and phylogenetic tree

The brochosome is a unique secretion produced by leafhoppers (Cicadellidae). In order to

determine the distribution of brochosome structural protein-coding genes in Hemiptera, the

brochosome structural protein-coding genes were used as queries to search for homologous

sequences in Hemiptera transcriptomes. We examined 116 Hemiptera species, encompassing

major families (Acanaloniidae, Achilidae, Aetalionidae, Aphrophoridae, Caliscelidae,

Cercopidae, Cicadellidae, Cicadidae, Cixiidae, Clastopteridae, Delphacidae, Derbidae,

Dictyopharidae, Epipygidae, Eurybrachidae, Flatidae, Fulgoridae, Issidae, Machaerotidae,

Melizoderidae, Membracidae, Myerslopiidae, Nogodinidae, Peloridiidae, Ricaniidae,

Tettigarctidae, Tettigometridae, Theaceae, Tropiduchidae). Transcriptome data of 116

Hemiptera species were downloaded from the NCBI Sequence Read Archive (Supplementary

Table 2), and assembled using SOAPdenovo-Trans (version 1.01)(Xie et al., 2014). The

resulting transcripts were filtered to remove potential contaminants. We compared four

identified BSM-encoded genes with these transcriptomes using the tblastn (E-value < 1.0e–5).

Heatmaps were generated using TBtools software(Chen et al., 2023). At least one homologous

gene for the four BSM genes was identified in 66 species. A species phylogenetic tree for these

66 Hemiptera species was constructed following the methods outlined in previous

studies(Johnson et al., 2018). OrthoFinder to identify orthogroups and infer the species tree,

obtaining 1683 single-copy orthologous genes after filtering (Emms & Kelly, 2019). MAFFT was

employed for sequence alignment, followed by trimming with trimAI (Capella-Gutiérrez et al.,

2009; Katoh & Standley, 2013). The best amino acid substitution model was selected using

ModelFinder based on the Bayesian Information Criterion (BIC) (Kalyaanamoorthy et al., 2017).

The maximum likelihood phylogenetic tree was constructed using IQ-TREE v1.6.8 with 1000

ultrafast bootstrap replicates (Minh et al., 2020). All analyses were conducted using PhyloSuite

v1.2.2, which integrates MAFFT, trimAI, ModelFinder, and IQ-TREE into a unified pipeline

(Capella-Gutiérrez et al., 2009; Emms & Kelly, 2019; Kalyaanamoorthy et al., 2017; Katoh &

Standley, 2013; Minh et al., 2020; Zhang et al., 2020).

The amino acid sequences of BSM2 and BSM3 were aligned using the Clustal W multiple

sequence alignment program(Larkin et al., 2007). AlphaFold2 was employed to predict the

three-dimensional structures of BSM2 and BSM3, and structural alignments were conducted

using PyMOL(DeLano, 2002; Jumper et al., 2021). In Membracoidea, the maximum likelihood

(ML) phylogenetic tree for BSM1-4 was constructed using MAFFT, trimAI, ModelFinder, and

IQ-TREE in PhyloSuite (Capella-Gutiérrez et al., 2009; Emms & Kelly, 2019; Kalyaanamoorthy

et al., 2017; Katoh & Standley, 2013; Minh et al., 2020; Xiang et al., 2023; Zhang et al., 2020).

Statistical analyses

All experiments were performed at least three independent replicates and statistical

analyses were performed with GraphPad Prism8.0 software. Statistical significance was

calculated by a two-tailed Student's t-test, one-way ANOVA, two-way ANOVA and/or unpaired

Student's t-test. P-values < 0.05 were considered statistically significant.

Acknowledgments

This work received support from the National Natural Science Foundation of China

(U23A6006, U20A2036, 32270150) and the Natural Science Foundation of Ningbo Municipality

(2023J112). The author acknowledges Dr. Yuan Cheng from the Instrumentation and Service

Center for Molecular Science (ISCMS) at Westlake University for assistance with optical

measurement and data interpretation. We are grateful for the support provided by the

Bioimaging Center of the Bioimaging Center, State Key Laboratory for Managing Biotic and

Chemical Threats to the Quality and Safety of Agro-products, Institute of Plant Virology, Ningbo

University, in TEM and SEM measurement and data interpretation. Additionally, we thank

Professor Shou-Wei Ding (University of California, Riverside, CA, United States) for his

valuable and constructive suggestions for manuscript improvement.

Author contributions:

671   W.W. J.-M.L., C.-X.Z., and J.-P.C. designed research; W.W. Q.M. Z.Y. Z.L. performed

672   research; W.W. and Z.Y. analyzed data; and W.W. H.-W.S., J.-M.L., C.-X.Z., and J.-P.C wrote

673   the paper. All authors gave final approval for publication. All authors have read and agreed to

674   the published version of the manuscript.

Data availability

677   All the data needed to understand and assess the conclusions of this research are available

678   in the article, Supplementary Materials, and GenBank (accession number PP273097,

679   PP273098, PP273099, PP273100). Any additional information required to reanalyse the data

680   reported in this paper is available from the lead contact upon request.

Declaration of interests

683   The authors declare no competing interests.

References

Andersen, S. O. (1979). Biochemistry of insect cuticle. *Annual review of entomology*, *24*(1), 29-59.

Banerjee, P., Burks, G. R., Bialik, S. B., Nassr, M., Bello, E., Alleyne, M., Freeman, B. D., Barrick, J. E., Schroeder, C. M., & Milliron, D. J. (2023). Nanostructure‐Derived Antireflectivity in Leafhopper Brochosomes. *Advanced Photonics Research*, *4*(7), 2200343. https://doi.org/10.1002/adpr.202200343

Bello, E., Chen, Y., & Alleyne, M. (2022). Staying Dry and Clean: An Insect's Guide to Hydrophobicity. *Insects*, *14*(1), 42. https://doi.org/10.3390/insects14010042

Bernhard, C. G. (1965). The insect corneal nipple array. A biological, broad-band impedance transformer that acts as an antireflection coating. *Acta physiol. scand.*, *63*(243), 1-79.

Blagodatski, A., Sergeev, A., Kryuchkov, M., Lopatina, Y., & Katanaev, V. L. (2015). Diverse set of Turing nanopatterns coat corneae across insect lineages. *Proceedings of the National Academy of Sciences*, *112*(34), 10750-10755. https://doi.org/10.1073/pnas.1505748112

Burks, G. R., Yao, L., Kalutantirige, F. C., Gray, K. J., Bello, E., Rajagopalan, S., Bialik, S. B., Barrick, J. E., Alleyne, M., Chen, Q., & Schroeder, C. M. (2022). Electron Tomography and Machine Learning for Understanding the Highly Ordered Structure of Leafhopper Brochosomes. *Biomacromolecules*, *24*(1), 190-200. https://doi.org/10.1021/acs.biomac.2c01035

Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T. (2009). trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*, *25*(15), 1972-1973.

Chan, I. Z., Rafi, F. Z., & Monteiro, A. (2019). Interacting effects of eyespot number and ultraviolet reflectivity on predation risk in Bicyclus anynana (Lepidoptera: Nymphalidae). *Journal of Insect Science*, *19*(6), 19.

Chang, S.-C., Shih, H.-T., & Lu, K.-H. (2019). Antifungal effect and chitinase activities of the froth of spittlebug Poophilus costalis (Walker)(Hemiptera: Cercopoidea: Aphrophoridae). *Journal of Asia-Pacific Entomology*, *22*(1), 269-273.

Chen, C., Wu, Y., Li, J., Wang, X., Zeng, Z., Xu, J., Liu, Y., Feng, J., Chen, H., He, Y., & Xia, R. (2023). TBtools-II: A "one for all, all for one" bioinformatics platform for biological big-data mining. *Molecular plant*, *16*(11), 1733-1742. https://doi.org/10.1016/j.molp.2023.09.010

Chotard, A., Ledamoisel, J., Decamps, T., Herrel, A., Chaine, A. S., Llaurens, V., & Debat, V. (2022). Evidence of attack deflection suggests adaptive evolution of wing tails in butterflies. *Proceedings of the Royal Society B: Biological Sciences*, *289*(1975), 20220562. https://doi.org/10.1098/rspb.2022.0562

Church, S. C., Bennett, A. T. D., Cuthill, I. C., & Partridge, J. C. (1998). Ultraviolet cues affect the foraging behaviour of blue tits. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *265*(1405), 1509-1514. https://doi.org/10.1098/rspb.1998.0465

Cinel, S. D., Hahn, D. A., & Kawahara, A. Y. (2020). Predator-induced stress responses in insects: A review. *Journal of insect physiology*, *122*, 104039. https://doi.org/10.1016/j.jinsphys.2020.104039

Clapham, P., & Hutley, M. (1973). Reduction of lens reflexion by the "Moth Eye" principle. *Nature*,
244(5414), 281-282.
Cronin, T. W., & Bok, M. J. (2016). Photoreception and vision in the ultraviolet. *Journal of*
Experimental Biology, *219*(18), 2790-2801. https://doi.org/10.1242/jeb.128769
Del-Claro, K., & Oliveira, P. S. (2000). Conditional outcomes in a neotropical treehopper-ant
association: temporal and species-specific variation in ant protection and homopteran
fecundity. *Oecologia*, *124*(2), 156-165. https://doi.org/10.1007/s004420050002
DeLano, W. L. (2002). Pymol: An open-source molecular graphics tool. *CCP4 Newsl. Protein*
Crystallogr, *40*(1), 82-92.
Dietrich, C., Rakitov, R., Holmes, J., & Black IV, W. (2001). Phylogeny of the major lineages of
Membracoidea (Insecta: Hemiptera: Cicadomorpha) based on 28S rDNA sequences.
Molecular Phylogenetics and Evolution, *18*(2), 293-305.
Emms, D. M., & Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome biology*, *20*, 1-14.
Endler, J. A. (1997). Some general comments on the evolution and design of animal
communication systems. *Philosophical Transactions of the Royal Society of London.*
Series   B: Biological   Sciences,   *340*(1292),   215-225.
https://doi.org/10.1098/rstb.1993.0060
Galloway, J. A. M., Green, S. D., Stevens, M., & Kelley, L. A. (2020). Finding a signal hidden
among noise: how can predators overcome camouflage strategies? *Philosophical*
Transactions of the Royal Society B: Biological Sciences, *375*(1802), 20190478.
https://doi.org/10.1098/rstb.2019.0478
Gouranton, J., & Maillet, P.-L. (1967). Origine et structure des brochosomes. *Journal de*
Microscopie, *6*, 53-64.
Hibino, H. (1996). Biology and Epidemiology of Rice Viruses. *Annual review of phytopathology*,
34(1), 249-274. https://doi.org/10.1146/annurev.phyto.34.1.249
Ho, S., Schachat, S. R., Piel, W. H., & Monteiro, A. (2016). Attack risk for butterflies changes
with eyespot number and size. *Royal Society open science*, *3*(1), 150614.
https://doi.org/10.1098/rsos.150614
Hooper, I. R., Vukusic, P., & Wootton, R. J. (2006). Detailed optical study of the transparent
wing membranes of the dragonfly Aeshna cyanea. *Optics express*, *14*(11), 4891-4897.
https://doi.org/10.1364/oe.14.004891
Hughes, A., Liggins, E., & Stevens, M. (2019). Imperfect camouflage: how to hide in a variable
world? *Proceedings of the Royal Society B: Biological Sciences*, *286*(1902), 20190646.
https://doi.org/10.1098/rspb.2019.0646
Johnson, K. P., Dietrich, C. H., Friedrich, F., Beutel, R. G., Wipfler, B., Peters, R. S., Allen, J.
M., Petersen, M., Donath, A., Walden, K. K. O., Kozlov, A. M., Podsiadlowski, L., Mayer,
C., Meusemann, K., Vasilikopoulos, A., Waterhouse, R. M., Cameron, S. L., Weirauch,
C., Swanson, D. R.,…Yoshizawa, K. (2018). Phylogenomics and the evolution of
hemipteroid insects. *Proceedings of the National Academy of Sciences*, *115*(50),
12775-12780. https://doi.org/10.1073/pnas.1815820115
Jones, M. P., Pierce, K. E., & Ward, D. (2007). Avian Vision: A Review of Form and Function
with Special Consideration to Birds of Prey. *Journal of exotic pet medicine*, *16*(2), 69-
87. https://doi.org/10.1053/j.jepm.2007.03.012

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool,
K., Bates, R., Žídek, A., & Potapenko, A. (2021). Highly accurate protein structure
prediction with AlphaFold. *Nature*, *596*(7873), 583-589.

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K., Von Haeseler, A., & Jermiin, L. S. (2017).
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature*
methods, *14*(6), 587-589.

Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Molecular biology and evolution*, *30*(4),
772-780.

Kryuchkov, M., Bilousov, O., Lehmann, J., Fiebig, M., & Katanaev, V. L. (2020). Reverse and
forward engineering of Drosophila corneal nanocoatings. *Nature*, *585*(7825), 383-389.
https://doi.org/10.1038/s41586-020-2707-9

Larkin, M. A., Blackshields, G., Brown, N. P., Chenna, R., McGettigan, P. A., McWilliam, H.,
Valentin, F., Wallace, I. M., Wilm, A., Lopez, R., Thompson, J. D., Gibson, T. J., &
Higgins, D. G. (2007). Clustal W and Clustal X version 2.0. *Bioinformatics*, *23*(21),
2947-2948. https://doi.org/10.1093/bioinformatics/btm404

Lei, C.-W., Chen, R.-Y., & Yang, H. (2020). Leafhopper Wing-Inspired Broadband
Omnidirectional Antireflective Embroidered Ball-Like Structure Arrays Using a
Nonlithography-Based Methodology. *Langmuir*, *36*(19), 5296-5302.
https://doi.org/10.1021/acs.langmuir.0c00634

Li, D., & Lim, M. L. M. (2005). Ultraviolet cues affect the foraging behaviour of jumping spiders.
Animal Behaviour, *70*(4), 771-776. https://doi.org/10.1016/j.anbehav.2004.12.021

Li, Z., Li, Y., Xue, A. Z., Dang, V., Holmes, V. R., Johnston, J. S., Barrick, J. E., Moran, N. A., &
Parsch, J. (2022). The Genomic Basis of Evolutionary Novelties in a Leafhopper.
Molecular biology and evolution, *39*(9), msac184.
https://doi.org/10.1093/molbev/msac184

Light, S., Basile, W., & Elofsson, A. (2014). Orphans and new gene origination, a structural and
evolutionary perspective. *Current Opinion in Structural Biology*, *26*, 73-83.
https://doi.org/10.1016/j.sbi.2014.05.006

Lim, U. T., & Ben-Yakir, D. (2020). Visual sensory systems of predatory and parasitic arthropods.
Biocontrol Science and Technology, *30*(7), 728-739.
https://doi.org/10.1080/09583157.2020.1752362

Lin, M., Yang, G., Wang, Z., & You, M. (2021). Brochosome detachment facilitates Empoasca
onukii (Hemiptera: Cicadellidae) adults escaping from spider web. *Acta Entomol Sin*,
64, 213-222.

Liu, S., Li, Z., Sui, Y., Schaefer, D. A., Alele, P. O., Chen, J., & Yang, X. (2015). Spider foraging
strategies dominate pest suppression in organic tea plantations. *BioControl*, *60*(6),
839-847. https://doi.org/10.1007/s10526-015-9691-2

Lyytinen, A., Brakefield, P. M., Lindström, L., & Mappes, J. (2004). Does predation maintain
eyespot plasticity in Bicyclus anynana? *Proceedings of the Royal Society of London.*
Series B: Biological Sciences, *271*(1536), 279-283.

Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., Von Haeseler,
A., & Lanfear, R. (2020). IQ-TREE 2: new models and efficient methods for

815 phylogenetic inference in the genomic era. *Molecular biology and evolution*, *37*(5),
816 1530-1534.

817 Monteiro, A., Rafi, F. Z., Chan, I. Z. W., & Fiedler, K. (2019). Interacting Effects of Eyespot
818 Number and Ultraviolet Reflectivity on Predation Risk in Bicyclus anynana (Lepidoptera:
819 Nymphalidae). *Journal of Insect Science*, *19*(6). https://doi.org/10.1093/jisesa/iez123

820 Morehouse, N. I., Buschbeck, E. K., Zurek, D. B., Steck, M., & Porter, M. L. (2017). Molecular
821 Evolution of Spider Vision: New Opportunities, Familiar Players. *Biological Bulletin*,
822 *233*(1), 21-38. https://doi.org/10.1086/693977

823 Nelson, A. S., & Mooney, K. A. (2022). The Evolution and Ecology of Interactions Between Ants
824 and Honeydew-Producing Hemipteran Insects. *Annual Review of Ecology, Evolution,*
825 *and Systematics*, *53*(1), 379-402. https://doi.org/10.1146/annurev-ecolsys-102220-
826 014840

827 Oraze, M. J., & Grigarick, A. A. (1989). Biological control of aster leafhopper (Homoptera:
828 Cicadellidae) and midges (Diptera: Chironomidae) by Pardosa ramulosa (Araneae:
829 Lycosidae) in California rice fields. *Journal of Economic Entomology*, *82*(3), 745-749.

830 Rakitov, R., & Gorb, S. N. (2013a). Brochosomal coats turn leafhopper (Insecta, Hemiptera,
831 Cicadellidae) integument to superhydrophobic state. *Proceedings of the Royal Society*
832 *B: Biological Sciences*, *280*(1752), 20122391. https://doi.org/10.1098/rspb.2012.2391

833 Rakitov, R., & Gorb, S. N. (2013b). Brochosomes protect leafhoppers (Insecta, Hemiptera,
834 Cicadellidae) from sticky exudates. *Journal of the Royal Society Interface*, *10*(87),
835 20130445. https://doi.org/10.1098/rsif.2013.0445

836 Rakitov, R., Moysa, A. A., Kopylov, A. T., Moshkovskii, S. A., Peters, R. S., Meusemann, K.,
837 Misof, B., Dietrich, C. H., Johnson, K. P., Podsiadlowski, L., & Walden, K. K. O. (2018).
838 Brochosomins and other novel proteins from brochosomes of leafhoppers (Insecta,
839 Hemiptera, Cicadellidae). *Insect biochemistry and molecular biology*, *94*, 10-17.
840 https://doi.org/10.1016/j.ibmb.2018.01.001

841 Rakitov, R. A. (1996). Post-moulting behaviour associated with Malpighian tubule secretions in
842 leafhoppers and treehoppers (Auchenorrhyncha: Membracoidea). *European Journal of*
843 *Entomology*, *93*(2), 167-184. <Go to ISI>://WOS:A1996VB52200006

844 Rakitov, R. A. (2000). Secretion of brochosomes during the ontogenesis of a leafhopper,
845 Oncometopia orbona (F.) (Insecta, Homoptera, Cicadellidae). *Tissue and Cell*, *32*(1),
846 28-39. https://doi.org/10.1054/tice.1999.0084

847 Rakitov, R. A. (2002). Structure and function of the Malpighian tubules, and related behaviors
848 in juvenile cicadas: evidence of homology with spittlebugs (Hemiptera: Cicadoidea &
849 Cercopoidea). *Zoologischer Anzeiger-A Journal of Comparative Zoology*, *241*(2), 117-
850 130.

851 Rakitov, R. A. (2009). Brochosomal Coatings of the Integument of Leafhoppers (Hemiptera,
852 Cicadellidae). In S. N. Gorb (Ed.), *Functional Surfaces in Biology* (pp. 113-137).
853 Springer Netherlands. https://doi.org/10.1007/978-1-4020-6697-9_8

854 Raut, H. K., Ganesh, V. A., Nair, A. S., & Ramakrishna, S. (2011). Anti-reflective coatings: A
855 critical, in-depth review. *Energy & Environmental Science*, *4*(10), 3779-3804.
856 https://doi.org/10.1039/c1ee01297e

857 Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of
858 image analysis. *Nature methods*, *9*(7), 671-675.

Sheikh, A. A., Rehman, N., & Kumar, R. (2017). Diverse adaptations in insects: A review. *Journal of entomology and zoology studies*, *5*(2), 343-350.

Siddique, R. H., Gomard, G., & Hölscher, H. (2015). The role of random nanostructures for the omnidirectional anti-reflection properties of the glasswing butterfly. *Nature Communications*, *6*(1), 6909. https://doi.org/10.1038/ncomms7909

Silberglied, R. E. (1979). Communication in the ultraviolet. *Annual Review of Ecology and Systematics*, *10*(1), 373-398. https://www.annualreviews.org/doi/abs/10.1146/annurev.es.10.110179.002105

Skelhorn, J., & Rowe, C. (2016). Cognition and the evolution of camouflage. *Proceedings of the Royal Society B: Biological Sciences*, *283*(1825), 20152890. https://doi.org/10.1098/rspb.2015.2890

Stavenga, D. G., Foletti, S., Palasantzas, G., & Arikawa, K. (2005). Light on the moth-eye corneal nipple array of butterflies. *Proceedings of the Royal Society B: Biological Sciences*, *273*(1587), 661-667. https://doi.org/10.1098/rspb.2005.3369

Stevens, M. (2007). Predator perception and the interrelation between different forms of protective coloration. *Proceedings of the Royal Society B: Biological Sciences*, *274*(1617), 1457-1464. https://doi.org/10.1098/rspb.2007.0220

Stevens, M., & Ruxton, G. D. (2018). The key role of behaviour in animal camouflage. *Biological Reviews*, *94*(1), 116-134. https://doi.org/10.1111/brv.12438

Stobbe, N., Dimitrova, M., Merilaita, S., & Schaefer, H. M. (2008). Chromaticity in the UV/blue range facilitates the search for achromatically background-matching prey in birds. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *364*(1516), 511-517. https://doi.org/10.1098/rstb.2008.0248

Swain, R. (1936). Notes on the oviposition and life-history of the leaf-hopper Oncometopia undata Fabr.(Homoptera: Cicadellidae).

Tautz, D., & Domazet-Lošo, T. (2011). The evolutionary origin of orphan genes. *Nature Reviews Genetics*, *12*(10), 692-702. https://doi.org/10.1038/nrg3053

Taylor, L. A., Maier, E. B., Byrne, K. J., Amin, Z., & Morehouse, N. I. (2014). Colour use by tiny predators: jumping spiders show colour biases during foraging. *Animal Behaviour*, *90*, 149-157. https://doi.org/10.1016/j.anbehav.2014.01.025

Thery, M., & Gomez, D. (2010). Insect colours and visual appearance in the eyes of their predators. *Advances in insect physiology*, *38*, 267-353.

Tovee, M. (1995). Ultra-violet photoreceptors in the animal kingdom: their distribution and function. *Trends in ecology & evolution*, *10*(11), 455-460. https://doi.org/10.1016/s0169-5347(00)89179-x

Vukusic, P., & Sambles, J. R. (2003). Photonic structures in biology. *Nature*, *424*(6950), 852-855. https://doi.org/10.1038/nature01941

Walker, S. E., & Rypstra, A. L. (2002). Sexual dimorphism in trophic morphology and feeding behavior of wolf spiders (Araneae: Lycosidae) as a result of differences in reproductive roles. *Canadian Journal of Zoology*, *80*(4), 679-688. https://doi.org/10.1139/z02-037

Wang, L., Li, Z., Shen, S., & Wong, T.-S. (2024). Geometric design of antireflective leafhopper brochosomes. *Proceedings of the National Academy of Sciences*, *121*(14), e2312700121.

Watson, G. S., Watson, J. A., & Cribb, B. W. (2017). Diversity of Cuticular Micro- and Nanostructures on Insects: Properties, Functions, and Potential Applications. *Annual Review of Entomology, Vol 62*, *62*, 185-205. https://doi.org/10.1146/annurev-ento-031616-035020

Wei, J., Jia, D., Mao, Q., Zhang, X., Chen, Q., Wu, W., Chen, H., & Wei, T. (2018). Complex interactions between insect-borne rice viruses and their vectors. *Current opinion in virology*, *33*, 18-23. https://doi.org/10.1016/j.coviro.2018.07.005

Wong, E. S. W., & Belov, K. (2012). Venom evolution through gene duplications. *Gene*, *496*(1), 1-7. https://doi.org/10.1016/j.gene.2012.01.009

Wu, W., Ye, Z., Mao, Q., Shan, H.-W., Li, J.-M., & Chen, J.-P. (2023). Combined Transcriptome and Proteome Analysis of the Protein Composition of the Brochosomes of the Leafhopper Nephotettix cincticeps. *Insects*, *14*(10), 784. https://doi.org/10.3390/insects14100784

Xi, W., Lee, Y.-J., Yu, S., Chen, Z., Shiomi, J., Kim, S.-K., & Hu, R. (2023). Ultrahigh-efficient material informatics inverse design of thermal metamaterials for visible-infrared-compatible camouflage. *Nature Communications*, *14*(1), 4694. https://doi.org/10.1038/s41467-023-40350-6

Xiang, C. Y., Gao, F., Jakovlić, I., Lei, H. P., Hu, Y., Zhang, H., Zou, H., Wang, G. T., & Zhang, D. (2023). Using PhyloSuite for molecular phylogeny and tree‐based analyses. *iMeta*, *2*(1), e87.

Xie, Y., Wu, G., Tang, J., Luo, R., Patterson, J., Liu, S., Huang, W., He, G., Gu, S., Li, S., Zhou, X., Lam, T.-W., Li, Y., Xu, X., Wong, G. K.-S., & Wang, J. (2014). SOAPdenovo-Trans: de novo transcriptome assembly with short RNA-Seq reads. *Bioinformatics*, *30*(12), 1660-1666. https://doi.org/10.1093/bioinformatics/btu077

Yan, B., Yu, X., Dai, R., Li, Z., & Yang, M. (2021). Chromosome-level genome assembly of Nephotettix cincticeps (Uhler, 1896)(Hemiptera: Cicadellidae: Deltocephalinae). *Genome Biology and Evolution*, *13*(11), evab236.

Yang, H., Peng, Y., Tian, J., Wang, J., Wei, B., Xie, C., & Wang, Z. (2018). Rice Field Spiders in China: A Review of the Literature. *Journal of Economic Entomology*, *111*(1), 53-64. https://doi.org/10.1093/jee/tox319

Yang, S. K., Sun, N., Stogin, B. B., Wang, J., Huang, Y., & Wong, T. S. (2017). Ultra-antireflective synthetic brochosomes. *Nature Communications*, *8*(1), 1285. https://doi.org/10.1038/s41467-017-01404-8

Yuan, F., Su, M., Li, T., Zhang, Y., Dietrich, C. H., Webb, M. D., & Wei, C. (2023). Functional and evolutionary implications of protein and metal content of leafhopper brochosomes. *Insect biochemistry and molecular biology*, *157*, 103962. https://doi.org/10.1016/j.ibmb.2023.103962

Zhang, D., Gao, F., Jakovlić, I., Zou, H., Zhang, J., Li, W. X., & Wang, G. T. (2020). PhyloSuite: An integrated and scalable desktop platform for streamlined molecular sequence data management and evolutionary phylogenetics studies. *Molecular ecology resources*, *20*(1), 348-355.

Zhang, G., Zhang, J., Xie, G., Liu, Z., & Shao, H. (2006). Cicada Wings: A Stamp from Nature for Nanoimprint Lithography. *Small*, *2*(12), 1440-1443. https://doi.org/10.1002/smll.200600255

946   Zou, Y., Araujo, D. P., Lim, M. L. M., & Li, D. (2011). Ultraviolet is a more important cue than
947   reflection in other wavelengths for a jumping spider to locate its spider prey. *Animal*
948   *Behaviour*, *82*(6), 1457-1463. https://doi.org/10.1016/j.anbehav.2011.09.031
949
950

Figure 1. Brochosomes are a distinctive coating on the cuticle surface of leafhopper.

(A) Male adult of the green rice leafhopper *N. cincticeps*. Bar, 1 mm. (B) Brochosomes on the surface of the forewing of *N. cincticeps*. Bar, 1 μm. (C, D) The morphologies of brochosomes by SEM (C) and TEM (D). Bar, 200 nm. (E) Alimentary tract and the Malpighian tubules of *N. cincticeps*. The leafhopper *N. cincticeps* have two pairs of Malpighian tubules, each divided into proximal segment, distal segment, and terminal segment. The brochosomes are synthesized in the distal segment, which is dilated and rod-shaped. Bar, 1 mm. (F) The distal segment epithelial cell displays an extensive rough endoplasmic reticulum and multiple Golgi regions (red arrow) containing developing brochosomes in its basal portion, as well as a number of secretory vacuoles with mature brochosomes near the cell border. Bar, 5 μm. (G) The initial stage of the development of brochosome. Bar, 500 mm. (H) The two brochosomes that are developing inside the primary vesicles are shown in close-up. Regular invaginations appear on the surface of the growing brochosome at the same time that its matrix separates into a looser core and a denser wall. Bar, 500 nm. (I) Larger vesicles containing numerous BS, formed by the fusion of multiple primary vesicles. The surface of the brochosomes has regular cell-like invaginations, and it is surrounded by amorphous flocculent material with a moderate electron

density. Bar, 1 μm. (J) A vesicle filled with mature brochosomes, each mature brochosome has

a spherical inner cavity and a well-swollen outer margin of the septa. Bar, 1 μm. E, F, G, H are

the enlargement of boxed area in D. Fc, filter chamber; Mg, midgut; Mt, Malpighian tubules;

mtd, distal segment of the Malpighian tubule; mtp, proximal segment of the Malpighian tubule;

mtt, terminal segment of the Malpighian tubule. All images are representative of at least three

replicates.

Figure 2. The distribution of brochosomes on the cuticle surface of leafhopper *N. cincticeps* are associated with predation by jumping spiders.

(A, B) Images of leafhopper *N. cincticeps* males and females in white (A) and ultraviolet light (B) at 5 and 25 days post-eclosion, respectively. Bar, 1 mm. (C, D) The distribution of brochosomes on the surface of a forewing (C) and morphological changes of the Malpighian

tubules (D) of *N. cincticeps* males and females at 5 and 25 days post-eclosion. Bar, 5 µm in

C; Bar, 1 mm in D. (E) Reflectance spectra of female and male forewing of *N. cincticeps* at 5

and 25 days post-eclosion. (F) Images of the jumping spider *P. paykulli*. Bar, 2 mm. (G-J)

Jumping spiders prefer leafhoppers with little brochosome covering as food. In predation

experiment, jumping spiders offered *N. cincticeps* male and female at 5 days post-eclosion

(G), male and female at 25 days post-eclosion (H), males at 5 and 25 days post-eclosion (I),

and females at 5 and 25 days post-eclosion (J). Data on predation times are displayed using

the traditional box and whisker shapes. All box plots with whiskers represent the data

distribution based on five number summary statistics (maximum, third quartile, median, first

quartile, minimum), each dot in box plot represents an independent experiment. **$P < 0.01$,

****$P < 0.0001$, ns no significance, Statistical significance was determined by unpaired t test

with Welch's correction method. Predation preference is shown in the pie chart. All images

are representative of at least three replicates.

Figure 3. Identification of brochosome structural proteins. (A) Morphology of brochosomes on the forewing of leafhopper *N. cincticeps* at 7 days post-microinjection with dsRNA mix targeting two non-overlapping regions of each BSM gene. Bar, 500 nm. (B) Gene structures of *BSM-1*, *BSM-2*, *BSM-3* and *BSM-4*. (C-F) Transcription levels of BSM-1 (C), BSM-2 (D), BSM-3 (E), and BSM-4 (F) at 7 days post-microinjection with dsRNA mix targeting two non-overlapping regions of each BSM gene. Each data point represents the result of one independent experiment. (G-J) The abundance of BSM transcripts in different tissues and whole bodies of *N. cincticeps* was determined by RT-qPCR. Notably, *BSM-1* (G), *BSM-2* (H), *BSM-3* (I), and *BSM-4* (J) exhibited specific expression in the Malpighian tubules. Each data point represents the result of one independent experiment. (K-N) The expression patterns of

1006   *BSM-1* (K), *BSM-2* (L), *BSM-3* (M) and *BSM-4* (N) transcripts were examined in male and

1007   female leafhopper at 5, 10, 15, 20, and 25 days post-eclosion. Results were obtained from 3

1008   independent experiments. For C-N, data shown are mean ± SD values. $*P < 0.05$; $**P < 0.01$;

1009 $***P < 0.001$; $****P < 0.0001$; ns no significance (C-F, two tailed Student's t-test; G-J, one-way

1010   ANOVA; K-N, two-way ANOVA). All images are representative of at least three replicates.

1011

1012

Figure 4. RNAi inhibits brochosome synthesis, alters brochosome morphology, and

influences predation in jumping spiders. (A) Images of leafhopper *N. cincticeps* males and

females in white and ultraviolet light after ds*GFP* or ds*BSM* treatment, respectively. Bar, 1 mm.

(B, C) Morphology of the brochosome on the forewings of leafhoppers after ds*GFP* (B) and

ds*BSM* (C) treatment. Bar, 2 μm. (D, E) Morphology of the brochosome in the distal segment

epithelial cells of Malpighian tubules after ds*GFP* (D) and ds*BSM* (E) treatment. Bar, 2 μm. (F)

The transcript levels of *BSM-1*, *BSM-2*, *BSM-3* and *BSM-4* at 7 days after ds*GFP* and ds*BSM*

treatment. Each data point represents the outcome of an individual independent experiment.

The presented data are expressed as mean ± SD values. Statistical significance is denoted as

*$P < 0.05$ and **$P < 0.01$, determined by two-way ANOVA. (G) Reflectance spectra of female

and male forewing of *N. cincticeps* at 7 days after ds*GFP* and ds*BSM* treatment. (H and I)

Jumping spiders prefer to prey on ds*BSM*-treated leafhoppers. Predation efficiency and

preference of jumping spiders on males (H) and females (I) *N. cincticeps* after ds*GFP* and

ds*BSM* treatment in the predation experiment. Data on predation times are displayed using the

traditional box and whisker shapes. All box plots with whiskers represent the data distribution

based on five number summary statistics (maximum, third quartile, median, first quartile,

minimum), each dot in box plot represents an independent experiment. **$P < 0.01$, ***$P < 0.001$,

Statistical significance was determined by unpaired t test with Welch's correction method.

Predation preference is shown in the pie chart. (J, K) The morphology of brochosomes is related

to their optical performance. Collect brochosomes treated with ds*GFP* and ds*BSM* separately,

apply them to brown planthopper wings (J) and quartz slides (K), and set up quartz slides or

brown planthopper wings treated solely with acetone as negative controls. All images are

representative of at least three replicates.

1036

1037

Figure 5. Brochosome coating on Cicadellidae cuticle surface is essential for their

optical qualities. (A) Images of forewings of four leafhopper species before and after acetone

treatment in white and ultraviolet light. Bar, 1mm. (B) Reflection spectra of the forewings of the

N. cincticeps before and after acetone treatment. (C) Phylogenetic associations between the

BSM and phylogeny of Hemipteran lineages. The left column describes the phylogeny of 76

representative species of Hemiptera with BSM proteins. In the right column, the presence of

BSM- encoded genes in different Hemiptera species is illustrated along with their homology

 analysis with *N. cincticeps*. White indicates the absence of related genes; the color gradient

1046 represents the degree of nucleotide sequence similarity. All images are representative of at

1047 least three replicates.

1048

1049

Fig. 6. Brochosomes serve as an anti-reflective camouflage coating on the cuticle

surface of leafhoppers. Brochosomes effectively reduce the reflection of various wavelengths

of light, particularly in the UV region. UV light is a crucial visual cue for numerous visual

predators to identify and locate prey. Brochosomes efficiently decrease the UV light reflection

on the surface of leafhoppers, thereby reducing their exposure risk to visual predators and

facilitating evasion of visual predation.