Brochosomes as an antireflective camouflage coating for leafhoppers

Wei Wu[1*], Qianzhuo Mao[1], Zhuangxin Ye[1], Zhenfeng Liao[2], Hong-Wei Shan[1,] Jun-Min Li[1],

Chuan-Xi Zhang[1], Jian-Ping Chen[1*]

[1]State Key Laboratory for Managing Biotic and Chemical Threats to the Quality and Safety of

Agro-products, Key Laboratory of Biotechnology in Plant Protection of Ministry of Agriculture

and Zhejiang Province, Institute of Plant Virology, Ningbo University, Ningbo 315211, China.

[2]State Key Laboratory for Sustainable Control of Pest and Disease, Institute of Virology and

Biotechnology, Zhejiang Academy of Agricultural Sciences, Hangzhou, 310021, China.

*Corresponding author. Email: jianpingchen@nbu.edu.cn (J.-P.C.);

wuwei_19861115@163.com (W.W.)

Abstract:

In nature, insects face immense predation pressure, where visual cues play a vital role in predators locating them. To counter this threat, insects employ a variety of nano- and microstructures on their cuticular layer to manipulate and interact with light, enhancing anti-reflective properties and providing camouflage or reducing detectability by predators. Leafhoppers have a unique extra-cuticular coating called brochosome, yet its anti-reflective functions and protein composition remain unclear. Our study demonstrates strong anti-reflective properties of brochosomes, effectively reducing reflectance on the cuticle surface, especially in the ultraviolet spectrum, to improve evasion from visual predators. Furthermore, we identify four novel structural proteins of the brochosome (BSM) for the first time. Inhibiting their synthesis by RNAi alters brochosome morphology, impacting the optical properties of the cuticle surface. Evolutionary origin analysis of BSM suggests that brochosomes likely originated from a process involving duplication–divergence. Our study reveals that leafhoppers employ a unique camouflage strategy by secreting brochosomes as anti-reflection nano-coatings, enabling them to evade natural predators and contributing to their evolutionary success.

Keywords:

Brochosome, Leafhopper, Camouflage, Antireflective, Predation, BSM

Introduction:

Predation exerts significant selective pressure on the evolution of various species, driving the development of anti-predator strategies across the animal kingdom(Galloway et al., 2020; Stevens & Ruxton, 2018). Camouflage is a widely employed strategy, allowing animals to blend into their surroundings by adjusting their coloration and patterns to reduce detection by visual predators(Hughes et al., 2019; Skelhorn & Rowe, 2016). Nevertheless, the presence of predators with multispectral vision, capable of perceiving non-visible light spectra like infrared and ultraviolet, can compromise traditional color and pattern-based camouflage(Galloway et al., 2020; Jones et al., 2007; Stevens, 2007; Thery & Gomez, 2010), paralleling the challenges of military camouflage under non-visible light(Xi et al., 2023). Consequently, achieving alignment between an animal's surface optical characteristics and its surroundings becomes a crucial requirement in visual camouflage strategies.

Insects, one of the most diverse animal groups on earth, contend with intense predation pressure(Cinel et al., 2020; Sheikh et al., 2017). Many of their predators, including birds, reptiles, and predatory arthropods, rely on visual cues, with some possessing highly developed ultraviolet (UV) vision, where UV light plays a critical role in locating insect prey(Lim & Ben-Yakir, 2020; Stobbe et al., 2008). Furthermore, the natural backgrounds where insects reside, such as leaves, tree bark, and soil, exhibit minimal light reflection(Endler, 1997). Consequently, light reflections originating from an insect's body significantly elevate its risk of exposure to predators(Tovee, 1995). To counter this challenge, the various nano- and microstructural features of the insect cuticle have excellent antireflective properties that affect insect body coloration, reduce surface light reflection and aid camouflage(Vukusic & Sambles, 2003; Watson et al., 2017). The first instances of anti-reflective structures were observed in the compound eyes of moths, characterized their corneas are covered by hexagonally arranged protrusions measuring approximately 200 nanometers apart (Bernhard, 1965; Blagodatski et al., 2015). Subsequent research has revealed that anti-reflective structures are widespread on insect body surfaces, primarily found on compound eyes and wings(Blagodatski et al., 2015; Chotard et al., 2022; Ho et al., 2016; Monteiro et al., 2019; Stavenga et al., 2005). These nano- and micro-structures on the cuticular layer are believed to possess anti-reflective properties, effectively minimizing light reflections from their bodies and

thus preventing detection by predators(Blagodatski et al., 2015; Chotard et al., 2022; Ho et al.,

2016; Monteiro et al., 2019; Stavenga et al., 2005).

Leafhoppers (Cicadellidae), one of the largest insect families with over 22,000 species,

possess unique extra-cuticular coating known as brochosome. Brochosomes, typically hollow,

honeycomb-like spheres with a diameter ranging from 0.2 to 0.6 micrometers, are synthesized

in the Malpighian tubules, secreted through the hindgut, and applied as a coating on the fresh

cuticle following molting(Rakitov, 2000; Rakitov, 2009). Brochosomes primarily comprise lipids

and proteins, with protein content ranging from 45% to 75%, and the exact proteins

responsible for brochosome formation remain unknown(Li et al., 2022; Rakitov et al., 2018;

Yuan et al., 2023). Some have proposed that brochosomes may serve as a protective layer on

the leafhopper's cuticle surface, with a potential role in enhancing the hydrophobicity of their

cuticle surface, providing defense against pathogens and predators(Rakitov & Gorb, 2013a;

Rakitov, 2009). Nevertheless, despite the known function of hydrophobicity and the shedding

of brochosomes in helping leafhopper escape spider webs, there is currently no definitive

biological evidence supporting other hypothesized functions of brochosomes(Lin et al., 2021;

Rakitov & Gorb, 2013a, 2013b). Recent studies have explored bio-inspired materials

mimicking the structure of brochosomes have shown outstanding antireflection properties in

recent studies(Lei et al., 2020; Yang et al., 2017). While these artificial materials do not

perfectly replicate the intricate structure of brochosomes, they offer valuable insights,

indicating that leafhoppers may use a unique strategy, setting them apart from other insects, to

achieve anti-reflection properties on their surfaces. Therefore, we hypothesize that

brochosomes could potentially serve as a camouflage coating on the leafhopper's cuticle

surface, helping it reduce surface optical reflections and stay hidden to avoid detection by

visual predators, ultimately escaping from predation.

In this study, we combined spectrophotometry, electron microscopy, transcriptome analysis,

proteome analysis, gene function validation, and leafhopper and jumping spider bioassays to

characterize the protein composition and the function of brochosome in leafhopper

camouflage. We focused our research on the rice green leafhopper, *Nephotettix cincticeps*, a

well-known agricultural pest with an extensive research history(Hibino, 1996; Wei et al., 2018;

Yan et al., 2021). Our results revealed the pronounced effectiveness of brochosome coverage

in reducing light reflection on leafhoppers *N. cincticeps*, especially within the scope of the

ultraviolet spectrum. By conducting RNAi experiments, we successfully identified four

structural proteins of brochosome for the first time. The suppression of their synthesis induced

changes in brochosome morphology, influencing the optical characteristics of the leafhopper's

cuticle surface. Furthermore, our analysis of the evolutionary origin of brochosomes indicates

that they likely originated through a process involving duplication–divergence. Our findings

support brochosomes as a unique camouflage coating that enables leafhoppers to evade

visual predators.

Results

Brochosomes are a distinctive coating on the cuticle surface of leafhopper *N.

cincticeps.*

The cuticle surface of rice green leafhopper *N. cincticeps* is coated with brochosomes

(Figure 1A and B), which are approximately 350nm in diameter and featuring a hollow sphere

with a honeycomb-like structure (Figure 1C and D). The leafhopper *N. cincticeps* possesses

two pairs of Malpighian tubules, each consisting of proximal segment, distal segment, and

terminal segment (Figure 1E). Brochosomes are synthesized in the distal segment of the

Malpighian tubule, which exhibits a swollen and rod-shaped appearance (Figure 1E). The

epithelial cells in this segment contain large spherical nuclei, an extensive rough endoplasmic

reticulum, and multiple Golgi regions (Figure 1E and F). Within the Golgi-derived vacuole, the

brochosomes undergo progressive development, transforming their initially spherical granules

into a honeycomb-like surface with closely spaced invaginations (Figure 1G-I). Mature

brochosomes are stored in secretory vacuoles at the boundaries of the cells and are

subsequently secreted into the lumen of the Malpighian tubules (Figure 1F).

The optical properties of the leafhopper's cuticle surface are intricately connected to

the brochosome coating.

To investigate the interplay between brochosome coverage and the optical features of *N.*

cincticeps cuticle surface, we observed the relationship for male and female adults at 5, 10, 15,

20, and 25 days post-eclosion. *N. cincticeps* exhibited a gradual shift in body color with

increasing eclosion time, manifesting clear distinctions between males and females (Figure

2A-figure supplement 1A and B). Males transformed from light green to dark green, while

females progressively transitioned from light green to translucent, displaying an iridescence

(Figure 2A-figure supplement 1A and B). Furthermore, the brightness of the leafhopper cuticle

surface under UV light intensified with the extension of post-eclosion time, and the UV

reflectance of female cuticle surface was significantly higher than that of males after 15 days

post-eclosion (Figure 2B-figure supplement 1A and B).

To validate the correlation between brochosome coverage, body color, and UV light

reflectance, we examined the distribution of brochosomes on the cuticle surface of male and

female *N. cincticeps* at various post-eclosion intervals using scanning electron microscopy

(SEM). SEM results unveiled a gradual reduction in brochosome coverage on the cuticle

surface of both genders as post-eclosion time progressed. Specifically, male coverage

decreased from 90% to approximately 40%, whereas female coverage descended from 90%

to around 10% (Figure 2C-figure supplement 1A-C). Microscopic analyses of the Malpighian

tubule revealed that, with prolonged post-eclosion time, the distal segment of the tubule

underwent gradual atrophy, particularly pronounced at 15 days post-eclosion. Notably, females

exhibited a more marked atrophy in the distal segment compared to males. By 25 days

post-eclosion, the distal segment of the Malpighian tubule in females became indistinguishable

from other segments (Figure 2D-figure supplement 1A and B). This denotes a progressive

reduction in post-eclosion brochosome synthesis, resulting in a corresponding decrease in

brochosome coverage on the cuticle surface. Furthermore, the distribution of brochosomes on

the leafhopper cuticle surface may be correlated with its optical characteristics. To substantiate

this hypothesis, we analyzed the light reflection characteristics of leafhopper forewings at

various post-eclosion time points. The findings revealed a positive correlation between

brochosome coverage on the forewings and light reflection values, underscoring the

remarkable anti-reflective attributes of the brochosome coating on leafhopper cuticle surface

(Figure 2E-figure supplement 1D and E). The brochosome coating efficiently diminishes the

reflectance of UV and visible light on the cuticle surface of *N. cincticeps*.

Preferential capture of leafhoppers with fewer brochosomes by jumping spiders

Given that UV light serves as a crucial visual cue for arthropod predators, especially

common visual hunters like jumping spiders(Cronin & Bok, 2016; Li & Lim, 2005; Morehouse

et al., 2017; Silberglied, 1979; Zou et al., 2011), the brochosome coating on leafhoppers'

cuticle surface significantly reduces UV light reflectance. This prompted us to investigate the

effect of brochosome coatings on reducing predation risk in leafhoppers. Predation

experiments were conducted using *Plexippus paykulli* (Figure 2F-figure supplement 2), a

common jumping spider in rice fields known to prey on leafhoppers(Yang et al., 2018).

Predation rates and times were scrutinized as parameters to assess predation

efficiency(Taylor et al., 2014; Walker & Rypstra, 2002). In the experimental group comprising

females and males at 5 days post-eclosion, *P. paykulli* did not exhibit a clear predation

preference (Figure 2G). However, in the experimental group involving females and males at 25

days post-eclosion, there was a higher probability of predation on female *N. cincticeps*

compare to male ones (Figure 2H). For male *N. cincticeps* in both the 5-day and 25-day

post-eclosion experimental groups, *P. paykulli* showed a stronger inclination towards preying

on individuals that were eclosed for 25 days, resulting in significantly shorter predation times

compared to those at 5 days post-eclosion (Figure 2I). These patterns were consistently

observed for female leafhoppers in both the 5-days and 25-days post-eclosion (Figure 2J).

The cumulative findings strongly suggest a distinct preference of the jumping spider *P. paykulli*

for individuals with lower brochosome coverage on the cuticle surface. Based on these

experiments, we proposed a hypothesis that the brochosome coating on the leafhopper cuticle

surface may mitigate predation risk by reducing surface light reflection, particularly in the UV

spectrum.

Identifying the major structural proteins of brochosomes.

Despite the initial discovery of brochosomes in the early 1950s and the confirmation in the

1960s that they are protein-lipid particles(Gouranton & Maillet, 1967), the specific protein

composition of brochosomes remains unknown. Recent suggested that they are primarily

composed of brochosomins (BSM) and brochosome-associated proteins (BSAP)(Li et al.,

2022; Rakitov et al., 2018). BSM, a novel class of secreted proteins with molecular weights

ranging from 21 to 40 kDa, is considered as the major structural component of

brochosomes(Rakitov et al., 2018). However, there remains uncertainty regarding the specific

proteins implicated in brochosome formation.

To further elucidate the essential protein component of brochosome and the relationship

between the brochosome coating and optical properties of leafhopper's cuticle surface,

transcriptomic and proteomic data derived from brochosome of *N. cincticeps* were

comprehensively analyzed. Based on our prior integrated analysis of the brochosome

transcriptome and proteome(Wu et al., 2023), in conjunction with literature search results, we

selected 50 candidate genes for functional analysis using RNAi-mediated gene silencing.

Through RNAi experiments, we successfully identified four genes encoding brochosome

structural proteins that led to significant morphological changes in brochosomes (Figure

3A-figure supplement 3). According to the coding sequence (CDS) length of these genes, the

four proteins were names as BSM-1 to BSM-4 (GenBank accession number PP273097,

PP273098, PP273099, PP273100) (Figure 3B). Comparative analysis with previous reported

brochosome proteins indicated that BSM2 and BSM3 may form a pair of paralogous genes

based on their highly conserved protein sequences (figure supplement 4A and C); although

BSM2 and BSM3 show lower sequence homology, their structural similarities is notably high

(figure supplement 4B). In contrast, BSM-1 and BSM-4 lack homologous genes within the

other reported leafhopper brochosome protein repertoire. Results of RT-qPCR showed that

microinjection of dsRNA effectively suppressed the expression of these four BSM-coding

genes (Figure 3C-F). SEM observations also indicated that the inhibition of these four

BSM-coding genes individually led to a reduction in brochosome coverage on the cuticle

surface of *N. cincticeps* (figure supplement 3). Specifically, knockdown of these genes resulted

in significant morphological changes in the brochosome, characterized by an increase in the

diameter of certain brochosomes and deformation of the honeycomb-like structures on the

surface (Figure 3*A*-figure supplement 3). In addition, investigation of temporal and spatial

expression patterns showed that all the four BSM-coding genes were specifically expressed in

the Malpighian tubules (Figure 3G-J) and exhibited relatively stable expression during the

early post-eclosion period, followed by a gradual decrease after 10 days (Figure 3K-N). By 25

days post-eclosion, the expression of BSM-coding genes declined to around 10% in females

and 30-40% in males, respectively (Figure 3K-N). This phenomenon is consistent with our

initial microscopic observations, suggesting that the gradual reduction in brochosome

synthesis contributing primarily to the decrease of brochosome coverage on the cuticle surface

of *N. cincticeps* after adulthood.

Brochosome coating diminishes light reflection and facilitates predator avoidance

To further investigate the function of brochosomes in leafhopper cuticle surface associated

with light reflection and predator avoidance, we implemented RNAi to simultaneously suppress

the expression of all four BSM-coding genes with ds*BSM* mixture injection. RT-qPCR results

demonstrated a substantial downregulation in the expression of these genes compared to the

ds*GFP* control (Figure 4F). Notably, under ultraviolet light, both male and female leafhoppers

exhibited a significant increase in UV reflection on their cuticle surface following ds*BSM*

treatment (Figure 4A). Spectral data corroborated these findings, showing elevated emission

values in the ultraviolet spectrum on the leafhopper's cuticle surface (Figure 4G). Electron

microscopy observations revealed pronounced alterations in brochosome morphology and

distribution after ds*BSM* treatment. Reduced distribution area and the accumulation of

deformed brochosomes were observed, differing from the uniform distribution in the ds*GFP*

treatment (Figure 4B and C-figure supplement 5A and B). Transmission electron microscopy of

the Malpighian tubules indicated numerous Golgi-derived vacuoles without brochosome

distribution in epithelial cells after ds*BSM* treatment (Figure 4D and E-figure supplement 5C

and D). Predation experiments with jumping spiders revealed an increased predation

preference for leafhoppers treated with ds*BSM*. For male leafhoppers, the proportion attacked

of the predator was 70%, with a significantly shorter predation time (71.3 s) compared to the

ds*GFP* control (100.4 s) (Figure 4H). Similarly, for females, there was a higher predation rate

(68.3%) and a significantly shorter predation time (79.6 s) for ds*BSM* treated groups compared

to the ds*GFP* ones (30% and 109.9 s, respectively) (Figure 4I).

Normal structure of the brochosome correlates with its antireflective properties

To further elucidate the correlation between brochosome morphology and its optical

performance, we collected brochosomes from the wings of leafhoppers treated with ds*GFP*

and ds*BSM*, respectively. Subsequently, these brochosomes were applied to the wings of

brown planthoppers and quartz slides. SEM observations revealed that the coverage of

brochosomes derived from both ds*GFP* and ds*BSM* treatments on the wings of brown

planthoppers and quartz slides were nearly identical (figure supplement 6). Spectral

measurements indicated that the application of brochosomes from both ds*GFP* and ds*BSM*

treatments effectively reduced the reflectance of the brown planthopper wings and quartz

slides. It should be noted that brochosomes from the ds*GFP* treatment exhibited significantly

higher anti-reflective performance compared to those from the ds*BSM* treatment (Figure 4J

and K). This indicates a close correlation between brochosome morphology and its optical

performance.

Brochosomes as a camouflage coating for the leafhoppers in the family Cicadellidae

Based on the forementioned findings, brochosomes of *N. cincticeps* can be considered as

an anti-reflective stealth coating against visual recognition by the predator *P. paykulli*.

However, it remains unknown whether brochosomes also serve as a stealth coating in other

leafhopper species. Therefore, we extended our investigation to include additional leafhopper

species in the family Cicadellidae, including *Recilia dorsalis*, *Empoasca onukii*, and

Psammotettix alienus. We carefully removed the brochosomes from the forewings of these

leafhopper species using acetone, and found a significant enhancement in UV reflectance on

the forewings of the leafhoppers (Figure 5A and B). In addition to *N. cincticeps*, we observed a

considerably higher brightness of acetone-treated forewings compared to the untreated ones

under UV light for *R. dorsalis*, *E. onukii*, and *P. alienus* (Figure 5A). This result suggests that

brochosomes as an anti-reflective stealth coating may be commonly existed in various species

of the family Cicadellidae.

Previous research indicated that brochosome as a unique secretion in Cicadellidae

(Rakitov & Gorb, 2013b; Rakitov et al., 2018). To investigate the conservation of genes linked

to brochosome synthesis in Cicadellidae, we systematically screened these four BSM coding

genes across 116 other hemipteran insect species (Supplementary Table 2). Results of

homology alignment reveal that BSM coding genes exhibit marked orphan gene

characteristics, with homologous genes exclusively present in Membracoidea. However, there

are variations in the species distribution of different BSM coding genes. In Membracoidea

species, homologous genes for BSM3 and 4 are observed, while BSM2 is identified in

Cicadellidae insects, and BSM1 exclusively in Deltocephalinae insects (Figure 5C).

Furthermore, the degree of homology of all BSM genes is positively correlated with taxonomic

classification. An intriguing observation is that in the majority of Cicadellidae species with

confirmed brochosome distribution, homologous gene distribution for BSM2 is detectable. In

Cicadellidae species without brochosome distribution and in the majority of Membracidae,

BSM2 genes are largely absent (Figure 5C). Since Membracidae evolved from Cicadellidae,

we hypothesize that BSM homologous gene duplication/loss may be a key factor in

brochosome formation/loss. In conclusion, our data suggest that in Cicadellidae, brochosome

is synthesized by a conserved group of BSM coding genes, making it a widely distributed

anti-reflective camouflage coating in Cicadellidae.

Discussion

Leafhoppers exhibit a distinctive grooming behavior, resulting in the deposition of a unique

extra-cuticular coating known as brochosome(Rakitov, 2009). However, the precise

characteristics and exact adaptive significance of this specialized coating remain largely

unknown. Although they are widely believed to confer hydrophobicity, protecting leafhoppers

from water and their own excreta(Rakitov & Gorb, 2013a), similar functionalities are attributed

to cuticular waxes on the external surfaces of various insects, including leafhoppers(Andersen,

1979; Bello et al., 2022). Consequently, the functionalities of brochosomes may extend

beyond this conventional understanding. In this study, we demonstrated that brochosomes

exhibit robust anti-reflective properties, markedly decreasing the reflectance of the leafhopper

cuticle surface, particularly in the ultraviolet spectral range. This substantial reduction is

visibility to visual predators, contributing to the evasion and predation avoidance(Vukusic &

Sambles, 2003; Watson et al., 2017). Thus, brochosomes function as a natural camouflage

coating on the leafhopper cuticle surface, playing a pivotal role in reducing leafhoppers'

visibility to visual predators and offering a distinct advantage for their survival in predation

scenarios as illustrated in Fig. 6.

The present work has elucidated the function of brochosome as a natural anti-reflective

camouflage coating on leafhoppers' extra-cuticular, with its coverage directly influencing the

anti-reflective properties of the cuticle surface (Figure 2A-E and Figure 4A-G). Previous

studies suggested that arthropods employ various nanostructures formed by proteins, lipids,

wax, or adhesive substances on their cuticle surfaces to achieve anti-reflectivity(Silberglied,

1979). These nanostructures, initially identified in the compound eyes of moths, have been

subsequently found to be widely distributed on the cuticle surfaces of insects(Bernhard, 1965;

Blagodatski et al., 2015; Chotard et al., 2022; Ho et al., 2016; Monteiro et al., 2019; Stavenga

et al., 2005). The nano- and microstructures on insect cuticle surfaces generate a gradual

change in refractive index near the surface, leading to antireflectivity(Raut et al., 2011). The

degree of reflection reduction depends on the shape of these nanostructures, studies in

Drosophila melanogaster have confirmed that modifying the morphology of nanostructures in

compound eyes directly impacts their optical performance(Kryuchkov et al., 2020). In this

study, a thorough examination of the structural protein composition of brochosome, coupled

with RNAi-mediated suppression of BSM coding gene expression, yielded brochosomes with

regular structure (ds*GFP* treatment) and structurally abnormal brochosomes (ds*BSM*

treatment) (Figure 4B-E). Subsequent in vitro experiments provided further evidence of the

correlation between the normal structure of brochosome and its anti-reflective performance

(Figure 4J and K). This correlation is likely attributed to the fact that the diameter of the hollow

pits in brochosome structure (approximately 100nm) is considerably smaller than most

spectral wavelengths(Burks et al., 2022; Rakitov, 2009). As light passes through these small

pits, it undergoes diffraction, while light passing through the ridges of brochosome induces

scattering. The interference of diffracted and scattered light between different pits and ridges

contributes to the observed extinction features in brochosome(Raut et al., 2011). Biomimetic

material and mathematical modeling analyses have clarified the relationship between the

extinction features of brochosome and the spacing between hollow pits(Banerjee et al., 2023;

Yang et al., 2017). Therefore, the regular morphology of brochosome is a robust guarantee for

its excellent anti-reflective performance.

Moreover, we noticed a distinct contrast in the anti-reflective capabilities when applying

brochosomes with regular and irregular structures onto different substrates (quartz glass and

brown planthopper wings), especially in the UV region (Figure 4J and K). This indicates that

the reduction in reflected light is not only related to the structure of brochosome but also to

other factors. SEM observations revealed that some brochosome is embedded in the grooves

of the leafhopper cuticle surface or brown planthopper wings, which is not observed on the

smooth surface of quartz glass (Figure 4B-figure supplement 6). When the wavelength of light

is higher than the size of the structure and the surface has a gradient refractive index, light

interacts entirely with the rough surface, causing the light to gradually bend(Raut et al., 2011).

Combined with the complex structure formed by brochosomes and nanostructures on the

cuticle surface, this result in multiple internal reflections of light within these structures,

increasing light absorption and significantly enhancing anti-reflective performance.

Furthermore, the aromatic ring structures, conjugated double bonds, and peptide bonds in the

proteins of both the cuticle and brochosomes may contribute to the absorption of ultraviolet

light, providing another potential reason for their excellent anti-reflective capabilities in the

ultraviolet spectral region. Therefore, the outstanding anti-reflective performance of

brochosome on the insect cuticle surface might be a result of the combined effect of

brochosome and cuticular nanostructures.

UV light serves as a crucial visual cue for various insect predators, enhancing foraging,

navigation, mating behavior, and prey identification(Cronin & Bok, 2016; Morehouse et al.,

2017; Silberglied, 1979), while the reflection of ultraviolet light from the insect cuticle surface

increases the risk of predation by disrupting camouflage (Tovee, 1995) . In this study, we

utilized the jumping spider *P. paykulli* to explore the impact of the brochosome coating on the

leafhopper's camouflage against predation. Hunting spiders, such as jumping spiders and wolf

spiders, are primary predators of leafhoppers(Liu et al., 2015; Oraze & Grigarick, 1989) and

the wavelength of UV light plays a crucial role as a visual cue for jumping spiders in identifying

and locating their prey compare to the other wavelengths(Li & Lim, 2005; Zou et al., 2011). Our

results revealed that *P. paykulli* displayed a preference for preying on *N. cincticeps* with a

lower brochosome coverage on their cuticle surface (Figure 2G-J and Figure 4H-I). Thus, we

hypothesize that the combination of brochosome and leafhopper cuticular microstructures

exhibits superior anti-reflective performance in the UV region, effectively reducing UV

reflection from leafhopper bodies and consequently diminishing the predator's ability to identify

them. Additionally, spectral measurement data indicated that the wing reflection spectrum of

the brown planthopper (Figure 4J) closely resembled the body spectrum of 25-day-old female

N. cincticeps (Figure 2E). It is possible that, in the same living environment, other Hemipteran

insects would be more conspicuous to visual predators than leafhoppers. Brochosome coating

could provide an advantage for leafhoppers to evade visual predators, thereby enhancing their

survival rates in interspecific competition. This adaptation could potentially favor the survival of

Cicadellidae in environments with high predation pressure, allowing them to thrive in a wider

range of environments. Additionally, within the leafhopper population, the brochosome

coverage on the cuticle surface of older individuals was notably lower than that of younger

individuals, making elderly leafhoppers more susceptible to be detected and captured by

predators (Figure 2F-I-figure supplement 1). This may contribute to eliminating elderly and

weaker individuals in the leafhopper population, conserving food resources, and protecting the

entire population. Therefore, we consider brochosome, as a form of anti-reflective camouflage

coating on the leafhopper's cuticle surface, represents a more advanced evolutionary strategy

against visual predation. It not only provides a significant advantage in terms of camouflage

and evasion from visual predators but also plays a crucial role in inter-species competition and

stably maintenance of the insect population, highlighting its importance in the survival and

evolution of leafhoppers.

In addition, we successfully identified four proteins essential for the structure of

brochosomes for the first time (Figure 3). These BSM-encoded genes exhibit typical orphan

gene characteristics, consisting with previous findings(Li et al., 2022; Rakitov et al., 2018;

Yuan et al., 2023). Orphan genes originating from processes like de novo evolution, horizontal

gene transfer (HGT), and duplication–divergence(Light et al., 2014; Tautz & Domazet-Lošo,

2011). To elucidate the evolutionary origin of the four identified BSM genes, we aligned them

with all prokaryotic genomes in NCBI and approximately 1000 arthropod genomes.

Interestingly, the homologous genes of these BSM genes were exclusively found in leafhopper

genomes in the family Cicadellidae, suggesting that the origin of these BSM genes might not

because of the de novo evolution or horizontal gene transfer. Combining the species

distribution of the four identified BSM genes (as well as their homology) (Figure 5C), we

hypothesize that the BSMs were potentially originated from a process involving duplication–

divergence. In this scenario, a new gene would be created through gene duplication or

transposition, undergoing fast adaptive evolution and losing similarity to the original gene(Light

et al., 2014; Tautz & Domazet-Lošo, 2011). For Membracoidea insects, the majority insects in

the family Cicadellidae (leafhoppers) can secrete brochosomes, while many Membracidae

(treehoppers) insects exhibit grooming behavior without observable distinct setae on the

cuticle surface(Rakitov, 1996). Considering that Membracidae are derived from Cicadellidae

(Dietrich et al., 2001), the formation of the symbiotic relationship between treehoppers and

ants might due to the loss of brochosome synthesis capability. Ants create a protective

environment around treehoppers, deterring threats and reducing predation pressure, providing

a relatively secure living environment, leading to lower predation pressure compared to

leafhoppers(Del-Claro & Oliveira, 2000; Nelson & Mooney, 2022). Consequently, the need for

the protective disguise represented by brochosomes is likely diminished. We speculate that

environmental changes serve as one of the major driving forces in the evolution of

brochosomes. Moreover, the distribution of paralogous genes BSM2 and BSM3 in

Membracoidea (Figure 5C-figure supplement 4C) suggests that gene duplication/loss may be

a primary factor causing the difference in brochosome synthesis capability between the insects

of the families Cicadellidae and Membracidae. In eukaryotes, gene duplication/loss is a

common mechanism for generating/losing gene functions, and this process promotes protein

diversity and functional redundancy through the production of multiple homologous genes

(Wong & Belov, 2012). Therefore, we propose that gene duplication may be a significant factor

contributing to the formation and morphological/functional diversity of brochosomes in

leafhoppers.

In summary, our exploration has underscored the indispensable role of brochosomes as a

vital anti-reflective camouflage coating in leafhoppers, granting them a significant edge in

eluding visual predators. The identification of the four BSM encoding genes, a groundbreaking

revelation in brochosome research, points towards the likely involvement of gene duplication in

the formation of these adaptive structures. Our study not only deepens our understanding of

leafhoppers' anti-predator tactics but also illuminates the distinct evolutionary trajectories

utilized by these insects within their ecological niche, thus contributing to the development of

biomimicry and the advancement of state-of-the-art camouflage technologies.

Materials and Methods

Insect

The leafhopper *N. cincticeps* and *R. dorsalis* adults were collected from a rice field in Jiaxing, Zhejiang Province, China in September 2020. *P. alienus* adults were collected from a wheat field in Shenyang, Liaoning Province, China in May 2020. *E. onukii* adult were collected from a tea-garden in Fuzhou, Fijian province, China in May 2020. The identity of the leafhopper species was confirmed using stereomicroscopy and mitochondrial cytochrome oxidase subunit 1 (CO1) sequence analysis (Supplementary Table 1). Collected leafhopper *N. cincticeps* and *R. dorsalis* are maintained in insect-proof greenhouses at 26 ± 1°C under a 16:8 h light:dark photoperiod and 50 ± 5% relative humidity on rice variety TaiChung Native 1(TN1).

The jumping spider *P. paykulli* were collected from a rice field in Ningbo, Zhejiang Province, China. Species identification was confirmed by assessing its morphological characteristics using stereomicroscopy and CO1 sequence analysis (Supplementary Table 1). Jumping spiders were fed about their own mass of leafhoppers *N. cincticeps* three times each week and examined 14-52 days following collection, as previously reported (Taylor et al., 2014).

Electron microscopy

The Malpighian tubules of *N. cincticeps* were dissected using an astereomicroscope, serially fixed overnight at 4°C with 2.5% glutaraldehyde in 0.1 mol/L PBS, and post-fixed for 2 hours at room temperature with 1% OsO4. Following that, samples were dehydrated in a series of ethanol solutions (50%, 70%, 80%, 90%, and 95%), then permeabilized with 100% ethanol and 100% acetone, followed by a series of epoxies in acetone (50%, 75%, and 100%). The pierced tissue was then immersed in epoxy resin and baked for more than 24 hours at 70°C. Ultrathin slices were cut and stained with uranyl acetate for 15 minutes and lead citrate for 5 minutes, and examined by Hitachi H7800 microscope (Hitachi, Japan).

Scanning electron microscopy was used to observe the morphology of leafhopper brochosomes. Samples were prepared as previously described(Rakitov & Gorb, 2013a). Leafhopper forewings were obtained by dissection under the stereomicroscope and glued onto SEM metal stubs with a double-sided carbon tape. The dried samples were coated with gold

for 1 min in a MC1000 Ion Sputter Coater (Hitachi, Japan) and viewed by a HITACHI Regulus

8100 SEM (Hitachi, Japan).

Spider predation experiment

Jumping spiders were starved for three days prior to the predation experiment to ensure

that participating spiders were hungry enough to attack leafhoppers in our predation

preference tests, but not so hungry that they would attack the first prey they encountered

(without being choosy)(Taylor et al., 2014). For the jumping spiders to prey on, we placed two

leafhoppers in a 9 cm diameter test arena (the bottom of the dish was covered with filter paper).

Before starting the test, the spiders were placed in the arena for 15 minutes to adapt, and then

they were released and permitted to prey on the leafhoppers. We directly observed the feeding

process of the jumping spiders and recorded when the spiders attacked the first leafhopper

and which leafhopper they preyed on. The predation finishes when the first leafhopper is

attacked or when the jumping spider does not attack any leafhopper within 5 minutes. Each

pair of jumping spider predation tests were independently replicated more than 60 times.

The distribution of cuticle surface brochosomes and the optical properties of male and

female insects at various periods post-eclosion

To investigate the distribution of brochosomes on the cuticle surface of adult *N. cincticeps*

at various post-eclosion periods, we first collected female and male adult leafhoppers at 5-, 10-,

15-, 20-, and 25-days post-eclosion and dissected the forewings under a Nikon SMZ25

microscope (Nikon, Japan). The forewings were gold-sprayed and examined using HITACHI

Regulus 8100 SEM (Hitachi, Japan). At each time point, 30 samples were randomly selected

for SEM imaging. A 10 μm × 10 μm area within the SEM images was used to assess

brochosome coverage, and ImageJ was employed for analysis(Schneider et al., 2012). Since

brochosomes are synthesized at the distal section of the Malpighian tubule and the

morphology of the Malpighian tubule in leafhoppers can also reflect brochosomes synthesis,

we used Nikon SMZ25 microscope (Nikon, Japan) to examine the morphology of the

Malpighian tubule in female and male adult leafhoppers at 5-, 10-, 15-, 20-, and 25-days

post-eclosion.

To examine the optical properties of male and female insects at various periods

post-eclosion, we first collected female and male adult leafhoppers at 5-, 10-, 15-, 20-, and

25-days post-eclosion, freeze-killed them and placed them under white or UV light for

observation through a Nikon SMZ25 microscope (Nikon, Japan). Following that, the forewings

of the leafhoppers were meticulously dissected, and the specular reflectance spectra of the

forewings (25 × 25 μm) were precisely measured utilizing a 20/30 PV UV-Vis-NIR

microspectrophotometer (CRAIC Technologies Inc., USA). UV-Vis-NIR reflectance spectra

(250 to 1000 nm) were obtained using a 10× UV-absorbing glass objective lens, with a

minimum of five replicates per sample.

Identification of brochosome structural protein-coding genes

Candidate brochosome structural protein-coding genes were identified through a combined

analysis of transcriptomic and proteomic data. To knock down the expression of these

candidates, RNAi was conducted by microinjecting double-stranded RNA (dsRNA) into the

abdomens of fifth instar nymphs. The dsRNAs targeting 500-1000 bp regions of Malpighian

tubule-specific expression genes or GFP were synthesized in vitro using the T7 RiboMAX

Express RNAi System (Promega, USA). The 50 fifth instar nymphs of *N. cincticeps* were

microinjected with 30 nl dsRNA (0.5 μg μl^{-1}) using the Nanoject II Auto-Nanoliter Injector

(Drummond, USA). Thereafter, they were transferred to healthy rice seedlings for recovery.

Forewings of treated leafhoppers were collected 7 days after microinjection (about 4-5 days

post-eclosion), sprayed with gold, and analyzed by HITACHI Regulus 8100 SEM (Hitachi,

Japan). Screening identified four brochosome structural protein-coding genes, and inhibiting

their expression resulted in significant changes in brochosome morphology. The full-length

sequence of these genes was obtained using by 5'-rapid amplification of cDNA ends (RACE)

and 3'-RACE using SMARTer RACE 5'/3' Kit (Clontech, USA).

To analyze the temporal expression profiles of four brochosome structural protein-coding

genes (GenBank accession number PP273097, PP273098, PP273099, PP273100), the whole

body, salivary gland, midgut, ovary, and testis were dissected from 200 adult insects, total

RNA in different tissue were extracted using TRIzol Reagent (Invitrogen, USA). The relative

expression of these genes in different tissues of *N. cincticeps* adults was detected by

RT-qPCR assay using a QuantStudio^{TM} 5 Real-Time PCR system (Thermo Fisher Scientific,

USA). The detected transcript levels were normalized to the transcript level of the

housekeeping gene elongation factor 1 alpha (EF1α) (GenBank accession number AB836665)

and estimated by the 2−△△Ct (cycle threshold) method. Using the same procedure, the

expression levels of these brochosome structural protein-coding genes were evaluated in

female and male adult leafhoppers at 5-, 10-, 15-, 20-, and 25-days post-eclosion.

Knocking down in vivo expression of BSM-coding genes in *N. cincticeps*

Following the prior procedure, dsRNA produced from four BSM-coding genes was injected

into the abdomen of fifth instar nymphs of *N. cincticeps* by microinjection, and a control group

injected with ds*GFP* was established. Seven days after injection, the forewings and Malpighian

tubules of the ds*BSM* and ds*GFP* treated leafhoppers were dissected, and the morphology of

brochosomes on the cuticle surface of leafhoppers and the synthesis of brochosomes in

Malpighian tubules were examined by SEM and TEM, respectively. To examine the optical

properties, the ds*BSM* and ds*GFP* treated leafhoppers, freeze-killed them and placed them

under white or UV light for observation through a Nikon SMZ25 microscope (Nikon, Japan).

Subsequently, the forewings of the leafhoppers were meticulously dissected, and the specular

reflectance spectra of the forewings (25 × 25 μm) were precisely measured utilizing a 20/30

PV UV-Vis-NIR microspectrophotometer (CRAIC Technologies Inc., USA). UV-Vis-NIR

reflectance spectra (250 to 1000 nm) were obtained using a 10× UV-absorbing glass objective

lens, with a minimum of five replicates per sample. Finally, the predation preference of the

jumping spiders on the ds*BSM* and ds*GFP* treated leafhoppers was examined by a predation

experiment.

In vitro measurements were conducted to assess the relationship between brochosome

morphology and their optical performance.

To collect approximately 500 *N. cincticeps* treated with ds*GFP* or ds*BSM*, dissect the

forewings of the leafhoppers and immerse them in acetone in a 50 mL centrifuge tube. Place

the centrifuge tubes on a room temperature orbital shaker at 50 rpm for 12 hours to separate

the forewings from the leafhoppers due to friction. Following this, centrifuge at 1000g for 10

minutes to separate the brochosomes. Resuspend the particles in fresh acetone, briefly

sonicate, then centrifuge at 1000g for 10 minutes. Repeat this process three times.

Resuspend the collected brochosomes from ds*GFP* and ds*BSM*-treated leafhopper forewings

in acetone and adjust the two suspension solutions to the same OD280 reading. Next, using a

pipette, carefully drop the brochosome solution onto quartz slides and brown planthopper

wings. Allow the acetone to completely evaporate at room temperature. At the same time, we

also set up quartz slides and brown planthopper wings that were treated with acetone alone as

negative controls. Prepared quartz slides and brown planthopper wings can be observed

under SEM to examine the distribution of brochosomes. Additionally, UV-Vis-NIR reflectance

spectra (250 to 1000 nm) were obtained using a 20/30 PV UV-Vis-NIR

microspectrophotometer (CRAIC Technologies Inc., USA), with a minimum of five replicates

per sample.

Removal of leafhopper forewing brochosome by acetone

Previous research has shown that acetone may efficiently remove brochosomes from the

cuticle surface of leafhoppers(Rakitov et al., 2018). We manually removed the brochosomes

from the forewings of leafhoppers using acetone and examined the imaging under white or UV

light for observation through a Nikon SMZ25 microscope (Nikon, Japan). UV-Vis-NIR

reflectance spectra (250 to 1000 nm) were obtained using a 20/30 PV UV-Vis-NIR

microspectrophotometer (CRAIC Technologies Inc., USA), with a minimum of five replicates

per sample.

Bioinformatics analysis and phylogenetic tree

The brochosome is a unique secretion produced by leafhoppers (Cicadellidae). In order to

determine the distribution of brochosome structural protein-coding genes in Hemiptera. The

brochosome structural protein-coding genes were used to queries to search for homologous

sequences in Hemiptera transcriptomes. We examined 116 Hemiptera species,

encompassing major families (Acanaloniidae, Achilidae, Aetalionidae, Aphrophoridae,

Caliscelidae, Cercopidae, Cicadellidae, Cicadidae, Cixiidae, Clastopteridae, Delphacidae,

Derbidae, Dictyopharidae, Epipygidae, Eurybrachidae, Flatidae, Fulgoridae, Issidae,

Machaerotidae, Melizoderidae, Membracidae, Myerslopiidae, Nogodinidae, Peloridiidae, Ricaniidae, Tettigarctidae, Tettigometridae, Theaceae, Tropiduchidae). Transcriptome data of 116 Hemiptera species were downloaded from the NCBI Sequence Read Archive (Supplementary Table 2), and assembled using SOAPdenovo-Trans (version 1.01)(Xie et al., 2014). The resulting transcripts were filtered to remove potential contaminants. We compared four identified BSM-encoded genes with these transcriptomes using the tblastn (E-value < 1.0e–5). Heatmaps were generated using TBtools software(Chen et al., 2023). At least one homologous gene for the four BSM genes was identified in 66 species. A species phylogenetic tree for these 66 Hemiptera species was constructed following the methods outlined in previous studies(Johnson et al., 2018).

The amino acid sequences of BSM2 and BSM3 were aligned using the Clustal W multiple sequence alignment program(Larkin et al., 2007). AlphaFold2 was employed to predict the three-dimensional structures of BSM2 and BSM3, and structural alignments were conducted using PyMOL(DeLano, 2002; Jumper et al., 2021). In Membracoidea, the maximum likelihood (ML) phylogenetic tree for BSM2 and BSM3 was constructed using IQtree in PhyloSuite(Minh et al., 2020; Xiang et al., 2023).

Statistical analyses

All experiments were performed at least three independent replicates and statistical analyses were performed with GraphPad Prism8.0 software. Statistical significance was calculated by a two-tailed Student's t-test, one-way ANOVA, two-way ANOVA and/or unpaired Student's t-test. P-values < 0.05 were considered statistically significant.

Acknowledgments

This work received support from the National Natural Science Foundation of China (U23A6006, U20A2036, 32270150) and the Natural Science Foundation of Ningbo Municipality (2023J112). The author acknowledges Dr. Yuan Cheng from the Instrumentation and Service Center for Molecular Science (ISCMS) at Westlake University for assistance with optical measurement and data interpretation. We are grateful for the support provided by the Bioimaging Center of the Bioimaging Center, State Key Laboratory for Managing Biotic and

Chemical Threats to the Quality and Safety of Agro-products, Institute of Plant Virology,

Ningbo University, in TEM and SEM measurement and data interpretation. Additionally, we

thank Professor Shou-Wei Ding (University of California, Riverside, CA, United States) for his

valuable and constructive suggestions for manuscript improvement.

Author contributions:

W.W. J.-M.L., C.-X.Z., and J.-P.C. designed research; W.W. Q.M. Z.Y. Z.L. performed

research; W.W. and Z.Y. analyzed data; and W.W. H.-W.S., J.-M.L., C.-X.Z., and J.-P.C wrote

the paper. All authors gave final approval for publication. All authors have read and agreed to

the published version of the manuscript.

Data availability

All the data needed to understand and assess the conclusions of this research are available

in the article, Supplementary Materials, and GenBank (accession number PP273097,

PP273098, PP273099, PP273100). Any additional information required to reanalyse the data

reported in this paper is available from the lead contact upon request.

Declaration of interests

The authors declare no competing interests.

References

Andersen, S. O. (1979). Biochemistry of insect cuticle. *Annual review of entomology*, *24*(1), 29-59.

Banerjee, P., Burks, G. R., Bialik, S. B., Nassr, M., Bello, E., Alleyne, M., Freeman, B. D., Barrick, J. E., Schroeder, C. M., & Milliron, D. J. (2023). Nanostructure-Derived Antireflectivity in Leafhopper Brochosomes. *Advanced Photonics Research*, *4*(7), 2200343. https://doi.org/10.1002/adpr.202200343

Bello, E., Chen, Y., & Alleyne, M. (2022). Staying Dry and Clean: An Insect's Guide to Hydrophobicity. *Insects*, *14*(1), 42. https://doi.org/10.3390/insects14010042

Bernhard, C. G. (1965). The insect corneal nipple array. A biological, broad-band impedance transformer that acts as an antireflection coating. *Acta physiol. scand.*, *63*(243), 1-79.

Blagodatski, A., Sergeev, A., Kryuchkov, M., Lopatina, Y., & Katanaev, V. L. (2015). Diverse set of Turing nanopatterns coat corneae across insect lineages. *Proceedings of the National Academy of Sciences*, *112*(34), 10750-10755. https://doi.org/10.1073/pnas.1505748112

Burks, G. R., Yao, L., Kalutantirige, F. C., Gray, K. J., Bello, E., Rajagopalan, S., Bialik, S. B., Barrick, J. E., Alleyne, M., Chen, Q., & Schroeder, C. M. (2022). Electron Tomography and Machine Learning for Understanding the Highly Ordered Structure of Leafhopper Brochosomes. *Biomacromolecules*, *24*(1), 190-200. https://doi.org/10.1021/acs.biomac.2c01035

Chen, C., Wu, Y., Li, J., Wang, X., Zeng, Z., Xu, J., Liu, Y., Feng, J., Chen, H., He, Y., & Xia, R. (2023). TBtools-II: A "one for all, all for one" bioinformatics platform for biological big-data mining. *Molecular plant*, *16*(11), 1733-1742. https://doi.org/10.1016/j.molp.2023.09.010

Chotard, A., Ledamoisel, J., Decamps, T., Herrel, A., Chaine, A. S., Llaurens, V., & Debat, V. (2022). Evidence of attack deflection suggests adaptive evolution of wing tails in butterflies. *Proceedings of the Royal Society B: Biological Sciences*, *289*(1975), 20220562. https://doi.org/10.1098/rspb.2022.0562

Cinel, S. D., Hahn, D. A., & Kawahara, A. Y. (2020). Predator-induced stress responses in insects: A review. *Journal of insect physiology*, *122*, 104039. https://doi.org/10.1016/j.jinsphys.2020.104039

Cronin, T. W., & Bok, M. J. (2016). Photoreception and vision in the ultraviolet. *Journal of Experimental Biology*, *219*(18), 2790-2801. https://doi.org/10.1242/jeb.128769

Del-Claro, K., & Oliveira, P. S. (2000). Conditional outcomes in a neotropical treehopper-ant association: temporal and species-specific variation in ant protection and homopteran fecundity. *Oecologia*, *124*(2), 156-165. https://doi.org/10.1007/s004420050002

DeLano, W. L. (2002). Pymol: An open-source molecular graphics tool. *CCP4 Newsl. Protein Crystallogr*, *40*(1), 82-92.

Dietrich, C., Rakitov, R., Holmes, J., & Black IV, W. (2001). Phylogeny of the major lineages of Membracoidea (Insecta: Hemiptera: Cicadomorpha) based on 28S rDNA sequences. *Molecular Phylogenetics and Evolution*, *18*(2), 293-305.

Endler, J. A. (1997). Some general comments on the evolution and design of animal communication systems. *Philosophical Transactions of the Royal Society of London.*

Series B: Biological Sciences, *340*(1292), 215-225. https://doi.org/10.1098/rstb.1993.0060

Galloway, J. A. M., Green, S. D., Stevens, M., & Kelley, L. A. (2020). Finding a signal hidden among noise: how can predators overcome camouflage strategies? *Philosophical Transactions of the Royal Society B: Biological Sciences*, *375*(1802), 20190478. https://doi.org/10.1098/rstb.2019.0478

Gouranton, J., & Maillet, P.-L. (1967). Origine et structure des brochosomes. *Journal de Microscopie*, *6*, 53-64.

Hibino, H. (1996). Biology and Epidemiology of Rice Viruses. *Annual review of phytopathology*, *34*(1), 249-274. https://doi.org/10.1146/annurev.phyto.34.1.249

Ho, S., Schachat, S. R., Piel, W. H., & Monteiro, A. (2016). Attack risk for butterflies changes with eyespot number and size. *Royal Society open science*, *3*(1), 150614. https://doi.org/10.1098/rsos.150614

Hughes, A., Liggins, E., & Stevens, M. (2019). Imperfect camouflage: how to hide in a variable world? *Proceedings of the Royal Society B: Biological Sciences*, *286*(1902), 20190646. https://doi.org/10.1098/rspb.2019.0646

Johnson, K. P., Dietrich, C. H., Friedrich, F., Beutel, R. G., Wipfler, B., Peters, R. S., Allen, J. M., Petersen, M., Donath, A., Walden, K. K. O., Kozlov, A. M., Podsiadlowski, L., Mayer, C., Meusemann, K., Vasilikopoulos, A., Waterhouse, R. M., Cameron, S. L., Weirauch, C., Swanson, D. R., . . . Yoshizawa, K. (2018). Phylogenomics and the evolution of hemipteroid insects. *Proceedings of the National Academy of Sciences*, *115*(50), 12775-12780. https://doi.org/10.1073/pnas.1815820115

Jones, M. P., Pierce, K. E., & Ward, D. (2007). Avian Vision: A Review of Form and Function with Special Consideration to Birds of Prey. *Journal of exotic pet medicine*, *16*(2), 69-87. https://doi.org/10.1053/j.jepm.2007.03.012

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., & Potapenko, A. (2021). Highly accurate protein structure prediction with AlphaFold. *Nature*, *596*(7873), 583-589.

Kryuchkov, M., Bilousov, O., Lehmann, J., Fiebig, M., & Katanaev, V. L. (2020). Reverse and forward engineering of Drosophila corneal nanocoatings. *Nature*, *585*(7825), 383-389. https://doi.org/10.1038/s41586-020-2707-9

Larkin, M. A., Blackshields, G., Brown, N. P., Chenna, R., McGettigan, P. A., McWilliam, H., Valentin, F., Wallace, I. M., Wilm, A., Lopez, R., Thompson, J. D., Gibson, T. J., & Higgins, D. G. (2007). Clustal W and Clustal X version 2.0. *Bioinformatics*, *23*(21), 2947-2948. https://doi.org/10.1093/bioinformatics/btm404

Lei, C.-W., Chen, R.-Y., & Yang, H. (2020). Leafhopper Wing-Inspired Broadband Omnidirectional Antireflective Embroidered Ball-Like Structure Arrays Using a Nonlithography-Based Methodology. *Langmuir*, *36*(19), 5296-5302. https://doi.org/10.1021/acs.langmuir.0c00634

Li, D., & Lim, M. L. M. (2005). Ultraviolet cues affect the foraging behaviour of jumping spiders. *Animal Behaviour*, *70*(4), 771-776. https://doi.org/10.1016/j.anbehav.2004.12.021

Li, Z., Li, Y., Xue, A. Z., Dang, V., Holmes, V. R., Johnston, J. S., Barrick, J. E., Moran, N. A., & Parsch, J. (2022). The Genomic Basis of Evolutionary Novelties in a Leafhopper.

Molecular biology and evolution, *39*(9), msac184. https://doi.org/10.1093/molbev/msac184

Light, S., Basile, W., & Elofsson, A. (2014). Orphans and new gene origination, a structural and evolutionary perspective. *Current Opinion in Structural Biology*, *26*, 73-83. https://doi.org/10.1016/j.sbi.2014.05.006

Lim, U. T., & Ben-Yakir, D. (2020). Visual sensory systems of predatory and parasitic arthropods. *Biocontrol Science and Technology*, *30*(7), 728-739. https://doi.org/10.1080/09583157.2020.1752362

Lin, M., Yang, G., Wang, Z., & You, M. (2021). Brochosome detachment facilitates Empoasca onukii (Hemiptera: Cicadellidae) adults escaping from spider web. *Acta Entomol Sin*, *64*, 213-222.

Liu, S., Li, Z., Sui, Y., Schaefer, D. A., Alele, P. O., Chen, J., & Yang, X. (2015). Spider foraging strategies dominate pest suppression in organic tea plantations. *BioControl*, *60*(6), 839-847. https://doi.org/10.1007/s10526-015-9691-2

Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., Von Haeseler, A., & Lanfear, R. (2020). IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. *Molecular biology and evolution*, *37*(5), 1530-1534.

Monteiro, A., Rafi, F. Z., Chan, I. Z. W., & Fiedler, K. (2019). Interacting Effects of Eyespot Number and Ultraviolet Reflectivity on Predation Risk in Bicyclus anynana (Lepidoptera: Nymphalidae). *Journal of Insect Science*, *19*(6). https://doi.org/10.1093/jisesa/iez123

Morehouse, N. I., Buschbeck, E. K., Zurek, D. B., Steck, M., & Porter, M. L. (2017). Molecular Evolution of Spider Vision: New Opportunities, Familiar Players. *Biological Bulletin*, *233*(1), 21-38. https://doi.org/10.1086/693977

Nelson, A. S., & Mooney, K. A. (2022). The Evolution and Ecology of Interactions Between Ants and Honeydew-Producing Hemipteran Insects. *Annual Review of Ecology, Evolution, and Systematics*, *53*(1), 379-402. https://doi.org/10.1146/annurev-ecolsys-102220-014840

Oraze, M. J., & Grigarick, A. A. (1989). Biological control of aster leafhopper (Homoptera: Cicadellidae) and midges (Diptera: Chironomidae) by Pardosa ramulosa (Araneae: Lycosidae) in California rice fields. *Journal of Economic Entomology*, *82*(3), 745-749.

Rakitov, R., & Gorb, S. N. (2013a). Brochosomal coats turn leafhopper (Insecta, Hemiptera, Cicadellidae) integument to superhydrophobic state. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1752), 20122391. https://doi.org/10.1098/rspb.2012.2391

Rakitov, R., & Gorb, S. N. (2013b). Brochosomes protect leafhoppers (Insecta, Hemiptera, Cicadellidae) from sticky exudates. *Journal of the Royal Society Interface*, *10*(87), 20130445. https://doi.org/10.1098/rsif.2013.0445

Rakitov, R., Moysa, A. A., Kopylov, A. T., Moshkovskii, S. A., Peters, R. S., Meusemann, K., Misof, B., Dietrich, C. H., Johnson, K. P., Podsiadlowski, L., & Walden, K. K. O. (2018). Brochosomins and other novel proteins from brochosomes of leafhoppers (Insecta, Hemiptera, Cicadellidae). *Insect biochemistry and molecular biology*, *94*, 10-17. https://doi.org/10.1016/j.ibmb.2018.01.001

Rakitov, R. A. (1996). Post-moulting behaviour associated with Malpighian tubule secretions in leafhoppers and treehoppers (Auchenorrhyncha: Membracoidea). *European Journal of Entomology*, *93*(2), 167-184. <Go to ISI>://WOS:A1996VB52200006

Rakitov, R. A. (2000). Secretion of brochosomes during the ontogenesis of a leafhopper, Oncometopia orbona (F.) (Insecta, Homoptera, Cicadellidae). *Tissue and Cell*, *32*(1), 28-39. https://doi.org/10.1054/tice.1999.0084

Rakitov, R. A. (2009). Brochosomal Coatings of the Integument of Leafhoppers (Hemiptera, Cicadellidae). In S. N. Gorb (Ed.), *Functional Surfaces in Biology* (pp. 113-137). Springer Netherlands. https://doi.org/10.1007/978-1-4020-6697-9_8

Raut, H. K., Ganesh, V. A., Nair, A. S., & Ramakrishna, S. (2011). Anti-reflective coatings: A critical, in-depth review. *Energy & Environmental Science*, *4*(10), 3779-3804. https://doi.org/10.1039/c1ee01297e

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nature methods*, *9*(7), 671-675.

Sheikh, A. A., Rehman, N., & Kumar, R. (2017). Diverse adaptations in insects: A review. *Journal of entomology and zoology studies*, *5*(2), 343-350.

Silberglied, R. E. (1979). Communication in the ultraviolet. *Annual Review of Ecology and Systematics*, *10*(1), 373-398. https://www.annualreviews.org/doi/abs/10.1146/annurev.es.10.110179.002105

Skelhorn, J., & Rowe, C. (2016). Cognition and the evolution of camouflage. *Proceedings of the Royal Society B: Biological Sciences*, *283*(1825), 20152890. https://doi.org/10.1098/rspb.2015.2890

Stavenga, D. G., Foletti, S., Palasantzas, G., & Arikawa, K. (2005). Light on the moth-eye corneal nipple array of butterflies. *Proceedings of the Royal Society B: Biological Sciences*, *273*(1587), 661-667. https://doi.org/10.1098/rspb.2005.3369

Stevens, M. (2007). Predator perception and the interrelation between different forms of protective coloration. *Proceedings of the Royal Society B: Biological Sciences*, *274*(1617), 1457-1464. https://doi.org/10.1098/rspb.2007.0220

Stevens, M., & Ruxton, G. D. (2018). The key role of behaviour in animal camouflage. *Biological Reviews*, *94*(1), 116-134. https://doi.org/10.1111/brv.12438

Stobbe, N., Dimitrova, M., Merilaita, S., & Schaefer, H. M. (2008). Chromaticity in the UV/blue range facilitates the search for achromatically background-matching prey in birds. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *364*(1516), 511-517. https://doi.org/10.1098/rstb.2008.0248

Tautz, D., & Domazet-Lošo, T. (2011). The evolutionary origin of orphan genes. *Nature Reviews Genetics*, *12*(10), 692-702. https://doi.org/10.1038/nrg3053

Taylor, L. A., Maier, E. B., Byrne, K. J., Amin, Z., & Morehouse, N. I. (2014). Colour use by tiny predators: jumping spiders show colour biases during foraging. *Animal Behaviour*, *90*, 149-157. https://doi.org/10.1016/j.anbehav.2014.01.025

Thery, M., & Gomez, D. (2010). Insect colours and visual appearance in the eyes of their predators. *Advances in insect physiology*, *38*, 267-353.

Tovee, M. (1995). Ultra-violet photoreceptors in the animal kingdom: their distribution and function. *Trends in ecology & evolution*, *10*(11), 455-460. https://doi.org/10.1016/s0169-5347(00)89179-x

Vukusic, P., & Sambles, J. R. (2003). Photonic structures in biology. *Nature*, *424*(6950),
852-855. https://doi.org/10.1038/nature01941

Walker, S. E., & Rypstra, A. L. (2002). Sexual dimorphism in trophic morphology and feeding
behavior of wolf spiders (Araneae: Lycosidae) as a result of differences in reproductive
roles. *Canadian Journal of Zoology*, *80*(4), 679-688. https://doi.org/10.1139/z02-037

Watson, G. S., Watson, J. A., & Cribb, B. W. (2017). Diversity of Cuticular Micro- and
Nanostructures on Insects: Properties, Functions, and Potential Applications. *Annual*
Review of Entomology, Vol 62, *62*, 185-205.
https://doi.org/10.1146/annurev-ento-031616-035020

Wei, J., Jia, D., Mao, Q., Zhang, X., Chen, Q., Wu, W., Chen, H., & Wei, T. (2018). Complex
interactions between insect-borne rice viruses and their vectors. *Current opinion in*
virology, *33*, 18-23. https://doi.org/10.1016/j.coviro.2018.07.005

Wong, E. S. W., & Belov, K. (2012). Venom evolution through gene duplications. *Gene*, *496*(1),
1-7. https://doi.org/10.1016/j.gene.2012.01.009

Wu, W., Ye, Z., Mao, Q., Shan, H.-W., Li, J.-M., & Chen, J.-P. (2023). Combined Transcriptome
and Proteome Analysis of the Protein Composition of the Brochosomes of the
Leafhopper Nephotettix cincticeps. *Insects*, *14*(10), 784.
https://doi.org/10.3390/insects14100784

Xi, W., Lee, Y.-J., Yu, S., Chen, Z., Shiomi, J., Kim, S.-K., & Hu, R. (2023). Ultrahigh-efficient
material informatics inverse design of thermal metamaterials for
visible-infrared-compatible camouflage. *Nature Communications*, *14*(1), 4694.
https://doi.org/10.1038/s41467-023-40350-6

Xiang, C. Y., Gao, F., Jakovlić, I., Lei, H. P., Hu, Y., Zhang, H., Zou, H., Wang, G. T., & Zhang,
D. (2023). Using PhyloSuite for molecular phylogeny and tree-based analyses. *iMeta*,
2(1), e87.

Xie, Y., Wu, G., Tang, J., Luo, R., Patterson, J., Liu, S., Huang, W., He, G., Gu, S., Li, S., Zhou,
X., Lam, T.-W., Li, Y., Xu, X., Wong, G. K.-S., & Wang, J. (2014). SOAPdenovo-Trans:
de novo transcriptome assembly with short RNA-Seq reads. *Bioinformatics*, *30*(12),
1660-1666. https://doi.org/10.1093/bioinformatics/btu077

Yan, B., Yu, X., Dai, R., Li, Z., & Yang, M. (2021). Chromosome-level genome assembly of
Nephotettix cincticeps (Uhler, 1896)(Hemiptera: Cicadellidae: Deltocephalinae).
Genome Biology and Evolution, *13*(11), evab236.

Yang, H., Peng, Y., Tian, J., Wang, J., Wei, B., Xie, C., & Wang, Z. (2018). Rice Field Spiders
in China: A Review of the Literature. *Journal of Economic Entomology*, *111*(1), 53-64.
https://doi.org/10.1093/jee/tox319

Yang, S. K., Sun, N., Stogin, B. B., Wang, J., Huang, Y., & Wong, T. S. (2017).
Ultra-antireflective synthetic brochosomes. *Nature Communications*, *8*(1), 1285.
https://doi.org/10.1038/s41467-017-01404-8

Yuan, F., Su, M., Li, T., Zhang, Y., Dietrich, C. H., Webb, M. D., & Wei, C. (2023). Functional
and evolutionary implications of protein and metal content of leafhopper brochosomes.
Insect biochemistry and molecular biology, *157*, 103962.
https://doi.org/10.1016/j.ibmb.2023.103962

Zou, Y., Araujo, D. P., Lim, M. L. M., & Li, D. (2011). Ultraviolet is a more important cue than
reflection in other wavelengths for a jumping spider to locate its spider prey. *Animal*
Behaviour, *82*(6), 1457-1463. https://doi.org/10.1016/j.anbehav.2011.09.031
840
841

Figure 1. Brochosomes are a distinctive coating on the cuticle surface of leafhopper.

(A) Male adult of the green rice leafhopper *N. cincticeps*. Bar, 1 mm. (B) Brochosomes on the surface of the forewing of *N. cincticeps*. Bar, 1 μm. (C, D) The morphologies of brochosomes by SEM (C) and TEM (D). Bar, 200 nm. (E) Alimentary tract and the Malpighian tubules of *N. cincticeps*. The leafhopper *N. cincticeps* have two pairs of Malpighian tubules, each divided into proximal segment, distal segment, and terminal segment. The brochosomes are synthesized in the distal segment, which is dilated and rod-shaped. Bar, 1 mm. (F) The distal segment epithelial cell displays an extensive rough endoplasmic reticulum and multiple Golgi regions (red arrow) containing developing brochosomes in its basal portion, as well as a number of secretory vacuoles with mature brochosomes near the cell border. Bar, 5 μm. (G) The initial stage of the development of brochosome. Bar, 500 mm. (H) The two brochosomes that are developing inside the primary vesicles are shown in close-up. Regular invaginations appear on the surface of the growing brochosome at the same time that its matrix separates into a looser core and a denser wall. Bar, 500 nm. (I) Larger vesicles containing numerous BS, formed by the fusion of multiple primary vesicles. The surface of the brochosomes has regular cell-like invaginations, and it is surrounded by amorphous flocculent material with a moderate

electron density. Bar, 1 µm. (J) A vesicle filled with mature brochosomes, each mature

brochosome has a spherical inner cavity and a well-swollen outer margin of the septa. Bar, 1

µm. E, F, G, H are the enlargement of boxed area in D. Fc, filter chamber; Mg, midgut; Mt,

Malpighian tubules; mtd, distal segment of the Malpighian tubule; mtp, proximal segment of

the Malpighian tubule; mtt, terminal segment of the Malpighian tubule. All images are

representative of at least three replicates.

Figure 2. The distribution of brochosomes on the cuticle surface of leafhopper *N. cincticeps* are associated with predation by jumping spiders.

(A, B) Images of leafhopper *N. cincticeps* males and females in white (A) and ultraviolet light (B) at 5 and 25 days post-eclosion, respectively. Bar, 1 mm. (C, D) The distribution of brochosomes on the surface of a forewing (C) and morphological changes of the Malpighian

tubules (D) of *N. cincticeps* males and females at 5 and 25 days post-eclosion. Bar, 5 μm in C;

Bar, 1 mm in D. (E) Reflectance spectra of female and male forewing of *N. cincticeps* at 5 and

25 days post-eclosion. (F) Images of the jumping spider *P. paykulli*. Bar, 2 mm. (G-J) Jumping

spiders prefer leafhoppers with little brochosome covering as food. In predation experiment,

jumping spiders offered *N. cincticeps* male and female at 5 days post-eclosion (G), male and

female at 25 days post-eclosion (H), males at 5 and 25 days post-eclosion (I), and females at 5

and 25 days post-eclosion (J). Data on predation times are displayed using the traditional box

and whisker shapes. All box plots with whiskers represent the data distribution based on five

number summary statistics (maximum, third quartile, median, first quartile, minimum), each dot

in box plot represents an independent experiment. **$P < 0.01$, ****$P < 0.0001$, ns no

significance, Statistical significance was determined by unpaired t test with Welch's correction

method. Predation preference is shown in the pie chart. All images are representative of at

least three replicates.

Figure 3. Identification of brochosome structural proteins. (A) Morphological of
brochosomes on the forewing of leafhopper *N. cincticeps* at 7 days post-microinjection of
dsRNA. Bar, 500 nm. (B) Gene structures of *BSM-1*, *BSM-2*, *BSM-3* and *BSM-4*. (C-F)
Transcript levels of *BSM-1* (C), *BSM-2* (D), *BSM-3* (E) and *BSM-4* (F) at 7 days
post-microinjection of dsRNA. Each data point represents the result of one independent
experiment. (G-J) The abundance of BSM transcripts in different tissues and whole bodies of
N. cincticeps was determined by RT-qPCR. Notably, *BSM-1* (G), *BSM-2* (H), *BSM-3* (I), and
BSM-4 (J) exhibited specific expression in the Malpighian tubules. Each data point represents
the result of one independent experiment. (K-N) The expression patterns of *BSM-1* (K), *BSM-2*
(L), *BSM-3* (M) and *BSM-4* (N) transcripts were examined in male and female leafhopper at 5,

897   15, and 25 days post-eclosion. Results were obtained from 3 independent experiments. For

898   C-N, data shown are mean ± SD values. *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001; ns

899   no significance (C-F, two tailed Student's t-test; G-J, one-way ANOVA; K-N, two-way ANOVA).

900   All images are representative of at least three replicates.

Figure 4. RNAi inhibits brochosome synthesis, alters brochosome morphology, and influences predation in jumping spiders. (A) Images of leafhopper *N. cincticeps* males and females in white and ultraviolet light after ds*GFP* or ds*BSM* treatment, respectively. Bar, 1 mm. (B, C) Morphology of the brochosome on the forewings of leafhoppers after ds*GFP* (B) and

ds*BSM* (C) treatment. Bar, 2 µm. (D, E) Morphology of the brochosome in the distal segment

epithelial cells of Malpighian tubules after ds*GFP* (D) and ds*BSM* (E) treatment. Bar, 2 µm. (F)

The transcript levels of *BSM-1*, *BSM-2*, *BSM-3* and *BSM-4* at 7 days after ds*GFP* and ds*BSM*

treatment. Each data point represents the outcome of an individual independent experiment.

The presented data are expressed as mean ± SD values. Statistical significance is denoted as

*$P < 0.05$ and **$P < 0.01$, determined by two-way ANOVA. (G) Reflectance spectra of female

and male forewing of *N. cincticeps* at 7 days after ds*GFP* and ds*BSM* treatment. (H and I)

Jumping spiders prefer to prey on ds*BSM*-treated leafhoppers. Predation efficiency and

preference of jumping spiders on males (H) and females (I) *N. cincticeps* after ds*GFP* and

ds*BSM* treatment in the predation experiment. Data on predation times are displayed using the

traditional box and whisker shapes. All box plots with whiskers represent the data distribution

based on five number summary statistics (maximum, third quartile, median, first quartile,

minimum), each dot in box plot represents an independent experiment. **$P < 0.01$, ***$P < 0.001$,

Statistical significance was determined by unpaired t test with Welch's correction method.

Predation preference is shown in the pie chart. (J, K) The morphology of brochosomes is

related to their optical performance. Collect brochosomes treated with ds*GFP* and ds*BSM*

separately, apply them to brown planthopper wings (J) and quartz slides (K), and set up a

control group with acetone application only. All images are representative of at least three

replicates.

Figure 5. Brochosome coating on Cicadellidae cuticle surface is essential for their optical qualities. (A) Images of forewings of four leafhopper species before and after acetone treatment in white and ultraviolet light. Bar, 1mm. (B) Reflection spectra of the forewings of the *N. cincticeps* before and after acetone treatment. (C) Phylogenetic associations between the BSM and phylogeny of Hemipteran lineages. The left column describes the phylogeny of 66 representative species of Hemiptera with BSM proteins. In the right column, the presence of BSM- encoded genes in different Hemiptera species is illustrated along with their homology

analysis with *N. cincticeps*. White indicates the absence of related genes; the color gradient

represents the degree of sequence homology. All images are representative of at least three

replicates.

Fig. 6. Brochosomes serve as an anti-reflective camouflage coating on the cuticle

surface of leafhoppers. Brochosomes effectively reduce the reflection of various

wavelengths of light, particularly in the UV region. UV light is a crucial visual cue for numerous

visual predators to identify and locate prey. Brochosomes efficiently decrease the UV light

reflection on the surface of leafhoppers, thereby reducing their exposure risk to visual

predators and facilitating evasion of visual predation.