Redistribution of fragmented mitochondria ensure symmetric organelle

partitioning and faithful chromosome segregation in mitotic mouse zygotes

Haruna Gekko[1†], Ruri Nomura[1†], Daiki Kuzuhara[2], Masato Kaneyasu[1], Genpei Koseki[1]

Deepak Adhikari[3], Yasuyuki Mio[2], John Carroll[3], Tomohiro Kono[4], Hiroaki Funahashi[1]

and Takuya Wakai[1]*

[1]Department of Animal Science, Graduate School of Environment and Life Science,

Okayama University, Okayama 700-8530, Japan

[2]Reproductive Centre, Mio Fertility Clinic, Tottori, 683-0008, Japan

[3]Development and Stem Cell Program and Department of Anatomy and Developmental

Biology, Monash Biomedicine Discovery Institute, Monash University, Melbourne,

Victoria 3800, Australia

[4]Department of Bioscience, Tokyo University of Agriculture, Tokyo 156-8502, Japan

[†]These authors equally contributed to this work.

*Correspondence: t2wakai@okayama-u.ac.jp

Impact statement:

Depletion of Dynamin-related protein 1, a key regulator of mitochondrial fission, in

mouse zygotes impair symmetric organelle partitioning and chromosome segregation

leading to early developmental arrest.

Funding information

1) Japan Society for the Promotion of Science (Grant reference: 17K17905, 20K06627

and 23K0886905): Takuya Wakai

2) National Health and Medical Research Council (NHMRC) Australia (Grant

reference: 1165627): Deepak Adhikari and John Carroll

The funders had no role in study design, data collection and interpretation, or the

decision to submit the work for publication.

Author information:

Haruna Gekko, Department of Animal Science, Graduate School of Environment and

Life Science, Okayama University, Okayama, Japan, Contribution: Formal analysis;

Investigation, No competing interests declared.

Ruri Nomura , Department of Animal Science, Graduate School of Environment and

Life Science, Okayama University, Okayama, Japan, Contribution: Formal analysis;

Investigation, No competing interests declared.

Daiki Kuzuhara, Reproductive Centre, Mio Fertility Clinic, Tottori, Japan, Contribution:

Investigation; Data curation, No competing interests declared.

Masato Kaneyasu, Department of Animal Science, Graduate School of Environment

and Life Science, Okayama University, Okayama, Japan, Contribution: Data curation;

Visualisation, No competing interests declared.

Genpei Koseki, Department of Animal Science, Graduate School of Environment and

Life Science, Okayama University, Okayama, Japan, Contribution: Formal analysis;

Data curation, No competing interests declared.

Deepak Adhikari, Development and Stem Cell Program and Department of Anatomy

and Developmental, Biology, Monash Biomedicine Discovery Institute, Monash

University, Melbourne, Australia, Contribution: Formal analysis; Investigation; Funding

acquisition

No competing interests declared.

Yasuyuki Mio, Reproductive Centre, Mio Fertility Clinic, Tottori, Japan, Contribution:

Resources, No competing interests declared.

John Carroll, Development and Stem Cell Program and Department of Anatomy and

Developmental, Biology, Monash Biomedicine Discovery Institute, Monash University,

Melbourne, Australia, Contribution: Writing – review & editing; Supervision; Funding

acquisition, No competing interests declared.

Tomohiro Kono, Department of Bioscience, Tokyo University of Agriculture, Tokyo

156-8502, Japan, Contribution: Supervision, No competing interests declared.

Hiroaki Funahashi, Department of Animal Science, Graduate School of Environment

and Life Science, Okayama University, Okayama, Japan, Contribution: Resources;

Supervision, No competing interests declared.

Takuya Wakai

Department of Animal Science, Graduate School of Environment and Life Science,

Okayama University, Okayama, Japan, Contribution: Conceptualization, Validation;

Formal analysis; Investigation; Data curation; Writing - original draft, Supervision,

Funding acquisition, No competing interests declared.

Abstract

In cleavage-stage embryos, preexisting organelles partition evenly into daughter blastomeres without significant cell growth after symmetric cell division. The presence of mitochondrial DNA within mitochondria and its restricted replication during preimplantation development makes their inheritance particularly important. While chromosomes are precisely segregated by the mitotic spindle, the mechanisms controlling mitochondrial partitioning remain poorly understood. In this study, we investigate the mechanism by which Dynamin-related protein 1 (Drp1) controls the mitochondrial redistribution and partitioning during embryonic cleavage. Depletion of Drp1 in mouse zygotes causes marked mitochondrial aggregation, and the majority of embryos arrest at the 2-cell stage. Clumped mitochondria are located in the center of mitotic Drp1-depleted zygotes with less uniform distribution, thereby preventing their symmetric partitioning. Asymmetric mitochondrial inheritance is accompanied by functionally inequivalent blastomeres with biased ATP and endoplasmic reticulum Ca^{2+} levels. We also find that marked mitochondrial centration in Drp1-depleted zygotes prevents the assembly of parental chromosomes, resulting in chromosome segregation defects and binucleation. Thus, mitochondrial fragmentation mediated by Drp1 ensure proper organelle positioning and partitioning into functional daughters during the first embryonic cleavage.

Keywords: Mitochondrial dynamics, dynamin-related protein 1, embryo cleavage, preimplantation development, organelle inheritance, ER calcium stores, chromosome segregation, binuclear formation, assisted reproductive technologies

Introduction

Preimplantation development of mammalian embryos consists of a series of symmetric

cell divisions without significant cell growth, known as embryonic cleavage, resulting

in approximately half of the cytoplasm in each resulting cell, called blastomere. The

segregation of maternal content between daughter blastomeres allows rapid division of

a single-cell zygote into multicellular organisms. To ensure the correct segregation of

cellular content to produce functional daughters, cells must duplicate and apportion

their various organelles with high accuracy (Carlton et al., 2020). Chromosomes are

assembled and segregated into daughter cells with the formation of mitotic spindles,

whereas membrane-bound organelles, such as the endoplasmic reticulum (ER) and

mitochondria, are typically generated from existing structures, and both daughters

receive a share of these components during cell division. This is particularly important

in the case of mitochondria because mitochondria are semi-autonomous organelles that

contain their own genome, mitochondrial DNA (mtDNA). Growing oocytes

exponentially increase their mtDNA, and a fully grown oocyte contains more than 200

000 copies of mtDNA (Mahrous et al., 2012). After fertilization, abundant preexisting

mtDNA is segregated among daughter cells without replication during preimplantation

development (Poulton et al., 2010). As mitochondria are the primary source of ATP

production via oxidative phosphorylation in oocytes and early embryos owing to low

glycolytic activity (Dumollard et al., 2007), proper inheritance must be ensured for cell

energetic homogeneity among blastomeres. Mitochondria also play an important role in

the regulation of intracellular Ca^{2+} signaling, which in turn controls cell metabolism and

cell death pathways (Rizzuto et al., 2012). Thus, defects in mitochondrial inheritance

are likely to have serious consequences for daughter blastomeres and manifest as the

failure of pre- or post-implantation development.

Mitochondria are highly dynamic organelles that move along cytoskeletal networks

and undergo continuous fusion and fission; these dynamics allow cells to respond and

adapt to various intracellular and extracellular changes and maintain cellular function

and homeostasis (Mishra and Chan, 2014; Westermann, 2010). Active partitioning of

mitochondria is observed in budding yeast fission, which undergoes asymmetric cell

division; mitochondria actively move along actin filaments into the daughter cell

(Mishra and Chan, 2014). In mammalian cells that divide symmetrically, the

partitioning of mitochondria is largely due to passive processes, as fragmented and

dispersed mitochondrial distribution allows for stochastic and roughly equal partitioning

into daughter cells. Mitochondrial fission is thus important for faithful inheritance and

proper intracellular distribution of the mitochondria (Westermann, 2010). Whether

active transport of mitochondria by the cytoskeleton contributes to mitochondrial

inheritance in dividing animal cells remains controversial. Microtubules function as

cytoskeletal platforms for the distribution and transport of mitochondria. During mitosis,

mitochondria decouple from spindle microtubules and disperse in the cell periphery,

allowing for passive segregation (Chung et al., 2016). Furthermore, recent studies have

shown that actin filaments take over as the dominant mitochondrial scaffold following

mitochondrial release from microtubules at the start of mitosis (Moore et al., 2021).

Myosin 19 (Myo19), a mitochondrial-localized myosin in vertebrates, acts as an

actin-based motor for mitochondrial dynamics. Notably, depletion of Myo19 causes

asymmetrical mitochondrial inheritance between daughter cells (Rohn et al., 2014),

though the molecular characteristics of Myo19 in mammalian oocytes/eggs has yet to

be shown.

Dynamin-related protein 1 (Drp1) is a key regulator of mitochondrial fission in

many eukaryotic organisms. Drp1 is recruited to the mitochondrial membrane where it

forms helical oligomers that induce membrane constriction and severing (Westermann,

2010). The indispensable role of Drp1-mediated mitochondrial fission in mammalian

oocytes has been documented using conditional knockout (KO) mice (Udagawa et al.,

2014). Drp1 KO oocytes undergo meiotic arrest due to severe organelle aggregation,

which complicates the functional analysis of Drp1 in mature oocytes and embryos.

More recently, mature oocytes have been successfully isolated from juvenile Drp1

conditional KO mice (Adhikari et al., 2022). After parthenogenetic activation of Drp1

KO oocytes, few embryos develop to the blastocyst stage. Nevertheless, the precise role

of mitochondrial fission in normal embryos remains elusive. To this end, we employed

Trim-Away, a method to degrade endogenous proteins via the proteasome-mediated

pathway (Clift et al., 2017), which acutely depletes large amounts of storage proteins in

the oocyte, allowing loss-of-function studies during the preimplantation development

from the zygotic stage. In this study, we show that Drp1 depletion in fertilized zygotes

causes marked mitochondrial aggregation and early embryonic arrest. Live imaging of

mitochondria during the first cleavage division reveals that loss of Drp1 disturbs the

spatiotemporal mitochondrial dynamics, resulting in an asymmetric mitochondrial

inheritance between daughter blastomeres with functional heterogeneities. We also find

that misplaced mitochondria impair the assembly of parental spindles leading to

binucleated blastomeres, which closely resemble a clinical phenotype of human

embryos in *in vitro* fertilization (IVF) procedures (Hardy et al., 1993; Meriano et al.,

2004; Seikkula et al., 2018)

Results

Fragmented mitochondria are redistributed during the first embryonic cleavage

and equally partitioned into daughter blastomeres

We first imaged mitochondria and chromosomes in zygotes expressing

mitochondrially-targeted GFP (mt-GFP) and histone H2B-mCherry (**Figure 1A; Figure**

1-Video 1). Mitochondria in interphase zygotes (27 h post-hCG) progressively

accumulated around the two pronuclei (**Figure 1B, left**), encircled metaphase (31 h

post-hCG) chromosomes after nuclear envelope breakdown (NEB, 30 h post-hCG).

During the transition from anaphase to telophase (32 h post-hCG), the cytokinetic

furrow constricts the mitochondrial ring that effectively divides the mitochondria

approximately equally into the two daughter blastomeres and mitochondria were

dispersed throughout the cytoplasm in the interphase 2-cell embryos (35 h post-hCG)

(**Figure 1B, right**). Immunofluorescence staining of microtubules also showed

mitochondria progressively surround the spindle and disperse back into the cytoplasm

after the first cleavage (**Figure 1-figure supplement 1A**). The accumulation of

mitochondria around the spindle is unique to the first cleavage division (**Figure**

1-figure supplement 1B, 1C), as live imaging of fluorescently labeled microtubules

(EB3-GFP) and mitochondria (mt-DsRed) confirmed spindle-associated accumulation

of mitochondria in one-cell zygotes, but no evidence of mitochondrial accumulation

was observed in the spindles of 2-cell or 4-cell stage embryos. By electron microscopy

(EM), we found that these accumulated mitochondria around the metaphase spindle

were highly fragmented (**Figure 1C**). The ERs were also abundantly distributed around

the spindle and were in partial contact with the mitochondria, and enriched bundles of

actin filaments were found around these organelles.

Interphase zygotes stained with fluorescence-labelled phalloidin displayed a

mitochondrial distribution pattern closely associated with F-actin cytoplasmic

meshwork (**Figure 1D; Figure 1-figure supplement 1D**). This cytoplasmic meshwork

was subsequently reorganized around the spindle at the metaphase and increased

co-localization with mitochondria (**Figure 1E**). In interphase 2-cell stage embryos, the

F-actin meshwork was seen again, albeit with significantly reduced overlap with

mitochondria. To test whether the F-actin is involved in mitochondrial distribution and

partitioning, zygotes were treated with 1 μM latrunculin A for 1 h just after NEB,

sufficient for F-actin depolymerization (**Figure 1-figure supplement 1E**). Disruption of

F-actin organization resulted in asymmetric mitochondrial inheritance (**Figure 1-figure**

supplement 1F, 1G), but also increased cell size asymmetry (**Figure 1-figure**

supplement 1H), and these two asymmetries were correlated (**Figure 1-figure**

supplement 1I). Given many cellular roles, disruption of F-actin per se was unsuitable

as a strategy for manipulating mitochondrial distribution, we have therefore targeted

Myo19 as a strategy for altering the mitochondrial distribution and ask if it is critical for

the early development. Asymmetric mitochondrial inheritance has been previously

reported in cells lacking Myo19 (Majstrowicz et al., 2021; Moore et al., 2021; Rohn et

al., 2014). Myo19 was expressed in both interphase and metaphase zygotes, where its

localization was consistent with the mitochondrial distribution (**Figure 1F**). Acute

depletion of Myo19 at the PN stage by Trim-Away was confirmed by Western blot

analysis, which was not observed in control zygotes injected with Trim21 mRNA and

control IgG (**Figure 1G, 1H**). Unexpectedly, no significant asymmetry was observed in

the mitochondrial distribution and partitioning between the daughter blastomeres in

Myo19-depleted 2-cell embryos (**Figure 1I, 1J**). Live imaging of mitochondria

(mt-GFP) and chromosomes (H2B-mCherry) the symmetric distribution and

partitioning of mitochondria during the first embryonic cleavage (**Figure 1-figure**

supplement 2A, 2B; Figure 1-Video 2). Moreover, Myo19 depletion did not affect

developmental competence of the embryos to the blastocyst stage (**Figure 1K**). Overall,

fragmented mitochondria are closely associated with cytoplasmic F-actin during the

first cleavage, but active transport by the actin motor is not directly involved in

mitochondrial partitioning.

Loss of Drp1 induces mitochondrial aggregation and disturbs subcellular organelle

compartments

The highly fragmented mitochondrial morphology appears to facilitate

mitochondrial partitioning during embryonic cleavage. To understand the underlying mechanism of this process, we examined the role of Drp1, a key regulator of mitochondrial fission. Western blot analysis revealed that Drp1 was expressed in PN zygotes, and comparable protein levels were found in the morula stage, whereas their expression was remarkably decreased at the blastocyst stage (**Figure 2A, 2B**). To demonstrate the role of Drp1-mediated mitochondrial fission in preimplantation embryos, we employed Trim-Away. Co-injection of Trim21 mRNA and an antibody against Drp1 into zygotes 20–22 h post hCG reduced the Drp1 protein to nearly undetectable levels by 5 h post-injection at the PN stage, whereas co-injection of Trim21 mRNA and control IgG failed to trigger protein degradation (**Figure 2C, 2D**).

Mitochondria in live 2-cell embryos (37/37) expressing mt-GFP were interspersed throughout the cytoplasm. In contrast, almost all Drp1 Trim-Away embryos (55/57) exhibited marked mitochondrial aggregation that was reversed by ectopic expression of mCherry-tagged Drp1 (mCh-Drp1) (50/53) (**Figure 2E, 2F**). In Drp1-depleted embryos, mitochondria became highly aggregated into a small number of large clusters, whereas in controls and Drp1 Trim-Away embryos expressing mCh-Drp1, mitochondria remained dispersed in a large number of small clusters (**Figures 2G, 2H**). We observed similar mitochondrial aggregation occurred in parthenogenetic Drp1 KO embryos (**Figure 2-figure supplement 1A**), as the mitochondrial cluster size in $Drp1^{\Delta/\Delta}$ parthenotes was greater than $Drp1^{fl/fl}$ control, albeit the number of clusters were comparable (**Figure 2-figure supplement 1B, 1C**).

EM images of control 2-cell embryos showed that rounded mitochondria with less

developed cristae were dispersed throughout the cytoplasm, whereas swollen or

partially elongated mitochondria with lamella cristae structures in the inner membrane

were observed in Drp1-depleted embryos (**Figure 2I**). Both mitochondrial major and

minor axes in Drp1-depleted embryos was significantly greater than those of control

embryos (**Figure 2J-L**). Quantification of the aspect ratio (long axis/short axis)

suggests that mitochondria are significantly elongated in Drp1-depleted embryos

(**Figure 2M**).

To clarify the effects of these morphological changes in mitochondria on their

energy production, we compared ATP levels between control and Drp1 Trim-Away

embryos at the 2-cell stages. To analyze intracellular ATP concentrations, we expressed

a fluorescence resonance energy transfer (FRET)-based ATP biosensor, ATeam AT1.03,

with an emission ratio of AT1.03 fluorescence (YFP/CFP) used to estimate ATP levels.

ATP levels in Drp1-depleted embryos were indistinguishable from those in the control

embryos (**Figure 2N**). No FRET signal was detected in embryos expressing a mutant

version of the ATP probe AT1.03RK that cannot bind ATP.

Recent studies have focused on the interactions between organelles at membrane

contact sites (MCSs) and their roles in maintaining cellular homeostasis (English and

Voeltz, 2013). The ER is a continuous membrane-bound organelle with MCSs such as

with the plasma membrane, mitochondria, Golgi, and peroxisomes. Since mitochondrial

dynamics are spatially coordinated at the ER–mitochondria MCSs (Friedman et al.,

2011), we surmised that mitochondrial aggregation due to the loss of Drp1 would

compromise the organization of ER MCSs. To visualize the distribution of mitochondria

with other organelles in live embryos, fluorescently tagged ER (ER-mCherry), Golgi

(Golgi-mCherry), and peroxisomes (PEX-mCherry) were co-expressed with mt-GFP.

The ER was partially confined to the regions of the mitochondrial aggregation in Drp1

Trim-Away embryos, which appeared to disturb the endogenous ER network (**Figure**

2-figure supplement 2A). Linscan of mt-GFP and ER-mCherry signals at each focal

plane of large mitochondrial clumps showed that ER is confined to mitochondrial

aggregation. In contrast, the subcellular distribution of the Golgi was not clearly altered

by Drp1 depletion (**Figure 2-figure supplement 2B**). Drp1 is also known to regulate

peroxisomal fission, as elongated peroxisomes are seen in Drp1-deficient cells (Koch et

al., 2003); however, the subcellular distribution of peroxisomes in mammalian embryos

has not been reported. The peroxisomes in normal 2-cell embryos showed a punctate

distribution in the cytoplasm, whereas these puncta partially aggregated following Drp1

depletion by Trim-Away (**Figure 2-figure supplement 2C**), although it appeared to be

independent of mitochondrial aggregation in terms of spatial positioning.

Drp1-mediated mitochondrial fragmentation is required for the symmetric

mitochondrial partitioning into two functional daughter blastomeres

To decipher the spatiotemporal regulation of mitochondrial dynamics following

Drp1 depletion, we imaged mitochondria and chromosomes in Drp1 Trim-Away

zygotes expressing mt-GFP and histone H2B-mCherry (**Figure 3A; Figure 3-Video 1**).

No significant delay in cell cycle progression was observed following Drp1 depletion

(data not shown) compared to control zygotes (Figure 1A). Clumped mitochondria were

located in the center of the metaphase zygotes which resulted in an increase in

mitochondrial asymmetry at anaphase (**Figure 3B)**. We postulated that this asymmetry

was due to non-uniformity in the distribution of mitochondria around the spindle. As

expected, live imaging of mitochondria (mt-DsRed) and spindle microtubules

(EB3-GFP) revealed compared with the uniform angular positioning of around the

spindle observed in control zygotes, mitochondria in Drp1-depleted zygotes lost their

uniform distribution (**Figure 3-figure supplement 1A and 1B**). The biased inheritance

of mitochondria was further validated by the quantification of mtDNA copy number in

isolated blastomeres from 2-cell embryos (**Figure 3C**). The inheritance ratio of mtDNA

in daughter blastomeres of control embryos was comparable, whereas Drp1 depletion

led to biased mtDNA inheritance. Although Drp1 depletion did not affect intracellular

ATP levels in the whole embryo (**Figure 2H**), we tested if the biased inheritance of

mitochondria led to differences in blastomere ATP levels. Using AT1.03, ATP

concentration and mitochondrial distribution were simultaneously visualized in

blastomeres of 2-cell embryos (**Figure 3-figure supplement 1C**). Notably, the bias in

ATP levels between blastomeres of 2-cell embryos was greater following the Drp1

depletion (**Figure 3D**). We then quantified mtDNA copy number in each blastomere and

found a positive correlation between mtDNA inheritance and ATP levels in the

individual blastomeres (**Figure 3E**).

The presence of ER-mitochondria MCSs may lead to mitochondria-associated ER

asymmetry during cleavage. Live imaging of mitochondria and ER showed that part of

the ER is confined to mitochondrial aggregation, resulting in asymmetric distribution at

the telophase of the first cleavage (**Figure 3F; Figure 3-figure supplement 1D; Figure**

3-Video 2) which was closely correlated with mitochondrial asymmetry (**Figure 3G**).

No significant asymmetry was observed in the distribution of Golgi or peroxisomes,

suggesting that these organelles are not directly confined by mitochondrial location. The

ER is the main reservoir of intracellular Ca^{2+} stores in non-excitable cells.

ER-mitochondria MCSs create localized domains of high Ca^{2+} concentration required

for Ca^{2+} transfer from the ER to the mitochondria, in which the propagation of Ca^{2+}

signals in the mitochondria controls energy metabolism (Rizzuto et al., 2012). To

explore whether alterations in the ER and mitochondrial distribution and the resulting

segregation of the inter-organelle contacts affect intracellular Ca^{2+} homeostasis in 2-cell

embryos, the Ca^{2+} response in each blastomere after exposure to thapsigargin (Tg) in

Ca^{2+} free medium, a specific inhibitor of sarco/ER Ca^{2+}-ATPase, was compared

between control and Drp1 Trim-Away embryos (**Figure 3H, 3I**). Drp1 depletion

significantly reduced thapsigargin-induced Ca^{2+}-release and also caused significant

variability between blastomeres of individual 2-cell embryos which was not seen in

controls (**Figure 3J**). These findings indicate that Drp1 depletion disrupts

mitochondrial-ER interaction leading to compromised ER Ca^{2+} stores and that

asymmetric inheritance of ER leads to further variability in Ca^{2+} stores between

daughter blastomeres.

Drp1 is required for the developmental competence of preimplantation embryos

We found that Drp1-depleted zygotes were predominantly arrested at the 2-cell

stage and few embryos developed to the blastocyst stage (**Figure 4A, 4B**). This is a

specific effect of Drp1 deletion because none of the internal control conditions

increased arrest at the 2-cell stage and arrest was completely reversed by microinjecting

Trim-away insensitive exogenous mCh-Drp1 mRNA (**Figure 4C**). Further, western blot

analysis confirmed expression of mCh-Drp1, in conditions where endogenous Drp1 was

degraded by Trim-Away (**Figure 4D**).

More than half of the Drp1 Trim-Away embryos were arrested at the 2-cell stage,

the vast majority of which did not enter the following M-phase, as assessed by the

presence of the nuclear envelope (data not shown). Phosphorylated histone H3 (Ser10),

localized in the heterochromatin region in G2-phase cells (Hendzel et al., 1997), was

detected in 89% (16/18) of arrested embryos following Drp1 depletion (**Figure 4-figure**

supplement 1A), suggesting that the majority of interphase arrests occur in the

G2-phase. Since G2/M interphase arrest presumably activates a DNA damage

checkpoint, we estimated DNA lesions by immunofluorescence of γH2AX, a widely

used marker for DNA damage 46 h post hCG (corresponding to the G2-stage of the

2-cell stage). As expected, the number of γH2AX foci in Drp1-depleted embryos was

significantly greater than that in control embryos (**Figure 4-figure supplement 1B, 1C**).

The accumulation of reactive oxygen species (ROS), which are generated as byproducts

of mitochondrial energy production, increases susceptibility to DNA damage.

Intracellular ROS levels, as estimated by H2DCFDA fluorescence, were highly

accumulated in mitochondria (**Figure 4-figure supplement 1D, 1E**). This is a specific

effect of Drp1 deletion because none of the internal control conditions increased arrest

at the 2-cell stage and arrest was completely reversed by microinjecting Trim-away

insensitive exogenous mCh-Drp1 mRNA.

Misplaced mitochondria during the first cleavage division impair the assembly of

parental chromosomes leading to binuclear formation

We observed that approximately 30% (17/53) of Drp1 Trim-Away embryos

consisted of 3 or 4 blastomeres despite the first cleavage (indicated by arrowheads in

Fig. 4B), suggesting that cleavage defects have occurred. Live imaging experiments

showed that Drp1 Trim-Away embryos frequently display chromosome segregation

defects. As a typical example, two pronuclei frequently fail to reach apposition but

continue into mitosis and subsequently enter anaphase leading to binucleate blastomeres

in the subsequent 2-cell embryo (**Figure 5A; Figure 5-Video 1**). Immunostaining of

metaphase zygotes showed that 55% (17/31) of Drp1-depleted zygotes formed two

independent spindles with clustered microtubule organizing centers (**Figure 5B**). Both

confocal and EM images showed that large mitochondrial clumps occupied the central

region in Drp1-depleted zygotes, to which two spindles were separately located (**Figure**

5-figure supplement 1A, 1B). Tracking of two sets of chromosomes with mitochondria

following Drp1 depletion revealed that the mitochondrial aggregation caused by

Drp1-depletion sits between the pronuclei and subsequent spindles such that the

distance from chromosomes to the centre of the zygote is increased (**Figure 5C, 5D**).

The duration from mitotic entry to chromosome segregation at anaphase was not

affected (not shown) but binucleation was observed in 74% (32/43) of Drp1-depleted

embryos and in none of the control zygotes (n = 51) (**Figure 5E**). In addition to the

binucleation of blastomeres via dual spindle formation, other cell division abnormalities

were apparent. In approximately half (24/43) of the zygotes, multiple cleavage furrows

formed between the different sets of chromosomes leading to transient formation of

three or four blastomeres, of which 42% (10/24) failed to complete cytokinesis and

collapsed back to form binucleated 2-cell embryos (**Figure 5-figure supplement 1C,**

1D; Figure 5-Video 2). Together, these results suggest that proper control of

mitochondrial positioning during mitosis is crucial for assembly into a single spindle,

and marked mitochondrial centration causes chromosome segregation defects, leading

to binuclear formation.

Discussion

Mechanisms underlying symmetric mitochondrial inheritance in preimplantation

embryos

To ensure proper partitioning of intracellular organelles for two functional daughter

cells, cells undergo complex and coordinated remodeling of their cytoskeleton and

membranes during cell division through mechanisms that are less well understood than

chromosomal dynamics. As mitochondria are unique organelles with their own

genomes, their biased inheritance is expected to have detrimental effects on daughter

cell function. We showed that Drp1 mediates mitochondrial fragmentation and

dispersion, enabling symmetric partitioning between daughter blastomeres

stochastically, which is indispensable for healthy embryo development (**Figure 5F**).

Mitochondria that form a morphologically complex network and interact with

microtubules at the interphase are rapidly fragmented upon mitotic entry, detached from

the microtubules, and redistributed throughout the cytoplasm (Ball and Singer, 1982).

Mitochondrial fission is activated via the phosphorylation of Drp1 during mitosis. Two

mitotic kinases, CDK1 and Aurora A, activate Drp1, leading to enhanced mitochondrial

fission (Yamano and Youle, 2011). As the cell exits mitosis, Drp1 is sumoylated and

targeted for degradation to prevent mitochondrial fragmentation (Figueroa-Romero et

al., 2009). Although it remains to be investigated whether these cell cycle-dependent

modifications of Drp1 also occur in early embryos, mitochondria appear to be

constitutively fragmented throughout the cell cycle. This unique mitochondrial

morphology not only facilitates symmetric partitioning in cell division without cell

growth, but also appears to be well-adapted to the rapid proliferation of single-cell

zygotes into multicellular organisms before cell lineages specification. In

asymmetrically dividing cells, functionally distinct mitochondria are unequally

apportioned to influence daughter cell fate (Katajisto et al., 2015). In contrast,

symmetric cell division is intended to produce identical daughter cells with comparable

fates. In this context, the highly fragmented mitochondrial morphology in early

embryos may contribute to avoid cell fate divergence, as well as fulfill metabolic

homogeneity among blastomeres. Interestingly, the apparent morphological changes in

mitochondria occur at the blastocyst stage, as elongated mitochondria are formed in

trophectoderm (TE) cells, which is likely to be associated with altered energy

metabolism in the TE lineage (Kumar et al., 2018). Considering that the transformation

from eggs to embryos is a fundamental process in animal development involving

changes such as the transition from meiotic to mitotic spindles and maternal to zygotic

gene expression, changes in mitochondrial morphology and function may also play a

role in regulating cell fate and function in the early embryo.

The cytoskeleton is a critical regulator of organelle positioning. In contrast to a

uniformly distributed mitochondria throughout the cytoplasm during somatic cell

division (Chung et al., 2016; Moore et al., 2021), mitochondria progressively

accumulated around pronuclei and encircle the metaphase spindle of the first embryonic

cleavage. Since the two pronuclei migrate from the periphery to the centre of the eggs

by the F-actin-dependent mechanism to unite chromosomes on the first mitotic spindle

(Chaigne et al., 2016; Scheffler et al., 2021), actin filaments may drive the

mitochondrial movement. In fact, mitochondrial distribution was closely associated

with cytoplasmic F-actin organization in zygotes, especially at the metaphase. Although

depolymerization of F-actin prior to metaphase led to asymmetric mitochondrial

partitioning, the cell division itself was also asymmetric, probably due to failure of

centering of the pronuclei. Given that the depletion of Myo19 did not affect the

mitochondrial inheritance between daughter blastomeres, actin-based motor may not

contribute the symmetric partitioning of mitochondria in cleavage embryos. This result

is mostly consistent with previous observations in mitochondrial Rho GTPase (Miro)

1-deleted zygotes. Miro1 is localized in the mitochondrial outer membrane and the

adaptor protein TRAK mediates the attachment of motor proteins to the mitochondria

(Lopez-Domenech et al., 2018). Miro also regulates Myo19, ensuring stability to

actin-mediated mitochondrial distribution (Lopez-Domenech et al., 2018), and thus loss

of Miro leads to asymmetric segregation of mitochondria during mitosis most likely due

to reduced levels of Myo19. In Miro1-deleted zygotes, mitochondrial dynamics was

partially disturbed, but this did not affect mitochondrial partitioning to daughter

blastomeres, and fertility was normal as a result (Lee et al., 2022). Thus, fragmented

mitochondria may be primarily important for symmetric mitochondrial partitioning in

cleavage embryos, as a less precise stochastic partitioning mechanism may be suitable

or sufficient for equally partitioning of mitochondria, which are present in abundance

and are located across the cell volume in oocytes/zygotes. Nevertheless, it is also worth

considering the possibility that actin accumulated in mitochondria may be indirectly

involved in mitochondrial dynamics via mitochondrial fission. For example, inverted

formin 2 controls actin polymerization and has been reported to be required for efficient

mitochondrial fission as an upstream function of Drp1 (Korobova et al., 2013).

The physiological roles of mitochondrial fission in preimplantation embryos

Inhibition of Drp1-mediated mitochondrial fission causes severe embryonic arrest.

Given the housekeeping cellular functions of mitochondria, dysregulation of

mitochondrial dynamics may exert profound effects on the developmental competence

of embryos. Mitochondrial dynamics are a highly regulated process, and when disrupted,

can lead to cellular and tissue pathologies. The importance of mitochondrial fission has

been intensively studied in animal cells, and emerging evidence suggests that

Drp1-mediated fission participates in diverse cellular processes, including ER contacts,

Ca^{2+} signaling, autophagy, and apoptosis (Kraus et al., 2021). Drp1-KO mice are

embryonic lethal, most likely due to brain, heart, and placental defects; however,

various tissues appear normal, and cultured cell lines continue to proliferate even

though mitochondria are unevenly segregated (Ishihara et al., 2009; Wakabayashi et al.,

2009). Cells lacking Drp1 have highly elongated mitochondria that cannot be divided

into transportable units, and typically accumulate in a limited area, leaving large parts

of the cell devoid of mitochondria. This imbalanced distribution of mitochondria may

fail to meet local energy demands and thus would be more manifest in large or extended

cells, such as neurons and oocytes/zygotes. Moreover, our data demonstrated that Drp1

depletion in cleavage embryos caused biased mitochondrial inheritance and energy

heterogeneity between blastomeres. Given the low levels of de novo mtDNA replication

during preimplantation development, these biases may be further magnified by cleavage

division progression. It is also noteworthy that we showed that mitochondrial

aggregation induced by the loss of Drp1 disturbs the spatial organization of ER. The

aggregated ER subdomains likely caused leakage of Ca^{2+} stores, which may

compromise the tethering functions at the ER MCSs. The precise mechanisms of 2-cell

stage arrest arising from depletion of Drp1 remain elusive. It has been reported that

Drp1 depletion during meiosis lead to premature chromosome segregation due to

spindle assembly checkpoint defects (Zhou et al., 2022). Drp1-deficient cells has also

been reported to cause replication stress, leading to DNA damage response and the

G2/M cell cycle arrest (Qian et al., 2012), which is more consistent with our

observations. The presence of a Chk1-dependent checkpoint leading to G2 arrest in

2-cell mouse embryos (Ladstatter and Tachibana-Konwalski, 2016) suggests that this

surveillance mechanism may be involve in the 2-cell arrest process in this study.

In the present study, we found frequent binucleation in Drp1-depleted embryos.

Binucleation of early embryos is relatively common in human IVF procedures, but the

mechanisms of its occurrence and association with the developmental competence of

the embryo are not fully understood (Gomes Paim and FitzHarris, 2020). Reichmann et

al. recently demonstrated that mouse zygotes form two functionally independent

spindles before anaphase, and spatially separate parental genomes during the first

cleavage (Reichmann et al., 2018). Notably, failure to align the two spindles produced

errors, resulting in binuclear formation in the 2-cell embryos. The present study

demonstrates that clustered mitochondria in the central space of zygotes interfere with

the assembly of two spindles, implying a potential contribution of the organelle

misplacement to binucleation in human embryos.

510   Mitochondrial dynamics and quality control

The present study revealed the response to defective mitochondrial fission at the

cellular level in early embryos, but the response at the organelle level remains to be

elucidated. Mitochondria in oocytes, with hundreds of thousands of mtDNA copies, are

maternally inherited after fertilization. The preservation and apportionment of healthy

mtDNA in oocytes and early embryos is important for the inheritance of normal mtDNA

in the offspring. Recent full-length mtDNA sequencing has identified de novo mtDNA

mutations in oocytes, and their frequency increases with age, implying mtDNA turnover

during meiotic arrest (Abrisch et al., 2020). Oocytes with severe mtDNA mutations can be eliminated (Fan et al., 2008), although the thresholds at which mtDNA mutations and the resulting mitochondrial dysfunction result in selection remain unclear. The autophagic elimination of mitochondria and mitophagy may also function as filters to segregate and eliminate pathogenic mtDNA mutations at the organelle level (Youle and van der Bliek, 2012). An attractive model has been proposed for the Drosophila female germline, as fragmentation of the mitochondrial network facilitates mitophagy and allows selective elimination of defective mitochondria containing mutant mtDNA (Lieber et al., 2019). Functional links between mitochondrial fission and removal of damaged mitochondria by mitophagy have also been documented in mammalian cells, as Drp1 deficiency causes mitochondrial dysfunction owing to the failure of a Drp1-dependent mechanism of mitophagy that removes damaged mitochondria within the cell (Twig et al., 2008). The highly fragmented mitochondrial structure in mammalian oocytes/zygotes allows mtDNA to be sequestered in small units, which may be beneficial for eliminating mutation-impaired mitochondria for mitophagy. Furthermore, symmetric segregation of fragmented mitochondria might randomize mutant mtDNA among daughter blastomeres during embryonic cleavage to maintain tissue homeostasis. Elucidating the behavior of mutant mitochondria in early embryos is clinically meaningful. Mitochondrial replacement therapy (MRT) is a promising approach to prevent the transmission of mtDNA disorders from mother to child, as pathogenic mutant mtDNA in mature oocytes or zygotes is replaced with normal mtDNA by nuclear transfer (Hyslop et al., 2016; Kang et al., 2016). A potential problem

with MRT is that a small fraction of patients' mtDNA is inevitably carried over into the

reconstructed oocytes or zygotes (approximately 1 to 4%, respectively). It has been

noted that these low levels of mutant mtDNA may predominantly expand during

post-implantation embryogenesis, suggesting a rapid shift to one mtDNA haplotype

during early development (Lee et al., 2012). Therefore, it is important to investigate the

role of mitochondrial dynamics in the regulation of mtDNA mutations in the

preimplantation embryo, and whether mitochondrial fission contributes to reducing the

mutational load.

Materials and methods

Preparation of zygotes

Zygotes were collected from the oviducts of 8- to 12-week-old BDF1 female mice

mated with 3- to 6-month-old BDF1 male mice. Females were superovulated by

intraperitoneal injection of 7.5 IU of equine chorionic gonadotropin (PMSG; ASKA

Pharmaceutical, Tokyo, Japan) followed by injection of human chorionic gonadotropin

(hCG; ASKA Pharmaceutical) 48 h later. Female mice were euthanized 20-24 h later

and fertilized zygotes were retrieved in Hepes-buffered KSOM (H-KSOM) medium

under paraffin oil, at 37°C in a humidified atmosphere containing 5% CO_2. All mice

were housed in a pathogen-free environment in filter-top cages and fed a standard diet.

Animal experiments were conducted according to the guidelines of the Animal Care and

Use Committee of Okayama University.

For experiments in Fig. 3C, Drp1$^{loxP/loxP}$ mice (Wakabayashi et al., 2009) were

crossed with transgenic mice that carried Gdf-9 promoter–mediated Cre recombinase.

After multiple rounds of crossing, homozygous $Drp1^{cKO}$ female mice lacking Drp1 in

oocytes ($Drp1^{loxP/loxP}$; Gdf9-Cre) or $Drp1^{\Delta/\Delta}$ oocytes were obtained. Mice that do not

carry the Cre transgene are referred to as $Drp1^{loxP/loxP}$ and were used as controls. For

collection of ovulated MII oocytes, mice were superovulated by sequential

intraperitoneal injections of 8-IU PMSG and 8-IU hCG at an interval of 48 h, and the

mice were culled 14 to 16 h after hCG injection to collect oocytes from oviducts.

Animal experimentation was approved by Monash University Animal Experimentation

Ethics Committee and was performed in accordance with Australian National Health

and Medical Research Council Guidelines on Ethics in Animal Experimentation.

Plasmids

EGFP, mCherry, and DsRed were cloned into pcDNA6 vector between the XhoI and

XbaI sites (pcDNA6/Myc-His B; Invitrogen, Waltham, MA) with a poly(A)-tail of 84

nucleotides (Wakai et al., 2014). The gene sequence encoding the mitochondrial

targeting sequence of Cox VIII was amplified by PCR and ligated to the EGFP and

DsRed-bearing pcDNA6 (mt-GFP and mt-DsRed). The gene sequences encoding the

ER-targeting sequence of calreticulin and the KDEL ER retention sequence were cloned

into mCherry (ER-mCherry)-bearing pcDNA6. For visualization of DNA and

microtubule growth, gene sequences encoding histones H2B and EB3 were ligated to

the pcDNA6 vector in frame with mCherry (H2B-mCherry) and EGFP (EB3-GFP),

respectively. The gene sequences encoding the amino-terminal 33 residues of

endothelial nitric-oxide synthase as a Golgi targeting sequence (Addgene, 14873) and

peroximal targeting signal 1 (Addgene, 54520) were cloned into mCherry-bearing

pcDNA6 (Golgi-mCherry and PEX-mCherry). The gene sequences encoding

N-terminally mCherry tagged Drp1 (Addgene, 49152) were inserted between the EcoRI

and PstI sites of the pCDNA vector (mCherry-Drp1). ATP biosensor, AT1.03 and

AT1.03RK in the pcDNA3.1 vector were kindly provided by Dr H. Imamura (Kyoto

University). Capped mRNA was synthesized with T7 polymerase (RiboMAX

Large-Scale RNA Production, Promega, Madison, WI) according to manufacturer

instructions. mRNAs were delivered into zygotes using a piezo-driven micropipette unit

(Prime Tech, Tsuchiura, Japan). For microinjection, mRNA solution was loaded into

glass micropipettes at the concentration of 200 ng/µl (mt-GFP, EB3-GFP, mt-DsRed,

H2B-mCherry, ER-mCherry, Golgi-mCherry, and PEX-mCherry), 500 ng/µl

(mCheery-Drp1) or 1000 ng/µl (AT1.03 and AT1.03RK). The volumes injected

typically ranged from 2 to 10 pl, which is 1–5% of the total volume of the zygotes.

Manipulation was carried out in H-KSOM medium containing 5 mg/mL cytochalasin B

(Sigma, St. Louis, MO) to increase post-microinjection survival.

Trim-Away

Trim-Away was performed as previously described with some modifications. In

brief, pCMV6-Entry vector harboring Trim-21 mouse ORF Clone (Origene, Rockville,

MD) was linearized by FseI, and mRNA was synthesized using T7 mMessage

mMachine and poly-A tailed with Poly(A) tailing kit (Thermo Fisher Scientific,

Waltham, MA). Mouse monoclonal anti-Drp1 antibody (BD Biosciences, Franklin

Lakes, NJ) or rabbit monoclonal anti-Myo19 antibody (Abcam, Cambridge, UK) was

concentrated using Amicon Ultra-0.5 100 KDa centrifugal filter devices (Millipore,

Bedford, MA) and the buffer was replaced with PBS containing 0.03% NP40 (Nakarai,

Kyoto, Japan). Pronuclear stage zygotes were injected with Trim21 mRNA (1000

ng/mL) and subsequently anti-Drp1 or anti-Myo19 antibodies. For the control

experiment, normal mouse immunoglobulin G (IgG) antibody (Santa Cruz

Biotechnology, Dallas, TX) and rabbit IgG (Cell Signaling Technology, Danvers, MA)

was used instead of an anti-Drp1 antibody and anti-Myo19 antibodies, respectively.

Western blotting

Whole cell lysates from zygotes/embryos were prepared by adding a 2× sample

buffer. Proteins were separated by SDS-PAGE and transferred to PVDF membranes

(Millipore). The membranes were then blocked and probed with anti-Drp1 (1:1000) or

anti-Myo19 (1:1000) antibodies for 1 h at room temperature. Goat anti-mouse antibody

conjugated to horseradish peroxidase (HRP) was used as a secondary antibody (1:2000)

for chemiluminescence detection (Clarity Western ECL Substrates, Bio-Rad, Hercules,

CA) according to the manufacturer's instructions. The signal was digitally captured

using a ChemiDoc XRS+ imaging system (Bio-Rad). The same membranes were

stripped at 50°C for 30 min (62.5 mM Tris, 2% SDS, and 100 mM 2-beta

mercaptoethanol) and re-probed with anti-β-actin monoclonal antibody (1:2000, Cell

Signaling Technology) as the internal control.

mtDNA copy number

The zona pellucida of 2-cell embryos was removed with H-KSOM medium

containing 0.1% protease (Sigma, St. Louis, MO). Individual blastomeres from the

zona-free embryos were obtained by repeatedly pipetting in Ca^{2+}-free and Ma^{2+}-free

H-KOSM medium. After washing in PBS, each blastomere was transferred into 5 µL of

lysis buffer (50 mM Tris-HCl, pH 8.5, with 0.5% Tween 20 and 100 µg proteinase K, at

55°C, followed by heat inactivation at 95°C for 10 min. The plasmid standard was

constructed by cloning a 194-bp fragment of the NADH-ubiquinone oxidoreductase

chain 1. The standard stock was serially diluted to prepare the standard curve (10^3 to

10^7). Real-time fluorescence-monitored quantitative PCR using the primer pair

5′-CCTATC ACCCTTGCCA-3′ and 5′-GAGGCTGTTGCTTGTG-3′ was performed on

a LightCycler 96 System (Roche, Basel, Switzerland). Each reaction of 20 µL consisted

of 10 µL of SYBR Green I (Roche), 0.5 µM each of the forward and reverse primers,

and 2 µL of DNA template. The cycling conditions were as follows: initial denaturation

at 95°C for 5 min followed by 45 cycles of denaturation at 95°C for 10 sec, annealing at

50°C for 20 sec, and elongation at 72°C for 8 sec. Standard curves were created for

each run, and the sample copy number was determined from the equation relating the Ct

value against copy number for the corresponding standard curve.

Immunofluorescence

Zygotes/embryos were fixed and permeabilized with 2% paraformaldehyde in

phosphate-buffered saline (PBS) containing 0.1% Triton X-100 for 40 min at room

temperature. After washing with PBS supplemented with 1% bovine serum albumin

(PBS–BSA), the zygotes/embryos were incubated overnight at 4°C with mouse

anti-α-tubulin (1:300, Sigma), rabbit monoclonal anti-Myo19 antibody (1:300, Abcam),

rabbit anti-pericentrin (1:500, Abcam) or mouse monoclonal Anti-γH2A.X (1:500,

Abcam) antibodies in PBS–BSA. Then zygotes/embryos were washed with PBS–BSA

and incubated with Alexa Fluor488-conjugated goat anti-rabbit IgG (1:300) for 1 h at

room temperature. DNA was stained with DAPI. Samples were placed in PBS under

mineral oil in glass bottom dishes (Matsunami, Osaka, Japan) and examined using a

laser-scanning confocal microscope (FV1200, Olympus, Tokyo, Japan) outfitted with a

63×1.4 NA oil immersion objective lens.

Mitochondrial and ROS staining

Mitochondrial distribution was analyzed by staining with 1 µM MitoTracker Red

CM-H2XRos (Thermo Fisher Scientific) in H-KSOM medium for 30 min at 37°C.

Intracellular ROS levels were analyzed using 5 µM H2DCFDA (Thermo Fisher

Scientific) in H-KSOM medium for 15 min at 37°C. The embryos were placed in drops

of H-KSOM medium under mineral oil in glass bottom dishes (Matsunami), and

fluorescence images of MitoTracker Red CMXros and H2DCFDA were obtained using

a laser-scanning confocal microscope (FV1200) fitted with a 63×1.4 NA

oil-immersion objective lens.

Electron microscopy

Zygotes/embryos were fixed in 2% glutaraldehyde in a 0.1 M phosphate buffer (pH

7.4) for 1 h and stored at 4°C until processed. After post-fixation in 2% osmium

tetroxide at 4°C, the specimens were dehydrated with a graded series of ethanol and

were embedded in epoxy resin Quetol-812. Semi-thin sections were cut for light

microscopy and stained with toluidine blue for further sectioning of areas. Thereafter,

ultra-thin sections were cut and stained with uranyl acetate and lead citrate and

examined by transmission electron microscopy (TEM) (JEM 1200EX, JEOL) at 80 kV.

Fluorescence resonance energy transfer and Ca^{2+} imaging

ATeam, a fluorescence resonance energy transfer-based ATP indicator, has been

used successfully to measure cellular ATP levels in live somatic cells (Imamura et al.,

2009). To estimate the relative changes in ATP levels, the emission ratio of AT1.03 and

AT1.03RK (YFP/CFP) was imaged using a CFP excitation filter, dichroic beam splitter,

and CFP and YFP emission filters (Chroma Technology, Rockingham, VT; ET436/20x,

89007bs, ET480/40m and ET535/30m). To measure cytoplasmic Ca^{2+}, embryos were

incubated with 1.25 μM Fluo-4 (Thermo Fisher Scientific) supplemented with 0.02%

pluronic acid (Thermo Fisher Scientific) for 20 min at room temperature. Fluo-4 was

excited with 480 nm wavelengths every 20 seconds and emitted light was collected at

wavelengths greater than 510 nm. To estimate Ca^{2+} levels in the ER, Ca^{2+} rise was

analyzed following addition of 10 μM thapsigargin (Sigma) in a Ca^{2+}-free medium. For

the monitoring of ATP and Ca^{2+} levels, embryos were attached to glass-bottomed dishes

(Matsunami) and placed on the stage of an inverted microscope. . The fluorescence

images were obtained by a scientific CMOS (sCMOS) camera (Hamamatsu Photonics,

Hamamatsu), and the rotation of excitation and emission filter wheels was controlled

using the MAC5000 filter wheel/shutter control box (Ludl Electronic Products Ltd,

Hawthorne, NY) and HCImage software.

Live cell confocal imaging

Spinning disk images of mitochondria (mt-GFP and mt-DsRed), chromosomes

(H2B-mCherry), microtubules (EB3-GFP), ER (ER-mCherry), Golgi (Golgi-mCherry)

and peroxisome (PEX-mCherry) and were obtained using a confocal scanner unit

(CellVoyager CV1000, Yokogawa, Tokyo, Japan) equipped with a C-Apochromat ×40,

1.2 NA water immersion objective. Images were typically acquired every 10-20 mins on

10 planes covering a 35-40 μm range (every 2.5-3.3 μm). Laser wavelengths of 488 and

561 nm were used for the excitation of EGFP and mCherry (DsRed), respectively (not

exceeding 0.2% laser intensity to minimize light-induced phototoxic stress).

Image analysis

Fiji (Schindelin et al., 2012) was used for quantification of fluorescence images.

For quantification of mitochondria at nuclear or spindle periphery in Figure 1B and

Figure 1-figure supplement 1C, the fluorescence images of nucleus (H2B-mCherry) of

spindle (EB3-GFP) were applied with a Gaussian filter (radius: 2 pixels) to remove

noise and segmented by intensity thresholding. The masked binary images were dilated

with 10 pixels width and the nucleus/spindle periphery was segmented by the

subtraction of the original image from the dilated image. The fluorescence intensity

ratio of mitochondrial layers at the periphery to that of outside periphery was measured

to estimate mitochondrial accumulation around the nucleus/spindle.

For quantitation of mitochondrial symmetry at the anaphase in Figure 1-figure

supplement 2B and Figure 3B, the cleavage plane of dividing blastomere was manually

outlined with respect to the middle of the anaphase chromosomes. Total mt-GFP

intensity in each blastomere was measured, and the greater value was divided by the

smaller value for the symmetry index. To analyze organelle symmetry at the telophase

in Figure 3F, total pixel intensity in the channel corresponding to mt-GFP, ER-mCherry,

Golgi-mCherry and PEX-mCherry in each blastomere was measured. The greater value

was divided by the smaller value; a ratio of 1 indicates symmetry, whereas higher

values indicate asymmetry.

To measure the size and count the number of mitochondrial clusters in Figure 2G,

2H and Figure 2- figure supplement 2B, 2C, the fluorescence images of mitochondria

(mt-GFP) were analyzed using a basic measurement function (analyze particles, size)

after adjusting intensity threshold (Yen algorism). Based on the average size of

mitochondria in EM images, mt-GFP signals larger than 0.15 μm^2 were counted as

mitochondrial clusters.

Uniformity analysis of mitochondrial distribution around spindles in y in Figure

3-figure supplement 1B was performed based on the angular distributions of

mitochondria (mt-DsRed) and microtubules (EB3-GFP) using "QuantEv Icy" plugin

(Pecot et al., 2018). The fluorescence images were applied with Gaussian filter (radius:

2 pixels) to reduce noise, and xy coordinates were converted to polar coordinates. The

fluorescence intensity profiles for mt-DsRed relative to the centroid of the EB3-GFP

signals was divided into sectors each 60 degree based on the long axis of the spindle.

The standard deviation of averaged mitochondrial intensity per 60° sector were

calculated, higher values were considered to indicate increased non-uniformity.

For quantitation of ROS enrichment in mitochondria in Figure 4-figure supplement

1E, the fluorescence images of mitochondria (MitoTracker Red CMXros) were

background-subtracted (rolling ball radius: 5 pixels) and applied with the Gaussian filter

(radius: 2 pixels) to remove noise. The inside mitochondria was masked by intensity

thresholding (Moments algorism). The outside mitochondria was segmented by

subtracting the masked image from the whole embryo area. The intensity ratio of

H2DCFDA fluorescence inside mitochondria to outside mitochondria was measured to

estimate ROS accumulation in mitochondria.

For tracking nuclei in zygotes in Figure 5C and 5D, XY coordinates of the centroid

of male and female chromosomes (H2B-mcherry) relative to the center of the zygote

were obtained using "Manual tracking" plugin. Only the zygotes with nuclei that

remained in the focal plane were analyzed.

Statistical analysis

All statistical analyses were performed using GraphPad Prism software. Values from

three or more experiments performed on different batches of cells were analyzed by

Statistical tests and p-values are indicated in figures and figure legends.

Resource availability

All unique/stable reagents and plasmids generated in this study are available from

the lead contact with a completed Materials Transfer Agreement. Further information

and requests for resources and reagents should be directed to and will be fulfilled by the

lead contact, Takuya Wakai (t2wakai@okayama-u.ac.jp).

Acknowledgements

We thank Dr. H. Imamura (Kyoto University) for sharing AT1.03 and AT1.03RK

plasmids. We are thankful to Dr. Hiromi Sesaki (Johns Hopkins University School of

Medicine) for sharing $Drp1^{fl/fl}$ mice line. We thank Ms. Sayaka Nakato and the staff of

Mio Fertility Clinic for technical assistance with CellVoyager CV1000. This work has

been partially supported by the Core-Facility Portal (CFPOU) at Okayama University

(DGP_13).

Figure Legends

Figure 1. Redistribution and symmetric partitioning of mitochondria during

embryonic cleavage.

(A) Representative time-lapse images (maximum-intensity Z projection) of

mitochondria and chromosomes in zygotes expressing mitochondrially-targeted

(mt-GFP) and histone H2B-mCherry during the first cleavage division. Time is relative

to the onset of NEBD. Scale bar, 20 μm.

(B) Quantification of mitochondrial accumulation around the nucleus at different cell

cycle phases by calculating the ratio of the mitochondrial fluorescence intensity at the

nuclear periphery to that of outside nuclear periphery.

(C) Representative electron microscopy (EM) images (n = 5) showing the accumulation

of mitochondria around the mitotic spindle. Higher-magnification images (31 sections)

of the representative boxed areas in the lower left and upper right, respectively.

Mitochondria (purple), endoplasmic reticulum (aqua), and actin filaments

(yellow-green) are pseudo-colored in the lower right panel. Scale bar left: 10 μm;

Lower left and upper right: 1 μm.

(D) Interphase or metaphase zygotes and interphase 2-cell embryos were stained with

Phalloidin-iFluor 488, MitoTracker Red CMXros and Hoechest 33342. Magnified

images in the box areas are shown in the right panels. An overall view of each embryo

is shown in Figure 1-figure supplement 1D.

(E) Quantification of the correlation between the localization of the F-actin

(Phalloidin-iFluor 488) and mitochondria (MitoTracker Red CMXros).

(F) Representative immunofluorescence images of Myo19. Interphase (n = 13) and metaphase (n = 11) zygotes were stained with MitoTracker Red CMXRos. After fixation and immunostaining with anti-Myo19 antibody and Hoechest 33342, single-section images crossing the mid-zone of the zygotes were imaged by confocal microscopy. Scale bar, 20 μm.

(G) Western blot analysis of zygotes overexpressing Trim21 and microinjected with control IgG or anti-Myo19 antibodies. Lysates of 100 zygotes were extracted 5 h after microinjection and were probed with antibodies specific to Myo19 and β-actin.

(H) Quantification of the relative Myo19 expression levels in (G) across 3 experimental replications following Myo19 depletion.

(J) Quantitation of mitochondrial mass inheritance after the first cleavage division of control and Myo19-depleted zygotes. Total mitochondrial fluorescence in each daughter blastomere of 2-cell embryos was measured, and the smaller value was divided by the greater value for the inheritance ratio.

(K) Developmental competence of control and Myo19-depleted embryos. Percentage of zygotes reaching the indicated developmental stage.

(I) Maximum-intensity Z projection of control and Myo19 Trim-Away embryos expressing mt-GFP and H2B-mCherry. Scale bar, 20 μm.

Data are represented as mean ± SD and P values calculated using two-tailed paired (B) or unpaired (E, H, K and J) Student's t test. $*p < 0.05$, $***p < 0.001$, $****p < 0.0001$; ns, not significant. Number of embryos are indicated in brackets.

Figure 1-figure supplement 1. Redistribution of mitochondria during the first

embryonic cleavage.

(A) Representative immunofluorescence images of microtubules (green), mitochondria

(red) and DNA (blue) during the first cleavage division. Metaphase II eggs were

collected 14 h after hCG. Zygotes were isolated 24 h after hCG and cultured to different

mitotic stages (n = 10–35). Interphase zygotes at the PN stage and 2-cell stage embryos

were fixed 25 and 35 h after hCG, respectively. Mitotic zygotes were fixed at 0.5

(prophase and prometaphase), 1 (metaphase), and 2 h (anaphase and telophase) after the

disappearance of the nuclear envelope was observed. Prior to fixation, zygotes were

stained with MitoTracker Red CMXRos for 0.5 h. After fixation and immunostaining

with anti-α-tubulin and DAPI, single-section images crossing the mid-zone of the

zygotes were imaged by confocal microscopy. Scale bar, 20 μm.

(B) Representative images (maximum-intensity Z projection) of mitochondria

(mt-DsRed) and microtubules (EB3-GFP) in metaphase blastomeres during the first to

third cleavage division. Scale bar, 20 μm.

(C) Quantification of mitochondrial accumulation around the spindle by calculating the

ratio of the mitochondrial fluorescence intensity at the spindle periphery to that of

outside spindle periphery.

(D) Interphase or metaphase zygotes and interphase 2-cell embryos were stained with

Phalloidin-iFluor 488, MitoTracker Red CMXros and Hoechest 33342. Magnified

images in the box areas are shown in Figure 1D.

(E) Confocal images of zygotes treated with DMSO (control) or Latrunculin A (LatA)

for 1 h just after NEB, fixed and stained for F-actin (Phalloidin-iFluor 488) and DNA

(Hoechest 33342). Magnified views of the cytoplasmic and cortical regions are shown

in the upper right and lower right, respectively. Scale bar, 10 μm.

(F) Zygotes were treated as in (E), then washed out, cultured to develop into 2-cell

embryos and stained with MitoTracker Red CMXros. Maximum-intensity Z projection

of the entire cell for mitochondria were imaged by confocal microscopy.

(G and H) Quantitation of mitochondrial mass inheritance (G) and blastomere size (H)

in control and LatA-treated 2-cell embryos. Total mitochondrial fluorescence and the

area in each daughter blastomere of two-cell embryos was measured, and the ratio was

calculated by dividing the smaller value by the larger value.

(I) Correlation between mitochondrial inheritance ratio and blastomere size ratio.

Data are represented as mean ± SD and P values calculated using two-tailed Student's t

test. $*p < 0.05$, $***p < 0.001$, $****p < 0.0001$; ns, not significant. Number of embryos

are indicated in brackets.

Figure 1-figure supplement 2. Redistribution of mitochondria during the first

cleavage in Myo10-depleted zygotes.

(A) Representative time-lapse images (maximum-intensity Z projection) of

mitochondria and chromosomes in anti-Myo19 microinjected Trim-Away zygotes

expressing mt-GFP and H2B-mCherry. Time is relative to the onset of NEBD. Dashed

lines indicate cleavage plane. Scale bar, 20 μm.

(B) Quantitation of mitochondrial asymmetry at anaphase in control and

Myo19-depleted zygotes. Total mitochondrial fluorescence in each dividing blastomere

was measured, and the greater value was divided by the smaller value for the symmetry

index.

Figure 1-Video 1. Symmetric mitochondrial partitioning during the first cleavage.

Representative time-lapse movies of mitochondria (mt-GFP) and chromosomes

(H2B-mCherry) in zygotes. Time is relative to the onset of NEB (hours_minutes).

Z-projection of 8 sections, 5 μm apart; time interval: 20 min.

Figure 1-Video 2. Mitochondrial dynamics during the first embryonic cleavage in

Myo19-depleted zygotes. Representative time-lapse movies of mitochondria (mt-GFP)

and chromosomes (H2B-mCherry) in zygotes. Time is relative to the onset of NEB

(hours_minutes). Z-projection of 8 sections, 5 μm apart; time interval: 20 min.

Figure 2. Depletion of Drp1 induces mitochondrial aggregation.

(A) Western blot analysis of preimplantation embryos at the different stages. Lysates of

30 cells were probed with antibodies specific to Drp1 and β-actin.

(B) Quantification of the relative Drp1 expression levels in (A) across 3 experimental

replications.

(C) Western blot analysis of zygotes overexpressing Trim21 and microinjected with

control IgG or anti-Drp1 antibodies (henceforth referred to simply as control and

Trim-Away, respectively), or zygotes microinjected with Trim21 mRNA, control IgG or

anti-Drp1 alone. Lysates were extracted 5 h after microinjection and immunoblotted for

the indicated proteins.

(D) Quantification of the relative Drp1 expression levels in (C) across 3 experimental

replications following Drp1 depletion.

(E) Representative images of in control and Drp1 Trim-Away embryos expressing

mt-GFP. Scale bar, 20 μm.

(F) A representative image of mitochondria (mt-GFP) in Drp1-depleted embryos

expressing exogenous Drp1 (mCh-Drp1). Note that mCh-Drp1 expression rescues the

mitochondrial aggregation following Drp1 depletion by Trim-Away to the normal

dispersed distribution. Scale bar, 20 μm.

(G and H) Comparisons of size (G) and number (H) of mitochondrial clusters (mt-GFP)

in (E) and (F).

(I) Representative EM images of control (n = 5, 17 sections) and Drp1-depleted (n = 6,

20 sections) 2-cell embryos. Right panels show higher-magnification images. Scale bar

left: 4 μm; right: 1 μm.

(J-M) Quantification of major and minor axes (J) of individual mitochondria observed

in (I). Comparisons of major axis (K), minor axis (L) and the aspect ratio (M) of

mitochondria in control and Drp1-depleted embryos.

(N) Levels of ATP in control and Drp1-depleted embryos were estimated using the

emission ratio of AT1.03 (YFP/CFP). The emission ratio of AT1.03RK (YFP/CFP),

which is unable to bind ATP, is shown as a negative control.

Data are represented as mean ± SD and P values calculated using two-tailed Student's t

test with Welch's correction. ****p < 0.0001. Number of embryos are indicated in

brackets.

Figure 2-figure supplement 1. Mitochondrial aggregation in KO pathenotes

913   (A) Representative images of mitochondria (mt-GFP) in $Drp1^{fl/fl}$ control (n =10) and

914    $Drp1^{\Delta/\Delta}$ (n = 8) parthenotes. Mature eggs were collected form 4- to 5-week-old $Drp1^{fl/fl}$

915   and Drp1^{cKO} mice. Scale bar, 20 μm. After microinjection of mt-GFP mRNA, oocytes

916   were exposed to 10 mM SrCl$_2$.

917   (B and C) Comparisons of size (B) and number (C) of mitochondrial clusters in (A).

Figure 2-figure supplement 2. Mitochondrial aggregation disturbs subcellular

organelle compartments.

(A-C) Sub-cellular distributions of mitochondria with ER (A), Golgi (B), and

peroxisome (C). Representative images of control and Drp1 Trim-Away embryos

co-expressing mt-GFP and ER-mCherry (control, n = 11; Trim-Away, n = 15),

Golgi-mCherry (control, n = 10; Trim-Away, n = 16) or PEX-mCherry (control, n = 19;

Trim-Away, n = 20). Magnified images in the box areas are shown in the right panels.

Scale bar, 20 μm. Bottom panels represent line scan analysis of ER-mCherry (A),

Golgi-mCherry (B) and PEX-mCherry at different focal planes of aggregated

mitochondria (mt-GFP) in Trim-Away embryos. The fluorescence intensity traces at the

dashed line across mitochondrial mass in focal images acquired at 5 μm intervals in the

Z-axis direction. Note that the intensity peak of ER-mCherry coincides with that of

mt-GFP in each focal plane.

Figure 3. Drp1 depletion causes asymmetric mitochondrial inheritance and

functions.

(A) Representative time-lapse images (maximum-intensity Z projection) of

mitochondria and chromosomes in anti-Drp1 microinjected Trim-Away zygotes

expressing mt-GFP and H2B-mCherry. Time is relative to the onset of NEBD. Dashed

lines indicate cleavage plane. Scale bar, 20 μm.

(B) Quantitation of mitochondrial asymmetry at anaphase in control and Drp1-depleted

zygotes. Total mitochondrial fluorescence in each dividing blastomere was measured,

and the greater value was divided by the smaller value for the symmetry index.

(C) mtDNA inheritance ratio after the first cleavage division of control and

Drp1-depleted zygotes. mtDNA copy number in each daughter blastomere of 2-cell

embryo was measured, and the ratio was calculated by dividing the greater value by the

smaller value.

(D) ATP inheritance ratio between two blastomeres was estimated from comparison of

emission ratio of AT1.03 (YFP/CFP) in the daughter blastomere A/B, where A is the

daughter blastomere with more mitochondria as confirmed by MitoTracker staining (see

Figure S3D).

(E) Correlation between mtDNA and ATP inheritance ratio between 2-cell blastomeres.

(F) Quantitation of organelle asymmetry at telophase was analyzed from time-lapse

images of embryos co-expressing mt-GFP with mCherry-tagged ER, Glogi, or

peroxisome. Total mitochondrial fluorescence in each blastomere was measured, and

the greater value was divided by the smaller value.

(G) Organelle symmetry ratios, as shown in (F), are plotted against the mitochondrial

symmetry ratio for each Drp1-depleted embryo. Note that mitochondrial and ER

inheritance ratios are highly correlated ($R^2 = 0.6734$).

(H-J) Endoplasmic reticulum Ca^{2+} content in each blastomere of control (H) and

Drp1-depleted (I) embryos was estimated following addition of 10 μM thapsigargin in

Ca^{2+} -free medium, and comparisons of the Fluo-4 fluorescent peaks are shown in a

graph (J).

Data are represented as mean ± SD and P values calculated using unpaired two-tailed t

tests with Welch's correction (B, C, D and F) or one-way ANOVA with Turkey's

multiple comparison test (J). *$p < 0.05$, ***$p < 0.001$, ****$p < 0.0001$; ns, not

significant. Number of embryos are indicated in brackets.

Figure 3-figure supplement 1. Drp1 depletion increases asymmetry in

mitochondrial inheritance between blastomeres.

(A) Representative time-lapse images (maximum-intensity Z projection) of

mitochondria and microtubules in control and anti-Drp1 microinjected Trim-Away

zygotes expressing EB3-GFP and mt-DsRed. Time is relative to the onset of anaphase.

Scale bar, 20 μm.

(B) Uniformity analysis of mitochondrial distribution around the spindle.

Representative angler distributions of mitochondria (red trace) and microtubules (green

trace) at the metaphase of control and Drp1 Trim-Away zygotes, as shown in (A). Based

on the angular distributions, standard deviation of averaged mitochondrial fluorescence

intensity per 60° sector were plotted in the graph to the right, higher values indicate

increased non-uniformity.

(C) Representative ratiometric images of ATeam AT1.03 and AT1.03RK fluorescence

(YFP/CFP) in control and Drp1-depleted 2-cell embryos. Mitochondrial distribution in

the same embryos is shown in the right panel. Arrowheads represent marked

mitochondrial aggregation in Drp1 Trim-Away embryos and blastomere A/B, where A is

the daughter blastomere with more mitochondria as indicated by MitoTracker staining.

(D) Representative time-lapse images (maximum-intensity Z projection) of

mitochondria and ER in control and Drp1 Trim-Away zygotes expressing mt-GFP and

ER-mCherry. Arrowheads represent regions of marked mitochondrial aggregation in

Drp1 Trim-Away embryos. Time is relative to the onset of NEBD. Scale bar, 20 μm.

Figure 3-Video 1. Asymmetric mitochondrial partitioning during the first cleavage of

Drp1-depleted zygotes, related to Fig. 2. Representative time-lapse movies of

mitochondria (mt-GFP) and chromosomes (H2B-mCherry) in Drp1 Trim-Away (right)

zygotes. Time is relative to the onset of NEB (hours_minutes). Z-projection of 8

sections, 5 μm apart; time interval: 20 min.

Figure 3-Video 2. Asymmetric partitioning of the ER confined to mitochondrial

aggregation regions in Drp1-depleted zygotes, related to Fig. 4 and Fig. S4.

Representative time-lapse movies of mitochondria (mt-GFP) and ER (ER-mCherry) in

control and Drp1 Trim-Away zygotes. Each of the movies in the upper row is displayed

in the lower row by LUT. Arrowheads represent regions of marked mitochondrial

aggregation. Time is relative to the onset of NEB (hours_minutes). Z-projection of 8

sections, 5 μm apart; time interval: 15 min.

1002

Figure 4. Drp1 depletion causes early embryonic arrest.

(A) Developmental competence of control and Drp1 Trim-Away zygotes. Following

Drp1 depletion by Trim-Away (co-injection of Trim21 mRNA and anti-Drp1 antibody)

or the control experiments in Fig. 2B, zygotes were cultured to the indicated stages.

(B) Representative images of embryos microinjected Trim21 mRNA with control IgG

or anti-Drp1 were cultured. Some embryos at E1.5 have three or four blastomeres

(arrowheads). Scale bar, 20 μm.

(C) Developmental competence of Drp1-depleted embryos is rescued by microinjection

of mCherry-Drp1 (mCh-Drp1) mRNA. Percentage of zygotes reaching the indicated

developmental stage. Note that the 2-cell stage arrest in control condition with RNA

buffer (water) upon Trim-Away depletion was reversed by microinjecting mCh-Drp1

mRNA.

(D) Western blot analysis of control and Drp1 Trim-Away zygotes that were

microinjected with water or mCh-Drp1 mRNA upon Trim-Away depletion. Lysates of

20 zygotes were extracted 5 h after microinjection and were probed with antibodies

specific to Drp1 and β-actin.

Data are represented as mean ± SD and P values calculated using one-way ANOVA

with Turkey's multiple comparison test. $*p < 0.05$, $***p < 0.001$, $****p < 0.0001$; ns,

not significant. Number of embryos are indicated in brackets.

1022

Figure 4-figure supplement 1. Drp1 depletion increases ROS in mitochondria and

DNA damage.

(A) Representative immunofluorescence images of phosphorylated (Ser10) histone H3

(green), known to localize in the heterochromatin region in G2-phase cells, and DAPI

(blue) in control (n = 18) and Drp1 Trim-Away (n = 20) embryos. Scale bars, 20 µm.

Note that the phosphorylated (Ser10) histone H3 signal accumulate in DAPI-intense

stained heterochromatin domains. Embryos were fixed at 46 h post hCG (corresponding

to the G2 stage of 2-cell embryos).

(B) Representative immunofluorescence images (maximum-intensity Z projection) of

γH2AX (green), MitoTracker Red CMXros (red) and DAPI (blue) in control and Drp1

Trim-Away embryos.

(C) Comparisons of the number of γH2AX foci in control and Drp1 Trim-Away

embryos.

(D) Representative images of H2DCFDA, a probe for intracellular reactive oxygen

species (ROS) and mitochondria (MitoTracker Red CMXros) in control and Drp1

Trim-Away embryos. Arrowheads represent regions of marked mitochondrial

aggregation in Drp1 Trim-Away embryos. Magnified images in the box areas are shown

in the right panels. Scale bar, 20 µm.

(E) Quantification of ROS enrichment levels in mitochondria by calculating the ratio of

H2DCFDA fluorescence intensity inside mitochondria to that outside mitochondria.

Data are represented as mean ± SD and P values calculated using unpaired two-tailed t

tests with Welch's correction. ***$p < 0.001$.

1045

Figure 5. Misplaced mitochondria cause chromosome segregation defects leading

to binucleation.

(A) Representative time-lapse images (maximum-intensity Z projection) of

mitochondria and chromosomes in Drp1 Trim-Away zygotes expressing mt-GFP and

H2B-mCherry. Time is relative to the onset of NEB. Scale bar, 20 μm.

(B) Representative immunofluorescence images of control (n = 10) and Drp1-depleted

(n = 31) zygotes at metaphase of the first cleavage. Pericentrin (red), microtubules

(green), and DNA (blue) were imaged by confocal microscopy. White arrowheads

indicate spindle poles. Scale bars, 10 μm.

(C) The position of chromosomes relative to the center of zygotes. The centroids of two

sets of chromosomes were tracked, and only the displacements projected in the

direction pointing from initial (NEB) to the final position (just before anaphase onset)

were indicated. Data for three representative zygotes were presented and each color

represents a zygote. Images of mitochondria at the final position automatically had their

thresholds stipulated using Fiji (Moments algorithm) and merged with images filled

inside the outline with the color corresponding to the chromosomes.

(D) Quantification of the distance of chromosomes to cell center just before anaphase

onset.

(E) Cell division abnormalities at the first cleavage. Note that binulceated blastomeres

in Drp1-depleted embryos result from various chromosome segregation defects (see

also Figure 5-figure supplement 1C and 1D).

(F) Schematic summary of results highlighting significant findings in this study. Drp1

mediate mitochondria fragmentation, ensuring symmetric mitochondrial inheritance

between daughter blastomeres. In normal embryos, fragmented mitochondria (green)

and the ER (pink) uniformly surround two spindles, allowing parental spindles (blue

and red) to assemble at the center of metaphase zygotes. Drp1 depletion increases

mitochondrial aggregation and asymmetric distribution. Misplaced mitochondria disturb

the organelle positioning during the first cleavage division leading to asymmetric

organelle inheritance and chromosome missegregation, which in turn results in early

embryonic arrest.

Data are represented as mean ± SD and P values calculated using unpaired two-tailed t

tests with Welch's correction. ***$p < 0.001$.

1078

Figure 5-figure supplement 1. Marked mitochondrial centration causes

chromosome segregation defects.

(A) A representative image of mitochondria (red), microtubules (green) and DNA (blue)

in Drp1-depleted metaphase zygotes (n = 22). Prior to fixation, zygotes were stained

with MitoTracker Red CMXRos for 0.5 h. Scale bars; 10 µm. The right panel represents

a schematic diagram showing mitochondria located in the center of two spindles based

on the left panel.

(B) Representative EM images (n = 6) showing mitochondria clustered in zygote center

and metaphase chromosomes, as surrounded by dashed lines, located in the vicinity of

the mitochondria. Representative higher-magnification images (31 sections) of the

boxed areas. Scale bar left: 10 μm; middle and right: 1 μm.

(C and D) Examples of two typical cleavage defects leading to binucleation in Drp1

Trim-Away zygotes. Time-lapse images (maximum-intensity Z projection) of

mitochondria (mtGFP) and chromosomes (H2B-mCherry) during the first cleavage

division show the mitochondrial aggregation sits between two sets of chromosomes

which transiently segregated into three (C) or four (D) blastomeres before forming

binucleated (arrowheads) 2-cell embryos. Time is relative to the onset of NEBD. Scale

bar, 20 μm.

1097

1098   **Figure 5-Video 1.** Binuclear formation in Drp1-depleted embryos, related to Fig. 5.

1099   Representative time-lapse movies of mitochondria (mt-GFP) and chromosomes

1100   (H2B-mCherry) in Drp1 Trim-Away zygotes. Time is relative to the onset of NEB

1101   (hours_minutes). Z-projection of 10 sections, 3.5 μm apart; time interval: 15 min.

1102

Figure 5-Video 2. Examples of two typical cleavage defects leading to binucleation in

Drp1 Trim-Away zygotes, related to Fig. 5 and Fig. S5. Time-lapse movies of

mitochondria (mt-GFP) and chromosomes (H2B-mCherry) in the zygote showed that

multiple cleavage furrows formed between the different sets of chromosomes leading to

transient formation of three or four blastomeres, which failed to complete cytokinesis

and collapsed back to form binucleated 2-cell embryos. Time is relative to the onset of

NEB (hours_minutes). Z-projection of 10 sections, 3.5 µm apart; time interval: 10 min.

1110

1111

1112

1113   **References**

1114   Abrisch, R.G., S.C. Gumbin, B.T. Wisniewski, L.L. Lackner, and G.K. Voeltz. 2020. Fission
1115   and fusion machineries converge at ER contact sites to regulate mitochondrial
1116   morphology. *J Cell Biol.* 219.

1117   Adhikari, D., I.W. Lee, U. Al-Zubaidi, J. Liu, Q.H. Zhang, W.S. Yuen, L. He, Y. Winstanley, H.
1118   Sesaki, J.R. Mann, R.L. Robker, and J. Carroll. 2022. Depletion of oocyte
1119   dynamin-related protein 1 shows maternal-effect abnormalities in embryonic
1120   development. *Sci Adv.* 8:eabl8070.

1121   Ball, E.H., and S.J. Singer. 1982. Mitochondria are associated with microtubules and not
1122   with intermediate filaments in cultured fibroblasts. *Proc Natl Acad Sci U S A.*
1123   79:123-126.

1124   Carlton, J.G., H. Jones, and U.S. Eggert. 2020. Membrane and organelle dynamics during
1125   cell division. *Nat Rev Mol Cell Biol.* 21:151-166.

1126   Chaigne, A., C. Campillo, R. Voituriez, N.S. Gov, C. Sykes, M.H. Verlhac, and M.E. Terret.
1127   2016. F-actin mechanics control spindle centring in the mouse zygote. *Nat Commun.*
1128   7:10253.

1129   Chung, J.Y., J.A. Steen, and T.L. Schwarz. 2016. Phosphorylation-Induced Motor Shedding
1130   Is Required at Mitosis for Proper Distribution and Passive Inheritance of
1131   Mitochondria. *Cell Rep.* 16:2142-2155.

1132   Clift, D., W.A. McEwan, L.I. Labzin, V. Konieczny, B. Mogessie, L.C. James, and M. Schuh.
1133   2017. A Method for the Acute and Rapid Degradation of Endogenous Proteins. *Cell.*
1134   171:1692-1706 e1618.

1135   Dumollard, R., M. Duchen, and J. Carroll. 2007. The role of mitochondrial function in the
1136   oocyte and embryo. *Curr Top Dev Biol.* 77:21-49.

1137   English, A.R., and G.K. Voeltz. 2013. Endoplasmic reticulum structure and interconnections
1138   with other organelles. *Cold Spring Harb Perspect Biol.* 5:a013227.

1139   Fan, W., K.G. Waymire, N. Narula, P. Li, C. Rocher, P.E. Coskun, M.A. Vannan, J. Narula,
1140   G.R. Macgregor, and D.C. Wallace. 2008. A mouse model of mitochondrial disease
1141   reveals germline selection against severe mtDNA mutations. *Science.* 319:958-962.

1142   Figueroa-Romero, C., J.A. Iniguez-Lluhi, J. Stadler, C.R. Chang, D. Arnoult, P.J. Keller, Y.
1143   Hong, C. Blackstone, and E.L. Feldman. 2009. SUMOylation of the mitochondrial
1144   fission protein Drp1 occurs at multiple nonconsensus sites within the B domain and

is linked to its activity cycle. *FASEB J.* 23:3917-3927.

Friedman, J.R., L.L. Lackner, M. West, J.R. DiBenedetto, J. Nunnari, and G.K. Voeltz. 2011.
ER tubules mark sites of mitochondrial division. *Science*. 334:358-362.

Gomes Paim, L.M., and G. FitzHarris. 2020. The impact of embryo binucleation depends
upon its origin. *Reproduction*. 160:V1-V4.

Hardy, K., R.M. Winston, and A.H. Handyside. 1993. Binucleate blastomeres in
preimplantation human embryos in vitro: failure of cytokinesis during early
cleavage. *J Reprod Fertil*. 98:549-558.

Hendzel, M.J., Y. Wei, M.A. Mancini, A. Van Hooser, T. Ranalli, B.R. Brinkley, D.P.
Bazett-Jones, and C.D. Allis. 1997. Mitosis-specific phosphorylation of histone H3
initiates primarily within pericentromeric heterochromatin during G2 and spreads
in an ordered fashion coincident with mitotic chromosome condensation.
Chromosoma. 106:348-360.

Hyslop, L.A., P. Blakeley, L. Craven, J. Richardson, N.M. Fogarty, E. Fragouli, M. Lamb, S.E.
Wamaitha, N. Prathalingam, Q. Zhang, H. O'Keefe, Y. Takeda, L. Arizzi, S.
Alfarawati, H.A. Tuppen, L. Irving, D. Kalleas, M. Choudhary, D. Wells, A.P.
Murdoch, D.M. Turnbull, K.K. Niakan, and M. Herbert. 2016. Towards clinical
application of pronuclear transfer to prevent mitochondrial DNA disease. *Nature*.
534:383-386.

Imamura, H., K.P. Nhat, H. Togawa, K. Saito, R. Iino, Y. Kato-Yamada, T. Nagai, and H. Noji.
2009. Visualization of ATP levels inside single living cells with fluorescence
resonance energy transfer-based genetically encoded indicators. *Proc Natl Acad Sci*
U S A. 106:15651-15656.

Ishihara, N., M. Nomura, A. Jofuku, H. Kato, S.O. Suzuki, K. Masuda, H. Otera, Y.
Nakanishi, I. Nonaka, Y. Goto, N. Taguchi, H. Morinaga, M. Maeda, R. Takayanagi,
S. Yokota, and K. Mihara. 2009. Mitochondrial fission factor Drp1 is essential for
embryonic development and synapse formation in mice. *Nat Cell Biol*. 11:958-966.

Kang, E., J. Wu, N.M. Gutierrez, A. Koski, R. Tippner-Hedges, K. Agaronyan, A.
Platero-Luengo, P. Martinez-Redondo, H. Ma, Y. Lee, T. Hayama, C. Van Dyken, X.
Wang, S. Luo, R. Ahmed, Y. Li, D. Ji, R. Kayali, C. Cinnioglu, S. Olson, J. Jensen, D.
Battaglia, D. Lee, D. Wu, T. Huang, D.P. Wolf, D. Temiakov, J.C. Belmonte, P. Amato,
and S. Mitalipov. 2016. Mitochondrial replacement in human oocytes carrying
pathogenic mitochondrial DNA mutations. *Nature*. 540:270-275.

Katajisto, P., J. Dohla, C.L. Chaffer, N. Pentinmikko, N. Marjanovic, S. Iqbal, R. Zoncu, W.
Chen, R.A. Weinberg, and D.M. Sabatini. 2015. Stem cells. Asymmetric apportioning
of aged mitochondria between daughter cells is required for stemness. *Science.*
348:340-343.

Koch, A., M. Thiemann, M. Grabenbauer, Y. Yoon, M.A. McNiven, and M. Schrader. 2003.
Dynamin-like protein 1 is involved in peroxisomal fission. *J Biol Chem.*
278:8597-8605.

Korobova, F., V. Ramabhadran, and H.N. Higgs. 2013. An actin-dependent step in
mitochondrial fission mediated by the ER-associated formin INF2. *Science.*
339:464-467.

Kraus, F., K. Roy, T.J. Pucadyil, and M.T. Ryan. 2021. Function and regulation of the
divisome for mitochondrial fission. *Nature.* 590:57-66.

Kumar, R.P., S. Ray, P. Home, B. Saha, B. Bhattacharya, H.M. Wilkins, H. Chavan, A.
Ganguly, J. Milano-Foster, A. Paul, P. Krishnamurthy, R.H. Swerdlow, and S. Paul.
2018. Regulation of energy metabolism during early mammalian development:
TEAD4 controls mitochondrial transcription. *Development.* 145.

Ladstatter, S., and K. Tachibana-Konwalski. 2016. A Surveillance Mechanism Ensures
Repair of DNA Lesions during Zygotic Reprogramming. *Cell.* 167:1774-1787 e1713.

Lee, H.S., H. Ma, R.C. Juanes, M. Tachibana, M. Sparman, J. Woodward, C. Ramsey, J. Xu,
E.J. Kang, P. Amato, G. Mair, R. Steinborn, and S. Mitalipov. 2012. Rapid
mitochondrial DNA segregation in primate preimplantation embryos precedes
somatic and germline bottleneck. *Cell Rep.* 1:506-515.

Lee, I.W., D. Adhikari, and J. Carroll. 2022. Miro1 depletion disrupts spatial distribution of
mitochondria and leads to oocyte maturation defects. *Front Cell Dev Biol.*
10:986454.

Lieber, T., S.P. Jeedigunta, J.M. Palozzi, R. Lehmann, and T.R. Hurd. 2019. Mitochondrial
fragmentation drives selective removal of deleterious mtDNA in the germline.
Nature. 570:380-384.

Lopez-Domenech, G., C. Covill-Cooke, D. Ivankovic, E.F. Halff, D.F. Sheehan, R. Norkett, N.
Birsa, and J.T. Kittler. 2018. Miro proteins coordinate microtubule- and
actin-dependent mitochondrial transport and distribution. *EMBO J.* 37:321-336.

Mahrous, E., Q. Yang, and H.J. Clarke. 2012. Regulation of mitochondrial DNA
accumulation during oocyte growth and meiotic maturation in the mouse.

Reproduction. 144:177-185.

Majstrowicz, K., U. Honnert, P. Nikolaus, V. Schwarz, S.J. Oeding, S.A. Hemkemeyer, and M.

Bahler. 2021. Coordination of mitochondrial and cellular dynamics by the

actin-based motor Myo19. *J Cell Sci*. 134.

Meriano, J., C. Clark, K. Cadesky, and C.A. Laskin. 2004. Binucleated and micronucleated

blastomeres in embryos derived from human assisted reproduction cycles. *Reprod*

Biomed Online. 9:511-520.

Mishra, P., and D.C. Chan. 2014. Mitochondrial dynamics and inheritance during cell

division, development and disease. *Nat Rev Mol Cell Biol*. 15:634-646.

Moore, A.S., S.M. Coscia, C.L. Simpson, F.E. Ortega, E.C. Wait, J.M. Heddleston, J.J.

Nirschl, C.J. Obara, P. Guedes-Dias, C.A. Boecker, T.L. Chew, J.A. Theriot, J.

Lippincott-Schwartz, and E.L.F. Holzbaur. 2021. Actin cables and comet tails

organize mitochondrial networks in mitosis. *Nature*. 591:659-664.

Pecot, T., L. Zengzhen, J. Boulanger, J. Salamero, and C. Kervrann. 2018. A quantitative

approach for analyzing the spatio-temporal distribution of 3D intracellular events in

fluorescence microscopy. *Elife*. 7.

Poulton, J., M.R. Chiaratti, F.V. Meirelles, S. Kennedy, D. Wells, and I.J. Holt. 2010.

Transmission of mitochondrial DNA diseases and ways to prevent them. *PLoS Genet*.

6.

Qian, W., S. Choi, G.A. Gibson, S.C. Watkins, C.J. Bakkenist, and B. Van Houten. 2012.

Mitochondrial hyperfusion induced by loss of the fission protein Drp1 causes

ATM-dependent G2/M arrest and aneuploidy through DNA replication stress. *J Cell*

Sci. 125:5745-5757.

Reichmann, J., B. Nijmeijer, M.J. Hossain, M. Eguren, I. Schneider, A.Z. Politi, M.J. Roberti,

L. Hufnagel, T. Hiiragi, and J. Ellenberg. 2018. Dual-spindle formation in zygotes

keeps parental genomes apart in early mammalian embryos. *Science*. 361:189-193.

Rizzuto, R., D. De Stefani, A. Raffaello, and C. Mammucari. 2012. Mitochondria as sensors

and regulators of calcium signalling. *Nat Rev Mol Cell Biol*. 13:566-578.

Rohn, J.L., J.V. Patel, B. Neumann, J. Bulkescher, N. McHedlishvili, R.C. McMullan, O.A.

Quintero, J. Ellenberg, and B. Baum. 2014. Myo19 ensures symmetric partitioning

of mitochondria and coupling of mitochondrial segregation to cell division. *Curr Biol*.

24:2598-2605.

Scheffler, K., J. Uraji, I. Jentoft, T. Cavazza, E. Monnich, B. Mogessie, and M. Schuh. 2021.

Two mechanisms drive pronuclear migration in mouse zygotes. *Nat Commun.*
12:841.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch,
C. Rueden, S. Saalfeld, B. Schmid, J.Y. Tinevez, D.J. White, V. Hartenstein, K.
Eliceiri, P. Tomancak, and A. Cardona. 2012. Fiji: an open-source platform for
biological-image analysis. *Nat Methods.* 9:676-682.

Seikkula, J., S. Oksjoki, S. Hurme, H. Mankonen, P. Polo-Kantola, and V. Jokimaa. 2018.
Pregnancy and perinatal outcomes after transfer of binucleated or multinucleated
frozen-thawed embryos: a case-control study. *Reprod Biomed Online.* 36:607-613.

Twig, G., A. Elorza, A.J. Molina, H. Mohamed, J.D. Wikstrom, G. Walzer, L. Stiles, S.E.
Haigh, S. Katz, G. Las, J. Alroy, M. Wu, B.F. Py, J. Yuan, J.T. Deeney, B.E. Corkey,
and O.S. Shirihai. 2008. Fission and selective fusion govern mitochondrial
segregation and elimination by autophagy. *EMBO J.* 27:433-446.

Udagawa, O., T. Ishihara, M. Maeda, Y. Matsunaga, S. Tsukamoto, N. Kawano, K. Miyado,
H. Shitara, S. Yokota, M. Nomura, K. Mihara, N. Mizushima, and N. Ishihara. 2014.
Mitochondrial fission factor Drp1 maintains oocyte quality via dynamic
rearrangement of multiple organelles. *Curr Biol.* 24:2451-2458.

Wakabayashi, J., Z. Zhang, N. Wakabayashi, Y. Tamura, M. Fukaya, T.W. Kensler, M. Iijima,
and H. Sesaki. 2009. The dynamin-related GTPase Drp1 is required for embryonic
and brain development in mice. *J Cell Biol.* 186:805-816.

Wakai, T., Y. Harada, K. Miyado, and T. Kono. 2014. Mitochondrial dynamics controlled by
mitofusins define organelle positioning and movement during mouse oocyte
maturation. *Mol Hum Reprod.* 20:1090-1100.

Westermann, B. 2010. Mitochondrial fusion and fission in cell life and death. *Nat Rev Mol*
Cell Biol. 11:872-884.

Yamano, K., and R.J. Youle. 2011. Coupling mitochondrial and cell division. *Nat Cell Biol.*
13:1026-1027.

Youle, R.J., and A.M. van der Bliek. 2012. Mitochondrial fission, fusion, and stress. *Science.*
337:1062-1065.

Zhou, C.J., X.Y. Wang, Y.H. Dong, D.H. Wang, Z. Han, X.J. Zhang, Q.Y. Sun, J. Carroll, and
C.G. Liang. 2022. CENP-F-dependent DRP1 function regulates APC/C activity
during oocyte meiosis I. *Nat Commun.* 13:7732.

1276

Figure 1

Figure 1–figure supplement 1

A Metaphase II | Interphase (PN) | Prophase | Prometaphase | Metaphase | Anaphase | Telophase | Interphase (2-cell)

B EB3-GFP / Mt-DsRed | 1st | 2nd | 3rd

Mt-DsRed

C Mitochondrial intensity ratio at spindle periphery

****
****
ns

(10) (11) (16)

1st 2nd 3rd
(Cleavage)

D Phalloidin | MitoTracker | Merge + DAPI

Interphase (PN)

Metaphase

Interphase (2-cell)

E DMSO | Lat A

Phalloidin + DAPI

F DMSO | Lat A

MitoTracker

G Mitochondrial inheritance ratio

*

(11) (15)

Control LatA

H Blastomere area ratio

***

(11) (15)

Control LatA

I R = 0.6065 P = 0.0165

Blastomere area ratio

Mitochondrial inheritance ratio

Figure 1–figure supplement 2

Figure 2

Figure 2–figure supplement 1

Figure 2–figure supplement 2

Figure 3

Figure 3–figure supplement 1

A Control

EB3-GFP Mt-DsRed

-15 min   0 min   +15 min   +30 min

Trim-Away

EB3-GFP Mt-DsRed

-15 min   0 min   +15 min   +30 min

B Angler distribution at metaphase

Control
Microtubules
Mitochondria

Trim-Away

0.012
0.01
0.008
0.006
0.004
0.002
0

*

s.d. of mitochondrial
intensity per 60°

(7)   (7)

Control   Trim-Away

C Ratio (YFP/CFP)

AT 1.03   AT1.03 RK   Mitotracker

Control

Trim-Away

B   A   B   A   B   A   B   A

D Control

NEBD   Metaphase   Telophase   Interphase

mt-GFP

0:00   1:30   1:45   2:15

ER-mCherry

0:00   1:30   1:45   2:15

Trim-Away

NEBD   Metaphase   Telophase   Interphase

mt-GFP

0:00   1:30   1:45   2:15

ER-mCherry

0:00   1:30   1:45   2:15

Figure 4

Figure 4–figure supplement 1

A — Control (46h) / Trim-Away (46h): DAPI, Phospho H3-ser10

B — Control (46h) / Trim-Away (46h): DAPI, γH2X, DAPI / γH2X / Mitochondria

C — No. γH2X foci (>0.5 μm²); (16) (16); Control, Trim-Away; ***

D — Control / Trim-Away: H2DCFDA, MitoTracker, Merge, Trans

E — H2DCFDA intensity ratio (inside/outside); (23) (20); Control, Trim-Away; ****

Figure 5

A Trim-away

NEB

mt-GFP
H2B-mCherry

-3:00   0:00   1:30   2:00   3:00

H2B-mCherry

B Control   Trim-away

DNA
αTublin
Pericentrin

C Control   Trim-Away

D (12)   (12)

***

Distance to the cell center (μm)

Control Trim-Away

E (51)   (43)

Proportion (%)

Control Trim-away

4 blastomeres with 1 nucleus

3 blastomeres (2 blastomeres
with 1 nucleus, 1 blastomere
with 2 nuclei)

2 blastomeres with 2 nuclei

2 blastomeres with 2 nuclei
(transient 3 blastomeres)

2 blastomeres with 2 nuclei
(transient 4 blastomeres)

Binucleation

2 blastomeres with 1 nucleus
(Normal 2-cell embryos)

F

Normal embryos

Mitochondrial
fragmentation

Mitochondrial
aggregation

Drp1-depleted embryos

Two-pronuclei
zygote

Uniform
mitochdnrial
distribution
around parental
spindles

Equal
partitioning of
mitochondria
into daugther
blastomeres

•Symmetric
mitochondrial
inheritance

•Parental
genome
mixing

Mitochondrial
centration
prevents
dual spindle
assembly

Mitochondrial
clumps disturb
organelle
partitioning

•Asymmetric
mitochondrial
inheritance

•Spatial
confinement of
the ER

•Binucleated
blastomeres

Figure 5–figure supplement 1