1

Redistribution of fragmented mitochondria ensure symmetric organelle partitioning and faithful chromosome segregation in mitotic mouse zygotes

Haruna Gekko[1†], Ruri Nomura[1†], Daiki Kuzuhara[2], Masato Kaneyasu[1], Genpei Koseki[1]

Deepak Adhikari[3], Yasuyuki Mio[2], John Carroll[3], Tomohiro Kono[4], Hiroaki Funahashi[1]

and Takuya Wakai[1]*

[1]Department of Animal Science, Graduate School of Environment and Life Science,

Okayama University, Okayama 700-8530, Japan

[2]Reproductive Centre, Mio Fertility Clinic, Tottori, 683-0008, Japan

[3]Development and Stem Cell Program and Department of Anatomy and Developmental

Biology, Monash Biomedicine Discovery Institute, Monash University, Melbourne,

Victoria 3800, Australia

[4]Department of Bioscience, Tokyo University of Agriculture, Tokyo 156-8502, Japan

[†]These authors equally contributed to this work.

*Correspondence: t2wakai@okayama-u.ac.jp

Impact statement:

Depletion of Dynamin-related protein 1, a key regulator of mitochondrial fission, in

mouse zygotes impair symmetric organelle partitioning and chromosome segregation

leading to early developmental arrest.

Funding information

1) Japan Society for the Promotion of Science (Grant reference: 17K17905 and

20K06627): Takuya Wakai

2) National Health and Medical Research Council (NHMRC) Australia (Grant

reference: 1165627): Deepak Adhikari and John Carroll

The funders had no role in study design, data collection and interpretation, or the

decision to submit the work for publication.

Author information:

Haruna Gekko, Department of Animal Science, Graduate School of Environment and

Life Science, Okayama University, Okayama, Japan, Contribution: Formal analysis;

Investigation, No competing interests declared.

Ruri Nomura , Department of Animal Science, Graduate School of Environment and

Life Science, Okayama University, Okayama, Japan, Contribution: Formal analysis;

Investigation, No competing interests declared.

Daiki Kuzuhara, Reproductive Centre, Mio Fertility Clinic, Tottori, Japan, Contribution:

Investigation; Data curation, No competing interests declared.

Masato Kaneyasu, Department of Animal Science, Graduate School of Environment

and Life Science, Okayama University, Okayama, Japan, Contribution: Data curation;

Visualisation, No competing interests declared.

Genpei Koseki, Department of Animal Science, Graduate School of Environment and

Life Science, Okayama University, Okayama, Japan, Contribution: Formal analysis;

Data curation, No competing interests declared.

Deepak Adhikari, Development and Stem Cell Program and Department of Anatomy

and Developmental, Biology, Monash Biomedicine Discovery Institute, Monash

University, Melbourne, Australia, Contribution: Formal analysis; Investigation; Funding

acquisition

No competing interests declared.

Yasuyuki Mio, Reproductive Centre, Mio Fertility Clinic, Tottori, Japan, Contribution:

Resources, No competing interests declared.

John Carroll, Development and Stem Cell Program and Department of Anatomy and

Developmental, Biology, Monash Biomedicine Discovery Institute, Monash University,

Melbourne, Australia, Contribution: Writing – review & editing; Supervision; Funding

acquisition, No competing interests declared.

Tomohiro Kono, Department of Bioscience, Tokyo University of Agriculture, Tokyo

156-8502, Japan, Contribution: Supervision, No competing interests declared.

Hiroaki Funahashi, Department of Animal Science, Graduate School of Environment

and Life Science, Okayama University, Okayama, Japan, Contribution: Resources;

Supervision, No competing interests declared.

Takuya Wakai

Department of Animal Science, Graduate School of Environment and Life Science,

Okayama University, Okayama, Japan, Contribution: Conceptualization, Validation;

Formal analysis; Investigation; Data curation; Writing - original draft, Supervision,

Funding acquisition, No competing interests declared.

Abstract

In cleavage-stage embryos, preexisting organelles partition evenly into daughter

blastomeres without significant cell growth after symmetric cell division. The presence

of mitochondrial DNA within mitochondria and its restricted replication during

preimplantation development makes their inheritance particularly important. While

chromosomes are precisely segregated by the mitotic spindle, the mechanisms

controlling mitochondrial partitioning remain poorly understood. In this study, we

investigate the mechanism by which Dynamin-related protein 1 (Drp1) controls the

mitochondrial redistribution and partitioning during embryonic cleavage. Deletion of

Drp1 in mouse zygotes causes marked mitochondrial aggregation, and the majority of

embryos arrest at the 2-cell stage. Clumped mitochondria are located in the center of

mitotic Drp1-depleted zygotes with less uniform distribution, thereby preventing their

symmetric partitioning. Asymmetric mitochondrial inheritance is accompanied by

functionally inequivalent blastomeres with biased ATP and endoplasmic reticulum Ca^{2+}

levels. We also find that marked mitochondrial centration in Drp1-depleted zygotes

prevents the assembly of parental chromosomes, resulting in chromosome segregation

defects and binucleation. Thus, mitochondrial fragmentation mediated by Drp1 ensure

proper organelle positioning and partitioning into functional daughters during the first

embryonic cleavage.

Keywords: Mitochondrial dynamics, dynamin-related protein 1, embryo cleavage,

preimplantation development, organelle inheritance, ER calcium stores, chromosome

segregation, binuclear formation, assisted reproductive technologies

Introduction

Preimplantation development of mammalian embryos consists of a series of symmetric cell divisions without significant cell growth, known as embryonic cleavage, resulting in approximately half of the cytoplasm in each resulting cell, called blastomere. The segregation of maternal content between daughter blastomeres allows rapid proliferation of a single-cell zygote into multicellular organisms. To ensure the correct segregation of cellular content to produce functional daughters, cells must duplicate and apportion their various organelles with high accuracy (Carlton et al., 2020). Chromosomes are assembled and segregated into daughter cells with the formation of mitotic spindles, whereas membrane-bound organelles, such as the endoplasmic reticulum (ER) and mitochondria, are typically generated from existing structures, and both daughters receive a share of these components during cell division. This is particularly important in the case of mitochondria because mitochondria are semi-autonomous organelles that contain their own genome, mitochondrial DNA (mtDNA). Growing oocytes exponentially increase their mtDNA, and a fully grown oocyte contains more than 200 000 copies of mtDNA (Mahrous et al., 2012). After fertilization, abundant preexisting mtDNA is segregated among daughter cells without replication during preimplantation development (Poulton et al., 2010). As mitochondria are the primary source of ATP production via oxidative phosphorylation in oocytes and early embryos owing to low glycolytic activity (Dumollard et al., 2007), proper inheritance must be ensured for cell energetic homogeneity among blastomeres.

Mitochondria also play an important role in the regulation of intracellular Ca^{2+} signaling,

which in turn controls cell metabolism and cell death pathways (Rizzuto et al., 2012).

Thus, defects in mitochondrial inheritance are likely to have serious consequences for

daughter blastomeres and manifest as the failure of pre- or post-implantation

development.

Mitochondria are highly dynamic organelles that move along cytoskeletal networks

and undergo continuous fusion and fission; these dynamics allow cells to respond and

adapt to various intracellular and extracellular changes and maintain cellular function

and homeostasis (Mishra and Chan, 2014; Westermann, 2010). Active partitioning of

mitochondria is observed in budding yeast fission, which undergoes asymmetric cell

division; mitochondria actively move along actin filaments into the daughter cell

(Mishra and Chan, 2014). In mammalian cells that divide symmetrically, the

partitioning of mitochondria is largely due to passive processes, as fragmented and

dispersed mitochondrial distribution allows for stochastic and roughly equal partitioning

into daughter cells. Mitochondrial fission is thus important for faithful inheritance and

proper intracellular distribution of the mitochondria (Westermann, 2010). Whether

active transport of mitochondria by the cytoskeleton contributes to mitochondrial

inheritance in dividing animal cells remains controversial. Microtubules function as

cytoskeletal platforms for the distribution and transport of mitochondria. During mitosis,

mitochondria decouple from spindle microtubules and disperse in the cell periphery,

allowing for passive segregation (Chung et al., 2016). Furthermore, recent studies have

shown that actin filaments took over as the dominant mitochondrial scaffold following

mitochondrial release from microtubules at the start of mitosis (Moore et al., 2021).

Myosin 19 (Myo19), a mitochondrial-localized myosin in vertebrates, acts as an

actin-based motor for mitochondrial dynamics. Notably, depletion of Myo19 causes

asymmetrical mitochondrial inheritance between daughter cells (Rohn et al., 2014),

though the molecular identification of Myo19 in mammalian oocytes/eggs has yet to be

shown.

Dynamin-related protein 1 (Drp1) is a key regulator of mitochondrial fission in

many eukaryotic organisms. Drp1 is recruited to the mitochondrial membrane where it

forms helical oligomers that induce membrane constriction and severing (Westermann,

2010). The indispensable role of Drp1-mediated mitochondrial fission in mammalian

oocytes has been documented using conditional knockout (KO) mice (Udagawa et al.,

2014). Drp1 KO oocytes undergo meiotic arrest due to severe organelle aggregation,

which hinders the functional importance of Drp1 in mature oocytes and embryos. More

recently, mature oocytes have been successfully isolated from juvenile Drp1 conditional

KO mice (Adhikari et al., 2022). After parthenogenetic activation of Drp1 KO oocytes,

few embryos develop to the blastocyst stage. Nevertheless, the precise role of

mitochondrial fission in normal embryos remains elusive. To this end, we employed

Trim-Away, a method to degrade endogenous proteins via the proteasome-mediated

pathway (Clift et al., 2017), which acutely depletes large amounts of storage proteins in

the oocyte, allowing loss-of-function studies during the preimplantation development

from the zygotic stage. In this study, we show that Drp1 depletion in fertilized zygotes

causes marked mitochondrial aggregation and early embryonic arrest. Live imaging of

mitochondria during the first cleavage division reveals that loss of Drp1 disturbs the

spatiotemporal mitochondrial dynamics, resulting in an asymmetric mitochondrial

inheritance between daughter blastomeres with functional heterogeneities. We also find

that misplaced mitochondria impair the assembly of parental spindles leading to

binucleated blastomeres, which closely resemble a clinical phenotype of human

embryos in *in vitro* fertilization (IVF) procedures.

Results

Fragmented mitochondria are redistributed during the first embryonic cleavage

and equally partitioned into daughter blastomeres

We first imaged mitochondria and chromosomes in zygotes expressing

mitochondrially-targeted GFP (mt-GFP) and histone H2B-mCherry (**Figure 1A; Figure**

1-Video 1). Mitochondria in interphase zygotes progressively accumulated around the

two pronuclei (**Figure 1B, left**), encircled metaphase chromosomes after nuclear

envelope breakdown (NEB). During the transition from anaphase to telophase, the

cytokinetic furrow constricts the mitochondrial ring that effectively divides the

mitochondria approximately equally into the two daughter blastomeres and

mitochondria were dispersed throughout the cytoplasm in the interphase 2-cell embryos

(**Figure 1B, right**). Immunofluorescence staining of microtubules also showed

mitochondria progressively surround the spindle and disperse back into the cytoplasm

after the first cleavage (**Figure 1-figure supplement 1A**). The accumulation of

mitochondria around the spindle is unique to the first cleavage division (**Figure**

1-figure supplement 1B and 1C), as live imaging of fluorescently labeled

microtubules (EB3-GFP) and mitochondria (mt-DsRed) confirmed spindle-associated

accumulation of mitochondria in one-cell zygotes, but no evidence of mitochondrial

accumulation was observed in the spindles of 2-cell or 4-cell stage embryos. By

electron microscopy (EM), we found that these accumulated mitochondria around the

metaphase spindle were highly fragmented (**Figure 1C**). The ERs were also abundantly

distributed around the spindle and were in partial contact with the mitochondria, and

enriched bundles of actin filaments were found around these organelles.

Interphase zygotes stained with fluorescence-labelled phalloidin displayed that

mitochondrial distribution is closely associated with F-actin cytoplasmic meshwork

(**Figure 1D; Figure 1-figure supplement 1D**). This cytoplasmic meshwork was

subsequently reorganized around the spindle and increased co-localization with

mitochondria (**Figure 1E**). In interphase 2-cell stage embryos, the F-actin meshwork

was seen again, albeit with significantly reduced overlap with mitochondria. To test

whether the F-actin is involved in mitochondrial distribution and partitioning, zygotes

were treated with 1 μM latrunculin A for 1 hr just after NEB, sufficient to F-actin

depolymerization (**Figure 1-figure supplement 1E**). Disruption of F-actin organization

resulted in asymmetric mitochondrial inheritance (**Figure 1-figure supplement 1F and**

1G), but also increased cell size asymmetry (**Figure 1-figure supplement 1H**), and

these two asymmetries were correlated (**Figure 1-figure supplement 1I**). Since the role

of F-actin in many cellular events, such as cytokinesis, preclude them as targets for

experimentally manipulating mitochondrial distribution, we have therefore targeted

Myo19 as a strategy to alter the mitochondrial distribution and ask if it is critical for the

early development. Asymmetric mitochondrial inheritance has been previously reported

in cells lacking Myo19 (Majstrowicz et al., 2021; Moore et al., 2021; Rohn et al., 2014).

Myo19 was expressed in both interphase and metaphase zygotes, where its localization

was consistent with the mitochondrial distribution (**Figure 1F**). Acute depletion of

Myo19 at the PN stage by Trim-Away was confirmed by Western blot analysis, which

was not observed in control zygotes injected with Trim21 mRNA and control IgG

(**Figure 1G**). Myo19 depletion did not affect developmental competence of the embryos

to the blastocyst stage (**Figure 1H**). Unexpectedly, no significant asymmetry was

observed in the mitochondrial distribution and partitioning between the daughter

blastomeres in Myo19-depleted 2-cell embryos (**Figure 1I and 1J**). Overall,

fragmented mitochondria are closely associated with cytoplasmic F-actin during the

first cleavage, but active transport by the actin motor is not directly involved in

mitochondrial partitioning.

Loss of Drp1 induces mitochondrial aggregation and disturbs subcellular organelle

compartments

The highly fragmented mitochondrial morphology appears to facilitate

mitochondrial partitioning during embryonic cleavage. To understand the underlying

mechanism of this process, we examined the role of Drp1, a key regulator of

mitochondrial fission. Western blot analysis revealed that Drp1 was expressed in PN

zygotes, and comparable protein levels were found in the morula stage, whereas their

expression was remarkably decreased at the blastocyst stage (**Figure 2A**). To

demonstrate the role of Drp1-mediated mitochondrial fission in preimplantation

embryos, we employed Trim-Away. Co-injection of Trim21 mRNA and an antibody

against Drp1 into zygotes 20–22 h post hCG reduced the Drp1 protein to nearly

undetectable levels by 5 h post-injection at the PN stage, whereas co-injection of

Trim21 mRNA and control IgG failed to trigger protein degradation (**Figure 2B**)**.**

Mitochondria in live 2-cell embryos (37/37) expressing mt-GFP were interspersed

throughout the cytoplasm. In contrast, almost all Drp1 Trim-Away embryos (55/57)

exhibited marked mitochondrial aggregation that was reversed by ectopic expression of

mCherry-tagged Drp1 (mCh-Drp1) (50/53) (**Figure 2C and 2D**). We confirmed a

similar, mitochondrial aggregation occurred in parthenogenetic embryos derived from

Drp1 (-/-) parthenotes (**Figure 2E**). EM images of control 2-cell embryos showed that

rounded mitochondria with less developed cristae were dispersed throughout the

cytoplasm, whereas swollen or partially elongated mitochondria were aggregated in

Drp1-depleted embryos (**Figure 2F**). The mean length of the long axis of mitochondria

in Drp1-depleted embryos (0.67 ± 0.32 μm) was significantly ($P < 0.0001$) greater than

that of control embryos (0.40 ± 0.12 μm) (**Figure 2G**). To clarify the effects of these

morphological changes in mitochondria on their energy production, we compared ATP

levels between control and Drp1 Trim-Away embryos at the 2-cell stages. To analyze

intracellular ATP concentrations, we expressed a fluorescence resonance energy transfer

(FRET)-based ATP biosensor, ATeam AT1.03, with an emission ratio of AT1.03

fluorescence (YFP/CFP) used to estimate ATP levels. ATP levels in Drp1-depleted

embryos were indistinguishable from those in the control embryos (**Figure 2H**). No

FRET signal was detected in embryos expressing a mutant version of the ATP probe

AT1.03RK that cannot bind ATP.

Recent studies have focused on the interactions between organelles at membrane

contact sites (MCSs) and their roles in maintaining cellular homeostasis (Abrisch et al.,

2020). The ER is a continuous membrane-bound organelle with MCSs with the plasma

membrane, mitochondria, Golgi, and peroxisomes. Since mitochondrial dynamics are

spatially coordinated at the ER–mitochondria MCSs, we surmised that mitochondrial

aggregation due to the loss of Drp1 would compromise the organization of ER MCSs.

To visualize the distribution of mitochondria with other organelles in live embryos,

fluorescently tagged ER (ER-mCherry), Golgi (Golgi-mCherry), and peroxisomes

(PEX-mCherry) were co-expressed with mt-GFP. The ER was partially confined to the

regions of the mitochondrial aggregation in Drp1 Trim-Away embryos, which appeared

to disturb the endogenous ER network (**Figure 2-figure supplement 1A**). In contrast,

the subcellular distribution of the Golgi was not clearly altered by Drp1 depletion

(**Figure 2-figure supplement 1B**). Drp1 is also known to regulate peroxisomal fission,

as elongated peroxisomes are seen in Drp1-deficient cells (Koch et al., 2003); however,

the subcellular distribution of peroxisomes in mammalian embryos has not been

reported. The peroxisomes in normal 2-cell embryos showed a punctate distribution in

the cytoplasm, whereas these puncta partially aggregated following the Drp1 depletion

by Trim-Away (**Figure 2-figure supplement 1C**), although it appeared to be

independent of mitochondria in terms of spatial positioning.

Drp1-mediated mitochondrial fragmentation is required for the symmetric mitochondrial partitioning into two functional daughter blastomeres

To decipher the spatiotemporal regulation of mitochondrial dynamics following the Drp1 depletion, we imaged mitochondria and chromosomes in Drp1 Trim-Away zygotes expressing mt-GFP and histone H2B-mCherry (**Figure 3A; Figure 3-Video 1**). Clumped mitochondria were located in the center of the metaphase zygotes which resulted in an increase in mitochondrial asymmetry at anaphase (**Figure 3B**). Live imaging of mitochondria (mt-DsRed) and spindle microtubules (EB3-GFP) revealed compared with the uniform angular positioning of around the spindle observed in control zygotes, mitochondria in Drp1-depleted zygotes lost their uniform distribution (**Figure 3-figure supplement 1A and 1B**), leading to the mitochondrial asymmetry. The biased inheritance of mitochondria was further validated by the quantification of mtDNA copy number in isolated blastomeres from 2-cell embryos (**Figure 3C**). The inheritance ratio of mtDNA in daughter blastomeres of control embryos was comparable, whereas Drp1 depletion led to biased mtDNA inheritance. Although Drp1 depletion did not affect intracellular ATP levels in the whole embryo (**Figure 2H**), we tested if the biased inheritance of mitochondria led to differences in blastomere ATP levels. Using AT1.03, ATP concentration and mitochondrial distribution were simultaneously visualized in blastomeres of 2-cell embryos (**Figure 3-figure supplement 1C**). Notably, the bias in ATP levels between blastomeres of 2-cell embryos was greater following the Drp1 depletion (**Figure 3D**). We then quantified

mtDNA copy number in each blastomere and found a positive correlation between

mtDNA inheritance and ATP levels in the individual blastomeres (**Figure 3E**).

The presence of ER-mitochondria MCSs may lead to mitochondria-associated ER

asymmetry during cleavage. Live imaging of mitochondria and ER showed that part of

the ER is confined to mitochondrial aggregation, resulting in asymmetric distribution at

the telophase of the first cleavage (**Figure 3F; Figure 3-figure supplement 1D; Figure**

3-Video 2) which was closely correlated with mitochondrial asymmetry (**Figure 3G**).

No significant asymmetry was observed in the distribution of Golgi or peroxisomes,

suggesting that these organelles are not directly confined by mitochondrial location. The

ER is the main reservoir of intracellular Ca^{2+} stores in non-excitable cells.

ER-mitochondria MCSs create localized domains of high Ca^{2+} concentration required

for Ca^{2+} transfer from the ER to the mitochondria, in which the propagation of Ca^{2+}

signals in the mitochondria controls energy metabolism (Rizzuto et al., 2012). To

explore whether alterations in the ER and mitochondrial distribution and segregation

affect intracellular Ca^{2+} homeostasis in 2-cell embryos, the Ca^{2+} response in each

blastomere after exposure to thapsigargin (Tg) in Ca^{2+} free medium, a specific inhibitor

of sarco/ER Ca^{2+}-ATPase, was compared between control and Drp1 Trim-Away

embryos (**Figure 3H and 3I**). Drp1 depletion significantly reduced

thapsigargin-induced Ca^{2+}-release and also caused significant variability between

blastomeres of individual 2-cell embryos which was not seen in controls (**Figure 3J**).

These findings indicate that Drp1 depletion disrupts mitochondrial-ER interaction

leading to compromised ER Ca^{2+} stores and that asymmetric inheritance of ER leads to

further variability in Ca^{2+} stores between daughter blastomeres.

Drp1 is required for the developmental competence of preimplantation embryos

We found that Drp1-depleted zygotes were predominantly arrested at the 2-cell

stage and few embryos developed to the blastocyst stage (**Figure 4A and 4B**). This is a

specific effect of Drp1 deletion because none of the internal control conditions

increased arrest at the 2-cell stage and arrest was completely reversed by microinjecting

Trim-away insensitive exogenous mCh-Drp1 mRNA (**Figure 4C**). Further, western blot

analysis confirmed expression of mCh-Drp1, in conditions where endogenous Drp1 was

degraded by Trim-Away (**Figure 4D**).

More than half of the Drp1 Trim-Away embryos were arrested at the 2-cell stage,

the vast majority of which did not enter the following M-phase, as assessed by the

presence of the nuclear envelope. Phosphorylated histone H3 (Ser10), localized in the

heterochromatin region in G2-phase cells, was detected in 89% (16/18) of arrested

embryos following Drp1 depletion (**Figure 4-figure supplement 1A**), suggesting that

the majority of interphase arrests occur in the G2-phase. Since G2/M interphase arrest

presumably activates a DNA damage checkpoint, we estimated DNA lesions by

immunofluorescence of γH2AX, a widely used marker for DNA damage 46 h post hCG

(corresponding to the G2-stage of the 2-cell stage). As expected, the number of γH2AX

foci in Drp1-depleted embryos was significantly greater than that in control embryos

(**Figure 4-figure supplement 1B and 1C**). The accumulation of reactive oxygen

species (ROS), which are generated as byproducts of mitochondrial energy production,

increases susceptibility to DNA damage. Intracellular ROS levels, as estimated by

H2DCFDA fluorescence, were highly accumulated in mitochondria (**Figure 4-figure**

supplement 1D).

Misplaced mitochondria during the first cleavage division impair the assembly of

parental chromosomes leading to binuclear formation

We observed that approximately 30% (17/53) of Drp1 Trim-Away embryos

consisted of 3 or 4 blastomeres despite the first cleavage (indicated by arrowheads in

Fig. 4B), suggesting that cleavage defects have occurred. Live imaging experiments

showed that Drp1 Trim-Away embryos frequently display chromosome segregation

defects. As a typical example, two pronuclei frequently fail to reach apposition but

continue into mitosis and subsequently enter anaphase leading to binucleate blastomeres

in the subsequent 2-cell embryo (**Figure 5A; Figure 5-Video 1**)**.** Immunostaining of

metaphase zygotes showed that 55% (17/31) of Drp1-depleted zygotes formed two

independent spindles with clustered microtubule organizing centers (**Figure 5B**). Both

confocal and EM images showed that large mitochondrial clumps occupied the central

region in Drp1-depleted zygotes, to which two spindles were separately located (**Figure**

5-figure supplement 1A and 1B). Tracking of two sets of chromosomes with

mitochondria following Drp1 depletion revealed that the mitochondrial aggregation

caused by Drp1-depletion sits between the pronuclei and subsequent spindles such that

the distance from chromosomes to the centre of the zygote is increased (**Figure 5C and**

5D). The duration from mitotic entry to chromosome segregation at anaphase was not

affected (not shown) but binucleation was observed in 74% (32/43) of Drp1-depleted

embryos and in none of the control zygotes (n = 51) (**Figure 5E**). In addition to the

binucleation of blastomeres via dual spindle formation, other cell division abnormalities

were apparent. In approximately half (24/43) of the zygotes, multiple cleavage furrows

formed between the different sets of chromosomes leading to transient formation of

three or four blastomeres, of which 42% (10/24) failed to complete cytokinesis and

collapsed back to form binucleated 2-cell embryos (**Figure 5-figure supplement 1C**

and 1D; Figure 5-Video 2). Together, these results suggest that proper control of

mitochondrial positioning during mitosis is crucial for assembly into a single spindle,

and marked mitochondrial centration causes chromosome segregation defects, leading

to binuclear formation.

Discussion

Mechanisms underlying symmetric mitochondrial inheritance in preimplantation

embryos

To ensure proper partitioning of intracellular organelles for two functional daughter

cells, cells undergo complex and coordinated remodeling of their cytoskeleton and

membranes during cell division through mechanisms that are less well understood than

chromosomal dynamics. As mitochondria are unique organelles with their own

genomes, their biased inheritance is expected to have detrimental effects on daughter

cell function. We showed that Drp1 mediates mitochondrial fragmentation and

dispersion, enabling symmetric partitioning between daughter blastomeres

stochastically, which is indispensable for healthy embryo development (**Figure 5F**).

Mitochondria that form a morphologically complex network and interact with

microtubules at the interphase are rapidly fragmented upon mitotic entry, detached from

the microtubules, and redistributed throughout the cytoplasm (Ball and Singer, 1982).

Mitochondrial fission is activated via the phosphorylation of Drp1 during mitosis. Two

mitotic kinases, CDK1 and Aurora A, activate Drp1, leading to enhanced mitochondrial

fission (Yamano and Youle, 2011). As the cell exits mitosis, Drp1 is sumoylated and

targeted for degradation to prevent mitochondrial fragmentation (Figueroa-Romero et

al., 2009). Although it remains to be investigated whether these cell cycle-dependent

modifications of Drp1 also occur in early embryos, mitochondria appear to be

constitutively fragmented throughout the cell cycle. This unique mitochondrial

morphology not only facilitates symmetric partitioning in cell division without cell

growth, but also appears to be well-adapted to the rapid proliferation of single-cell

zygotes into multicellular organisms before cell lineages specification. In

asymmetrically dividing cells, functionally distinct mitochondria are unequally

apportioned to influence daughter cell fate (Katajisto et al., 2015). In contrast,

symmetric cell division is intended to produce identical daughter cells with comparable

fates. In this context, the highly fragmented mitochondrial morphology in early

embryos may contribute to avoid cell fate divergence, as well as fulfill metabolic

homogeneity among blastomeres. Interestingly, the apparent morphological changes in

mitochondria occur at the blastocyst stage, as elongated mitochondria are formed in

trophectoderm (TE) cells, which is likely to be associated with altered energy

metabolism in the TE lineage (Kumar et al., 2018). Considering that the transformation

from eggs to embryos is a fundamental process in animal development involving

changes such as the transition from meiotic to mitotic spindles and maternal to zygotic

gene expression, changes in mitochondrial morphology and function may also play a

role in regulating cell fate and function in the early embryo.

The cytoskeleton is a critical regulator of organelle positioning. In contrast to a

uniformly distributed mitochondria throughout the cytoplasm during somatic cell

division (Chung et al., 2016; Moore et al., 2021), mitochondria progressively

accumulated around pronuclei and encircle the metaphase spindle of the first embryonic

cleavage. Since the two pronuclei migrate from the periphery to the central of the eggs

by the F-actin-dependent mechanism to unite chromosomes on the first mitotic spindle

(Chaigne et al., 2016; Scheffler et al., 2021), actin filaments may drive the

mitochondrial movement. In fact, mitochondrial distribution was closely associated

with cytoplasmic F-actin organization in zygotes, especially at the metaphase. Although

depolymerization of F-actin prior to metaphase led to asymmetric mitochondrial

partitioning, the cell division itself was also asymmetric, probably due to failure of

centering of the pronuclei. Given that the depletion of Myo19 did not affect the

mitochondrial inheritance between daughter blastomeres, actin-based motor may not

contribute the symmetric pertaining of mitochondria in cleavage embryos. This result is

mostly consistent with previous observations in mitochondrial Rho GTPase (Miro)

1-deleted zygotes. Miro1 is localized in the mitochondrial outer membrane and the

adaptor protein TRAK mediates the attachment of motor proteins to the mitochondria

(Lopez-Domenech et al., 2018). Miro also regulates Myo19, ensuring stability to

actin-mediated mitochondrial distribution (Lopez-Domenech et al., 2018), and thus loss

of Miro leads to asymmetric segregation of mitochondria during mitosis most likely due

to reduced levels of Myo19. In Miro1-deleted zygotes, mitochondrial dynamics was

partially disturbed, but this did not affect mitochondrial partitioning to daughter

blastomeres, and fertility was normal as a result (Lee et al., 2022). Thus, fragmented

mitochondria may be primarily important for symmetric mitochondrial partitioning in

cleavage embryos, as a less precise stochastic partitioning mechanism may be suitable

or sufficient for equally partitioning of mitochondria, which are present in abundance

and are located across the cell volume in oocytes/zygotes. Otherwise, other mechanisms

may exist that tether mitochondria to the cytoskeletal network in these cells.

The physiological roles of mitochondrial fission in preimplantation embryos

Inhibition of Drp1-mediated mitochondrial fission causes severe embryonic arrest.

Given the housekeeping cellular functions of mitochondria, dysregulation of

mitochondrial dynamics may exert profound effects on the developmental competence

of embryos. Mitochondrial dynamics are a highly regulated process, and when disrupted,

can lead to cellular and tissue pathologies. The importance of mitochondrial fission has

been intensively studied in animal cells, and emerging evidence suggests that

Drp1-mediated fission participates in diverse cellular processes, including ER contacts,

Ca^{2+} signaling, autophagy, and apoptosis (Kraus et al., 2021). Drp1-KO mice are

embryonic lethal, most likely due to brain, heart, and placental defects; however,

various tissues appear normal, and cultured cell lines continue to proliferate even

though mitochondria are unevenly segregated (Ishihara et al., 2009; Wakabayashi et al.,

2009). Cells lacking Drp1 have highly elongated mitochondria that cannot be divided

into transportable units, and typically accumulate in a limited area, leaving large parts

of the cell devoid of mitochondria. This imbalanced distribution of mitochondria may

fail to meet local energy demands and thus would be more manifest in large or extended

cells, such as neurons and oocytes/zygotes. Moreover, our data demonstrated that Drp1

depletion in cleavage embryos caused biased mitochondrial inheritance and energy

heterogeneity between blastomeres. Given the low levels of de novo mtDNA replication

during preimplantation development, these biases may be further magnified by cleavage

division progression. It is also noteworthy that we showed that the loss of Drp1 induced

mitochondrial aggregation disturbs the spatial organization of ER. The aggregated ER

subdomains likely caused leakage of Ca^{2+} stores, which may compromise the tethering

functions at the ER MCSs. Furthermore, we found that marked mitochondrial

aggregation in Drp1-depleted 2-cell embryos caused the ectopic accumulation of ROS,

which presumably led to an increase in DNA lesions. A hyperfused mitochondrial

network in Drp1-deficient cells has also been reported to cause replication stress and

chromosomal instability during the cell cycle, leading to DNA damage response (Qian

et al., 2012). Although the precise mechanisms of 2-cell stage arrest arising from

depletion of Drp1 remain elusive, the presence of a Chk1-dependent checkpoint leading

to G2 arrest in 2-cell mouse embryos (Ladstatter and Tachibana-Konwalski, 2016)

suggests that the surveillance mechanism may involve this process.

In the present study, we found frequent binucleation in Drp1-depleted embryos.

Binucleation of early embryos is relatively common in human IVF procedures, but the

mechanisms of its occurrence and association with the developmental competence of

the embryo are not fully understood (Gomes Paim and FitzHarris, 2020). Reichmann et

al. recently demonstrated that mouse zygotes form two functionally independent

spindles before anaphase, and spatially separate parental genomes during the first

cleavage (Reichmann et al., 2018). Notably, failure to align the two spindles produced

errors, resulting in binuclear formation in the 2-cell embryos. The present study

demonstrates that clustered mitochondria in the central space of zygotes interfere with

the assembly of two spindles, implying a potential contribution of the organelle

misplacement to binucleation in human embryos.

Mitochondrial dynamics and quality control

The present study revealed the response to defective mitochondrial fission at the

cellular level in early embryos, but the response at the organelle level remains to be

elucidated. Mitochondria in oocytes, with hundreds of thousands of mtDNA copies, are

maternally inherited after fertilization. The preservation and apportionment of healthy

mtDNA in oocytes and early embryos is important for the inheritance of normal mtDNA

in the offspring. Recent full-length mtDNA sequencing has identified de novo mtDNA

mutations in oocytes, and their frequency increases with age, implying mtDNA turnover

during meiotic arrest (Abrisch et al., 2020). Oocytes with severe mtDNA mutations can

be eliminated (Fan et al., 2008), although the thresholds at which mtDNA mutations and

the resulting mitochondrial dysfunction result in selection remain unclear. The

autophagic elimination of mitochondria and mitophagy may also function as filters to

segregate and eliminate pathogenic mtDNA mutations at the organelle level (Youle and

van der Bliek, 2012). An attractive model has been proposed for the Drosophila female

germline, as fragmentation of the mitochondrial network facilitates mitophagy and

allows selective elimination of defective mitochondria containing mutant mtDNA

(Lieber et al., 2019). Functional links between mitochondrial fission and removal of

damaged mitochondria by mitophagy have also been documented in mammalian cells,

as Drp1 deficiency causes mitochondrial dysfunction owing to the failure of a

Drp1-dependent mechanism of mitophagy that removes damaged mitochondria within

the cell (Twig et al., 2008). The highly fragmented mitochondrial structure in

mammalian oocytes/zygotes allows mtDNA to be sequestered in small units, which may

be beneficial for eliminating mutation-impaired mitochondria for mitophagy.

Furthermore, symmetric segregation of fragmented mitochondria might randomize

mutant mtDNA among daughter blastomeres during embryonic cleavage to maintain

tissue homeostasis. Elucidating the behavior of mutant mitochondria in early embryos is

clinically meaningful. Mitochondrial replacement therapy (MRT) is a promising

approach to prevent the transmission of mtDNA disorders from mother to child, as

pathogenic mutant mtDNA in mature oocytes or zygotes is replaced with normal

mtDNA by nuclear transfer (Hyslop et al., 2016; Kang et al., 2016). A potential problem

with MRT is that a small fraction of patients' mtDNA is inevitably carried over into the

reconstructed oocytes or zygotes (approximately 1 to 4%, respectively). It has been

noted that these low levels of mutant mtDNA may predominantly expand during

post-implantation embryogenesis, suggesting a rapid shift to one mtDNA haplotype

during early development (Lee et al., 2012). Therefore, it is important to investigate the

role of mitochondrial dynamics in the regulation of mtDNA mutations in the

preimplantation embryo, and whether mitochondrial fission contributes to reducing the

mutational load.

Materials and methods

Preparation of zygotes

Zygotes were collected from the oviducts of 8- to 12-week-old BDF1 female mice

mated with 3- to 6-month-old BDF1 male mice. Females were superovulated by

intraperitoneal injection of 7.5 IU of equine chorionic gonadotropin (PMSG; ASKA

Pharmaceutical, Tokyo, Japan) followed by injection of human chorionic gonadotropin

(hCG; ASKA Pharmaceutical) 48 h later. Female mice were euthanized 20-24 h later

and fertilized zygotes were retrieved in Hepes-buffered KSOM (H-KSOM) medium

under paraffin oil, at 37°C in a humidified atmosphere containing 5% CO_2. All mice

were housed in a pathogen-free environment in filter-top cages and fed a standard diet.

Animal experiments were conducted according to the guidelines of the Animal Care and

Use Committee of Okayama University.

For experiments in Fig. 3C, Drp1$^{loxP/loxP}$ mice (Wakabayashi et al., 2009) were

crossed with transgenic mice that carried Gdf-9 promoter–mediated Cre recombinase.

After multiple rounds of crossing, homozygous Drp1^{cKO} female mice lacking Drp1 in

oocytes (Drp1$^{loxP/loxP}$; Gdf9-Cre) or Drp1$^{Δ/Δ}$ oocytes were obtained. Mice that do not

carry the Cre transgene are referred to as Drp1$^{loxP/loxP}$ and were used as controls. For

collection of ovulated MII oocytes, mice were superovulated by sequential

intraperitoneal injections of 8-IU PMSG and 8-IU hCG at an interval of 48 hours, and

the mice were culled 14 to 16 hours after hCG injection to collect oocytes from oviducts.

Animal experimentation was approved by Monash University Animal Experimentation

Ethics Committee and was performed in accordance with Australian National Health

and Medical Research Council Guidelines on Ethics in Animal Experimentation.

Plasmids

EGFP, mCherry, and DsRed were cloned into pcDNA6 vector between the XhoI and

XbaI sites (pcDNA6/Myc-His B; Invitrogen, Waltham, MA) with a poly(A)-tail of 84

nucleotides (Wakai et al., 2014). The gene sequence encoding the mitochondrial

targeting sequence of Cox VIII was amplified by PCR and ligated to the EGFP and

DsRed-bearing pcDNA6 (mt-GFP and mt-DsRed). The gene sequences encoding the

ER-targeting sequence of calreticulin and the KDEL ER retention sequence were cloned

into mCherry (ER-mCherry)-bearing pcDNA6. For visualization of DNA and

microtubule growth, gene sequences encoding histones H2B and EB3 were ligated to

the pcDNA6 vector in frame with mCherry (H2B-mCherry) and EGFP (EB3-GFP),

respectively. The gene sequences encoding the amino-terminal 33 residues of

endothelial nitric-oxide synthase as a Golgi targeting sequence (Addgene, 14873) and

peroximal targeting signal 1 (Addgene, 54520) were cloned into mCherry-bearing

pcDNA6 (Golgi-mCherry and PEX-mCherry). The gene sequences encoding

N-terminally mCherry tagged Drp1 (Addgene, 49152) were inserted between the EcoRI

and PstI sites of the pCDNA vector (mCherry-Drp1). ATP biosensor, AT1.03 and

AT1.03RK in the pcDNA3.1 vector were kindly provided by Dr H. Imamura (Kyoto

University). Capped mRNA was synthesized with T7 polymerase (RiboMAX

Large-Scale RNA Production, Promega, Madison, WI) according to manufacturer

instructions. mRNAs were delivered into zygotes using a piezo-driven micropipette unit

(Prime Tech, Tsuchiura, Japan). For microinjection, mRNA solution was loaded into

glass micropipettes at the concentration of 200 ng/μl (mt-GFP, EB3-GFP, mt-DsRed,

H2B-mCherry, ER-mCherry, Golgi-mCherry, and PEX-mCherry), 500 ng/μl

(mCheery-Drp1) or 1000 ng/μl (AT1.03 and AT1.03RK). The volumes injected

typically ranged from 2 to 10 pl, which is 1–5% of that of the oocytes. Manipulation

was carried out in H-KSOM medium containing 5 mg/mL cytochalasin B (Sigma, St.

Louis, MO) to increase post-microinjection survival.

Trim-Away

Trim-Away was performed as previously described with some modifications. In

brief, pCMV6-Entry vector harboring Trim-21 mouse ORF Clone (Origene, Rockville,

MD) was linearized by FseI, and mRNA was synthesized using T7 mMessage

mMachine and poly-A tailed with Poly(A) tailing kit (Thermo Fisher Scientific,

Waltham, MA). Mouse monoclonal anti-Drp1 antibody (BD Biosciences, Franklin

Lakes, NJ) or rabbit monoclonal anti-Myo19 antibody (Abcam, Cambridge, UK) was

concentrated using Amicon Ultra-0.5 100 KDa centrifugal filter devices (Millipore,

Bedford, MA) and the buffer was replaced with PBS containing 0.03% NP40 (Nakarai,

Kyoto, Japan). Pronuclear stage zygotes were injected with Trim21 mRNA (1000

ng/mL) and subsequently anti-Drp1 or anti-Myo19 antibodies. For the control

experiment, normal mouse immunoglobulin G (IgG) antibody (Santa Cruz

Biotechnology, Dallas, TX) and rabbit IgG (Cell Signaling Technology, Danvers, MA)

was used instead of an anti-Drp1 antibody and anti-Myo19 antibodies, respectively.

Western blotting

Whole cell lysates from zygotes/embryos were prepared by adding a 2× sample

buffer. Proteins were separated by SDS-PAGE and transferred to PVDF membranes

(Millipore). The membranes were then blocked and probed with anti-Drp1 (1:1000) or

anti-Myo19 (1:1000) antibodies for 1 h at room temperature. Goat anti-mouse antibody

conjugated to horseradish peroxidase (HRP) was used as a secondary antibody (1:2000)

for chemiluminescence detection (Clarity Western ECL Substrates, Bio-Rad, Hercules,

CA) according to the manufacturer's instructions. The signal was digitally captured

using a ChemiDoc XRS+ imaging system (Bio-Rad). The same membranes were

stripped at 50°C for 30 min (62.5 mM Tris, 2% SDS, and 100 mM 2-beta

mercaptoethanol) and re-probed with anti-β-actin monoclonal antibody (1:2000, Cell

Signaling Technology) as the internal control.

mtDNA copy number

The zona pellucida of 2-cell embryos was removed with H-KSOM medium

containing 0.1% protease (Sigma, St. Louis, MO). Individual blastomeres from the

zona-free embryos were obtained by repeatedly pipetting in Ca^{2+}-free and Ma^{2+}-free

H-KOSM medium. After washing in PBS, each blastomere was transferred into 15 μL

of lysis buffer (50 mM Tris-HCl, pH 8.5, with 0.5% Tween 20 and 100 μg proteinase K,

at 55°C, followed by heat inactivation at 95°C for 10 min. The plasmid standard was

constructed by cloning a 194-bp fragment of the NADH-ubiquinone oxidoreductase

chain 1. The standard stock was serially diluted to prepare the standard curve (10^3 to

10^7). Real-time fluorescence-monitored quantitative PCR using the primer pair

5′-CCTATC ACCCTTGCCA-3′ and 5′-GAGGCTGTTGCTTGTG-3′ was performed on

a LightCycler 96 System (Roche, Basel, Switzerland). Each reaction of 20 μL consisted

of 10 μL of SYBR Green I (Roche), 0.5 μM each of the forward and reverse primers,

and 5 μL of diluted DNA template. The cycling conditions were as follows: initial

denaturation at 95°C for 5 min followed by 45 cycles of denaturation at 95°C for 10 sec,

annealing at 50°C for 20 sec, and elongation at 72°C for 8 sec. Standard curves were

created for each run, and the sample copy number was determined from the equation

relating the Ct value against copy number for the corresponding standard curve.

Immunofluorescence

Zygotes/embryos were fixed and permeabilized with 2% paraformaldehyde in

phosphate-buffered saline (PBS) containing 0.1% Triton X-100 for 40 min at room

temperature. After washing with PBS supplemented with 1% bovine serum albumin

(PBS–BSA), the zygotes/embryos were incubated overnight at 4°C with mouse

anti-α-tubulin (1:300, Sigma), rabbit monoclonal anti-Myo19 antibody (1:300, Abcam),

rabbit anti-pericentrin (1:500, Abcam) or mouse monoclonal Anti-γH2A.X (1:500,

Abcam) antibodies in PBS–BSA. Then zygotes/embryos were washed with PBS–BSA

and incubated with Alexa Fluor488-conjugated goat anti-rabbit IgG (1:300) for 1 h at

room temperature. DNA was stained with DAPI. Samples were placed on the medium

under mineral oil in glass bottom dishes (Matsunami, Osaka, Japan) and examined

using a laser-scanning confocal microscope (FV1200, Olympus, Tokyo, Japan) outfitted

with a 63 × 1.4 NA oil immersion objective lens.

Mitochondrial and ROS staining

Mitochondrial distribution was analyzed by staining with 1 μM MitoTracker Red

CM-H2XRos (Thermo Fisher Scientific) in H-KSOM medium for 30 min at 37°C.

Intracellular ROS levels were analyzed using 5 μM H2DCFDA (Thermo Fisher

Scientific) in H-KSOM medium for 15 min at 37°C. The embryos were placed in drops

of H-KSOM medium under mineral oil in glass bottom dishes (Matsunami), and

fluorescence images of MitoTracker Red CMXros and H2DCFDA were obtained using

a laser-scanning confocal microscope (FV1200) fitted with a 63 × 1.4 NA

oil-immersion objective lens.

Electron microscopy

Zygotes/embryos were fixed in 2% glutaraldehyde in a 0.1 M phosphate buffer (pH

7.4) for 1 h and stored at 4°C until processed. After post-fixation in 2% osmium

tetroxide at 4°C, the specimens were dehydrated with a graded series of ethanol and

were embedded in epoxy resin Quetol-812. Semi-thin sections were cut for light

microscopy and stained with toluidine blue for further sectioning of areas. Thereafter,

ultra-thin sections were cut and stained with uranyl acetate and lead citrate and

examined by transmission electron microscopy (TEM) (JEM 1200EX, JEOL) at 80 kV.

Fluorescence resonance energy transfer and Ca^{2+} imaging

ATeam, a fluorescence resonance energy transfer-based ATP indicator, has been

used successfully to measure cellular ATP levels in live somatic cells (Imamura et al.,

2009). To estimate the relative changes in ATP levels, the emission ratio of AT1.03 and

AT1.03RK (YFP/CFP) was imaged using a CFP excitation filter, dichroic beam splitter,

and CFP and YFP emission filters (Chroma Technology, Rockingham, VT; ET436/20x,

89007bs, ET480/40m and ET535/30m). To measure cytoplasmic Ca^{2+}, embryos were

incubated with 1.25 μM Fluo-4 (Thermo Fisher Scientific) supplemented with 0.02%

pluronic acid (Thermo Fisher Scientific) for 20 min at room temperature. Fluo-4 was

excited with 480 nm wavelengths every 20 seconds and emitted light was collected at

wavelengths greater than 510 nm. To estimate Ca^{2+} levels in the ER, Ca^{2+} rise was

analyzed following the addition of 10 μM thapsigargin (Sigma) in a Ca^{2+}-free medium.

For the monitoring of ATP and Ca^{2+} levels, embryos were attached to glass-bottomed

dishes (Matsunami) and placed on the stage of an inverted microscope. . The

fluorescence images were obtained by a scientific CMOS (sCMOS) camera

(Hamamatsu Photonics, Hamamatsu), and the rotation of excitation and emission filter

wheels was controlled using the MAC5000 filter wheel/shutter control box (Ludl

Electronic Products Ltd, Hawthorne, NY) and HCImage software.

Live cell confocal imaging

Spinning disk images of mitochondria (mt-GFP and mt-DsRed), chromosomes

(H2B-mCherry), microtubules (EB3-GFP), ER (ER-mCherry), Golgi (Golgi-mCherry)

and peroxisome (PEX-mCherry) and were obtained using a confocal scanner unit

(CellVoyager CV1000, Yokogawa, Tokyo, Japan) equipped with a C-Apochromat ×40,

1.2 NA water immersion objective. Images were typically acquired every 10-20 mins on

10 planes covering a 35-40 μm range (every 2.5-3.3 μm). Laser wavelengths of 488 and

561 nm were used for the excitation of EGFP and mCherry (DsRed), respectively (not

exceeding 0.2% laser intensity to minimize light-induced phototoxic stress).

Image analysis

Fiji (Schindelin et al., 2012) was used for quantification of fluorescence images.

For quantification of mitochondria at nuclear or spindle periphery in Figure 1B and

Figure-1 figure supplement 1C, the fluorescence images of nucleus (H2B-mCherry) of

spindle (EB3-GFP) were applied with a Gaussian filter (radius: 2 pixels) to remove

noise and segmented by intensity thresholding. The masked binary images were dilated

with 10 pixels width and the nucleus/spindle periphery was segmented by the

subtraction of the original image from the dilated image. The fluorescence intensity

ratio of mitochondrial layers at the periphery to that of outside periphery was measured

to estimate mitochondrial accumulation around the nucleus/spindle.

For quantitation of mitochondrial symmetry at the anaphase in Figure 3B, the

cleavage plane of dividing blastomere was manually outlined with respect to the middle

of the anaphase chromosomes. Total mito-GFP intensity in each blastomere was

measured, and the greater value was divided by the smaller value for the symmetry

index. To analyze organelle symmetry at the telophase in Figure 3F, total pixel intensity

in the channel corresponding to mito-GFP, ER-mCherry, Golgi-mCherry and

PEX-mCherry in each blastomere was measured. The greater value was divided by the

smaller value; a ratio of 1 indicates symmetry, whereas higher values indicate

asymmetry.

Uniformity analysis of mitochondrial distribution around spindles in y in Figure

3-figure supplement 1B was performed based on the angular distributions of

mitochondria (mt-DsRed) and microtubules (EB3-GFP) using "QuantEv Icy" plugin

(Pecot et al., 2018). The fluorescence images were applied with a Gaussian filter to

reduce noise, and xy coordinates were converted to polar coordinates. The fluorescence

intensity profiles for mt-DsRed relative to the centroid of the EB3-GFP signals was

divided into sectors each 60 degree based on the long axis of the spindle. The standard

deviation of averaged mitochondrial intensity per $60°$ sector were calculated, higher

values were considered to indicate increased non-uniformity.

For tracking nuclei in zygotes in Figure 5C and 5D, XY coordinates of the centroid

of male and female chromosomes (H2B-macherry) relative to the center of the zygote

were obtained using "Manual tracking" plugin. Only the zygotes with nuclei that

remained in the focal plane were analyzed.

Statistical analysis

All statistical analyses were performed using GraphPad Prism software. Values from

three or more experiments performed on different batches of cells were analyzed by

Statistical tests and p-values are indicated in figures and figure legends.

Resource availability

All unique/stable reagents and plasmids generated in this study are available from

the lead contact with a completed Materials Transfer Agreement. Further information

and requests for resources and reagents should be directed to and will be fulfilled by the

lead contact, Takuya Wakai (t2wakai@okayama-u.ac.jp).

Acknowledgements

We thank Dr. H. Imamura (Kyoto University) for sharing AT1.03 and AT1.03RK

plasmids. We thank Sayaka Nakato for technical assistance. We thank the Division of

Instrumental Analysis, Okayama University for the experiments using FV1200 and

gratefully thank the staff of Mio Fertility Clinic for technical assistance with

CellVoyager CV1000.

Figure Legends

Figure 1. Redistribution and symmetric partitioning of mitochondria during

embryonic cleavage.

(A) Representative time-lapse images (maximum-intensity Z projection) of

mitochondria and chromosomes in zygotes expressing mitochondrially-targeted

(mt-GFP) and histone H2B-mCherry during the first cleavage division. Time is relative

to the onset of NEBD. Scale bar, 20 μm.

(B) Quantification of mitochondrial accumulation around the nucleus at different cell

cycle phases by calculating the ratio of the mitochondrial fluorescence intensity at the

nuclear periphery to that of outside nuclear periphery.

(C) Representative electron microscopy (EM) images (n = 5) showing the accumulation

of mitochondria around the mitotic spindle. Higher-magnification images (31 sections)

of the representative boxed areas in the lower left and upper right, respectively.

Mitochondria (purple), endoplasmic reticulum (pink), and actin filaments

(yellow-green) are pseudo-colored in the lower right panel. Scale bar left: 10 μm;

Lower left and upper right: 1 μm.

(D) Interphase or metaphase zygotes and interphase 2-cell embryos were stained with

Phalloidin-iFluor 488, MitoTracker Red CMXros and Hoechest 33342. Magnified

images in the box areas are shown in the right panels. An overall view of each embryo

is shown in Figure 1-figure supplement 1D.

(E) Quantification of the correlation between the localization of the F-actin

(Phalloidin-iFluor 488) and mitochondria (MitoTracker Red CMXros).

(F) Representative immunofluorescence images of Myo19. Interphase (n = 13) and

metaphase (n = 11) zygotes were stained with MitoTracker Red CMXRos. After

fixation and immunostaining with anti-Myo19 antibody and Hoechest 33342,

single-section images crossing the mid-zone of the zygotes were imaged by confocal

microscopy. Scale bar, 20 μm.

(G) Western blot analysis of zygotes overexpressing Trim21 and microinjected with

control IgG or anti-Myo19 antibodies. Lysates of 100 zygotes were extracted 5 h after

microinjection and were probed with antibodies specific to Myo19 and β-actin.

(H) Developmental competence of control and Myo19-depleted embryos. Percentage of

zygotes reaching the indicated developmental stage.

(I) Maximum-intensity Z projection of control and Myo19 Trim-Away embryos

expressing mt-GFP and H2B-mCherry. Scale bar, 20 μm.

(J) Quantitation of mitochondrial mass inheritance after the first cleavage division of

control and Myo19-depleted zygotes. Total mitochondrial fluorescence in each daughter

blastomere of 2-cell embryos was measured, and the smaller value was divided by the

greater value for the inheritance ratio.

Data are represented as mean ± SD and P values calculated using two-tailed paired (B)

or unpaired (E, H and J) Student's t test. $*p < 0.05$, $***p < 0.001$, $****p < 0.0001$; ns,

not significant. Number of embryos are indicated in brackets.

Figure 1-figure supplement 1. Redistribution of mitochondria during the first

cleavage.

(A) Representative immunofluorescence images of microtubules (green), mitochondria

(red) and DNA (blue) during the first cleavage division. Metaphase II eggs were

collected 14 h after hCG. Zygotes were isolated 24 h after hCG and cultured to different

mitotic stages (n = 10–35). Interphase zygotes at the PN stage and 2-cell stage embryos

were fixed 25 and 35 h after hCG, respectively. Mitotic zygotes were fixed at 0.5

(prophase and prometaphase), 1 (metaphase), and 2 h (anaphase and telophase) after the

disappearance of the nuclear envelope was observed. Prior to fixation, zygotes were

stained with MitoTracker Red CMXRos for 0.5 h. After fixation and immunostaining

with anti-α-tubulin and DAPI, single-section images crossing the mid-zone of the

zygotes were imaged by confocal microscopy. Scale bar, 20 μm.

(B) Representative images (maximum-intensity Z projection) of mitochondria

(mt-DsRed) and microtubules (EB3-GFP) in metaphase blastomeres during the first to

third cleavage division. Scale bar, 20 μm.

(C) Quantification of mitochondrial accumulation around the spindle by calculating the

ratio of the mitochondrial fluorescence intensity at the spindle periphery to that of

outside spindle periphery.

(D) Interphase or metaphase zygotes and interphase 2-cell embryos were stained with

Phalloidin-iFluor 488, MitoTracker Red CMXros and Hoechest 33342. Magnified

images in the box areas are shown in Figure 1D.

(E) Confocal images of zygotes treated with DMSO (control) or Latrunculin A (LatA)

for 1 hr just after NEB, fixed and stained for F-actin (Phalloidin-iFluor 488) and DNA

(Hoechest 33342). Magnified views of the cytoplasmic and cortical regions are shown

in the upper right and lower right, respectively. Scale bar, 10 μm.

(F) Zygotes were treated as in (E), then washed out, cultured to develop into 2-cell

embryos and stained with MitoTracker Red CMXros. Maximum-intensity Z projection

of the entire cell for mitochondria were imaged by confocal microscopy.

(G and H) Quantitation of mitochondrial mass inheritance (G) and blastomere size (H)

in control and LatA-treated 2-cell embryos. Total mitochondrial fluorescence and the

area in each daughter blastomere of two-cell embryos was measured, and the ratio was

calculated by dividing the smaller value by the larger value.

(I) Correlation between mitochondrial inheritance ratio and blastomere size ratio.

Data are represented as mean ± SD and P values calculated using two-tailed Student's t

test. *$p < 0.05$, ***$p < 0.001$, ****$p < 0.0001$; ns, not significant. Number of embryos

are indicated in brackets.

Figure 1-Video 1. Symmetric mitochondrial partitioning during the first cleavage.

Representative time-lapse movies of mitochondria (mt-GFP) and chromosomes

(H2B-mCherry) in zygotes. Time is relative to the onset of NEB (hours_minutes).

Z-projection of 8 sections, 5 μm apart; time interval: 20 min.

Figure 2. Depletion of Drp1 induces mitochondrial aggregation.

(A) Western blot analysis of preimplantation embryos at the different stages. Lysates of

30 cells were probed with antibodies specific to Drp1 and β-actin.

(B) Western blot analysis of zygotes overexpressing Trim21 and microinjected with

control IgG or anti-Drp1 antibodies (henceforth referred to simply as control and

Trim-Away, respectively), or zygotes microinjected with Trim21 mRNA, control IgG or

anti-Drp1 alone. Lysates were extracted 5 h after microinjection and immunoblotted for

the indicated proteins.

(C) Representative images of in control (n = 37) and Drp1 Trim-Away (n = 57) embryos

expressing mt-GFP. Scale bar, 20 μm.

(D) A representative image of mitochondria (mt-GFP) in Drp1-depleted embryos (n

=53) expressing exogenous Drp1 (mCh-Drp1).). Note that mCh-Drp1 expression

rescues the mitochondrial aggregation following Drp1 depletion by Trim-Away to the

normal dispersed distribution. Scale bar, 20 μm.

(E) Representative images of mitochondria (mt-GFP) in Drp1$^{fl/fl}$ control (n =10) and

Drp1$^{\Delta/\Delta}$ (n = 8) parthenotes. Mature eggs were collected form 4- to 5-week-old Drp1$^{fl/fl}$

and Drp1^{cKO} mice. Scale bar, 20 μm. After microinjection of mt-GFP mRNA, oocytes

were exposed to 10 mM SrCl$_2$.

(F) Representative EM images of control (n = 5, 17 sections) and Drp1-depleted (n = 6,

20 sections) 2-cell embryos. Right panels show higher-magnification images. Scale bar

left: 4 μm; right: 1 μm.

(G) Major axis of mitochondria in control and Drp1-depleted embryos was measured

from EM images (F).

(H) Levels of ATP in control and Drp1-depleted embryos were estimated using the

emission ratio of AT1.03 (YFP/CFP). The emission ratio of AT1.03RK (YFP/CFP),

which is unable to bind ATP, is shown as a negative control.

Data are represented as mean ± SD and P values calculated using two-tailed Student's t

test with Welch's correction. ****p < 0.0001. Number of embryos are indicated in

brackets.

Figure 2-figure supplement 1. Mitochondrial aggregation disturbs subcellular

organelle compartments.

(A-C) Sub-cellular distributions of mitochondria with ER (A), Golgi (B), and

peroxisome (C). Representative images of control and Drp1-depleted embryos

co-expressing mt-GFP and ER-mCherry (control, n = 11; Trim-Away, n = 15),

Golgi-mCherry (control, n = 10; Trim-Away, n = 16) or PEX-mCherry (control, n = 19;

Trim-Away, n = 20). Magnified images in the box areas are shown in the right panels.

Scale bar, 20 μm.

Figure 3. Drp1 depletion causes asymmetric mitochondrial inheritance and

functions.

(A) Representative time-lapse images (maximum-intensity Z projection) of

mitochondria and chromosomes in anti-Drp1 microinjected Trim-Away zygotes

expressing mt-GFP and H2B-mCherry. Time is relative to the onset of NEBD. Dashed

lines indicate cleavage plane. Scale bar, 20 μm.

(B) Quantitation of mitochondrial asymmetry at anaphase in control and Drp1-depleted

zygotes. Total mitochondrial fluorescence in each dividing blastomere was measured,

and the greater value was divided by the smaller value for the symmetry index.

(C) mtDNA inheritance ratio after the first cleavage division of control and

Drp1-depleted zygotes. mtDNA copy number in each daughter blastomere of 2-cell

embryo was measured, and the ratio was calculated by dividing the greater value by the

smaller value.

(D) ATP inheritance ratio between two blastomeres was estimated from comparison of

emission ratio of AT1.03 (YFP/CFP) in the daughter blastomere A/B, where A is the

daughter blastomere with more mitochondria as confirmed by MitoTracker staining (see

Figure S3D).

(E) Correlation between mtDNA and ATP inheritance ratio between 2-cell blastomeres.

(F) Quantitation of organelle asymmetry at telophase was analyzed from time-lapse

images of embryos co-expressing mt-GFP with mCherry-tagged ER, Glogi, or

peroxisome. Total mitochondrial fluorescence in each blastomere was measured, and

the greater value was divided by the smaller value.

(G) Organelle symmetry ratios, as shown in (F), are plotted against the mitochondrial

symmetry ratio for each Drp1-depleted embryo. Note that mitochondrial and ER

inheritance ratios are highly correlated ($R^2 = 0.6734$).

(H-J) Endoplasmic reticulum Ca^{2+} content in each blastomere of control (H) and

Drp1-depleted (I) embryos was estimated following the addition of 10 μM thapsigargin

in Ca^{2+} -free medium, and comparisons of the Fluo-4 fluorescent peaks are shown in a

graph (J).

Data are represented as mean ± SD and P values calculated using unpaired two-tailed t

tests with Welch's correction (B, C, D and F) or one-way ANOVA with Turkey's

multiple comparison test (J). *p < 0.05, ***p < 0.001, ****p < 0.0001; ns, not

significant. Number of embryos are indicated in brackets.

Figure 3-figure supplement 1. Drp1 depletion increases asymmetry in

mitochondrial inheritance between blastomeres.

(A) Representative time-lapse images (maximum-intensity Z projection) of

mitochondria and microtubules in control and anti-Drp1 microinjected Trim-Away

zygotes expressing EB3-GFP and mt-DsRed. Time is relative to the onset of anaphase.

Scale bar, 20 μm.

(B) Uniformity analysis of mitochondrial distribution around the spindle.

Representative angler distributions of mitochondria (red trace) and microtubules (green

trace) at the metaphase of control and Drp1 Trim-Away zygotes, as shown in (A). Based

on the angular distributions, standard deviation of averaged mitochondrial fluorescence

intensity per 60° sector were plotted in the graph to the right, higher values indicate

increased non-uniformity.

(C) Representative ratiometric images of ATeam AT1.03 and AT1.03RK fluorescence

(YFP/CFP) in control and Drp1-depleted 2-cell embryos. Mitochondrial distribution in

the same embryos is shown in the right panel. Arrowheads represent marked

mitochondrial aggregation in Drp1 Trim-Away embryos and blastomere A/B, where A is

the daughter blastomere with more mitochondria as indicated by MitoTracker staining.

(D) Representative time-lapse images (maximum-intensity Z projection) of

mitochondria and ER in control and Drp1 Trim-Away zygotes expressing mt-GFP and

ER-mCherry. Arrowheads represent regions of marked mitochondrial aggregation in

Drp1 Trim-Away embryos. Time is relative to the onset of NEBD. Scale bar, 20 μm.

Figure 3-Video 1. Asymmetric mitochondrial partitioning during the first cleavage of

Drp1-depleted zygotes, related to Fig. 2. Representative time-lapse movies of

mitochondria (mt-GFP) and chromosomes (H2B-mCherry) in Drp1 Trim-Away (right)

zygotes. Time is relative to the onset of NEB (hours_minutes). Z-projection of 8

sections, 5 μm apart; time interval: 20 min.

Figure 3-Video 2. Asymmetric partitioning of the ER confined to mitochondrial

aggregation regions in Drp1-depleted zygotes, related to Fig. 4 and Fig. S4.

Representative time-lapse movies of mitochondria (mt-GFP) and ER (ER-mCherry) in

control and Drp1 Trim-Away zygotes. Each of the movies in the upper row is displayed

in the lower row by LUT. Arrowheads represent regions of marked mitochondrial

aggregation. Time is relative to the onset of NEB (hours_minutes). Z-projection of 8

sections, 5 μm apart; time interval: 15 min.

Figure 4. Drp1 depletion causes early embryonic arrest.

(A) Developmental competence of control and Drp1 Trim-Away zygotes. Following

Drp1 depletion by Trim-Away (co-injection of Trim21 mRNA and anti-Drp1 antibody)

or the control experiments in Fig. 2B, zygotes were cultured to the indicated stages.

(B) Representative images of embryos microinjected Trim21 mRNA with control IgG

or anti-Drp1 were cultured. Some embryos at E1.5 have three or four blastomeres

(arrowheads). Scale bar, 20 μm.

(C) Developmental competence of Drp1-depleted embryos is rescued by microinjection

of mCherry-Drp1 (mCh-Drp1) mRNA. Percentage of zygotes reaching the indicated

developmental stage. Note that the 2-cell stage arrest in control condition with RNA

buffer (water) upon Trim-Away depletion was reversed by microinjecting mCh-Drp1

mRNA.

(D) Western blot analysis of control and Drp1 Trim-Away zygotes that were

microinjected with water or mCh-Drp1 mRNA upon Trim-Away depletion. Lysates of

20 zygotes were extracted 5 h after microinjection and were probed with antibodies

specific to Drp1 and β-actin.

Data are represented as mean ± SD and P values calculated using one-way ANOVA

with Turkey's multiple comparison test. $*p < 0.05$, $***p < 0.001$, $****p < 0.0001$; ns,

not significant. Number of embryos are indicated in brackets.

Figure 4-figure supplement 1. Drp1 depletion induces the ectopic accumulation of

ROS and DNA damage.

(A) Representative immunofluorescence images of phosphorylated (Ser10) histone H3

(green), known to localize in the heterochromatin region in G2-phase cells, and DAPI

(blue) in control (n = 18) and Drp1 Trim-Away (n = 20) embryos. Scale bars, 20 μm.

Embryos were fixed at 46 h post hCG (corresponding to the G2 stage of 2-cell

embryos).

(B) Representative immunofluorescence images (maximum-intensity Z projection) of

γH2AX (green), MitoTracker Red CMXros (red) and DAPI (blue) in control and Drp1

Trim-Away embryos.

(C) Comparisons of the number of γH2AX foci in control and Drp1 Trim-Away

embryos.

(D) Representative images of H2DCFDA, a probe for intracellular reactive oxygen

species (ROS) and mitochondria (MitoTracker Red CMXros) in control (n = 29) and

Drp1 Trim-Away embryos (n = 35). Arrowheads represent regions of marked

mitochondrial aggregation in Drp1 Trim-Away embryos. Magnified images in the box

areas are shown in the right panels. Scale bar, 20 μm.

Data are represented as mean ± SD and P values calculated using unpaired two-tailed t

tests with Welch's correction. ***$p < 0.001$.

Figure 5. Misplaced mitochondria cause chromosome segregation defects leading

to binucleation.

(A) Representative time-lapse images (maximum-intensity Z projection) of

mitochondria and chromosomes in Drp1 Trim-Away zygotes expressing mt-GFP and

H2B-mCherry. Time is relative to the onset of NEB. Scale bar, 20 μm.

(B) Representative immunofluorescence images of control (n = 10) and Drp1-depleted

(n = 31) zygotes at metaphase of the first cleavage. Pericentrin (red), microtubules

(green), and DNA (blue) were imaged by confocal microscopy. White arrowheads

indicate spindle poles. Scale bars, 10 μm.

(C) The position of chromosomes relative to the center of zygotes. The centroids of two

sets of chromosomes were tracked, and only the displacements projected in the

direction pointing from initial (NEB) to the final position (just before anaphase onset)

were indicated. Data for three representative zygotes were presented and each color

represents a zygote. Images of mitochondria at the final position automatically had their

thresholds stipulated using Fiji (Moments algorithm) and merged with images filled

inside the outline with the color corresponding to the chromosomes.

(D) Quantification of the distance of chromosomes to cell center just before anaphase

onset.

(E) Cell division abnormalities at the first cleavage. Note that binulceated blastomeres

in Drp1-depleted embryos result from various chromosome segregation defects (see

also Figure 5-figure supplement 1C and 1D).

(F) Schematic summary of results highlighting significant findings in this study. Drp1

mediate mitochondria fragmentation, ensuring symmetric mitochondrial inheritance

between daughter blastomeres. In normal embryos, fragmented mitochondria (green)

and the ER (pink) uniformly surround two spindles, allowing parental spindles (blue

and red) to assemble at the center of metaphase zygotes. Drp1 depletion increases

mitochondrial aggregation and asymmetric distribution. Misplaced mitochondria disturb

the organelle positioning during the first cleavage division leading to asymmetric

organelle inheritance and chromosome missegregation, which in turn results in early

embryonic arrest.

Data are represented as mean ± SD and P values calculated using unpaired two-tailed t

tests with Welch's correction. ***$p < 0.001$.

1004

Figure 5-figure supplement 1. Marked mitochondrial centration causes

chromosome segregation defects.

(A) A representative image of mitochondria (red), microtubules (green) and DNA (blue)

in Drp1-depleted metaphase zygotes (n = 22). Prior to fixation, zygotes were stained

with MitoTracker Red CMXRos for 0.5 h. Scale bars; 10 μm. The right panel represents

a schematic diagram showing mitochondria located in the center of two spindles based

on the left panel.

(B) Representative EM images (n = 6) showing mitochondria clustered in zygote center

and metaphase chromosomes, as surrounded by dashed lines, located in the vicinity of

the mitochondria. Representative higher-magnification images (31 sections) of the

boxed areas. Scale bar left: 10 μm; middle and right: 1 μm.

(C and D) Examples of two typical cleavage defects leading to binucleation in Drp1

Trim-Away zygotes. Time-lapse images (maximum-intensity Z projection) of

mitochondria (mtGFP) and chromosomes (H2B-mCherry) during the first cleavage

division show the mitochondrial aggregation sits between two sets of chromosomes

which transiently segregated into three (C) or four (D) blastomeres before forming

binucleated (arrowheads) 2-cell embryos. Time is relative to the onset of NEBD. Scale

bar, 20 μm.

1023

1024   **Figure 5-Video 1.** Binuclear formation in Drp1-depleted embryos, related to Fig. 5.

1025   Representative time-lapse movies of mitochondria (mt-GFP) and chromosomes

1026   (H2B-mCherry) in Drp1 Trim-Away zygotes. Time is relative to the onset of NEB

1027   (hours_minutes). Z-projection of 10 sections, 3.5 µm apart; time interval: 15 min.

1028

Figure 5-Video 2. Examples of two typical cleavage defects leading to binucleation in

Drp1 Trim-Away zygotes, related to Fig. 5 and Fig. S5. Time-lapse movies of

mitochondria (mt-GFP) and chromosomes (H2B-mCherry) in the zygote showed that

multiple cleavage furrows formed between the different sets of chromosomes leading to

transient formation of three or four blastomeres, which failed to complete cytokinesis

and collapsed back to form binucleated 2-cell embryos. Time is relative to the onset of

NEB (hours_minutes). Z-projection of 10 sections, 3.5 µm apart; time interval: 10 min.

1036

1037

1038

References

Abrisch, R.G., S.C. Gumbin, B.T. Wisniewski, L.L. Lackner, and G.K. Voeltz. 2020. Fission
and fusion machineries converge at ER contact sites to regulate mitochondrial
morphology. *J Cell Biol*. 219.

Adhikari, D., I.W. Lee, U. Al-Zubaidi, J. Liu, Q.H. Zhang, W.S. Yuen, L. He, Y. Winstanley, H.
Sesaki, J.R. Mann, R.L. Robker, and J. Carroll. 2022. Depletion of oocyte
dynamin-related protein 1 shows maternal-effect abnormalities in embryonic
development. *Sci Adv*. 8:eabl8070.

Ball, E.H., and S.J. Singer. 1982. Mitochondria are associated with microtubules and not
with intermediate filaments in cultured fibroblasts. *Proc Natl Acad Sci U S A*.
79:123-126.

Carlton, J.G., H. Jones, and U.S. Eggert. 2020. Membrane and organelle dynamics during
cell division. *Nat Rev Mol Cell Biol*. 21:151-166.

Chaigne, A., C. Campillo, R. Voituriez, N.S. Gov, C. Sykes, M.H. Verlhac, and M.E. Terret.
2016. F-actin mechanics control spindle centring in the mouse zygote. *Nat Commun*.
7:10253.

Chung, J.Y., J.A. Steen, and T.L. Schwarz. 2016. Phosphorylation-Induced Motor Shedding
Is Required at Mitosis for Proper Distribution and Passive Inheritance of
Mitochondria. *Cell Rep*. 16:2142-2155.

Clift, D., W.A. McEwan, L.I. Labzin, V. Konieczny, B. Mogessie, L.C. James, and M. Schuh.
2017. A Method for the Acute and Rapid Degradation of Endogenous Proteins. *Cell*.
171:1692-1706 e1618.

Dumollard, R., M. Duchen, and J. Carroll. 2007. The role of mitochondrial function in the
oocyte and embryo. *Curr Top Dev Biol*. 77:21-49.

Fan, W., K.G. Waymire, N. Narula, P. Li, C. Rocher, P.E. Coskun, M.A. Vannan, J. Narula,
G.R. Macgregor, and D.C. Wallace. 2008. A mouse model of mitochondrial disease
reveals germline selection against severe mtDNA mutations. *Science*. 319:958-962.

Figueroa-Romero, C., J.A. Iniguez-Lluhi, J. Stadler, C.R. Chang, D. Arnoult, P.J. Keller, Y.
Hong, C. Blackstone, and E.L. Feldman. 2009. SUMOylation of the mitochondrial
fission protein Drp1 occurs at multiple nonconsensus sites within the B domain and
is linked to its activity cycle. *FASEB J*. 23:3917-3927.

Gomes Paim, L.M., and G. FitzHarris. 2020. The impact of embryo binucleation depends

upon its origin. *Reproduction*. 160:V1-V4.

Hyslop, L.A., P. Blakeley, L. Craven, J. Richardson, N.M. Fogarty, E. Fragouli, M. Lamb, S.E.
Wamaitha, N. Prathalingam, Q. Zhang, H. O'Keefe, Y. Takeda, L. Arizzi, S.
Alfarawati, H.A. Tuppen, L. Irving, D. Kalleas, M. Choudhary, D. Wells, A.P.
Murdoch, D.M. Turnbull, K.K. Niakan, and M. Herbert. 2016. Towards clinical
application of pronuclear transfer to prevent mitochondrial DNA disease. *Nature*.
534:383-386.

Imamura, H., K.P. Nhat, H. Togawa, K. Saito, R. Iino, Y. Kato-Yamada, T. Nagai, and H. Noji.
2009. Visualization of ATP levels inside single living cells with fluorescence
resonance energy transfer-based genetically encoded indicators. *Proc Natl Acad Sci*
U S A. 106:15651-15656.

Ishihara, N., M. Nomura, A. Jofuku, H. Kato, S.O. Suzuki, K. Masuda, H. Otera, Y.
Nakanishi, I. Nonaka, Y. Goto, N. Taguchi, H. Morinaga, M. Maeda, R. Takayanagi,
S. Yokota, and K. Mihara. 2009. Mitochondrial fission factor Drp1 is essential for
embryonic development and synapse formation in mice. *Nat Cell Biol*. 11:958-966.

Kang, E., J. Wu, N.M. Gutierrez, A. Koski, R. Tippner-Hedges, K. Agaronyan, A.
Platero-Luengo, P. Martinez-Redondo, H. Ma, Y. Lee, T. Hayama, C. Van Dyken, X.
Wang, S. Luo, R. Ahmed, Y. Li, D. Ji, R. Kayali, C. Cinnioglu, S. Olson, J. Jensen, D.
Battaglia, D. Lee, D. Wu, T. Huang, D.P. Wolf, D. Temiakov, J.C. Belmonte, P. Amato,
and S. Mitalipov. 2016. Mitochondrial replacement in human oocytes carrying
pathogenic mitochondrial DNA mutations. *Nature*. 540:270-275.

Katajisto, P., J. Dohla, C.L. Chaffer, N. Pentinmikko, N. Marjanovic, S. Iqbal, R. Zoncu, W.
Chen, R.A. Weinberg, and D.M. Sabatini. 2015. Stem cells. Asymmetric apportioning
of aged mitochondria between daughter cells is required for stemness. *Science*.
348:340-343.

Koch, A., M. Thiemann, M. Grabenbauer, Y. Yoon, M.A. McNiven, and M. Schrader. 2003.
Dynamin-like protein 1 is involved in peroxisomal fission. *J Biol Chem*.
278:8597-8605.

Kraus, F., K. Roy, T.J. Pucadyil, and M.T. Ryan. 2021. Function and regulation of the
divisome for mitochondrial fission. *Nature*. 590:57-66.

Kumar, R.P., S. Ray, P. Home, B. Saha, B. Bhattacharya, H.M. Wilkins, H. Chavan, A.
Ganguly, J. Milano-Foster, A. Paul, P. Krishnamurthy, R.H. Swerdlow, and S. Paul.
2018. Regulation of energy metabolism during early mammalian development:

TEAD4 controls mitochondrial transcription. *Development*. 145.

Ladstatter, S., and K. Tachibana-Konwalski. 2016. A Surveillance Mechanism Ensures

Repair of DNA Lesions during Zygotic Reprogramming. *Cell*. 167:1774-1787 e1713.

Lee, H.S., H. Ma, R.C. Juanes, M. Tachibana, M. Sparman, J. Woodward, C. Ramsey, J. Xu,

E.J. Kang, P. Amato, G. Mair, R. Steinborn, and S. Mitalipov. 2012. Rapid

mitochondrial DNA segregation in primate preimplantation embryos precedes

somatic and germline bottleneck. *Cell Rep*. 1:506-515.

Lee, I.W., D. Adhikari, and J. Carroll. 2022. Miro1 depletion disrupts spatial distribution of

mitochondria and leads to oocyte maturation defects. *Front Cell Dev Biol*.

10:986454.

Lieber, T., S.P. Jeedigunta, J.M. Palozzi, R. Lehmann, and T.R. Hurd. 2019. Mitochondrial

fragmentation drives selective removal of deleterious mtDNA in the germline.

Nature. 570:380-384.

Lopez-Domenech, G., C. Covill-Cooke, D. Ivankovic, E.F. Halff, D.F. Sheehan, R. Norkett, N.

Birsa, and J.T. Kittler. 2018. Miro proteins coordinate microtubule- and

actin-dependent mitochondrial transport and distribution. *EMBO J*. 37:321-336.

Mahrous, E., Q. Yang, and H.J. Clarke. 2012. Regulation of mitochondrial DNA

accumulation during oocyte growth and meiotic maturation in the mouse.

Reproduction. 144:177-185.

Majstrowicz, K., U. Honnert, P. Nikolaus, V. Schwarz, S.J. Oeding, S.A. Hemkemeyer, and M.

Bahler. 2021. Coordination of mitochondrial and cellular dynamics by the

actin-based motor Myo19. *J Cell Sci*. 134.

Mishra, P., and D.C. Chan. 2014. Mitochondrial dynamics and inheritance during cell

division, development and disease. *Nat Rev Mol Cell Biol*. 15:634-646.

Moore, A.S., S.M. Coscia, C.L. Simpson, F.E. Ortega, E.C. Wait, J.M. Heddleston, J.J.

Nirschl, C.J. Obara, P. Guedes-Dias, C.A. Boecker, T.L. Chew, J.A. Theriot, J.

Lippincott-Schwartz, and E.L.F. Holzbaur. 2021. Actin cables and comet tails

organize mitochondrial networks in mitosis. *Nature*. 591:659-664.

Pecot, T., L. Zengzhen, J. Boulanger, J. Salamero, and C. Kervrann. 2018. A quantitative

approach for analyzing the spatio-temporal distribution of 3D intracellular events in

fluorescence microscopy. *Elife*. 7.

Poulton, J., M.R. Chiaratti, F.V. Meirelles, S. Kennedy, D. Wells, and I.J. Holt. 2010.

Transmission of mitochondrial DNA diseases and ways to prevent them. *PLoS Genet*.

6.

Qian, W., S. Choi, G.A. Gibson, S.C. Watkins, C.J. Bakkenist, and B. Van Houten. 2012.
Mitochondrial hyperfusion induced by loss of the fission protein Drp1 causes
ATM-dependent G2/M arrest and aneuploidy through DNA replication stress. *J Cell*
Sci. 125:5745-5757.

Reichmann, J., B. Nijmeijer, M.J. Hossain, M. Eguren, I. Schneider, A.Z. Politi, M.J. Roberti,
L. Hufnagel, T. Hiiragi, and J. Ellenberg. 2018. Dual-spindle formation in zygotes
keeps parental genomes apart in early mammalian embryos. *Science*. 361:189-193.

Rizzuto, R., D. De Stefani, A. Raffaello, and C. Mammucari. 2012. Mitochondria as sensors
and regulators of calcium signalling. *Nat Rev Mol Cell Biol*. 13:566-578.

Rohn, J.L., J.V. Patel, B. Neumann, J. Bulkescher, N. McHedlishvili, R.C. McMullan, O.A.
Quintero, J. Ellenberg, and B. Baum. 2014. Myo19 ensures symmetric partitioning
of mitochondria and coupling of mitochondrial segregation to cell division. *Curr Biol*.
24:2598-2605.

Scheffler, K., J. Uraji, I. Jentoft, T. Cavazza, E. Monnich, B. Mogessie, and M. Schuh. 2021.
Two mechanisms drive pronuclear migration in mouse zygotes. *Nat Commun*.
12:841.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch,
C. Rueden, S. Saalfeld, B. Schmid, J.Y. Tinevez, D.J. White, V. Hartenstein, K.
Eliceiri, P. Tomancak, and A. Cardona. 2012. Fiji: an open-source platform for
biological-image analysis. *Nat Methods*. 9:676-682.

Twig, G., A. Elorza, A.J. Molina, H. Mohamed, J.D. Wikstrom, G. Walzer, L. Stiles, S.E.
Haigh, S. Katz, G. Las, J. Alroy, M. Wu, B.F. Py, J. Yuan, J.T. Deeney, B.E. Corkey,
and O.S. Shirihai. 2008. Fission and selective fusion govern mitochondrial
segregation and elimination by autophagy. *EMBO J*. 27:433-446.

Udagawa, O., T. Ishihara, M. Maeda, Y. Matsunaga, S. Tsukamoto, N. Kawano, K. Miyado,
H. Shitara, S. Yokota, M. Nomura, K. Mihara, N. Mizushima, and N. Ishihara. 2014.
Mitochondrial fission factor Drp1 maintains oocyte quality via dynamic
rearrangement of multiple organelles. *Curr Biol*. 24:2451-2458.

Wakabayashi, J., Z. Zhang, N. Wakabayashi, Y. Tamura, M. Fukaya, T.W. Kensler, M. Iijima,
and H. Sesaki. 2009. The dynamin-related GTPase Drp1 is required for embryonic
and brain development in mice. *J Cell Biol*. 186:805-816.

Wakai, T., Y. Harada, K. Miyado, and T. Kono. 2014. Mitochondrial dynamics controlled by

mitofusins define organelle positioning and movement during mouse oocyte

maturation. *Mol Hum Reprod.* 20:1090-1100.

Westermann, B. 2010. Mitochondrial fusion and fission in cell life and death. *Nat Rev Mol*

Cell Biol. 11:872-884.

Yamano, K., and R.J. Youle. 2011. Coupling mitochondrial and cell division. *Nat Cell Biol.*

13:1026-1027.

Youle, R.J., and A.M. van der Bliek. 2012. Mitochondrial fission, fusion, and stress. *Science.*

337:1062-1065.

1178

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
genetic reagent (*Mus musculus*)	DBA/2 (♂)	Charles River Laboratories	RRID:MGI:2159768	
genetic reagent (*Mus musculus*)	Mouse: C57BL/6N (♀)	Charles River Laboratories	RRID:MGI:2159965	
genetic reagent (*Mus musculus*)	Mouse: B6D2F1 mice	Department of Animal Resources, Okayama University	RRID:MGI:5649818	
genetic reagent (*Mus musculus*)	*Drp1$^{fl/fl}$* mice	PMID: 19752021		
genetic reagent (*Mus musculus*)	*Drp1cKO* mice	PMID: 35704569		
antibody	Mouse monoclonal anti-Drp1	BD Biosciences	Cat# 611739; RRID:AB_399215	WB(1:1000) Trim-Away experiment (1000 ng/mL)
antibody	Mouse monoclonal anti-β-actin	Cell Signalling Technology	Cat# 3700; RRID:AB_2242334	WB(1:2000)
antibody	Rabbit polyclonal anti-Myo19	Abcam	Cat# ab174286; RRID:	IF(1:300) WB(1:1000)
antibody	Rabbit polyclonal anti-Pericentrin	Abcam	Cat# ab4448; RRID:AB_304461	IF(1:500)
antibody	Mouse monoclonal Anti-γH2A.X	Abcam	Cat# ab22551; RRID:AB_447150	IF(1:500)

antibody	HRP anti-Mouse	Cell Signaling Technology	Cat# 7076; RRID:AB_330924	WB(1:2000)
antibody	HRP anti-Rabbit	Cell Signaling Technology	Cat# 7074; RRID:AB_2099233	WB(1:2000)
antibody	Alexa Fluor 488 anti-Mouse IgG	Thermo Fisher Scientific	Cat# A-11001; RRID:AB_2534069	IF(1:300)
antibody	Alexa Fluor 568 anti-Rabbit IgG	Thermo Fisher Scientific	Cat# A-11011; RRID:AB_143157	IF(1:500)
antibody	Normal mouse IgG	Santa Cruz Biotechnology	Cat# sc-2025; RRID:AB_737182	Trim-Away experiment (1000 ng/mL)
antibody	Normal Rabbit IgG	Cell Signaling Technology	Cat# 2729; RRID:AB_1031062	Trim-Away experiment (1000 ng/mL)
recombinant DNA reagent	pcDNA6-Mt-GFP(plasmid)	PMID: 25113836		
sequence-based reagent	pcDNA6-Mt-DsRed (plasmid)	PMID: 25113836		
recombinant DNA reagent	pcDNA6-ER-mCherry (plasmid)	PMID: 25113836		
recombinant DNA reagent	pcDNA6-H2B-mCherry (plasmid)	PMID: 25113836		
recombinant DNA reagent	pcDNA6-EB3-GFP (plasmid)	PMID: 25113836		

recombinant DNA reagent	pcDNA3-AT1.03 (plasmid)	PMID: 19720993		
recombinant DNA reagent	pcDNA3-AT1.03 RK (plasmid)	PMID: 19720993		
recombinant DNA reagent	pCMV-Trim21 (plasmid)	ORIGENE	Cat#: MR207378	
sequence-based reagent	pcDNA6-Golgi-mCherry cloning (sense)	Eurofins	PCR primers	5'-ATCAAGCTTGCCACCATGGGCAACTTGAAG-3'
sequence-based reagent	pcDNA6-Golgi-mCherry cloning (antisense)	Eurofins	PCR primers	5'-TATCTCGAGACCACCTCCACCTCCTCCA-3'
sequence-based reagent	pcDNA6-PEX-mCherry cloning (sense)	Eurofins	PCR primers	5'-GATCTCGAGCTCAAGCTTCGAATTCTGCAG-3'
sequence-based reagent	pcDNA6-PEX-mCherry cloning (antisense)	Eurofins	PCR primers	5'-TATCTAGAGTCGCGGCCGCTACAGCTTG-3'
sequence-based reagent	pcDNA6-mCherry-Drp1 infusion cloning (sense)	Eurofins	PCR primers	5'-CAGTGTGGTGGAATTGCCACCATGGTGAGCAAG-3'
sequence-based reagent	pcDNA6-mCherry-Drp1 infusion cloning (antisense)	Eurofins	PCR primers	5'-ACTGTGCTGGATATCTCACCAAAGATGAGTCTCC-3'
sequence-based reagent	mtDNA qPCR (left)	Eurofins	PCR primers	5'-CCTATCACCCTTGCCA-3

sequence-based reagent	mtDNA qPCR (right)	Eurofins	PCR primers	5'-GAGGCTGT TGCTTGTG-3'
commercial assay or kit	RiboMAX™ Large Scale RNA Production Systems	Promega	Cat# P1300	
commercial assay or kit	Ribo m7G Cap Analog	Promega	Cat# P1712	
commercial assay or kit	T7 mMessage mMachine Kit	Thermo Fisher Scientific	Cat# AM1344	
commercial assay or kit	Poly(A) Tailing Kit	Thermo Fisher Scientific	Cat# AM1350	
chemical compound, drug	Hyaluronidase	Sigma-Aldrich	Cat# D2650	
chemical compound, drug	Protease	Sigma-Aldrich	Cat# P5147	
chemical compound, drug	DMSO	Sigma-Aldrich	Cat# D2650	
chemical compound, drug	Cytochalasin B	Sigma-Aldrich	Cat# C6762	
chemical compound, drug	Polyvinylpyrroli done (PVP) Solution	Irvine Scientific	Cat# 90123	
chemical compound, drug	Thapsigargin	Thermo Fisher Scientific	Cat# T9033	

chemical compound, drug	MitoTracker™ Red CMXRos	Thermo Fisher Scientific	Cat# M7512	
chemical compound, drug	CM-H2-DCFDA	Thermo Fisher Scientific	Cat# C6827	
chemical compound, drug	Fluo-4, AM	Thermo Fisher Scientific	Cat# F14201	
chemical compound, drug	Pluronic F-127	Thermo Fisher Scientific	Cat# P6867	
chemical compound, drug	Triton X-100	Sigma-Aldrich	Cat# T9284	
chemical compound, drug	DAPI	Sigma-Aldrich	Cat# D9542	
software, algorithm	Fiji	PMID: 22743772		
software, algorithm	QuantEv Icy plugin	PMID: 30091700		
software, algorithm	HCImage	Hamamatsu photonics		
software, algorithm	Prism 8	GraphPad Prism		

eLife Version of Record checklist

4   Please refer to this checklist when you are uploading your Version of Record.

6 ## Article file

Title is approx. 120 characters or fewer. It is one clear and concise sentence and does not include punctuation or multiple parts*. The biological system is indicated in the title and/or abstract. *For articles in **Medicine or Epidemiology and Global Health,** two-part titles are allowed, and the study design is in the title.	☒
Authors: all authors are listed with the correct names, in the desired order with all affiliations relevant to the study.	☒
Abstract: normally a single 150–200 word paragraph* If the biological system not mentioned in the title, it is included early in the abstract. *For Medical submissions:** up to 250 words. Subheadings: Background; Methods; Results; Conclusions; Funding; Clinical trial number (for submissions reporting results of a trial only).	☒
Article file: uploaded as a .docx or .tex file. **LaTex:** for .tex files, all .cls, .sty, and .bib files are uploaded as Latex Support files and a clean, compiled PDF is uploaded as a Related Manuscript file.	☒
Tables are editable and included within the article file so that they can be typeset during the production process. Where possible, bold and/or italics are used instead of colours to highlight cells or data.	☒
Key Resources Table: follows the eLife template. The Key Resources Table is incorporated within the article file at the very beginning of the Materials and Methods section.	☒
Appendices: if applicable, included at the end of the main article file or uploaded in an editable format as a second article file.	☒

8 ## Figure, figure supplement and video files

Figures: each main figure is provided as an individual file in one of the accepted formats (PDF, EPS, AI, JPEG, PNG or 16 bit TIF). No figure numbers, titles, or author names appear in the figure files (e.g., "Figure 1, Holmes et al." is not permitted).	☒
Appendix figures: provided as an individual file in one of the accepted formats (PDF, EPS, AI, JPEG, PNG or 16 bit TIF) and labelled in the format "Appendix 1–figure 1", "Appendix 1–figure 2", "Appendix 2–figure 1" etc.	☒
Figure supplements: each figure supplement is provided as an individual file in one of the accepted formats (PDF, EPS, AI, JPEG, PNG or 16 bit TIF) and fits on one page.	☒
Figure supplements are **linked to a main figure** and labelled in the format "Figure 1–figure supplement 1", "Figure 1–figure supplement 2", "Figure 2–figure supplement 1" etc.	☒
Principles of Colour Universal Design (Masataka, Okabe and Kei Ito, J*Fly) used when preparing figures, to ensure maximum accessibility for all types of colour vision.	☒
Rich Media/Videos: uploaded in one of the accepted formats (AVI, WMV, MOV, MP4, or H264). Rich media files can be independent or linked to main figure (e.g. Animation 1, Video 1, Figure 1–video 1) .	☒
Figure permissions: the legend for any adapted, redrawn, or reproduced figure includes: the reference of the original publication (with original figure number); the licence of the original publication; and that the copyright holder has granted permissions to publish under a CC BY 4.0 licence.	☒

Additional submission files

MDAR checklist A completed version of the MDAR checklist (PDF; Word) has been uploaded, which is suitable for publication in the event of acceptance. Any data information included in this file is also included in the Data Availability Statement.	☒
Source data files: **All data generated and analysed should be shared in accordance with our Data availability policy.**	☒
Source data files under 100MB have been uploaded and are labelled correctly. – Where linked to a figure, figure supplement or table, source data files follow this naming convention: 'Figure 1–Source Data 1', 'Figure 1–Figure Supplement 1–Source Data 1', 'Table 1–Source Data 1'. If the data is linked to more than one display item (e.g. Figure 2 and Figure 3), only link it to **one** of these (e.g. 'Figure 2–Source Data 1'). – Where linked to the manuscript as a whole, source data files are labelled 'Source Data 1', 'Source Data 2' etc. (Datasets which include many files larger than 100MB need to be deposited to an external repository. See our Data availability policy for more information.)	☒
Each source data file has a title (and an optional legend) provided within the article file.	☒
If gels or blots are depicted, source data files are required for the relevant figure(s) and figure supplement(s). Zipped folder(s) containing **both** the original files of the full raw unedited gels or blots **and** the uncropped gels or blots with relevant bands clearly labelled have been uploaded and the files have been labelled correctly (see above).	☒
Source code files: Software and code should be shared in accordance with our Software sharing policy.	☒
Source code files are labelled correctly. – Where linked to a figure, figure supplement or table, source code files follow this naming convention: 'Figure 1–Source Code 1', 'Figure 1–Figure Supplement 1–Source Code 1', 'Table 1–Source Code 1'. If the code is linked to more than one display item (e.g. Figure 2 and Figure 3), the label only links it to **one** of these (e.g. 'Figure 2–Source Code 1'). – Where linked to the manuscript as a whole, source code files are labelled 'Source Code 1', 'Source Code 2' etc.	☒
Each source code file has a title (and an optional legend) provided within the article file.	☒
Supplementary files: Supplementary files are reserved for information not central to the narrative which falls outside of the other formats specified above, such as long lists of strains or plasmids. Supplementary files are uploaded individually in the most useful format (i.e. which allows for re-use of the information contained in the file). Each Supplementary File is uploaded individually using the 'Supplementary File' format and is correctly labelled "Supplementary File 1", "Supplementary File 2", etc.	☒
Answers for the _eLife_ Digest [recommended]: if interested, an 'Answers for the eLife Digest' file has been uploaded, as described in this blog post.	☒
Potential Striking Image [recommended]: is at least 1800 x 900 pixels, provided in png, tiff or jpeg formats. Image is a single panel and does not include labels or text. All Striking Images must be available to use under the terms of a Creative Commons Attribution License. Please see here, or look at our social media channels for more ideas about the types of images we look for.	☒

Submission information

Title and Abstract match the title and abstract provided in the Article file and follow the same guidelines.	☒

Impact Statement: a single, complete sentence under 40 words that does not begin with phrases like: "We show…", "This study…" or "Our work…"	☒
Author details are complete for all authors (name, current affiliation, current email address, city, country and competing interest) and authors are listed in the same order as the Article File.	☒
Author contributions: at least one box checked for every author.	☒
Permissions: state whether any previously published material have been adapted or reproduced. If so, the reference of the original publication (with original figure number); the licence of the original publication; and whether the copyright holder has granted permissions to publish under a CC BY 4.0 licence have been stated.	☒
Data availability statement describes where and how the data and code for this study has been made publicly available, including DOI or accession numbers.	☒
Animal ethics statement states the institutional review board or ethics committee that approved the study and/or the guidelines that have been followed. Statement includes any protocol or reference numbers associated with the ethical approval of your work.	☒
Human ethics statement explains that informed consent, and consent to publish, was obtained, or if this was not necessary, please state this and explain the reasons. If necessary, the consent process can be described in depth in the Materials and Methods. The specific ethical approval obtained and/or guidelines that were followed, including the name of the institutional review board or ethics committee that approved the study, where applicable, along with any relevant approval identifier are included.	☒
Funding information is complete, with grant numbers where possible, and the authors associated with the specific funding sources.	☒
Cell line source has been stated i.e. where the cell lines were obtained from.	☒
Cell identity has been **authenticated** and include details of the authentication method used (such as STR profiling). Alternatively, the identity was verified by other means if cell line authentication is not commercially available.	☒
Confirm that the **mycoplasma contamination testing** is negative.	☒

15
16

Figure 1

Figure 1–figure supplement 1

Figure 2

A

PN 2-cell 4-cell 8-cell Morula Blastocyst   kDa

Drp1 —— 70

β-Actin —— 50

B

Trim21	−	+	−	+	+
anti-Drp1	−	−	+	−	+
Control IgG	−	−	−	+	−

kDa

Drp1 —— 70

β-Actin —— 50

C

Control   Trim-Away

D

mt-GFP   mCh-Drp1

Trim-Away + mCh-Drp1

E

Drp1fl/f   *Drp1Δ/Δ*

F

Control

Trim-Away

G

(327)   (289)

****

Major axis of mitochondria (μm)

Control   Trim-Away

H

AT1.03   AT1.03RK

(15) (13)   (15) (9)

ns

Ratio (YFP / CFP)

Control Trim-Away   Control Trim-Away

Figure 2–figure supplement 1

Figure 3

Figure 3–figure supplement 1

Figure 4

Figure 4–figure supplement 1

Figure 5

A **Trim-away**

mt-GFP / H2B-mCherry

NEB

-3:00 0:00 1:30 2:00 3:00

H2B-mCherry

B **Control** / **Trim-away**

DNA / αTublin / Pericentrin

C **Control** / **Trim-Away**

D

Distance to the cell center (µm)

(12) (12)

***

Control Trim-Away

E

Proportion (%)

(51) (43)

Control Trim-away

- 4 blastomeres with 1 nucleus
- 3 blastomeres (2 blastomeres with 1 nucleus, 1 blastomere with 2 nuclei)
- 2 blastomeres with 2 nuclei
- 2 blastomeres with 2 nuclei (transient 3 blastomeres)
- 2 blastomeres with 2 nuclei (transient 4 blastomeres)

Binucleation

- 2 blastomeres with 1 nucleus (Norma 2-cell embryos)

F

Normal embryos

Mitochondrial fragmentation

Mitochondrial aggregation

Drp1-depleted embryos

Two-pronuclei zygote

Uniform mitochodnrial distribution around parental spindles

Equal partitioning of mitochondria into daugther blastomeres

- Symmetric mitochondrial inheritance
- Parental genome mixing

Mitochondrial centration prevents dual spindle assembly

Mitochondrial clumps disturb organelle partitioning

- Asymmetric mitochondrial inheritance
- Spatial confinement of the ER
- Binucleated blastomeres

Figure 5–figure supplement 1