Positive feedback between demographic and selective fluctuations can greatly amplify the

random fluctuation of population size

Yuseob Kim

Division of EcoScience, Ewha Womans University, Seoul, Korea 03760

yuseob@ewha.ac.kr

Running title: eco-evolutionary feedback for size fluctuations

Keyword: fluctuation, storage effect, population size, positive feedback, eco-evolutionary

Correspondence:

Yuseob Kim, Science Hall B308, Ewhayeodae-gil 52, Seodaemun-gu, Seoul, Korea 03760

Tel: +82 10 5138 5175

ABSTRACT

Population sizes fluctuate over time probably due to random variability in the external

environment. However, the severity of such fluctuations should depend on the characteristics of

a species shaped in its evolutionary history. Previous studies have suggested that species are likely

to evolve to minimize demographic fluctuations because an allele causing a smaller variance in

offspring number is advantageous. However, this study finds that evolution in the opposite

direction, favoring a mutation causing larger fitness fluctuation, occurs in a simple eco-

evolutionary model under a randomly changing environment. This requires that (1) the mutant

allele is under fluctuating selection within a subset, the field, of the population but neutral in

another subset, the refuge, and (2) the field-to-refuge ratio of the carrying capacity is positively

correlated to the fitness of the mutant allele. A general condition for the fixation of such a mutation

was derived to depend on the relative strengths of demographic and fitness fluctuations and the

mutational effect on the carrying capacity. Multi-locus simulations revealed that positive feedback

between demography and selection accelerates the sequential fixations of fluctuation-amplifying

mutations, leading to a drastic size fluctuation. This study therefore offers an unconventional

explanation for large demographic fluctuations.

INTRODUCTION

Natural populations experience temporal changes in abiotic and biotic environments. A randomly
variable environment is expected to produce fluctuations in various aspects of demography, most
importantly the number of reproducing individuals (i.e., population size), as well as fluctuation in
the fitness effects of non-neutral or phenotype-changing alleles segregating in the population. If
fitness differences under fluctuation are large, evolutionary changes may occur rapidly at the
same speed as short-term ecological/demographic fluctuations. Recent developments in
evolutionary biology have revealed that such rapid evolutionary changes are widespread (Bell
2010; Thompson 2013; Messer *et al.* 2016). One may then ask whether mutual interaction
between demographic and selective fluctuations creates qualitatively novel eco-evolutionary
dynamics that cannot arise if one of the demographic or population genetic variables is held
constant.

Considering that a fluctuation in population size reflects a fluctuation in the reproductive success
of individuals and that long-term reproductive success is maximized if its variance in the offspring
number over time is minimized (Slatkin 1974; Gillespie 1977; Tuljapurkar 1982), one may predict
that a population will evolve to reduce size fluctuation as much as possible (Pfister 1998). For
example, a variant increasing the size rather than the number of eggs when plentiful resources
are available may be positively selected as the offspring can survive better during an unfavorable
period. Then, when a large fluctuation in population density, such as mast seeding or insect
outbreak, does occur despite the advantage of a small temporal variance in offspring number, this
may be seen as a non-adaptive response of the population to a heavy fluctuation of the biotic
and/or abiotic environment (Myers 1988; Dwyer *et al.* 2004; White 2008; Bogdziewicz *et al.*
2020).

However, recent studies suggested that a population may evolve toward a larger fluctuation in its
size. Liu *et al.* (2019) found that a reproductive strategy increasing the temporal variance of
offspring number can be advantageous in populations reproducing in overlapping generations
under various patterns of environmental fluctuations. Kim (2023) showed that a mutation
amplifying the cyclic fluctuation of population size can be positively selected if this population is
partially protected from fluctuating selection (i.e. fitness fluctuation occurs only in a subset of the
population) at a given time. An additional condition promoting positive selection in the latter
study was that the oscillation of fitness is positively correlated with that of population size and
the regulation of population density is not strong enough so that the population size can over- or

undershoot the carrying capacity. It is argued that the condition for partial protection from fluctuating selection is commonly satisfied in many species. The amplitude of fluctuation in selective pressure is likely to be heterogeneous over the geographic range of a spatially structured species. In plants and animals that have their life cycles divided into several stages and reproduce in overlapping generations, it is difficult to imagine a selective pressure uniformly acting on all individuals in the population regardless of the stage or age. For example, if a trait under fluctuating selection is expressed in the insect larval stage, alleles affecting this trait currently carried by adults are not visible to the selection. Similarly, alleles affecting traits that appear after germination remain neutral in the seed bank of a plant species. Such a subset of the population protected from fluctuating selection may be called a "refuge." The presence of a refuge of various kinds was found to generate the advantage of rare alleles, a form of balancing selection known as the storage effect (Chesson & Warner 1981; Ellner & Hairston 1994; Reinhold 2000; Gulisija & Kim 2015; Gulisija *et al.* 2016; Bertram & Masel 2019; Yamamichi *et al.* 2023). However, Kim (2023) showed that if the population size is oscillating the negative frequency-dependent selection may not persist but turn into directional selection so that polymorphism is no longer maintained, unless fitness fluctuation is strong relative to population size fluctuation.

The theoretical results above — under what conditions balancing or directional selection on the alleles of fluctuating fitness will occur — were obtained from a model assuming cyclical/seasonal fluctuation in the environment and no reproduction in the refuge. A model with seasonal fluctuation is suitable for cases in which an environmental (seasonal) cycle spans multiple generations. However, for many species, one generation is not shorter than a seasonal cycle. Consequently, this study built a simple mathematical model in which the environment fluctuates randomly over time, which is probably relevant to a wider range of species including plants that exhibit mast seeding. In addition, the demographic assumption (no reproduction) of the refuge in Kim (2023), which was for specifically modeling a seed bank (Turelli *et al.* 2001), was relaxed so that it may represent a separate subpopulation undergoing reproduction and fluctuation in its carrying capacity. A general condition for balancing versus directional selection was analytically derived for the one-locus evolutionary dynamics under random environmental fluctuation. This one-locus model was then extended to a multi-locus model. The simulation of the latter showed that mutant alleles causing wider fluctuation in offspring number can reach fixation at multiple loci in a wide range of conditions. These fixations were found to occur at faster rates with increasing numbers of loci, as fluctuating population size and fluctuating selection reinforced each other in positive feedback. This result shows that eco-evolutionary dynamics arising under

the general condition of heterogeneous fluctuating selection can produce highly variable

demography.

METHODS

Eco-evolutionary model

Consider a haploid population that reproduces in discrete generations. It is divided into two

subpopulations, termed the field and the refuge. A fluctuating environment causes fluctuations in

the carrying capacities of both the field and refuge, which are given by $K_F^{(t)}$ and $K_R^{(t)}$

respectively for generation t. It is assumed that the carrying capacities of both subpopulations

depend on common environmental factors that fluctuate randomly over time but that

subpopulations may differ in the strengths of the dependence. The sizes of subpopulations, $N_F^{(t)}$

and $N_R^{(t)}$, after the reproduction step in generation t, may or many not match the corresponding

carrying capacities depending on the strength of population density regulation (see below).

Before a new generation starts, each individual in both subpopulations is assumed to migrate to

or remain in the field with probability 1 - r and migrate to or remain in the refuge with probability

r. As a result, the proportion of the refuge in the total population is always r at the start of a

generation. Next, the size of the field (refuge) changes from $\dot{N}_F^{(t)}$ ($\dot{N}_R^{(t)}$) at the beginning to $N_F^{(t)}$

($N_R^{(t)}$) at the end of generation t, where $\mathrm{E}[\dot{N}_F^{(t)}] = (1-r)(N_F^{(t-1)} + N_R^{(t-1)})$ and $\mathrm{E}[\dot{N}_R^{(t)}] =$

$r(N_F^{(t-1)} + N_R^{(t-1)})$.

It is assumed that the population is initially fixed for the A_1 allele at a locus. Then, a new mutant

allele A_2, subject to fluctuating selection in the field but neutral in the refuge, arises. The fitness of

an individual carrying A_2 relative to A_1 is given by $1 + S_t$ in the field, where $\mathrm{E}[S_t] = 0$ and

$\mathrm{Var}[S_t] > 0$. Notably, while the arithmetic mean of the fitness of A_2 over time is $\mathrm{E}[1 + S_t] = 1$, its

geometric mean is less than 1. Therefore, natural selection is expected to eliminate this mutant

allele from a population in a simple demographic model (containg the field only).

Let $n_{1\mathrm{F}}$ and $n_{2\mathrm{F}}$ be the numbers of individuals in the field carrying A_1 and A_2 alleles before

reproduction for generation t (thus $n_{1\mathrm{F}} + n_{2\mathrm{F}} = \dot{N}_F^{(t)}$). In the reproduction step, two modes of

natural selection (soft vs. hard selection; Wallace (1975)) are considered regarding the effects of

mutations on the population size. First, under soft selection, very strong density regulation is

assumed so that the final size of the field is constrained to $K_F^{(t)}$. Each parent produces a Poisson

number of progenies with mean $WK_F^{(t)}/\ddot{N}_F^{(t)}$, where W is 1 for A_1 and $1 + S_t$ for A_2, and $\ddot{N}_F^{(t)} =$

$n_{1F} + (1 + S_t)n_{2F}$. (Reproduction in Poisson numbers was needed to accommodate genetic drift;

see below.) Second, under hard selection, fitness-changing mutations lead to changes in not only

allele frequencies but also in the population size; they may increase or decrease the absolute

reproductive output of the carriers such that the population size can increase over or decrease

below the carrying capacity, which is now interpreted as the exeped population size assuming

that all individuals carry the A_1 allele. To this effect, each parent is now assumed to produce a

Poisson number of progenies with mean $WK_F^{(t)}/\dot{N}_F^{(t)}$. Namely, when $q = n_{2F}/(n_{1F} + n_{2F})$ is the

relative frequency of A_2 at the beginning of a generation, the size of the field is expected to reach

$(1 + qS_t)K_F^{(t)}$. Reproduction in the refuge occurs similarly but without selection: each individual

produces a Poisson offspring number with mean $K_R^{(t)}/\dot{N}_R^{(t)}$.

Finally, how the environment fluctuates over time and how demographic and selective parameters

— $K_F^{(t)}$, $K_R^{(t)}$, and S_t — respond need to be specified. The environmental condition is assumed

to fluctuate randomly in each generation. More specifically, fluctuations occur such that $K_F^{(t)} =$

$K_{F0}e^{U_t}$ and $K_R^{(t)} = K_{R0}e^{V_t}$ where U_t and V_t are drawn each generation from a joint normal

distribution satisfying $\mathrm{E}[U_t] = \mathrm{E}[V_t] = 0$ and $\mathrm{Cov}[U_t, V_t] \geq 0$. The oscillation of carrying

capacities is therefore symmetrical in the log scale, which is generally predicted in demographic

models and observed in nature (Tuljapurkar & Orzack 1980; Myers 1988). S_t fluctuates such

that $\mathrm{E}[S_t] = 0$, $\mathrm{Cov}[S_t, U_t] \geq 0$, and $\mathrm{Cov}[S_t, V_t] \geq 0$. Assuming non-zero correlations among the

three parameters means that the population sizes and fitness respond to the fluctuation of

common environmental variables.

Stochastic simulation

Stochastic simulations of the above single-locus eco-evolutionary model and its extension to the

multi-locus model were implemented in Mathematica ver. 11.3 (notebook files available at DOI

10.5281/zenodo.125116522). An individual has L loci that are arranged linearly with probability

c for recombination between adjacent loci per generation. Simulation keeps track of two sets of

lists: one containing the sets of haplotypes carried by non-zero individuals in the field and in the

refuge (H_F and H_R) and the other containing the corresponding numbers of individiuals (N_F and

N_R). Generation t begins after the migration step (re-distribution of individuals produced in the

previous generation into the field and refuge) is completed. Haplotype numbers are first updated

by mutation and recombination. A Poisson variate with mean $NL\mu$, where N is the current size of

a subpopulation at hand, is drawn as the total number of mutation events and each event is

assigned to haplotypes and loci proportional to their current frequencies. Similarly, for $L > 1$, a

Poisson variate with mean $N(L\text{-}1)c$ is drawn as the total number of recombination events in the

subpopulation. For each recombination event, two haplotypes are chosen proportional to their

frequencies and one of the L-1 intervals is chosen to be the position of crossing-over between the

two haplotypes. Newly created or lost haplotypes during mutation and recombination steps are

tracked by updating (H_F, H_R) and (N_F, N_R). Then, the updated lists are further updated in the step

of reproduction according to their fitness. Under soft selection, the number of individuals with

haplotype i born in the next generation is Poisson distributed with mean $W_i \dot{N}_{F,i}^{(t)} K_F^{(t)} / \sum_j W_j \dot{N}_{F,j}^{(t)}$

in the field and $\dot{N}_{R,i}^{(t)} K_R^{(t)} / \sum_j \dot{N}_{R,j}^{(t)}$ in the refuge, where W_i is the relative fitness of haplotype i.

Under hard selection, the corresponding mean number in the field is $W_i \dot{N}_{F,i}^{(t)} K_F^{(t)} / \sum_j \dot{N}_{F,j}^{(t)}$. Finally,

when there are N_F and N_R individuals in the field and refuge, Poisson variates with means rN_F and

$(1\text{-}r)N_R$ are drawn to determine the numbers of migrants from field to refuge and from refuge to

field, respectively. Migrants are chosen proportional to the haplotype frequencies and then

merged with the residents of the destination. This completes a single-generation iteration in

updating haplotype lists.

RESULTS

Analytic results: the fate of the A_2 allele

This section aims to predict the fate of the rare A_2 allele — whether it will be lost, fixed, or

increased to intermediate frequencies and maintained polymorphic in the population — as a

function of the demographic and fitness parameters. First, soft selection with strong density

regulation, under which $N_F^{(t)} = K_F^{(t)}$ and $N_R^{(t)} = K_R^{(t)}$, and no genetic drift are assumed. Let n_1

and n_2 be the total numbers of individuals (counted after reproduction/selection and before

migration) carrying A_1 and A_2 in generation t – 1 and n'_1 and n'_2 be their expected numbers in

generation t. Then, with $n_1 \gg n_2$,

$$\frac{n'_2}{n_2} = (1-r)\frac{K_F^{(t)}}{\dot{N}_F^{(t)}}(1+S_t) + r\frac{K_R^{(t)}}{\dot{N}_R^{(t)}} \approx (1-r)\frac{K_F^{(t)}}{\dot{N}_F^{(t)}}(1+S_t) + r\frac{K_R^{(t)}}{\dot{N}_R^{(t)}} = \frac{(1+S_t)K_F^{(t)}+K_R^{(t)}}{K_F^{(t-1)}+K_R^{(t-1)}}. \qquad (1)$$

This equation takes it into account that the change in the absolute number of A_2 is determined

first by demography, as the size of field changes from $\dot{N}_F^{(t)}$ to $K_F^{(t)}$ and that of refuge from $\dot{N}_R^{(t)}$

to $K_R^{(t)}$, and also by fitness $1 + S_t$. Whether the copy number of rare allele A_2 is expected to

increase or decrease is determined by the expectation of $\log\left[\frac{n_2'}{n_2}\right]$ over demographic and fitness

fluctuations. It is shown in Appendix A that, the condition for the A_2 allele being positively selected

(i.e. invading the population initially fixed for A_1), $\mathrm{E}\left[\log\left[\frac{n_2'}{n_2}\right]\right] > 0$, is simplified to

$$\Phi = \frac{\mathrm{Cov}[S,U] - \mathrm{Cov}[S,V]}{\mathrm{Var}[S]} = \frac{\mathrm{Cov}[S,\log[K_F/K_R]]}{\mathrm{Var}[S]} > \frac{1}{2\rho} \tag{2}$$

where $\rho = K_{R0}/K_{F0}$ (note that $K_F^{(t)} = K_{F0}e^{U_t}$ and $K_R^{(t)} = K_{R0}e^{V_t}$) represents the relative size of

the refuge. Therefore, the rare A_2 allele is positively selected if the covariance between its fitness

and the log-ratio of the field to refuge size is large relative to the amplitude of the fitness

fluctuation. This requires that the demographic fluctuation of the field be sufficiently large, as

already shown in Kim (2023), and that of the refuge is either small or out of phase with the

fluctuation of the field. This condition is easier to meet with the increasing size of the refuge.

In a similar manner, the condition for the A_1 allele to invade a population initially fixed for A_2 under

soft selection becomes

$$\Phi < 1 + \frac{1}{2\rho} \tag{3}$$

(Appendix A). From inequalities (2) and (3), it is predicted that the A_1 and A_2 alleles coexist in the

population if $1/(2\rho) < \Phi < 1 + 1/(2\rho)$ under soft selection. With $\Phi < 1/(2\rho)$, the A_2 allele is

eliminated from the population, as the fitness fluctuation that reduces its geometric mean in the

field is more important than population size fluctuation. With $\Phi > 1 + 1/(2\rho)$, the A_2 allele is

predicted to reach fixation in the population.

Using the same approach, the fate of the A_2 allele under hard selection can be analyzed. Note that

the carrying capacity of the field is given by $(1 + qS_t)K_F^{(t)}$ with hard selection where q is the

frequency of A_2. The frequency change of A_2 when rare is therefore effectively identical to that in

soft selection. Therefore, the condition for A_2 invading the population fixed for A_1 is again $\Phi >$

$1/(2\rho)$. However, the condition for the fixation of A_2 is drastically different from that under soft

selection. It is shown in Appendix A that the condition for $\mathrm{E}\left[\log\left[\frac{n_1'}{n_1}\right]\right] > 0$ for rare A_1 is exactly

the condition for $\mathrm{E}\left[\log\left[\frac{n_2'}{n_2}\right]\right] < 0$ for rare A_2. Therefore, the condition for the A_1 allele to invade

the population fixed for A_2 is approximately $\Phi < 1/(2\rho)$, which means that the fixation of A_2 is

ensured if $\Phi > 1/(2\rho)$: once positively selected after its appearance by mutation, the A_2 allele is

expected to increase in frequency all the way to fixation. This leads to a fluctuation amplification

in the size of the field, from $K_F^{(t)}$ to $(1 + S_t)K_F^{(t)}$. This result is consistent with the finding in Kim

(2023) that, under weak density regulation of the population size, an allele with a wider

oscillation in absolute fitness reaches fixation in the presence of refuge.

One-locus simulation

To validate and explore the dynamics of the above eco-evolutionary model, the one-locus

stochastic simulation was performed. The demographic and fitness fluctuations were given by

$U_t = a_U Z_1 + b_U Z_2$, $V_t = a_V Z_1 + b_V Z_3$, and $S_t = a_S Z_1 + b_S Z_4$, where the Z_is are standard normal

variates that were drawn independently each generation. Therefore, population sizes and fitness

were modeled to respond to the fluctuation of a common environmental determinant represented

by Z_1. In this setting, $\Phi = (a_S a_U - a_S a_V)/(a_S^2 + b_S^2)$.

Simulations were first performed with soft selection. Initially the numbers of A_1 and A_2 individuals

changed deterministically once the demographic and fitness parameters (U, V, and S) for a given

generation were sampled from their joint distribution, effectively assuming that the sizes of the

field and refuge were finite and fluctuating but very large so that genetic drift can be ignored (Fig.

1A). Various values of a_S and b_S were chosen so that Φ ranged from 0.3 to 2, while ρ was set to 1.

Starting from 0.5, the frequency of the A_2 allele at the 5,000th generation was recorded for 1000

replicates per parameter set. As predicted, with $\Phi \leq 0.5$, the frequency of A_2 approached 0, thus

confirming negative selection, and with $\Phi \geq 1.5$ the frequency approached 1, thus confirming

positive directional selection. The distribution of the allele frequency was approximately uniform

between 0 and 1 when $\Phi = 1$.

Next, genetic drift and bidirectional mutation between two alleles were added to the above

simulations (Fig. 1B). This weakened the strengths of all modes of selection. With $\Phi = 1$, more

simulation runs ended with allele frequencies close to 0 and 1. However, the proportion of runs

ending with intermediate allele frequencies was still larger than that observed when the alleles

were set as neutral (dashed curve in Fig. 1B). Therefore, an evolutionary force to maintain

polymorphism beyond the level under the neutrality, namely balancing selection, was confirmed.

Similarly, the fate of the A_2 allele under hard selection was examined. Stochastic simulation

confirmed the approximation above predicting that A_2 is positively selected towards fixation if

$\Phi > \frac{1}{2\rho} = 0.5$, although genetic drift and mutation weakened the trend (Fig 1C,D). Positive

selection on both rare and common A_2 under the same condition means that this allele may not

remain polymorphic for long in the population. Indeed, across all values of Φ there were fewer

simulation runs ending with intermediate frequencies of A_2 compared to the simulation of neutral

alleles subject to genetic drift and bidirectional mutations only (dashed curve in Fig. 1D).

Therefore, balancing selection that was observed with soft selection did not arise with hard

selection.

Figure 1. Distribution of A_2 allele frequencies (p) at the 5,000th generation in the stochastic simulations

of soft (A, B) and hard (C, D) selection. The initial frequency of A_2 is 0.5 in both field and refuge.

Reproduction occurred while $K_F^{(t)}$, $K_F^{(t)}$ and S_t varied stochastically without (A, C) or with (B, D) genetic

drift and bidirectional mutation ($\mu = 5 \times 10^{-5}$). Proportions of simulation replicates (out of 1000) with the

frequencies in the interval $0 \le p < 0.1, 0.1 \le p < 0.2, ..., 0.8 \le p < 0.9$, and $0.9 \le p \le 1$ were plotted and

connected by lines. Fluctuations of population sizes are given by $K_{R0} = K_{F0} = 1000$ ($\rho = 1$), $a_U =$

$0.3, b_U = 0.1, a_V = 0.05, b_V = 0.1$. Parameters of fluctuation were chosen to yield $\Phi = 0.3, 0.5, 0.77, 1, 1.2,$

$1.5,$ and 2: $(a_S, b_S) = (0.07, 0.23), (0.1, 0.2), (0.14, 0.16), (0.2, 0.1), (0.2, 0.04), (0.15, 0.05),$ and $(0.1, 0.05)$.

In the case of hard selection with $\Phi = 1$ and no genetic drift (panel C), simulation ran only upto 2,500

generations: longer simulations or larger Φ resulted in all replicates with final frequencies > 0.9.

Multi-locus simulation

To investigate how the eco-evolutionary dynamics shown in the single-locus model above play out when there are many such loci in the genome, new sets of simulation were performed in which a haploid individual is modeled to have L linked loci, each carrying either the A_1 or A_2 allele described above. The simulation tracks the numbers of individuals of different haplotypes (see METHODS). Epistasis is not considered; the relative fitness of haplotype i is given by $W_i = \prod_{l=1}^{L}(1 + S_{t,l})$, where $S_{t,l}$ is 0 if the lth locus of the haplotype carries A_1 and $a_S Z_1 + b_S Z_4$ if A_2. Z_1 is drawn once for all loci while Z_4 is drawn separately for different loci. Individuals randomly pair and undergo recombination by crossing-over such that the probability of recombination between adjacent loci is c per generation. The simulation starts with a population fixed for A_1 at all loci. Then A_2 arises by mutations with probability μ/locus/generation.

Figure 2. Changes of mutant (A_2) allele frequencies and population size ($N_F^{(t)} + N_R^{(t)}$) in a multi-locus simulation of hard selection with L = 8, c = 0.02, K_{R0} = K_{F0} = 1000, μ = 2×10^{-5}, and Φ = 1 (a_S = 0.15, b_S = 0, a_U = 0.15, a_V = 0, b_U = 0, b_V = 0.1). Allele frequency trajectories of 8 loci were plotted in different colors.

Simulations can be run under two different parameter domains. First, the demographic and selective parameters that lead to balancing selection in the one-locus simulation (soft selection

with $1/(2\rho) < \Phi < 1 + 1/(2\rho))$ were chosen. In limited sets of simulation (L = 4, or 10), the

observed level of per-locus polymorphism was above the level in equivalent simulations with

neutral alleles (S_t = 0), thus confirming the operation of multi-locus balancing selection. This

result is described in Appendix B. This study does not further explore this but focuses on the

second (and wider) parameter domain, namely $\Phi > 1 + 1/(2\rho)$ with soft selection and $\Phi >$

$1/(2\rho)$ with hard selection, in which the fixation of the A_2 allele was observed in the single-locus

simulation. As expected, substitutions occurred sequentially until A_2 reached fixation at all L loci

(Fig. 2 and Fig. S1 for example). Times (in generations) when the frequency of A_2 first exceeded

0.9 at each locus were recorded and sorted into T_1 to T_L. There is a large difference in substitution

dynamics between soft and hard selection. With soft selection, the mean waiting time, $\bar{T} = (T_1 +$

$\cdots + T_L)/L$, increased moderately as L increased from 1 to 8 (Fig. 3A). This may be explained by a

reduction in the efficacy of positive selection on A_2 at a given locus when selection at other loci

results in genome-wide reduction in the effective population size.

Figure 3. Mean fixation time ($\bar{T}$) with varying number of loci (A; L = 1, 2, 4, and 8) and the coefficient of variation in fixation times with L = 8 in multi-locus simulation. Error bars show two times the standard errors. A total of 100 replicates of simulations were run under soft selection with Φ = 2 (cyan; a_S = 0.1, b_S = 0, a_U = 0.25, b_U = a_V = b_V = 0.05), hard selection with Φ = 1 (brown; a_S = 0.1, b_S = 0, a_U = 0.1, a_V = 0, b_U = b_V = 0.1), or hard selection with Φ = 0.75 (orange; a_S = 0.2, b_S = 0, a_U = 0.25, a_V = 0.1, b_U = b_V = 0.05). Other parameters are identical to those in Figure 3.

287   With hard selection, however, $\bar{T}$ decreased as L increased. Furthermore, the coefficient of

288   variation, the standard deviation of T_i's divided by $\bar{T}$, was significantly smaller with hard selection

289   than with soft selection (Fig. 3B). As expected for hard selection, the amplitude of the population

size fluctuation progressively increased as the number of loci fixed for A_2 increased (Fig. 2). These results clearly demonstrate that positive feedback between demographic and selective fluctuations under hard selection led to a great amplification of population size fluctuation; an initial fluctuation in $K_F^{(t)}$ positively selected the mutant allele A_2 at one locus whose fitness fluctuates in positive correlation with $K_F^{(t)}$, leading to the fixation of A_2 and therefore a larger fluctuation of the field (given by $(1 + S_t)K_F^{(t)}$), which makes it easier for the A_2 allele at another locus to be positively selected. This process could continue to increase the fluctuation of the field until its carrying capacity is given by $(1 + S_t)^L K_F^{(t)}$, where L is the number of loci available in the genome for mutation into A_2. There was also a moderate effect of the recombination rate in accelerating the multi-locus fixation of A_2 with hard selection (Table S1). Fixation times generally increased with decreasing recombination rates for both soft and hard selection, suggesting that tighter linkage increased the effect of clonal interference.

DISCUSSION

A key assumption in this study is that a tradeoff commonly occurs for the fitness effects of a mutant allele between different time points in the environmental fluctuation. Namely, a mutant phenotype that increases the reproductive output relative to others when the environment is generally favorable is likely to decrease it relative to others during an unfavorable period. Such a mutant allele increases the variance of the offspring number over time while the mean may remain unchanged. For example, as postulated by Slatkin (1974), a mutation that makes a female bird lay 10 eggs in one batch instead of spreading them over time in a randomly and rapidly fluctuating environment will increase the variance of her fitness while the mean remains the same. Modeling the fluctuating fitness of a mutant by $1 + S_t$ reflects such a tradeoff. Classical theory predicts that such a mutant (termed an arithmetic-mean-preserving or AMF allele in Kim (2023)) is eliminated from a population as its geometric mean fitness is less than 1. However, it was demonstrated in this study that the AMF mutant is positively selected if a correlation between demographic and fitness fluctuations exist in the presence of refuge. Analysis given in Appendix A shows that the arithmetic mean fitness of an AMF allele is effectively increased above 1 due to correlated demographic fluctuation in the field. However, without a refuge, its geometric mean cannot exceed 1. Namely, as pointed out in Kim (2023), by dampening the fluctuation of the AMF allele's effective fitness in the total population the refuge turns the heightened arithmetic mean into the geometric mean greater than 1.

One may also model the fitness of a mutant by $\exp[S_t]$, making the mutant (termed a geometric-
mean-preserving or GMF allele) quasi-neutral as its geometric mean fitness is identical to that of
the wild-type allele even without demographic fluctuation. Analyzing the model of cyclic
environmental oscillation, Kim (2023) showed that the parameter range of positive selection for
a GMF allele is wider than that for an AMF allele. Therefore, the major results obtained in this
study, both balancing and positive directional selection on a fitness-amplifying allele, are probably
easier to achieve with GMF mutations. However, a GMF allele may be considered simply a
beneficial allele as the arithmetic mean of offspring number is greater than the wild-type allele.
Such a mutation is probably rare and was not considered in this study.

The refuge in the current model may correspond to various demographic elements. It directly
models a patch in a spatially structured population that is protected from fluctuating selection.
However, given that many studies modeling the storage effects of a certain life-cycle stage yield
equations similar to eq. (1), namely in the form $p'/p = (1-r)f + r$ where p is the frequency of
the rare-type, r is the proportion of the refuge (e.g., adult stage when selection acts on the
juveniles, seed bank, or diapausing eggs), and f is the relative fitness of the type outside the refuge
(Chesson & Warner 1981; Ellner & Hairston 1994; Turelli *et al.* 2001; Gremer & Venable 2014),
this study should apply to such species if $K_R^{(t)}$ is given an appropriate interpretation. A major
result in this study is that an AMF allele is positively selected if correlation of its fitness fluctuation
with the size of the field is larger than that with the size of the refuge. This requirement is
important for sex-limited gene expression. Previous studies suggested that if a gene is expressed
in one sex the other sex could be a refuge in a dioecious species (Reinhold 2000; Kim 2023).
However, as equal numbers of male and female gametes are used in reproduction, the population
sizes of reproductive males and females are effectively equal while the sex ratio of the adults may
fluctuate. Therefore, Φ is zero as the size of the field relative to the refuge does not fluctuate at all.
Hence, positive selection on the fitness-amplifying mutation shown in this study should not
happen for sex-limited genes, unless GMF mutants are considered.

While the eco-evolutionary model of this study can result in both balancing and directional
selection on size-fluctuation-amplifying alleles, this study focused on the latter. The condition for
the fixation of those alleles is less restrictive compared to that for balanced polymorphism,
because it requires relatively weaker selection (smaller variance in relative fitness) for a given
demographic fluctuation to satisfy $\Phi > 1 + 1/(2\rho)$ with soft selection and $\Phi > 1/(2\rho)$ with
hard selection. Demographic fluctuation that drives this evolutionary process should apply
equally to all loci in the genome. Therefore, fixations occur at all available loci on which proper

mutations, i.e. the A_2 allele, arise, as observed in the multi-locus simulation. With soft selection,

positive selection at each locus is contributed by the same degree of demographic fluctuation

regardless of the evolutionary changes at other loci. However, with hard selection, fluctuation in

the size of the field affecting a given locus intensifies as more fluctuation-amplifying mutations

reach fixation at other loci. Since the carrying capacity of the field is effectively $(1 + S_t)^l K_F^{(t)}$,

where l is the number of loci where A_2 reached fixation, and wider fluctuation generates stronger

positive selection on the rare A_2 (larger $E[\Delta_2]$ in Appendix A), fixation at one locus facilitates the

subsequent fixations at other loci. This positive feedback between demographic fluctuation and

positive selection drastically increases the amplitude of fluctuation in the total population size

(Fig. 2). How much the amplitude will increase should depend on the number of loci at which

mutations subject to hard selection arise. There should be multiple independent ways in which

changes in mutants' relative fitness can be translated into changes in the mean absolute fitness of

the entire population. For various life history traits, one may imagine phenotypes (smaller eggs,

smaller adult body, earlier onset of first reproduction, etc.) that increase the absolute number of

surviving offspring during a favorable, resource-rich period but decrease the number in a bad

period while not affecting the amount of resources that the carriers of the alternative, ancestral

alleles acquire. Then, many loci may be available for hard selection, the combined effect of which

may lead to a very large fluctuation in population size.

The size of a population and its fluctuation should be governed not only by extrinsic factors (the

environmental variables such as temperature and parasite prevalence) but also by intrinsic

factors (species-specific responses to those extrinsic factors). The intrinsic factors such as various

life-history traits are the product of the species' evolutionary history. For example, phenotypic

plasticity responding to environmental cues can reduce the amplitude of the population size

fluctuation, thus acting as a mechanism of population regulation (Caswell 1983; Reed *et al.* 2010),

which according to the classical theory is adaptive as it enhances long-term reproductive success

by increasing the geometric mean fitness (Gillespie 1977; Pfister 1998). If life history traits

evolved in the direction toward suppressing random or cyclic fluctuations in the population size,

fluctuations observed in nature should be interpreted as the effect of variable environments that

the evolved mechanisms of population regulation were unable to contain. However, this study

suggests that evolution can proceed in the opposite direction. A large random fluctuation in

population size can be the product of adaptive evolution driven by positive feedback between

demographic and selective fluctuations. When the degrees of population size fluctuation are

different across species, the eco-evolutionary dynamics proposed in this study might be

particularly important for those exhibiting large fluctuations. Interestingly, annual desert plants

experiencing wider fitness fluctuations after germination have greater proportions of their

population remaining in the seed bank (Gremer & Venable 2014). This observation is in

agreement with the analytic results in this study because a larger seed bank corresponds to a

larger value of ρ, which makes it easier for the fixation of the fluctuation-amplifying alleles (i.e.,

lowering the threshold of Φ for positive selection).

ACKNOWLEDGMENT

This study was supported by the National Research Foundation (NRF) grant 2020R1A2C1009261

funded by the Korean government.

Conflict of interest statement

The author declares no conflict of interest

Literature Cited

Bell, G. (2010). Fluctuating selection: the perpetual renewal of adaptation in variable
environments. *Philos Trans R Soc Lond B Biol Sci*, 365, 87-97.

Bertram, J. & Masel, J. (2019). Different mechanisms drive the maintenance of polymorphism at
loci subject to strong versus weak fluctuating selection. *Evolution*, 73, 883-896.

Bogdziewicz, M., Ascoli, D., Hacket-Pain, A., Koenig, W.D., Pearse, I., Pesendorfer, M. *et al.* (2020).
From theory to experiments for testing the proximate mechanisms of mast seeding: an
agenda for an experimental ecology. *Ecology Letters*, 23, 210-220.

Caswell, H. (1983). Phenotypic plasticity in life-history traits: demographic effects and
evolutionary consequences. *American Zoologist*, 23, 35-46.

Chesson, P.L. & Warner, R.R. (1981). Environmental variability promotes coexistence in lottery
competitive systems. *The American Naturalist*, 117, 923-943.

Dwyer, G., Dushoff, J. & Yee, S.H. (2004). The combined effects of pathogens and predators on
insect outbreaks. *Nature*, 430, 341-345.

Ellner, S. & Hairston, N.G. (1994). Role of overlapping generations in maintaining genetic variation
in a fluctuating environment. *The American Naturalist*, 143, 403-417.

Gillespie, J.H. (1977). Natural selection for variances in offspring numbers: a new evolutionary
principle. *The American Naturalist*, 111, 1010-1014.

Gremer, J.R. & Venable, D.L. (2014). Bet hedging in desert winter annual plants: optimal
germination strategies in a variable environment. *Ecol Lett*, 17, 380-387.

Gulisija, D. & Kim, Y. (2015). Emergence of long-term balanced polymorphism under cyclic
selection of spatially variable magnitude. *Evolution*, 69, 979-992.

Gulisija, D., Kim, Y. & Plotkin, J.B. (2016). Phenotypic plasticity promotes balanced polymorphism
in periodic environments by a genomic storage effect. *Genetics*, 202, 1437-1448.

Kim, Y. (2023). Partial protection from fluctuating selection leads to evolution towards wider
population size fluctuation and a novel mechanism of balancing selection. *Proceedings of*
the Royal Society B, 290, 20230822.

Liu, M., Rubenstein, D.R., Liu, W.-C. & Shen, S.-F. (2019). A continuum of biological adaptations to
environmental fluctuation. *Proceedings of the Royal Society B*, 286, 20191623.

Messer, P.W., Ellner, S.P. & Hairston, N.G. (2016). Can population genetics adapt to rapid evolution?
Trends in Genetics, 32, 408-418.

Myers, J.H. (1988). Can a general hypothesis explain population cycles of forest Lepidoptera? In:
Advances in Ecological Research. Elsevier, pp. 179-242.

Pfister, C.A. (1998). Patterns of variance in stage-structured populations: evolutionary predictions
and ecological implications. *Proceedings of the National Academy of Sciences*, 95, 213-218.

Reed, T.E., Waples, R.S., Schindler, D.E., Hard, J.J. & Kinnison, M.T. (2010). Phenotypic plasticity and
population viability: the importance of environmental predictability. *Proceedings of the*
Royal Society B: Biological Sciences, 277, 3391-3400.

Reinhold, K. (2000). Maintenance of a genetic polymorphism by fluctuating selection on sex-
limited traits. *Journal of Evolutionary Biology*, 13, 1009-1014.

Slatkin, M. (1974). Hedging one's evolutionary bets. *Nature*, 250, 704-705.

Thompson, J.N. (2013). *Relentless evolution*. University of Chicago Press.

Tuljapurkar, S. & Orzack, S.H. (1980). Population dynamics in variable environments I. Long-run
growth rates and extinction. *Theoretical Population Biology*, 18, 314-342.

Tuljapurkar, S.D. (1982). Population dynamics in variable environments. III. Evolutionary
dynamics of r-selection. *Theoretical population biology*, 21, 141-165.

Turelli, M., Schemske, D.W. & Bierzychudek, P. (2001). Stable two-allele polymorphisms
maintained by fluctuating fitnesses and seed banks: protecting the blues in Linanthus
parryae. *Evolution*, 55, 1283-1298.

Wallace, B. (1975). Hard and soft selection revisited. *Evolution*, 465-473.

White, T. (2008). The role of food, weather and climate in limiting the abundance of animals.
Biological Reviews, 83, 227-248.

Yamamichi, M., Letten, A.D. & Schreiber, S.J. (2023). Eco-evolutionary maintenance of diversity in
fluctuating environments. *Ecology Letters*, 26, S152-S167.

Appendix A. Derivation of conditions for balancing and directional selection on A_2

Here only deterministic changes in allele frequencies are tracked as a very large population is
assumed. At the end of reproduction but before migration into subpopulations at generation t - 1,
there are n_1 and n_2 copies (individuals) of A_1 and A_2 in the entire population. After migration, the
numbers of A_1 and A_2 copies are $n_{1F} = (1-r)n_1$ and $n_{2F} = (1-r)n_2$ in the field and $n_{1R} =$
rn_1 and $n_{2R} = rn_2$ in the refuge. Therefore, the relative frequencies of alleles are identical
between the field and refuge before reproduction at the beginning of generation t.

Soft selection is first considered. Then, each A_2 allele produces $(1 + S_t)K_F^{(t)}/\ddot{N}_F^{(t)}$ copies of
descendants, where $\ddot{N}_F^{(t)} = n_{1F} + (1 + S_t)n_{2F} = (1-r)(n_1 + (1 + S_t)n_2)$. Since $n_1 \gg n_2$ after A_2
appears by mutation, $n_1 + (1 + S_t)n_2 \approx n_1$ and $\ddot{N}_F^{(t)} \approx \dot{N}_F^{(t)} = (1-r)(n_1 + n_2) = (1-$
$r)\left(K_F^{(t-1)} + K_R^{(t-1)}\right)$. Similar approximations apply to the refuge. Let n'_2 be the expected number
of A_2 in generation t. This leads to eq. (1). Then, with $K_F^{(t-1)} = K_{F0}e^U$, $K_R^{(t-1)} = K_{R0}e^V$, $K_F^{(t)} =$
$K_{F0}e^{U'}$, and $K_R^{(t)} = K_{R0}e^{V'}$,

471
$$\Delta_2 \equiv \log\left[\frac{n'_2}{n_2}\right] = \log[e^{U'}(1 + S) + \rho e^{V'}] - \log[e^U + \rho e^V],$$

472 where $S = S_t$ and $\rho = K_{R0}/K_{F0}$. If $E[\Delta_2] > 0$, where the expectation is over demographic and
473 fitness fluctuations, the rare allele A_2 is expected to increase its copy number over time. Applying
474 the Taylor expansion up to the second order,

475
$$\Delta_2 \approx \frac{1}{1+\rho}(S + U' - U) - \frac{\rho}{1+\rho}(V' - V) - \frac{1}{2(1+\rho)^2}S^2 + \frac{\rho}{2(1+\rho)^2}\left(U'^2 + V'^2 - U^2 - V^2\right)$$

476
$$+ \frac{\rho}{(1+\rho)^2}(SU' - SV' - U'V' + UV)$$

Since all random variables in the above equation have the mean at zero and (U, V) and (U', V')
are from the same joint probability distribution,

$$\mathrm{E}[\Delta_2] \approx -\frac{1}{2(1+\rho)^2}\mathrm{Var}[S] + \frac{\rho}{(1+\rho)^2}\left(\mathrm{Cov}[S,U] - \mathrm{Cov}[S,V]\right) = \frac{\mathrm{Var}[S]}{(1+\rho)^2}\left(\rho\Phi - \frac{1}{2}\right),$$

where

$$\Phi = \frac{\mathrm{Cov}[S,U] - \mathrm{Cov}[S,V]}{\mathrm{Var}[S]} = \frac{\mathrm{Cov}[S, \log[K_F/K_R]]}{\mathrm{Var}[S]}.$$

Then, the condition for the A_2 allele being positively selected (i.e. invading the population initially

fixed for A_1), is $\mathrm{E}[\Delta_2] > 0$ or $\Phi > \frac{1}{2\rho}$.

A similar result can be obtained for the condition for the A_1 allele to invade a population initially

fixed for A_2. From

$$\frac{n_1'}{n_1} \approx \frac{K_F^{(t)}}{K_F^{(t-1)} + K_R^{(t-1)}}\frac{1}{1+S_t} + \frac{K_R^{(t)}}{K_F^{(t-1)} + K_R^{(t-1)}} = \frac{e^{U'} + (1+S_t)\rho e^{V'}}{(1+S_t)(e^U + \rho e^V)}$$

for $n_1 \ll n_2$, the dynamics of the rare A_1 allele depends on the expectation of

$$\Delta_{1,\mathrm{soft}} \equiv \log\left[\frac{n_1'}{n_1}\right] = \log\left[e^{U'} + (1+S)\rho e^{V'}\right] - \log[(1+S)(e^U + \rho e^V)].$$

Then, using quadratic approximation

$$\Delta_{1,\mathrm{soft}} \approx \frac{1}{1+\rho}(-S + U' - U) + \frac{\rho}{1+\rho}(V' - V) + \frac{1+2\rho}{2(1+\rho)^2}S^2$$

$$+ \frac{\rho}{2(1+\rho)^2}\left(U'^2 + V'^2 - U^2 - V^2\right) + \frac{\rho}{(1+\rho)^2}(-SU' + SV' - U'V' + UV)$$

one obtains

$$\mathrm{E}[\Delta_{1,\mathrm{soft}}] \approx \frac{1+2\rho}{2(1+\rho)^2}\mathrm{Var}[S] - \frac{\rho}{(1+\rho)^2}\left(\mathrm{Cov}[S,U] - \mathrm{Cov}[S,V]\right) = \frac{1+2\rho}{2(1+\rho)^2}\mathrm{Var}[S]\left(1 - \frac{2\rho}{2\rho+1}\Phi\right).$$

The condition for the rare A_1 allele to increase in frequency is $\mathrm{E}[\Delta_{1,\mathrm{soft}}] > 0$ or

$$\Phi < 1 + \frac{1}{2\rho}.$$

With hard selection, the frequency change of A_2 when rare is again given by eq. (1). Therefore, the

condition for A_2 invading the population fixed for A_1 ($\mathrm{E}[\Delta_2] > 0$ above) is no different from that

for soft selection. However, because the carrying capacity of the field fixed for A_2 at generation t

is given by $(1 + S_t)K_F^{(t)}$ under hard selection, the opposite condition for A_1 invading A_2 is given

by

501
$$\frac{n_1'}{n_1} = (1-r)\frac{(1+S_t)K_F^{(t)}}{\dot{N}_F^{(t)}}\frac{1}{1+S_t} + r\frac{K_R^{(t)}}{\dot{N}_R^{(t)}} = \frac{K_F^{(t)}+K_R^{(t)}}{(1+S_{t-1})K_F^{(t-1)}+K_R^{(t-1)}}$$

502 Then,

503
$$\Delta_{1,hard} \equiv \log\left[\frac{n_1'}{n_1}\right] = \log[e^{U'} + \rho e^{V'}] - \log[(1+S)e^U + \rho e^V]$$

504 and, again, the quadratic approximation is obtained as

505
$$\Delta_{1,hard} \approx \frac{1}{1+\rho}(-S + U' + \rho V' - U - \rho V) + \frac{1}{2(1+\rho)^2}\left(S^2 + \rho U'^2 + \rho V'^2 - \rho U^2 - \rho V^2\right)$$

506
$$+ \frac{\rho}{(1+\rho)^2}(-SU' + SV' - U'V' + UV)$$

507 Finally,

508
$$E[\Delta_{1,hard}] \approx \frac{1}{2(1+\rho)^2}\mathrm{Var}[S] - \frac{\rho}{(1+\rho)^2}(\mathrm{Cov}[S,U] - \mathrm{Cov}[S,V]) = \frac{1}{2(1+\rho)^2}\mathrm{Var}[S](1-2\rho\Phi).$$

The condition for the fixation of A_2 (the failure of A_1 to invade) is $E[\Delta_{1,hard}] < 0$ or $\Phi > \frac{1}{2\rho}$.

Now, a heuristic argument can be given for positive selection (both soft and hard) on A_2, which is
expected to be eliminated from a population whose size does not fluctuate. Positive selection
occurs because the mean number of offspring per parent, i.e. absolute fitness, of the mutant in the
field is given by $W_{2f} = \frac{K_F^{(t)}}{\dot{N}_F^{(t)}}(1+S_t)$ with fluctuation in population size (eq. 1). With positive
correlation between $K_F^{(t)}$ and S_t, the fluctuation of W_{2f} becomes greater than that of $1+S_t$
and, more importantly, its arithmetic mean can now exceed 1. However, the geometric mean of
W_{2f} is still not greater than 1 due to its fluctuation over time. The presence of a refuge then
reduces the variance of mutant's fitness averaged across the whole population such that the
geometric mean of this average fitness becomes greater than 1. In summary, the refuge turns the
heightened arithmetic mean raised by the demographic fluctuation of the field into an elevation
of the geometric mean in the total population (see also Kim (2023))

Appendix B. Multi-locus simulation for balanced polymorphism

Considering that (1) the negative frequency-dependent selection on a variant with fitness
fluctuation was caused primarily by correlated fluctuation in the carrying capacity (population
size) of the field and that (2) this demographic fluctuation should be "felt" at all loci in the genome,

one may expect that such balancing selection can occur simultaneously at many loci harboring

such variants. To verify if this is true, simulations were run under soft selection and

demographic/selective parameters that promoted polymorphism in the single-locus model (Φ =

1 for ρ = 1) with L = 1, 4, or 10. Exemplary trajectories of allele frequencies are shown in Fig. S2.

Each run lasted at least 10^5 generations, over which the mean of expected heterozygosity (H =

$2p(1-p)$) and the proportions of time A_2 spent in frequency between 0.1 and 0.5, P_{15}, and

between 0.5 and 0.9, P_{59}, were recorded per locus (Table S2). In all cases with fluctuating fitness

($a_S > 0$), polymorphism per locus was above the level observed in equivalent simulations with

neutral alleles (S_t = 0), thus confirming multi-locus balancing selection. For cases of stronger

selection (larger $\text{Var}[S_t]$), the level of polymorphism per locus declined as L increased from 1 to

10. The combined effects of the number of loci and the rate of recombination were not easy to

interpret as they depended on the parameters of selection (for sets of fitness/demographic

parameters that yield the same Φ = 1, H responded differently with varying L and c). One possible

explanation for generally lower variation with 10 loci instead of 1 or 4 is that the simultaneous

segregation of alleles at many loci increased the variance of the offspring numbers, thus

decreasing the effective population size and weakening the negative frequency-dependent

selection that is needed for protected polymorphism. Further exploration of multi-locus

polymorphism at a larger scale (in both number of loci and population size) may not be feasible

with the current simulation approach based on haplotype frequencies.

Table S1. Mean time to the fixation ($q > 0.9$) of A_2 in the multi-locus simulation

mode	L	c	$\bar{T}$	c. v.*
	1	-	5863	-
soft selection, $\Phi = 2$	5	10^{-4}	9287	0.66 ± 0.21
($a_S = 0.1$, $b_S = 0$, $a_U = 0.25$,	5	10^{-3}	8212	0.67 ± 0.20
$a_V = 0.05$, $b_U = b_V = 0.05$)	5	0.01	8621	0.70 ± 0.26
	5	0.1	8162	0.65 ± 0.26
	1	-	6242	-
hard selection, $\Phi = 0.75$	5	10^{-4}	3776	0.34 ± 0.17
($a_S = 0.2$, $b_S = 0$, $a_U = 0.25$,	5	10^{-3}	3228	0.32 ± 0.16
$a_V = 0.1$, $b_U = b_V = 0.05$)	5	0.01	3030	0.31 ± 0.20
	5	0.1	3459	0.28 ± 0.19
	1	-	5177	-
hard selection, $\Phi = 1$	5	10^{-4}	4648	0.35 ± 0.15
($a_S = 0.15$, $b_S = 0$, $a_U = 0.15$,	5	10^{-3}	4018	0.35 ± 0.17
$b_U = 0.05$, $a_V = 0$, $b_V = 0.1$)	5	0.01	3969	0.32 ± 0.20
	5	0.1	3467	0.37 ± 0.21

549   *Coefficient of variation for 5 consecutive times (T_1, ..., T_5) observed in a single simulation run,

550   averaged over replicates ($\pm$ standard deviation).

551   All results are based on 100 replicates for each parameter set

554 Table S2. The long-term level of polymorphism in the multi-locus simulation

mode	L	c	H	P_{15}	P_{59}
neutral ($a_S = b_S = 0$, $a_U = 0.2$, $a_V = b_U = b_V = 0.1$)	1	-	0.0639	0.069	0.074
soft selection, $\Phi = 1$ ($a_S = 0.1$, $b_S = 0$, $a_U = 0.2$, $a_V = b_U = b_V = 0.1$)	1	-	0.132	0.15	0.16
	4	10^{-4}	0.0960	0.099	0.12
	4	10^{-3}	0.103	0.12	0.12
	4	0.01	0.0977	0.11	0.12
	4	0.1	0.136	0.11	0.14
	10	10^{-4}	0.080	0.087	0.095
	10	10^{-3}	0.102	0.12	0.12
	10	0.01	0.109	0.15	0.10
	10	0.1	0.107	0.13	0.12
neutral ($a_S = b_S = 0$, $a_U = 0.3$, $a_V = 0.05$, $b_U = b_V = 0.1$)	1	-	0.0619	0.056	0.079
soft selection, $\Phi = 1$ ($a_S = 0.2$, $b_S = 0.1$, $a_U = 0.3$, $a_V = 0.05$, $b_U = b_V = 0.1$)	1	-	0.177	0.24	0.18
	4	10^{-4}	0.162	0.19	0.19
	4	10^{-3}	0.130	0.21	0.11
	4	0.01	0.168	0.22	0.18
	4	0.1	0.183	0.25	0.19
	10	10^{-4}	0.120	0.14	0.14
	10	10^{-3}	0.141	0.17	0.17
	10	0.01	0.0934	0.076	0.14
	10	0.1	0.105	0.093	0.15
neutral ($a_S = b_S = 0$, $a_U = 0.05$, $a_V = b_U = b_V = 0$)	1	-	0.0660	0.085	0.063
soft selection, $\Phi = 1$ ($a_S = 0.05$, $b_S = 0$, $a_U = 0.05$, $a_V = b_U = b_V = 0$)	1	-	0.0816	0.11	0.081
	10	10^{-4}	0.0728	0.085	0.079
	10	10^{-3}	0.0982	0.12	0.11
	10	0.01	0.0953	0.12	0.096
	10	0.1	0.0952	0.13	0.092

- Simulations were run for 5×10^5, 2×10^5, and 10^5 generations with L = 1, 4, and 10, respectively.

- Other parameters: $K_{R0} = K_{F0}$ = 1,000 ($\rho = 1$), $\mu = 2 \times 10^{-5}$.

Figure S1

Figure S1 (continued)

Figure S1 legend: Examples of simulation runs in which the mutant allele (A_2) reaches fixation

sequentially at all 8 loci used. The change of population size is shown below that of allele

frequencies, plotted in different colors for different loci. Parameter values are shown on the

right side of graphs. Other parameters: $K_{R0} = K_{F0} = 1000$, $\mu = 2\times10^{-5}$.

Figure S2

Figure S2 legend: Exemplary trajectories of allele (A_2) frequencies in simulations. Allele

frequencies are plotted in different colors for different loci. The number of loci (L) and

recombination rate (c) are shown on the right side of graphs. Other parameters: $K_{R0} = K_{F0} =$

1000, $\mu = 2 \times 10^{-5}$, $\Phi = 1$ ($a_S = 0.2$ [0 for neutral], $b_S = 0.1$, $a_U = 0.3$, $b_U = 0.1$, $a_V = 0.05$, $b_V = 0.1$).