The reduction of nuptial gifts in sclerosomatid Opiliones coincides

with increased behavioural sexual conflict

Tyler A. Brown[a,*,1]

Emily Marinko[a,2]

Mercedes Burns[a]

[a] Department of Biological Sciences, University of Maryland Baltimore County,

Baltimore, MD, U.S.A.

*Corresponding author: Tyler A. Brown[1]

tabrownresearch@gmail.com

Department of Biological Sciences

University of Maryland, Baltimore County

1000 Hilltop Circle, Baltimore, MD 21250

[1] Present Address: Department of Biological Sciences, University of North Carolina at

Charlotte, Charlotte, NC, U.S.A.

[2] Present Address: School of Life Sciences, University of Nevada Las Vegas, Las

Vegas, NV, U.S.A.

Abstract:

Nuptial gifts serve to increase donor fitness through a variety of pathways. Some nuptial

gifts deliver benefits similar to those of antagonistic male behaviours: functioning to secure

additional copulations, increase sperm transfer or storage, or increase paternity share. These

commonalities may result in evolutionary transitions between solicitous and coercive strategies,

wherein behavioural sexual conflict could function to secure mates in lieu of nuptial gifts. In

temperate leiobunine harvesters (Arachnida: Opiliones), nuptial gifts have been repeatedly lost,

resulting in two primary mating syndromes: an ancestral, sacculate state in which males

endogenously produce high-investment nuptial gifts and females lack pregenital barriers, and a

derived, nonsacculate state in which females have pregenital barriers and males produce

significantly reduced, low-investment nuptial gifts. In this study, we investigated whether

behavioural sexual conflict compensates for reduced nuptial gifts in nonsacculate harvesters by

comparing the intensity of pre-, peri-, and postcopulatory mating behaviour between the

nonsacculate species *Leiobunum vittatum* and *L. euserratipalpe* and the sacculate species *L.*

aldrichi and *L. bracchiolum*. We additionally sought to establish an automated behavioural

analysis pipeline by developing analogues for metrics traditionally scored manually. Our results

revealed significantly different, potentially coercive, behaviour in nonsacculate species, indicating

that the loss and reduction of pre- and pericopulatory nuptial gifts may contribute to increased

behavioural antagonism. Mating behaviour also differed significantly between *L. vittatum* and *L.*

euserratipalpe, indicating there are multiple suites of potentially antagonistic behaviours.

Together, these results suggest that multiple behavioural strategies may be effective substitutes

for nuptial gifts in leiobunine Opiliones, although the mechanisms through which male fitness is

increased requires further research.

Highlights:

- Temperate Opiliones have repeatedly evolved reduced nuptial gift investment
- Reduced-gift (nonsacculate) species possess morphology indicative of antagonism
- We tested if mating behaviour differed between nonsacculate and sacculate species
- Nonsacculates displayed significantly higher behavioural antagonism than sacculates
- Nonsacculate species differed in their suites of antagonistic behaviours

Keywords:

- Antagonism
- Mate guarding
- Mating behaviour
- Nuptial gift
- Opiliones
- Sexual conflict

Introduction

<u>Nuptial gift background</u>

Animals employ a multitude of tactics to increase their fitness during mating interactions, including direct intrasexual competition, mate choice, elaborate signals, and physical or chemical coercion (Buchinger & Li 2023; Davranoglue et al. 2023; Edward & Chapman 2011; Órfão et al. 2023; Rosvall 2011; Schaefer & Ruxton 2015; Shuker & Kvarnemo 2021). Of these myriad strategies, nuptial gifts are among the most unique in their almost altruistic appearance. Nuptial gifts are any materials, beyond gametes, which are provided by a donor to a recipient before or during copulation in order to increase the donor's fitness (Gwynne 2008; Lewis & South 2012; Lewis et al. 2014). These gifts might be consumed orally or received internally during copulation and can range from exogenous materials (such as captured prey wrapped in silk) to endogenous materials (such as proteinaceous fluids) (Gao et al. 2019; Lewis & South 2012). The fitness consequences to the recipient remains intentionally unspecified under this definition, and extensive research shows nuptial gifts can range from being highly beneficial (Albo et al. 2017; Brown & Barry 2016; Gwynne 2008; Lehmann & Lehmann 2016; O'Hara & Brown 2021; Toft & Albo 2015; Voigt et al. 2005; Welke & Schneider 2011) to neutral (Martínez Villar et al. 2021) to highly costly (Arnqvist & Rowe 2005; Sakaluk et al. 2019; Sirot et al. 2015; Wolfner 2002). Predicting these consequences, however, can be difficult due to the highly variant nature of many nuptial gifts, which often vary with source, delivery mechanism, and both within and between individuals (Kahn et al. 2018; Lewis & South 2012; Wedell 1994). Gifts may also fluctuate with environmental conditions, rearing conditions, donor nutrition, or competition intensity (Fox et al. 2006; Ghislandi et al. 2018; Jarrige et al. 2015; Křemenová et al. 2021; McPherson et al. 2022; Rapkin et al. 2016), and males may even adjust gifts based on recipient traits (Jarrige et al. 2015).

Donor fitness benefits have long been debated with little conclusive support for a single hypothesis (Arnqvist & Rowe 2005; Gwynne 2008; Lewis & South 2012; Lewis et al. 2014;

Sakaluk 2000; Thornhill 1976; Vahed 1998, 2007). Lewis and South (2012) present a revised

approach to describing nuptial gift benefits, suggesting eight non-exclusive benefits through which

donor fitness may be increased (increasing mate attraction, copulation success, insemination

success, sperm transfer, sperm storage, paternity share, female fecundity, or offspring fitness).

Behavioural modifications can serve many of these functions, often increasing copulation and

insemination success, sperm storage and transfer, and paternity share (Lewis & South 2012;

Gwynne 2008; Arnqvist & Rowe 2005). .

The presence of beneficial nuptial gifts does not necessarily signal an interaction lacking

conflict (Arnqvist & Rowe 2005; Gwynne 2008; Lewis & South 2012; Sirot et al. 2015). Mating

interactions can be costly to females and males – direct physical or physiological harm, resistance

costs, investment in gametes or nuptial gifts, or simple energy expenditure and predation risks

can each render mating a potentially costly encounter (Arnqvist & Rowe 2005; Boulton et al. 2018;

del Castillo & Gwynne 2007; Kokko & Jennions 2014; Lehtonen et al. 2012; Małek et al. 2019;

Perry & Tse 2013; Prokop & Okrouhlik 2021; Sentenská et al. 2020; Tatarnic et al. 2014). The

costs of producing or procuring nuptial gifts often lead to selective pressures to maximise gift

efficiency and efficacy through various strategies, including deceiving females or plastically

altering gifts in response to female quality (Albo et al. 2019; Beyer et al. 2021; Ghislandi et al.

2018; Martínez-Villar et al. 2023; Pandulli-Alonso et al. 2021; Solano-Brenes et al. 2021;

Stålhandske 2002).

Trade-offs between reproductive tactics, such as those between nuptial gifts and extended

clasping or deception, have been increasingly documented across arthropod species (del Castillo

& Gwynne 2007; Lehmann et al. 2016; Liu et al. 2015; Miller et al. 2019; Simmons & Emlen 2006;

Simmons et al. 2017; South et al. 2011; Yamane et al. 2010). Vahed et al. (2014) demonstrate

this, providing convincing evidence that prolonged clasping time can function as a substitute for

nuptial gifts. This has subsequently been shown in other orthopterans, suggesting that these

nuptial gifts function in some capacity to increase sperm transfer (Gwynne et al. 2008; Lehmann

et al. 2016; Vahed et al. 2014). This is also evident in *Neopanorpa* scorpionflies, in which the sole

species that has lost nuptial gifts (*N. longiprocessa*) has evolved enhanced clasping structures

(Tong & Hua 2019; Zhong & Hua 2013). In this species, mating interactions are characterised by

intense female struggle while males attempt to clasp prior to intromission. *Leiobunum* Opiliones

show a highly similar pattern of ancestral nuptial gifts and derived nuptial gift reduction in

conjunction with enhanced clasping traits, although to date, behavioural studies that directly

compare multiple species are lacking, and to our knowledge this is the first study to do so.

Study Species

North American sclerosomatid harvesters (Opiliones: Sclerosomatidae), also known as

"daddy-longlegs", form a widespread family of arachnids with a mating system ideal for studying

the role of nuptial gifts in reproductive behaviour (reproductive biology reviewed in Machado &

Burns 2023). *Leiobunum* species possess highly variable genitalia in both females and males,

with genital morphology being one of the primary methods of species identification in both sexes

(Shultz 2018). All *Leiobunum* females have a moveable, "hinged" genital operculum that covers

their genital chamber and must be opened prior to intromission (Shultz & Pinto-da-Rocha 2007;

Shultz 2018). Some species have additional pregenital barriers and sclerotization on the

operculum which engages an interior sternum, effectively locking the genital opening, but these

only occur in species which have also lost nuptial gift sacs (Burns et al. 2013; Shultz 2018).

Moveable female barriers are often indicative of sexual conflict, and Eberhard (2010) notes that

these barriers may be particularly likely to occur in systems undergoing sexually antagonistic

coevolution (though they are not entirely impossible under cryptic female choice). Species in

which males have lost nuptial gift sacs also have unique physical and mechanical penis

properties, with some species possessing robust penises capable of applying significant force to

female structures and other species occupying an intermediary state (Burns & Shultz 2015, 2016).

Burns and colleagues have described this repeated correlated evolution of species-specific

female and male reproductive morphology, revealing two primary reproductive syndromes (Burns

et al. 2012, 2013): 1) An ancestral, sacculate state in which males have terminal nuptial gift sacs

on their penises and females do not have pregenital barriers; and 2) a derived, nonsacculate state

in which males do not have terminal nuptial gift sacs on their penises and females have pregenital

barriers.

Both sacculate and nonsacculate *Leiobunum* harvesters produce nuptial gifts to some

degree, although the presence of terminal nuptial gift sacs increases the quantity transferred to

females during mating interactions and allows for precopulatory transfer (Machado & Burns 2023;

Machado & Macías-Ordóñez 2007). Sacculate females receive nuptial gift fluid before and during

copulation, potentially allowing for better precopulatory assessment of males (as expected under

cryptic female choice; Eberhard 1996), while nonsacculate females can only receive it after

committing themselves to copulation (Figure 1). In addition to these morphological differences,

sacculate and nonsacculate species also differ in the biochemistry of their nuptial gifts; sacculate

species have nuptial gifts with significantly higher proportions of essential amino acids compared

to nonsacculate gifts, and gift composition corresponded to the presence/absence of gift sacs

rather than phylogeny (Kahn et al. 2018). Essential amino acids are those that cannot be

produced by an organism and must come from their food, likely making them more costly for a

male to include in a gift (Kahn et al. 2015). These are also likely particularly valuable for females

to receive due to their contribution towards oogenesis (Jarrige et al. 2015; Kahn et al. 2018;

Wheeler 1996). This transition from high- to low-value nuptial gifts correlates with morphological

transitions between sacculate and nonsacculate states and is consistent with the expectations of

a system high in sexual conflict, in which males reduce investment in a highly-costly gift in favour

of coercion. Alternatively, it is possible that the reduction of nuptial gifts is instead related to a shift

in female preferences, with nonsacculate females evaluating males on behavioural cues rather

than biochemical or nutritional cues (Eberhard 1996).

The *L. vittatum* and *L. calcar* species-groups have species with pedipalps that are

structurally enhanced to facilitate more effective clasping (Burn et al. 2013; Ingianni et al. 2011).

Notably, these exaggerated features only occur in nonsacculate species with female pregenital

barriers, though they do not contact these structures during mating. Enhanced-clasping pedipalps

evolved independently in these groups, and in all instances this occurred after the transition to a

nonsacculate state (Burns et al. 2013). These clasping-structure modifications may allow males

to clasp and mate with females for longer than is ideal for the female (and therefore circumventing

potential female preferences), and longer intromission periods are highly associated with

increased sperm transfer, mate guarding, and sexual conflict in a number of species (Arnqvist &

Rowe 2005). Alternatively, the reduction of nuptial gifts in nonsacculate species may be

compensated for with an extended period of evaluation and interaction that is facilitated by better

clasping structures, and clasping alone is not necessarily indicative of conflict. Critical information

may be conveyed during this process, allowing an organism to evaluate a potential mate. Genital

clasping by males may provide stimulation to females (Eberhard 2010), and some species with

exaggerated genital clasping structures do not necessarily have high levels of sexual conflict

(Briceño & Eberhard 2009; Briceño et al. 2015; Myers et al. 2016; Peretti & Aisenberg 2015).

However, non-genital clasping structures, such as the pedipalps of Opiliones, are potentially more

indicative of sexual conflict (Arnqvist & Rowe 2005; Rowe & Arnqvist 2012).

Leiobunum vittatum (Say 1821)

B

♂

Precopulation

♀

C

Unreceptive female
angling body

D

Copulation   *Leiobunum verrucosum* (Wood 1868)

Figure 1: Primary mating phases and body orientations of *Leiobunum spp.* **A)** Male (left) *L.*

vittatum clasping female (right) precopulation. Photo courtesy of Joe Warfel. **B)** Precopulatory

phase, during which consumption of precopulatory nuptial gift (when present in the species)

occurs. **C)** Precopulatory posture of an unreceptive female. Female's anterior is angled

downwards to prevent male access to the ventrally located genital operculum. **D)** Copulatory

phase, during which consumption of pericopulatory nuptial gift occurs. **B-D)** Images are semi-

diagrammatic and the legs are not pictured to increase clarity (*L. verrucosum*, adapted from

Figure 1 in Burns et al. 2013).

Although there is morphological and chemical evidence indicative of a potential shift
towards increased conflict, comparative behavioural evidence in the genus is limited. Fowler-Finn
and colleagues (2014, 2018, 2019) have thoroughly documented the mating behaviour of the
nonsacculate *L. vittatum* as well as several other species, but no direct, comparative, behavioural
work between sacculate and nonsacculate species has been conducted. The presence of two
nuptial gifts in some *Leiobunum* species is unique, and the multiple independent transitions to a
derived, nonsacculate state provide an ideal system for studying the role of behaviour in the
reduction of nuptial gifts and whether sexual conflict may compensate for this change. In this
study, we tested whether nonsacculate leiobunine species displayed significantly different
reproductive behaviour than sacculate *Leiobunum* species. We hypothesised that the reduction
of nuptial gifts in nonsacculate species of *Leiobunum* would correspond with significantly different
mating behaviour when compared to sacculate species. Further, we expected nonsacculate
males to compensate for this reduction in gift value with increased coercive behaviour and overall
higher levels of sexual conflict. Alternatively, behavioural changes may be driven by shifts in
female preferences and mate assessment pathways. To test this, we performed mating trials
using two sacculate and two nonsacculate *Leiobunum* species and compared the intensity and
frequency of ostensibly antagonistic behaviour to search for differences between the two
reproductive syndromes.

Methods

<u>Study Species</u>

Leiobunine harvesters have a broad, Holarctic distribution and require extensive
taxonomic revision (Hedin et al. 2012), however, the "late-season" (defined as overwintering as
eggs) leiobunines of eastern North America form a well-supported monophyletic group (Burns et
al. 2012). We therefore restricted our study species to this clade, within which there are three
major phylogenetically-supported groups of harvesters. From these we selected one

representative species per group, as well as one species which was not a member of any major

group (Table 1; see Figure A1 for the full phylogeny).

Species	Species-group	Penis Morphology	Pedipalp Morphology	Nuptial Gift Chemistry	Behavioral Predictions
L. euserratipalpe	*calcar*	Nonsacculate	Unmodified pedipalps	Low investment	High conflict
L. vittatum	*vittatum*	Nonsacculate	Male pedipalps specialized for clasping	Low investment	High conflict
L. bracchiolum	*politum*	Sacculate	Unmodified pedipalps	High investment	Low conflict
L. aldrichi	-	Sacculate	Unmodified pedipalps	High investment	Low Conflict

Table 1: Morphological and biochemical phenotypes of the four species in this study, as well as

overall behaviour expected in trials (Burns et al. 2012; Kahn et al. 2018; Shultz 2018). High conflict

would be indicated by reduced latency to make a clasping attempt, elevated contact duration,

elevated time within one leg span, elevated clasping duration, elevated guarding duration,

elevated female body angle, reduced association distance, and reduced pursuit angle. Low

conflict would be indicated by the inverse of these metrics.

In sacculate species, males first use their pedipalps to clasp females face-to-face by the

coxae of leg II (Figure 1A and D). The male then extends its penis to the female's mouth, allowing

the female to consume the endogenously-produced precopulatory nuptial gift that accumulates in

the terminal sacs after being produced by accessory glands (Figure 1B). The male redirects its

penis to the female's genital operculum, which the female then opens to allow intromission. At

this stage, the female consumes the endogenously-produced, pericopulatory nuptial gift directly

from the male's papilla as its mouth aligns with male's haematodocha (Figure 1D; Burns et al.

2012, 2013, Machado & Macías-Ordóñez 2007). In nonsacculate species, this behavioural

sequence proceeds much the same, save for two notable differences: 1) the precopulatory nuptial

gift does not occur, and 2) the male may use its penis to pry the female genital operculum open

should the female resist copulation (Burns et al. 2012, 2013).

Study Populations

We collected harvesters as juveniles in the spring and summer of 2021, 2022, and 2023.

The exact timing and location of collection depended upon the species, as each species matures

at different times, and some are restricted in their geographic range across central Maryland.

Leiobunum vittatum were collected in June and July of 2021 at Rockburn Branch Park

(39.219471, -76.761198) and the University of Maryland, Baltimore County (39.247397, -

76.710568); *L. aldrichi* were collected in June and July of 2021 at Sugarloaf Mountain (39.261630,

-77.395560) and Plummer's Island (39.261630, -77.395560); and *L. euserratipalpe* were collected

in April and May of 2022 at Schooley Mill Park (39.166325, -76.961830); and *L. bracchiolum* were

initially collected in May and June of 2021 at Rockburn Branch park. Due to a low rate of

successful clasps, additional *L. bracchiolum* were collected in May and June of 2023 at the same

location to ensure sufficient data for analyses which required successful clasps.

Following collection, we separated harvesters by species and sex and housed them communally

at 24° C, 80% humidity, and 12:12 h light:dark periods in 37 cm x 22 cm x 24.5 cm plastic terraria.

Each terrarium contained ~ 2 cm of Eco Earth coconut fibre substrate (Zoo Med), petri dishes

with TetraMin Select tropical flakes fish food (Tetra), moist cotton balls, and leaves, sticks, and

bark gathered from the collection locations for shelter. All terraria were moistened daily and

checked for mould, and all were cleaned weekly. Upon maturation, we maintained harvesters in

similar conditions until mating trials were complete, at which point we housed them communally

by sex until death.

<u>Behaviour</u>

Behavioural trials consisted of pairing male and female conspecifics and recording mating

behaviour. We conducted all behavioural trials in a 29.21 cm x 29.21 cm acrylic arena lined with

white cardstock to reduce reflections and allow for easy cleaning. The arena was housed in a

room kept at 24º C and illuminated with white light (leiobunine Opiliones have poor vision and

lighting is unlikely to impact mating behaviour (Machado & Macías-Ordóñez 2007; Shultz & Pinto-

da-Rocha 2007) and we video recorded interactions dorsally and laterally with Microsoft LifeCam

Studio cameras. Following the methods used in studies by Fowler-Finn and colleagues (2014,

2018, 2019), we conducted trials as follows: We first sterilised the arena with 95% ethanol before

placing a male and female harvester in the arena. Initially, harvesters were isolated for 5 minutes

under transparent plastic containers to allow acclimation to the arena without contact with one

another. Following acclimation, we removed the containers and the trial began. Trials lasted for

either 20-minutes or until the mating interaction ended. Interactions were considered concluded

when contact was broken between individuals and the male made no attempt to chase the female.

If the interaction had not ended by the 60-minute mark, the trial was concluded.

We aimed to complete 20 trials per species, though the exact number per species varied

due to unexpected deaths and exclusions due to inactivity (Appendix Table A1). Trials were

included in analyses if the male made an attempt to hook its pedipalps around the female's coxae

within the first 20 minutes. If no mating attempt was made at this point, we removed the harvesters

from the arena, excluded them from future trials, and did not include the trial in analyses. The

majority of males (86.36%) made an attempt to clasp within the first 20 minutes, meaning only 12

of 88 potential trials were excluded due to inactivity. The majority of interactions (86.67%) had

concluded before the imposed 60-minute limit, with only 11 of 75 trials artificially ended.

Interestingly, seven of these 11 trials were with *L. euserratipalpe* pairs.

<u>Video Analyses</u>

285   We used the automated tracking program TRex (Walter & Couzin 2021) to track individual

286 identities and coordinates to calculate the distance between individuals, total time individuals were

287 within one male leg span (measured using the longest leg length specific to the male in each trial),

288 and total clasping duration (Table 2). We additionally explored a method to assess the degree to

289 which males followed females between first contact and first mating attempt. We used the centroid

290 coordinates to first calculate male and female trajectory vectors and magnitudes before using

291 these to derive the angle between vectors. This metric describes the "pursuit angle" between male

292 and female movement; values closer to 0 degrees indicate the individuals were moving in the

293 exact same direction, which we would expect if the male were chasing the female along parallel

294 vectors (Table 2). Conversely, values closer to 180 degrees indicate the individuals were moving

295 in the exact opposite direction. Some behaviours, particularly those occurring on the ventral side

296 of the harvesters or those involving legs, could not be reliably scored with TRex and we instead

297 used manual scoring to record contact time, female body angle, instances of investigatory leg

298 tapping bouts and instances of females fleeing males (Table 2).

Metric (Category)	Definition	Expectations under High Conflict
Clasp, attempted	Male attempts to hook its pedipalps behind the female's leg II coxae.	-
Clasp, successful	Male successfully hooks its pedipalps behind the female's leg II coxae.	-
Clasping time	Time during which the male maintains a clasp with its pedipalps hooked behind the female's leg II coxae.	Increased: The length of clasping may indicate conflict if males continue to clasp females beyond the female's ideal duration. Increased clasping duration is often associated with increased sperm transfer, mate guarding, and antagonistic interactions (Arqnvist & Rowe 2005).
Contact time, pre-attempt	Any contact between individuals before a successful clasp has occurred.	Decreased: Longer pre-attempt contact times may indicate solicitous, courtship-like behaviour because it provides individuals with a greater opportunity to assess a potential mate. Shorter times provide less opportunity for assessment and courtship, and so would be expected (Dmitriew & Blanckenhorn 2012; Fowler et al. 2022)
Contact time, total	Any contact between individuals throughout the entire trial.	Increased: *Leiobunum* harvesters primarily rely on tactile and chemical senses and contact between individuals is an important communication channel. Increased association time between males and females can reduce female fitness in many species and may therefore be indicative of conflict. Alternatively, females may benefit from increased time to assess a potential mate and from beneficial male guarding (Bokides et al. 2012; Iglesias-Carrasco et al. 2019).
Distance	Distance between the female and male body centroid post-initial contact.	Decreased: An automated analog of total contact time, see above.
Female body angle	Maximum angle formed by the female's genital operculum and the substrate during a male clasping attempt or success.	Increased: The female genital opening is ventrally located, and angling the anterior portion of its body downwards may make it more difficult for the male to access the genital operculum. Genital repositioning has been documented in several other arthropods with obligate reproductive clasping (Burke et al. 2015; Golov et al. 2018; Maroni et al. 2023).
Flee count	Instances of the female rapidly moving away from male contact.	Increased: Upon contact, unreceptive females may rapidly move away from the male to avoid reproduction attempts.
Guarding time	Any instance of contact lasting greater than 5 seconds post-successful clasp. Subsequent clasps were scored as such, rather than as guarding.	Increased: Mate guarding is strongly associated with sexual conflict, allowing males to prevent remating and ensure paternity. Though guarding is not typically beneficial for females, beneficial mate guarding has been documented, albeit rarely, and could function to prevent fitness-reducing harassment from other males (Alcock 1994; Arnqvist & Rowe 2005; Sherman 1983).
Latency to attempt	Time between the initial contact and the first clasping attempt.	Decreased: Shorter latency allows for less mate assessment and fewer courtship behaviours and has been associated with high conflict in other arthropods (Dmitriew & Blanckenhorn 2012; Fowler et al. 2022).
Pursuit angle	Angle formed by the male and female velocity vectors post-initial contact.	Decreased: Angles near 0 degrees indicate the female is moving away from the male while the male follows and is indicative of the male pursuing a potentially unreceptive female. Angles near 180 degrees indicate individuals are moving towards one another, which would indicative a receptive female (Bokides et al. 2012; Iglesias-Carrasco et al. 2019).
Tap bout count	Bouts of investigatory leg taps performed during an interaction.	Decreased: *Leiobunum* Opiliones primarily rely on tactile and chemical sensing, performing investigatory taps with their antenniform second pair of legs. Tapping may be solicitous or courtship-like if it provides an assessment signal between individuals (Shultz & Pinto-da-Rocha 2007; Wulff et al. 2018).
Time within 1 leg span	Time during which the distance between male and female body centroids was less than the length of the male's longest leg.	Increased: An automated analog of total contact time, see above.

Table 2: Behavioural metrics recorded during all trials. Rationale for each metric is noted in text

with supporting citations.

To ensure TRex performed accurately and matched manual tracking expectations, we

additionally scored all trials with JWatcher (Blumstein & Daniel 2007). For all attempted-clasp

trials, we scored the total time the harvesters were in contact, the latency between first contact

and first mating attempt, the number of female fleeing instances, and the number of "tapping

bouts" by the male (Table 2). For successful-clasp trials, we additionally scored the time between

first mating attempt and first successful clasp, the total time the individuals were in contact, the

total time the male clasped the female as a proxy for intromission time (as per Fowler-Finn et al.

2014, 2018, 2019), and the total time the male guarded the female post-copulation (Table 2). In

leiobunine harvesters, mate guarding consists of the male keeping in constant contact with at

least one leg grasping the female's leg; because they primarily rely on tactile sensing using

antenniform legs and their vision is particularly poorly-developed, this contact is likely required

(Machado & Macías-Ordóñez 2007; Shultz & Pinto-da-Rocha 2007; Fowler-Finn et al. 2018,

2019). We considered post-copulatory contact lasting greater than 5 seconds to constitute mate

guarding (Fowler-Finn et al. 2014, 2018, 2019).

Finally, in all attempted-clasp trials, we used the image analysis program FIJI-ImageJ

(Schindelin et al. 2012) to measure the maximum angle the female's genital operculum formed

with the substrate of the arena (Figure 1C). Due to the degree to which some mating pairs moved

during their interactions, we could not effectively automate this metric to obtain a continuous

measure of the angle, thus we elected to use the maximum angle reached during a clasp attempt

(Table 2).

<u>Statistical Analyses</u>

Due to our selected study species being closely related, we first tested for phylogenetic

signal across all our behavioural metrics. Previous studies have not found evidence for signal

among morphological traits (Burns & Shultz 2015); and significant evidence suggests that

behavioural traits are among the least likely to display patterns of phylogenetic signal because of

their lability (Blomberg et al. 2003). We imported our behavioural data to R (v4.3.1; R Core Team

2023) and calculated Pagel's lambda (Pagel 1999) using the *phytools* package (Revell 2012). We

used a pruned version (consisting of our four study species) of a maximum clade credibility tree

constructed from mitochondrial and nuclear sequences by Burns et al 2013. We found extremely

low levels of phylogenetic signal (Likelihood ratio test versus Brownian motion: All $P > 0.6$;

Appendix Table A2) and after consideration with previous evidence (Burns & Shultz 2016, 2015;

Blomberg et al. 2003) proceeded with frequentist statistical analyses.

Leiobunine harvesters primarily rely on tactile senses; sweeping and tapping their

antenniform second pair of legs to investigate their surroundings (Shultz & Pinto-da-Rocha 2007;

Willemart & Hebets 2012). We therefore evaluated normality using the Shapiro-Wilk test and

followed this with Kruskal-Wallis tests for nonparametric data to determine whether the individuals

used in our study differed significantly between species in the number of sensory legs and total

legs. Because Opiliones can autotomize their legs, severing them to escape potential predators

(Gnaspini & Hara 2007), not all individuals in the study had the same number of total legs or

sensory legs. We incorporated this variability by constructing multiple linear regression models

with behavioural metrics as response variables (Table 2) and leg count metrics (number of male

legs, number of male sensory legs, number of female legs, and number of female sensory legs)

as predictor variables. The results of these multiple linear regression models allowed us to

determine whether the loss of legs impacted behavioural sexual conflict.

For all behavioural metrics, we used the Shapiro-Wilk test for normality and found at least

one species in all but one metric to be non-normally distributed. Due to this, we used the Kruskal-

Wallis test to compare the four species for all metrics except the average distance between

individuals. For both attempted- and successful-clasp trials this metric was normally distributed

after a logarithmic transformation, and so these data were analysed with a one-way ANOVA after

transformation. We followed any significant groupwise comparisons with planned pairwise

comparisons using Dunn's multiple comparisons test or Šídák's multiple comparisons test, where

appropriate, between sacculate and nonsacculate species (*L. bracchiolum x L. vittatum*; *L.*

bracchiolum x L. euserratipalpe; *L. aldrichi x L. vittatum*; and *L. aldrichi x L. euserratipalpe*).

Following inspection of several metrics (pursuit angle, attempt latency, clasping duration, and

maximum female body angle), we decided to compare the two nonsacculate species (*L. vittatum*

and *L. euserratipalpe*) to one another with post-hoc Dunn's tests, corrected for multiple

comparisons.

We concluded our statistical analyses with multiple logistic regression models to detect

which behavioural metrics best predict clasping success. We built models with success/failure to

secure a clasp as the response variable and the behavioural metrics as the predictor variables.

We constructed a total-species model using all data, a sacculate-species model using only *L.*

bracchiolum and *L. aldrichi* data, and species models using species-level data. We could not build

nonsacculate-species or species-level models for *L. vittatum* and *L. euserratipalpe* due to each

species having only one failed clasp attempt, thus leading to perfect separation in our models.

<u>Ethical Note</u>

We collected *L. vittatum*, *L. euserratipalpe*, *L. aldrichi*, and *L. bracchiolum* from populations

throughout their natural range in central Maryland. The Washington Biologists' Field Club

provided collection permission for Plummer's Island, while collection permits were not required

for any other location. All source populations used for collections were well-established, with the

species in question being clearly abundant and unlikely to be affected by our collections. The

harvesters were then held in the lab following ASAB/ABS guidelines for the treatment of animals

used in behavioural research. Harvesters were reared collectively whenever possible in

accordance with their gregarious behaviour in the field. Behavioural trials were generally not

harmful to individuals – females experienced harassment from males during trials, but it was likely

far less frequent than harassment experienced in natural conditions. Following trials, individuals

were maintained in the laboratory until natural death, typically within one to two months.

Results

<u>Overall Contact and Association Metrics</u>

We found one significant difference when comparing the number of legs between species

(total legs, female *L. bracchiolum* versus female *L. aldrichi*; Appendix Table A3), however, this

metric was not a significant predictor in our multiple linear regression models. Due to this, we did

not incorporate leg data into the remainder of the analyses. We found significant differences

between species for all contact and association metrics, although the specific relationships varied.

Overall, females and males of nonsacculate species (*L. vittatum* and *L. euserratipalpe*) associated

significantly more closely than females and males of sacculate species (*L. bracchiolum* and *L.*

aldrichi; Figure 2 & A2).

In attempted-clasp trials, nonsacculate males and females were in physical contact for

significantly longer than sacculate males and females. In these trials, median contact time was

between three and four times longer in nonsacculate species than in sacculate species (Figure

2A). When comparing only successful-clasp trials, both nonsacculate species retained

significantly longer contact time when compared to *L. aldrichi*, however, *L. bracchiolum* was not

significantly different (Figure 2B). When comparing the number of bouts of investigatory tapping

performed during a mating trial, we found that *L. vittatum* did so significantly less than *L.*

bracchiolum in both attempted-clasp ($P = 0.001$) and successful clasp trials ($P = 0.005$; Figure 3).

Surprisingly, we also found that *L. vittatum* had significantly fewer bouts than its fellow

nonsacculate in both trial categories (attempted-clasp trials: $P = 0.0341$; successful clasp trials:

$P = 0.0221$), and that *L. bracchiolum* had significantly more tapping bouts than its fellow sacculate

in attempted-clasp trials ($P = 0.0208$; Figure 3). There were no clear patterns when comparing

flee instances between species, though *L. bracchiolum* females rarely fled males and had a

median flee count of 0 instances (IQR 0 - 2 instances), resulting in L. bracchiolum having a

significantly lower flee count than *L. aldrichi* in attempted-clasp (*P* > 0.0001) and successful clasp

trials (*P* = 0.0121; Figure A3).

Figure 2: Association metrics between male and female pairs. Boxes represent medians plotted

with IQR, symbols represent individual trial values, and whiskers correspond to the minimum and

maximum trial values. The results from group comparison tests are shown in the upper right-hand

corners, while significant results from post-hoc multiple comparisons tests are noted by red

brackets with asterisks (* = *P* < 0.05; ** = *P* < 0.01). The simplified phylogeny indicates

reproductive morphology; red tips indicate nonsacculate species and black tips represent

sacculate species (phylogeny adapted from Burns et al. 2013). **A)** Total contact time in seconds

between males and females during attempted-clasp trials. **B)** Total contact time in seconds

between males and females during successful-clasp trials.

Figure 3: Instances of investigatory tapping bouts. Boxes represent medians plotted with IQR, symbols represent individual trial values, and whiskers correspond to the minimum and maximum trial values. The results from group comparison tests are shown in the upper right-hand corners, while significant results from post-hoc multiple comparisons tests are noted by red brackets with asterisks (* = P < 0.05; ** = P < 0.01; *** = P < 0.001). The simplified phylogeny indicates reproductive morphology; red tips indicate nonsacculate species and black tips represent sacculate species (phylogeny adapted from Burns et al. 2013). **A)** Instances of tapping bouts between males and females during attempted-clasp trials. **B)** Instances of tapping bouts between males and females during successful-clasp trials.

Using automated tracking, we found that in both attempted- and successful-clasp trials males were within one leg span of females significantly more often in nonsacculate species than in the nonsacculate *L. aldrichi*. This matched the contact patterns found in the successful-clasp trials manual scoring, however, for attempted-clasp trials we failed to find significance comparing *L. bracchiolum* to nonsacculates. This discrepancy in contact time is likely due to using the body centroids for position in TRex – we only considered females to be within the male's leg span if the female's body centroid was within the given male leg span, meaning leg-leg contact may occur without being scored. During attempted-clasp trials, nonsacculate females and males were also

significantly closer to one another when compared to *L. aldrichi*, though this pattern did not hold

for successful-clasp trials.

<u>Precopulatory Phase</u>

Across all species, we found significant differences in the latency between first contact

and first clasping attempt (Figure 4). *Leiobunum euserratipalpe* median attempt latency was 6.0

s (IQR 3.0 - 10.0 s), significantly lower than *L. vittatum* (76.5 s, IQR 20.75 - 605.3 s) and *L. aldrichi*

(128.0 s, IQR 7.25 - 462.0 s), but was not significantly different from that of *L. bracchiolum* (18.0

s, IQR 4.0 - 327.5 s). All species displayed marked variation in latency to attempt a clasp except

L. euserratipalpe, which had a standard deviation of 10.72 seconds (*L. bracchiolum* SD = 230.7

s; *L. aldrichi* SD = 363.5 s; *L. vittatum* SD = 330.4 s). The longest attempt latency for *L.*

euserratipalpe was just 41 s, with all but three males making an attempt within 10 s of their initial

contact with a female (Figure 4).

Figure 4: Latency in seconds between first contact and first clasp attempt during successful-

clasp trials. Boxes represent medians plotted with IQR, symbols represent within-trial values, and

whiskers correspond to the minimum and maximum trial values. The result from a Kruskal-Wallis

test is shown in the upper right-hand corner, while significant results from Dunn's multiple

comparisons tests are noted by red brackets with asterisks (** = *P* < 0.01). The simplified

phylogeny indicates reproductive morphology; red tips indicate nonsacculate species and black

tips represent sacculate species (phylogeny adapted from Burns et al. 2013).

Finally, we found significant differences in pursuit angle between species for attempted-

and successful-clasp trials. In trials with attempted clasps, the median pursuit angle for *L. vittatum*

(82.64 degrees, IQR 82.05 - 85.45 degrees) was significantly smaller than that of *L. bracchiolum*

(87.75 degrees, IQR 85.79 - 88.86 degrees) (Figure 5A). *Leiobunum vittatum* also had a

significantly smaller angle than that of *L. euserratipalpe* (88.71 degrees, IQR 85.67 - 89.92

degrees) despite both being nonsacculate. Though in successful-clasp trials the *L. vittatum*

pursuit angle (82.30 degrees, IQR 82.00 - 84.66 degrees) was similar to both sacculate species,

it remained significantly smaller than *L. euserratipalpe* (89.04 degrees, IQR 85.53 - 89.89

degrees) (Figure 5B).

Figure 5: Angle between male and female trajectories between first contact and first clasp

attempt. Boxes represent medians plotted with IQR, symbols represent within-trial means, and

whiskers correspond to the minimum and maximum trial means. The simplified phylogeny

indicates reproductive morphology; red tips indicate nonsacculate species and black tips

represent sacculate species (phylogeny adapted from Burns et al. 2013). The results from

Kruskal-Wallis tests are shown in the upper right-hand corner, while significant results from

Dunn's multiple comparisons tests are noted by red brackets with asterisks (** = P < 0.01). **A)**

Attempted-clasp trials. **B)** Successful-clasp trials.

<u>Copulatory Phase</u>

We found significant differences when comparing the overall clasping duration of each

species using a Kruskal-Wallis test, although there were no clear patterns between sacculate and

nonsacculate species (Figure 6). During successful-clasp trials *L. vittatum* males clasped females

significantly longer than either sacculate species, with a median time of 676.8 s (IQR 310.6 -

1271.0 s) compared to 118.7 s (IQR 98.47 - 179.1 s) for *L. bracchiolum* and 260.5 s (IQR 17.11 -

662.8 s) for *L. aldrichi* (Figure 6). Interestingly it appeared that *L. euserratipalpe* clasped far less

than expected, and an unplanned post-hoc test comparing the species to *L. vittatum* revealed that

L. euserratipalpe clasped for significantly less time than its nonsacculate counterpart (Figure 6).

The median clasping duration for *L. euserratipalpe*, 155.7 s (IQR 87.22 - 370.7 s), was more

aligned with the clasping durations of sacculate species, and was approximately half that of *L.*

aldrichi, contrary to our predictions.

Figure 6: Time in seconds that males clasping females during trials with successful clasps. Boxes represent medians plotted with IQR, symbols represent within-trial values, and whiskers correspond to the minimum and maximum trial values. The result from a Kruskal-Wallis test is shown in the upper right-hand corner, while significant results from Dunn's multiple comparisons tests are noted by red brackets with asterisks (* = $P < 0.05$; ** = $P < 0.01$). The simplified phylogeny indicates reproductive morphology; red tips indicate nonsacculate species and black tips represent sacculate species (phylogeny adapted from Burns et al. 2013).

Comparing the degree to which females angled their body (which blocks genital operculum access, Figure 1C) revealed significant differences across all species, but these analyses resulted in mixed results when comparing sacculate and nonsacculate species. *Leiobunum vittatum* was again significantly different from its nonsacculate counterpart, *L. euserratipalpe*, and the sacculate *L. aldrichi*, but it was not significantly different from *L. bracchiolum*. In all trials with attempted clasps, *L. vittatum* females angled their bodies to a median of 46.39 degrees (IQR 33.54 - 90.0 degrees) with four females reaching a full 90 degrees perpendicular to the substrate (Figure 7A).

Leiobunum euserratipalpe females displayed a median angle of 0 degrees + (IQR 0 - 39.54

degrees) and rarely angled their bodies, with only six of 15 females doing so and only one female

reaching 90 degrees. While *L. bracchiolum* and *L. aldrichi* females angled their bodies more often

than *L. euserratipalpe* females, no females reached 90 degrees, and their median angles were

28.98 degrees (IQR 0 - 49.35 degrees) and 25.5 degrees + (IQR 0 - 42.84 degrees) respectively.

The contact time (between first touch and first clasp attempt) and the total time within one leg

span were significant predictors of clasping success for both the total-species full model and the

total-species reduced model (Table 3). Species-level models for *L. bracchiolum* and *L. aldrichi*

were non-significant, while nonsacculate, *L. vittatum*, and *L. euserratipalpe* models were not

possible due to few failed clasping attempts.

Figure 7: Maximum angle females formed between genital operculum and substrate. Boxes

represent medians plotted with IQR, symbols represent within-trial values, and whiskers

correspond to the minimum and maximum trial values. The simplified phylogeny indicates

reproductive morphology; red tips indicate nonsacculate species and black tips represent

sacculate species (phylogeny adapted from Burns et al. 2013). The results from Kruskal-Wallis

tests are shown in the upper right-hand corner, while significant results from Dunn's multiple

comparisons tests are noted by red brackets with asterisks (* = *P* < 0.05; ** = *P* < 0.01). **A)**

Attempted-clasp trials. **B)** Successful-clasp trials.

Model/Factor	AICc	Pseudo-R²	Estimate	SE	|Z|	P-value
Total-species full model	72.95	0.388	-	-	-	-
Intercept	-	-	-3.247	8.348	0.389	0.6973
Initial latency	-	-	-0.002	0.002	1.125	0.2607
Pre-attempt contact time	-	-	**-0.004**	**0.002**	**2.482**	**0.0131**
Average distance	-	-	0.114	0.093	1.226	0.2200
Time within 1 leg span	-	-	**0.003**	**0.001**	**2.257**	**0.0240**
Female body angle	-	-	0.032	0.017	1.885	0.0594
Pursuit angle	-	-	0.016	0.095	0.168	0.8665
Flee count			0.606	0.438	1.385	0.1661
Tap count			-0.216	0.125	1.721	0.0853
Total-species reduced model	68.87	0.319	-	-	-	-
Intercept	-	-	-1.113	0.645	1.725	0.0845
Pre-attempt contact time	-	-	**-0.003**	**0.001**	**2.685**	**0.0072**
Time within 1 leg span	-	-	**0.002**	**7e-04**	**2.844**	**0.0045**
Female body angle	-	-	0.024	0.014	1.690	0.0910
Flee count	-	-	0.755	0.391	1.928	0.0538

Table 3: Multiple logistic regression modelling of clasping success with behavioural metrics as

predictor variables for the total-species full model and the total-species reduced model. Predictor

variables were iteratively removed beginning with the variable with the greatest *P*-value until no

variables with *P* > 0.1 remained. Significant predictor variables are in bold.

<u>Postcopulatory</u>

Finally, when analysing the duration of mate guarding – the primary indicator of

postcopulatory conflict in leiobunine harvesters – we found significant differences across species

(Figure 8). Nonetheless, there were few pairwise significant differences, potentially due to

significant intraspecific variation in all species except *L. aldrichi*. *Leiobunum aldrichi* males

guarded for a median time of just 59.0 s + (IQR 0 - 270.5 s), while *L. bracchiolum* and *L. vittatum*

males guarded for 474.0 s (IQR 0 - 857.0 s) and 351 s (IQR 18.0 - 1145.0 s), respectively (Figure

8). *Leiobunum euserratipalpe* males guarded females for significantly longer than *L. aldrichi* males

did, with its median guarding times being 1209.0 s (IQR 209.8 - 2919.0 s) (Figure 8). Although

male *L. euserratipalpe* tended to guard females for extended periods, they did not always do so

– five males guarded for less than 320.0 s, and two of these did not guard at all. Due to its broad

distribution of guarding duration, we did not have sufficient statistical power to resolve any

potential differences between *L. euserratipalpe*, *L. vittatum*, and *L. bracchiolum*, and further study

may allow the resolution of these differences.

Figure 8: Time in seconds that males guarded females during successful-clasp trials. Boxes

represent medians plotted with IQR, symbols represent within-trial values, and whiskers

correspond to the minimum and maximum trial values. The result from a Kruskal-Wallis test is

shown in the upper right-hand corner, while significant results from Dunn's multiple comparisons

tests are noted by red brackets with asterisks (** = $P < 0.01$). The simplified phylogeny indicates

reproductive morphology; red tips indicate nonsacculate species and black tips represent

sacculate species (phylogeny adapted from Burns et al. 2013).

Discussion

<u>Behavioural comparisons between sacculate and nonsacculate species</u>

Our results indicate that the nonsacculate harvesters *L. vittatum* and *L. euserratipalpe*

display significantly different mating behaviour when compared to the sacculate species *L. aldrichi*

and *L. bracchiolum* (Table 4). This variation in mating behaviour corresponds to the penis

morphology and reproductive syndromes of each species; *L. vittatum* and *L. euserratipalpe* lack

precopulatory nuptial gifts and possess antagonistic morphology, while *L. aldrichi* and *L.*

bracchiolum give precopulatory nuptial gifts and lack antagonistic morphology (Burns et al. 2013).

Across all mating trials, nonsacculate males repeatedly displayed significantly longer median

association times, were significantly closer to females, and performed fewer investigatory tapping

bouts (Figures 2, 3, & A2). Male *L. euserratipalpe* attempted mating clasps significantly faster

upon first touch than male *L. aldrichi* and *L. vittatum* (Figure 4), while male *L. vittatum* pursued

females more vigorously and female *L. vittatum* also displayed higher levels of resistance

behaviour (Figures 5, 7, & A3). Hallmark conflict behaviours such as extended mate clasping and

mate guarding, were not found to be elevated across both nonsacculate species, with instead *L.*

vittatum displaying extended clasping behaviour (Figure 6) and *L. euserratipalpe* displaying

extended guarding behaviour (Figure 8). Altogether, these data indicate that nonsacculate

harvesters display significantly higher levels of antagonistic behaviour, possibly to compensate

for the loss of precopulatory nuptial gifts and the reduction of pericopulatory nuptial gifts (Table

4).

Metric	*L. vittatum*	*L. euserratipalpe*	*L. bracchiolum*	*L. aldrichi*
Conflict prediction	High	High	Low	Low
Latency to 1st attempt	No Support	Support	No Support	Support
Pursuit angle	Support	No Support	Mixed	No Support
Contact time	Support	Support	Mixed	Support
Tap Count	Support	No Support	Support	No Support
Flee Count	No Support	Mixed	Mixed	No Support
Time within 1 leg span	Support	Support	No Support	Support
Pairwise distance	Mixed	Mixed	No Support	Mixed
Female body angle	Support	No Support	Mixed	Mixed
Clasping time	Support	No Support	Support	Support
Guarding time	No Support	Support	No Support	Support
Supporting Metrics	6-7 of 10	4-6 of 10	2-6 of 10	5-7 of 10

Table 4: Summary of our predictions for behavioural sexual conflict intensity and the degree to

which the results of each metric matched what we expected for each species (see Table 2 for

metric specific expectations). "Support" indicates that the species matched our predictions in both

attempted-clasp and successful-clasp trials; "Mixed" indicates metrics in which this occurred in

only one of the two trial categories (attempted-clasp and successful-clasp), and "No support"

indicates that the species did not match our expectations in either trial category.

With these results we provide the first direct behavioural comparison between sacculate

and nonsacculate leiobunine harvesters. Despite extensive evidence of a shift from a solicitous

suite of traits to an antagonistic suite, the role of behaviour had not yet been investigated with

respect to sexual conflict in this system. Previous studies have revealed various losses of

solicitous traits such as penile sacs, precopulatory nuptial gifts, and high essential amino acid

content in pericopulatory nuptial gifts (Burns et al. 2013; Kahn et al. 2018). Meanwhile, these and

other studies have also shown gains of antagonistic traits, such as enhanced penile prying force,

female reproductive barriers, and modified clasping structures (Burns et al. 2013; Burns & Shultz

2015, 2016; Karachiwalla et al. 2020). While these changes arose independently across

leiobunine harvesters, many have reached convergent states. Our results indicate that these

morphological and biochemical changes have been accompanied by marked behavioural

changes as well.

Due to leiobunine harvesters primarily relying on their antenniform legs for sensory cues

(Shultz & Pinto-da-Rocha 2007; Willemart & Hebets 2012) we expected that males would keep

females within their leg span throughout antagonistic mating interactions to facilitate control. In

this study, nonsacculate species maintained this state for two to three times longer than sacculate

species. The distance and contact time between individuals are not clear indicators of sexual

conflict in isolation – both can be expected to occur in solicitous and antagonistic interactions

(though short contact time would primarily indicate a lack of antagonism due to extended clasping

and guarding being hallmark characteristics of behavioural conflict; Arnqvist & Rowe 2005;

Bokides et al. 2012; Iglesias-Carrasco et al. 2019). However, in conjunction with the other metrics

in the study, including female body angle, clasping time, and guarding time, it seems more likely

to be associated with antagonism. Across mammals, reptiles, fish, and arthropods, many studies

have shown cohabitation with males or proximity to males can each impact female fitness

negatively through harassment (Arnqvist & Rowe 2005; Krupa et al. 1990; Kunz et al. 2021;

Pilastro et al. 2003; Plath et al. 2003; Rönn et al. 2006; Schlupp et al. 2001; Schütz & Taborsky

2005; Shine et al. 2005). In particular, even if copulation does not occur, close association with

males can lower female fitness, whether due to direct stress from male activity or from selecting

low-quality refuge instead of a high-quality habitat with males as an avoidance strategy (Bokides

et al. 2012; Iglesias-Carrasco et al. 2019). Furthermore, the exceptionally low latency between

first contact and first attempt in *L. euserratipalpe* might be driven by a variety of factors, although

many which might normally influence latency are controlled in this experiment, such as female

age or sexual history, male starvation, and the presence of male rivals (Burke & Bondurianksy

2018; Dmitriew & Blanckenhorn 2012; Fowler et al. 2022). Low latency has previously been

associated with high conflict (Blanckenhorn et al. 2020; Skwierzyńska & Plesnar-Bielak 2018),

and in concert with the elevated guarding duration in *L. euserratipalpe,* this low latency appears

to indicate a species with high conflict.

<u>Clasping, guarding, and conflict</u>

Extended clasping time (beyond what might be ideal for female fitness) is a classic

indicator of sexual conflict found in myriad arthropod and vertebrate systems (Arnqvist & Rowe

2005; Johns et al. 2009; Rowe & Arnqvist 2012; Rueda-Solano et al. 2022; Sigvardt et al. 2017).

Indeed, males of some species, including those in this study, are entirely unable to ensure

copulation when unable to clasp the female, providing significant motivation for unreceptive

females to prevent clasping (Machado & Macías-Ordóñez et al. 2007; Myers et al. 2016).

Extended clasping can function as a multifaceted advantage for males, the primary explanatory

hypotheses being the Ejaculate Transfer (ET) hypothesis (Edvardsson & Canal 2006; Mazzi et

al. 2009; Simmons 2002) and the Extended Mate Guarding (EMG) hypothesis (Alcock 1994;

Mazzi et al. 2009). In the ET hypothesis, males extend clasping time to maximise the removal of

sperm from previous males and the transfer of ejaculate materials such as sperm, seminal fluids,

and mating plugs (Arnqvist & Rowe 2005; Edward et al. 2015; Mazzi et al. 2009; Macías-Ordóñez

et al. 2010; Simmons 2002; Townsend et al. 2019; Vahed et al. 2014). In the EMG hypothesis,

males instead maximise clasping time as a function of mate guarding in order to reduce the

potential for female remating (Mazzi et al. 2009; Zatz et al. 2011). In both these nonexclusive

hypotheses, the ultimate result may be quite similar to those achieved by nuptial gifts (Lewis &

South 2012). Mating plugs, sperm removal, and clasping as mate guarding have all been

observed in tropical harvesters (Machado & Burns 2023; Macías-Ordóñez et al. 2010; Townsend

et al. 2015, 2019; Zatz et al. 2011), and so the behavioural conflict over clasping time in these

North American harvesters might be attributable to either hypothesis or some combination of the

two.

Female harvesters often angle their body anterior-down into the substrate, blocking

access to their genital operculum. This potentially makes it more difficult for males to achieve or

maintain a clasp, though this does not seem to be particularly effective for females in our study

based on our logistic regression models (Figure 1C, Table 3; Fowler-Finn et al. 2014, 2018, 2019;

Machado & Macías-Ordóñez 2007). If females orient their bodies in this manner, males cannot

access the genital operculum and therefore cannot initiate intromission (Figure 1C). Many

arthropod species have evolved morphological anti-clasping adaptations (Arnqvist & Rowe 1995,

2005; Karlsson Green et al. 2013) similarly, body angling might be considered a concordant

behavioural anti-clasping adaptation – actions to move the genitals away from males are common

among arthropods with high levels of sexual conflict (Akinyemi et al. 2021; Burke et al. 2015;

Golov et al. 2018; Machado & Macías-Ordóñez 2007; Maroni et al. 2023; Zweerus et al. 2021).

While our regression modelling did not include female body angle as a significant predictor, it is

possible that the enhanced clasping morphology found in male *L. vittatum* allows them to

effectively overcome this female behaviour, and further investigation is required.

Alternatively, there is some evidence that extended clasping time in *L. vittatum* may be

attributable to female antagonism rather than male, which would also serve to explain the

discrepancy between clasping times in the two nonsacculate species. First suggested by Fowler-

Finn et al. (2014), female *L. vittatum* may clasp the male's penis using the genital operculum,

allowing the female to receive more nuptial gift than is optimal for the male to provide. Once

clasping has been achieved and intromission has begun, male *L. vittatum* occasionally appear to

struggle to end copulation (Fowler-Finn et al. 2014). This phenomenon is supported by genital

morphology; female *L. vittatum* have well-developed opercular musculature and sclerotization, in

stark contrast to the male's relatively fragile penis (Burns & Shultz 2016). Burns and Shultz (2016)

additionally cite reports of males with broken penises in the field, an injury consistent with

struggles to disengage females. Female aggression can negatively correlate with nuptial gift

quality and quantity (Kuriwada & Kasuya 2012; Toft & Albo 2016), and in *L. vittatum* this

manipulation by females might be the direct result of the deterioration of nuptial gift quality and

quantity (Khan et al. 2018; Machado & Burns 2023; Machado & Macías-Ordóñez 2007). Because

females of nonsacculate species can only receive nuptial gift fluid during intromission, rather than

both before and during, there may be a tangible benefit to females remaining in copula as long

as possible. It may be that this acts as a cost-mitigating strategy for females, although it seems

unlikely that females would prefer this as opposed to simply avoiding the encounter altogether

due to the elevated levels of precopulatory resistance from females. The potential transition from

male pre-copulatory antagonism to female copulatory antagonism would provide a fascinating

dichotomy of sexual conflict deserving further research.

Mate guarding is another strong indicator of potential sexual conflict commonly found in

arthropods (Alcock 1994; Arnqvist & Rowe 2005; Simmons 2002). Mate guarding may allow

males additional control over female mating decisions by discouraging remating, preventing

sperm removal by rival males, and reducing sperm competition – all factors which might

alternatively (or additionally) be influenced by nuptial gifts (Alcock 1994; Arnqvist & Rowe 2005;

Elias et al. 2014; Frankino & Sakaluk 1994; Gór et al. 2023; Hagg et al. 2024; Lewis & South

2012; Machado & Burns 2023; Sakaluk 1991; Simmons 2002). It is important to note, however,

that mate guarding can be beneficial if the guarding male reduces future harassment or predation

for the female, though in these species female resistance to guarding should be less prevalent

(Rodríguez-Muñoz et al. 2011; Sherman 1983). Sperm competition and cryptic female choice are

distinct possibilities in leiobunine harvesters – all females have a sperm-storing spermatheca,

while some species have a complex, multi-chambered spermatheca potentially affording the

female control over sperm fate (Karachiwalla et al. 2020; Shultz & Pinto-da-Rocha 2007). These

potential confounding factors would only further incentivize males to guard females and prevent

remating, as suggested by both models and experimental evidence (Bateman et al. 2001; Del

Matto et al. 2021; Gór et al. 2023; Harts & Kokko 2013). Many male Opiliones, including

Leiobunum species, engage in extended mate guarding which typically lasts until oviposition

(Fowler-Finn et al. 2014, 2018, 2019; Machado & Burns 2023, Macías-Ordóñez 1997; Wijnhoven

2011; Zatz et al. 2011). This is likely an extremely costly behaviour for males because these

harvesters often aggregate in high density, leading to elevated levels of male-male competition

(Macías-Ordóñez 1997). The few field studies of *Leiobunum* mating behaviour have observed

male *Leiobunum spp.* defending ovipositing females from upwards of 6 rival males, although

evidence for late-season male sex-ratio biases and the dearth of field studies likely leads to this

being an underestimate (Bacon, R. & Brown, T. A., pers. observ; Macías-Ordóñez 1997;

Wijnhoven 2011). Our study did not find *L. vittatum* guarding time to be significantly longer than

sacculate species, but a key missing factor may be the absence of competitors. In previous

studies, male *L. vittatum* were found to be most aggressive when in the presence of both males

and females during their reproductive period (Macías-Ordóñez 1997). While this aggression was

often male-directed, it remains possible that the absence of male competitors resulted in less

guarding than expected.

<u>Multiple mating strategies</u>

The *calcar* species-group and *vittatum* species-group have convergently evolved similar

nuptial gift chemistries and penis morphologies (Burns et al. 2013; Kahn et al. 2018), yet the

groups vary significantly in sexual and somatic morphology and biomechanics (Burns & Shultz

(2015, 2016). Female genital operculum morphology differs in reproductive barrier structure

between groups as well (Burns et al. 2013; Ingianni et al. 2011). Despite females of both groups

possessing unique, species-specific pregenital barriers formed through sclerotization of the

operculum and sternum, the mechanism through which these structures engage varies between

groups (Burns et al. 2013; Shultz 2018). *Leiobunum euserratipalpe* does not have sexually

dimorphic palpal morphology, while *L. vittatum* does (Ingianni et al. 2011; Shultz 2018). This

variation in somatic morphology correlates with the significant differences in clasping duration

revealed in this study – *L. vittatum* males clasp longer than *L. euserratipalpe* and possess clasping

structures indicative of this. *Leiobunum euserratipalpe* males, lacking enhanced palpal clasping

structures, do not have particularly long clasping durations and instead guard females for

extended periods postcopulation.

These variable suites of antagonistic traits indicate the evolutionary forces driving the

transition to antagonism may differ between the *calcar* and *vittatum* species-groups, and further

research investigating these mechanisms is needed. There are many temperate species within

the Opiliones family Sclerosomatidae with documented transitions towards nonsacculate

morphology, but to date no tropical species with these transitions have been reported (Burns et

al. 2013; Burns & Tsurusaki 2016; Machado & Burns 2023; Martens 2020). While there is a dearth

of tropical sexual conflict research overall, neotropical Opiliones have been fairly well studied,

albeit with a focus on non-Sclerosomatidae species (Bacon et al. 2023; Burns & Machado 2023;

Pinto-da-Rocha et al. 2007). This conspicuous lack of ostensibly antagonistic structures in tropical

Sclerosomatidae might indicate that harsher climes drive the transition towards antagonism in

temperate sclerosomatids. These data, in conjunction with previous morphological, phylogenetic,

and biochemical work (Burns et al. 2013; Burns & Shultz 2015, 2016; Kahn et al. 2018), suggest

nonsacculate *Leiobunum* harvesters have a frequently coercive mating system driven by sexual

conflict, but this is not to say that every mating interaction is inherently antagonistic.

An alternative explanation for the behaviour we observed is cryptic female choice (CFC),

which has been proposed to explain many common morphological and behavioural patterns

typically assumed to be due to sexually antagonistic coevolution (Eberhard 1996, 2010). Under

CFC, females are expected to exhibit peri- or post-copulatory preferences for particular male

genital morphologies, such as those better able to stimulate the female (Eberhard 1996, 2010).

The female may then preferentially store sperm, fertilise eggs, or reduce its remating rate to the

benefit of the male (Eberhard 1996, 2010). Eberhard (2011) presents an excellent framework for

effectively testing whether male stimulation of the female occurs by experimentally "blinding" a

female to potential stimulation. Myers et al. (2016) performed such an experiment on the stick

insect *Clitarchus hookeri*, which exhibits mating behaviour similar to Opiliones in that males must

clasp females for mating to occur, and females have a moveable genital operculum. Myers et al.

(2016) demonstrate through female desensitisation and male clasper abrasion that this clasping

is not antagonistic and likely serves to signal to females to open their genital operculum and allow

intromission. Though they note that, unlike in the present study, unmodified females typically do

not resist unmodified males, their study nonetheless presents an excellent example of necessary

future work in *Leiobunum*.

Conclusions

While we intended to expedite video analysis of reproductive behaviour through a

combination of automated tracking and manual scoring, we found the benefits of incorporating

automated tracking to be minimal. Our subsequent attempts using alternative tracking programs

resulted in similar difficulties, and many fine-scale behaviours, such as those involving leg or

pedipalp movements, were not feasible to score with automated tracking and required manual

scoring. Automated tracking may still provide benefits when used across many studies because

trained models can be reused within a species, but manual tracking likely remains preferable for

species with fine-scale behaviour or rapid positional changes, or when de novo training is

required.

Our study demonstrates that mating behaviour significantly differs between sacculate and

nonsacculate species, providing another line of evidence that sexual conflict has shaped

reproduction in these Opiliones. Our data suggests that nonsacculate harvesters may resort to

antagonistic behaviours in lieu of nuptial gifts to increase fitness through some combination of

copulation success, insemination success, sperm transfer, sperm storage, and paternity share.

Importantly, genital modification studies and economic studies of the costs of mating remain to

be done in *Leiobunum* harvesters. Though it remains possible that other mechanisms, such as

cryptic female choice, play an important role in this genus, it is unlikely for sexual conflict to be

entirely absent in a system in which there is benefit for males to circumvent female preference

(Snow et al. 2019).

A significant area that remains unclear is the effect of reproductive phenology on sexual

conflict. North American leiobunine harvesters have significant variation in the timing of major life

events such as maturation, oviposition, and senescence, as well as in the length of the

reproductive period (Curtis & Machado 2007; Machado & Macías-Ordóñez 2007). Additionally,

male-male competition and female antagonism are further factors which likely contribute to the

evolution and maintenance of male antagonism in leiobunine Opiliones. How these myriad factors

interact and impact sexual conflict is not well established, and the evidence of significant variation

in mating behaviour in leiobunine harvesters provides the perfect opportunity for further

behavioural research.

Author Contributions: Tyler A. Brown: Conceptualization, methodology, investigation,

formal analysis, visualisation, writing – original draft, writing – review & editing, funding

acquisition. **Emily Marinko**: Investigation, formal analysis, writing – review & editing.

Mercedes Burns: Conceptualization, methodology, formal analysis, writing – review &

editing, supervision, funding acquisition.

Declaration of Interest: The authors declare they have no competing interests.

Data Availability: Mendeley Data doi: https://doi.org/10.17632/gsvd5z9g2b.2

Acknowledgements

We would like to thank Ryan Bacon, Sai Wai, Emily Redmond, Lian Jackson, and Shelby Hawk for assistance with specimen collection and husbandry. We would also like to thank Sophia Nawaz and Megan Ramirez Cuenca for data processing assistance and Dr. Andrew Gordus for his helpful advice regarding automated tracking implementation. T. A. B. and M. B. were supported by funds from NSF IOS 00116. This work was also funded in part by the American Arachnological Society and Washington Biologists' Field Club, which also provided access to Plummer's Island.

Appendix

O penile sacs present, female barrier absent
● penile sacs absent, female barrier present
◑ penile sacs present, female barrier present
◐ penile sacs absent, female barrier absent

798
799 **Figure A1:** Bayesian Maximum Clade Credibility tree of the *Leiobunum* species used in this study,

800 adapted from Burns & Shultz (2015), where scale = base substitutions/site. Parsimonious

801 character state combinations of penile sac presence and female pregenital barrier presence are

802 mapped on branches.

Figure A2: Association metrics between male and female pairs. Boxes represent medians plotted

with IQR, symbols represent individual trial values, and whiskers correspond to the minimum and

maximum trial values. The results from group comparison tests are shown in the upper right-hand

corners, while significant results from post-hoc multiple comparisons tests are noted by red

brackets with asterisks (* = *P* < 0.05). The simplified phylogeny indicates reproductive

morphology; red tips indicate nonsacculate species and black tips represent sacculate species

(phylogeny adapted from Burns et al. 2013). **A)** Average distance in centimetres between male

and female pairs during attempted-clasp trials. **B)** Average distance in centimetres between male

and female pairs during successful-clasp trials. **C)** Total time in seconds the female was within

one leg span of the male during attempted-clasp trials. **D)** Total time in seconds the female was

within one leg span of the male during successful-clasp trials.

Figure A3: Instances of females fleeing from males during reproductive trials. Boxes represent

medians plotted with IQR, symbols represent individual trial values, and whiskers correspond to

the minimum and maximum trial values. The results from group comparison tests are shown in

the upper right-hand corners, while significant results from post-hoc multiple comparisons tests

are noted by red brackets with asterisks (* = P < 0.05; **** = P < 0.000). The simplified phylogeny

indicates reproductive morphology; red tips indicate nonsacculate species and black tips

represent sacculate species (phylogeny adapted from Burns et al. 2013). **A)** Attempted-clasp

trials. **B)** Successful-clasp trials.

Metric	Result	Normality	N(total)	N(Lbr)	N(Lald)	N(Lvitt)	N(Leus)
Latency	Attempts	No	75	28	18	14	15
Contact Time	Attempts	No	75	28	18	14	15
	Successes	No	53	13	13	13	14
Pre-Attempt Contact Time	Attempts	No	75	28	18	14	15
	Successes	No	53	13	13	13	14
Flee Count	Attempts	No	75	28	18	14	15
	Successes	No	53	13	13	13	14
Tap Count	Attempts	No	75	28	18	14	15
	Successes	No	53	13	13	13	14
Pursuit Angle	Attempts	No	74	28	18	14	14
	Successes	No	52	13	13	13	13
Time within 1 Leg Span	Attempts	No	74	28	18	14	14
	Successes	No	52	13	13	13	13
Average Distance	Attempts	Log	74	28	18	14	14
	Successes	Log	52	13	13	13	13
Maximum Female Angle	Attempts	No	75	28	18	14	15
	Successes	No	53	13	13	13	14
Clasping Time	Successes	No	53	13	13	13	14
Guarding Time	Successes	No	53	13	13	13	14

Table A1: Behavioural metrics recorded during this study (species codes: *Leiobunum vittatum*: Lvitt; *L. euserratipalpe*: Leus; *L. bracchiolum*: Lbr; *L. aldrichi*: Lald). Shapiro-Wilk test for normality results are shown for all metrics save average distance, which was log-normal based on a Shapiro-Wilk test for lognormality. Sample sizes for each metric follow. Note that for pursuit angle, time within one leg span, and average distance, one *L. euserratipalpe* trial could not be used due to video corruption preventing TRex scoring.

Metric	Trial Type	Pagel's λ	logL(λ)	LR(λ=0)	*P*-value
Latency to 1st Attempt	Attempts	6.61e-05	-29.451	-6.88e-05	1
Pursuit Angle	Attempts	6.61e-05	-9.263	-2.32e-05	1
	Successes	6.61e-05	-9.831	-3.37e-05	1
Contact Time	Attempts	6.61e-05	-30.302	-1.02e-04	1
	Successes	0.999	-28.626	0.187	0.6658
Flee Count	Attempts	6.61e-05	-2.328	-6.79e-05	1
	Successes	6.61e-05	-2.328	-6.79e-05	1
Tap Count	Attempts	0.999	-2.799	0.208	0.6481
	Successes	0.999	-2.799	0.208	0.6481
Pre-Attempt Contact Time	Attempts	6.61e-05	-25.851	-5.73e-05	1
	Successes	6.61e-05	-24.774	-2.57e-05	1
Time within 1 Leg Span	Attempts	6.61e-05	-29.705	-1e-04	1
	Successes	0.999	-28.167	0.189	0.6634
Average Distance	Attempts	6.61e-05	-6.717	-9.68e-05	1
	Successes	0.999	-6.788	0.229	0.6325
Maximum Female Body Angle	Attempts	0.999	-19.163	0.057	0.8111
	Successes	6.61e-05	-18.764	-6.09e-05	1
Clasping Time	Successes	6.61e-05	-28.053	-1.97e-05	1
Guarding Time	Successes	6.61e-05	-30.947	-7.55e-06	1

835 **Table A2:** Degree of phylogenetic signal λ within the various behavioural metrics. Pagel's lambda

836 (Pagel 1999) was calculated using the phytools package (Revell 2012). We used a pruned version

837 (consisting of our four study species) of a maximum clade credibility tree constructed from

838 mitochondrial and nuclear sequences by Burns et al (2013). These models were then compared

839 to Brownian motion predictions with a likelihood ratio test to produce our P-values.

MLR Model/Factor	F (df)	Estimate	SE	|t|	P value
Total-species model	3.672 (4, 48)	-	-	-	**0.0109**
Intercept	-	**1378**	**453.6**	**3.038**	**0.0038**
# male legs	-	**-290.4**	**81.68**	**3.556**	**0.0009**
# female legs	-	92.59	72.23	1.282	0.2061
# male sensory legs	-	**489.2**	**155.6**	**3.143**	**0.0029**
# female sensory legs	-	**-257.7**	**114.7**	**2.247**	**0.0292**
L. vittatum model	1.186 (4, 8)	-	-	-	0.3867
L. euserratipalpe model	1.272 (4, 8)	-	-	-	0.3493
L. bracchiolum model	0.795 (4, 8)	-	-	-	0.5607
L. aldrichi model	0.239 (4, 8)	-	-	-	0.9087

842 **Table A3:** Multiple linear regression of clasping duration with leg number metrics as predictor

843 variables for the total-species model and species-level models. Significant predictor variables are

844 in bold; no predictor variables were dropped from modelling.

References

Akinyemi, A. O., Subramanian, S., Mfuti, D. K., Pope, T. W., Tamiru, A., & Kirk, W. D. J.

(2021). Mating behaviour, mate choice and female resistance in the bean flower thrips

(*Megalurothrips sjostedti*). *Scientific Reports*, *11*(1), 14504.

https://doi.org/10.1038/s41598-021-93891-5

Albo, M. J., Franco-Trecu, V., Wojciechowski, F. J., Toft, S., & Bilde, T. (2019).

Maintenance of deceptive gifts in a natural spider population: Ecological and

demographic factors. *Behavioral Ecology*, *30*(4), 993–1000.

https://doi.org/10.1093/beheco/arz040

Albo, M. J., Macías-Hernández, N., Bilde, T., & Toft, S. (2017). Mutual benefit from

exploitation of female foraging motivation may account for the early evolution of gifts in

spiders. *Animal Behaviour*, *129*, 9–14. https://doi.org/10.1016/j.anbehav.2017.05.001

Alcock, J. (1994). Postinsemination Associations Between Males and Females in Insects:

The Mate-Guarding Hypothesis. *Annual Review of Entomology*, *39*, 1–21.

Arnqvist, G., & Rowe, L. (1995). Sexual conflict and arms races between the sexes: A

morphological adaptation for control of mating in a female insect. *Proceedings of the*

Royal Society of London. Series B: Biological Sciences, *261*(1360), 123–127.

https://doi.org/10.1098/rspb.1995.0126

Arnqvist, G., & Rowe, L. (2005). *Sexual Conflict*. Princeton University Press.

Bacon, R., Washington, D., Johnson, M.-K., & Burns, M. (2023). The Geography of Sexual

Conflict: A Synthetic Review. *The American Naturalist*, *201*(3), 429–441.

https://doi.org/10.1086/722797

Bateman, P. W., Gilson, L. N., & Ferguson, J. W. H. (2001). Investment in Mate Guarding

May Compensate for Constraints on Ejaculate Production in the Cricket *Gryllodes*

sigillatus. *Ethology*, *107*(12), 1087–1098. https://doi.org/10.1046/j.1439-

0310.2001.00756.x

Beyer, M., Mangliers, J., & Tuni, C. (2021). Silk-borne chemicals of spider nuptial gifts elicit

female gift acceptance. *Biology Letters*, *17*(11), 20210386.

https://doi.org/10.1098/rsbl.2021.0386

Blanckenhorn, W. U., Baur, J., Busso, J. P., Giesen, A., Gourgoulianni, N., Van

Koppenhagen, N., Roy, J., Schäfer, M. A., Wegmann, A., & Rohner, P. T. (2020).

Sexual size dimorphism is associated with reproductive life history trait differentiation in

coexisting sepsid flies. *Oikos*, *129*(8), 1152–1162. https://doi.org/10.1111/oik.07036

Blomberg, S. P., Garland, T., & Ives, A. R. (2003). Testing for Phylogenetic Signal in

Comparative Data: Behavioral Traits are More Labile. *Evolution*, *57*(4), 717–745.

https://doi.org/10.1111/j.0014-3820.2003.tb00285.x

Blumstein, D. T., & Daniel, J. C. (2007a). *Quantifying behavior the JWatcher way*. Sinauer

Associates.

Bokides, D., Lou, Y., & Hamilton, I. M. (2012). A model of sexual selection and female use

of refuge in a coercive mating system. *Proceedings of the Royal Society B: Biological*

Sciences, 279(1741), 3209–3216. https://doi.org/10.1098/rspb.2012.0246

Boulton, R. A., Zuk, M., & Shuker, D. M. (2018). An Inconvenient Truth: The Unconsidered

Benefits of Convenience Polyandry. *Trends in Ecology & Evolution*, *33*(12), 904–915.

https://doi.org/10.1016/j.tree.2018.10.002

Briceño, R. D., & Eberhard, W. G. (2009). Experimental demonstration of possible cryptic

female choice on male tsetse fly genitalia. *Journal of Insect Physiology, 55*(11), 989–

996. https://doi.org/10.1016/j.jinsphys.2009.07.001

Briceño, R. D., Eberhard, W. G., Chinea-Cano, E., Wegrzynek, D., & Rolo, T. D. S. (2016).

Species-specific differences in the behavior of male tsetse fly genitalia hidden in the

female during copulation. *Ethology Ecology & Evolution, 28*(1), 53–76.

https://doi.org/10.1080/03949370.2014.1002114

Brown, W. D., & Barry, K. L. (2016). Sexual cannibalism increases male material

investment in offspring: Quantifying terminal reproductive effort in a praying mantis.

Proceedings of the Royal Society B: Biological Sciences, *283*(1833), 20160656.

https://doi.org/10.1098/rspb.2016.0656

Buchinger, T. J., & Li, W. (2023). Chemical communication and its role in sexual selection

across Animalia. *Communications Biology*, *6*(1), 1178. https://doi.org/10.1038/s42003-

023-05572-w

Burke, N. W., & Bonduriansky, R. (2018). The fitness effects of delayed switching to sex in

a facultatively asexual insect. *Ecology and Evolution*, *8*(5), 2698–2711.

https://doi.org/10.1002/ece3.3895

Burke, N. W., Crean, A. J., & Bonduriansky, R. (2015). The role of sexual conflict in the

evolution of facultative parthenogenesis: A study on the spiny leaf stick insect. *Animal*

Behaviour, *101*, 117–127. https://doi.org/10.1016/j.anbehav.2014.12.017

Burns, M., Hedin, M., & Shultz, J. W. (2012). Molecular phylogeny of the leiobunine

harvestmen of eastern North America (Opiliones: Sclerosomatidae: Leiobuninae).

Molecular Phylogenetics and Evolution, *63*(2), 291–298.

https://doi.org/10.1016/j.ympev.2011.12.025

Burns, M. M., Hedin, M., & Shultz, J. W. (2013). Comparative Analyses of Reproductive

Structures in Harvestmen (Opiliones) Reveal Multiple Transitions from Courtship to

Precopulatory Antagonism. *PLoS ONE*, *8*(6), e66767.

https://doi.org/10.1371/journal.pone.0066767

Burns, M., & Shultz, J. W. (2015). Biomechanical Diversity of Mating Structures among

Harvestmen Species Is Consistent with a Spectrum of Precopulatory Strategies. *PLOS*

ONE, *10*(9), e0137181. https://doi.org/10.1371/journal.pone.0137181

Burns, M., & Shultz, J. W. (2016). Mechanical properties of male genitalia in *Leiobunum*

harvestmen (Opiliones: Sclerosomatidae). *Journal of Arachnology*, *44*(2), 199–209.

https://doi.org/10.1636/JoA-S-15-010

Burns, M., & Tsurusaki, N. (2016). Male Reproductive Morphology Across Latitudinal Clines

and Under Long-Term Female Sex-Ratio Bias. *Integrative and Comparative Biology*,

56(4), 715–727. https://doi.org/10.1093/icb/icw017

Curtis, D. J., & Machado, G. (2007). Ecology. In R. Pinto da Rocha, G. Machado, & G.

Giribet (Eds.), *Harvestmen: The Biology of Opiliones* (pp. 280–308). Harvard University

Press.

930 Davranoglou, L., Taylor, G. K., & Mortimer, B. (2023). Sexual selection and predation drive

931   the repeated evolution of stridulation in Heteroptera and other arthropods. *Biological*

932   *Reviews*, *98*(3), 942–981. https://doi.org/10.1111/brv.12938

933 del Castillo, R. C., & Gwynne, D. T. (2007). Increase in song frequency decreases

934   spermatophore size: Correlative evidence of a macroevolutionary trade-off in katydids

935   (Orthoptera: Tettigoniidae). *Journal of Evolutionary Biology*, *20*(3), 1028–1036.

936   https://doi.org/10.1111/j.1420-9101.2006.01298.x

937 Del Matto, L., De Macedo Rego, R., & Santos, E. (2021). Mate-guarding duration is mainly

938   influenced by the risk of sperm competition and not by female quality in a golden orb-

939   weaver spider. *PeerJ*, *9*, e12310. https://doi.org/10.32942/OSF.IO/2U8TC

940 del Sol, J. F., Hongo, Y., Boisseau, R. P., Berman, G. H., Allen, C. E., & Emlen, Douglas. J.

941   (2021). Population differences in the strength of sexual selection match relative

942   weapon size in the Japanese rhinoceros beetle, *Trypoxylus dichotomus* (Coleoptera:

943   Scarabaeidae). *Evolution*, *75*(2), 394–413. https://doi.org/10.1111/evo.14101

944 Dmitriew, C., & Blanckenhorn, W. U. (2012). The Role of Sexual Selection and Conflict in

945   Mediating Among-Population Variation in Mating Strategies and Sexually Dimorphic

946   Traits in *Sepsis punctum*. *PLoS ONE*, *7*(12), e49511.

947   https://doi.org/10.1371/journal.pone.0049511

948 Eberhard, W. G. (1996). *Female Control: Sexual Selection by Cryptic Female Choice*.

949   Princeton University Press. https://doi.org/10.2307/j.ctvs32rx1

950 Eberhard, W. G. (2010). Evolution of genitalia: Theories, evidence, and new directions.

951   *Genetica, 138*(1), 5–18. https://doi.org/10.1007/s10709-009-9358-y

952 Eberhard, W. G. (2011). Experiments with genitalia: A commentary. Trends in Ecology &

953   *Evolution, 26*(1), 17–21. https://doi.org/10.1016/j.tree.2010.10.009

Edvardsson, M., & Canal, D. (2006). The effects of copulation duration in the bruchid beetle

Callosobruchus maculatus. *Behavioral Ecology*, *17*(3), 430–434.

https://doi.org/10.1093/beheco/arj045

Edward, D. A., & Chapman, T. (2011). The evolution and significance of male mate choice.

Trends in Ecology & Evolution, *26*(12), 647–654.

https://doi.org/10.1016/j.tree.2011.07.012

Edward, D. A., Stockley, P., & Hosken, D. J. (2015). Sexual Conflict and Sperm

Competition. *Cold Spring Harbor Perspectives in Biology*, *7*(4), a017707.

https://doi.org/10.1101/cshperspect.a017707

Elias, D. O., Sivalinghem, S., Mason, A. C., Andrade, M. C. B., & Kasumovic, M. M. (2014).

Mate-guarding courtship behaviour: Tactics in a changing world. *Animal Behaviour*, *97*,

25–33. https://doi.org/10.1016/j.anbehav.2014.08.007

Fowler, E. K., Leigh, S., Bretman, A., & Chapman, T. (2022). Plastic responses of males

and females interact to determine mating behavior. *Evolution*, *76*(9), 2116–2129.

https://doi.org/10.1111/evo.14568

Fowler-Finn, K., Boyer, S., Ikagawa, R., Jeffries, T., Kahn, P., Larsen, E., Lee, D., &

Smeester, M. (2018). Variation in Mating Dynamics across Five Species of Leiobunine

Harvestmen (Arachnida: Opiliones). *Biology*, *7*(2), 36.

https://doi.org/10.3390/biology7020036

Fowler-Finn, K., Boyer, S. L., Ikagawa, R., Jeffries, T., Kahn, P. C., Larsen, E. M., Lee, D.,

Sasson, D., & Smeester, M. (2019). Qualitative and quantitative comparisons of mating

behaviour across multiple populations and six species of leiobunine harvestmen

(Arachnida: Opiliones). *Behaviour*, *156*(3–4), 363–390.

https://doi.org/10.1163/1568539X-00003545

Fowler-Finn, K. D., Triana, E., & Miller, O. G. (2014). Mating in the harvestman *Leiobunum*

vittatum (Arachnida: Opiliones): from premating struggles to solicitous tactile

engagement. *Behaviour*, *151*(12–13), 1663–1686. https://doi.org/10.1163/1568539X-

00003209

Fox, C. W., Stillwell, R. C., Wallin, W. G., & Hitchcock, L. J. (2006). Temperature and host

species affect nuptial gift size in a seed-feeding beetle. *Functional Ecology*, *20*(6),

1003–1011. https://doi.org/10.1111/j.1365-2435.2006.01197.x

Frankino, A. W., & Sakaluk, S. K. (1994). Post copulatory mate guarding delays

promiscuous mating by female decorated crickets. *Animal Behaviour*, *48*, 1479–1481.

https://doi.org/10.1006/anbe.1994.1389

Gao, Q., Turnell, B. R., Hua, B., & Shaw, K. L. (2019). The effect of nuptial gift number on

fertilization success in a Hawaiian swordtail cricket. *Behavioral Ecology and*

Sociobiology, *73*(7), 92. https://doi.org/10.1007/s00265-019-2705-9

Ghislandi, P. G., Pekár, S., Matzke, M., Schulte-Döinghaus, S., Bilde, T., & Tuni, C. (2018).

Resource availability, mating opportunity and sexual selection intensity influence the

expression of male alternative reproductive tactics. *Journal of Evolutionary Biology*,

31(7), 1035–1046. https://doi.org/10.1111/jeb.13284

Gnaspini, P., & Hara, M. R. (2007). Defense Mechanisms. In R. Pinto da Rocha, G.

Machado, & G. Giribet (Eds.), *Harvestmen: The Biology of Opiliones* (pp. 374–399).

Harvard University Press.

Golov, Y., Harari, A., Rillich, J., & Ayali, A. (2018). Precopulatory behavior and sexual

conflict in the desert locust. *PeerJ*, *6*, e4356. https://doi.org/10.7717/peerj.4356

Gór, Á., Lang, Z., Pásztor, K., Szigeti, V., Vajna, F., & Kis, J. (2023). Mate-guarding

success depends on male investment in a butterfly. *Ecology and Evolution*, *13*(9),

e10533. https://doi.org/10.1002/ece3.10533

Gwynne, D. T. (2008). Sexual Conflict over Nuptial Gifts in Insects. *Annual Review of*

Entomology, *53*(1), 83–101. https://doi.org/10.1146/annurev.ento.53.103106.093423

Hagg, L., LaMere, C. J., & Zuk, M. (2024). Postcopulatory song as a mate-guarding tactic in the Pacific field cricket, *Teleogryllus oceanicus*. *Animal Behaviour*, *207*, 201–207. https://doi.org/10.1016/j.anbehav.2023.11.001

Harts, A. M. F., & Kokko, H. (2013). Understanding Promiscuity: When is Seeking Additional Mates Better than Guarding an Already Found One?: Mate-Guarding, Monogamy, and Promiscuity. *Evolution*, *67*(10), 2838–2848. https://doi.org/10.1111/evo.12163

Hedin, M., Tsurusaki, N., Macías-Ordóñez, R., & Shultz, J. W. (2012). Molecular systematics of sclerosomatid harvestmen (Opiliones, Phalangioidea, Sclerosomatidae): Geography is better than taxonomy in predicting phylogeny. *Molecular Phylogenetics and Evolution*, *62*(1), 224–236. https://doi.org/10.1016/j.ympev.2011.09.017

Iglesias-Carrasco, M., Fox, R. J., Vega-Trejo, R., Jennions, M. D., & Head, M. L. (2019). An experimental test for body size-dependent effects of male harassment and an elevated copulation rate on female lifetime fecundity and offspring performance. *Journal of Evolutionary Biology*, *32*(11), 1262–1273. https://doi.org/10.1111/jeb.13526

Ingianni, E. A., McGhee, C. R., & Shultz, J. W. (2011). Taxonomy of the *Leiobunum calcar* species-group (Opiliones: Sclerosomatidae: Leiobuninae). *Journal of Arachnology*, *39*(3), 454–481. https://doi.org/10.1636/Ha11-42.1

Jarrige, A., Body, M., Giron, D., Greenfield, M. D., & Goubault, M. (2015). Amino acid composition of the bushcricket spermatophore and the function of courtship feeding: Variable composition suggests a dynamic role of the nuptial gift. *Physiology & Behavior*, *151*, 463–468. https://doi.org/10.1016/j.physbeh.2015.08.009

Johns, J. L., Roberts, J. A., Clark, D. L., & Uetz, G. W. (2009). Love bites: Male fang use during coercive mating in wolf spiders. *Behavioral Ecology and Sociobiology*, *64*(1), 13–18. https://doi.org/10.1007/s00265-009-0812-8

Kahn, P. C., Cao, D. D., Burns, M., & Boyer, S. L. (2018). Nuptial gift chemistry reveals

convergent evolution correlated with antagonism in mating systems of harvestmen

(Arachnida, Opiliones). *Ecology and Evolution*, *8*(14), 7103–7110.

https://doi.org/10.1002/ece3.4232

Karachiwalla, Z., deCarvahlo, T., & Burns, M. (2023). Spermathecal Variation in Temperate

Opiliones. *Integrative And Comparative Biology*, *63*(1), 188–197.

https://doi.org/10.1093/icb/icaa120

Karlsson Green, K., Kovalev, A., Svensson, E. I., & Gorb, S. N. (2013). Male clasping

ability, female polymorphism and sexual conflict: Fine-scale elytral morphology as a

sexually antagonistic adaptation in female diving beetles. *Journal of The Royal Society*

Interface, *10*(86), 20130409. https://doi.org/10.1098/rsif.2013.0409

Kokko, H., & Jennions, M. D. (2014). The Relationship between Sexual Selection and

Sexual Conflict. *Cold Spring Harbor Perspectives in Biology*, *6*(9), a017517–a017517.

https://doi.org/10.1101/cshperspect.a017517

Křemenová, J., Bartonička, T., Balvín, O., Massino, C., Reinhardt, K., Sasínková, M., Weig,

A. R., & Otti, O. (2021). Male diet affects female fitness and sperm competition in

human- and bat-associated lineages of the common bedbug, *Cimex lectularius*.

Scientific Reports, *11*(1), 15538. https://doi.org/10.1038/s41598-021-94622-6

Krupa, J. J., Leopold, W. R., & Sih, A. (1990). Avoidance of Male Giant Water Striders By

Females. *Behaviour*, *115*(3–4), 247–253. https://doi.org/10.1163/156853990X00590

Kunz, J. A., Duvot, G. J., Van Noordwijk, M. A., Willems, E. P., Townsend, M., Mardianah,

N., Utami Atmoko, S. S., Vogel, E. R., Nugraha, T. P., Heistermann, M., Agil, M.,

Weingrill, T., & Van Schaik, C. P. (2021). The cost of associating with males for

Bornean and Sumatran female orangutans: A hidden form of sexual conflict?

Behavioral Ecology and Sociobiology, *75*(1), 6. https://doi.org/10.1007/s00265-020-

02948-4

Kuriwada, T., & Kasuya, E. (2012). Nuptial gifts protect male bell crickets from female

aggressive behavior. *Behavioral Ecology*, *23*(2), 302–306.

https://doi.org/10.1093/beheco/arr186

Lehmann, G. U. C., Heller, K.-G., & Mai, R. (2016). Prolonged copulations as an alternative

to male nuptial gift investment in the bushcricket *Letana inflata* (Orthoptera:

Tettigoniidae). *Journal of Ethology*, *34*(2), 191–194. https://doi.org/10.1007/s10164-

016-0457-9

Lehmann, G. U. C., & Lehmann, A. W. (2016). Material benefit of mating: The bushcricket

spermatophylax as a fast uptake nuptial gift. *Animal Behaviour*, *112*, 267–271.

https://doi.org/10.1016/j.anbehav.2015.12.022

Lehtonen, J., Jennions, M. D., & Kokko, H. (2012). The many costs of sex. *Trends in*

Ecology & Evolution, *27*(3), 172–178. https://doi.org/10.1016/j.tree.2011.09.016

Lewis, S. M., Vahed, K., Koene, J. M., Engqvist, L., Bussière, L. F., Perry, J. C., Gwynne,

D., & Lehmann, G. U. C. (2014). Emerging issues in the evolution of animal nuptial

gifts. *Biology Letters*, *10*(7), 20140336. https://doi.org/10.1098/rsbl.2014.0336

Lewis, S., & South, A. (2012). The Evolution of Animal Nuptial Gifts. In H. J. Brockmann, T.

J. Roper, M. Naguib, J. C. Mitani, & L. W. Simmons (Eds.), *Advances in the Study of*

Behavior (Vol. 44, pp. 53–97). Elsevier. https://doi.org/10.1016/B978-0-12-394288-

3.00002-2

Liu, X., Hayashi, F., Lavine, L. C., & Yang, D. (2015). Is diversification in male reproductive

traits driven by evolutionary trade-offs between weapons and nuptial gifts?

Proceedings of the Royal Society B: Biological Sciences, *282*(1807), 20150247.

https://doi.org/10.1098/rspb.2015.0247

Machado, G., & Burns, M. (2023). Reproductive biology of harvestmen (Arachnida:

Opiliones): a review of a rapidly evolving research field. *Current Zoology*, *zoac102*, 1–

21. https://doi.org/10.1093/cz/zoac102

Machado, G., & Macías-Ordóñez, R. (2007). Reproduction. In R. Pinto da Rocha, G.

    Machado, & G. Giribet (Eds.), *Harvestmen: The Biology of Opiliones* (pp. 414–454).

    Harvard University Press.

Macías-Ordóñez, R. (1997). *The mating system of* Leiobunum vittatum *(Say 1891)*

    (Arachnida: Opiliones: Palpitores): Resource defense polygyny in the striped

    harvestman (Doctoral dissertation). Bethlehem, PA: Lehigh University.

Macías-Ordóñez, R., Machado, G., Pérez-González, A., & Shultz, J. W. (2010). Genitalic

    evolution in Opiliones. In *The Evolution of Primary Sexual Characters in Animals* (p.

    537). Oxford University Press.

Małek, D. K., Dańko, M. J., & Czarnoleski, M. (2019). Does seed size mediate sex-specific

    reproduction costs in the *Callosobruchus maculatus* bean beetle? *PLOS ONE*, *14*(12),

    e0225967. https://doi.org/10.1371/journal.pone.0225967

Maroni, P. J., Bryant, K. A., & Tatarnic, N. J. (2023). Female genital concealment and a

    corresponding male clasping apparatus in Australian ripple bugs (Hemiptera: Veliidae).

    Arthropod Structure & Development, *74*, 101254.

    https://doi.org/10.1016/j.asd.2023.101254

Martens, J. (2020). A harvestman with elaborate palpal pliers, *Thunbergia gretae n. gen. n.*

    sp. from China (Opiliones: Sclerosomatidae: Gagrellinae). *Zoologischer Anzeiger*, *287*,

    160–166. https://doi.org/10.1016/j.jcz.2020.03.010

Martínez Villar, M., Germil, M., Pavón-Peláez, C., Tomasco, I. H., Bilde, T., Toft, S., & Albo,

    M. J. (2023). Lack of Female Preference for Nuptial Gifts May Have Led to Loss of the

    Male Sexual Trait. *Evolutionary Biology*, *50*(3), 318–331.

    https://doi.org/10.1007/s11692-023-09606-3

Martínez Villar, M., Trillo, M. C., & Albo, M. J. (2021). Ineffective nuptial gifts suggest

    female emancipation from sensory exploitation. *Behavioral Ecology and Sociobiology*,

    75(3), 61. https://doi.org/10.1007/s00265-021-02994-6

Mazzi, D., Kesäniemi, J., Hoikkala, A., & Klappert, K. (2009). Sexual conflict over the

duration of copulation in *Drosophila montana*: Why is longer better? *BMC Evolutionary*

Biology, *9*(1), 132. https://doi.org/10.1186/1471-2148-9-132

McPherson, S., Wada-Katsumata, A., Hatano, E., Silverman, J., & Schal, C. (2022).

Nutritional condition affects tergal gland secretion and courtship success of male

cockroaches. *PLOS ONE*, *17*(8), e0271344.

https://doi.org/10.1371/journal.pone.0271344

Miller, C. W., Joseph, P. N., Kilner, R. M., & Emberts, Z. (2019). A weapons–testes trade-

off in males is amplified in female traits. *Proceedings of the Royal Society B: Biological*

Sciences, *286*(1908), 20190906. https://doi.org/10.1098/rspb.2019.0906

Myers, S. S., Buckley, T. R., & Holwell, G. I. (2016). Male genital claspers influence female

mate acceptance in the stick insect *Clitarchus hookeri*. *Behavioral Ecology and*

Sociobiology, *70*(9), 1547–1556. https://doi.org/10.1007/s00265-016-2163-6

O'Hara, M. K., & Brown, W. D. (2021). Sexual Cannibalism Increases Female Egg

Production in the Chinese Praying Mantid (*Tenodera sinensis*). *Journal of Insect*

Behavior, *34*(3), 127–135. https://doi.org/10.1007/s10905-021-09776-y

Órfão, I., Carvalho, C., Rodrigues, I., Ascensão, L., Pedaccini, M., Vicente, L., Barbosa, M.,

& Varela, S. A. M. (2023). The role of intrasexual competition on the evolution of male-

male courtship display: A systematic review. *PeerJ*, *10*, e14638.

https://doi.org/10.7717/peerj.14638

Pagel, M. (1999). Inferring the historical patterns of biological evolution. *Nature*, *401*(6756),

877–884. https://doi.org/10.1038/44766

Pandulli-Alonso, I., Tomasco, I. H., & Albo, M. J. (2022). The handsome liar: Male spiders

offering worthless gifts can benefit increasing mating duration. *Ethology*, *128*(3), 215–

222. https://doi.org/10.1111/eth.13258

Peretti, A. V., & Aisenberg, A. (Eds.). (2015). *Cryptic Female Choice in Arthropods:*

Patterns, Mechanisms and Prospects. Springer International Publishing.

https://doi.org/10.1007/978-3-319-17894-3

Perry, J. C., & Tse, C. T. (2013). Extreme Costs of Mating for Male Two-Spot Ladybird

Beetles. *PLoS ONE*, *8*(12), e81934. https://doi.org/10.1371/journal.pone.0081934

Pilastro, A., Benetton, S., & Bisazza, A. (2003). Female aggregation and male competition

reduce costs of sexual harassment in the mosquitofish *Gambusia holbrooki*. *Animal*

Behaviour, *65*(6), 1161–1167. https://doi.org/10.1006/anbe.2003.2118

Pinto da Rocha, R., Machado, G., & Giribet, G. (2007). *Harvestmen: The biology of*

Opiliones. Harvard University Press.

Plath, M., Parzefall, J., & Schlupp, I. (2003). The role of sexual harassment in cave and

surface dwelling populations of the Atlantic molly, *Poecilia mexicana* (Poeciliidae,

Teleostei). *Behavioral Ecology and Sociobiology*, *54*(3), 303–309.

https://doi.org/10.1007/s00265-003-0625-0

Prokop, P., & Okrouhlík, J. (2021). Metabolic cost of holding nuptial food gifts for male

spiders. *Ecological Entomology*, *46*(3), 684–690. https://doi.org/10.1111/een.13008

Rapkin, J., Jensen, K., Lane, S. M., House, C. M., Sakaluk, S. K., & Hunt, J. (2016).

Macronutrient intake regulates sexual conflict in decorated crickets. *Journal of*

Evolutionary Biology, *29*(2), 395–406. https://doi.org/10.1111/jeb.12794

Revell, L. J. (2012). phytools: An R package for phylogenetic comparative biology (and

other things). *Methods in Ecology and Evolution*, *3*(2), 217–223.

https://doi.org/10.1111/j.2041-210X.2011.00169.x

Rodríguez-Muñoz, R., Bretman, A., & Tregenza, T. (2011). Guarding Males Protect

Females from Predation in a Wild Insect. *Current Biology, 21*(20), 1716–1719.

https://doi.org/10.1016/j.cub.2011.08.053

Rönn, J., Katvala, M., & Arnqvist, G. (2006). The costs of mating and egg production in

    Callosobruchus seed beetles. *Animal Behaviour*, *72*(2), 335–342.

    https://doi.org/10.1016/j.anbehav.2005.10.024

Rosvall, K. A. (2011). Intrasexual competition in females: Evidence for sexual selection?

    Behavioral Ecology, *22*(6), 1131–1140. https://doi.org/10.1093/beheco/arr106

Rowe, L., & Arnqvist, G. (2012). Sexual Selection and the Evolution of Genital Shape and

    Complexity in Water Striders: Evolution of Water Strider Genitalia. *Evolution, 66*(1),

    40–54. https://doi.org/10.1111/j.1558-5646.2011.01411.x

Rueda-Solano, L. A., Vargas-Salinas, F., Pérez-González, J. L., Sánchez-Ferreira, A.,

    Ramírez-Guerra, A., Navas, C. A., & Crawford, A. J. (2022). Mate-guarding behaviour

    in anurans: Intrasexual selection and the evolution of prolonged amplexus in the

    harlequin toad *Atelopus laetissimus*. *Animal Behaviour*, *185*, 127–142.

    https://doi.org/10.1016/j.anbehav.2021.12.003

Sakaluk, S. K. (1991). Post-copulatory mate guarding in decorated crickets. *Animal

    Behaviour*, *41*(2), 207–216. https://doi.org/10.1016/S0003-3472(05)80472-5

Sakaluk, S. K. (2000). Sensory exploitation as an evolutionary origin to nuptial food gifts in

    insects. *Proceedings of the Royal Society of London. Series B: Biological Sciences*,

    267(1441), 339–343. https://doi.org/10.1098/rspb.2000.1006

Sakaluk, S. K., Duffield, K. R., Rapkin, J., Sadd, B. M., & Hunt, J. (2019). The troublesome

    gift: The spermatophylax as a purveyor of sexual conflict and coercion in crickets. In M.

    Naguib, L. Barrett, S. D. Healy, J. Podos, L. W. Simmons, & M. Zuk (Eds.), *Advances

    in the Study of Behavior* (Vol. 51, pp. 1–30). Elsevier.

    https://doi.org/10.1016/bs.asb.2018.12.001

Schaefer, H. M., & Ruxton, G. D. (2015). Signal Diversity, Sexual Selection, and

    Speciation. *Annual Review of Ecology, Evolution, and Systematics*, *46*(1), 573–592.

    https://doi.org/10.1146/annurev-ecolsys-112414-054158

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,

Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J.,

Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: An open-source

platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682.

https://doi.org/10.1038/nmeth.2019

Schlupp, I., McKnab, R., & Ryan, M. (2001). Sexual Harassment as a Cost for Molly

Females: Bigger Males Cost Less. *Behaviour*, *138*(2), 277–286.

https://doi.org/10.1163/15685390151074438

Schütz, D., & Taborsky, M. (2005). Mate Choice and Sexual Conflict in the Size Dimorphic

Water Spider *Argyroneta aquatica* (Araneae, Argyronetidae). *Journal of Arachnology*,

33(3), 767–775. https://doi.org/10.1636/S03-56.1

Sentenská, L., Šedo, O., & Pekár, S. (2020). Biting and binding: An exclusive coercive

mating strategy of males in a philodromid spider. *Animal Behaviour*, *168*, 59–68.

https://doi.org/10.1016/j.anbehav.2020.08.001

Sherman, K. J. (1983). The adaptive significance of postcopulatory mate guarding in a

dragonfly, *Pachydiplax longipennis*. *Animal Behaviour, 31*(4), 1107–1115.

https://doi.org/10.1016/S0003-3472(83)80018-9

Shine, R., Wall, M., Langkilde, T., & Mason, R. T. (2005). Do Female Garter Snakes Evade

Males to Avoid Harassment or to Enhance Mate Quality? *The American Naturalist*,

165(6), 660–668. https://doi.org/10.1086/429591

Shuker, D. M., & Kvarnemo, C. (2021). The definition of sexual selection. *Behavioral*

Ecology, *32*(5), 781–794. https://doi.org/10.1093/beheco/arab055

Shultz, J. W. (2018). A Guide to the Identification of the Harvestmen (Arachnida: Opiliones)

of Maryland. *Northeastern Naturalist*, *25*(1), 21–49.

https://doi.org/10.1656/045.025.0102

Shultz, J. W., & Pinto-da-Rocha, R. (2007). Morphology and Functional Anatomy. In R.

Pinto da Rocha, G. Machado, & G. Giribet (Eds.), *Harvestmen: The Biology of*

Opiliones (pp. 14–61). Harvard University Press.

Sigvardt, Z. M. S., Rogers, D. C., & Olesen, J. (2017). Functional morphology of amplexus

(clasping) in spinicaudatan clam shrimps (Crustacea, Branchiopoda) and its evolution

in bivalved branchiopods: A video-based analysis. *Journal of Morphology*, *278*(4), 523–

546. https://doi.org/10.1002/jmor.20649

Simmons, L. W. (2002). *Sperm Competition and Its Evolutionary Consequences in the*

Insects. Princeton University Press. https://doi.org/10.1515/9780691207032

Simmons, L. W., & Emlen, D. J. (2006). Evolutionary trade-off between weapons and

testes. *Proceedings of the National Academy of Sciences*, *103*(44), 16346–16351.

https://doi.org/10.1073/pnas.0603474103

Simmons, L. W., Lüpold, S., & Fitzpatrick, J. L. (2017). Evolutionary Trade-Off between

Secondary Sexual Traits and Ejaculates. *Trends in Ecology & Evolution*, *32*(12), 964–

976. https://doi.org/10.1016/j.tree.2017.09.011

Sirot, L. K., Wong, A., Chapman, T., & Wolfner, M. F. (2015). Sexual Conflict and Seminal

Fluid Proteins: A Dynamic Landscape of Sexual Interactions. *Cold Spring Harbor*

Perspectives in Biology, *7*(2), a017533. https://doi.org/10.1101/cshperspect.a017533

Skwierzyńska, A. M., & Plesnar-Bielak, A. (2018). Proximate mechanisms of the differences

in reproductive success of males bearing different alleles of Pgdh – a gene involved in

a sexual conflict in bulb mite. *Journal of Evolutionary Biology*, *31*(5), 657–664.

https://doi.org/10.1111/jeb.13250

Snow, S. S., Alonzo, S. H., Servedio, M. R., & Prum, R. O. (2019). Female resistance to

sexual coercion can evolve to preserve the indirect benefits of mate choice. *Journal of*

Evolutionary Biology, 32(6), 545–558. https://doi.org/10.1111/jeb.13436

Solano-Brenes, D., Costa-Schmidt, L. E., Albo, M. J., & Machado, G. (2021). Differential

allocation in a gift-giving spider: Males adjust their reproductive investment in response

to female condition. *BMC Ecology and Evolution*, *21*(1), 140.

https://doi.org/10.1186/s12862-021-01870-1

South, A., Stanger-Hall, K., Jeng, M.-L., & Lewis, S. M. (2011). Correlated Evolution of

Female Neoteny and Flightlessness with Male Spermatophore Production in Fireflies

(Coleoptera: Lampyridae): Female Flight Linked to Male Spermatophore Production in

Fireflies. *Evolution*, *65*(4), 1099–1113. https://doi.org/10.1111/j.1558-

5646.2010.01199.x

Stålhandske, S. (2002). Nuptial gifts of male spiders function as sensory traps. *Proceedings*

of the Royal Society of London. Series B: Biological Sciences, *269*(1494), 905–908.

https://doi.org/10.1098/rspb.2001.1917

Tatarnic, N. J., Cassis, G., & Siva-Jothy, M. T. (2014). Traumatic Insemination in Terrestrial

Arthropods. *Annual Review of Entomology*, *59*(1), 245–261.

https://doi.org/10.1146/annurev-ento-011613-162111

Thornhill, R. (1976). Sexual Selection and Nuptial Feeding Behavior in *Bittacus apicalis*

(Insecta: Mecoptera). *The American Naturalist*, *110*(974), 529–548.

https://doi.org/10.1086/283089

Toft, S., & Albo, M. J. (2015). Optimal numbers of matings: The conditional balance

between benefits and costs of mating for females of a nuptial gift-giving spider. *Journal*

of Evolutionary Biology, *28*(2), 457–467. https://doi.org/10.1111/jeb.12581

Toft, S., & Albo, M. J. (2016). The shield effect: Nuptial gifts protect males against pre-

copulatory sexual cannibalism. *Biology Letters*, *12*(5), 20151082.

https://doi.org/10.1098/rsbl.2015.1082

Tong, X., & Hua, B.-Z. (2019). Nuptial feeding and genital coupling of *Neopanorpa*

scorpionflies (Insecta: Mecoptera: Panorpidae) with notal organs of various lengths.

Contributions to Zoology, *88*(4), 498–512. https://doi.org/10.1163/18759866-20191409

Townsend, V. R., Bertram, M. S., & Milne, M. A. (2015). Variation in ovipositor morphology

among laniatorean harvestmen (Arachnida: Opiliones). *Zoomorphology*, *134*(3), 487–

497. https://doi.org/10.1007/s00435-015-0269-4

Townsend, V. R., Pérez-González, A., & Proud, D. N. (2019). Putative mating plugs of

harvestmen (Opiliones, Laniatores). *Zoologischer Anzeiger*, *278*, 101–109.

https://doi.org/10.1016/j.jcz.2018.12.005

Vahed, K. (1998). The function of nuptial feeding in insects: A review of empirical studies.

Biological Reviews of the Cambridge Philosophical Society, *73*(1), 43–78.

https://doi.org/10.1017/S0006323197005112

Vahed, K. (2007). All that Glisters is not Gold: Sensory Bias, Sexual Conflict and Nuptial

Feeding in Insects and Spiders. *Ethology*, *113*(2), 105–127.

https://doi.org/10.1111/j.1439-0310.2006.01312.x

Vahed, K., Gilbert, J. D. J., Weissman, D. B., & Barrientos-Lozano, L. (2014). Functional

Equivalence of Grasping Cerci and Nuptial Food Gifts in Promoting Ejaculate Transfer

in Katydids: Nuptial Gifts and Long Copulation are Analogous. *Evolution*, *68*(7), 2052–

2065. https://doi.org/10.1111/evo.12421

Voigt, C. C., Michener, R., & Kunz, T. H. (2005). The Energetics of Trading Nuptial Gifts for

Copulations in Katydids. *Physiological and Biochemical Zoology*, *78*(3), 417–423.

https://doi.org/10.1086/430224

Walter, T., & Couzin, I. D. (2021). TRex, a fast multi-animal tracking system with

markerless identification, and 2D estimation of posture and visual fields. *eLife*, *10*,

e64000. https://doi.org/10.7554/eLife.64000

Wedell, N. (1994). Variation in nuptial gift quality in bush crickets (Orthoptera:

Tettigoniidae). *Behavioral Ecology*, *5*(4), 418–425.

https://doi.org/10.1093/beheco/5.4.418

Welke, K. W., & Schneider, J. M. (2012). Sexual cannibalism benefits offspring survival.

Animal Behaviour, *83*(1), 201–207. https://doi.org/10.1016/j.anbehav.2011.10.027

Wheeler, D. (1996). The Role of Nourishment in Oogenesis. *Annual Review of Entomology*,

41, 407–431. https://doi.org/10.1146/annurev.en.41.010196.002203

Wijnhoven, H. (2011). Notes on the biology of the unidentified invasive harvestman

Leiobunum sp. (Arachnida: Opiliones). *Arachnologische Mitteilungen*, *41*, 17–30.

https://doi.org/10.5431/aramit4103

Willemart, R. H., & Hebets, E. A. (2012). Sexual Differences in the Behavior of the

Harvestman *Leiobunum vittatum* (Opiliones, Sclerosomatidae) Towards Conspecific

Cues. *Journal of Insect Behavior*, *25*(1), 12–23. https://doi.org/10.1007/s10905-011-

9268-6

Wolfner, M. F. (2002). The gifts that keep on giving: Physiological functions and

evolutionary dynamics of male seminal proteins in *Drosophila*. *Heredity*, *88*(2), 85–93.

https://doi.org/10.1038/sj.hdy.6800017

Yamane, T., Okada, K., Nakayama, S., & Miyatake, T. (2010). Dispersal and ejaculatory

strategies associated with exaggeration of weapon in an armed beetle. *Proceedings of*

the Royal Society B: Biological Sciences, *277*(1688), 1705–1710.

https://doi.org/10.1098/rspb.2009.2017

Zatz, C., Werneck, R. M., Macías-Ordóñez, R., & Machado, G. (2011). Alternative mating

tactics in dimorphic males of the harvestman *Longiperna concolor* (Arachnida:

Opiliones). *Behavioral Ecology and Sociobiology*, *65*(5), 995–1005.

https://doi.org/10.1007/s00265-010-1103-0

Zhong, W., & Hua, B. (2013). Mating Behaviour and Copulatory Mechanism in the

Scorpionfly *Neopanorpa longiprocessa* (Mecoptera: Panorpidae). *PLoS ONE, 8*(9),

e74781. https://doi.org/10.1371/journal.pone.0074781

Zweerus, N. L., Van Wijk, M., Schal, C., & Groot, A. T. (2021). Experimental evidence for

female mate choice in a noctuid moth. *Animal Behaviour*, *179*, 1–13.

https://doi.org/10.1016/j.anbehav.2021.06.022

Table 2

Metric (Category)	Definition	Expectations under High Conflict
Clasp, attempted	Male attempts to hook its pedipalps behind the female's leg II coxae.	-
Clasp, successful	Male successfully hooks its pedipalps behind the female's leg II coxae.	-
Clasping time	Time during which the male maintains a clasp with its pedipalps hooked behind the female's leg II coxae.	Increased: The length of clasping may indicate conflict if males continue to clasp females beyond the female's ideal duration. Increased clasping duration is often associated with increased sperm transfer, mate guarding, and antagonistic interactions (Arqnvist & Rowe 2005).
Contact time, pre-attempt	Any contact between individuals before a successful clasp has occurred.	Decreased: Longer pre-attempt contact times may indicate solicitous, courtship-like behaviour because it provides individuals with a greater opportunity to assess a potential mate. Shorter times provide less opportunity for assessment and courtship, and so would be expected (Dmitriew & Blanckenhorn 2012; Fowler et al. 2022)
Contact time, total	Any contact between individuals throughout the entire trial.	Increased: *Leiobunum* harvesters primarily rely on tactile and chemical senses and contact between individuals is an important communication channel. Increased association time between males and females can reduce female fitness in many species and may therefore be indicative of conflict. Alternatively, females may benefit from increased time to assess a potential mate and from beneficial male guarding (Bokides et al. 2012; Iglesias-Carrasco et al. 2019).
Distance	Distance between the female and male body centroid post-initial contact.	Decreased: An automated analog of total contact time, see above.
Female body angle	Maximum angle formed by the female's genital operculum and the substrate during a male clasping attempt or success.	Increased: The female genital opening is ventrally located, and angling the anterior portion of its body downwards may make it more difficult for the male to access the genital operculum. Genital repositioning has been documented in several other arthropods with obligate reproductive clasping (Burke et al. 2015; Golov et al. 2018; Maroni et al. 2023).
Flee count	Instances of the female rapidly moving away from male contact.	Increased: Upon contact, unreceptive females may rapidly move away from the male to avoid reproduction attempts.
Guarding time	Any instance of contact lasting greater than 5 seconds post-successful clasp. Subsequent clasps were scored as such, rather than as guarding.	Increased: Mate guarding is strongly associated with sexual conflict, allowing males to prevent remating and ensure paternity. Though guarding is not typically beneficial for females, beneficial mate guarding has been documented, albeit rarely, and could function to prevent fitness-reducing harassment from other males (Alcock 1994; Arnqvist & Rowe 2005; Sherman 1983).
Latency to attempt	Time between the initial contact and the first clasping attempt.	Decreased: Shorter latency allows for less mate assessment and fewer courtship behaviours and has been associated with high conflict in other arthropods (Dmitriew & Blanckenhorn 2012; Fowler et al. 2022).
Pursuit angle	Angle formed by the male and female velocity vectors post-initial contact.	Decreased: Angles near 0 degrees indicate the female is moving away from the male while the male follows and is indicative of the male pursuing a potentially unreceptive female. Angles near 180 degrees indicate individuals are moving towards one another, which would indicative a receptive female (Bokides et al. 2012; Iglesias-Carrasco et al. 2019).
Tap bout count	Bouts of investigatory leg taps performed during an interaction.	Decreased: *Leiobunum* Opiliones primarily rely on tactile and chemical sensing, performing investigatory taps with their antenniform second pair of legs. Tapping may be solicitous or courtship-like if it provides an assessment signal between individuals (Shultz & Pinto-da-Rocha 2007; Wulff et al. 2018).
Time within 1 leg span	Time during which the distance between male and female body centroids was less than the length of the male's longest leg.	Increased: An automated analog of total contact time, see above.