The developing Human Connectome Project fetal functional MRI release: Methods and data structures

Vyacheslav R. Karolis[1,2,*,†], Lucilio Cordero-Grande[1,3,*], Anthony N. Price[1], Emer Hughes[1], Sean P. Fitzgibbon[2], Vanessa Kyriakopoupou[1], Alena Uus[1], Nicholas Harper[1], Denis Prokopenko[1], Devi Bridglal[1], Jucha Willers Moore[1,4], Sian Wilson[1], Maximilian Pietsch[1], Daan Christiaens[1], Maria Deprez[1], Logan Williams[1,4], Emma Robinson[1,4], Antonis Makropoulos[1], Seyedeh-Rezvan Farahibozorg[2], Jonathan O'Muircheartaigh[1,4,6], Mary A. Rutherford[1], Daniel Rueckert[7,8], A. David Edwards[1], Tomoki Arichi[1,4,9], Stephen M. Smith[2], Eugene Duff[2,10], & Joseph V. Hajnal[1]

Affiliations
1. Centre for the Developing Brain, School of Biomedical Engineering and Imaging Sciences, King's College London, London, UK.
2. Wellcome Centre for Integrative Neuroimaging, FMRIB, Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, UK.
3. Biomedical Image Technologies, ETSI Telecomunicación, Universidad Politécnica de Madrid & CIBER-BBN, ISCIII, Madrid, Spain
4. MRC Centre for Neurodevelopmental Disorders, King's College London, London, UK.
5. Department of Biomedical Computing, School of Biomedical Engineering and Imaging Sciences, King's College London, London, UK.
6. Department of Forensic and Neurodevelopmental Sciences, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK
7. Biomedical Image Analysis Group, Department of Computing, Imperial College London, London, UK.
8. Klinikum rechts der Isar, Technical University of Munich, Germany.
9. Paediatric Neurosciences, Evelina London Children's Hospital, Guy's and St Thomas' NHS Foundation Trust, London, UK.
10. Department of Brain Sciences, Faculty of Medicine, Imperial College London, London, UK.

* - these authors contributed equally.
† - corresponding author: slava.karolis@kcl.ac.uk

Abstract

Recent advances in fetal fMRI present a new opportunity for neuroscience to study functional human brain connectivity at the time of its emergence. Progress in the field however has been hampered by the lack of openly available datasets that can be exploited by researchers across disciplines to develop methods that would address the unique challenges associated with imaging and analysing functional brain in utero, such as unconstrained head motion, dynamically evolving geometric distortions, or inherently low signal-to-noise ratio. Here we describe the developing Human Connectome Project's release of the largest open access fetal fMRI dataset to date, containing 275 scans from 255 fetuses and spanning the period of 20.86 to 38.29 post-menstrual weeks. We present a systematic approach to its pre-processing, implementing multi-band soft SENSE reconstruction, dynamic distortion corrections via phase unwrapping method, slice-to-volume reconstruction and a tailored temporal filtering model, with attention to the prominent sources of structured noise in the *in utero* fMRI. The dataset is accompanied with an advanced registration infrastructure, enabling group-level data fusion, and contains outputs from the main intermediate processing steps. This allows for various levels of data exploration by the imaging and neuroscientific community, starting from the development of robust pipelines for anatomical and temporal corrections to methods for elucidating the development of functional connectivity in utero. By providing a high-quality template for further method development and benchmarking, the release of the dataset will help to advance fetal fMRI to its deserved and timely place at the forefront of the efforts to build a life-long connectome of the human brain.

Keywords:

Fetal fMRI, open-access data resource, dynamic distortion correction, slice-to-volume motion correction, temporal filtering, network analysis

1. Introduction.

Already at birth, brain activity appears to be organised into a "connectome" of distributed networks (Doria et al., 2010; Fitzgibbon et al., 2020; Gao et al., 2015) that underpin complex behaviours and cognitive functions later in life. Multiple lines of evidence now also point to the critical importance of the fetal period for healthy development (Bergman et al., 2007; Boukhris et al., 2016; Brown et al., 1995; Laplante et al., 2008; O'Donnell et al., 2009; Rakers et al., 2020; Sandman et al., 2012; Skranes & Løhaugen, 2022; Zerbo et al., 2015). There is thus an increasing need for data on how functional connections become established in the prenatal brain.

Advances in fetal fMRI present an opportunity for neuroscientific studies of the functional human brain at the time of its emergence (Ferrazzi et al., 2014; Schöpf et al., 2012). Imaging an in-utero brain, however, poses numerous unique challenges (Ferrazzi et al., 2014; Seshamani et al., 2016; Sobotka et al., 2022; Taymourtash et al., 2019). Unconstrained motion, non-rigid maternal tissues surrounding the fetal brain, a greater distance between the head and receiver coil are among the MRI-adverse factors that lower signal-to-noise ratio (SNR) and increase the level of structured artifacts. Motion-free periods are empirically rare in *in utero* fMRI acquisitions given that motion associated with maternal respiration and fetal movement itself systematically induce fetal head displacements. When extreme, these displacements may prohibit the reconstruction of particular volumes in the timeseries (Sobotka et al., 2022; Taymourtash et al., 2021). When mild to moderate, they can still cause significant spin-history effects (Ferrazzi et al., 2014; Seshamani et al., 2016), manifested by spatially non-stationary image intensity changes (*travelling waves*), not readily amenable to established methods of data temporal filtering (Griffanti et al., 2014; Salimi-Khorshidi et al., 2014). The interaction between the magnetic properties of the moving fetal head and the adjacent maternal tissues induces dynamic changes in the "static" magnetic field (B0) inhomogeneity that results in temporally evolving distortion of fetal brain geometry (Cordero-Grande et al., 2018). The high contrast of maternal tissues may also induce leakage artifacts in the multi-band (MB) sensitivity encoded (SENSE) MRI data reconstruction. Overall, the *in utero* setting introduces multiple challenges to be navigated *en route* to artifact-controlled characterisation of emerging brain functional connectivity.

Finding solutions to these challenges, as well as the development of tailored methods for fetal brain connectivity analyses, requires a community-wide effort. Progress in this direction however is being hampered by the lack of openly available datasets that could be exploited by researchers across disciplines. The developing Human Connectome Project (dHCP) closes this gap by releasing the first open-access and largest-to-date fetal fMRI dataset (Data and code availability, Resource 1) of 275 scans from 255 individuals (gestational age (GA): 20.86 - 38.29 weeks), processed using tailored methods and accompanied by an advanced registration infrastructure between imaging and template spaces. The dataset complements the open-access dataset of neonatal fMRI data (Fitzgibbon et al., 2020), together allowing for detailed investigations of connectivity in the perinatal brain. In this paper we set forth details of this endeavour to process the fetal fMRI data from the stage of frequency-to-image reconstruction all the way to the level when they can be utilised for group-level analyses. To aid future method benchmarking, we make available the outputs from all the main pre-processing stages. This allows for various levels of data exploration by the imaging and

neuroscientific community, starting from the development of robust pipelines for anatomical,
distortion and temporal corrections to the analyses investigating the development of the
prenatal functional connectivity. The aim of this paper is threefold: 1) describe the
organisation of the dHCP fetal fMRI release data, 2) provide detailed descriptions and
motivations for the pre-processing methods implemented in the released data; 3)
demonstrate its capacity for performing group-level analyses. We interleave descriptions of
methods and results for a clearer exposition of analytical approaches and their contribution
to the improvement of data quality.
56
57
58   2.   Methods and Results
59
60 2.1. Acquisition parameters and data sample
61
Participants were prospectively recruited as part of the dHCP, a cross-sectional open science
initiative funded by the European Research Council and approved by the UK National
Research Ethics Authority (14/LO/1169). Written informed consent was obtained from all
participating families prior to imaging. Resting-state fMRI data were acquired with a Philips
Achieva 3T system (Best, NL) using a 32-channel cardiac coil at St Thomas' Hospital London.
Scanner software was R3.2.2 with a custom patch. Each data set consists of 350 volumes (48
slices each), acquired using a single-shot echo planar imaging (EPI) (TR/TE = 2200/60)
sequence, with slice matrix = 144 x 144, isotropic resolution = 2.2 mm, MB factor = 3, and
SENSE factor = 1.4 (Price, 2019). Brains of all fetuses were reported by a neuroradiologist as
showing appropriate appearances on the T2-weighted anatomical scan for their GA with no
acquired lesions or congenital malformations of clinical significance.
73
74 A total of 277 completed fetal fMRI scans were acquired. Two were excluded from public data
75 release due to poor data quality across all modalities (T2-weighted, diffusion, and fMRI). The
76 remaining 275 fMRI sessions were acquired from 255 unique individual subjects (137 male,
77 116 female, 2 unrecorded, GA: 20.86 - 38.29 weeks). Following an initial visual assessment, 1
78 scan was excluded from further processing due to an incomplete field-of-view.
79
With one exception, all fMRI scans were complemented by a T2-weighted anatomical scan
obtained at the same scanning session. Of these, 248 were successfully pre-processed using
the dHCP anatomical pipeline, generating brain tissue segmentations. In addition, brain masks
(released with the dCHP anatomical data) and cortical quasi-probabilistic segmentations (Data
and code availability, Resource 2) of anatomical scans generated using a 3D U-Net based tool
(Uus et al., 2023) were available for all cases with an anatomical scan.
86
87 2.2.   Overview of data processing stages and data structure
88
89 Figure 1 graphically presents the overview and inter-dependencies of processing stages,
90 derived data, and complementary data/images utilised in the process. The processing stages
91 comprise of MB-SENSE image reconstruction, dynamic distortion correction, motion
92 correction, and temporal filtering.
93

94　Of relevance, the released data contains items in two imaging spaces: the native acquisition
95　space and the anatomically consistent (tissue) space. The anatomically consistent space
96　results from distortion and motion correction processes, which involve warping, moving, and
97　rotating the original image slices, so that the resulting voxel timeseries represent the
98　temporal signal evolution at specific tissue locations.

99
100　*Figure 1. Overview of the processing stages and data structure of fetal dHCP fMRI. ${sub} and ${ses} refer to participant's*
101　*and session IDs, respectively. Combination of the paths induced by this data structure with a specific ID allows for accessing*
102　*required images in the database.*

103　## 2.3.　MB-SENSE image reconstruction
104
We used the soft SENSE reconstruction proposed as part of ESPIRIT (Uecker et al., 2014) for
considering motion or fat-shift induced model inconsistencies, which was extended to
account for MB acquisition (Zhu et al., 2016). Sensitivities were obtained from a single-band
(SB) dataset with matched readout (included in the release). Nyquist ghosting correction
parameters were obtained for all slices in the field-of-view by using the calibration
information collected with the SB data. Three image components were reconstructed with
soft SENSE, one corresponding to the target reconstruction and the other two to artefactual
information. Spatially adaptive regularisation maps (Fuderer et al., 2004) were constructed
for each image component by combining SB reconstructions and the corresponding
eigenvalue maps from ESPIRIT.
115
116　## 2.4.　Spatial Corrections
117
118　### 2.4.1.　Dynamic distortion correction
119
A major challenge of fetal imaging is overcoming the effects of fetal motion and its interaction
with the changing *in utero* and maternal environment. The presence of gas bubbles in the gut,
changes in maternal body pose during respiration and other incidental movements all cause
susceptibility-induced B0 inhomogeneities resulting in highly unpredictable spatial and
temporal signal fluctuations, particularly in the fetal brain boundaries. Therefore, spatially and

temporally resolved dynamic shot-by-shot B0 field correction is required for improved
imaging, especially when imaging at 3T where the aforementioned inhomogeneities are
amplified. As a result, the efficiency of registration-based methods (Hutter et al., 2018;
Kuklisova-Murgasova et al., 2018; Oubel et al., 2012) may be compromised in this scenario.
We instead opted to use the phase information in the reconstructed gradient echo EPI images
acquired for fMRI, as it allows direct estimation of dynamic distortion and therefore
separation of distortion and motion correction problems. As residual B0 dynamic evolution is
proportional to the phase of the observed signal, its estimation can be posed as a phase
unwrapping problem. The solution is obtained by a global phase unwrapping method based
on a 4D weighted least squares formulation (Ghiglia & Romero, 1994) with weights
constructed by the combination of magnitude information and local phase gradients. A global
method is used due to its robustness to local deviations of the phase-based distortion model
due to structural noise. Its application has been effective in removing clear distortions in
many individuals, with an example shown in Figure 2A.

Figure 2. Spatial corrections in the fetal dHCP dataset. Images are in radiological orientation (right is left). A) Distortion
correction: Visual example of image data before and after dynamic distortion corrections; B) Motion correction: TSNR
comparison between slice-to-volume (s2v) and volume-to-volume (v2v) corrections for an exemplar subject. Positive values
indicate higher TSNR for s2v; C) Motion correction: Distributions of averaged (across image) v2v TSNR, slice-to-volume (s2v)
TSNR and the difference in TSNR between slice-to-volume and volume-to-volume approaches (s2v-v2v). For the latter
positive values indicate higher TSNR for the slice-to-volume approach.

147
FMRI data can be severely compromised by motion between the acquisition of different
slices. Estimation of rigid-body fetal head motion at the slice level is challenging and
correction is ill-posed in the absence of orthogonal stacks (as are typically acquired for
anatomical imaging). To address these challenges, we have followed a multi-scale strategy.
First, volume-to-volume motion estimates are obtained by standard multiresolution
longitudinal registration. Then, these are used to initialise slice-to-volume motion estimation
and correction with motion states defined jointly for simultaneously excited slices. Motion
parameters are obtained by registering the collected information corresponding to each

motion state against a motion-compensated temporal average target. In a final step, motion
compensated reconstructions of each fMRI volume are obtained by using the conjugate
gradient algorithm for inverting a forward model of the observed fMRI data that considers
previously estimated motion parameters and second order regularisation in the slice
direction. The whole procedure is based on the framework introduced in (Cordero-Grande et
al., 2018b).
162
The advantage of using slice-to-volume alignment compared to volume-to-volume alignment
(still the default approach in much ex utero image pre-processing) can be observed both at
individual and group levels. Figure 2B shows the map of temporal signal-to-noise ratio (TSNR,
calculated as signal mean divided by its standard deviation over timeseries) difference
between both approaches for an exemplary individual. Figure 2C shows average-across-map
group-level TSNR distributions for each approach and their difference, indicating the benefit
of the slice-to-volume alignment.
170
171 ## 2.5. Native-to-template mappings
172
At this stage, the spatially corrected fMRI data were integrated into the dHCP volumetric
registration infrastructure (Figure 3), adopting an approach previously implemented for the
neonatal dHCP data release (Fitzgibbon et al., 2020). This allows users to flexibly manipulate
the data and map between template space and native spaces of all modalities. The remainder
of this section will focus on the detailed description of how this infrastructure has been built.
A reader interested in processing functional data only can proceed to Section 2.6, without
missing out on important details.
180
The building blocks of this infrastructure are: 1) linear mapping between native (i.e.,
individual) functional and T2 spaces, 2) non-linear mapping between native T2 and age-
matched template spaces, and 3) non-linear mapping between each pair of age-adjacent
templates. The first two building blocks are included in the release, whereas the between-
template mappings are available at the dHCP fetal weekly structural atlas repository (Uus et
al., 2023 ; Data and code availability, Resource 2). The age-matched template corresponded
to that of the subject's age when rounded to the nearest integer, except for subjects whose
GA was > 36.5 weeks, which were all assigned to the 36-week-old template, the oldest
template in the dHCP structural atlas.
190
All linear mappings, including those that preceded estimation of non-linear transformations,
were computed using FSL FLIRT. All non-linear warps were estimated using ANTs. The warps
were converted into the FSL *fnirt* format, which can be concatenated using the *convertwarp*
tool from the FSL library (Smith et al., 2004) in order to create composite displacement warps,
incorporating both linear and non-linear components. These composite warps would provide
mappings between distant imaging spaces while avoiding multiple image interpolation steps.
197
To lower the computational burden for the dataset users, two types of composite warps and
their inverse have been made available in the release: 1) mapping between native functional
and age-matched template spaces and 2) between native functional and 36-week-old
template spaces, which could be used for the group-level data synthesis. In addition, the
release includes a forward mapping between native functional and the 36-week-old template

space optimised for the group-level fMRI analysis, the creation of which is described Section
2.5.4. All linear and non-linear transformations between image spaces reside in
derivatives/dhcp_fmri_pipeline/sub-{sub}/ses-{ses}/xfm folders of the released data. Details of
the procedures used to obtain the required mappings are included below.

207
208
209

210
211 *Figure 3. Registration infrastructure associated with the fetal dHCP fMRI data. Concatenated transformations between*
212 *native fMRI and template spaces are part of the release. Concatenated transformations between template spaces (within-*
213 *fetal and fetal-to-neonatal) are available at g-node (Data and code availability, Resource 2).*

214
2.5.1. Native functional to structural mapping
216 For cross-modal mapping between native fMRI and T2 spaces, a modified standard
217 procedure, as implemented in the FSL's *epi_reg* tool, was used. It consists of two stages: a)
218 rigid whole-brain linear registration (using global search and mutual information as a cost
219 function), followed by b) refinement using boundary-based registration (BBR) with local
220 absolute differences cost function.

221
As the fetal brain is surrounded by maternal tissues, including high-intensity amniotic fluid
which has an intensity similar to that of cerebrospinal fluid (CSF), accurate masking is required
to ensure good alignment, particularly for whole-brain registration. The loose masks used for
motion correction of fMRI data, obtained using a spherical Markov Random Field deformable
model based on (Cordero-Grande et al., 2011), were employed to initialise alignment.
Subsequently, iterations were performed between function-to-structure alignment and back-
projection of high-quality T2 masks into the native functional space.

229
The details of the procedure are as follows. The alignment of a functional mean image to an
anatomical scan was initialised using the masks used for fMRI motion correction. The results
were visually checked for being approximately correct and, if necessary, were recomputed
after manual mask adjustments. The following steps were then repeated twice. First,

anatomical masks were projected into the functional space by inverting the mapping
obtained from the initial alignment. These new masks were applied and the N4 corrections
for intensity non-uniformity (bias field) within the new mask were applied to the mean fMRI
image. The corrected image was then used to re-compute the rigid alignment.
238
The output from the second iteration of the above procedure was then used to initialise BBR.
For the cases (N=26) which lacked the white matter (WM) segmentation required for BBR,
binarised cortical quasi-probability maps were used instead, thresholded by minimising the
difference between the volume of the thresholded maps and the volume of the cortical mask
obtained by the dHCP anatomical pipeline segmentation in the rest of the sample. The results
of whole-brain and BBR were visually checked and compared against each other by one of the
co-authors (VRK). The BBR alignment was retained if it was rated (based on visual inspection)
as performing similarly or better than the whole-brain alignment.
247
248 Out of 273 cases with an anatomical scan, 8 scans failed the function-to-structure alignment.
249 All of these failed cases had very poor MB-SENSE reconstruction quality and thus were
250 dropped from the subsequent pre-processing. The BBR alignment was retained in 254 out of
251 265 cases. Examples of the comparison between whole-brain and boundary-based
252 registrations are presented in Figure 4.
253

254
Figure 4. Example of whole-brain vs boundary-based mapping between native fMRI and T2 spaces. Images are in
radiological orientation (right is left). The outline of the native structural cortical segmentation is overlaid in cyan. The
differences between the two approaches are typically subtle, compare regions pointed by arrows following the whole-brain
registration and corresponding regions following BBR.

2.5.2. T2 to age-matched template mapping
The registrations between native T2 images and age-matched templates were initialised by
12-degrees-of-freedom linear registration using FSL FLIRT (Jenkinson et al., 2002). Non-linear
alignment was obtained using ANTs SyN diffeomorphic multi-channel registration (Avants et
al., 2011), with T2-weighted and cortical probability maps serving as moving sources (i.e., the
images to be aligned) and corresponding maps of the dHCP fetal template as the targets.
265

2.5.3. Template to template mapping

The registration procedure between age-adjacent fetal templates and between fetal and neonatal 36w templates mirrored the registration procedure for individual T2 images to the age-matched template, i.e., was initialised with FLIRT and non-linearly refined with ANTs using T2 and cortical probability maps as registration channels. The older template in each pair was used as the target.

2.5.4. Fetal-to-neonatal template mapping

To create the mappings between dHCP neonatal (Serag et al., 2012; Data and code availability, Resource 3) and fetal templates, a non-linear transformation was computed between fetal and neonatal 36-week-old templates, followed by creation of composite warps to achieve a direct mapping between arbitrary neonatal and fetal GA template spaces. For the purpose of this study, the mappings between each week fetal template and the "standard" 40-week-old neonatal template were explicitly computed (Data and code availability, Resource 2).

2.5.5. fMRI-optimised standard space for group-level analyses

To compensate for possible residual distortions or misalignment, the registration of the fMRI data to the standard group space was further optimised for group-level analyses. For this sake, mean-across-time native fMRI data were mapped into the 36w template space and a grand-average (i.e., across all subjects) of the mapped data was computed, as well as the grand averages of T2 images and WM and cortical segmentations. We then used ANTs to run a multi-channel non-linear alignment. The subject's fMRI mean and the grand-average of all subjects' fMRI means were used respectively, as moving sources and target for the first channel of alignment and the cross-subject means of T2 volumes, and WM and cortical segmentations were used as both moving sources and targets in three other channels. In other words, the deformations were driven by the fMRI image intensities, with identical structural anatomical features between the moving source and the target used to provide spatially varying (contingent on the anatomy) constraints for the scale of deformations allowed. The weight for each structural data channel was assigned to be 10% of the fMRI data channel. The obtained warps were concatenated with warps mapping functional data to the 36-week-old template, allowing for a direct mapping between native fMRI and standard fMRI-optimised spaces.

2.6. Temporal filtering

As the standards for temporal filtering of fetal fMRI are yet to be established, the development of a regression-based temporal filtering model was accompanied by an in-depth study of the effect that various types of deconfounding regressors may have on data characteristics, thereby promoting a better understanding of signal properties. We started with an identification of sources for prominent artefacts in the fetal fMRI data and split them into 3 coherent classes. We then selected candidate deconfounding regressors for each class and defined an order in which they were incorporated into a temporal filtering model. After temporal filtering, the data underwent quality assessment, identifying cases suitable for group-level analysis. Finally, we performed a post-hoc study of the unique effects that each set of deconfounding regressors had on the data metrics. For the latter, we used TSNR and seed-to-brain correlations, with the left and right thalami as seeds.

313
We define 3 broad classes of artefacts. The first class pertains to factors that may affect
spatial similarity of the volumes across time. Broadly speaking, here we deal with the effect of
imperfections in spatial corrections on the evolution of the signal in the temporal domain.
Potential artefacts attributed to this class are: a) poor motion correction in the presence of
large fetal motion; b) signal leakages not fully suppressed during MB-SENSE reconstruction; c)
residuals of dynamic distortion correction; d). low-frequency temporal drifts in signal
intensity, related to gradient system instabilities.
321
The second class of artefacts constitute motion-induced changes in the temporal evolution of
the signal, potentially causing biology-unrelated covariance structure. Within this group,
particular attention is required for spin history artefacts. Their manifestation is likely to take
the form of spatially non-stationary image intensity changes (*travelling waves*) (Ferrazzi et al.,
2014), which are particularly difficult to address using traditional spatial ICA denoising (which
presumes spatial invariance of the signal source).
328
The third class pertains to the sequential MB data sampling scheme used in the dHCP which
induces variable temporal gaps between the acquisition of two spatially adjacent slices. In all
cases except for the first and last slices from two adjacent MB stacks, this gap corresponds to
the time in between two consecutive RF excitations. In contrast, the temporal gap is on the
order of the repetition time (TR) for the slices at the edges of the MB stacks, which can result
in abrupt signal changes between them.
335
336 These broad classes of potential artefacts lead to the usage of three groups of temporal
337 deconfounding regressors, specific to each class.
338
2.6.1. Group 1. Temporal regressors for spatial inconsistency artefacts
This group consisted of detrending regressors, volume censoring ("scrubbing") regressors
(Power et al., 2012) and two novel types of voxelwise regressors, which we call folding and
density maps respectively. Together they constitute a minimal set of denoising regressors
which were used both in combination with other regressors to define more complex
denoising models and on their own to residualise the data prior to obtaining data-derived
regressors using spatial ICA (see below, Group 2 regressors).
346
347 *Detrending regressors.* These were formed by the first 10 columns of the discrete cosine
348 transform matrix, which for the current data corresponded to a high-pass filter with cut-off at
349 0.0067Hz (150 secs).
350
Volume censoring ("scrubbing") regressors. This was implemented by spike regressors
(Lemieux et al., 2007; Satterthwaite et al., 2013), which in effect equates a censored volume
to a temporal average. The criterion for censoring was based on the spatial similarity of a
volume to that average. As a metric for the latter, we utilised the root mean square
difference (RMSD) measured in terms of the ratio to the grand average of data intensity.
More precisely,
357

$$d_t = \sqrt{\frac{1}{n}\sum_{i=1}^{n}\left(\frac{f_{t,i} - M_i}{M}\right)^2}$$

where $f_{t,i}$ is the image intensity in voxel location i, $i = 1, 2, .., n$, and volume t, M_i is the temporal median of $f_{t,i}$, and M is the spatial mean of M_i.

A volume t was censored if $d_t - median(d_t) > .05$. The threshold was determined empirically in order to capture not only global perturbations in image characteristics due to excessive motion but also identifying the volumes with strong local artefacts, caused either by incomplete volume sampling or strong leakage artefacts. The volumes that immediately followed an above-threshold volume were also censored, due to higher likelihood of spin-history impacting the integrity and motion parameter estimates of the next volume (Friston et al., 1996).

Folding maps. We utilised the timecourses of the voxels that were simultaneously acquired with a target voxel as deconfounding regressors for potential leakage artefacts in that voxel. For this, we applied a positional shift to the fMRI timecourses, such that the timecourse of a voxel simultaneously acquired with the timecourse of a target voxel was assigned to the latter voxel's location. Given MB = 3, two 4D maps of this type were formed, which we called folding maps.

As noted previously, there is a distinction between acquisition and anatomically consistent spaces (as in Figure 1). For this reason, distortion and motion correction operators must be applied to the folding maps in order to bring them from the acquisition space into the anatomically consistent space. As a result, voxels in the resultant folding maps represent weighted averages of the original maps in accordance with the transformations and interpolation schema applied to the data.

Density maps. These were designed to remove temporal dependencies potentially induced by the distortion correction procedure. Specifically, in addition to spatial coordinate transformations, the distortion correction involves a Jacobian modulation which compensates for the compression/spread of the signal in the phase encoding direction. Modulation is a voxel-wise multiplicative factor which corrects image intensities and should not, in principle, be correlated with the temporal evolution of the signal after correction; if this is the case it can be assumed that it is likely to be artefactual. To generate the denoising map that carries this information, we created a 4D image of uniform intensity with geometry matched to the geometry of the data in the acquisition space. We then applied distortion- and motion-correction transformations to simultaneously bring the image into the anatomically consistent space and scale it with the applied modulation.

2.6.2. Group 2. Temporal regressors for motion-induced signal changes
This group included two types of regressors: 1) time-courses of non-grey-matter (GM) tissues (WM and CSF), derived in a data-driven manner and 2) 4D voxel-wise maps representing the evolution of motion parameters over time.

Data-derived regressors. Data-derived non-GM maps regressors (e.g., Behzadi et al., 2007)
were obtained by performing probabilistic ICA (FSL MELODIC) analysis (Beckmann & Smith,
2004) on the data within regions combining subarachnoid and ventricular CSF and WM that
were residualised with respect to Group 1 regressors. The mask combining these tissues was
eroded by one voxel to reduce partial voluming. For cases in which the dHCP anatomical data
failed the quality control assessment, missing segmented WM and CSF masks were generated
by projecting the segmentations of age-matched subjects into the native functional space of
that particular case and averaging them to produce an approximate map. Depending on the
particular denoising model (as described below), the ICA dimensionality was set to 6, 24, or
automatically determined using FSL MELODIC implementation of the Minka algorithm (Minka,
2000)/capped at 90 components (whichever was smaller).

Motion parameter (MP)-based voxelwise maps. Following slice-to-volume motion
correction, each of the 16 MB excitations (i.e., 48 slices / MB factor of 3) in a volume obtains
its own set of 6 motion re-alignment parameters. The utility of this information on the motion
evolution is compromised by the fact that they describe transformations applied to the slices
of the volumes residing in the acquisition space. In other words, each parameter is specific
here for a set of 3 slices, but not voxel-specific. To account for this, we created 4D maps for
each parameter, where all voxels within slices belonging to a particular MB group were
assigned the MP that was used to map them into the anatomically consistent space. After
that, the motion-correction operations were applied to these maps to bring them into the
anatomically consistent space. The resultant denoising map no longer contains identical
values within a slice, since the value for each voxel is derived from interpolation.

12 maps of this type were created. The first 6 corresponded to 3 rotation angles and 3
translation values and the other 6 represented corresponding differentials in the slice
direction. The slice differentials were calculated as a difference $S_{t+1} - S_t$, obtained by
rearranging the MP timeseries in the anatomically consistent space, $V_1, V_2, \dots, V_{350}$ (where
any V_k is a $n \times 1$ vector, n is the number of voxels, and 350 is the number of timepoints),
into the simultaneously excited slices-to-slices timeseries $S_1, S_2, \dots, S_{350 \times 16}$ (where any S_k is
a $n/16 \times 1$ vector, with 16 the number of excitations in the acquisition of a volume, i.e.,
they comprise slices in the same MB excitation). The reason for creating maps of
simultaneously excited slices' differentials was to enrich MP-based regressors with dynamics
in the intrinsic dimension for spin-history effects.

The above maps were used to create a broader set of MP-based regressors, unique for each
voxel. By defining the temporal evolution of the set of 6 MP and their differentials in the slice
direction at a voxel in location i as M_6^i and $\Delta^s M_6^i$, respectively, and first- and second-order
differentials in volume-to-volume direction as Δ^v and Δ_2^v, respectively, the complete set of 60
regressors for each voxel was defined as $cat\left\{M_6^i, \left(M_6^i\right)^2, \Delta^v M_6^i, \left(\Delta^v M_6^i\right)^2, \Delta_2^v M_6^i, \left(\Delta_2^v M_6^i\right)^2,\right.$
$\left.\Delta^s M_6^i, \left(\Delta^s M_6^i\right)^2, \Delta^v(\Delta^s M_6^i), \left(\Delta^v(\Delta^s M_6^i)\right)^2\right\}$, where $cat\{\}$ denotes column-wise
concatenation. This set was transformed first by z-scoring each column and then by using
principal component analysis, considering only non-censored volumes, with the number of
components retained to account for 99% of variance in the regressor set.

Selection of Group 2 regressors – signal implanting test. Having a broad choice of Group 2
regressors potentially creates an overfitting problem, especially when considering that the
temporal filtering model can combine both MP-based and data-derived regressors. Overly
aggressive temporal filtering can inadvertently remove the biologically relevant signal
fluctuations of interest, which could seriously hamper further analysis given the inherently
low SNR of fetal fMRI data, especially at the single-subject level.

To guide the selection of the best performing denoising models, we designed a surrogate test,
which we call the signal implanting test. This test is based on injecting a biologically-plausible
signal into the data, and observe which composition of temporal regressors allows for: 1)
most accurate recovery of this signal and 2) maximally preventing signal loss. For this, we
utilised the neonatal group-level maps (Fitzgibbon et al., 2020) and their timecourses,
estimated by regressing the group maps against individual data in the neonatal subjects from
the dHCP neonatal cohort, aged between 37 and 43 weeks. The timecourses were paired at
random with the fetal subjects and the group maps were projected into the fetal native
functional spaces. The product of the timecourses and the group maps was then scaled to
either 3% or 6% of the temporal standard deviation of the real data for each particular
subject and added to the data, constituting an "implanted signal". The data were then
residualised with respect to a candidate set of denoising regressors, and the spatial maps
were regressed against the denoised data to obtain estimates of their timecourses; the
correlation between their vectorised product and vectorised implanted signal were taken as a
single measure of the quality of signal recovery. Similarly, the same procedure was applied to
the timecourses that were removed from the data with implanted signal during denoising. A
high correlation between removed and implanted signal would indicate that the model
inadvertently removes signal from the data. We will refer to the latter measure as signal loss
statistics.

We tested several models that differed in the number of data-derived and MP-based
regressors (Figure 5A). All candidate models were complemented with Group 1 regressors.
Figure 5B shows the results of the test. Overall, the analysis indicates that MP-based
denoising models with a limited number of non-GM data-derived regressors performed best
with respect to the signal recovery statistics. Considering both signal recovery and signal loss
statistics, the MP-based model with 6 non-GM regressors (highlighted) performed the best.
Firstly, it performed better than the others with respect to signal recovery statistics. Secondly,
the model also showed a similar performance to the model with 0 non-GM regressors with
respect to signal loss statistics whereas the model with 24 non-GM regressors showed
significant losses.

In three cases, the application of the best-performing model to the real data resulted in a
numerical overflow. These cases were of a very poor quality and had a large number (> 210)
of volumes to censor; for this reason, no attempt was made to correct the issue
retrospectively. Consequently, the fully processed dHCP dataset consists of 263 sessions (247
unique individual subjects, 132 male, 113 female, 2 unrecorded).

Visual inspection of the filtered data reveals a notable reduction of the travelling waves
pattern, indicating that MP-based voxel-wise regressors are effective in ameliorating the spin-
history effects. Quantification of this phenomenon is not straightforward, but it can be

observed qualitatively when temporally demeaned data are rearranged into slices-to-slices timeseries (Figure 5C).

Figure 5. Group 2 regressors. A) Scope of temporal denoising models assessed using the signal implanting test. "F" stands for the "full" set for a particular (MP-based or data-derived) type of regressors. Correspondingly, the "F/0" model includes full set of MP-based and no data-derived regressors, the "0/F" model includes the full set of data-derived and no MP-based regressors, with intermediate models in between. The "greedy" model includes maximal number of regressors of both the MP-based and data-derived type. The sample distributions for the number of regressors in the extreme and greedy (F/F) models are plotted underneath, calculated for 266 scan sessions that passed the function-to-structure alignment. As the full set of data-derived regressors is estimated from the data, implanting different volumes of signal (3% or 6%) may change the output of the Minka algorithm/capping at 90. The sample distribution of MP-based regressors is calculated from the spatial means, given that .99 variance may be represented by a different number of components across voxels. B) Results of the signal implanting test. Model ordering as in A. Left - signal recovery statistics. Right - same for the signal loss statistics. The best performing model is highlighted. C) Segment of slices-to-slices timeseries before (Group1 regressors only) and after application of Group 2 regressors. Ticks along x-axis mark TRs, i.e., 16 slices. Note the vertical stripes, repeating at the rate of approximately 2-3 TRs and likely to be caused by mother's breathing cycle, that are significantly reduced after application of Group 2 regressors.

2.6.3. Group 3. Temporal regressors for sampling scheme artefacts

A set of regressors addressing artefacts related to the difference in temporal separation between first and last slice of adjacent MB stacks compared to other neighbouring slices were derived using single-subject spatial ICA with dimensionality set to 30 components run on the timeseries following their residualisation with respect to the model that included Group 1 and (best performing) Group 2 regressor sets. Prior to this, the data were slightly smoothed using a FWHM=1 mm 3D Gaussian filter. Here we sought to identify slab-like spatial patterns

repeating roughly across the MB stacks width (i.e., 16 slices). Two observers (VRK and DB)
independently performed the rating of spatial component maps, with three scores allowed:
'to remove', or 'to be equivocal', or 'to keep'. One observer started the rating from the first
subject in the subject list, and the other started the rating from the middle subject. To ensure
that similar subjective criteria were applied across the cohort, each observer used the rating
of the first 20 subjects to get accustomed to the task. The ratings for these subjects were
reviewed again after the observer completed the rating of the whole dataset. The criterion
for consensus to regress out component timecourses was defined as either 'to remove' by
both raters or 'to remove' by one and 'to be equivocal' by the other.
529
Figure 6A shows examples of components that were rated by consensus as representing a
sampling artefact. The average number of such components per subject was 7.6, sd = 3.96
(Figure 6B). Residualisation of the data with respect to the timecourses of these components
was performed on top of residualisation with respect to Group 1 and Group 2 regressors.
Because the timecourses of the independent components are not constrained to be
orthogonal, the regression coefficients for the timecourses to regress out were estimated in
combination with other components' timecourses. This procedure generated the most
thoroughly processed data in the release.
538

539
540 *Figure 6. Group 3 regressors. Images are in radiological orientation (right is left). A) Examples of components with double-*
541 *slab-like spatial structure in an exemplary subject; B) Distribution of the number of components rated to be removed per*
542 *subject.*

543 ### 2.6.4. Quality control and data selection for group-level analyses
544
545 To rate the quality of individual pre-processed data and select cases suitable for group-level
546 analyses, we used both visual assessment and parametric measures.
547
548 For visual assessment, the timeseries were rated by an observer (VRK) using a 5-point scale, 0
549 – fail, 1 – bad quality (seemingly unusable), 2 – seemingly usable, 3 – reasonably good, 4 –
550 good. This was complemented with the two parametric measures, mean (over voxels) *tsnr*
551 and *dvars*, disregarding censored volumes. Specifically, mean-over-voxels *tsnr* was defined as:
552

553
$$tsnr = \frac{1}{n}\sum_{i=1}^{n}\frac{M_i}{SD_i}$$

554   where:

$$M_i = \frac{1}{T}\sum_t f_{t,i}$$

$$SD_i = \sqrt{\frac{1}{T}\sum_t (f_{t,i} - M_i)^2}$$

557     $$t \in R^T$$

559   and

$$dvars = \frac{1}{N}\sum_t \sqrt{\frac{1}{n}\sum_{i=1}^{n}(f_{t,i} - f_{t-1,i})^2}$$

$$t, t-1 \in R^T$$

where t is the order number of a volume, R^T is the set of T uncensored volumes, N is the
number of consecutive pairs of uncensored volumes, $t, t-1 \in R^T$, and n is the number of
voxels. Both measures were calculated in the functional native space, within a binarised
anatomical segmentation image, which provides a tight mask by excluding non-brain tissues
(e.g., subarachnoid CSF).

The decision rule was as follows: all cases rated >1 (high score) were considered as suitable
for analysis EXCEPT when they were EITHER positive *dvars* outliers OR negative *tsnr* outliers
with respect to all cases rated > 1. Specifically, the outlier O_{dvars} for *dvars* and O_{tsnr} for *tsnr*
were defined as:

$$O_{dvars} > P_{75} + 1.5IQR$$
$$O_{tsnr} < P_{25} - 1.5IQR$$
where P_{25} and P_{75} are 25[th] and 75[th] percentiles of the cases rated > 1, and IQR is their
interquartile range.

Furthermore, all cases rated <=1 (low score) were also considered as suitable for an analysis,
if BOTH their *dvars* AND *tsnr* were, respectively, $< P_{75}$ AND $> P_{25}$ of cases rated > 1. All
ratings as well as their *dvars* and *tsnr* measures are available at Data and code availability,
Resource 4.

Figure 7A-B shows that visual assessment of the pre-processed data, *dvars* and *tsnr* were all in
reasonable agreement. Based on the predetermined criteria, 217 out of 263 scans passed the
QC (114 male, 101 female, 2 unrecorded). Figure 7C demonstrates an example of the QC
report, available for each individual subject (Data and code availability, Resource 4).

587
588 *Figure 7. Quality assessment. A) Average dvars and tsnr for high (> 1) and low (<=1) scores in visual quality assessment. B)*
589 *Relationship between average tsnr and dvars; C) QC report for an exemplar subject, showing carpet plots of z-scored*
590 *intensity, spatial profiles of TSNR and temporal evolution of dvars for the data after motion and distortion correction (type:*
591 *mcdc, without censoring) and for the fully processed data (type: preproc, including censoring).*

592 ### 2.6.5. Group-level statistics of deconfounding regressor groups
593
Figure 8 shows the summary statistics of all 263 fully pre-processed cases for the temporal
filtering regressor groups with respect to the loss of the effective degrees-of-freedom (DOFs,
= number of regressors, Figure 8A) and TSNR (Figure 8B). On average, the fully pre-processed
data lose approximately 32% of effective DOFs, and gain more than 2 times TSNR, compared
to the data that underwent spatial distortion and slice-to-volume motion correction only. The
loss in DOFs is significantly higher than reported in adult imaging (e.g., approx. 15% for
minimally pre-processed data in the Human Connectome Project (HCP) - Smith et al., 2013),
but much of it occurs via volume censoring, which has no analogy in HCP.
602
In order to evaluate the local effects of the temporal filtering models, we utilise the dataset
capacity for group-level analyses, afforded by the released registration infrastructure,
projecting TSNR maps (smoothed with a FWHM = 3 mm Gaussian kernel in the native space)
into the fMRI-optimised group space. The aligned maps were then modelled with linear
regression, with demeaned age as a co-variate, providing estimates of average TSNR
characteristics in the sample and their association with age.
609
The mean TSNR maps (Figure 8C) and the maps of the TSNR differences (Figure 8D) between
temporal filtering models help appreciate the effect of each regressor group. The patterns are

presented in terms of fitted beta coefficients for intercept and after conversion into t-
statistics for both intercept and age effect. In terms of beta coefficients, both the average
TSNR and its gain from application of the more complex denoising model was the highest in
WM, which reflects the fact that mean intensity (i.e., numerator in the TSNR formula) is
higher in the WM than in the GM and CSF. This points to the multiplicative effect of noise,
supported by the observation that WM TSNR is no longer greater than GM TSNR when
converted into t-statistic values, taking into account across-subject variability.
619
As TSNR is not informative with respect to the effects of deconfounding regressors on
estimates of brain functional connectivity, we investigated the latter in the context of seed-
to-brain correlation maps, using the left and right thalami as seeds. The analysis was
performed in the sample of individuals GA > 24.5, which passed quality control criteria. This
comprised 201 subjects (106 male, 93 female, 2 unrecorded). Individual correlation maps
were computed for the left and right thalamus separately in the native functional space
(following FWHM = 3 mm smoothing) and projected into left-right symmetrised space (Data
and code availability, Resource 4) for group-level analysis. To enable computing of the left-
right subject's average maps, the maps of the left thalamus connectivity were left-right
flipped. We then modelled the average maps using linear regression with demeaned age as a
covariate.
631
Figure 8E-F shows the results for the fitting. In summary, using a more complex denoising
model results in a decrease of correlations throughout the brain. However, perhaps with
exception for Group 1 regressors (spatial correlation -0.20 between "mcdc" and "Group1-
mcdc" maps in Figure 8E), this decrease was proportionally scaled, i.e., more prominent in
areas showing a stronger correlation to the seed timecourse prior to model application
(spatial correlation -0.95 between "Group1" and "Group1&2-Group1" maps and -0.73
between "Group1&2" and "Group1&2&3-Group1&2" maps). A similar proportional effect was
observed for the age effect (Figure 8F; spatial correlations = -0.04, -0.92 and -0.62,
respectively). The most profound decrease for both intercept and age-effect was observed
when including Group 2 regressors.
642
643
644

2.7. Group-ICA analysis.

To demonstrate the dataset capacity for multi-variate network analysis, we performed
estimation of functional modes ("networks") in the same sample as in the analysis of thalamic
connectivity. All individual data were smoothed in native space using a FWHM=3 mm
Gaussian kernel. To account for heterogeneity inherent to a sample with large maturational
differences, we first concatenated the timeseries of age-matched individuals (as defined for
age-matched template registration) and compressed them to 1400 principal components for
each age. After transforming within-component (across voxels) values by z-scoring, the
components were concatenated with similarly defined components from other age groups
and compressed to a sample-average set of 1400 principal components which was then fed
into FSL melodic ICA to obtain whole-sample decomposition into 25 components. In other
words, the procedure ensured a balanced contribution of each age to the final factorisation,
whereas z-scoring of values within each component ensures that factorisation attends to
common patterns irrespective of their absolute scale, thereby implicitly taking into account
that these patterns may have different prominence across ages.

Figure 9A shows functional modes obtained in these data, which predominantly comprise of
components localised to one brain region (hereafter - dominant node). Despite this, there
was a notable prevalence of component pairs (7 pairs, grouped together in the figure), where
dominant nodes are a left-right reflection of each other. Out of these 7 pairs, 3 pairs (ICs 14 vs
21, 10 vs 16, and 6 vs 19) showed symmetry not only in the location of a dominant node, but
also in the location of secondary non-negligible nodes. Another notable observation is that 3
components (ICs 15, 20, and 24, highlighted in bold green in Figure 9A) were characterised by
a node pair, one dominant and another smaller, located in the homologous regions of the two
hemispheres, and another (IC 3) showed one medial component with approximately equal
bilateral representation.

To evaluate the structure of whole-brain functional architecture, we derived the temporal
time courses of the obtained 25 ICs by spatial regression (Beckmann et al., 2009) against
individual fMRI data in native functional space. For each subject individually, we computed
the correlation matrix between component timecourses, excluding timecourses of IC0 which
is of a distinctively vascular origin. We then combined matrices from all subjects in order to fit
each element with linear regression, using demeaned age as a covariate. The procedure
resulted in two matrices, one containing estimates of an average strength of association
between each pair of components' timecourses and the other providing estimates of their
age-related changes. The elements of these two matrices were converted into t-values,
thereby accounting for between-subject variability. Normalised Laplacian embedding of each
matrix (positively thresholded to make this estimable) was then used to represent the
relationships between components in terms of a Euclidean distance between their

coordinates in the embedded space. We then applied hierarchical clustering using the Ward
method (Ward, 1963), based on the coordinates in the first 3 non-null dimensions.
696
The results of clustering (Figure 9B) demonstrate a weakness of association between
contralateral nodes in terms of average strength of their correlation; the overall tendency for
this metric was to form clusters of unilateral nodes, sometimes representing a symmetrical
image of each other (e.g., ICs 17,22,18 vs ICs 8,13,23 and ICs 6,10 vs ICs 16,19) and/or
representing coherent functional systems, such as right dorsal fronto-parietal (ICs 7,14), left-
dominated sensori-motor (ICs 2,3,20,21), left (ICs 5,9,12) and right (ICs 1,4,11) frontal.
703
704 Clusters that mix both uni- and contralateral components are more evident in the analysis of
705 age-related changes (Figure 9C). Of particular interest is a cluster (ICs 10, 15, 16, 19, 24) that
706 combines areas belonging to semantic and language processing. Notably, it includes only a
707 left but not right anterior temporal node, which aligns well with a previous report on
708 connectivity of these areas in the adult brain (Hurley et al., 2015).

709
710 *Figure 9. Modes of covariation (fetal "resting-state networks"). Images are in radiological orientation (right is left). A)*
711 *Spatial maps grouped by their anatomical location. The dotted-line boxes show pairs of maps showing a distinctive*
712 *interhemispheric symmetry. Components with non-negligible bilateral homotopic representation are highlighted in bold*
713 *green. B) Full (lower triangle)- partial (upper triangle) correlation matrix and hierarchical clustering based on full*
714 *correlations transformed into t-statistics. C) Same as B) for the age-related changes in correlations.*

715
716

3. Discussion

Despite its rapid progress (De Asis-Cruz et al., 2021; Ferrazzi et al., 2014; Jakab et al., 2014; Ji et al., 2022; Karolis et al., 2023; Schöpf et al., 2012, 2014; Seshamani et al., 2016; Sobotka et al., 2022; Taymourtash et al., 2021, 2023; Thomason et al., 2015; van den Heuvel et al., 2018), fetal fMRI remains a relatively novel field, where pre-processing standards are yet to be established and which until now lacked an open-access resource available to researchers. In this paper, we introduce an open-access fetal fMRI data repository and present a systematic approach for pre-processing, spanning stages from image reconstruction to group-level analysis. We implemented state-of-the-art methods for the spatial reconstruction and correction of the data, including MB- SENSE reconstruction, dynamic distortion and slice-to-volume corrections, and a tailored temporal filtering model with attention to the prominent sources of structured noise. The final dataset consists of 263 fully processed cases, with 217 cases determined to be suitable for the connectivity analyses. We thereby provide both a resource and a high-quality template to facilitate further development and benchmarking of methods for fetal fMRI. In the remainder of the discussion, we focus on the potential areas for improvement.

Studying fetal samples using fMRI is perhaps an exemplary case where the age of a subject simultaneously represents the most important variable-of-interest and the most prominent confound. The latter manifestations encompass, to name a few, signal-altering differences resulting from rapidly changing tissue properties, drastically different morphology, and three-fold differences in effective resolution. These factors entail various challenges for the progress of the fetal fMRI field. Firstly, they create obstacles for reproducible experiments. The standard practice of scanning one participant repeatedly (Duff et al., 2022; Noble et al., 2017; Shah et al., 2016; Shehzad et al., 2009) in fetal imaging settings is practically complicated and ethically dubious, as this approach can render meaningful results only if there is no significant temporal gap between scanning sessions. Secondly, they entail difficulties in defining sound quantitative metrics, appropriate for the entire age range, that would enable comparative testing between various processing approaches. Finally, they also complicate the inter-subject data fusion for group-level analysis and defining ground truths that could guide pre-processing approaches in order to harmonise data quality across ages and fit a broad range of potential goals for a study. As an example, the immature fetal WM is thought to contain its own functional units within temporary developmental structures such as the subplate and constitutes a metabolically active area, especially at younger ages (Colonnese & Phillips, 2018; Tolner et al., 2012). Consequently, studying the WM BOLD signal at younger ages may require a different approach to the temporal filtering problem.

Our approach to the temporal filtering follows the trend that was set by the seminal work by (Power et al., 2012) and (Satterthwaite et al., 2013), evaluating the effect of motion on resting-state connectivity in adults. This work demonstrated an improved precision of connectivity estimates following application of sophisticated denoising models enriched with motion-related or data-derived deconfounding regressors (Power et al., 2014; Pruim et al., 2015; Satterthwaite et al., 2013; Yan et al., 2013). Here, by virtue of using the recovery of an artificially implanted cortical signal as a metric to assess efficiency of temporal filtering models, we sought to optimise the data for the analysis of cortical networks. We were

conscious of the possibility that the usage of complex models may inadvertently remove
relevant signal information, i.e., that selectivity may be prioritised at the expense of
sensitivity, which may be suboptimal in the fetal fMRI low-SNR setting. Namely, we found that
the data-derived regressors (obtained using ICA on non-GM timeseries), which represent a
popular choice for *ex utero* data (Kiviniemi et al., 2003; Kochiyama et al., 2005; Salimi-
Khorshidi et al., 2014) including paediatric data (Fitzgibbon et al., 2020), are particularly prone
to undesired implanted signal removal in the fetal fMRI.

Correspondingly, our approach to temporal filtering relied on a rich set of MP-based
regressors, helping us to achieve a notable reduction of artefacts, including difficult-to-tackle
spin-history artefacts. In spite of taking these measures, we observed proportionally scaled
decreases in TSNR and estimates of thalamic connectivity following the application of
complex MP-enriched models. These appear to suggest that in real data signal and noise do
not fully comply to the additivity assumption of the signal implanting test. The proportional-
like decrease in the strength of the age effect following application of MP-based and data-
derived regressors points to the same possibility. A case in point here is the age-related
changes in connectivity between the thalamus and sensorimotor and parietal areas. On the
one hand, age-related increases in these functional connections are expected based on
current knowledge of prenatal brain development from post-mortem studies (Kostović et al.,
2019; Kostović & Judaš, 2010). On the other hand, these changes are best detectable in the
data before any temporal filtering, indicating contribution of age-related structured noise to
this contrast. Taken together, these findings suggest several far-reaching implications. Firstly,
they suggest that magnitude of the structured noise may scale with age and therefore the
strength of the age effect cannot be accepted as a sole criterion for optimisation of the model
for temporal filtering. Secondly, in the absence of a ground truth, the intricate entanglement
of signal and noise complicates the determination of the temporal filtering complexity for a
right sensitivity-selectivity trade-off. Thirdly, more sophisticated motion modelling could be
required to increase selectivity of temporal regressors to structured noise.

We therefore invite dataset users to develop temporal filtering models that could better fit
the requirements of their analyses and/or investigate the ways to boost their statistical
power, including alternatives not considered in this paper, such as more aggressive spatial
smoothing and lowpass temporal filtering. To facilitate this, we release a rich set of
complementary data in the form of denoising maps and provide Python code that can be
used with the regressors in the released database or adapted to include new ones (Data
availability statement, Resource 4). We also remark that the implementation of data
reconstruction and spatial corrections was developed to be compatible with proprietary
format of raw data collected by the scanner used in the dHCP, considered not only fetal fMRI
but many other MRI scanning domains, and was optimized for the local computing hardware
infrastructure. Therefore, we have decided not to provide a ready-to-use code release for
data reconstruction and spatial corrections to be run externally (Data and code availability,
Resource 4).

Another area for improvement relates to incorporation of deep learning models for improving
brain extraction, spatial corrections and temporal denoising in the proposed pipeline, with
recent approaches mature enough, in particular, for the brain extraction problem (e.g.,
(Rutherford et al., 2022)). A particularly challenging scenario is encountered in scans where

the brain's orientation varies significantly in time. In the current implementation, such differences could result in an orientation flip in some volumes of the reconstructed timeseries, which had to be censored for downstream processing. The extreme situation where a fetus may turn upside-down within an fMRI scan is far from being manageable with current analysis techniques, which exemplifies that data rejection criteria and quality control also merit future efforts for improvement and standardisation across studies.

Finally, detailed investigations are also needed to determine how the large age-dependent variability in brain characteristics can analytically be incorporated into building group-level normative models of functional brain development. Fetal fMRI data reveals idiosyncratic properties (Karolis et al., 2023), apparent in various forms, including when results of group-level ICA factorisation between fetal and neonatal samples are contrasted. Analysis in the fetal brain typically reveals a dominance of a single-node functional modes and a paucity of spatially distributed patterns (Ji et al., 2022; Karolis et al., 2023), whereas in the neonatal brain the latter are not uncommon, especially those left-right symmetrically organised (Eyre et al., 2021; Fitzgibbon et al., 2020; Gao et al., 2015). This may reflect a genuine property of the fetal brain's functional organisation, its immaturity and perhaps the functional state of the brain while *in utero*, but may also simply be a reflection of a lower signal-to-noise ratio or may suggest that the method, that implicates estimation of a group-level 'mean' model in a developmentally heterogeneous sample, is not appropriate for these data (Karolis et al., 2023). In this paper we adopted pre-processing steps in group-ICA analysis that aimed to ameliorate the challenges of data fusion across ages. The obtained results were more prone to revealing signatures of distributed spatial patterns than previously reported (e.g., Ji et al., 2022; Karolis et al., 2023). Furthermore, they reveal a range of symmetrically located network pairs, tentatively pointing to an emergence of a bilaterally organised functional architecture (Thomason et al., 2013). However, despite these elements of qualitative convergence with the neonatal data, harmonisation of in- and ex-utero cohorts for a combined analysis remains a task for future research.

In conclusion, research into functional brain development in utero is strongly motivated by a large body of evidence that shows the fetal period has critical importance for individual development. This information can be obtained non-invasively using fetal fMRI, but much work is required in order to establish the standards of data acquisition, pre-processing and analysis in this challenging domain. The current dataset represents a unique resource that aims to provide a firm foundation for advancing fetal fMRI from its current status as a niche research field to its deserved and timely place at the forefront of the community-wide efforts to build a life-long connectome of the human brain.

Data and code availability.

The data and code presented in this paper are available in the following locations:
1. dHCP fetal release: https://nda.nih.gov/edit_collection.html?id=3955
2. The dHCP volumetric template can be accessed at: https://gin.g-node.org/kcl_cdb/fetal_brain_mri_atlas. Folder "structural" contains T2-weighted spatiotemporal atlas, folder "composite_warps" contains fsl-style warps between different ages of template; folder "composite_warps_fetal2neonatal" contains

mappings between fetal template and the 2 neonatal templates spaces. Folder
"cnn_cortex_probability" contains quasi-probability maps of cortical segmentations.
3. The neonatal templates and mappings between different ages are available at:
https://git.fmrib.ox.ac.uk/seanf/dhcp-resources
4. Other materials associated with this paper, such as sample demographics,
supplementary data, QC reports, and the python code used for performing temporal
filtering, are available at: https://gin.g-
node.org/kcl_cdb/dHCP_fetal_fMRI_release_paper. Enquiries about sharing the
existing code for image reconstruction, distortion and motion corrections and
guidance on its application should be directed to co-author LCG.

Acknowledgments

The Developing Human Connectome Project was funded by the European Research Council
under the European Union Seventh Framework Programme (FP/20072013)/ERC Grant
Agreement no. 319456. The Wellcome centre for Integrative Neuroimaging is supported by
core funding from the Wellcome Trust [203139/Z/16/Z]. The authors also acknowledge
support in part from the Wellcome Engineering and Physical Sciences Research Council
(EPSRC) Centre for Medical Engineering at King's College London [WT 203148/Z/16/Z], the
Medical Research Council (MRC) Centre for Neurodevelopmental Disorders [MR/N026063/1],
and the Department of Health through an NIHR Comprehensive Biomedical Research Centre
Award (to Guy's and St. Thomas' National Health Service (NHS) Foundation Trust in
partnership with King's College London and King's College Hospital NHS Foundation Trust).
VRK and TA were supported by a MRC translation support award [MR/V036874/1]. VRK also
acknowledges support by EEU-AIMS (European Autism Interventions)/EU AIMS-2-TRIALS, an
Innovative Medicines Initiative Joint Undertaking under Grant Agreement No. 777394. TA was
also supported by a MRC Clinician Scientist Fellowship [MR/P008712/1] and a MRC Senior
Clinical Fellowship [MR/Y009665/1]. JOM is supported by a Sir Henry Dale Fellowship jointly
funded by the Wellcome Trust and the Royal Society [206675/Z/17/Z]. LCG acknowledges
funding from MCIN/AEI/10.13039/501100011033/FEDER, EU under Project PID2021-
129022OA-I00 as well as Universidad Politécnica de Madrid for providing computing
resources on Magerit Supercomputer. SRF's research is supported by the Royal Academy of
Engineering under the Research Fellowship programme [RF2122-21-310]. JWM was
supported by PhD funding from the UK Medical Research Council [MR/P502108/1].

References:

Avants, B. B., Tustison, N. J., Song, G., Cook, P. A., Klein, A., & Gee, J. C. (2011). A
reproducible evaluation of ANTs similarity metric performance in brain image
registration. *NeuroImage*, *54*(3), 2033–2044.
https://doi.org/10.1016/j.neuroimage.2010.09.025
Beckmann, C. F., & Smith, S. M. (2004). Probabilistic independent component analysis for
functional magnetic resonance imaging | IEEE Journals & Magazine | IEEE Xplore.
IEEE Transactions on Medical Imaging, *23*(2), 137–152.
https://doi.org/10.1109/TMI.2003.822821

Beckmann, C., Mackay, C., Filippini, N., & Smith, S. (2009). Group comparison of resting-state
FMRI data using multi-subject ICA and dual regression. *NeuroImage*, *47*, S148.
https://doi.org/10.1016/S1053-8119(09)71511-3
Behzadi, Y., Restom, K., Liau, J., & Liu, T. T. (2007). A component based noise correction
method (CompCor) for BOLD and perfusion based fMRI. *NeuroImage*, *37*(1), 90–101.
https://doi.org/10.1016/j.neuroimage.2007.04.042
Bergman, K., Sarkar, P., O'Connor, T. G., Modi, N., & Glover, V. (2007). Maternal Stress
During Pregnancy Predicts Cognitive Ability and Fearfulness in Infancy. *Journal of the*
American Academy of Child & Adolescent Psychiatry, *46*(11), 1454–1463.
https://doi.org/10.1097/chi.0b013e31814a62f6
Boukhris, T., Sheehy, O., Mottron, L., & Bérard, A. (2016). Antidepressant Use During
Pregnancy and the Risk of Autism Spectrum Disorder in Children. *JAMA Pediatrics*,
170(2), 117. https://doi.org/10.1001/jamapediatrics.2015.3356
Brown, A. S., Susser, E. S., Lin, S. P., Neugebauer, R., & Gorman, J. M. (1995). Increased Risk
of Affective Disorders in Males after Second Trimester Prenatal Exposure to the
Dutch Hunger Winter of 1944–45. *British Journal of Psychiatry*, *166*(5), 601–606.
https://doi.org/10.1192/bjp.166.5.601
Colonnese, M. T., & Phillips, M. A. (2018). Thalamocortical function in developing sensory
circuits. *Current Opinion in Neurobiology*, *52*, 72–79.
https://doi.org/10.1016/j.conb.2018.04.019
Cordero Grande, L., Price, A., Ferrazzi, G., Hutter, J., Christiaens, D., Hughes, E., & Hajnal, J.
(2018). Spin And Field Echo (SAFE) dynamic field correction in 3T fetal EPI.
Proceedings of International Society for Magnetic Resonance Imaging, 26, 208.
https://archive.ismrm.org/2018/0208.html
Cordero-Grande, L., Hughes, E. J., Hutter, J., Price, A. N., & Hajnal, J. V. (2018). Three-
dimensional motion corrected sensitivity encoding reconstruction for multi-shot
multi-slice MRI: Application to neonatal brain imaging: Aligned Multi-Shot Multi-Slice
MRI. *Magnetic Resonance in Medicine*, *79*(3), 1365–1376.
https://doi.org/10.1002/mrm.26796
Cordero-Grande, L., Vegas-Sánchez-Ferrero, G., Casaseca-de-la-Higuera, P., Alberto San-
Román-Calvar, J., Revilla-Orodea, A., Martín-Fernández, M., & Alberola-López, C.
(2011). Unsupervised 4D myocardium segmentation with a Markov Random Field
based deformable model. *Medical Image Analysis*, *15*(3), 283–301.
https://doi.org/10.1016/j.media.2011.01.002
De Asis-Cruz, J., Andersen, N., Kapse, K., Khrisnamurthy, D., Quistorff, J., Lopez, C., Vezina,
G., & Limperopoulos, C. (2021). Global Network Organization of the Fetal Functional
Connectome. *Cerebral Cortex*, *31*(6), 3034–3046.
https://doi.org/10.1093/cercor/bhaa410
Doria, V., Beckmann, C. F., Arichi, T., Merchant, N., Groppo, M., Turkheimer, F. E., Counsell,
S. J., Murgasova, M., Aljabar, P., Nunes, R. G., Larkman, D. J., Rees, G., & Edwards, A.
D. (2010). Emergence of resting state networks in the preterm human brain.
Proceedings of the National Academy of Sciences, *107*(46), 20015–20020.
https://doi.org/10.1073/pnas.1007921107
Duff, E., Zelaya, F., Almagro, F. A., Miller, K. L., Martin, N., Nichols, T. E., Taschler, B.,
Griffanti, L., Arthofer, C., Douaud, G., Wang, C., Okell, T. W., Bethlehem, R. A. I.,
Eickel, K., Günther, M., Menon, D. K., Williams, G., Facer, B., Lythgoe, D. J., …
Consortium, on behalf of the C.-C. (2022). Reliability of multi-site UK Biobank MRI

brain phenotypes for the assessment of neuropsychiatric complications of SARS-CoV-
2 infection: The COVID-CNS travelling heads study. *PLOS ONE*, *17*(9), e0273704.
https://doi.org/10.1371/journal.pone.0273704
Eyre, M., Fitzgibbon, S. P., Ciarrusta, J., Cordero-Grande, L., Price, A. N., Poppe, T., Schuh, A.,
Hughes, E., O'Keeffe, C., Brandon, J., Cromb, D., Vecchiato, K., Andersson, J., Duff, E.
P., Counsell, S. J., Smith, S. M., Rueckert, D., Hajnal, J. V., Arichi, T., … Edwards, A. D.
(2021). The Developing Human Connectome Project: Typical and disrupted perinatal
functional connectivity. *Brain*, *144*(7), 2199–2213.
https://doi.org/10.1093/brain/awab118
Ferrazzi, G., Kuklisova Murgasova, M., Arichi, T., Malamateniou, C., Fox, M. J., Makropoulos,
A., Allsop, J., Rutherford, M., Malik, S., Aljabar, P., & Hajnal, J. V. (2014). Resting State
fMRI in the moving fetus: A robust framework for motion, bias field and spin history
correction. *NeuroImage*, *101*, 555–568.
https://doi.org/10.1016/j.neuroimage.2014.06.074
Fitzgibbon, S. P., Harrison, S. J., Jenkinson, M., Baxter, L., Robinson, E. C., Bastiani, M., Bozek,
J., Karolis, V., Cordero Grande, L., Price, A. N., Hughes, E., Makropoulos, A., Passerat-
Palmbach, J., Schuh, A., Gao, J., Farahibozorg, S.-R., O'Muircheartaigh, J., Ciarrusta, J.,
O'Keeffe, C., … Andersson, J. (2020). The developing Human Connectome Project
(dHCP) automated resting-state functional processing framework for newborn
infants. *NeuroImage*, *223*, 117303.
https://doi.org/10.1016/j.neuroimage.2020.117303
Friston, K. J., Williams, S., Howard, R., Frackowiak, R. S. J., & Turner, R. (1996). Movement-
Related effects in fMRI time-series. *Magnetic Resonance in Medicine*, *35*(3), 346–355.
https://doi.org/10.1002/mrm.1910350312
Fuderer, M., van den Brink, J., & Jurrissen, M. (2004). SENSE reconstruction using feed
forward regularization. *Proc Int Soc Magn Reson Med*, *11*.
Gao, W., Alcauter, S., Smith, J. K., Gilmore, J. H., & Lin, W. (2015). Development of human
brain cortical network architecture during infancy. *Brain Structure and Function*,
220(2), 1173–1186. https://doi.org/10.1007/s00429-014-0710-3
Ghiglia, D. C., & Romero, L. A. (1994). Robust two-dimensional weighted and unweighted
phase unwrapping that uses fast transforms and iterative methods. *JOSA A*, *11*(1),
107–117. https://doi.org/10.1364/JOSAA.11.000107
Griffanti, L., Salimi-Khorshidi, G., Beckmann, C. F., Auerbach, E. J., Douaud, G., Sexton, C. E.,
Zsoldos, E., Ebmeier, K. P., Filippini, N., Mackay, C. E., Moeller, S., Xu, J., Yacoub, E.,
Baselli, G., Ugurbil, K., Miller, K. L., & Smith, S. M. (2014). ICA-based artefact removal
and accelerated fMRI acquisition for improved resting state network imaging.
NeuroImage, *95*, 232–247. https://doi.org/10.1016/j.neuroimage.2014.03.034
Hurley, R. S., Bonakdarpour, B., Wang, X., & Mesulam, M.-M. (2015). Asymmetric
Connectivity between the Anterior Temporal Lobe and the Language Network.
Journal of Cognitive Neuroscience, *27*(3), 464–473.
https://doi.org/10.1162/jocn_a_00722
Hutter, J., Christiaens, D. J., Schneider, T., Cordero-Grande, L., Slator, P. J., Deprez, M., Price,
A. N., Tournier, J.-D., Rutherford, M., & Hajnal, J. V. (2018). Slice-level diffusion
encoding for motion and distortion correction. *Medical Image Analysis*, *48*, 214–229.
https://doi.org/10.1016/j.media.2018.06.008
Jakab, A., Schwartz, E., Kasprian, G., Gruber, G. M., Prayer, D., Schöpf, V., & Langs, G.
(2014). Fetal functional imaging portrays heterogeneous development of emerging

human brain networks. *Frontiers in Human Neuroscience, 8*.
https://doi.org/10.3389/fnhum.2014.00852

Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved Optimization for the
Robust and Accurate Linear Registration and Motion Correction of Brain Images.
NeuroImage, 17(2), 825–841. https://doi.org/10.1006/nimg.2002.1132

Ji, L., Hendrix, C. L., & Thomason, M. E. (2022). Empirical evaluation of human fetal fMRI
preprocessing steps. *Network Neuroscience, 6*(3), 702–721.
https://doi.org/10.1162/netn_a_00254

Karolis, V., Fitzgibbon, S., Cordero-Grande, L., Farahibozorg, S., Price, A., Hughes, E., Fetit, A.,
Kyriakopoulou, V., Pietsch, M., Rutherford, M., Rueckert, D., Hajnal, J., Edwards, A.,
O'Muircheartaigh, J., Duff, E., & Arichi, T. (2022). *Maturational networks of fetal*
brain activity [Preprint]. Neuroscience. https://doi.org/10.1101/2022.06.14.495883

Karolis, V. R., Fitzgibbon, S. P., Cordero-Grande, L., Farahibozorg, S.-R., Price, A. N., Hughes,
E. J., Fetit, A. E., Kyriakopoulou, V., Pietsch, M., Rutherford, M. A., Rueckert, D.,
Hajnal, J. V., Edwards, A. D., O'Muircheartaigh, J., Duff, E. P., & Arichi, T. (2023).
Maturational networks of human fetal brain activity reveal emerging connectivity
patterns prior to ex-utero exposure. *Communications Biology, 6*(1), 661.
https://doi.org/10.1038/s42003-023-04969-x

Kiviniemi, V., Kantola, J.-H., Jauhiainen, J., Hyvärinen, A., & Tervonen, O. (2003).
Independent component analysis of nondeterministic fMRI signal sources.
NeuroImage, 19(2), 253–260. https://doi.org/10.1016/S1053-8119(03)00097-1

Kochiyama, T., Morita, T., Okada, T., Yonekura, Y., Matsumura, M., & Sadato, N. (2005).
Removing the effects of task-related motion using independent-component analysis.
NeuroImage, 25(3), 802–814. https://doi.org/10.1016/j.neuroimage.2004.12.027

Kostović, I., & Judaš, M. (2010). The development of the subplate and thalamocortical
connections in the human foetal brain: Human foetal cortical circuitry. *Acta*
Paediatrica, 99(8), 1119–1127. https://doi.org/10.1111/j.1651-2227.2010.01811.x

Kostović, I., Sedmak, G., & Judaš, M. (2019). Neural histology and neurogenesis of the human
fetal and infant brain. *NeuroImage, 188*, 743–773.
https://doi.org/10.1016/j.neuroimage.2018.12.043

Kuklisova-Murgasova, M., Lockwood Estrin, G., Nunes, R. G., Malik, S. J., Rutherford, M. A.,
Rueckert, D., & Hajnal, J. V. (2018). Distortion Correction in Fetal EPI Using Non-Rigid
Registration With a Laplacian Constraint. *IEEE Transactions on Medical Imaging,*
37(1), 12–19. https://doi.org/10.1109/TMI.2017.2667227

Laplante, D. P., Brunet, A., Schmitz, N., Ciampi, A., & King, S. (2008). Project Ice Storm:
Prenatal Maternal Stress Affects Cognitive and Linguistic Functioning in 5½-Year-Old
Children. *Journal of the American Academy of Child & Adolescent Psychiatry, 47*(9),
1063–1072. https://doi.org/10.1097/CHI.0b013e31817eec80

Lemieux, L., Salek-Haddadi, A., Lund, T. E., Laufs, H., & Carmichael, D. (2007). Modelling large
motion events in fMRI studies of patients with epilepsy. *Magnetic Resonance*
Imaging, 25(6), 894–901. https://doi.org/10.1016/j.mri.2007.03.009

Minka, T. (2000). Automatic Choice of Dimensionality for PCA. *Advances in Neural*
Information Processing Systems, 13.
https://proceedings.neurips.cc/paper/2000/hash/7503cfacd12053d309b6bed5c89de
212-Abstract.html

Noble, S., Spann, M. N., Tokoglu, F., Shen, X., Constable, R. T., & Scheinost, D. (2017).
Influences on the Test–Retest Reliability of Functional Connectivity MRI and its

Relationship with Behavioral Utility. *Cerebral Cortex*, *27*(11), 5415–5429. https://doi.org/10.1093/cercor/bhx230

O'Donnell, K., O'Connor, T. G., & Glover, V. (2009). Prenatal Stress and Neurodevelopment of the Child: Focus on the HPA Axis and Role of the Placenta. *Developmental Neuroscience*, *31*(4), 285–292. https://doi.org/10.1159/000216539

Oubel, E., Koob, M., Studholme, C., Dietemann, J.-L., & Rousseau, F. (2012). Reconstruction of scattered data in fetal diffusion MRI. *Medical Image Analysis*, *16*(1), 28–37. https://doi.org/10.1016/j.media.2011.04.004

Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2012). Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *NeuroImage*, *59*(3), 2142–2154. https://doi.org/10.1016/j.neuroimage.2011.10.018

Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2014). Methods to detect, characterize, and remove motion artifact in resting state fMRI. *NeuroImage*, *84*, 320–341. https://doi.org/10.1016/j.neuroimage.2013.08.048

Price, A. (2019). The developing Human Connectome Project (dHCP): Fetal acquisition protocol. *Proceedings of the 27th Annual Meeting of the Organisation for Human Brain Mapping, 2479*.

Pruim, R. H. R., Mennes, M., van Rooij, D., Llera, A., Buitelaar, J. K., & Beckmann, C. F. (2015). ICA-AROMA: A robust ICA-based strategy for removing motion artifacts from fMRI data. *NeuroImage*, *112*, 267–277. https://doi.org/10.1016/j.neuroimage.2015.02.064

Rakers, F., Rupprecht, S., Dreiling, M., Bergmeier, C., Witte, O. W., & Schwab, M. (2020). Transfer of maternal psychosocial stress to the fetus. *Neuroscience & Biobehavioral Reviews*, *117*, 185–197. https://doi.org/10.1016/j.neubiorev.2017.02.019

Rutherford, S., Sturmfels, P., Angstadt, M., Hect, J., Wiens, J., van den Heuvel, M. I., Scheinost, D., Sripada, C., & Thomason, M. (2022). Automated Brain Masking of Fetal Functional MRI with Open Data. *Neuroinformatics*, *20*(1), 173–185. https://doi.org/10.1007/s12021-021-09528-5

Salimi-Khorshidi, G., Douaud, G., Beckmann, C. F., Glasser, M. F., Griffanti, L., & Smith, S. M. (2014). Automatic denoising of functional MRI data: Combining independent component analysis and hierarchical fusion of classifiers. *NeuroImage*, *90*, 449–468. https://doi.org/10.1016/j.neuroimage.2013.11.046

Sandman, C. A., Davis, E. P., Buss, C., & Glynn, L. M. (2012). Exposure to Prenatal Psychobiological Stress Exerts Programming Influences on the Mother and Her Fetus. *Neuroendocrinology*, *95*(1), 8–21. https://doi.org/10.1159/000327017

Satterthwaite, T. D., Elliott, M. A., Gerraty, R. T., Ruparel, K., Loughead, J., Calkins, M. E., Eickhoff, S. B., Hakonarson, H., Gur, R. C., Gur, R. E., & Wolf, D. H. (2013). An improved framework for confound regression and filtering for control of motion artifact in the preprocessing of resting-state functional connectivity data. *NeuroImage*, *64*, 240–256. https://doi.org/10.1016/j.neuroimage.2012.08.052

Schöpf, V., Kasprian, G., Brugger, P. C., & Prayer, D. (2012). Watching the fetal brain at 'rest'. *International Journal of Developmental Neuroscience*, *30*(1), 11–17. https://doi.org/10.1016/j.ijdevneu.2011.10.006

Schöpf, V., Schlegl, T., Jakab, A., Kasprian, G., Woitek, R., Prayer, D., & Langs, G. (2014). The relationship between eye movement and vision develops before birth. *Frontiers in Human Neuroscience*, *8*. https://doi.org/10.3389/fnhum.2014.00775

Serag, A., Aljabar, P., Ball, G., Counsell, S. J., Boardman, J. P., Rutherford, M. A., Edwards, A.
D., Hajnal, J. V., & Rueckert, D. (2012). Construction of a consistent high-definition
spatio-temporal atlas of the developing brain using adaptive kernel regression.
NeuroImage, *59*(3), 2255–2265. https://doi.org/10.1016/j.neuroimage.2011.09.062
Seshamani, S., Blazejewska, A. I., Mckown, S., Caucutt, J., Dighe, M., Gatenby, C., &
Studholme, C. (2016). Detecting default mode networks in utero by integrated 4D
fMRI reconstruction and analysis. *Human Brain Mapping*, *37*(11), 4158–4178.
https://doi.org/10.1002/hbm.23303
Shah, L. M., Cramer, J. A., Ferguson, M. A., Birn, R. M., & Anderson, J. S. (2016). Reliability
and reproducibility of individual differences in functional connectivity acquired
during task and resting state. *Brain and Behavior*, *6*(5), e00456.
https://doi.org/10.1002/brb3.456
Shehzad, Z., Kelly, A. M. C., Reiss, P. T., Gee, D. G., Gotimer, K., Uddin, L. Q., Lee, S. H.,
Margulies, D. S., Roy, A. K., Biswal, B. B., Petkova, E., Castellanos, F. X., & Milham, M.
P. (2009). The Resting Brain: Unconstrained yet Reliable. *Cerebral Cortex*, *19*(10),
2209–2229. https://doi.org/10.1093/cercor/bhn256
Skranes, J., & Løhaugen, G. (2022). Prenatal illicit drug and polysubstance exposure and
impact on developmental outcome. In *Handbook of Substance Misuse and
Addictions: From Biology to Public Health.* (pp. 2833–2861). Springer International
Publishing.
Smith, S. M., Beckmann, C. F., Andersson, J., Auerbach, E. J., Bijsterbosch, J., Douaud, G.,
Duff, E., Feinberg, D. A., Griffanti, L., Harms, M. P., Kelly, M., Laumann, T., Miller, K.
L., Moeller, S., Petersen, S., Power, J., Salimi-Khorshidi, G., Snyder, A. Z., Vu, A. T., …
Glasser, M. F. (2013). Resting-state fMRI in the Human Connectome Project.
NeuroImage, *80*, 144–168. https://doi.org/10.1016/j.neuroimage.2013.05.039
Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E. J., Johansen-
Berg, H., Bannister, P. R., De Luca, M., Drobnjak, I., Flitney, D. E., Niazy, R. K.,
Saunders, J., Vickers, J., Zhang, Y., De Stefano, N., Brady, J. M., & Matthews, P. M.
(2004). Advances in functional and structural MR image analysis and implementation
as FSL. *NeuroImage*, *23*, S208–S219.
https://doi.org/10.1016/j.neuroimage.2004.07.051
Sobotka, D., Ebner, M., Schwartz, E., Nenning, K.-H., Taymourtash, A., Vercauteren, T.,
Ourselin, S., Kasprian, G., Prayer, D., Langs, G., & Licandro, R. (2022). Motion
correction and volumetric reconstruction for fetal functional magnetic resonance
imaging data. *NeuroImage*, *255*, 119213.
https://doi.org/10.1016/j.neuroimage.2022.119213
Taymourtash, A., Kebiri, H., Tourbier, S., Schwartz, E., Nenning, K.-H., Licandro, R., Sobotka,
D., Lajous, H., de Dumast, P., Cuadra, M. B., & Langs, G. (2021). *4D iterative
reconstruction of brain fMRI in the moving fetus*.
https://doi.org/10.48550/ARXIV.2111.11394
Taymourtash, A., Schwartz, E., Nenning, K.-H., Sobotka, D., Diogo, M., Kasprian, G., Prayer,
D., & Langs, G. (2019). Quantifying Residual Motion Artifacts in Fetal fMRI Data. In Q.
Wang, A. Gomez, J. Hutter, K. McLeod, V. Zimmer, O. Zettinig, R. Licandro, E.
Robinson, D. Christiaens, E. A. Turk, & A. Melbourne (Eds.), *Smart Ultrasound
Imaging and Perinatal, Preterm and Paediatric Image Analysis* (pp. 171–180).
Springer International Publishing. https://doi.org/10.1007/978-3-030-32875-7_19

Taymourtash, A., Schwartz, E., Nenning, K.-H., Sobotka, D., Licandro, R., Glatter, S., Diogo, M. C., Golland, P., Grant, E., Prayer, D., Kasprian, G., & Langs, G. (2023). Fetal development of functional thalamocortical and cortico–cortical connectivity. *Cerebral Cortex*, *33*(9), 5613–5624. https://doi.org/10.1093/cercor/bhac446

Thomason, M. E., Dassanayake, M. T., Shen, S., Katkuri, Y., Alexis, M., Anderson, A. L., Yeo, L., Mody, S., Hernandez-Andrade, E., Hassan, S. S., Studholme, C., Jeong, J.-W., & Romero, R. (2013). Cross-Hemispheric Functional Connectivity in the Human Fetal Brain. *Science Translational Medicine*, *5*(173). https://doi.org/10.1126/scitranslmed.3004978

Thomason, M. E., Grove, L. E., Lozon, T. A., Vila, A. M., Ye, Y., Nye, M. J., Manning, J. H., Pappas, A., Hernandez-Andrade, E., Yeo, L., Mody, S., Berman, S., Hassan, S. S., & Romero, R. (2015). Age-related increases in long-range connectivity in fetal functional neural connectivity networks in utero. *Developmental Cognitive Neuroscience*, *11*, 96–104. https://doi.org/10.1016/j.dcn.2014.09.001

Tolner, E. A., Sheikh, A., Yukin, A. Y., Kaila, K., & Kanold, P. O. (2012). Subplate Neurons Promote Spindle Bursts and Thalamocortical Patterning in the Neonatal Rat Somatosensory Cortex. *The Journal of Neuroscience*, *32*(2), 692–702. https://doi.org/10.1523/JNEUROSCI.1538-11.2012

Uecker, M., Lai, P., Murphy, M. J., Virtue, P., Elad, M., Pauly, J. M., Vasanawala, S. S., & Lustig, M. (2014). ESPIRiT—an eigenvalue approach to autocalibrating parallel MRI: Where SENSE meets GRAPPA. *Magnetic Resonance in Medicine*, *71*(3), 990–1001. https://doi.org/10.1002/mrm.24751

Uus, A., Kyriakopoulou, V., Cordero Grande, L., Christiaens, D., Pietsch, M., Price, A., Wilson, S., Patkee, P., Karolis, V., Schuh, A., Gartner, A., Williams, L., Hughes, E., Arichi, T., O'Muircheartaigh, J., Hutter, J., Robinson, E., Tournier, J.-D., Rueckert, D., … Edwards, A. D. (n.d.). *Multi-channel spatio-temporal MRI atlas of the normal fetal brain development from the developing Human Connectome Project.* [dataset]. G-Node. https://doi.org/10.12751/g-node.ysgsy1

Uus, A. U., Kyriakopoulou, V., Makropoulos, A., Fukami-Gartner, A., Cromb, D., Davidson, A., Cordero-Grande, L., Price, A. N., Grigorescu, I., Williams, L. Z. J., Robinson, E. C., Lloyd, D., Pushparajah, K., Story, L., Hutter, J., Counsell, S. J., Edwards, A. D., Rutherford, M. A., Hajnal, J. V., & Deprez, M. (2023). BOUNTI: Brain vOlumetry and aUtomated parcellatioN for 3D feTal MRI. *eLife*, *12*. https://doi.org/10.7554/eLife.88818.1

van den Heuvel, M. I., Turk, E., Manning, J. H., Hect, J., Hernandez-Andrade, E., Hassan, S. S., Romero, R., van den Heuvel, M. P., & Thomason, M. E. (2018). Hubs in the human fetal brain network. *Developmental Cognitive Neuroscience*, *30*, 108–115. https://doi.org/10.1016/j.dcn.2018.02.001

Ward Jr., J. H. (1963). Hierarchical Grouping to Optimize an Objective Function. *Journal of the American Statistical Association*, *58*(301), 236–244. https://doi.org/10.1080/01621459.1963.10500845

Wu, Z., Nunes, R. G., Malik, S. J., Estrin, G. L., Hughes, E. J., Malamateniou, C., Counsell, S. J., Rutherford, M. A., & Hajnal, J. V. (n.d.). *Fetal imaging with EPI – FOV, SNR and distortion correction*.

Yan, C.-G., Cheung, B., Kelly, C., Colcombe, S., Craddock, R. C., Di Martino, A., Li, Q., Zuo, X.-N., Castellanos, F. X., & Milham, M. P. (2013). A comprehensive assessment of regional variation in the impact of head micromovements on functional

connectomics. *NeuroImage*, *76*, 183–201.
https://doi.org/10.1016/j.neuroimage.2013.03.004
Zerbo, O., Qian, Y., Yoshida, C., Grether, J. K., Van De Water, J., & Croen, L. A. (2015).
Maternal Infection During Pregnancy and Autism Spectrum Disorders. *Journal of*
Autism and Developmental Disorders, *45*(12), 4015–4025.
https://doi.org/10.1007/s10803-013-2016-3
1189