1

A novel monomeric amyloid β-activated signaling pathway regulates brain development via inhibition of microglia

Hyo Jun Kwon, Devi Santhosh, and Zhen Huang[*]

Departments of Neurology and Neuroscience

University of Wisconsin-Madison

Madison, WI 53706, USA

* Author for correspondence

Tel.: 608-263-2469

Email: zhuang3@wisc.edu

Running title: Monomeric Aβ pathway inhibits microglia in developing brain

ABSTRACT

Amyloid β (Aβ) forms aggregates in the Alzheimer's disease brain and is well known for its pathological

roles. Recent studies show that it also regulates neuronal physiology in the healthy brain. Whether Aβ

also regulates glial physiology in the normal brain, however, has remained unclear. In this article, we

describe the discovery of a novel signaling pathway activated by the monomeric form of Aβ that plays

essential roles in the regulation of microglial activity and the assembly of neocortex during

development. We find that activation of this pathway depends on the function of amyloid precursor

(APP) and heterotrimeric G proteins in microglia and inhibits microglial immune activation at

transcriptional and post-transcriptional levels. Genetic disruption of this pathway during neocortical

development results in microglial dysregulation and excessive matrix proteinase activation, leading to

basement membrane degradation, neuronal ectopia, and laminar disruption. These results uncover a

previously unknown function of Aβ as a negative regulator of brain microglia and substantially elucidate

the molecular mechanisms underlying this regulation. Considering the prominence of Aβ and

neuroinflammation in the pathology of Alzheimer's disease, they also highlight a potentially overlooked

role of Aβ monomer depletion in the development of the disease.

INTRODUCTION

Aβ, a core component of amyloid plaques in the Alzheimer's disease brain, is well known to form

oligomers under disease conditions. Studies have shown that the oligomers formed by Aβ are highly

toxic, with wide-ranging effects including inhibition of neurotransmitter release, depletion of synaptic

vesicle pools, disruption of postsynaptic organization and function, and impairment of multiple forms of

synaptic plasticity (Gulisano et al., 2018; He et al., 2019; Kim et al., 2013; Lauren et al., 2009; Lazarevic et

al., 2017; Parodi et al., 2010; Puzzo et al., 2008; Shankar et al., 2008; Walsh et al., 2002; Yang et al.,

2015; Zott et al., 2019). These effects likely significantly underpin the pathogenic role of Aβ in

Alzheimer's disease and contribute to neuron loss and cognitive decline in patients. Besides its

pathological roles, recent studies show that Aβ is also produced in the healthy brain by neurons in a

neural activity-dependent manner and regulates the normal physiology of neurons (Cirrito et al., 2005;

Fogel et al., 2014; Galanis et al., 2021; Garcia-Osta and Alberini, 2009; Gulisano et al., 2018; Gulisano et

al., 2019; Morley et al., 2010; Palmeri et al., 2017; Puzzo et al., 2008; Zhou et al., 2022). For example,

consistent with studies showing that Aβ monomers and low molecular weight oligomers positively

regulate synaptic function and plasticity, administration of these molecules in vivo has been found to

improve learning and memory in animals (Fogel et al., 2014; Garcia-Osta and Alberini, 2009; Gulisano et

al., 2018; Gulisano et al., 2019; Morley et al., 2010; Palmeri et al., 2017; Puzzo et al., 2008). Recent

studies have further shown that Aβ monomers directly promote synapse formation and function and

homeostatic plasticity, processes crucial to normal cognitive function (Galanis et al., 2021; Kamenetz et

al., 2003; Zhou et al., 2022). Together, these findings have provided crucial insights into the

physiological roles that Aβ plays in regulating normal neuronal function in the brain. However, it

remains unclear if Aβ also regulates the physiology of glia, nonneuronal cells that also play important

roles in normal brain function.

Microglia and astrocytes, two of the major glial cell types in the brain, are known to play critical

roles in the normal development, function, and plasticity of the brain circuitry (Barres, 2008; Schafer and

Stevens, 2015). They coordinately regulate, among others, the spatiotemporally specific expression of

immune cytokines in the brain that regulate numerous processes in brain circuit development, function,

and plasticity (Zipp et al., 2023). For example, in the thalamus, a key relay station in the visual pathway,

populations of astrocytes have been found to activate the expression of interleukin-33 in a neural

activity-dependent manner, induce activity-dependent elimination of supernumerary synapses, and

promote the maturation of the visual circuitry in early postnatal life (Vainchtein et al., 2018). In the

adult hippocampus, in contrast, astrocytes have been found to activate the expression of interleukin-33

under neuronal activity blockade and induce homeostatic synaptic plasticity that maintains circuit

activity balance (Wang et al., 2021). In the striatum and the neocortex, not only have astrocytes but

also have microglia been observed to activate the expression of TNFα upon changes in neural circuit

activity and induce homeostatic synaptic plasticity that dampens circuit perturbation (Heir et al., 2024;

Lewitus et al., 2016; Stellwagen and Malenka, 2006). In the clinic, the induction of microglial release of

cytokines such as TNFα also underpins the application of repetitive transcranial magnetic stimulation, a

noninvasive brain stimulation technique frequently used to induce cortical plasticity and treat

pharmaco-resistant depression (Eichler et al., 2023). Furthermore, in neurodegenerative diseases such

as Alzheimer's disease, glial activation and brain cytokine elevation are key pathologic factors in disease

development (Colonna and Butovsky, 2017; Patani et al., 2023). Elevated TNFα expression by microglia

also underlies interneuron deficits and autism-like phenotype linked to maternal immune activation (Yu

et al., 2022). Thus, the precise regulation of glial cytokine expression in the brain plays a key role in the

normal development and function of the brain and its dysregulation is linked to common

neurodevelopmental and neurodegenerative diseases. However, how glial cytokine expression is

mechanistically regulated by cell-cell communication in the brain have remained largely unknown.

In this article, we report the discovery of a novel microglial signaling pathway activated by $A\beta$,

the neuron-produced peptide at the center of Alzheimer's disease, that plays a crucial role in precisely

regulating the levels of microglial cytokine expression and activity and ensuring the proper assembly of

neuronal laminae during cerebral cortex development. We first came across this pathway in our study

of the function of *ric8*a, a gene encoding a molecular chaperone for heterotrimeric G proteins (Gabay et

al., 2011; Ma et al., 2012; Ma et al., 2017; Tall et al., 2003). We found that deletion of *ric8a* during

cortical development resulted in cortical basement membrane degradation, neuronal ectopia, and

laminar disruption. However, unlike in classic models of cobblestone lissencephaly, these phenotypes

resulted not from *ric8a* deficiency in brain neural cell types, but from deficiency in microglia. The

phenotypes also resemble those in mutants of amyloid precursor protein (APP) family and pathway

genes. Indeed, we found that *app* deficiency in microglia also underpins ectopia formation in *app* family

gene mutants. Furthermore, we found that APP and Ric8a form a functional pathway in microglia that is

specifically activated by the monomeric form of $A\beta$ and this pathway normally inhibits the

transcriptional and post-transcriptional expression of immune cytokines by microglia.

RESULTS

Cortical ectopia in *ric8a-emx1-cre* mutants results from non-neural deficiency

To study of the function of *ric8a* in neocortical development, we deleted a conditional *ric8a* allele (Ma et

al., 2012; Ma et al., 2017) using *emx1-cre,* a *cre* line designed to target dorsal forebrain neural

progenitors in mice (Gorski et al., 2002). We found it result in ectopia formation exclusively in the lateral

cortex of the perinatal mutant brain (**Fig. 1a-d**). Birth-dating showed that the ectopia consisted of both

early and late-born neurons (**Supplemental Fig. 1**). Consistent with this observation, neurons in the

ectopia also stained positive for both Ctip2 and Cux1, genes specific to lower and upper-layer neurons,

respectively. Interestingly, in cortical areas without ectopia, radial migration of early- and late-born

neurons appeared largely normal as shown by birth-dating as well as Cux1 and Ctip2 staining

(**Supplemental Fig. 2**). This suggests that cell-autonomous defects in neurons are unlikely the cause of

the ectopia. At E16.5, clear breaches in the pial basement membrane of the developing cortex were

already apparent (**Supplemental Fig. 3**). However, unlike classic models of cobblestone lissencephaly,

where radial glial fibers typically retract, radial glial fibers in *ric8a* mutants instead extended beyond the

breaches. This argues against radial glial cell adhesion defects since they would be predicted to retract.

Furthermore, in areas without ectopia, we also observed normal localization of Cajal-Retzius cells,

expression of Reelin, and splitting of the preplate, arguing against primary defects in Cajal-Retzius cells.

In cobblestone lissencephaly, studies show that ectopia result from primary defects in radial glial

maintenance of the pial basement membrane (Beggs et al., 2003; Graus-Porta et al., 2001; Moore et al.,

2002; Satz et al., 2010). In *ric8a* mutants, we observed large numbers of basement membrane breaches

at E14.5, almost all associated with ectopia (**Supplemental Fig. 4**). In contrast, at E13.5, although we

also observed significant numbers of breaches, none was associated with ectopia. This indicates that

basement membrane breaches similarly precede ectopia in *ric8a* mutants. However, at E12.5, despite a

complete lack of basement membrane breaches, we observed increased numbers of laminin-positive

debris across the lateral cortex, both beneath basement membrane segments with intact laminin

staining and beneath segments with disrupted laminin staining, the latter presumably sites of future

breach (**Supplemental Fig. 5**). As a major basement membrane component, the increased amounts of

laminin debris suggest increased degradative activity within the developing cortex. Thus, these results

indicate that excessive basement membrane degradation, but not defective maintenance, is likely a

primary cause of cortical ectopia in *ric8a* mutants.

To determine the cell type(s) genetically responsible for cortical basement membrane

degradation and ectopia in *ric8a* mutants, we employed a panel of *cre* lines (**Fig. 1e-h'**). To target Cajal-

Retzius cells, we employed *wnt3a-cre* (Yoshida et al., 2006) but found *ric8a* deletion using *wnt3a-cre* did

not result in ectopia. To target postmitotic excitatory and inhibitory neurons, we employed *nex-cre*

(Goebbels et al., 2006) and *dlx5/6-cre* (Stenman et al., 2003) respectively but similarly found neither

result in ectopia. These results point to *ric8a* requirement in cell types other than post-mitotic neurons.

To test the involvement of neural progenitors, we employed *nestin-cre* (Graus-Porta et al., 2001).

Previous studies show that deletion of $\beta1$ *integrin* and related genes by *emx1-cre* and *nestin-cre* results

in similar ectopia phenotypes (Belvindrah et al., 2006; Graus-Porta et al., 2001; Huang et al., 2006;

Niewmierzycka et al., 2005). To our surprise, deletion of *ric8a* by *nestin-cre* did not result in ectopia (Ma

et al., 2017) (**Fig. 1g-g'**). Since *nestin-cre*-mediated deletion in neural progenitors is inherited by post-

mitotic neurons and astrocytes, this indicates that the combined deletion of *ric8a* from all these cell

types does not lead to ectopia. The onset of *nestin-cre* expression is, however, developmentally slightly

later than that of *emx1-cre* (Gorski et al., 2002). To assess the potential contribution of this temporal

difference, we employed *foxg1-cre*, a *cre* line expressed in forebrain neural progenitors starting from

E10.5 (Hebert and McConnell, 2000). We found that *ric8a* deletion using *foxg1-cre* still failed to produce

ectopia (**Fig. 1h-h'**). Thus, these results strongly argue against the interpretation that *ric8a* deficiency in

neural cell lineages is responsible for basement membrane degradation and ectopia in *ric8a* mutants.

During embryogenesis, the neural tube undergoes epithelial-mesenchymal transition giving rise

to neural crest cells (Leathers and Rogers, 2022). This process involves region-specific basement

membrane breakdown that resembles the *ric8a* mutant phenotype. To determine if ectopic epithelial-

mesenchymal transition plays a role, we examined potential changes in neuro-epithelial cell fates in the

mutant cortex. We found that cortical neural progenitors expressed Pax6, Nestin, and Vimentin

normally (**Supplemental Fig. 6**). Cell proliferation in the ventricular zone was also normal. Furthermore,

although *ric8a* regulates asymmetric cell division in invertebrates (Afshar et al., 2004; Couwenbergs et

al., 2004; David et al., 2005; Hampoelz et al., 2005; Wang et al., 2005)*,* we observed no significant

defects in mitotic spindle orientation at the ventricular surface. Additionally, no ectopic expression of

neural crest markers or Wnt pathway activation was observed (**Supplemental Fig. 7**). Altogether, these

results further indicate that non-neural-cell deficiency is responsible for ectopia formation in *ric8a*

mutants.

Microglial *ric8a* deficiency is responsible for ectopia formation

To assess the role of non-neural cell types, we turned our attention to microglia since RNA-seq studies

show that brain microglia express *emx1* at a significant level (Zhang et al., 2014). To determine if *emx1-*

cre is expressed and active in microglia, we isolated microglia from *ric8a-emx1-cre* mutants. We found

that *emx1-cre* mediated *ric8a* deletion indeed resulted in altered cytokine expression in microglia

(**Supplemental Fig. 8**). This indicate that *emx1-cre* is expressed and active in microglia and deletes *ric8a*.

To determine the specific significance of *ric8a* deletion from microglia alone, we next employed a

microglia-specific *cx3cr1-cre* (Yona et al., 2013). Like *emx1-cre* mutants, *ric8a-cx3cr1-cre* mutant

microglia also showed elevated cytokine secretion and transcription in comparison to control microglia

upon stimulation by lipopolysaccharide (LPS) (**Fig. 2a-b**). Similar results were also obtained with

stimulation by polyinosinic-polycytidylic acid (poly I:C), an intracellular immune activator. Thus, these

results indicate that *ric8a* deficiency in microglia results in broad increases in microglial sensitivity to

immune stimulation.

To determine if microglial *ric8a* deficiency alone is sufficient to cause cortical ectopia in vivo, we

examined *ric8a-cx3cr1-cre* mutants but found that it did not affect either basement membrane integrity

or cortical layering (**Fig. 2c**). We reasoned that this may be related to the fact that *ric8a* mutant

microglia only show heightened activity upon stimulation but not under basal unstimulated conditions

(**Fig. 2a-b**) but elevated microglial activity may be needed for basement membrane degradation and

ectopia formation. To test this possibility, we employed in utero LPS administration to activate microglia

during cortical development. We found that over 50% of *ric8a-cx3cr1-cre* mutant neonates showed

ectopia when administered LPS at E11.5-12.5 (10 of 19 mutant neonates examined) (**Fig. 2d**). In

contrast, no cortical ectopia were observed in any of the 32 littermate controls that were similarly

administered LPS. This indicates that only the combination of microglial *ric8a* deficiency and immune

activation leads to ectopia formation. In *emx1-cre* mutants, ectopia develop without LPS administration

(**Fig. 1**). We suspect that this may be due to concurrent *ric8a* deficiency in neural cell types, which may

result in deficits that mimic immune stimulation. To test this, we next additionally deleted *ric8a* from

neural cells in the *ric8a-cx3cr1-cre* microglial mutant background by introducing *nestin-cre*. We have

shown that *ric8a* deletion by *nestin-cre* alone does not result in ectopia (**Fig. 1g-g'**). However, we found

that, like deletion by *emx1-cre*, *ric8a* deletion by the dual *cre* combination of *cx3cr1-cre* and *nestin-cre*

also resulted in ectopia in all double *cre* mutants (6 of 6 examined) (**Fig. 2f**). Thus, these results indicate

that elevated immune activation of *ric8a* deficient microglia during cortical development is responsible

for ectopia formation.

Microglial *app* deficiency also results in ectopia formation

The exclusive localization of ectopia to the lateral cortex in *ric8a* mutants is reminiscent of the

phenotypes observed in APP family and pathway mutants including *app/aplp1/2* triple (Herms et al.,

2004) and *apbb1/2* double knockouts (Guenette et al., 2006). Independent studies also point to a role

of non-neural cells in ectopia formation in these mutants. For example, unlike the triple knockout,

specific *app* knockdown in cortical neurons during development results in under- but not over-migration

of neurons (Young-Pearse et al., 2007). To test the role of microglia in ecotopia formation in APP

pathway mutants, we first analyzed *app* mutant microglia. To this end, we employed *cx3cr1-cre* to

delete a conditional allele of *app* from microglia and found that microglia cultured from *app-cx3cr1-cre*

mutants showed reduced TNFα and IL-6 secretion as well as muted IL-6 transcription upon stimulation

(**Fig. 3a, Supplemental Fig. 9**). This indicates that *app* plays a previously unrecognized, cell-autonomous

role in microglia in regulating microglial activity. Microglia exhibit attenuated immune response

following chronic stimulation, especially when carrying strong loss-of-function mutations in anti-

inflammatory pathways (Chamberlain et al., 2015; Sayed et al., 2018). We suspect that the attenuated

response by *app* mutant microglia may result from similar effects following in vitro culture. To test

effects of *app* mutation under conditions that more closely resemble in vivo conditions, we next isolated

fresh, unelicited peritoneal macrophages and acutely analyzed their response to immune stimulation.

We found that *app* mutant macrophages showed significantly elevated secretion of all cytokines tested

(**Fig. 3b**). At the transcriptional level, mRNA induction was also increased for all cytokines (**Fig. 3c**).

Thus, like that of *ric8a*, the normal function of *app* also appears to be to suppress the inflammatory

activation of microglia.

To determine if microglial *app* deficiency is also responsible for ectopia formation in *app* triple

knockout mutants, we next asked if activating microglia in microglia-specific *app* mutants similarly

results in pial ectopia during cortical development. To this end, we administered LPS in utero at E11.5-

12.5 to *app-cx3cr1-cre* mutant animals as we did to *ric8a-cx3cr1-cre* mutants above. We found that,

while none of the 81 littermate controls administered LPS showed ectopia, a significant number of

mutant neonates showed ectopia (6 of 31 neonates examined, ~19%) (**Fig. 3e**). Thus, *app* deficient

microglia, when activated, also results in cortical ectopia during development. The reduced severity of

the ectopia observed, as compared to that in *ric8a/cx3cr1-cre* mutants, likely in part results from the

reduced LPS dosage (by ~3 folds) we had to use in these animas due to the enhanced immune sensitivity

of their strain genetic background. Other *app* gene family members are also expressed in microglia

(Zhang et al., 2014) and may in addition compensate for the loss of APP. Thus, these results indicate

that *app* normally plays a cell-autonomous role in microglia that negatively regulate microglial activation

and its loss of function underlies ectopia formation. The similarities between *app* and *ric8a* mutant

phenotypes suggest that they form a previously unknown anti-inflammatory pathway in microglia.

Monomeric Aβ suppresses microglial inflammatory activation via APP and Ric8a

The possibility that *app* and *ric8a* may form a novel anti-inflammatory pathway in microglia raises

questions on the identity of the ligands for the pathway. Several molecules have been reported to bind

to APP and/or activate APP-dependent pathways (Fogel et al., 2014; Milosch et al., 2014; Rice et al.,

2012), among which Aβ is note-worthy for its nanomolar direct binding affinity (Fogel et al., 2014;

Shaked et al., 2006). Aβ oligomers and fibrils have been shown by numerous studies to be pro-

inflammatory, while non-fibrillar Aβ lack such activity (Halle et al., 2008; Huang, 2023, 2024; Lorton et

al., 1996; Muehlhauser et al., 2001; Tan et al., 1999). In contrast, when employed under conditions that

favor the monomer conformation, Aβ inhibits T cell activation (Grant et al., 2012). This suggests that,

unlike Aβ oligomers, Aβ monomers may be anti-inflammatory. To test this possibility, we dissolved

Aβ40 peptides in DMSO, which has been shown to preserve the monomeric conformation (LeVine,

2004; Stine et al., 2011). We found that Aβ monomers as prepared potently suppressed the secretion of

large numbers of cytokines (**Fig. 4a**, **Supplemental Fig. 10**) and showed similar effects on microglia no

matter if they were activated by LPS or poly I:C (**Fig. 4b**). We also found that the Aβ monomers similarly

strongly inhibited the induction of cytokines at the transcriptional level (**Fig. 4c, Supplemental Fig. 10**).

In addition, we observed these effects with Aβ40 peptides from different commercial sources. Thus,

these results indicate that monomeric Aβ possesses a previously unreported anti-inflammatory activity

against microglia that strongly inhibits microglial inflammatory activation.

To determine whether monomeric Aβ signals through APP, we employed *app-cx3cr1-cre* mutant

microglia. We found that, unlike that of control microglia, Aβ monomers failed to suppress the

secretion of all tested cytokines by *app* mutant microglia (**Fig. 4d, Supplemental Fig. 10**). Interestingly,

this blockade appeared to be specific to *app* since Aβ monomers still significantly suppressed cytokine

secretion by *aplp2* mutant microglia. At the transcriptional level, Aβ monomers also failed to suppress

cytokine induction in *app* mutant microglia (**Fig. 4e, Supplemental Fig. 10**). Together, these results

indicate that APP is functionally required in microglia for Aβ monomer inhibition of cytokine expression

at both transcriptional and post-transcriptional levels. Cultured microglia from *app-cx3cr1*-cre mutants

showed attenuated immune activation (**Fig. 3**). To assess whether this may affect the efficacy of Aβ

monomer inhibition, we next tested the response of fresh, unelicited macrophages. We found that, like

that of control microglia, cytokine secretion by control macrophages was also strongly suppressed by Aβ

monomers (**Fig. 4f, Supplemental Fig. 10**). However, even though *app* mutant macrophages showed

elevated response to immune stimulation in comparison to control macrophages, they still failed to

respond to Aβ monomers and displayed levels of cytokine secretion that were indistinguishable from

those of DMSO-treated cells (**Fig. 4f, Supplemental Fig. 10**). Thus, these results further indicate that

APP function is required in microglia for mediating the anti-inflammatory effects of Aβ monomers.

The similarity of *ric8a* ectopia to *app* ectopia phenotype (**Figs. 2** & **3**) also suggests that Ric8a

functions in the same pathway as APP in mediating Aβ monomer anti-inflammatory signaling in

microglia. This is consistent with previous studies showing that heterotrimeric G proteins are coupled to

APP and mediate APP intracellular signaling in vitro and vivo (Fogel et al., 2014; Milosch et al., 2014;

Nishimoto et al., 1993; Ramaker et al., 2013) and that Ric8a is a molecular chaperone essential for the

post-translational stability of heterotrimeric G proteins (Gabay et al., 2011; Tall et al., 2003). To directly

test if Ric8a is part of this pathway, we next employed *ric8a-cx3cr1-cre* mutant microglia. We found

that, indeed, like that of *app* mutant microglia, Aβ monomers also failed to suppress the secretion of

TNFα and IL-6 by *ric8a* mutant microglia (**Fig. 4g**). This indicates that heterotrimeric G proteins function

is likely required in the same pathway of APP in microglia for the suppression of TNFα and IL-6 secretion.

However, unlike APP, we found that Ric8a appears to be dispensable for Aβ monomer regulation of

other cytokines. For example, unlike that of TNFα and IL-6, Aβ monomers still suppressed IL-1β

secretion by *ric8a* mutant microglia (**Supplemental Fig. 10**). It also appears to be dispensable for the

regulation of cytokine transcription since Aβ monomers similarly suppressed IL-6 transcriptional

induction in both control and *ric8a* mutant microglia. These results suggest that heterotrimeric G

proteins function may only mediate some of the anti-inflammatory signaling of monomeric Aβ. Thus,

APP and Ric8a-regulated heterotrimeric G proteins form part of a novel anti-inflammatory pathway

activated by monomeric Aβ in microglia.

Elevated matrix metalloproteinases (MMPs) cause basement membrane degradation

We have shown that heightened microglial activation due to mutation in the Aβ monomer-activated

APP/Ric8a pathway results in basement membrane degradation and ectopia during cortical

development. To further test this interpretation, we sought to test the prediction that inhibition of

microglial activation in these mutants suppressed the formation ectopia. To this end, we employed

dorsomorphin and S3I-201, inhibitors targeting Akt, Stat3, and other mediators in pro-inflammatory

signaling (Lee et al., 2016; Qin et al., 2012). Consistent with their anti-inflammatory activity, we found

that dorsomorphin and S3I-201 both suppressed astrogliosis associated with neuroinflammation in the

cortex of *ric8a-emx1-cre* mutants (**Supplemental Fig. 11**). Furthermore, they also suppressed the

formation of ectopia in *ric8a-emx1-cre* mutants, reducing both the number and the size of the ectopia

observed (**Fig. 5a-f**). Most strikingly, the combined administration of dorsomorphin and S3I-201 nearly

eliminated all ectopia in *ric8a-emx1-cre* mutants (**Fig. 5d, 5e**). Thus, these results indicate that excessive

inflammatory activation of microglia is responsible for ectopia formation in *ric8a* mutants.

Under neuroinflammatory conditions, brain cytokines frequently induce MMPs, which lead to

breakdown of the extracellular matrix and contribute to disease pathology (Pagenstecher et al., 1998;

Wang et al., 2000). Since *ric8a* mutant microglia are hyperactive in inflammatory cytokine production,

we wonder if induction of MMPs may underlie the laminin degradation and cortical basement

membrane break observed in *ric8a* mutants. To test this, we examined the activities of MMP2 and

MMP9 in the developing cortex using gelatin gel zymography. We found that the activity of MMP9 in

the mutant cortex was significantly increased (**Fig. 5i, Supplemental Fig. 12**). In contrast, that activity of

MMP2 remained unaffected. Similarly, at the protein level, we found that the immunoreactivity for

MMP9 was increased in *ric8a-emx1-cre* mutants (**Fig. 5g-h**). To test if the increased MMP activity is

responsible for the ectopia observed, we next employed BB94, a broad-spectrum inhibitor of MMPs.

We found that BB94 administration significantly suppress both the number and the size of the ectopia in

ric8a mutants (**Fig. 5j-m**). To narrow down the identity of MMPs responsible, we further employed an

inhibitor specific for MMP9 and 13 and found that it similarly suppressed both the number and the size

of the ectopia (**Fig. 5l-m**). Furthermore, consistent with its near complete suppression of cortical

ectopia (**Fig. 5a-f**), we found that the co-administration of dorsomorphin and S3I-201 also reduced

MMP9 activity in the mutant cortex to the control level (**Supplemental Fig. 12**). Thus, these results

indicate this Aβ monomer-activated anti-inflammatory pathway normally promotes cortical

development through suppressing microglial activation and MMP induction.

DISCUSSION

The spatiotemporal expression of immune cytokines by glial cells in the brain plays critical roles in the normal development, function, and plasticity of the brain circuitry (Barres, 2008; Schafer and Stevens, 2015; Zipp et al., 2023). In this article, we have identified a novel microglial anti-inflammatory pathway activated by monomeric Aβ that inhibits microglial cytokine expression and plays essential roles in the normal development of the cerebral cortex. We have found that this pathway is mediated by APP and heterotrimeric G proteins in microglia and its activation leads to the inhibition of microglial cytokine induction at transcriptional and post-transcriptional levels (**Figs. 1-4**). We further show that a key function of this pathway is to suppress the activity of MMP9 during corticogenesis and disruption of this regulation results in cortical basement membrane degradation and neuronal ectopia development (**Figs. 1-3, 5**). Furthermore, we find that this pathway is activated specifically by the monomeric form of Aβ (**Fig. 4**), identifying, for the first time, an isoform specific activity of Aβ against microglia. These results provide novel insights into the neuron-glia communication mechanisms that coordinate the regulation of immune cytokines, key regulators of Hebbian and non-Hebbian synaptic plasticity, by glial cells in the brain. The discovery of the novel activity of monomeric Aβ as a negative microglial regulator may also facilitate the further elucidation of Alzheimer's disease pathogenesis.

Microglial activity regulation during cortical development

Among the glial cell populations in the brain, astrocytes and oligodendrocyte are both born within the nervous system at the end of cortical neurogenesis. As such, they play limited roles in the early steps of cortical development. In contrast, microglia are not only of a distinct non-neural lineage that originates from outside the nervous system but also begin to populate the brain at the onset of corticogenesis (Ginhoux et al., 2010; Hattori et al., 2023). As such, they play unique roles throughout cortical

development. Indeed, microglial activity has been found to regulate the size of the cortical neural

precursor pool (Cunningham et al., 2013). Microglia-secreted cytokines have also been found to

promote both neurogenesis and oligodendrogenesis (Shigemoto-Mogami et al., 2014). As such, the

precise regulation of microglial activity is critical to the normal development of the neocortex from an

early stage. In this study, we have shown that immune over-activation of microglia due to disruption of

a monomeric Aβ-activated pathway results in excessive cortical matrix proteinase activation, leading

basement membrane degradation and neuronal ectopia. Previous studies have shown that reductions

in the expression of microglial immune and chemotaxis genes instead lead to the failure of microglia to

populate the brain (Iyer et al., 2022). These results together thus highlight the importance of precisely

regulating the level of microglial activity during brain development. The dramatic destructive effects of

microglial hyperactivity we have observed during corticogenesis also foreshadow the critical roles it

plays in brain dysfunction and disease at later stages of life.

In this study, we have also shown that the anti-inflammatory regulation of microglia in

corticogenesis depends on a pathway composed of APP and Ric8a-regulated heterotrimeric G proteins.

This has revealed new insight into the intercellular signaling mechanisms regulating microglial activity in

the brain. Heterotrimeric G proteins are well-known mediators of G protein-coupled receptor (GPCR)

signaling. In this study, we have found that they also function in the same pathway as APP. To our

knowledge, ours is the first study to report an in vivo anti-inflammatory function of this pathway in

microglia and has significantly advanced knowledge in microglial biology. This is also consistent with

previous studies showing that heterotrimeric G proteins interact with the APP cytoplasmic domain and

mediate APP signaling from invertebrates to mammals in several other cell types (Fogel et al., 2014;

Milosch et al., 2014; Nishimoto et al., 1993; Ramaker et al., 2013). In this study, we have in addition

shown that this pathway is specifically activated by the monomeric form of Aβ, a peptide produced by

neurons in the brain (Cirrito et al., 2005), providing further insight into the biological function of this

pathway. In the early cortex, neurogenesis is just beginning, and most neurons born are in an immature

state. It is unclear if this pathway is activated by Aβ at this stage. However, studies have shown that

other APP ligands such as pancortin, a member of the olfactomedin family proteins known to inhibit

innate immunity (Liu et al., 2010), are expressed in the cortex at this stage (Rice et al., 2012). It will be

interesting to determine if these innate immune regulators play a role in regulating this pathway.

Neuronal activity, glial cytokine expression, and brain circuit plasticity

Activity-dependent competitive and homeostatic plasticity is a foundational rule that regulates the

development, maturation, and function of neural circuits across brain regions. Studies have shown that

glial cells, through regulating the spatiotemporal expression of immune cytokines, play a pivotal role in

this process. In the developing thalamus, by activating interleukin-33 expression in an activity-

dependent manner, astrocytes have been found to promote the segregation of eye-specific axonal

projection and the maturation of the visual circuitry (He et al., 2022; Vainchtein et al., 2018). In the

visual cortex, astrocytic expression of TNFα similarly mediates activity-dependent homeostatic upscaling

of cortical synapses following peripheral monocular deprivation (Barnes et al., 2017; Heir et al., 2024;

Kaneko et al., 2008). In this study, we have shown that Aβ monomers inhibit expression of cytokines by

brain microglia via a novel APP/heterotrimeric G protein-mediated pathway. Aβ is primarily produced

by neurons in the brain in a neural activity-dependent manner and form oligomers when large quantities

are produced (Cirrito et al., 2005). Aβ oligomers, in contrary to monomers, are proinflammatory and

increase glial cytokine expression (Halle et al., 2008; Huang, 2023; Lorton et al., 1996; Muehlhauser et

al., 2001; Tan et al., 1999). These findings thus suggest that different levels of neural circuit activity in

the brain may differentially regulate glial cytokine expression through inducing different levels of Aβ.

High levels of neural activity may lead to high levels of Aβ and the formation of Aβ oligomers that

activate glial cytokine production, while low levels of neural activity may produce low levels of Aβ,

maintain Aβ as monomers, and inhibit glial cytokine production. Thus, Aβ in the brain may not only be a

reporter of the levels of neural circuit activity but may also serve as an agent that directly mediate

activity level-dependent plasticity. Following sensory deprivation, for example, Aβ levels may be

lowered due to loss of sensory stimulation. This may lead to the relief of monomeric Aβ inhibition of

cytokines such as TNFα and as a result trigger homeostatic upscaling of cortical synapses in the visual

cortex (Barnes et al., 2017; Heir et al., 2024; Kaneko et al., 2008). In contrary, when neural activity levels

are high, large quantities of Aβ may be produced, leading to formation of Aβ oligomers that may in turn

induce expression of cytokines such as IL-33 that promote synaptic pruning. A large body of evidence

strongly indicates that Aβ and related pathways indeed mediate homeostatic and competitive plasticity

in the visual and other systems of the brain (Galanis et al., 2021; Huang, 2023, 2024; Kamenetz et al.,

2003; Kim et al., 2013). Our discovery of the Aβ monomer-activated pathway has therefore provided

novel insights into a universal mechanism that senses neural circuit activity pattern and translates it into

homeostatic and competitive synaptic changes in the brain, a mechanism with fundamental roles in

cognitive function.

In this study, we have also found that the matrix proteinase MMP9 is a key downstream effector

of microglial activity in the developing cortex. We find that microglial hyperactivity results in increased

levels of MMP9, leading to cortical basement membrane degradation and neuronal ectopia and

inhibiting MMP9 directly or indirectly suppresses the phenotype. This suggests that the regulation of

MMP9 may be a key mechanism by which glial cells regulate brain development and plasticity. Indeed,

independent studies have shown that, in the visual cortex, MMP9 is also a pivotal mediator of TNFα-

dependent homeostatic upscaling of central synapses following monocular deprivation (Akol et al.,

2022; Kaneko et al., 2008; Kelly et al., 2015; Spolidoro et al., 2012). In the Xenopus tectum, MMP9 has

similarly been found to be induced by neural activity and promote visual activity-induced dendritic

growth (Gore et al., 2021). Importantly, in both wildtype and amblyopic animals, light reintroduction

after dark exposure has been found to reactivate plasticity in the adult visual cortex via MMP9,

uncovering a potential treatment for common visual conditions (Murase et al., 2017; Murase et al.,

2019). These results therefore highlight a conserved glia/cytokine/MMP9-mediated mechanism that

regulates brain development and plasticity from embryogenesis to adulthood. In ocular dominance

plasticity, MMP9 is activated at perisynaptic regions (Murase et al., 2017; Murase et al., 2019). MMP9

mRNA translation has been also observed in dendrites (Dziembowska et al., 2012). In the *ric8a* mutant

cortex, we find that MMP9 activity is increased. Further studies are required to determine the cellular

sources of MMP9 and how its activity is regulated.

Aβ monomer anti-inflammatory activity and Alzheimer's disease

Aβ is well known as a component of the amyloid plaques in the Alzheimer's disease brain. It is a unique

amphipathic peptide that can, dependent on concentration and other conditions, remain as monomers

or form oligomers. Studies on Aβ have historically focused on the neurotoxic effects of Aβ oligomers

and their proinflammatory effects on glia (Gulisano et al., 2018; Halle et al., 2008; He et al., 2019;

Huang, 2023; Kim et al., 2013; Lauren et al., 2009; Lazarevic et al., 2017; Lorton et al., 1996;

Muehlhauser et al., 2001; Parodi et al., 2010; Puzzo et al., 2008; Shankar et al., 2008; Tan et al., 1999;

Walsh et al., 2002; Yang et al., 2015; Zott et al., 2019). In this study, we have found that, in contrary to

Aβ oligomers, Aβ monomers instead possess a previously unknown anti-inflammatory activity that acts

through a unique microglial pathway. We have further found that genetic disruption of this pathway in

corticogenesis results microglial hyperactivity, leading to neuronal ectopia and large disruption of

cortical structural organization. To our knowledge, ours is the first study to uncover this overlooked

anti-inflammatory activity of Aβ monomers. It is in alignment with recent studies showing that Aβ

monomers are also directly protective to neurons and positively regulate synapse development and

function (Galanis et al., 2021; Giuffrida et al., 2009; Plant et al., 2003; Ramsden et al., 2002; Zhou et al.,

2022). Assuming a set amount of Aβ peptides, the formation of Aβ oligomers and aggregates in the

brain would, by chemical law, be predicted to result in the depletion of Aβ monomers (Dear et al., 2020;

Michaels et al., 2020). Thus, in the Alzheimer's disease brain, besides the obvious formation of Aβ

aggregates, there may also be a less visible depletion of Aβ monomers taking place at the same time,

which may, like Aβ oligomers, also contribute to the development of neuroinflammation and neuronal

damage (Huang, 2023). In support of this interpretation, high soluble brain Aβ42, which likely also

means high levels of Aβ monomers in the brain, have been found in clinical studies to preserve cognition

in patients of both familial and sporadic Alzheimer's disease, in spite of increasing amyloidosis detected

in their brains (Espay et al., 2021; Sturchio et al., 2022; Sturchio et al., 2021). In our study, we have also

found that the effects of microglial disinhibition are mediated by MMP9. Importantly, in

neurodegenerative diseases, MMP9 has been similarly found to be a key determinant regulatings the

selective degeneration of neuronal cell types (Kaplan et al., 2014; Tran et al., 2019). MMP9 levels are

also upregulated in the plasma in both mild cognitive impairment and Alzheimer's disease patients

(Bruno et al., 2009; Lorenzl et al., 2008; Tsiknia et al., 2022). In addition, in several motor neuron

disease models, reducing MMP9 has been found to protect neurons and delay the loss of motor

function (Kaplan et al., 2014; Spiller et al., 2019). Thus, our study has not only uncovered a potentially

overlooked role of Aβ monomer depletion in the development of Alzheimer's disease but also identified

downstream effectors. Elucidating the roles these factors play may reveal new insight into the

pathogenesis of Alzheimer's disease.

METHODS

Generation of *ric8a* conditional allele

Standard molecular biology techniques were employed for generating the conditional *ric-8a* allele.

Briefly, genomic fragments, of 4.5 and 2.5 kilobases and flanking exons 2-4 of the *ric-8a* locus at the 5'

and 3' side, respectively, were isolated by PCR using high fidelity polymerases. Targeting plasmid was

constructed by flanking the genomic fragment containing exons 2-4 with two loxP sites together with a

neomycin positive selection cassette, followed by 5' and 3' genomic fragments as homologous

recombination arms and a *pgk-DTA* gene as a negative selection cassette. ES cell clones were screened

by Southern blot analysis using external probes at 5' and 3' sides. For derivation of conditional allele, the

neomycin cassette was removed by crossing to an *actin-flpe* transgenic line after blastocyst injection and

germ line transmission. The primer set for genotyping *ric-8a* conditional allele, which produces a

wildtype band of ~110bp and a mutant band of ~200bp, is: 5'- cctagttgtgaatcagaagcacttg-3' and 5'-

gccatacctgagttacctaggc-3'. Animals homozygous for the conditional *ric-8a* allele are viable and fertile,

without obvious phenotypes.

Mouse breeding and pharmacology

emx1-cre, nestin-cre, foxg1-cre, cx3cr1-cre, floxed app as well as the *BAT-lacZ* reporter mouse lines were

purchased from the Jackson Lab. *nex-cre* and *wnt3a-cre* were as published (Goebbels et al., 2006; Yoshida

et al., 2006). *cre* transgenes were introduced individually into the *ric8a or app* conditional mutant

background for phenotypic analyses and *ric8a or app* homozygotes without *cre* as well as heterozygotes

with *cre* (littermates) were both analyzed as controls. For BB94 and MMP9/13 inhibitor injection, pregnant

females were treated daily from E12.5 to E14.5 at 30 μg (BB94) or 37.5 μg (MMP9/13 inhibitor) per g of

body weight. For dorsomorphin and S3I-201 injection, pregnant females were treated on E12.5 at 7.5 and 25

µg per g of body weight, respectively. For sham treatment, pregnant females were treated on E12.5 with

100 µls of DMSO. BrdU was injected at 100 µg per g of body weight, and embryos were collected 4 hours

later for cell proliferation analysis, or alternatively, pups were sacrificed at P5 for neuronal migration

analysis and at P17 for other analysis. For LPS treatment, pregnant females were injected intraperitoneally

with 400ng (*ric8a* genetic background) or 150ng (*app* genetic background) LPS per g of body weight on both

E11.5 and E12.5. Animal use was in accordance with institutional guidelines.

Immunohistochemistry

Vibratome sections from brains fixed in 4% paraformaldehyde were used. The following primary antibodies

were used at respective dilutions/concentrations: mouse anti-BrdU supernatant (clone G3G4,

Developmental Studies Hybridoma Bank (DSHB), University of Iowa, IA; 1:40), mouse anti-RC2 supernatant

(DSHB; 1:10), mouse anti-Nestin supernatant (DSHB; 1:20), mouse anti-Vimentin supernatant (DSHB; 1:10),

mouse anti-Pax6 supernatant (DSHB; 1:20), moue anti-Reelin (Millipore, 1:500), mouse anti-chondroitin

sulfate (CS-56, Sigma, 1:100), rat anti-Ctip2 (Abcam, 1:500), rabbit anti-phospho Histone H3 (Ser10)

(Millipore; 1:400), rabbit anti-Cux1 (CDP) (Santa Cruz; 1:100), rabbit anti-laminin (Sigma; 1:2000), rabbit anti-

GFAP (Dako;1:1000), rabbit anti-ALDH1L1 (Abcam, 1:500), rabbit anti-MMP9 (Abcam, 1:1000), goat anti-

MMP2 (R&D Systems; 5 µg/ml), rabbit anti-Calretinin (Chemicon, 1:2000), mouse anti-S100β (Thermo

Scientific; 1:100), rabbit anti-S100β (Thermo Scientific; 1:200), and rabbit anti-phospho-Smad1/5

(Ser463/465) (41D10; Cell Signaling, 1:200). FITC and Cy3 conjugated secondary antibodies were purchased

from Jackson ImmunoResearch Laboratories (West Grove, PA). Peroxidase conjugated secondary antibodies

were purchased from Santa Cruz Biotech. Staining procedures were performed as described previously

(Huang et al., 2006), except for anti-Ric-8a, MMP9, and phospho-Smad1/5 staining, in which a tyramide

signal amplification (TSA) plus Cy3 kit (PerkinElmer, Waltham, MA) was used per manufacturer's instruction.

Sections were mounted with Fluoromount G medium (Southern Biotech, Birmingham, AB) and analyzed

under a Nikon *eclipse* Ti microscope or an Olympus confocal microscope.

Microglia culture and assay

Cerebral hemispheres were dissected from individual neonates, mechanically dissociated, split into 3-4

wells each and cultured in DMEM-F12 (Lonza) containing 10% fetal bovine serum (FBS) (Invitrogen).

Microglial cells were harvested by light trypsinization that removes astroglial sheet on day 13-15. For

experiments other than assaying IL-1β secretion, microglia were treated with LPS at 20ng/ml for 3 hours

or at 5ng/ml overnight and, if applicable, DMSO or Aβ40 (ApexBio and Genscript) was applied at the

same time as LPS. For assaying IL-1β secretion, microglia were primed with LPS at 200ng/ml for 5-6

hours before treatment with 3mM ATP for 15 minutes. In these experiments, DMSO or Aβ40 was

applied at the same time as ATP if applicable. Supernatants were collected and used for cytokine ELISA

assays per manufacturer's instructions (Biolegend). Total RNAs were prepared from collected cells using

Trizol (Invitrogen) and cDNAs were synthesized using a High-capacity cDNA reverse transcription kit

(Applied Biosystems). Quantitative PCR was performed using a GoTaq qPCR master mix per

manufacturer's instructions (Promega). All gene expression levels were normalized against that of

GAPDH.

Quantitative and statistical analysis

The sample size was estimated to be 3-9 animals each genotype (every 4[th] of 50 μm coronal sections, 7-

10 sections each animal) for ectopia analysis, 3-5 animals each genotype (3-4 sections each animal) for

immunohistochemical analysis, and 4-6 animals each genotype for gel zymography and Western blot

analysis, as has been demonstrated by previous publications to be adequate for similar animal studies.

Matching sections were used between controls and mutants. NIS-Elements BR 3.0 software (Nikon) was

used for quantifying the numbers and sizes of neuronal ectopia, the numbers of laminin positive debris,

as well as the numbers of astrocytes. ImageJ software (NIH) was used for quantifying the intensity of

immunostainings. Statistics was performed using Student's t test when comparing two conditions, or

one-way ANOVA followed by Tukey's post hoc test when comparing three or more conditions. All data

are represented as means $\pm$ s.e.m.

ACKNOWLEDGMENTS

Z. H. thanks Dr. L. F. Reichardt for supporting the initial generation of *ric8a* mutant ES cells, Dr. E. A.

Grove (Chicago) for providing the *wnt3a-cre* strain, the late Dr. B. A. Barres (Stanford) for critiques and

input, and Drs. W. L. Murphy and E. Bresnick (UW-Madison) for access to a plate reader and a qPCR

machine. We also thank the late Dr. D. Oertel (UW-Madison) for critical reading and editing and Dr. L

Puglielli (UW-Madison) for critical reading of a previous version of the manuscript. This work was

supported by funds from the Departments of Neurology and Neuroscience, UW-Madison, and a Basil

O'Connor award from the March of Dimes foundation to Z.H.

AUTHOR INFORMATION

Z. H. designed experiments, generated *ric8a* conditional ES cells, performed microglial and related

experiments, and wrote the manuscript. H. J. K., D. S., and Z. H. performed other experiments and

analyzed data.

REFERENCES

Afshar, K., Willard, F.S., Colombo, K., Johnston, C.A., McCudden, C.R., Siderovski, D.P., and Gonczy, P. (2004). RIC-8 is required for GPR-1/2-dependent Galpha function during asymmetric division of C. elegans embryos. Cell *119*, 219-230.

Akol, I., Kalogeraki, E., Pielecka-Fortuna, J., Fricke, M., and Lowel, S. (2022). MMP2 and MMP9 Activity Is Crucial for Adult Visual Cortex Plasticity in Healthy and Stroke-Affected Mice. The Journal of neuroscience : the official journal of the Society for Neuroscience *42*, 16-32.

Barnes, S.J., Franzoni, E., Jacobsen, R.I., Erdelyi, F., Szabo, G., Clopath, C., Keller, G.B., and Keck, T. (2017). Deprivation-Induced Homeostatic Spine Scaling In Vivo Is Localized to Dendritic Branches that Have Undergone Recent Spine Loss. Neuron *96*, 871-882 e875.

Barres, B.A. (2008). The mystery and magic of glia: a perspective on their roles in health and disease. Neuron *60*, 430-440.

Beggs, H.E., Schahin-Reed, D., Zang, K., Goebbels, S., Nave, K.A., Gorski, J., Jones, K.R., Sretavan, D., and Reichardt, L.F. (2003). FAK deficiency in cells contributing to the basal lamina results in cortical abnormalities resembling congenital muscular dystrophies. Neuron *40*, 501-514.

Belvindrah, R., Nalbant, P., Ding, S., Wu, C., Bokoch, G.M., and Muller, U. (2006). Integrin-linked kinase regulates Bergmann glial differentiation during cerebellar development. Molecular and cellular neurosciences *33*, 109-125.

Bruno, M.A., Mufson, E.J., Wuu, J., and Cuello, A.C. (2009). Increased matrix metalloproteinase 9 activity in mild cognitive impairment. J Neuropathol Exp Neurol *68*, 1309-1318.

Chamberlain, L.M., Holt-Casper, D., Gonzalez-Juarrero, M., and Grainger, D.W. (2015). Extended culture of macrophages from different sources and maturation results in a common M2 phenotype. J Biomed Mater Res A *103*, 2864-2874.

Cirrito, J.R., Yamada, K.A., Finn, M.B., Sloviter, R.S., Bales, K.R., May, P.C., Schoepp, D.D., Paul, S.M., Mennerick, S., and Holtzman, D.M. (2005). Synaptic activity regulates interstitial fluid amyloid-beta levels in vivo. Neuron *48*, 913-922.

Colonna, M., and Butovsky, O. (2017). Microglia Function in the Central Nervous System During Health and Neurodegeneration. Annu Rev Immunol *35*, 441-468.

Couwenbergs, C., Spilker, A.C., and Gotta, M. (2004). Control of embryonic spindle positioning and Galpha activity by C. elegans RIC-8. Current biology : CB *14*, 1871-1876.

Cunningham, C.L., Martinez-Cerdeno, V., and Noctor, S.C. (2013). Microglia regulate the number of neural precursor cells in the developing cerebral cortex. The Journal of neuroscience : the official journal of the Society for Neuroscience *33*, 4216-4233.

David, N.B., Martin, C.A., Segalen, M., Rosenfeld, F., Schweisguth, F., and Bellaiche, Y. (2005). Drosophila Ric-8 regulates Galphai cortical localization to promote Galphai-dependent planar orientation of the mitotic spindle during asymmetric cell division. Nature cell biology *7*, 1083-1090.

Dear, A.J., Michaels, T.C.T., Meisl, G., Klenerman, D., Wu, S., Perrett, S., Linse, S., Dobson, C.M., and Knowles, T.P.J. (2020). Kinetic diversity of amyloid oligomers. Proceedings of the National Academy of Sciences of the United States of America *117*, 12087-12094.

Dziembowska, M., Milek, J., Janusz, A., Rejmak, E., Romanowska, E., Gorkiewicz, T., Tiron, A., Bramham, C.R., and Kaczmarek, L. (2012). Activity-dependent local translation of matrix metalloproteinase-9. The Journal of neuroscience : the official journal of the Society for Neuroscience *32*, 14538-14547.

Eichler, A., Kleidonas, D., Turi, Z., Fliegauf, M., Kirsch, M., Pfeifer, D., Masuda, T., Prinz, M., Lenz, M., and Vlachos, A. (2023). Microglial Cytokines Mediate Plasticity Induced by 10 Hz Repetitive Magnetic Stimulation. The Journal of neuroscience : the official journal of the Society for Neuroscience *43*, 3042-3060.

Espay, A.J., Sturchio, A., Schneider, L.S., and Ezzat, K. (2021). Soluble Amyloid-beta Consumption in Alzheimer's Disease. J Alzheimers Dis *82*, 1403-1415.

Fogel, H., Frere, S., Segev, O., Bharill, S., Shapira, I., Gazit, N., O'Malley, T., Slomowitz, E., Berdichevsky, Y., Walsh, D.M.*, et al.* (2014). APP homodimers transduce an amyloid-beta-mediated increase in release probability at excitatory synapses. Cell reports *7*, 1560-1576.

Gabay, M., Pinter, M.E., Wright, F.A., Chan, P., Murphy, A.J., Valenzuela, D.M., Yancopoulos, G.D., and Tall, G.G. (2011). Ric-8 proteins are molecular chaperones that direct nascent G protein alpha subunit membrane association. Science signaling *4*, ra79.

Galanis, C., Fellenz, M., Becker, D., Bold, C., Lichtenthaler, S.F., Muller, U.C., Deller, T., and Vlachos, A. (2021). Amyloid-Beta Mediates Homeostatic Synaptic Plasticity. The Journal of neuroscience : the official journal of the Society for Neuroscience *41*, 5157-5172.

Garcia-Osta, A., and Alberini, C.M. (2009). Amyloid beta mediates memory formation. Learn Mem *16*, 267-272.

Ginhoux, F., Greter, M., Leboeuf, M., Nandi, S., See, P., Gokhan, S., Mehler, M.F., Conway, S.J., Ng, L.G., Stanley, E.R.*, et al.* (2010). Fate mapping analysis reveals that adult microglia derive from primitive macrophages. Science *330*, 841-845.

Giuffrida, M.L., Caraci, F., Pignataro, B., Cataldo, S., De Bona, P., Bruno, V., Molinaro, G., Pappalardo, G., Messina, A., Palmigiano, A.*, et al.* (2009). Beta-amyloid monomers are neuroprotective. The Journal of neuroscience : the official journal of the Society for Neuroscience *29*, 10582-10587.

Goebbels, S., Bormuth, I., Bode, U., Hermanson, O., Schwab, M.H., and Nave, K.A. (2006). Genetic
targeting of principal neurons in neocortex and hippocampus of NEX-Cre mice. Genesis *44*, 611-621.

Gore, S.V., James, E.J., Huang, L.C., Park, J.J., Berghella, A., Thompson, A.C., Cline, H.T., and Aizenman,
C.D. (2021). Role of matrix metalloproteinase-9 in neurodevelopmental deficits and experience-
dependent plasticity in Xenopus laevis. eLife *10*.

Gorski, J.A., Talley, T., Qiu, M., Puelles, L., Rubenstein, J.L., and Jones, K.R. (2002). Cortical excitatory
neurons and glia, but not GABAergic neurons, are produced in the Emx1-expressing lineage. The Journal
of neuroscience : the official journal of the Society for Neuroscience *22*, 6309-6314.

Grant, J.L., Ghosn, E.E., Axtell, R.C., Herges, K., Kuipers, H.F., Woodling, N.S., Andreasson, K., Herzenberg,
L.A., Herzenberg, L.A., and Steinman, L. (2012). Reversal of paralysis and reduced inflammation from
peripheral administration of beta-amyloid in TH1 and TH17 versions of experimental autoimmune
encephalomyelitis. Sci Transl Med *4*, 145ra105.

Graus-Porta, D., Blaess, S., Senften, M., Littlewood-Evans, A., Damsky, C., Huang, Z., Orban, P., Klein, R.,
Schittny, J.C., and Muller, U. (2001). Beta1-class integrins regulate the development of laminae and folia
in the cerebral and cerebellar cortex. Neuron *31*, 367-379.

Guenette, S., Chang, Y., Hiesberger, T., Richardson, J.A., Eckman, C.B., Eckman, E.A., Hammer, R.E., and
Herz, J. (2006). Essential roles for the FE65 amyloid precursor protein-interacting proteins in brain
development. The EMBO journal *25*, 420-431.

Gulisano, W., Melone, M., Li Puma, D.D., Tropea, M.R., Palmeri, A., Arancio, O., Grassi, C., Conti, F., and
Puzzo, D. (2018). The effect of amyloid-beta peptide on synaptic plasticity and memory is influenced by
different isoforms, concentrations, and aggregation status. Neurobiol Aging *71*, 51-60.

Gulisano, W., Melone, M., Ripoli, C., Tropea, M.R., Li Puma, D.D., Giunta, S., Cocco, S., Marcotulli, D.,
Origlia, N., Palmeri, A.*, et al.* (2019). Neuromodulatory Action of Picomolar Extracellular Abeta42
Oligomers on Presynaptic and Postsynaptic Mechanisms Underlying Synaptic Function and Memory. The
Journal of neuroscience : the official journal of the Society for Neuroscience *39*, 5986-6000.

Halle, A., Hornung, V., Petzold, G.C., Stewart, C.R., Monks, B.G., Reinheckel, T., Fitzgerald, K.A., Latz, E.,
Moore, K.J., and Golenbock, D.T. (2008). The NALP3 inflammasome is involved in the innate immune
response to amyloid-beta. Nat Immunol *9*, 857-865.

Hampoelz, B., Hoeller, O., Bowman, S.K., Dunican, D., and Knoblich, J.A. (2005). Drosophila Ric-8 is
essential for plasma-membrane localization of heterotrimeric G proteins. Nat Cell Biol *7*, 1099-1105.

Hattori, Y., Kato, D., Murayama, F., Koike, S., Asai, H., Yamasaki, A., Naito, Y., Kawaguchi, A., Konishi, H.,
Prinz, M.*, et al.* (2023). CD206(+) macrophages transventricularly infiltrate the early embryonic cerebral
wall to differentiate into microglia. Cell reports *42*, 112092.

He, D., Xu, H., Zhang, H., Tang, R., Lan, Y., Xing, R., Li, S., Christian, E., Hou, Y., Lorello, P.*, et al.* (2022).
Disruption of the IL-33-ST2-AKT signaling axis impairs neurodevelopment by inhibiting microglial
metabolic adaptation and phagocytic function. Immunity *55*, 159-173 e159.

He, Y., Wei, M., Wu, Y., Qin, H., Li, W., Ma, X., Cheng, J., Ren, J., Shen, Y., Chen, Z.*, et al.* (2019). Amyloid
beta oligomers suppress excitatory transmitter release via presynaptic depletion of phosphatidylinositol-
4,5-bisphosphate. Nat Commun *10*, 1193.

Hebert, J.M., and McConnell, S.K. (2000). Targeting of cre to the Foxg1 (BF-1) locus mediates loxP
recombination in the telencephalon and other developing head structures. Developmental biology *222*,
296-306.

Heir, R., Abbasi, Z., Komal, P., Altimimi, H.F., Franquin, M., Moschou, D., Chambon, J., and Stellwagen, D.
(2024). Astrocytes Are the Source of TNF Mediating Homeostatic Synaptic Plasticity. The Journal of
neuroscience : the official journal of the Society for Neuroscience *44*.

Herms, J., Anliker, B., Heber, S., Ring, S., Fuhrmann, M., Kretzschmar, H., Sisodia, S., and Muller, U.
(2004). Cortical dysplasia resembling human type 2 lissencephaly in mice lacking all three APP family
members. The EMBO journal *23*, 4106-4115.

Huang, Z. (2023). A Function of Amyloid-beta in Mediating Activity-Dependent Axon/Synapse
Competition May Unify Its Roles in Brain Physiology and Pathology. J Alzheimers Dis *92*, 29-57.

Huang, Z. (2024). Evidence that Alzheimer's Disease Is a Disease of Competitive Synaptic Plasticity Gone
Awry. J Alzheimers Dis *99*, 447-470.

Huang, Z., Shimazu, K., Woo, N.H., Zang, K., Muller, U., Lu, B., and Reichardt, L.F. (2006). Distinct roles of
the beta 1-class integrins at the developing and the mature hippocampal excitatory synapse. J Neurosci
26, 11208-11219.

Iyer, H., Shen, K., Meireles, A.M., and Talbot, W.S. (2022). A lysosomal regulatory circuit essential for the
development and function of microglia. Sci Adv *8*, eabp8321.

Kamenetz, F., Tomita, T., Hsieh, H., Seabrook, G., Borchelt, D., Iwatsubo, T., Sisodia, S., and Malinow, R.
(2003). APP processing and synaptic function. Neuron *37*, 925-937.

Kaneko, M., Stellwagen, D., Malenka, R.C., and Stryker, M.P. (2008). Tumor necrosis factor-alpha
mediates one component of competitive, experience-dependent plasticity in developing visual cortex.
Neuron *58*, 673-680.

Kaplan, A., Spiller, K.J., Towne, C., Kanning, K.C., Choe, G.T., Geber, A., Akay, T., Aebischer, P., and
Henderson, C.E. (2014). Neuronal matrix metalloproteinase-9 is a determinant of selective
neurodegeneration. Neuron *81*, 333-348.

Kelly, E.A., Russo, A.S., Jackson, C.D., Lamantia, C.E., and Majewska, A.K. (2015). Proteolytic regulation of
synaptic plasticity in the mouse primary visual cortex: analysis of matrix metalloproteinase 9 deficient
mice. Front Cell Neurosci *9*, 369.

Kim, T., Vidal, G.S., Djurisic, M., William, C.M., Birnbaum, M.E., Garcia, K.C., Hyman, B.T., and Shatz, C.J.
(2013). Human LilrB2 is a beta-amyloid receptor and its murine homolog PirB regulates synaptic
plasticity in an Alzheimer's model. Science *341*, 1399-1404.

Lauren, J., Gimbel, D.A., Nygaard, H.B., Gilbert, J.W., and Strittmatter, S.M. (2009). Cellular prion protein mediates impairment of synaptic plasticity by amyloid-beta oligomers. Nature *457*, 1128-1132.

Lazarevic, V., Fienko, S., Andres-Alonso, M., Anni, D., Ivanova, D., Montenegro-Venegas, C., Gundelfinger, E.D., Cousin, M.A., and Fejtova, A. (2017). Physiological Concentrations of Amyloid Beta Regulate Recycling of Synaptic Vesicles via Alpha7 Acetylcholine Receptor and CDK5/Calcineurin Signaling. Front Mol Neurosci *10*, 221.

Leathers, T.A., and Rogers, C.D. (2022). Time to go: neural crest cell epithelial-to-mesenchymal transition. Development *149*.

Lee, Y., Park, B.H., and Bae, E.J. (2016). Compound C inhibits macrophage chemotaxis through an AMPK-independent mechanism. Biochemical and biophysical research communications *469*, 515-520.

LeVine, H., 3rd (2004). Alzheimer's beta-peptide oligomer formation at physiologic concentrations. Anal Biochem *335*, 81-90.

Lewitus, G.M., Konefal, S.C., Greenhalgh, A.D., Pribiag, H., Augereau, K., and Stellwagen, D. (2016). Microglial TNF-alpha Suppresses Cocaine-Induced Plasticity and Behavioral Sensitization. Neuron *90*, 483-491.

Liu, W., Yan, M., Liu, Y., Wang, R., Li, C., Deng, C., Singh, A., Coleman, W.G., Jr., and Rodgers, G.P. (2010). Olfactomedin 4 down-regulates innate immunity against Helicobacter pylori infection. Proceedings of the National Academy of Sciences of the United States of America *107*, 11056-11061.

Lorenzl, S., Buerger, K., Hampel, H., and Beal, M.F. (2008). Profiles of matrix metalloproteinases and their inhibitors in plasma of patients with dementia. Int Psychogeriatr *20*, 67-76.

Lorton, D., Kocsis, J.M., King, L., Madden, K., and Brunden, K.R. (1996). beta-Amyloid induces increased release of interleukin-1 beta from lipopolysaccharide-activated human monocytes. Journal of neuroimmunology *67*, 21-29.

Ma, S., Kwon, H.J., and Huang, Z. (2012). Ric-8a, a guanine nucleotide exchange factor for heterotrimeric G proteins, regulates bergmann glia-basement membrane adhesion during cerebellar foliation. The Journal of neuroscience : the official journal of the Society for Neuroscience *32*, 14979-14993.

Ma, S., Santhosh, D., Kumar, T.P., and Huang, Z. (2017). A Brain-Region-Specific Neural Pathway Regulating Germinal Matrix Angiogenesis. Developmental cell *41*, 366-381 e364.

Michaels, T.C.T., Saric, A., Curk, S., Bernfur, K., Arosio, P., Meisl, G., Dear, A.J., Cohen, S.I.A., Dobson, C.M., Vendruscolo, M.*, et al.* (2020). Dynamics of oligomer populations formed during the aggregation of Alzheimer's Abeta42 peptide. Nat Chem *12*, 445-451.

Milosch, N., Tanriover, G., Kundu, A., Rami, A., Francois, J.C., Baumkotter, F., Weyer, S.W., Samanta, A., Jaschke, A., Brod, F.*, et al.* (2014). Holo-APP and G-protein-mediated signaling are required for sAPPalpha-induced activation of the Akt survival pathway. Cell death & disease *5*, e1391.

Moore, S.A., Saito, F., Chen, J., Michele, D.E., Henry, M.D., Messing, A., Cohn, R.D., Ross-Barta, S.E., Westra, S., Williamson, R.A.*, et al.* (2002). Deletion of brain dystroglycan recapitulates aspects of congenital muscular dystrophy. Nature *418*, 422-425.

Morley, J.E., Farr, S.A., Banks, W.A., Johnson, S.N., Yamada, K.A., and Xu, L. (2010). A physiological role for amyloid-beta protein:enhancement of learning and memory. J Alzheimers Dis *19*, 441-449.

Muehlhauser, F., Liebl, U., Kuehl, S., Walter, S., Bertsch, T., and Fassbender, K. (2001). Aggregation-Dependent interaction of the Alzheimer's beta-amyloid and microglia. Clin Chem Lab Med *39*, 313-316.

Murase, S., Lantz, C.L., and Quinlan, E.M. (2017). Light reintroduction after dark exposure reactivates plasticity in adults via perisynaptic activation of MMP-9. eLife *6*.

Murase, S., Winkowski, D., Liu, J., Kanold, P.O., and Quinlan, E.M. (2019). Homeostatic regulation of perisynaptic matrix metalloproteinase 9 (MMP9) activity in the amblyopic visual cortex. eLife *8*.

Niewmierzycka, A., Mills, J., St-Arnaud, R., Dedhar, S., and Reichardt, L.F. (2005). Integrin-linked kinase deletion from mouse cortex results in cortical lamination defects resembling cobblestone lissencephaly. The Journal of neuroscience : the official journal of the Society for Neuroscience *25*, 7022-7031.

Nishimoto, I., Okamoto, T., Matsuura, Y., Takahashi, S., Okamoto, T., Murayama, Y., and Ogata, E. (1993). Alzheimer amyloid protein precursor complexes with brain GTP-binding protein G(o). Nature *362*, 75-79.

Pagenstecher, A., Stalder, A.K., Kincaid, C.L., Shapiro, S.D., and Campbell, I.L. (1998). Differential expression of matrix metalloproteinase and tissue inhibitor of matrix metalloproteinase genes in the mouse central nervous system in normal and inflammatory states. Am J Pathol *152*, 729-741.

Palmeri, A., Ricciarelli, R., Gulisano, W., Rivera, D., Rebosio, C., Calcagno, E., Tropea, M.R., Conti, S., Das, U., Roy, S.*, et al.* (2017). Amyloid-beta Peptide Is Needed for cGMP-Induced Long-Term Potentiation and Memory. The Journal of neuroscience : the official journal of the Society for Neuroscience *37*, 6926-6937.

Parodi, J., Sepulveda, F.J., Roa, J., Opazo, C., Inestrosa, N.C., and Aguayo, L.G. (2010). Beta-amyloid causes depletion of synaptic vesicles leading to neurotransmission failure. The Journal of biological chemistry *285*, 2506-2514.

Patani, R., Hardingham, G.E., and Liddelow, S.A. (2023). Functional roles of reactive astrocytes in neuroinflammation and neurodegeneration. Nat Rev Neurol *19*, 395-409.

Plant, L.D., Boyle, J.P., Smith, I.F., Peers, C., and Pearson, H.A. (2003). The production of amyloid beta peptide is a critical requirement for the viability of central neurons. The Journal of neuroscience : the official journal of the Society for Neuroscience *23*, 5531-5535.

Puzzo, D., Privitera, L., Leznik, E., Fa, M., Staniszewski, A., Palmeri, A., and Arancio, O. (2008). Picomolar amyloid-beta positively modulates synaptic plasticity and memory in hippocampus. The Journal of neuroscience : the official journal of the Society for Neuroscience *28*, 14537-14545.

Qin, H., Yeh, W.I., De Sarno, P., Holdbrooks, A.T., Liu, Y., Muldowney, M.T., Reynolds, S.L., Yanagisawa,
L.L., Fox, T.H., 3rd, Park, K*., et al.* (2012). Signal transducer and activator of transcription-3/suppressor of
cytokine signaling-3 (STAT3/SOCS3) axis in myeloid cells regulates neuroinflammation. Proceedings of
the National Academy of Sciences of the United States of America *109*, 5004-5009.

Ramaker, J.M., Swanson, T.L., and Copenhaver, P.F. (2013). Amyloid precursor proteins interact with the
heterotrimeric G protein Go in the control of neuronal migration. The Journal of neuroscience : the
official journal of the Society for Neuroscience *33*, 10165-10181.

Ramsden, M., Henderson, Z., and Pearson, H.A. (2002). Modulation of Ca2+ channel currents in primary
cultures of rat cortical neurones by amyloid beta protein (1-40) is dependent on solubility status. Brain
research *956*, 254-261.

Rice, H.C., Townsend, M., Bai, J., Suth, S., Cavanaugh, W., Selkoe, D.J., and Young-Pearse, T.L. (2012).
Pancortins interact with amyloid precursor protein and modulate cortical cell migration. Development
139, 3986-3996.

Satz, J.S., Ostendorf, A.P., Hou, S., Turner, A., Kusano, H., Lee, J.C., Turk, R., Nguyen, H., Ross-Barta, S.E.,
Westra, S*., et al.* (2010). Distinct functions of glial and neuronal dystroglycan in the developing and adult
mouse brain. J Neurosci *30*, 14560-14572.

Sayed, F.A., Telpoukhovskaia, M., Kodama, L., Li, Y., Zhou, Y., Le, D., Hauduc, A., Ludwig, C., Gao, F.,
Clelland, C*., et al.* (2018). Differential effects of partial and complete loss of TREM2 on microglial injury
response and tauopathy. Proceedings of the National Academy of Sciences of the United States of
America *115*, 10172-10177.

Schafer, D.P., and Stevens, B. (2015). Microglia Function in Central Nervous System Development and
Plasticity. Cold Spring Harb Perspect Biol *7*, a020545.

Shaked, G.M., Kummer, M.P., Lu, D.C., Galvan, V., Bredesen, D.E., and Koo, E.H. (2006). Abeta induces
cell death by direct interaction with its cognate extracellular domain on APP (APP 597-624). FASEB J *20*,
1254-1256.

Shankar, G.M., Li, S., Mehta, T.H., Garcia-Munoz, A., Shepardson, N.E., Smith, I., Brett, F.M., Farrell,
M.A., Rowan, M.J., Lemere, C.A*., et al.* (2008). Amyloid-beta protein dimers isolated directly from
Alzheimer's brains impair synaptic plasticity and memory. Nat Med *14*, 837-842.

Shigemoto-Mogami, Y., Hoshikawa, K., Goldman, J.E., Sekino, Y., and Sato, K. (2014). Microglia enhance
neurogenesis and oligodendrogenesis in the early postnatal subventricular zone. The Journal of
neuroscience : the official journal of the Society for Neuroscience *34*, 2231-2243.

Spiller, K.J., Khan, T., Dominique, M.A., Restrepo, C.R., Cotton-Samuel, D., Levitan, M., Jafar-Nejad, P.,
Zhang, B., Soriano, A., Rigo, F*., et al.* (2019). Reduction of matrix metalloproteinase 9 (MMP-9) protects
motor neurons from TDP-43-triggered death in rNLS8 mice. Neurobiol Dis *124*, 133-140.

Spolidoro, M., Putignano, E., Munafo, C., Maffei, L., and Pizzorusso, T. (2012). Inhibition of matrix
metalloproteinases prevents the potentiation of nondeprived-eye responses after monocular
deprivation in juvenile rats. Cereb Cortex *22*, 725-734.

Stellwagen, D., and Malenka, R.C. (2006). Synaptic scaling mediated by glial TNF-alpha. Nature *440*,
1054-1059.

Stenman, J., Toresson, H., and Campbell, K. (2003). Identification of two distinct progenitor populations
in the lateral ganglionic eminence: implications for striatal and olfactory bulb neurogenesis. The Journal
of neuroscience : the official journal of the Society for Neuroscience *23*, 167-174.

Stine, W.B., Jungbauer, L., Yu, C., and LaDu, M.J. (2011). Preparing synthetic Abeta in different
aggregation states. Methods Mol Biol *670*, 13-32.

Sturchio, A., Dwivedi, A.K., Malm, T., Wood, M.J.A., Cilia, R., Sharma, J.S., Hill, E.J., Schneider, L.S., Graff-
Radford, N.R., Mori, H.*, et al.* (2022). High Soluble Amyloid-beta42 Predicts Normal Cognition in
Amyloid-Positive Individuals with Alzheimer's Disease-Causing Mutations. J Alzheimers Dis.

Sturchio, A., Dwivedi, A.K., Young, C.B., Malm, T., Marsili, L., Sharma, J.S., Mahajan, A., Hill, E.J.,
Andaloussi, S.E., Poston, K.L.*, et al.* (2021). High cerebrospinal amyloid-beta 42 is associated with normal
cognition in individuals with brain amyloidosis. EClinicalMedicine *38*, 100988.

Tall, G.G., Krumins, A.M., and Gilman, A.G. (2003). Mammalian Ric-8A (synembryn) is a heterotrimeric
Galpha protein guanine nucleotide exchange factor. The Journal of biological chemistry *278*, 8356-8362.

Tan, J., Town, T., Paris, D., Mori, T., Suo, Z., Crawford, F., Mattson, M.P., Flavell, R.A., and Mullan, M.
(1999). Microglial activation resulting from CD40-CD40L interaction after beta-amyloid stimulation.
Science *286*, 2352-2355.

Tran, N.M., Shekhar, K., Whitney, I.E., Jacobi, A., Benhar, I., Hong, G., Yan, W., Adiconis, X., Arnold, M.E.,
Lee, J.M.*, et al.* (2019). Single-Cell Profiles of Retinal Ganglion Cells Differing in Resilience to Injury
Reveal Neuroprotective Genes. Neuron *104*, 1039-1055 e1012.

Tsiknia, A.A., Sundermann, E.E., Reas, E.T., Edland, S.D., Brewer, J.B., Galasko, D., Banks, S.J., and
Alzheimer's Disease Neuroimaging, I. (2022). Sex differences in Alzheimer's disease: plasma MMP-9 and
markers of disease severity. Alzheimers Res Ther *14*, 160.

Vainchtein, I.D., Chin, G., Cho, F.S., Kelley, K.W., Miller, J.G., Chien, E.C., Liddelow, S.A., Nguyen, P.T.,
Nakao-Inoue, H., Dorman, L.C.*, et al.* (2018). Astrocyte-derived interleukin-33 promotes microglial
synapse engulfment and neural circuit development. Science *359*, 1269-1273.

Walsh, D.M., Klyubin, I., Fadeeva, J.V., Cullen, W.K., Anwyl, R., Wolfe, M.S., Rowan, M.J., and Selkoe, D.J.
(2002). Naturally secreted oligomers of amyloid beta protein potently inhibit hippocampal long-term
potentiation in vivo. Nature *416*, 535-539.

Wang, H., Ng, K.H., Qian, H., Siderovski, D.P., Chia, W., and Yu, F. (2005). Ric-8 controls Drosophila neural
progenitor asymmetric division by regulating heterotrimeric G proteins. Nat Cell Biol *7*, 1091-1098.

Wang, X., Jung, J., Asahi, M., Chwang, W., Russo, L., Moskowitz, M.A., Dixon, C.E., Fini, M.E., and Lo, E.H.
(2000). Effects of matrix metalloproteinase-9 gene knock-out on morphological and motor outcomes
after traumatic brain injury. The Journal of neuroscience : the official journal of the Society for
Neuroscience *20*, 7037-7042.

Wang, Y., Fu, W.Y., Cheung, K., Hung, K.W., Chen, C., Geng, H., Yung, W.H., Qu, J.Y., Fu, A.K.Y., and Ip,
N.Y. (2021). Astrocyte-secreted IL-33 mediates homeostatic synaptic plasticity in the adult hippocampus.
Proceedings of the National Academy of Sciences of the United States of America *118*.

Yang, Y., Kim, J., Kim, H.Y., Ryoo, N., Lee, S., Kim, Y., Rhim, H., and Shin, Y.K. (2015). Amyloid-beta
Oligomers May Impair SNARE-Mediated Exocytosis by Direct Binding to Syntaxin 1a. Cell reports *12*,
1244-1251.

Yona, S., Kim, K.W., Wolf, Y., Mildner, A., Varol, D., Breker, M., Strauss-Ayali, D., Viukov, S., Guilliams, M.,
Misharin, A.*, et al.* (2013). Fate mapping reveals origins and dynamics of monocytes and tissue
macrophages under homeostasis. Immunity *38*, 79-91.

Yoshida, M., Assimacopoulos, S., Jones, K.R., and Grove, E.A. (2006). Massive loss of Cajal-Retzius cells
does not disrupt neocortical layer order. Development *133*, 537-545.

Young-Pearse, T.L., Bai, J., Chang, R., Zheng, J.B., LoTurco, J.J., and Selkoe, D.J. (2007). A critical function
for beta-amyloid precursor protein in neuronal migration revealed by in utero RNA interference. The
Journal of neuroscience : the official journal of the Society for Neuroscience *27*, 14459-14469.

Yu, D., Li, T., Delpech, J.C., Zhu, B., Kishore, P., Koshi, T., Luo, R., Pratt, K.J.B., Popova, G., Nowakowski,
T.J.*, et al.* (2022). Microglial GPR56 is the molecular target of maternal immune activation-induced
parvalbumin-positive interneuron deficits. Sci Adv *8*, eabm2545.

Zhang, Y., Chen, K., Sloan, S.A., Bennett, M.L., Scholze, A.R., O'Keeffe, S., Phatnani, H.P., Guarnieri, P.,
Caneda, C., Ruderisch, N.*, et al.* (2014). An RNA-sequencing transcriptome and splicing database of glia,
neurons, and vascular cells of the cerebral cortex. The Journal of neuroscience : the official journal of the
Society for Neuroscience *34*, 11929-11947.

Zhou, B., Lu, J.G., Siddu, A., Wernig, M., and Sudhof, T.C. (2022). Synaptogenic effect of APP-Swedish
mutation in familial Alzheimer's disease. Sci Transl Med *14*, eabn9380.

Zipp, F., Bittner, S., and Schafer, D.P. (2023). Cytokines as emerging regulators of central nervous system
synapses. Immunity *56*, 914-925.

Zott, B., Simon, M.M., Hong, W., Unger, F., Chen-Engerer, H.J., Frosch, M.P., Sakmann, B., Walsh, D.M.,
and Konnerth, A. (2019). A vicious cycle of beta amyloid-dependent neuronal hyperactivation. Science
365, 559-565.

Figures and legends

Fig. 1. Deletion of *ric8a* using *emx1-cre* results in cortical ectopia due to non-neural deficits.

(**a-d**) Nissl staining of control (ctrl, **a**&**c**) and mutant (mt, **b**&**d**) anterior motor (**a-b**) and posterior

somatosensory (**c-d**) cortex at P0.

(**e**-**e'**) Laminin (LN, in green) and nuclear (DAPI, in blue) staining of control cortices at P0. A continuous

basement membrane is observed at the pia, beneath which cells are well organized in the cortical wall.

(**f-f'**) Staining of *ric8a-emx1-cre* mutant cortices at P0. Basement membrane breach and neuronal

ectopia are observed following *ric8a* deletion by *emx1-cre*, a *cre* line expressed in cortical radial glial

progenitors beginning at E10.5.

(g-g′) Staining of *ric8a-nestin-cre* mutant cortices at P0. No obvious basement membrane breach or

neuronal ectopia is observed following *ric8a* deletion by *nestin-cre*, a *cre* line expressed in cortical

progenitors beginning around E12.5.

(h-h′) Staining of *ric8a-foxg1-cre* mutant cortices at P0. No obvious basement membrane breach or

neuronal ectopia is observed following *ric8a* deletion by *foxg1-cre*, a *cre* line expressed in forebrain

neural progenitors from E9.0.

Scale bars, 640μm for (**a-b**), 400μm for (**c-d**), and 100μm for (**e-h′**).

Fig. 2. *ric8a* deficiency in microglia is responsible for cortical ectopia.

(**a**) TNFα, IL-1β, and IL-6 secretion (pg/ml) in control and *ric8a/cx3cr1-cre* mutant microglia following LPS

stimulation. *, *P* < 0.05; **, *P* < 0.01; ***, *P* < 0.001; n = 6-8 each group.

(**b**) TNFα, IL-1 β, and IL-6 mRNA expression in control and *ric8a/cx3cr1-cre* mutant microglia following

LPS stimulation. *, *P* < 0.05; ***, *P* < 0.001; n = 5-6 each group.

(**c-d**) Nuclear (DAPI, in grey) staining of *ric8a-cx3cr1-cre* mutant cortices at P0 in the absence (**c**) or

presence (**d**) of LPS treatment during embryogenesis.

(**e-f**) Nuclear (DAPI, in grey) staining of *ric8a-nestin-cre* single *cre* (**e**) and *ric8a-nestin-cre+cx3cr1-cre*

double *cre* (**f**) mutant cortices at P0.

Scale bar in (**c**), 100 μm for (**c-f**).

Fig. 3. *app* deficiency results in hypersensitive microglia and cortical ectopia.

(a) TNFα and IL-1β secretion (pg/ml) in cultured control and *app/cx3cr1-cre* mutant microglia following

LPS stimulation. *, *P* < 0.05; n = 7-9 each group.

(**b**) TNFα, IL-1β, IL-6, and MCP1 secretion (pg/ml) in fresh unelicited control and *app/cx3cr1-cre* mutant

peritoneal macrophages following LPS stimulation. ***, *P* < 0.001; n = 7-10 each group.

(**c**) TNFα, IL-1 β, IL-6, and IL-23 mRNA expression in fresh unelicited control and *app/cx3cr1-cre* mutant

peritoneal macrophages following LPS stimulation. **, *P* < 0.01; ***, *P* < 0.001; n = 6 each group

(**d-e**) Nuclear (DAPI, in blue) staining of control (**D**) and LPS-treated *app/cx3cr1-cre* mutant (**E**) cortices at

P0. Note cortical ectopia in the mutant cortex (arrowhead).

Scale bar in (**d**), 200μm for (**d-e**).

Fig. 4. Monomeric Aβ40 suppresses microglia via an APP and Ric8a.

(a) TNFα, IL-6, IL-1β, and MCP1 secretion (pg/ml) by wildtype microglia following LPS stimulation in the

absence or presence of Aβ40 (200 or 500nM). *, $P < 0.05$; ***, $P < 0.001$; n = 8-14 each group.

(b) TNFα and IL-1β secretion (pg/ml) by wildtype microglia following poly I:C stimulation in the absence

or presence of Aβ40 (500nM). *, $P < 0.05$; **, $P < 0.01$; n = 6-7 each group.

(c) IL-6 and IL-1β mRNA induction in wildtype microglia following LPS stimulation in the absence or

presence of Aβ40 (500nM). *, $P < 0.05$; n = 6 each group.

(d) TNFα and IL-6 secretion (pg/ml) by control and *app/cx3cr1-cre* mutant microglia following LPS

stimulation in the absence or presence of Aβ40 (200nM). **, $P < 0.01$; ***, $P < 0.001$; n = 8 each group.

(e) IL-6 and IL-1β mRNA induction in control and *app/cx3cr1-cre* mutant microglia following LPS

stimulation in the absence or presence of Aβ40 (200nM). *, $P < 0.05$; **, $P < 0.01$; n = 6 each group.

(f) TNFα and IL-6 secretion (pg/ml) by control and *app/cx3cr1-cre* mutant peritoneal macrophages

following LPS stimulation in the absence or presence of Aβ40 (500nM). *, $P < 0.05$; n = 6-7 each group

(g) TNFα and IL-6 secretion (pg/ml) by control and *ric8a/cx3cr1-cre* mutant microglia following LPS

stimulation in the absence or presence of Aβ40 (200nM). ***, $P < 0.001$; n = 12-14 each group.

Fig. 5. Inhibition of both microglial inflammatory activation and cortical MMP9 activity suppresses

basement membrane breach and neuronal ectopia.

(a-d) Nuclear (DAPI, in grey) staining of untreated (**a**), anti-inflammatory drug dorsomorphin (DM) (**b**),

Stat3 inhibitor S3I-201 (S3I) (**c**), and DM/S3I (**d**) dual treated mutant cortices at P0.

(e-f) Quantitative analysis of ectopia number (**e**) and size (**f**) in the neonatal mutant cortex after DMSO,

DM, S3I, and DM/S3I dual treatment at E12.5. *, $P < 0.05$; ***, $P < 0.001$; all compared to untreated

mutants. The reduction in ectopia size after dual treatment is not statistically significant, likely due to

the small number of ectopias that remained.

(**g-h**) MMP9 (in red) staining of control (**g**) and mutant cortices (**h**) at E13.5. Quantification shows

statistically significant increases in mutants (control, 24.8 ± 0.2 AU (Arbitrary Units); mutant, 35.7 ± 1.7

AU; *P* = 0.002; n = 6).

(**i**) Gel zymography of control and mutant cortical lysates at E13.5. Increased levels of MMP9 but not of

MMP2 were observed in mutants (control, 1.00 ± 0.06 AU; mutant, 3.72 ±1.86 AU; *P* = 0.028; n = 4)

(**j-k'**) Laminin (in green) and nuclear (DAPI, in blue) staining of mutant cortices untreated (**H**) or treated

(**I**) with BB94.

(**l-m**) Quantitative analysis of ectopia number and size following MMP inhibitor BB94 or MMP9/13

inhibitor treatment. *, *P* < 0.05; ***, *P* < 0.001; all compared to untreated mutants.

Supplemental Fig. 1. Birth-dating of early and late-born neurons in *ric8a-emx1-cre* mutant cortices.

(**a-c**) BrdU (in red) staining in control (**a**) and mutant (**b**) cortices at P5 after administration at E12.5.

Quantification is shown in (**c**). No statistically significant differences were observed between control

and mutant neurons in regions without ectopia.

(**d-f**) BrdU staining in control (**d**) and mutant (**e**) cortices at P5 after administration at E15.5.

Quantification is shown in (**f**). Neuronal migration appears slightly delayed in mutants as compared to

controls. *, $P < 0.05$; **, $P < 0.01$; n = 5.

Supplemental Fig. 2. Lamina-specific neuronal markers are normal outside ectopia in *ric8a-emx1-cre*

mutant cortices.

(a-b") Cux1 (in red) and nuclear (DAPI, in blue) staining of control (**a-a**") and mutant (**b-b**") cortices at P0

in a region without ectopia. No obvious changes in the expression pattern of Cux1, an upper layer

neuronal marker, were observed in the mutant cortex, except in areas with ectopia (see panel (**g**)).

(**c**-**d**") Ctip2 (in red) and nuclear (DAPI, in blue) staining of control (**c**-**c**") and mutant (**d**-**d**") cortices at P0.

No obvious changes in the expression pattern of Ctip2, a deep layer neuronal marker, were observed in

the mutant cortex, except in areas with ectopia.

(**e-f**) Quantification of cortical neurons positive for Cux1 (**e**) and Ctip2 (**f**) in matching cortical regions at

P0. No significant differences were observed in the density of Cux1 (control, 218.1 ± 1.7 per 100 μm

cortical width; mutant, 216.4 ± 4.3 per 100 μm cortical width; $P = 0.36$, n = 12) or Ctip2 (control, $157.8 \pm$

5.0 per field; mutant, 161.9 ± 5.9 per field; $P = 0.31$, n = 12) positive neurons between controls and

mutants.

(**g**) Cux1 (in red) staining of mutant cortices at P0 in a region with ectopia.

Scale bar in (**a**), 200 μm for all panels.

Supplemental Fig. 3. Neuronal ectopia in *ric8a-emx1-cre* mutants result from pial basement

membrane breach during embryogenesis.

(a-a") Laminin (LN, in green), radial glial marker RC2 (in red), and nuclear (DAPI, in blue) staining of

control cortices at E16.5. A continuous basement membrane is observed at the pia, where radial glial

endfeet are anchored.

(**b**-**b"**) Laminin, RC2, and nuclear staining of *ric8a/emx1-cre* mutant cortices at E16.5. Neuronal ectopias

are consistently observed at sites of basement membrane breakage (arrowheads in **b**). Radial glial fibers

at these sites extend beyond the pia (inset in **b'**).

(**c**-**e**) Calretinin (CR, in green) and nuclear (DAPI, in blue) staining of control (**c**) and mutant (**d-e**) cortices

at E16.5. A continuous row of Calretinin positive Cajal-Retzius cells is observed in the marginal zone of

control cortices (**c**). By contrast, in mutants, Cajal-Retzius cells are absent at large ectopias (**d**). However,

they appear passively displaced by over-migrating neurons at small ectopias (arrowhead in **e**).

(**f-g**) Reelin (Reln, in red) and nuclear (DAPI, in blue) staining of control (**f**) and *ric8a/emx1-cre* mutant (**g**)

cortices at E15.5. Strong Reelin expression is observed in Cajal-Retzius cells in the marginal zone of both

control and mutant cortices.

(**h-i**) Chondroitin sulfate proteoglycan (CS56, in red) staining of control (**h**) and mutant (**i**) cortices at

E14.5. Normal preplate splitting is observed in mutants.

Scale bar in (**a**), 200μm for (**a**-**g**) and 500μm for (**h-i**).

1004

1005

Supplemental Fig. 4. Basement membrane breaches precede neuronal ectopia in *ric8a-emx1-cre* mutant

cortices.

(a-a") Laminin (LN, in green) and nuclear (DAPI, in blue) staining of control cortices at E13.5. A

continuous basement membrane is observed at the pia, beneath which cells are well organized in the

cortical wall.

(**b-b"**) Laminin and nuclear staining of *ric8a-emx1-cre* mutant cortices at E13.5. In a subset of mutants, a

small disruption of basement membrane is observed (bracket and inset in **b**), but not yet associated with

ectopia (arrowhead in **b'**).

(c-c") Laminin (LN, in green) and nuclear (DAPI, in blue) staining of control cortices at E13.5.

(**d-d"**) Laminin and nuclear staining of *ric8a-emx1-cre* mutant cortices at E13.5. Although at E13.5 we

observe basement membrane defects in the absence of neuronal ectopia (see **b-b"**), when there are

neuronal ectopia, they are always associated with basement membrane breakage.

(**e-e"**) Laminin and nuclear staining of control cortices at E14.5.

(**f-f"**) Laminin and nuclear staining of *ric-8a-emx1-cre* mutant cortices at E14.5. Neuronal ectopia at

E14.5 are also always associated with basement membrane breakage.

Scale bar in (**a**), 100 μm for all panels.

1022

1023

1024 **Supplemental Fig. 5. Signs of basement membrane degradation before breach formation at E12.5.**

1025 (**a-b**) Laminin (in green) staining of control (**a**) and *ric8a-emx1-cre* mutant (**b**) cortices at E12.5. Increased

1026 numbers of laminin positive debris were observed in mutants (compare insets), even though breaches

1027 had yet to form.

1028 (**c**) Quantitative analysis shows significant increases.

1029

1030

Supplemental Fig. 6. Cortical radial glial identity and proliferation are unaffected in *ric8a-emx1-cre*

mutants.

(a-b') Pax6 (in red) and nuclear (DAPI, in blue) staining of control (**a&a'**) and mutant (**b&b'**) cortices at

E12.5.

(**c-d'**) Pax6 (in red) and nuclear (DAPI, in blue) staining of control (**c&c'**) and mutant (**d&d'**) cortices at

E14.5. No ectopic Pax6 positive cells were observed at either E12.5 or E14.5.

(**e-f**) Nestin (in red) and nuclear (DAPI, in blue) staining of control (**e**) and mutant (**f**) cortices at E14.5.

(**g-h**) Vimentin (in red) and nuclear (DAPI, in blue) staining of control (**g**) and mutant (**h**) cortices at E14.5.

(**i-j**) BrdU staining (in red) in control (**i**) and mutant (**j**) cortices at E13.5.

(**k-l**) BrdU staining (in red) in control (**k**) and mutant (**l**) cortices at E15.5.

(**m-n**) Phospho-histone 3 (PH3, in green) and nuclear (DAPI, in blue) staining of control (**m**) and mutant

(**n**) cortices at E14.5.

(**o-p**) Cleavage plane distribution of radial glial mitosis in control (**o**) and mutant (**p**) cortices at E14.5. No

significant differences were observed ($P > 0.4$, n = 3 animals each genotype; 73 cells for controls and 76

cells for mutants).

(**q-r**) Cleavage plane distribution of radial glial mitosis in control (**q**) and mutant (**r**) cortices at E15.5. No

significant differences were observed ($P > 0.1$, n = 3 animals each genotype; 70 cells for controls and 59

cells for mutants).

Scale bar in (**a**), 100 μm for (**a-b'**) and 200 μm for (**c-n**).

1050

1051

1052 **Supplemental Fig. 7. Wnt pathway activity is normal in *ric8a-emx1-cre* mutant cortices.**

1053 X-gal staining of BAT-lacZ expression in *ric8a-emx1-cre* control (**a**) and mutant (**b**-**d**) cortices at E13.5. No

1054 obvious differences are observed between controls and three different mutants at this stage.

1055

1056

1057

1058

1059 **Supplemental Fig. 8. *emx1-cre* is active in microglia.**

1060 TNFα secretion (pg/ml) (**a**) and basal TNFα and IL-1β mRNA expression (**b**) in control and *ric8a-emx1-cre*

1061 mutant microglia. *, *P* < 0.05; n = 5-8 each group.

1062

1063

Supplemental Fig. 9. Cytokine secretion and transcriptional induction in *app-cx3cr1-cre* mutant

microglia.

(a) TNFα and IL-6 secretion (pg/ml) in control and *app-cx3cr1-cre* mutant microglia following overnight

LPS stimulation. *, $P < 0.05$; **, $P < 0.01$; n = 9-13 each group.

(**b**) TNFα, IL-1β and IL-6 mRNA expression in control and *app-cx3cr1-cre* mutant microglia following

overnight 3-hr LPS stimulation. *, $P < 0.05$; n = 6-7 each group.

1070

1071

Supplemental Fig. 10. Effects of monomeric Aβ on cytokine secretion and transcription in control and

1072

mutant microglial lineage cells.

1073

(a) TNFα and IL-6 secretion (pg/ml) in wildtype microglia following LPS stimulation in the absence or

presence of Aβ40 (50nM). *, $P < 0.05$; **, $P < 0.01$; n = 25 each group for TNFα and 11 each group for IL-

6.

(**b**) TNFα and MCP1 secretion (pg/ml) in wildtype microglia following LPS stimulation in the absence or

presence of Aβ40 (500nM) from Genscript. Effects on IL-1β secretion in **Fig. 4b** was also performed with

Genscript Aβ40. All other experiments in Fig. 7 were performed with ApexBio Aβ40. *, $P < 0.05$; **, $P <$

0.01; n = 5-7 each group.

(**c**) TNFα, IL-23, and IL-10 mRNA expression in wildtype microglia following LPS stimulation in the

absence or presence of Aβ40 (400nM). *, $P < 0.05$; n = 6 each group

(d) IL-1β secretion (pg/ml) in control and *app/cx3cr1-cre* mutant microglia following LPS stimulation in

the absence or presence of Aβ40. *, $P < 0.05$; n = 8-12 each group.

(e) TNFα (pg/ml) in control or *aplp2/cx3cr1-cre* mutant microglia following LPS stimulation in the

absence or presence of Aβ40 (400nM). *, $P < 0.05$; n = 9-13 each group.

(f) IL-10 and IL-23 mRNA expression in control and *app/cx3cr1-cre* mutant microglia following LPS

stimulation in the absence or presence of Aβ40 (400nM). *, $P < 0.05$; n = 6 each group.

(g) IL-1β secretion (pg/ml) in fresh unelicited control and *app/cx3cr1-cre* mutant peritoneal

macrophages following LPS stimulation in the absence or presence of Aβ40 (400nM). *, $P < 0.05$; n = 12

each group.

(**h**) IL-1β secretion (pg/ml) in f control and *ric8a/cx3cr1-cre* mutant microglia following LPS stimulation in

the absence or presence of Aβ40 (500nM). *, $P < 0.05$; ***, $P < 0.001$; n = 7-8 each group.

(**i**) IL-6 mRNA expression in control and *ric8a/cx3cr1-cre* mutant microglia following LPS stimulation in

the absence or presence of Aβ40 (200nM). *, $P < 0.05$; n = 6 each group.

1096

1097

1098

1099

1100

Supplemental Fig. 11. Suppression of astrogliosis in *ric8a-emx1-cre* mutant cortices by anti-

inflammatory drugs, dorsomorphin (DM) and S3I-201 (S3I).

(a-e) GFAP (in red) and nuclear (DAPI, in blue) staining of neonatal control (**E**) and mutant cortices

without treatment (**F**) or mutant cortices after dorsomorphin (DM, **G**), S3I-201 (S3I, **H**), or dual (DM+S3I,

I) treatment at E12.5. Note, GFAP is normally expressed in the neonatal hippocampus (HC) (dashed line

in (a)).

(f) Quantitative analysis of GFAP-positive astrocyte numbers in the neonatal mutant cortex after

treatment at E12.5.

1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

Supplemental Fig. 12. Suppression of MMP9 expression in *ric8a-emx1-cre* mutant cortices by anti-

inflammatory drugs, dorsomorphin (DM) and S3I-201 (S3I).

(**a**) MMP9 (in red) staining in control cortices at E13.5.

(**b**) MMP9 (in red) staining in mutant cortices at E13.5 after DM and S3I dual treatment at E12.5.

(**c**) Quantitative analysis of MMP9 expression. No significant differences are observed in mutants after

inhibitor treatment in comparison to controls (P = 0.44, n = 6).

(**d**) Gel zymography of E13.5 control and mutant cortical lysates following DM/S3I treatment at E12.5.

Similar levels of MMP9 are observed between controls and mutants. Quantification also showed no

significant differences in normalized MMP9 levels (P = 0.46, n = 4).

1125