1

Expanding salivary biomarker detection by creating a synthetic neuraminic acid sensor via chimeragenesis

Samuel J. Verzino[1], Sharona A. Priyev[1], Valeria A. Sánchez Estrada[1], Gemma X. Crowley[1], Alexandra Rutkowski[1], Amelia C. Lam[1], Elizabeth S. Nazginov[1], Paola Kotemelo[1], Agustina Bacelo[1,2], Desiree T. Sukhram[1], Frank X. Vázquez[2], Javier F. Juárez[1]

[1]Dept. of Biological Sciences and [2]Dept. of Chemistry, Saint John's University, 8000 Utopia Parkway, Queens, NY (USA)

ABSTRACT

Accurate and timely diagnosis of oral squamous cell carcinoma (OSCC) is crucial in preventing its progression to advanced stages with a poor prognosis. As such, the construction of sensors capable of detecting previously established disease biomarkers for the early and non-invasive diagnosis of this and many other conditions has enormous therapeutic potential. In this work, we apply synthetic biology techniques for the development of a whole-cell biosensor (WCB) that leverages the physiology of engineered bacteria *in vivo* to promote the expression of an observable effector upon detection of a soluble molecule. To this end, we have constructed a bacterial strain expressing a novel chimeric transcription factor (Sphnx) for the detection of N-acetylneuraminic acid (Neu5Ac), a salivary biomolecule correlated with the onset of OSCC. This WCB serves as the proof-of-concept of a platform that can eventually be applied to clinical screening panels for a multitude of oral and systemic medical conditions whose biomarkers are present in saliva.

INTRODUCTION

Biosensors are analytical devices in which biological elements convert a physicochemical stimulus into a detectable signal[1,2]. Among the many existing variants, whole-cell biosensors (WCBs) stand out due to their versatility and engineering potential[1]. These are detection systems in which a physiologically active cell sustains an array of biomolecular elements (receptors, transporters, enzymes, regulators, etc.) capable of signal recognition, integration, and production of an observable output[1]. Among the organisms that can serve as WCBs, there are bacterial species which either produce or limit the production of reporter proteins in the presence of a target analyte[3,4]. Microbial WCBs have demonstrated potential in fields such as environmental monitorization, food safety, and medicine[3,5]. Microorganisms can grow rapidly in relatively inexpensive media and produce all the necessary biological elements for analyte detection, providing a low-cost and portable alternative to other detection systems[6]. Unlike conventional sensors that may require large and complex instruments as well as laborious sample processing, WCBs can detect a bioavailable substance directly from the environment with minimal preparation, enabling their incorporation into *in situ* detection hardware that may revolutionize point-of-care diagnosis[6]. Multiplexing strategies, which enable detection of several signals simultaneously, combined with improved safety controls (i.e. biocontainment strategies) increase the opportunities for these biosensors to be deployed in both clinical and diagnostic settings[3,4]. Additionally, the robust and versatile nature of bacterial biosensors makes them suitable for use in environments where other technologies would not be able to operate, such as strenuous pH and osmotic conditions[3]. However, the use of WCBs is not exempt from certain drawbacks, such as slow reaction times, potentially impractical dynamic and detection ranges, and sensitivity and

selectivity issues. However, each year an increasingly large corpus of publications proposes

solutions to these problems, making biosensors more competitive and reliable[7].

WCBs benefit from the synergies resulting from the combination of native systems,

evolved during millions of years to detect the miscellany of molecules surrounding microbes, with

the ever growing toolkit that synthetic biology provides (such as the creation of engineered genetic

circuits[8–11]). In WCBs based on transcription factors (TFs) the presence of a soluble analyte

becomes associated with a conformational change of a protein, so that the binding state of the

polypeptide to DNA gets altered, modifying the transcription profile of a regulated reporter[12].

These biosensors benefit the most from the use of monogenic transcriptional repressors[8,9], control

systems encoded by a single gene whose product prevents transcription in the absence of a small

soluble inducer molecule. The catalog of characterized TFs capable of interacting with small

soluble molecules is expanding, yet still limited, which represents one of the main obstacles in the

field[13,14]. To expand the panoply of molecules that can be detected, we can employ synthetic

biology, whose technologies have enabled us to better understand protein modularity such that we

can better engage in regulator customization and *de novo* creation[15,16]. In this paper, we centered

our efforts on the detection of a salivary biomarker that, to the best of our knowledge, was not the

inducer / inhibitor of any characterized TF and thus required the creation of a synthetic regulator.

Saliva is a biological fluid resulting from the secretions of several glands (among them the

parotid, submandibular and sublingual), crevicular fluid from the gingiva, cell residues, microbes

and associated structures such as biofilms, plus secretions from the respiratory tract[17,18]. Human

saliva contains a vast array of metabolites informative of oral and systemic health[19], making its

frequent or continuous sampling by portable and wearable equipment extremely coveted.

Translation to dental practices of such sampling technologies may increase their role at the

forefront of early disease prevention and diagnosis[20]. Electrochemical intraoral biodevices capable

of detecting salivary biomarkers presently exist, yet they are limited in the scope of molecules they

detect and lack adaptability to sense new ones[21]. Biological detection systems based on engineered

genetic circuits offer a plug-and-play potential, in which the substitution of one TF by another,

plus its cognate operator, is often enough to repurpose a pre-existing sensor. Bioengineering

solutions enabling point-of-care diagnostics will greatly benefit patients by improving at-home

health monitorization[22]. This approach is especially relevant due to the widespread acceptance and

use of consumer technology within society, which has normalized wearable biosensors connected

to a phone or a smartwatch[23]. Diagnostic screenings that do not require invasive procedures (i.e. a

blood sample or a biopsy) are preferred by patients, which makes saliva a prime biological fluid

candidate to be sampled for health and disease markers[24,25]. In this work we focused our attention

on N-acetylneuraminic acid (neuraminic acid, Neu5Ac), the most common of the sialic acids

originally isolated from salivary glands[26–28]. Neu5Ac fulfills key roles in human physiology[29] as

well as in commensal and pathogenic bacteria[26]. Moreover, changes in salivary sialylation might

become a suitable early indicator of changes associated with malignant pathologies, given their

correlation to periodontal disease (PD) and oral cancer[30]. The oral microbiome of patients with PD

include bacteria such as *Tannerella forsythia*, which engage in glycan harvesting behavior: they

cleave Neu5Ac molecules from glycoproteins to stealthily evade their host's immune system[31].

Additionally, abnormally high concentrations of Neu5Ac have been associated with the onset of

oral cancer[32–34]: a group of neoplasms which affect any region in the oral cavity, pharynx, and

salivary glands[35]. Oral cancer is often used interchangeably with oral squamous cell carcinoma

(OSCC), which accounts for more than 90% of oral malignant epithelial neoplasms[35]. As one of

the most prevalent cancers in the world[36], untreated OSCC has very poor prognosis and, as such,

early detection is essential for successful treatment[37]. Previous evaluations of sialic acid levels in

the saliva of healthy individuals compared to patients with oral cancer provided a guide for the

range of salivary Neu5Ac that can be observed. The concentration range identified in the saliva of

healthy adults was approximately 21.65 ± 5.71 mg/dl (0.52 - 0.88 mM), while confirmed oral pre-

cancer individuals displayed 59.75 ± 7.29 mg/dl (1.7 - 2.2 mM). Histopathologically confirmed

OSCC patients showed a wide range of concentrations, 204.85 ± 60.38 mg/dl (4.6 - 8.6 mM),

always superior to the other two groups[38]. Additional studies have employed multivariate

regression analysis comparing both free and protein-bound salivary sialic acid concentrations

between OSCC patients and healthy individuals. While free sialic acid levels were found to be

significantly elevated in patients diagnosed with oral cancer, differences in protein-bound sialic

acid levels were not found to be statistically significant[39]. These observations are further evidence

that total salivary sialic acid concentration can be utilized as a biomarker for OSCC. It is important

to note that there is at least one study in which Neu5Ac concentrations in saliva were decreased in

OSCC patients, having been hypothesized that a downregulation of the pathways modulating

Neu5Ac production might have occurred in these subjects[40]. Beyond its connection to OSCC, it

has also been proposed that salivary Neu5Ac could interfere with SARS-Cov-2 (COVID-19)

infection, acting as a competitive inhibitor blocking the viral antigens recognized by the host

receptors[41]. Each of these factors made quantification of salivary Neu5Ac a desirable target for

detection by biosensors.

The absence of a native TF capable of detecting Neu5Ac motivated us to construct a

synthetic monogenic repressor for its integration in a genetic circuit that would enable the strain

bearing it to behave as a WCB. We favor these regulators due to their simplicity, compact nature,

and ease of integration into synthetic circuits[8,9,42]. The process of constructing our custom

Neu5Ac-sensing regulators is based on streamlined gene fusion / chimeragenesis methods

developed by our laboratory to detect soluble molecules[15] . Here, we describe a Neu5Ac TF-based

WCB that employs engineered bacterial machinery to fulfill the functions of the three basic

modules present in any detection system: a sensing module, a processing unit, and an actuator. The

living WCB can intake a Neu5Ac signal by expressing a novel synthetic transcription factor,

Sphnx, that bridges detection and processing (a common feature of bioreceptors[23]). A separate

reporter plasmid completes the system providing a gene encoding a fluorescent protein (sfGFP,

henceforth stylized as GFP) whose expression is regulated by the synthetic TF Sphnx so that

detectable changes in fluorescence occur upon the detection of soluble Neu5Ac. Sphnx is a

member of a new combinatorial library of TFs assembled via a simplified chimeragenesis process.

Important clues on inter-domain communication of modular proteins have started to be gathered

by comparing Sphnx to non-functional members of the TF collection. Protein tertiary structure

modeling of functional and non-functional regulators has contributed to establish comparisons

between them helping to elucidate intricate and frequently difficult to predict dynamics that may

condition the success or failure of any chimeric TF.

RESULTS AND DISCUSSION

Design and Construction of Synthetic Transcription Factors. The functionality of the WCBs

that we built in this work hinged on the obtention of a TF capable of binding Neu5Ac with efficacy

and precision. The Neu5Ac-induced monogenic transcriptional repressor we intended to assemble

included three modules: a N-terminus (N_t) DNA Binding Domain (DBD), a linker sequence

(LNK), and a C-terminus (C_t) Ligand Binding Domain (LBD). The DBD specifically recognizes

an operator sequence in the promoter it regulates, driving the expression of the reporter gene

(*sfgfp*). The LBD is tasked with the recognition of Neu5Ac, providing specificity to the TF. Finally,

the LNK sequence acts as a bridge, providing the right amount of flexibility/rigidity necessary for

the transfer of information between DBD and LBD: steric changes in the LBD associated with

Neu5Ac binding get translated into steric changes in the DBD effecting a change of its oligomeric

state that results in the unbinding of the TF from its operator and thus in a de-repression of

transcription. By using a gene fusion / chimeragenesis approach we generated a library of

repressors with a common N_t-DBD-LNK-LBD-C_t architecture.

Our selected DBD was sourced from the paradigm of transcriptional regulators: the *lac*

repressor or LacI[43]. Though naturally induced by lactose/IPTG, we exclusively employ the DBD

of LacI recognizing the operator *lacO* present in the P_{lac} promoter (as defined in[15]). We decided to

use DBD-LacI because the *lac* repressor originated as a modular protein with two domains, and

has been previously used in the construction of several functional chimeras[44–46]. It is important to

remark that the exclusive use of DBD-LacI, as opposed to the full protein, ensures that chimeric

proteins carrying this domain will recognize their target operator, yet makes them incapable of

responding to native inducers.

Candidate LBD selection was accomplished through bibliographical research. Since the

LBD was the domain that recognized the molecule of interest for which we built the biosensing

strain, it could be sourced from any whole protein or protein domain as opposed to being co-opted

from a previously documented regulator. One of our main sources of the LBDs we employ in our

chimeragenesis pipeline are periplasmic binding proteins (PBPs)[47]. These are standalone subunits

of transporters that act as molecular shuttles that assist in the traffic of small solutes from the outer

to the inner membrane in Gram negative bacteria[48]. PBPs share a common ancestor with the LBD

domains of the LacI / GalR family of regulators[45]. To build the synthetic TF we required for this

project, we needed to identify LBD candidates that strongly bound our soluble molecule of interest

(Neu5Ac). PBPs SatA and SiaP were selected for their binding affinity and specificity for the

transport of Neu5Ac. SatA is a product of the *satABCD* (*s*ialic *a*cid *t*ransport) operon that encodes

an ATP Binding Cassette (ABC) transporter[49] from *Haemophilus ducreyi*[50]. An alternative

satABCD system present in *Corynebacterium glutamicum* has been demonstrated to be essential

for its growth using Neu5Ac as sole carbon source[51]. SiaP is a product of the *siaPQM* operon,

which encodes a Tripartite ATP-independent periplasmic (TRAP) transporter in *Haemophilus*

influenzae[52,53]. SiaP was the first TRAP-associated PBP whose structure was elucidated (in *Vibrio*

cholerae, *Pasteurella multocida* and *Fusobacterium nucleatum*[52,54]). Structural comparisons and

thermodynamic studies of SatA and SiaP suggest that similar affinities for Neu5Ac are achieved

in the two PBPs through distinct mechanisms: one enthalpically driven and the other entropically

driven. SiaP, as well as other TRAP transporters, are hypothesized to command an enthalpically

driven ligand binding strategy due to the abundance of charged residues (Arg, Lys, Asp, and Glu)

at their binding pocket. SatA, alternatively, is hypothesized to bind its substrate in what is

considered an entropic approach because of the numerous hydrophobic and polar residues (Tyr,

Phe, Gly, Leu, Asn, and Ser) at its binding pocket[55]. The structure of *H. influenzae* SiaP in the

presence of Neu5Ac (PDB 3B50) reveals the ligand bound in a deep cavity with its carboxyl group

forming a salt bridge with a highly conserved Arg residue[52]. Initial studies of SiaP isolated from

H. influenzae displayed a binding K_d of ≈120 nM when affinity for Neu5Ac was tested[53].

Subsequent studies established a value closer to K_d ≈58 nM[52]. Similar binding affinity of SatA

complexes has been identified with a K_d ≈133 nM[55]. The collection of empirical data supporting

the ability of SatA and SiaP to bind Neu5Ac made them ideal candidates to become the LBD of a

chimeric TF responsive to Neu5Ac.

The final piece of the chimeric gene puzzle consisted on establishing a connection between

DBD and LBD domains. This task was performed by elements called linker (LNK) regions that

tend to be less structured than the domains they connect[56]. In order to maximize the chances of

transmitting the information of Neu5Ac binding to the DBD, so that it can detach from *lacO* and

enable transcription, we decided to connect DBD and LBD via two different strategies: i) a direct

linkage without a polypeptide linker (condition identify as LNK1 for uniformity purposes), or ii),

by one out of four linkers characterized by having different biophysical properties (identified as

LNK2 to LNK5; *Supp. Table 1*). It is important to note that while LNK1 does not contain a linker

per se, the LBDs we chose, by virtue of being PBPs, are proteins that would be exported to the

periplasm of the cell under natural circumstances, and thus contain a native signal peptide at their

N-terminus. This kind of peptide appears to be unstructured for the most part, barring a short alpha-

helix region[57]. We decided to include the signal peptides of both SiaP and SatA due to existent

examples of functional chimeras containing similar structures, such as the glucose-inducible

SLCP$_{GL}$[46] and the benzoate-inducible CbnR-ABE44898-OD[15] TFs. At this stage, we had identified

all the elements we needed to start assembling our new chimeric TFs.

dsDNA fragments encoding the DBD and all LNK-LBD variants were obtained as detailed

in the *Materials and Methods* section. To initiate the construction of a library containing all the

desired chimeric TFs, we performed several scarless assembly reactions with all the domain-

encoding fragments and linearized expression vector pUC19RBS, followed by transformation in

NEB5-alpha competent cells. Resulting clones were isolated and individually sequenced. This

approach was performed twice. Chimeric combinations that were not present in the libraries were

re-attempted to be individually cloned five times with no success, at which point we proceeded to

put pause to their construction and move the project forward, subcloning the TF collection we had

generated up to that point in time into the low-copy vector pCKTRBS, what generated the set of

plasmids we collectively identified as pCKT-*Chimera*. We hypothesized that chimeras

unsuccessfully cloned at this stage may be potentially deleterious for the cells. It is important to

note that the expression of any heterologous protein is *per se* taxing on bacterial metabolism, and

that TFs might be especially burdensome[58,59]. Chimeric constructions including DBD-LacI and

SiaP/SatA seem to provoke a certain toxicity to the cell, albeit to different degrees depending on

the specific domain combination. Indirect evidence supporting this hypothesis was observed when

constructions expressing the chimeric TFs were transformed into *E. coli* MC4100 to assess their

behavior in M63 minimal media with glycerol as the sole carbon source. Under these resource-

limited conditions, stressed bacteria have more difficulties in thriving, and MC4100 cells bearing

pCKT-*Chimera* vectors displayed a very modest and erratic growth consistent with our

assumptions (data not shown). *Table 1* contains the chimeric TFs built and assayed presented in

this study, as well as a simplified interpretation of their functionality when grown in rich media.

Screening of transcription factors. A functional *in vivo* screening was implemented using GFP

as reporter. Plasmid pHC_DYOLacI-R (*Supp. Table 2*) was transformed into candidate strains

expressing the chimeric TFs from pCKTRBS vectors under the control of TetR-aTc. These two

plasmids expressed in the same strain constituted a genetic circuit (*Figure 1a*) that enabled a

controlled expression of the chimera as well as its controlled induction. This circuit behaves as a

B-or-not-A universal two input gate[60] (*Figure 1b*), where there are two inputs (A, aTc

supplementation; B, Neu5Ac supplementation) and one output (detection of GFP fluorescence).

Based on the addition of the inputs we can define three states in the genetic circuit (*Figure 1c*):

Basal (aTc$^-$/Neu5Ac$^-$), no input added, GFP is expressed from its promoter in pHC_DYOLacI-R;

Repression (aTc$^+$/Neu5Ac$^-$), the presence of aTc induces TetR which is incapable of remaining

attached to P_{tetO} and thus the chimera is expressed from its corresponding pCKT-*Chimera* vector

repressing the expression of GFP; *De-repression* (aTc$^+$/Neu5Ac$^+$), the chimera is expressed but if

it is functional it is induced by Neu5Ac, de-repressing the promoter driving the expression of GFP.

By comparing the behavior of the strain expressing the aforementioned circuit under these

conditions, we can rule out non-functional synthetic TFs. In the case that the chimeric protein was

not a functional repressor, relative GFP-associated fluorescence (Rel.fluor) could be detected in

the *Repression* state (ΔRel.fluor$_{basal-repressed}$ tending to 0). On the other hand, if the chimera was

able to bind to *lacO* on the reporter promoter yet incapable to recognize Neu5Ac with its LBD, or

to transmit a LBD to DBD conformational change resulting in its detachment from the operator,

we would have obtained a functional super-repressor[61] unable to respond to its inducer and, thus,

not a valid tool to detect Neu5Ac (ΔRel.fluor$_{(de-repressed)-repressed}$ tending to 0). *Figure 1d* illustrates

the expected GFP production along a growth curve for a strain expressing the screening genetic

circuit that included an ideal chimera (ΔRel.fluor$_{(de-repressed)-repressed}$ >0). It is important to note that

even though Neu5Ac is a monosaccharide, its supplementation to rich growth media did not

significantly increase the OD$_{600}$ of the assayed strains, at least up to the tested 10 mM concentration

(*Supp. Figure 1*). Regardless of this observation, GFP-associated fluorescence data displayed in

this publication is always expressed as relative fluorescence (arbitrary units) corrected by the

OD$_{600}$ of the culture, in order to account for growth disparities among different strains and

conditions. It is also important to note that repression by LacI, or LacI-derived chimeras, is rarely

absolute so that a certain level of leakage, and thus of background reporter gene expression, is

always expected[46,62].

All the chimeric TFs included in *Table 1* were screened for their ability to bind DNA and

thus to repress the expression of the GFP reporter, as well as to be induced by Neu5Ac enabling

production of the fluorescent protein. All members of the collection of MG1655 (pCKT-*Chimera*,

pHC_DYOLacI-R) strains bearing the genetic circuit described in *Figure 1* were tested *in vivo* to

assess their ability to behave as Neu5Ac biosensors (*Supp. Table 4*). Among the chimeric proteins

expressed by this strain collection, we included the TF Lumos, which did not contain any of the

expected LNK1-5 options but an unexpected 6 amino acid linker (LNK6) resulting from an

abnormal assembly reaction in which the last 18 bases of *DBD-lacI* were duplicated (*Supp. Table*

1). Given the stable nature of the amino acid string, as well as our agnostic approach to which

LNK sequence could be the best performing one, we decided to maintain this construction in our

pool. TFs Kunst (LacI-LNK3-SiaP), Scorpio (LacI-LNK4-SiaP), Lumos (LacI-LNK6-SiaP), and

Haus (LacI-LNK2-SatA) showed no repression abilities. These TFs seem to not be able to stably

bind to their operator boxes in the reporter plasmid, potentially due to an inability to acquire their

dimeric form. The strain bearing Haus also presented a diminished growth, which strongly

suggests the potential toxicity of this transcription factor for the host cell.

Bastet (LacI-LNK5-SiaP) was intriguing due to the possibility that it was able to repress

and de-repress for the first 10 hours of growth, yet its behavior requires a deeper study given the

observed dispersion in the analyzed data. This observation suggests that the strain might be

unstable due to the potential toxicity of the protein. Also of note, Harpy (LacI-LNK1-SatA)

seemed to be able to repress at a certain level and potentially de-repress after the 18 hour mark.

Another TF that presented this similar and unexpected induction profile but in more consistent

basis was Siren (LacI-LNK1-SiaP), which displayed an intriguing function as a proper repressor

inducible by Neu5Ac but only in the stationary phase of growth (*Supp. Figure 2*). In the future we

intend to study this protein in depth, since its repression/de-repression profile is very attractive for

a whole-cell biosensor setup in which a bacterial culture sustaining itself at stationary phase is

continuously monitoring an input for an extended period of time. Service of WCBs over time is a

known challenge associated with these technologies, and engineering of genetic circuits

incorporating regulators such as Siren, is one of the strategies that are being explored to account

for this problem[3]. A candidate biosensor strain expressing Sphnx (LacI-LNK2-SiaP) exhibited the

most suitable repression and induction profile compatible with that of a TF capable of repressing

P_{lac} while de-repressing it in a detectable manner upon binding to Neu5Ac. To summarize this

initial evaluation of the functionality of our candidate TF-based WCBs, and judging by the

difficulty to obtain them, their lack of functionality, and the reduced sample size, SatA-based

fusions seem to be less stable and more prone to not be responsive to Neu5Ac. Further studies and

the creation of expanded chimeric libraries, however, are required to confirm this hypothesis. At

this injunction, we focused our efforts on better characterization of Sphnx, since it was the most

promising TF we had assembled (estimated as the relative fluorescence difference under de-

repressed and repressed conditions along the growth curve or (ΔRel.fluor$_{(de-repressed)-repressed}$)

(*Supp.Table 4*).

In order to validate our preliminary observations, we systematically compared GFP

production over time of MG1655 (pCKT-Sphnx, pHC_DYOLacI-R) cells expressing Sphnx either

in the presence or absence of Neu5Ac. *Figure 2a* displays the results of time course experiments

over a period of 25 hours. GFP production induced by the presence of Neu5Ac was observed from

the first hour of growth, with the Neu5Ac-induced cultures showing a tendency to display higher

fluorescence than the repressed ones. At this point our attention was captured by Kunst as a model

for a non-functional chimera in opposition to Sphnx. *Supp. Figure 3* shows the behavior of Kunst,

which exhibited a tendency to repress P_{lac} at the very beginning of a 40 h time course, yet after the

3 h mark did not show any difference between induced and uninduced conditions. A preliminary

evaluation of the structural differences between Sphnx and Kunst is included in its own section of

Results and Discussion.

In an effort to validate the *in vivo* observations confirming Sphnx functionality, we

analyzed the induction of P_{lac} driving the expression of GFP in MG1655 (pCKT-Sphnx,

pHC_DYOLacI-R) by qRT-PCR experiments (*Supp. Figure 4)* performed as indicated in

Materials and Methods. The expression levels of *sfgfp* calculated as the ratio of the induced

condition over the repressed condition were 1.58 ± 0.24 (*n* = 8, error expressed as SEM),

statistically significant and consistent with an induction of the expression in the presence of

Neu5Ac.

Further *in vitro* experiments were also performed to confirm the interaction between P_{lac}

and the DBD-LacI of Sphnx. Electrophoretic mobility shift assays (EMSA; *Materials and*

Methods) showed that cell extracts from NEB5-alpha (pCKT-Sphnx) grown for 5 h were capable

of retarding the migration of a P_{lac}-*sfgfp* probe compared to control extracts that were not

expressing Sphnx (data not shown). Neu5Ac behaved as the inducer of Sphnx, since binding of

the protein to the probe was diminished in a concentration-dependent manner in the presence of

the sialic acid (*Figure 2b*). On the other hand, cell extracts from NEB5-alpha (pCKT-Siren) grown

for 5 h were able to bind the probe but not to unbind in the presence of Neu5Ac, confirming our

in vivo observations. Previous publications report a similar behavior: DBDs from bacterial

transcriptional repressors tend to retain their ability to bind to their operator sequences even when

integrated in a new chimeric TF[63–65], another testimony to the high modularity of these proteins.

Our initial *in vitro* DNA-protein interaction studies require future validation with purified chimeras

in order to calculate the kinetics of the Sphnx-*lacO* and Sphnx-Neu5Ac-*lacO* interactions, yet they

represent a promising step forward towards characterizing the interaction between our synthetic

TF and its cognate operator.

At this juncture of the project, both *in vivo* and *in vitro* experiments strongly suggested that

Sphnx was a transcriptional repressor capable of interacting with *lacO* and being de-repressed by

neuraminic acid. To better understand the dynamics of the Sphnx-Neu5Ac interaction we analyzed

the production of GFP by MG1655 (pCKT-Sphnx, pHC_DYOLacI-R) cultures at 14 hours of

growth (*Materials and Methods*) when exposed to different concentrations of neuraminic acid

around the average salivary concentration for an adult male ($\approx$1 mM)[66]. The mean fold difference

of GFP expression between the repressed and induced conditions was 2.25 ± 0.74 for cultures

induced with 0.5 mM Neu5Ac, 2.54 ± 0.52 for cultures induced with 1.0 mM Neu5Ac, 3.81 ± 1.15

for cultures induced with 2.0 mM Neu5Ac, and 5.21 ± 2.00 for cultures induced with 5.0 mM

Neu5Ac ($n = 8$, error expressed as SEM). These results were commensurable with those observed

for modular chimeric TFs in the presence by their ligands: $SLCP_{GL}$ (LacI-GGBP-OD) induced by

glucose[46], CbnR-ABE44898-OD and LmrR-BzdB1_nSP induced by benzoate[15]. *Figure 2c* shows

the levels of GFP fluorescence observed in basal and repressed cultures as well as in cultures

induced by 0.5, 1.0, 2.0 and 5.0 mM neuraminic acid. These assays corroborated that relative

fluorescence between basal (aTc$^-$/Neu5Ac$^-$) and repressed (aTc$^+$/Neu5Ac$^-$) conditions were

significant. Cultures in which the genetic circuit used to test Sphnx were de-repressed

(aTc$^+$/Neu5Ac$^+$) displayed statistically significant values compared to the repressed cultures when

in the presence of 1.0 mM and 2.0 mM Neu5Ac. Induction by 0.5 and 5.0 mM was not significant.

It is important to note that relative fluorescence values in the 5.0 mM-induced biological replicas

were widely dispersed, suggesting that an overabundance of Neu5Ac might interfere with the

stability of the host strain bearing both plasmids, producing some deleterious effect associated to

the expression of the heterologous Sphnx and GFP proteins in the assayed conditions, which in

practical means may entail the testing of a few sample dilutions when using these biosensors for

out-of-the-lab applications. These results showing Sphnx induction are promising, yet they

underline the need to refine chimeric TFs resulting from the assembly of pre-existing modules to

optimize their induction range as well as gene dose and reporter genes used to study their behaviour

in vivo and *in vitro*, as it has been previously proposed[14]. Newly created synthetic TFs tend to have

limited transfer function and specificity, and thus to require further engineering for their field

deployment in biosensors[67]. In this regard, we are optimistic about the future improvement of the

chimeras we have assembled: the estimated K_d of SiaP and SatA for neuraminic acid are in the nM

range[52,53,55] (calculated *in vitro*), whereas in this work we present data in the mM range (calculated

in vivo). This suggests that we have not reached the full potential of our LBDs and there is an

opportunity to optimize Sphnx's structure to better benefit from the high specificity and affinity

of our LBD. It is also important to note that Sphnx is currently at the same point in evolutionary

history when DBD-LacI was fused to the ancestral version of its current LBD, long before

communication between DBD and LBD had been refined and perfected by natural selection[45]. A

systematic exploration of promoter and RBS variants can provide a better solution to compensate

for some of these deficiencies[14]. Regarding the use of alternative reporters that may shed more

light on the induction profile of Sphnx, we acknowledge that the use of GFP has some inherent

limitations in prokaryotes (e.g. widely varying maturation times dependent on strain and metabolic

state[68]), which has motivated us to initiate the development of better reporting systems.

Modeling of the 3D structure of chimeric transcription factors. Another element we are

starting to explore is the importance of linkers for the functionality of modular proteins, an often-

neglected design element[69]. Since the beginning of our journey characterizing the chimeric TFs,

we found extremely intriguing the fact that Sphnx (LacI-LNK2-SiaP, N_t-DBD-EKEKEK-LBD-

C_t) and Kunst (LacI-LNK3-SiaP, N_t-DBD-GSGSGS-LBD-C_t), which are nearly identical proteins

(*Supp. Materials Appendix*), possessed opposed behaviours when interacting with neuraminic

acid. The recent release of the AlphaFold 3 software[70] has enabled us to predict the structures of

multimeric chimeric proteins in the presence of their dsDNA operator, allowing us to model the

proteins under conditions closer to those found in actual WCBs. Sphnx and Kunst were both

modeled as dimers using AlphaFold 3, along with a DNA double helix containing the *lacO* (5'-

ATTGTGAGCGGATAACAATT-3' and its complementary sequence) recognized by the DBD-

LacI they both carry. Our goal was to predict structures that could help explain why Sphnx binds

Neu5Ac while Kunst does not, despite only differing by six amino acids. *Figure 3a* shows the

predicted Kunst DBD (green) bound to the DNA double helix (gray) and the formation of a dimer

between the PBP (cyan). The structure shows a mostly disordered linking region (orange),

including the GSGSGS (LNK3) linker (violet). The predicted Sphnx model (*Figure 3b*), on the

other hand, shows that the connecting region adopts a helical structure, including the EKEKEK

(LNK2) linker, and appears to form contacts between both the DBD and PBP. Rotating the Kunst

structure by 90° (*Figure 3c*) we can observe that the GSGSGS (linker does not interact with the

protein or DNA. In contrast, rotating the Sphnx structure by 90° (*Figure 3d*) further shows that the

linker forms contacts between the DBD and PBP domains, with interactions occurring across

different monomer chains. Further analysis shows that the helical linker in Sphnx forms

electrostatic salt bridges between itself and both the DBD and PBP domains. *Figure 3e* shows the

acidic residues on the linker (Glu) forming contacts with basic residues (Lys) on the DBD and

PBP domains. Specifically, the predicted structure suggests that the residues expected to form salt

bridges include Glu67-Lys2 on the DBD and Glu69-Lys152 on the PBP. These salt bridges likely

stabilize the helical linker, affecting the flexibility and stability of the polypeptide, an effect that

has been previously observed in other proteins[71]. From these predicted structures, we hypothesize

that these salt bridges stabilize helicity in the flexible linker in a way that positively affects the

PBP function by interacting with the DBD and the PBP, explaining the ligand binding functionality

of Sphnx compared to non-binding Kunst protein. Zooming out to see the larger structure (*Figure*

3f), aromatic residues near the linker salt bridges are visible. These residues are key for testing the

new predicted structures and designing new chimeric proteins in the future. We intend to purify

Sphnx and Kunst to perform fluorescence analysis of the protein-ligand interactions, including

fluorescence quenching and fluorescence anisotropy studies. The construction of 3D models, using

the latest version of AlphaFold, will continue to enable us to explore which mutations might

enhance ligand binding in a way that was previously unavailable, also allowing us to model

conditions closer to the actual multimeric *in vivo* conditions.

<u>CONCLUSIONS</u>

This work presents MG1655 (pCKT-Sphnx, pHC_DYOLacI-R), the first iteration of a

transcription factor-based whole-cell biosensor strain capable of detecting the salivary biomarker

N-acetylneuraminic acid and relaying its presence through the production of a fluorescent reporter,

the traditional litmus test for a biosensor circuit[14]. The keystone of the genetic circuit enabling the

detection of the soluble molecule is a new transcriptional repressor: Sphnx. The apparent absence

of a native Neu5Ac-binding transcription factor led the authors to create a custom-made regulator

via a straightforward synthetic biology approach that enables the generation of chimeras capable

of repressing a known promoter and being de-repressed by a molecule of interest. Neuraminic acid

is an attractive analyte to monitor since its enhanced levels in saliva are correlated with conditions

that range from periodontal disease[66] (oral microbes harvest sialic acid to evade our immune

system[31]) to the onset of oral squamous cell carcinoma[32–34]. To the best of our knowledge, Sphnx

is the first published transcriptional repressor capable of being induced by neuraminic acid.

This study supports the idea that the ability to construct custom-made TFs, affordably and

on-demand, is instrumental in expanding the panoply of small soluble molecules that can be

detected by biosensors in a field limited by the availability of sensing modules[14,45]. It is important

to note that synthetic TFs such as Sphnx do not represent the closing of a chapter, but the origin

of a new source of regulators whose performance can be improved via directed engineering of

their ligand binding pockets[72,73], an approach that can now benefit from high-throughput

approaches[74]. It is widely acknowledged that biosensor specificity and biosensor-analyte reaction

curves need to be optimized for the practical application of the WCB[67]. Once the biosensor strain

is optimized we will test it first on synthetic saliva preparations[75] with a gradient of Neu5Ac

concentrations and, eventually, with clinical salivary samples. The MG1655 (pCKT-Sphnx,

pHC_DYOLacI-R) strain represents the initial step in the obtention of a functional WCB for the

detection of neuraminic acid, a necessary launchpad to achieve the goal of eventual clinical

deployment of the sensor.

The modular assembly of fusion genes such as Sphnx sheds light on the importance of

inter-domain communication for multi-domain protein functionality[76] and exemplifies how a

reduced number of independent modules can be the playground of natural selection to synergically

originate new functionalities in the cell[77]. The results presented in this work have challenged our

expectations regarding chimeric transcription factors: while we expected most of them to be able

to repress expression from their regulated promoter, we found several instances in which the chimeric product was unable to bind to its cognate operator boxes. Our current hypothesis is that different linkage alternatives condition the 3D structure of the multimodular protein to such an extent that the dimerization step necessary to bind the operator boxes cannot be achieved. We are looking forward to expanding these observations in the future. Our study presents the interesting contrast between the inducibility of Sphnx as opposed to Kunst, and particularly the unexpected role that a potential LBD-mediated stabilization of the linker may play in the de-repressibility of the TF dimer. *In vitro* assays using purified proteins, as well as further modeling simulation analysis and, eventually, the crystallization of these chimeras will contribute to illuminate the complex dynamics necessary for modular proteins to fulfill their function.

In summary, this work presents a new biosensor for N-acetylneuraminic acid, a salivary biomarker relevant to oral health, which is based on a synthetic TF whose construction adds to our understanding on the role that different domains play in the functionality of modular proteins.

MATERIALS AND METHODS

464

465 **Bacterial strains and growth conditions**. Bacterial strains and plasmids used are listed in *Supp.*

466 *Table 2*. *E. coli* strains were grown at 37°C in LB (Lysogenic Broth[78]) medium unless otherwise

467 indicated. When required, *E. coli* cells were grown in M63 minimal medium[79] using the necessary

468 nutritional supplements and 30 mM glycerol (Sigma, St. Louis, MO) as a carbon source.

469 Antibiotics were added at the following concentrations: 100 μg·ml^{-1} ampicillin/carbenicillin and

470 25 μg·ml^{-1} chloramphenicol (Sigma, St. Louis, MO). For protein expression from pCKTRBS-

471 derived vectors, cultures were induced with 0.5 μg·ml^{-1} anhydrotetracycline (aTc) (Clontech,

472 Mountain View, CA). When experimental conditions required it, cultures were supplemented with

473 N-acetylneuraminic acid (neuraminic acid, Neu5Ac; Sigma, St. Louis, MO) at a 1 mM

474 concentration unless otherwise indicated.

475 Culture growth was routinely monitored in a BioPhotometer D30 (BioSpectrometer Basic Kinetic

476 and Fluorescence; Eppendorf, Hamburg, Germany). When it was required, bacterial growth and

477 GFP production were recorded in a Varioskan LUX Multimode Microplate Reader (Thermo

478 Fisher, Waltham, MA). Cultures were pipetted on a 96-well plate (Flat bottom Polystyrene Black

479 with clear bottom; Corning, Corning, NY) covered with a Breathe-Easy BEM-1 gas-permeable

480 membrane (Diversified Biotech, Dedham, MA) and incubated in a plate reader, where OD$_{600}$ and

481 fluorescence in the range of GFP (excitation 485 nm, emission 528 nm) were either tracked along

482 time courses (up to 40 h) or subjected to point measurements. NEB5-alpha (pHC_DYOLacI-R)

483 and MG1655 were included in every plate as GFP-positive and GFP-negative reference strains,

484 respectively. For standard time course experiments, overnight precultures of the strains were used

485 to inoculate (starting OD$_{600}$ 0.025) fresh growth media (supplemented with aTc and Neu5Ac when

486 indicated) in 96-well plates (180 μl total volume per well). For single point measurements,

overnight precultures of the strains (plus aTc and Neu5Ac when required) were used to inoculate

5 ml of fresh media supplemented in the same manner as the precultures. After 14 h of growth,

180 µl samples of the cultures were transferred to 96-well plates to enable OD_{600} and fluorescence

readings.

Statistical analysis as well as data visualization were performed with R (https://www.r-

project.org/) in the RStudio (RStudio Inc., Boston, MA) and Visual Studio Code environments

(Microsoft Corp., Redmond, WA).

Molecular biology techniques. Molecular biology techniques were performed following

commonly used standard protocols and as per manufacturers' instructions[80]. PCR reactions took

place in a 6321 Mastercycler PRO Vapo Protect Thermal Cycler (Eppendorf, Hamburg, Germany).

Routine separation of DNA and RNA in agarose gels was performed in an RunOne Electrophoresis

system (EmbiTec, San Diego, CA). Agarose gels were imaged in an UVP UVSolo Touch (Analytic

Jena, Jena, Germany). Plasmid DNA was purified with a Qiaprep Spin Miniprep Kit (Qiagen,

Hilden, Germany). DNA fragments were purified with DNA Clean-up and Concentration Kit

(Zymo research, Irvine, CA). The oligonucleotides employed for PCR amplification of the cloned

fragments and other molecular biology techniques are summarized in *Supp. Table 3* and were

supplied by IDT (Coralville, IA). All cloned inserts and DNA fragments were confirmed either via

Sanger sequencing[81] performed by Azenta Inc. (Burlington, MA), or Oxford-Nanopore whole-

plasmid sequencing performed by Plasmidsaurus (Eugene, OR). Commercially available NEB5-

alpha chemically competent cells (NEB, Ipswich, MA) were used for routine transformations.

Alternatively, electrocompetent *E. coli* cells were generated and transformed immediately (Gene

Pulser; Bio-Rad, Hercules, CA)[80]. Nucleotide sequence analyses were done at the National Center

for Biotechnology Information server (www.ncbi.nlm.nih.gov) and Benchling Biology Software

(www.benchling.com). Cloning was routinely performed via Gibson assembly[82] unless otherwise

indicated.

Domain selection and cloning. DBD-LacI is located on *lacI* 5' region, from the start codon to

base 198, encoding a peptide of 66 amino acids that comprises the N-terminus of our chimeric

proteins. This domain was obtained via PCR amplification from plasmid pCKTRBS-LacIwt[15]

using the oligonucleotide pair SV00001/SV00002. SatA from *H. ducreyi* (Uniprot Q7VL18) and

SiaP from *H. influenzae* (Uniprot P44542) were ordered as synthetic dsDNA fragments (IDT,

Coralville, IA). Several variants of the LBDs SatA and SiaP were amplified via PCR so that they

would include 5' overhangs including linkers (LNK2-5) or no linker whatsoever (LNK1). SatA

was amplified with primers SV00030-SV00034 / SV00004 and SiaP with primers SV00015-

SV00019 / JFJ0018.

Genes encoding the chimeric TFs were originally assembled in pUC19[83]. The vector was amplified

via divergent PCR with primers JFJ00019/JFJ00020, which introduce a consensus RBS for *E. coli*,

originating plasmid pUC19RBS. Tripartite Gibson assemblies including linear pUC19RBS, DBD-

LacI, and LNK-LBD were performed and the resulting constructions transformed into NEB5-alpha

competent cells. After a carbenicillin selection in solid media, a subset of the resulting clones was

isolated and their plasmids (pUC19-*Chimera* family, *Supp. Table 2*) purified and sequenced to

validate the presence of chimeric TFs. The resulting TF genes cloned into pUC19RBS were

amplified using SV00024/SV00025 for SiaP-derived chimeras and SV00024/SV00026 for their

SatA counterparts. These primers amplify our TF genes from the 5' DBD-LacI, just downstream

of the consensus RBS site, to the stop codon of the LBD. Low copy plasmid pCKTRBS[15] was

amplified via divergent PCR using primer pairs SV00022/SV00021 for SiaP-derived chimeras and

SV00022/SV00023 for their SatA counterparts. These primers introduce 5' and 3' overhang

sequences providing homology to our dsDNA chimeric gene fragments, and including in their 5'

side our consensus RBS site. Bipartite Gibson assemblies including linear pCKTRBS and the

fusion genes encoding the chimeras were performed, and the resulting assembly products

transformed into NEB5-alpha competent cells. After a carbenicillin selection in solid media,

resulting clones were isolated and their plasmids sequenced to confirm the presence of chimera-

encoding genes. Resulting pCKT-*Chimera* plasmids were purified from NEB5-alpha (pCKT-

Chimera) strains and electroporated into MG1655 cells. After confirming the presence of the

plasmid in the new host, MG1655 (pCKT-*Chimera*) were electroporated with the reporter plasmid

pHC_DYOLacI-R. Validated MG1655 (pCKT-*Chimera*, pHC_DYOLacI-R) strains were used for

in vivo and *in vitro* analysis of the functionality of the novel chimeric TFs when required. The

sequences of all the oligonucleotides used in this work can be found in *Supp. Table 3*.

Electrophoretic mobility shift assay (EMSA). In order to obtain cell extracts enriched in the

Sphnx/Siren proteins, NEB5-alpha (pCKT-Sphnx) and NEB5-alpha (pCKT-Siren) cells were

grown on LB supplemented with aTc for 5 hours at 37˚C in a shaker incubator. Cells were

recovered by centrifugation at 4°C (15' at 4,000 g) and then resuspended in 20 mM Tris-HCl (pH

7.0) to be sonicated at 70 amp for 30 seconds (30 cycles of 1 second; Branson Digital Sonifier,

Danbury, CN). A subsequent centrifugation at 4°C (20' at 20,000 g) was performed to eliminate

cell debris from the sample. The recovered supernatant was then incubated with the DNA probe,

a 251 bp dsDNA fragment amplified from plasmid pHC_DYOLacI-R with primers

SV00028/SV00029, containing P_{lac} (including *lacO*). Cell extract (5 µl) and DNA probe (5 µl)

were incubated together with reaction buffer (20% glycerol, 100 mM KCl, 500 µg·ml^{-1} bovine

serum albumin, 20mM Tris-HCl pH 7.5). The reaction mixture is incubated for 30' at 30 ˚C. After

incubation of the retardation mixtures, 2 µl of Gel loading dye without SDS (NEB, Ipswich, MA)

were added to each reaction. Mixtures were subsequently fractionated by electrophoresis in precast

6% polyacrylamide gels (Novex TBE Gels; Invitrogen, Waltham, MA) buffered with 0.5X TBE

(45 mM Tris borate, 1mM EDTA). Gels were run at constant voltage (100 V) for 90' in a Mini

Gel Tank (Invitrogen, Waltham, MA) and afterwards stained for 1 h in 50 ml of 0.5X TBE buffer

supplemented with 5 µl of SYBR Gold dye (Invitrogen, Waltham, MA) while gently shaken. Post

staining, gels were imaged in a UVP UVSolo Touch (Analytic Jena, Jena, Germany).

qRT-PCR analysis. Quantitative Real Time Polymerase Chain Reaction (qRT-PCR) was used to

assay the transcription of our reporter gene (*sfgfp*). MG1655 (pCKT-*Chimera*, pHCDYOLacI-R)

strains were precultured overnight in of 3 ml LB supplemented with the corresponding selection

antibiotics. On the next day, 100 µl of overnight preculture were used as inoculum for several

cultures, 10 ml each, of fresh LB (plus selection antibiotics). Each one of the cultures was

supplemented so that it would represent one of these three different conditions: *Basal* (aTc$^-$,

Neu5Ac$^-$), *Repressed* (aTc$^+$,Neu5Ac$^-$), and *De-repressed* (aTc$^+$, Neu5Ac$^+$). Where experimental

conditions required it, cultures were supplemented with 0.5 µg·ml^{-1} anhydrotetracycline (aTc)

(Clontech, Mountain View, CA) and 1 mM neuraminic acid (Neu5Ac; Sigma, St. Louis, MO). For

every independent qRT-PCR experiment, three biological replicates were created across each of

these three conditions. These cultures were placed in a shaker incubator at 37˚C for 5 hours and

pelleted at 4˚C and 4,000 g. Pellets were either frozen at -80˚C for future use or immediately

processed for total RNA extraction. Cells from each individual pellet were set for lysis by

resuspension in 1 ml of IBI reagent, a phenol and guanidine isothiocyanate mixture (IBI Scientific,

Dubuque, IA). 200 µl of chloroform (Fisher Scientific, Pittsburgh, PA) was added to the lysate

and the mixture was vigorously shaken to homogeneity, followed by centrifugation (benchtop

minifuge) at 4˚C and 14,500 rpm for 15 minutes that led to a phase separation. 500 µl of the upper

aqueous phase was mixed with an equal volume of 100% ethanol. At this point, total RNA was

purified from DNA/RNA phase with Direct-zol RNA Miniprep Kit (Zymo Research, Irvine, CA)

where the TRIzol reagent steps are replaced with the phenol and chloroform phase separation.

Total RNA eluted with ddH$_2$O was then retrotranscribed using qScript Flex cDNA Synthesis Kit

(QuantaBio, Beverly, MA). A volume totaling 1 µg RNA was incubated with random hexamers

and then supplemented with retrotranscriptase enzyme mix to synthesize cDNA. cDNA resulting

from the last step was used as a template for the amplification of a 256 bp fragment with

oligonucleotides SV00027A/SV00027 in a CFX96 Real-Time PCR System (Bio-Rad, Hercules,

CA). Primers had been designed to amplify a region of pHC_DYOLacI comprising the beginning

of *sfgfp*, so that when the tested chimera was being expressed it repressed the expression of GFP

unless Neu5Ac was present. Cq values were analyzed as a measure of gene expression using the

$\Delta\Delta$Cq ($2^{-\Delta\Delta Ct}$) method[84].

Protein Structure Prediction using AlphaFold 3. The 3D structures of the target proteins, Sphnx

(LacI-LNK2-SiaP, N$_t$-DBD-EKEKEK-LBD-C$_t$) and Kunst (LacI-LNK3-SiaP, N$_t$-DBD-

GSGSGS-LBD-C$_t$), were predicted using AlphaFold 3[70], developed by Google DeepMind

(London, UK). The predictions were performed using the publicly available AlphaFold 3 web

server (https://golgi.sandbox.google.com/). Both Sphnx and Kunst proteins were modeled as

dimers, along with the DNA double helix containing *lacO* (5'-ATTGTGAGCGGATAACAATT-

3' and its complementary sequence). Post-prediction, the structures were visualized and further

analyzed using VMD[85] to determine key protein-protein interactions.

Acknowledgments and Funding

This material is based upon work supported by St. John's University (SJU) via J.F.J.'s starting package as well as St. John's SEED grants (2020-2021, 2021-2022). Analysis and visualization of predicted structures was performed by F.X.V and A.B. on hardware resources supported by the National Institutes of Health under award number NIH SC2GM131992. This work is solely the responsibility of the authors and does not necessarily represent the official views of St. John's University. We want to acknowledge L. Bryan, D. Howarth, T. Mustacchio, M. Ruggiu, and W. Munyao (SJU) for their support and time sharing equipment, reagents, and protocols.

Author contributions

S.J.V. and J.F.J. developed the concepts, conceived of the study plan, and wrote the manuscript. S.J.V. designed and performed the experiments with assistance from S.A.P., V.A.S.E., G.X.C., A.R., A.C.L., E.S.N., P.K., and D.T.S. J.F.J provided project administration, supervision, resources, and data analysis, the latter with V.A.S.E. and S.J.V.'s assistance. A.B. performed 3D structure modeling and analysis of protein interactions under the supervision of F.X.V., who wrote the corresponding section of the manuscript.

Competing interests

The authors declare no competing interests.

 TABLES

Table 1. Chimeric transcription factors discussed in this work.

Name		TF behaviour		Comments
Common	**Systematic**	**Repressor**	**De-repressible**	
Siren	LacI-LNK1-SiaP	Yes	Yes	Functional only in late stationary phase
Sphnx	LacI-LNK2-SiaP	Yes	Yes	Biosensor candidate
Kunst	LacI-LNK3-SiaP	Yes	No	-
Scorpio	LacI-LNK4-SiaP	No	No	-
Bastet	LacI-LNK5-SiaP	No	No	Repression in early hours of growth.
Lumos	LacI-LNK6-SiaP	No	No	Duplicated sequence acts as Linker.
Harpy	LacI-LNK1-SatA	Yes	Unreliable	Unstable strain
Haus	LacI-LNK2-SatA	No	No	-

Figure 1

Figure 1. Screening strategy to find functional chimeric transcription factors. a) Biosensor genetic circuit for the *in vivo* testing of the functionality of cloned chimeric TFs. MG1655 (pCKT-*Chimera*, pHC_DYOLacI-R) cells express a chimeric TF under the control of TetR. The chimeric product can repress in *trans* the expression of a GFP reporter unless Neu5Ac is present. *tetR* / TetR (*grey arrow / circle*); chimeric TF gene / protein (*gold-blue arrow / circle*); *gfp* /GFP (*green arrow / circle*); anhydrotetracycline, aTc; neuraminic acid, Neu5Ac. **b)** Standard representation of the B-or-not-A universal two input gate that governs the expression of GFP in the circuit. **c)** States of the biosensor genetic circuit (top to bottom). *Basal*, the absence of aTc (aTc$^-$) enables TeR to repress the expression of the chimeric repressor: regardless of the presence or absence of Neu5Ac GFP is expressed, and a fluorescent signal is identified. *Repressed*, the presence of aTc (aTc$^+$) induces TetR which is unable to repress the promoter driving the transcription of the chimeric TF, which in turn represses the promoter driving the expression of *gfp* so that no fluorescent signal is detected. *De-repressed*, the presence of aTc (aTc$^+$) ensures the production of the chimeric TF which, if functional, gets in turn induced by Neu5Ac, so that it cannot repress *gfp* and the GFP-associated fluorescence is detected again. **d)** Expected profile of fluorescent production over time by a bacterial culture expressing the biosensor genetic circuit and a functional chimeric TF. Relative fluorescence (GFP-associated fluorescence corrected by cell density measured as OD$_{600}$). *Positive control* (*green line*), corresponds to the maximum GFP-associated fluorescence that can be observed, corresponding to a NEB5-alpha (pHC_DYOLacI-R) strain in which no repressor of the promoter driving GFP expression is present. *Negative control* (*orange line*), displays background fluorescence by a strain that does not carry *gfp*. *Basal* (*grey line*), *Repressed* (*red line*), *De-repressed* (*blue line*): GFP production associated to a MG1655 (pCKT-*Chimera*, pHC_DYOLacI-R) culture expressing a chimera in the basal, repressed, and de-repressed states, respectively.

Figure 2

Figure 2. *In vivo* validation of the functionality of Sphnx (LacI-LNK2-SiaP). a) Time course showing relative fluorescence of MG1655 (pCKT-Sphnx, pHC_DYOLacI-R) cells grown in a multiwell plate reader. Promoter activity is measured as $\log_2$ fold of the relative fluorescence (GFP-associated fluorescence in arbitrary units / OD_{600}) of the strain growing in the de-repressed (aTc^+, $Neu5Ac^+$; *blue boxplots*) or repressed (aTc^+, $Neu5Ac^-$; *red boxplots*) states compared to the basal expression of the reporter (aTc^-, $Neu5Ac^-$). Boxplots with whiskers represent data dispersion of the average values of biological replicas (n = 8-10). It can be appreciated how in order to approach the basal activity ($\log_2$ fold = 0), the addition of neuraminic acid is necessary when the chimera gets expressed by the supplementation of aTc. Growth conditions and fluorescence assays performed as described in *Materials and Methods*.

b) Sphnx and Siren protein interact with their cognate promoter P_{lac}. EMSA assays were performed as indicated under *Materials and Methods*. Interaction between NEB5-alpha (pCKT-*Chimera*) cell extracts expressing either Sphnx or Siren chimeric TFs and a dsDNA probe (251 bp) including the P_{lac} promoter. A fixed concentration of cell extract and DNA probe was exposed to growing concentrations of neuraminic acid (Neu5Ac). Lane numbers refer to the Neu5Ac concentration (mM) used for each reaction. The positions in the gel associated to unbound and bound probes are identified with orange and blue dashed rectangles, respectively.

c) Relative fluorescence of MG1655 (pCKT-Sphnx, pHC_DYOLacI-R) cultures after induction with different concentrations of neuraminic acid and growth in a multiwell plate reader for 14 hours. Promoter activity is measured as relative fluorescence (GFP-associated fluorescence in arbitrary units / OD_{600}) of the strain growing in the *basal* (aTc^-, $Neu5Ac^-$; *brown*), repressed (aTc^+, $Neu5Ac^-$; *red*) or de-repressed (aTc^+, $Neu5Ac^+$; *blue shades*) conditions. The concentrations of neuraminic acid (mM) added to the de-repressed conditions are specified under the corresponding plots. Boxplots with whiskers represent data dispersion of the average values of biological replicas (n = 8). The average value for every condition corresponds to a white circle. Growth conditions, induction protocol, and fluorescence assays performed as described in *Methods*. Statistically significant differences between selected conditions are marked with stars (p value < 0.05).

Figure 3

Figure 3. Modeled structures of chimeric protein dimers with double helix DNA obtained with Alphafold 3. For each structure, one monomer is shown using the New Cartoon representation and the other using the space-filling Surf representation. The DBD is shown in *green*, the PBP in *cyan*, and the connecting chain in *orange*, with the engineered linker in *violet*. The Kunst (**a**) and Sphnx (**b**) dimers are shown looking down the axis of the double helix, along with the rotated Kunst (**c**) and Sphnx (**d**) structures. The next panels contain the zoomed-in helical protein (**e**) linker (*violet*) in contact with the DBD (green) and PBP through the formation of salt bridges, with acidic residues shown in red and basic residues in blue. The zoomed-out protein (**f**) highlights the location of aromatic residues (*white*) that can be used to explore the validity of predicted structures using fluorescence anisotropy.

BIBLIOGRAPHY

1. Pancrazio, J. J., Whelan, J. P., Borkholder, D. A., Ma, W. & Stenger, D. A. Development and application of cell-based biosensors. *Ann. Biomed. Eng.* **27**, 697–711 (1999).

2. Su, L., Jia, W., Hou, C. & Lei, Y. Microbial biosensors: a review. *Biosens. Bioelectron.* **26**, 1788–1799 (2011).

3. Moraskie, M. *et al.* Microbial whole-cell biosensors: Current applications, challenges, and future perspectives. *Biosens. Bioelectron.* **191**, 113359 (2021).

4. O'Connor, G. *et al.* Whole-Cell Biosensors as Tools for the Detection of Quorum-Sensing Molecules: Uses in Diagnostics and the Investigation of the Quorum-Sensing Mechanism. *Adv. Biochem. Eng. Biotechnol.* (2015) doi:10.1007/10_2015_337.

5. Chang, H.-J., Voyvodic, P. L., Zúñiga, A. & Bonnet, J. Microbially derived biosensors for diagnosis, monitoring and epidemiology. *Microb. Biotechnol.* **10**, 1031–1035 (2017).

6. Wu, Y., Wang, C.-W., Wang, D. & Wei, N. A Whole-Cell Biosensor for Point-of-Care Detection of Waterborne Bacterial Pathogens. *ACS Synth. Biol.* **10**, 333–344 (2021).

7. Goode, J. A., Rushworth, J. V. H. & Millner, P. A. Biosensor Regeneration: A Review of Common Techniques and Outcomes. *Langmuir* **31**, 6267–6276 (2015).

8. Zhang, F. & Keasling, J. Biosensors and their applications in microbial metabolic engineering. *Trends Microbiol.* **19**, 323–329 (2011).

9. He, W., Yuan, S., Zhong, W.-H., Siddikee, M. A. & Dai, C.-C. Application of genetically engineered microbial whole-cell biosensors for combined chemosensing. *Appl. Microbiol. Biotechnol.* **100**, 1109–1119 (2016).

10. Saltepe, B., Kehribar, E. Ş., Su Yirmibeşoğlu, S. S. & Şafak Şeker, U. Ö. Cellular Biosensors with Engineered Genetic Circuits. *ACS Sens* **3**, 13–26 (2018).

11. Berepiki, A., Kent, R., Machado, L. F. M. & Dixon, N. Development of High-Performance

Whole Cell Biosensors Aided by Statistical Modeling. *ACS Synth. Biol.* **9**, 576–589 (2020).

12. Schleif, R. DNA binding by proteins. *Science* **241**, 1182–1187 (1988).

13. Rogers, J. K. *et al.* Synthetic biosensors for precise gene control and real-time monitoring of

metabolites. *Nucleic Acids Res.* **43**, 7648–7660 (2015).

14. Tellechea-Luzardo, J., Stiebritz, M. T. & Carbonell, P. Transcription factor-based biosensors

for screening and dynamic regulation. *Front Bioeng Biotechnol* **11**, 1118702 (2023).

15. Juárez, J. F., Lecube-Azpeitia, B., Brown, S. L., Johnston, C. D. & Church, G. M. Biosensor

libraries harness large classes of binding domains for construction of allosteric transcriptional

regulators. *Nat. Commun.* **9**, 3101 (2018).

16. Koch, M., Pandi, A., Borkowski, O., Batista, A. C. & Faulon, J.-L. Custom-made

transcriptional biosensors for metabolic engineering. *Curr. Opin. Biotechnol.* **59**, 78–84

(2019).

17. Kaufman, E. & Lamster, I. B. The diagnostic applications of saliva--a review. *Crit. Rev. Oral*

Biol. Med. **13**, 197–212 (2002).

18. Lee, Y.-H. & Wong, D. T. Saliva: an emerging biofluid for early detection of diseases. *Am.*

J. Dent. **22**, 241–248 (2009).

19. Wren, M. E., Shirtcliff, E. A. & Drury, S. S. Not all biofluids are created equal: chewing over

salivary diagnostics and the epigenome. *Clin. Ther.* **37**, 529–539 (2015).

20. Khanna, P. & Walt, D. R. Salivary diagnostics using a portable point-of-service platform: a

review. *Clin. Ther.* **37**, 498–504 (2015).

21. Bandodkar, A. J. & Wang, J. Non-invasive wearable electrochemical sensors: a review.

Trends Biotechnol. **32**, 363–371 (2014).

22.   Campuzano, S., Pedrero, M., Torrente-Rodríguez, R. M. & Pingarrón, J. M. Affinity-based

674   wearable electrochemical biosensors: Natural versus biomimetic receptors. *Anal. Sens.*

675   (2022) doi:10.1002/anse.202200087.

676   23.   Kim, J., Campbell, A. S., de Ávila, B. E.-F. & Wang, J. Wearable biosensors for healthcare

677   monitoring. *Nat. Biotechnol.* **37**, 389–406 (2019).

678   24.   Omar, E. Current concepts and future of noninvasive procedures for diagnosing oral

679   squamous cell carcinoma--a systematic review. *Head Face Med.* **11**, 6 (2015).

680   25.   Wang, S., Yang, M., Li, R. & Bai, J. Current advances in noninvasive methods for the

681   diagnosis of oral squamous cell carcinoma: a review. *Eur. J. Med. Res.* **28**, 53 (2023).

682   26.   Haines-Menges, B. L., Whitaker, W. B., Lubin, J. B. & Boyd, E. F. Host Sialic Acids: A

683   Delicacy for the Pathogen with Discerning Taste. *Microbiol Spectr* **3**, (2015).

684   27.   Blix, G. Concerning the carbohydrate groups of submaxillary mucin. *Hoppe Seylers Z.*

685   *Physiol. Chem.* **240**, 43–54.

686   28.   Lundblad, A. Gunnar Blix and his discovery of sialic acids. Fascinating molecules in

687   glycobiology. *Ups. J. Med. Sci.* **120**, 104–112 (2015).

688   29.   Röhrig, C. H., Choi, S. S. H. & Baldwin, N. The nutritional role of free sialic acid, a human

689   milk monosaccharide, and its application as a functional food ingredient. *Crit. Rev. Food Sci.*

690   *Nutr.* **57**, 1017–1038 (2017).

691   30.   Vajaria, B. N. *et al.* Salivary glyco-sialylation changes monitors oral carcinogenesis.

692   *Glycoconj. J.* **31**, 649–659 (2014).

693   31.   Phansopa, C. *et al.* Characterization of a sialate-O-acetylesterase (NanS) from the oral

694   pathogen Tannerella forsythia that enhances sialic acid release by NanH, its cognate sialidase.

695   *Biochem. J* **472**, 157–167 (2015).

32.   Arthisri, A. S., Sathiyamoorthy, A., Meenakshi, B. & Chandran, C. R. Ratio of Salivary Sialic

Acid to Fucose as Tumor Markers in Potentially Malignant Disorders and Oral Cancer.

Contemp. Clin. Dent. **11**, 131–135 (2020).

33.   Chaudhari, V., Pradeep, G. L., Prakash, N. & Mahajan, A. M. Estimation of salivary sialic

acid in oral premalignancy and oral squamous cell carcinoma. *Contemp. Clin. Dent.* **7**, 451–

456 (2016).

34.   Chittemsetti, S., Manchikatla, P. K. & Guttikonda, V. Estimation of serum sialic acid in oral

submucous fibrosis and oral squamous cell carcinoma. *J. Oral Maxillofac. Pathol.* **23**, 156

(2019).

35.   Markopoulos, A. K. Current aspects on oral squamous cell carcinoma. *Open Dent. J.* **6**, 126–

130 (2012).

36.   Coletta, R. D., Yeudall, W. A. & Salo, T. Grand Challenges in Oral Cancers. *Front Oral*

Health **1**, 3 (2020).

37.   Warnakulasuriya, S. & Greenspan, J. S. *Textbook of Oral Cancer: Prevention, Diagnosis and*

Management. (Springer Nature, 2020).

38.   Jacob, T. V. *et al.* A non-invasive study to estimate and compare salivary sialic acid level as

tumor marker in patients with pre-cancer and oral cancer. *J. Cancer Res. Ther.* **12**, 634–639

(2016).

39.   Sanjay, P. R., Hallikeri, K. & Shivashankara, A. R. Evaluation of salivary sialic acid, total

protein, and total sugar in oral cancer: a preliminary report. *Indian J. Dent. Res.* **19**, 288–291

(2008).

40.   Sridharan, G., Ramani, P., Patankar, S. & Vijayaraghavan, R. Evaluation of salivary

metabolomics in oral leukoplakia and oral squamous cell carcinoma. *J. Oral Pathol. Med.* **48**,

299–306 (2019).

41. Robson, B. Bioinformatics studies on a function of the SARS-CoV-2 spike glycoprotein as

the binding of host sialic acid glycans. *Comput. Biol. Med.* **122**, 103849 (2020).

42. Stanton, B. C. *et al.* Systematic transfer of prokaryotic sensors and circuits to mammalian

cells. *ACS Synth. Biol.* **3**, 880–891 (2014).

43. Jacob, F. & Monod, J. Genetic regulatory mechanisms in the synthesis of proteins. *J. Mol.*

Biol. **3**, 318–356 (1961).

44. Meinhardt, S. *et al.* Novel insights from hybrid LacI/GalR proteins: family-wide functional

attributes and biologically significant variation in transcription repression. *Nucleic Acids Res.*

40, 11139–11154 (2012).

45. Fukami-Kobayashi, K., Tateno, Y. & Nishikawa, K. Parallel evolution of ligand specificity

between LacI/GalR family repressors and periplasmic sugar-binding proteins. *Mol. Biol.*

Evol. **20**, 267–277 (2003).

46. Sakaguchi-Mikami, A., Taniguchi, A., Sode, K. & Yamazaki, T. Construction of a novel

glucose-sensing molecule based on a substrate-binding protein for intracellular sensing.

Biotechnol. Bioeng. **108**, 725–733 (2011).

47. Giuliani, S. E. *et al.* Environment sensing and response mediated by ABC transporters. *BMC*

Genomics **12 Suppl 1**, S8 (2011).

48. Fukamizo, T., Kitaoku, Y. & Suginta, W. Periplasmic solute-binding proteins: Structure

classification and chitooligosaccharide recognition. *Int. J. Biol. Macromol.* **128**, 985–993

(2019).

49. Locher, K. P. Mechanistic diversity in ATP-binding cassette (ABC) transporters. *Nat. Struct.*

Mol. Biol. **23**, 487–493 (2016).

50. Post, D. M. B., Mungur, R., Gibson, B. W. & Munson, R. S., Jr. Identification of a novel

sialic acid transporter in *Haemophilus ducreyi*. *Infect. Immun.* **73**, 6727–6735 (2005).

51. Gruteser, N., Marin, K., Krämer, R. & Thomas, G. H. Sialic acid utilization by the soil

bacterium Corynebacterium glutamicum. *FEMS Microbiol. Lett.* **336**, 131–138 (2012).

52. Müller, A. *et al.* Conservation of structure and mechanism in primary and secondary

transporters exemplified by SiaP, a sialic acid binding virulence factor from *Haemophilus*

influenzae. *J. Biol. Chem.* **281**, 22212–22222 (2006).

53. Severi, E. *et al.* Sialic acid transport in *Haemophilus influenzae* is essential for

lipopolysaccharide sialylation and serum resistance and is dependent on a novel tripartite

ATP-independent periplasmic transporter. *Mol. Microbiol.* **58**, 1173–1185 (2005).

54. Gangi Setty, T., Cho, C., Govindappa, S., Apicella, M. A. & Ramaswamy, S. Bacterial

periplasmic sialic acid-binding proteins exhibit a conserved binding site. *Acta Crystallogr. D*

Biol. Crystallogr. **70**, 1801–1811 (2014).

55. Gangi Setty, T. *et al.* Molecular characterization of the interaction of sialic acid with the

periplasmic binding protein from. *J. Biol. Chem.* **293**, 20073–20084 (2018).

56. Oertel-Buchheit, P., Reinbolt, J., John, M., Granger-Schnarr, M. & Schnarr, M. A LexA

mutant repressor with a relaxed inter-domain linker. *Protein Sci.* **7**, 512–515 (1998).

57. Tsirigotaki, A., De Geyter, J., Šoštaric, N., Economou, A. & Karamanou, S. Protein export

through the bacterial Sec pathway. *Nat. Rev. Microbiol.* **15**, 21–36 (2017).

58. Glick, B. R. Metabolic load and heterologous gene expression. *Biotechnol. Adv.* **13**, 247–261

(1995).

59. Liu, Q., Schumacher, J., Wan, X., Lou, C. & Wang, B. Orthogonality and Burdens of

Heterologous AND Gate Gene Circuits in *E. coli*. *ACS Synth. Biol.* **7**, 553–564 (2018).

60. Logic Gates. *The Department of Mathematics and Computer Science at Oxford College of Emory University* https://mathcenter.oxford.emory.edu/site/cs170/gates/.

61. Lewis, M. The lac repressor. *C. R. Biol.* **328**, 521–548 (2005).

62. Deans, T. L., Cantor, C. R. & Collins, J. J. A tunable genetic switch based on RNAi and repressor proteins for regulating gene expression in mammalian cells. *Cell* **130**, 363–372 (2007).

63. Langdon, R. C., Burr, T., Pagan-Westphal, S. & Hochschild, A. A chimeric activator of transcription that uses two DNA-binding domains to make simultaneous contact with pairs of recognition sites. *Mol. Microbiol.* **41**, 885–896 (2001).

64. Durante-Rodríguez, G. *et al.* Identification of a missing link in the evolution of an enzyme into a transcriptional regulator. *PLoS One* **8**, e57518 (2013).

65. Durante-Rodríguez, G., Mancheño, J. M., Díaz, E. & Carmona, M. Refactoring the λ phage lytic/lysogenic decision with a synthetic regulator. *Microbiologyopen* **5**, 575–581 (2016).

66. Oktay, S. *et al.* Serum and saliva sialic acid in periodontitis patients with and without cardiovascular disease. *Pathophysiol. Haemost. Thromb.* **37**, 67–71 (2010).

67. Snoek, T. *et al.* Evolution-guided engineering of small-molecule biosensors. *Nucleic Acids Res.* **48**, e3 (2020).

68. Hebisch, E., Knebel, J., Landsberg, J., Frey, E. & Leisner, M. High variation of fluorescence protein maturation times in closely related *Escherichia coli* strains. *PLoS One* **8**, e75991 (2013).

69. Chen, X., Zaro, J. L. & Shen, W.-C. Fusion protein linkers: property, design and functionality. *Adv. Drug Deliv. Rev.* **65**, 1357–1369 (2013).

70. Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with AlphaFold

3. *Nature* (2024) doi:10.1038/s41586-024-07487-w.

71. Meuzelaar, H., Vreede, J. & Woutersen, S. Influence of Glu/Arg, Asp/Arg, and Glu/Lys Salt

Bridges on α-Helical Stability and Folding Kinetics. *Biophys. J.* **110**, 2328–2341 (2016).

72. Raman, S., Taylor, N., Genuth, N., Fields, S. & Church, G. M. Engineering allostery. *Trends*

Genet. **30**, 521–528 (2014).

73. Malisi, C. *et al.* Binding pocket optimization by computational protein design. *PLoS One* **7**,

e52505 (2012).

74. Lerma Romero, J. A. *et al.* Binding pocket stabilization by high-throughput screening of yeast

display libraries. *Front Mol Biosci* **9**, 1023131 (2022).

75. Levine, M. J. Development of artificial salivas. *Crit. Rev. Oral Biol. Med.* **4**, 279–286 (1993).

76. Persico, M. *et al.* From Protein Communication to Drug Discovery. *Curr. Top. Med. Chem.*

15, 2019–2031 (2015).

77. Long, M., Betrán, E., Thornton, K. & Wang, W. The origin of new genes: glimpses from the

young and old. *Nat. Rev. Genet.* **4**, 865–875 (2003).

78. Bertani, G. Lysogeny at mid-twentieth century: P1, P2, and other experimental systems. *J.*

Bacteriol. **186**, 595–600 (2004).

79. Miller, J. H. *Experiments in Molecular Genetics*. (1972).

80. Sambrook, J. & Russell, D. W. *Molecular Cloning: A Laboratory Manual*. (CSHL Press,

2001).

81. Sanger, F., Nicklen, S. & Coulson, A. R. DNA sequencing with chain-terminating inhibitors.

Proc. Natl. Acad. Sci. U. S. A. **74**, 5463–5467 (1977).

82. Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred kilobases.

Nat. Methods **6**, 343–345 (2009).

83. Yanisch-Perron, C., Vieira, J. & Messing, J. Improved M13 phage cloning vectors and host

812   strains: nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* **33**, 103–119 (1985).

813   84. Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time

814   quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402–408 (2001).

815   85. Humphrey, W., Dalke, A. & Schulten, K. VMD: visual molecular dynamics. *J. Mol. Graph.*

816   **14**, 33–8, 27–8 (1996).

817