# Closely matched comparisons suggest that separable processes mediate contextual size illusions

Dietrich S. Schwarzkopf*, Xinran A. Yu, & Livia F. Fischer

*School of Optometry & Vision Science, University of Auckland, New Zealand*

* Corresponding author: s.schwarzkopf@auckland.ac.nz

**Abstract**

Previous research suggests the magnitudes of the Ebbinghaus, Delboeuf, Ponzo, and tilt illusions all depend on the cortical distance between the neural representations of target stimuli and the surrounding context. However, several psychophysical studies found no compelling association between these illusions, calling this hypothesis into question. Here we ask if these discrepant reports could arise from methodological differences between these studies. We ran a battery of visual size illusion and basic discrimination tasks with carefully matched geometric properties, using a classic forced choice design. In our small, homogenous sample, the Ebbinghaus and Delboeuf illusion magnitudes were strongly correlated, consistent with the idea that they reflect the same underlying mechanism when other sources of individual differences are minimized. Ponzo illusion magnitude also correlated with these two illusions, albeit to a lesser extent, indicative that both shared and independent factors govern these illusions. Interestingly, the classic arrowhead version of the Mueller-Lyer illusion did not correlate with any of the other illusions or even with the objective ability to discriminate line length. This suggests that an altogether separate process underlies this perceptual effect. We further demonstrate that presenting stimuli briefly with central fixation critically affects measurements of the Ebbinghaus illusion. Taken together, our findings suggest that different tasks probe separable processes determining illusion measurements. They further highlight the importance of the experimental design when testing relationships between perceptual effects and their links to neural processing.

**Keywords**

Spatial vision, perception, illusions, cortical distance, primary visual cortex, psychophysics, experimental design, correlations, discrimination, subjective

**Introduction**

The human visual system is not a digital camera. Our interpretation of visual space is subject to many contextual effects that can manifest as perceptual illusions. These effects can arise from basic limitations in neural architecture, such as the spatial resolution in peripheral vision being limited due to fewer neuronal resources than in central vision. In contrast, other illusions reflect the tasks the visual system has evolved to solve. For instance, navigating and interacting with a three-dimensional world involves interpreting the size of objects and their distance from the observer. Faced with a two-dimensional image that deliberately violates real-world relationships can lead to perceptual distortions.

*Mechanisms mediating contextual illusions*

Several illusions of spatial context have been linked to the functional architecture and/or response properties of the visual cortex. Starting with a seminal study (Murray et al., 2006), numerous reports have established that activity in human primary visual cortex (V1) reflects the apparent, not the veridical, eccentricity of stimuli (Fang et al., 2008; He et al., 2015; Ni et al., 2014), the apparent size of retinal afterimages (Sperandio et al., 2012), the repulsive perceptual effect of adapting to different stimulus sizes (Pooresmaeili et al., 2013), and the perceived separation in a dot variant of the Mueller-Lyer illusion (Ho & Schwarzkopf, 2022). One study also suggested that effective connectivity between cortical representations of target and context stimuli correlates with the magnitude of the tilt illusion (Song, Schwarzkopf, Lutti, et al., 2013). In contrast to studies on the neural representation of illusions, we found that individual differences in the surface area of V1 correlate with the magnitude of the Ebbinghaus and Ponzo illusions (Schwarzkopf et al., 2011; Schwarzkopf & Rees, 2013) and the tilt illusion (Song, Schwarzkopf, & Rees, 2013). We found hints of a similar link also for the Delboeuf illusion (Moutsiana et al., 2016), although that study merely inferred illusion magnitude from raw perceptual biases, rather than measuring it via direct comparisons. Moreover, the relationship was significant only for some stimulus parameters. Taken together, these findings allude to a possible role of cortical magnification in mediating these contextual illusions: the cortical distance between target and context stimuli in V1 explains a substantial amount of the variance in the magnitude and direction of these visuospatial context illusions (Mareschal et al., 2010; Schwarzkopf & Rees, 2013; Urale & Schwarzkopf, 2023). Other indications that the Ebbinghaus illusion is at least partly mediated by early stages of visual processing are that it is substantially reduced when targets and inducers are presented to different eyes (Song et al., 2011) and it partly persists when inducers are masked from awareness by continuous flash suppression (Chen et al., 2018). These findings suggest a role for early, monocular channels.

The Ebbinghaus is often described as a "size contrast" effect whereby the perceived size of a target appears to change depending on the sizes of surrounding contextual stimuli (Massaro & Anderson, 1971). However, this interpretation is partly based on a naïve understanding of the effect in its classic configuration: a disc surrounded by a context of small discs (inducers),

appears larger than the same disc surrounded by large inducers (Figure 1A). However, this interpretation misses the fact that when testing each inducer configuration separately against a no-context control, the illusion is in fact an underestimation of target size for most stimulus parameters (Roberts et al., 2005; Schwarzkopf & Rees, 2013; Urale & Schwarzkopf, 2023). Rather than size contrast, the illusion seems in large parts a geometric effect of the inducers, based on the interaction between contours (Todorović & Jovanović, 2018). The parameters exerting the most profound influence on the illusion are the distance between the target and the inducers, and how much of the surround is covered by inducers (Jaeger & Grasso, 1993; Jaeger & Klahs, 2015; Roberts et al., 2005; Urale & Schwarzkopf, 2023). This has led to the suggestion that the Delboeuf illusion is in fact a special case of the Ebbinghaus: an Ebbinghaus illusion with an annulus of very small, densely packed inducers is effectively a Delboeuf stimulus, yielding a nigh identical profile of illusion magnitudes (Roberts et al., 2005). According to this interpretation, results for these two illusions should be highly similar – that is, strongly correlated – across observers. Other studies have indeed found these two illusions to be highly similar, indicative of a role of contour attraction as the mediating mechanism (Sherman & Chouinard, 2016). The Ebbinghaus illusion is however more complex than the Delboeuf; the surrounding context consists of multiple stimuli of varying sizes, which are likely to interact with each other as well as the target. Moreover, there have been various suggestions of purportedly higher-level modulators of the Ebbinghaus illusion, such as the similarity between targets and inducers (Choplin & Medin, 1999; Deni & Brigner, 1997). While some of these effects are conceivably due to low-level feature interactions, certain studies suggest that semantic similarity may also play a role (Coren & Enns, 1993). The illusion effect likely involves both low-level contour interaction and higher-level processes related to size judgement (Jaeger & Klahs, 2015; Weintraub, 1979). If true, this implicates factors beyond local geometric interactions mediated by V1 circuits. This in turn predicts that a substantial proportion of the variance in the Ebbinghaus illusion is unrelated to that of the Delboeuf illusion.

Similar questions arise about the Ponzo illusion (Figure 1C). Researchers often consider the Ponzo to reflect the interpretation of distance in depth from flat two-dimensional images, a failure of size constancy (Gregory, 1966, 2008). Such a process conceivably involves higher-order processing well beyond primary visual cortex, because it requires interpreting three-dimensional scenes and relies on neurons with larger receptive fields than those in V1. This hypothesis is borne out by findings that the Ponzo illusion exhibits complete interocular transfer (Song et al., 2011) and that it is wiped out when the context is rendered invisible through continuous flash suppression (Chen et al., 2018). Research, including our own, has also shown that both the classic Ponzo, a corridor-type illusion with real-world scenes, and various steps in between, all show an inversion effect: the illusion is substantially stronger if the context implies the ground rather than the ceiling (Altan et al., 2023; Brislin, 1974; Poom, 2020). One study also found that a version of the Ponzo illusion, in which the local geometry of the target remained constant, irrespective of whether it suggested a near or far interpretation, yielded the most pronounced illusion compared to other versions

(Grzeczkowski et al., 2018). While there were still inevitable low-level differences between targets in this version, all these findings are better explained by a higher-level interpretation of the images than by a lower-level geometric interaction. While we found that the Ponzo illusion correlates with V1 surface area, just like the Ebbinghaus, this correlation was probably driven by an influential outlier and is therefore not robust (Schwarzkopf et al., 2011). Curiously, we also found no direct correlation between the Ebbinghaus and Ponzo illusion magnitudes themselves (but see e.g., Mazuz et al., 2023). Nevertheless, geometric parameters could theoretically play a role in the Ponzo. In the classic Ponzo configuration, the target which is typically perceived as smaller (and nearer to the observer) is generally positioned farther from the contextual stimuli; this conforms with the effect of target-inducer distance in the Ebbinghaus and Delboeuf illusions (Roberts et al., 2005; Urale & Schwarzkopf, 2023). Even in real-world corridor-type versions of the Ponzo, the distance between the "near/smaller" target and landmarks in the contextual image is greater. Interestingly, bandpass filtering the contextual images across different spatial frequency ranges does not impact the illusion's magnitude (Yildiz et al., 2021). This suggests that while a purely low-level mechanism for this illusion is unlikely, it does not rule out a contribution of such processes.

Finally, in the classic Mueller-Lyer illusion, the lines are perceptually lengthened or shortened depending on the direction of arrowheads at the line ends (Figure 1D). This effect is often thought to involve higher-order analysis of three-dimensional scenes, such as interpreting the two configurations as edges/corners near or far from the observer (Gregory, 1966). However, evidence against this are the facts that the Brentano version of this illusion manifests as a perceived position shift and correspondingly the mid-point of the line, conflicting with the interpretation of lines as a cue for depth. The illusion is also effective for simple dot variants devoid of any lines (Ho & Schwarzkopf, 2022). The perception of this dot variant is reflected in V1 responses (Ho & Schwarzkopf, 2022), while the Brentano version is reduced when adapting to low, but not high, spatial frequency patterns (Carrasco et al., 1986). Nevertheless, the current literature cannot conclusively rule out a role of higher-order processing in this illusion, especially in the classic line-and-arrowhead version, which tends to produce stronger effects than stripped down versions like the dot variant (Coren, 1970). As with the Ponzo and the Ebbinghaus, it is likely that multiple separate mechanisms also contribute to the Mueller-Lyer.

Given this diversity, it seems inconceivable that there could be a common factor to all spatial contextual illusions. This has not stopped researchers from searching for such factors, however (Axelrod et al., 2017; Cretenoud et al., 2021; Grzeczkowski et al., 2017; Jastrzębowska et al., 2023; Makowski et al., 2023). Interestingly, the magnitude of some of these illusions may be partly heritable (Coren & Porac, 1979) and related to personality traits (Makowski et al., 2023). A potential shared factor underlying this could be visual *abilities,* like spatial acuity and sensitivity thresholds (Halpern et al., 1999), which could have downstream effects on illusion magnitudes (Coren & Porac, 1987). Yet, like for illusion magnitudes, correlations have been found only for few basic visual abilities (Cappe et al., 2014). A large psychophysical study on simultaneous contrast tasks found that the magnitudes of these

illusions are mostly uncorrelated between stimulus dimensions (Bosten & Mollon, 2010), consistent with our own findings comparing size and luminance perception (Song, Schwarzkopf, & Rees, 2013). Often only versions of the same illusion (e.g., different configurations of the Ebbinghaus) share any significant variance at all (Grzeczkowski et al., 2017). However, many of these reports used highly distinct stimulus parameters, making no effort to match conditions. Such an experimental design is reasonable for asking if there are any correlations between illusions in the broadest sense. For effects that arise in higher decision-making stages of cognition the exact stimulus parameters may be largely irrelevant. However, when testing hypotheses about the involvement of circuits in the early visual system, this is extremely problematic. If both the Delboeuf and Ebbinghaus illusion depend on the cortical distance between target and inducers, it is imperative that any comparison between these illusions controls for extraneous factors, rather than presenting each illusion at different visual field locations with different geometric parameters.

Experiments investigating these visual cortical processes therefore also require brief stimulus presentation and stable eye fixation. Both are essential for ensuring that stimuli are presented in fixed locations of the visual field and thus target the appropriate neuronal populations in retinotopic visual cortex. Yet, most studies exploring the relationship between different illusions used unconstrained experimental designs – again, a reasonable approach for studying the higher-order cognitive aspects of these tasks, but highly problematic for studying low-level perceptual processing.

*Measuring illusion magnitude*

The ways how illusion magnitudes are measured is also critical. Any behavioural measures, including those quantifying perceptual experience, reflect the output of a whole brain working in concert. This ranges from multiple steps of image processing, making perceptual inferences, and eventually producing a motor output to report the judgement. This is another aspect in which no illusion is plausibly mediated solely by a single process, as any of these component processes can modulate the outcome. Numerous paradigms exist for quantifying the magnitude of an observer's perceptual illusions, each of which involves cognitive confounds to some degree: this includes adjustment/reproduction tasks (e.g., Coren & Girgus, 1972; Ho & Schwarzkopf, 2022; Jastrzębowska et al., 2023), forced-choice procedures (e.g., Schwarzkopf & Rees, 2013), confidence ratings (Gallagher et al., 2019), and drift diffusion models (Sánchez-Fuenzalida et al., 2023). Adjustment tasks may be particularly prone to demand effects because they rely on a subjective criterion for when there is a perceptual match between a test and reference stimulus. Reiterating an argument made by Morgan and colleagues (Morgan et al., 2013), one can imagine the difficulty of training a laboratory rat to perform an adjustment task. In contrast, the task is trivial to explain to human observers and may in fact be more natural and ecologically valid for how humans usually make perceptual judgements. And even simply by virtue of giving observers an extended time window to consider their response, adjustment likely involves higher cognition.

However, it is now widely recognised that even classic forced-choice procedures – considered
unbiased for measuring objective performance like acuity or detection thresholds (Green &
Swets, 1966) – are susceptible to decision-making confounds. This realization has led to the
development of more complex procedures based on comparing the 'difference between
differences' to disentangle perceptual from cognitive biases (Finlayson et al., 2017; Gallagher
et al., 2019; Jogan & Stocker, 2014; Manning et al., 2017; Morgan et al., 2012; Morgan et al.,
2013; Patten & Clifford, 2015).

It could be argued that forced-choice tasks, at least when using brief stimulus presentations,
involve a memory component. Because the stimulus has already disappeared by the time the
observer can respond, observers must rely on an internal representation of the stimuli to
make a perceptual judgement. This could involve iconic or working memory, although with
brief stimuli and speeded responses the former is perhaps more likely. Judging which of two
simultaneous stimuli is larger is not all that different from judging the orientation of a briefly
flashed grating – yet few people regard the latter task as a test of memory function.
Nevertheless, one could consider this a higher-level process than reproducing a perceptual
match in an adjustment task. On balance, however, the long decision window to inspect
stimuli and make an adjustment still allows more time for cognition to play a role than
speeded decisions in a forced choice task. And even if there were other high-level mechanisms
at play in forced choice tasks, briefly presented stimuli inevitably provide better control over
eye fixation and stimulus properties.

Therefore, the classic two-alternative forced choice (2AFC) design in which observers must
make a perceptual judgement between a constant reference and a variable test stimulus
remains a widely used method for quantifying illusion magnitudes. It constitutes a
compromise between disentangling cognitive confounds for measuring perceptual bias and
efficiency. Provided the observer's expectations or preconceived notions of what the
measurement reflects (such as suggestions that an illusion might be stronger/weaker in
certain demographics) can be minimised, the 2AFC design is arguably still an adequate method.
One way to achieve this is by randomly interleaving different conditions, randomising
when/where reference and test stimuli are presented to control for response bias, and using
a speeded design to prevent the observer from relying on non-perceptual cues in their
decision. As explained, this design importantly also make it relatively straightforward to
control gaze and thus the low-level stimulus parameters.

This design differs from most of the studies that failed to find correlations between contextual
illusions because these used adjustment procedures (Cretenoud et al., 2021; Cretenoud,
Francis, et al., 2020; Cretenoud, Grzeczkowski, et al., 2020; Grzeczkowski et al., 2017;
Jastrzębowska et al., 2023). Adjustment tasks require prolonged stimulus presentation, during
which observers tend to shift their gaze between stimuli while determining a perceptual
match. Some previous experiments were also conducted online or outside lab environments,
allowing minimal control over stimulus characteristics and eye gaze. Generally, adjustment

tasks can be done in considerably less time than forced-choice paradigms. However, this
confounds any conclusions we can draw from these findings.

For example, one of the studies that reported similar estimates for an adjustment and a 2AFC
design (Cretenoud et al., 2021) found correlations between 0.6-0.67, meaning that between
half to two thirds of the variance remained unexplained. It is conceivable that the correlation
in these experiments largely reflects the higher-level decision processes but that the residuals
– where the two measurements diverge – are due to the underlying stimulus-dependent
processing by the visual system. While this is not explicitly stated in the methods, it is likely
that these experiments did not control eye gaze and used unlimited stimulus presentations in
the 2AFC tasks, further confounding any conclusions we can draw from these experiments
about early visual processing. The illusion strength doubtless generalizes across tasks up to a
point, but the exact parameters – such as the retinal location of the stimuli – have a profound
effect (Jaeger & Grasso, 1993; Jaeger & Klahs, 2015; Roberts et al., 2005; Urale & Schwarzkopf,
2023). Relatedly, a study on another geometric illusion, the vertical-horizontal (bisection)
illusion, demonstrated that gaze stability affects illusion strength (Chouinard et al., 2017).

*The present study*

Here we set out to quantify the individual differences in these contextual size modulations,
specifically the Ebbinghaus, Delboeuf, Ponzo, and Mueller-Lyer illusions. We tested a battery
of these size illusions, using a classic two-alternative forced choice (2AFC) procedure with brief
stimulus presentations and fixation instructions. We also included two objective
discrimination tasks for judging the width of discs and the length of lines, both without any
illusion-inducing contexts. This was based on our previous research, which indicated
associations between the objective acuities and subjective illusion magnitudes and associated
spatial selectivity of visual neuronal populations (Moutsiana et al., 2016; Song et al., 2015;
Song, Schwarzkopf, & Rees, 2013). The stimulus conditions across all these tasks were closely
matched. For example, target stimuli were identical filled discs for testing the Ebbinghaus,
Delboeuf, and Ponzo illusions as well as the corresponding disc width discrimination tasks, and
the length of the lines in the Mueller-Lyer illusion and the corresponding line length
discrimination task matched the diameters of the discs in the other tasks (Figure 1A-F). In the
same vein, the distances between targets and inducers were kept comparable across the
different illusions. This design therefore maximised the chance of finding associations
between these perceptual effects; the influence of any extraneous stimulus differences should
be minimised.

We tested a series of preregistered hypotheses aimed to better understand the mechanisms
mediating these illusions. Critically, under the proposition that the Ebbinghaus and Delboeuf
illusions are effectively the same phenomenon (Roberts et al., 2005), we predicted that these
two effects are strongly correlated, directly contradicting other recent reports (Jastrzębowska
et al., 2023). Of note, a similar approach to ours – to minimize stimulus differences – also

suggested a strong correlation between the Ebbinghaus and Delboeuf illusions when using an adjustment task (Sherman & Chouinard, 2016).

In a second experiment, we further assessed the effects of task and stimulus parameters on the Ebbinghaus illusion magnitude. We compared illusion magnitudes across three conditions: an adjustment task, a 2AFC task with unlimited stimulus presentation and free eye movements, and a classic 2AFC task with brief stimulus presentation and fixation instructions, as used in our earlier studies investigating brain-perception links for this illusion (Schwarzkopf & Rees, 2013). We also compared the effect of using two different stimulus parameters because the stimulus parameters described in our earlier study contained an error (see Materials and Methods for details), meaning researchers seeking to replicate those experiments following the methods would inadvertently use different stimuli than our original study (Jastrzębowska et al., 2023).

**Materials and Methods**

Procedures were preregistered on the Open Science Framework for both Experiment 1 (https://osf.io/bhgud) and Experiment 2 (https://osf.io/5ey8h).

*Participants*

Thirteen observers (ages 18-45, 10 female, 3 left-handed) were recruited for this study, both from the staff and student body of the University of Auckland as well as the wider community. Three of these were the authors, but the remaining observers were naïve to the purpose of the experiments. One observer was excluded from the whole study based on preregistered criteria (see below). Another observer was excluded from Experiment 2 only. Observers gave written informed consent, and all procedures were approved by the University of Auckland Human Participants Research Ethics Committee. Observers received a NZ$25 supermarket voucher for their participation in both experiments.

*Experiment 1*

*Stimuli*

Stimuli were closely matched across the different experimental conditions. In all tests, the fixation dot was a white disc (luminance: 140 cd/m$^2$) with diameter of 0.15° visual angle, presented in the centre of a black screen (luminance: 0.6 cd/m$^2$). All other stimuli were a dark grey (luminance: 6.2 cd/m$^2$). The line width of any unfilled discs and line stimuli was 0.1°. All target stimuli were centred at an eccentricity of 4.65° to the left and right of fixation. The reference stimulus size (disc diameter or line length) was always 2.06°. The width or length of test stimuli varied on a binary logarithmic scale. The reference and test stimuli could appear either on the left or right of fixation. The main differences between the experimental conditions were in the design of the contextual stimuli (Figure 1A-F):

1. *Disc width discrimination:* Only the target discs were presented. The stimulus design was identical to the control condition in our earlier study (Schwarzkopf & Rees, 2013), but note that the methods in that article incorrectly described the radius of the disc stimuli as diameters.

2. *Ebbinghaus illusion:* The target stimuli were filled discs, surrounded by other filled discs, the inducers, that were placed in equal angular steps, starting from the 3 o'clock position. Sixteen small inducers (0.52° diameter) were centred 1.86° from the centre of the reference stimulus. Six large inducers (4.14° diameter) were centred 4.34° from the centre of the test stimulus. These stimuli were also identical to the illusion stimuli in our earlier study (Schwarzkopf & Rees, 2013).

3. *Delboeuf illusion:* The target stimuli were filled discs, surrounded by rings. The ring around the reference and test stimuli had a radius of 1.6° and 4.34°, respectively.

4. *Ponzo illusion:* The target stimuli were filled discs. They were placed within a context of two converging lines that could either converge towards the left or right. The lines started and ended 5.9° horizontally from the vertical meridian in both directions. To converge, their ends were positioned above and below the horizontal meridian at a distance 4.3° and 1.24°, respectively. The reference and test stimuli were at the narrower and wider ends, respectively.

5. *Line length discrimination:* The target stimuli were vertical lines. They bisected horizontal lines (1.3° long) at each end.

6. *Mueller-Lyer illusion:* The target stimuli were vertical lines. At each end, they were joined to two 0.65° long lines angled at ±45° relative to vertical, forming an arrowhead. In the test and reference stimuli, the lines were directed towards or away from the horizontal meridian, respectively.

*Procedure*

After giving consent, participants were seated in front of a computer screen (Dell P2214H 21.5", dimensions: 47.6 cm x 26.8 cm, resolution: 1920 x 1080 pixels, controlled by an integrated Intel UGD Graphics 770 card) at a viewing distance of 57 cm, with their heads stabilised by a chin and forehead rest. The experiment was controlled by a Dell Latitude laptop 5410 running MATLAB R2023b, using Psychtoolbox 3 (Brainard, 1997), running under 64-bit Windows 10 Education (version 22H2) operating system. The general experimental design was a 2AFC task with an identical trial structure for all illusion and discrimination conditions. We matched stimulus parameters between the conditions as much as reasonably feasible. The task involved making a perceptual judgement about which target disc was larger or which line was longer, respectively. There were always two stimuli presented simultaneously within each trial, with one being a constant reference (diameter/length of discs/lines: 2.06°) and the other a test stimulus that varied across trials.

The experimental conditions were presented in the following sequence: participants were pseudo-randomised to start with either line judgements (line length discrimination and Mueller-Lyer illusion) or the disc judgements (disc width discrimination, and Ebbinghaus,

Delboeuf, and Ponzo illusions) was. However, the objective discrimination tasks for these judgements always came first before testing the corresponding illusions. This familiarized observers – many of whom had little practice with visual psychophysics – with the task under simpler conditions before doing it under subjective modulations of size perception. The order of the three disc illusions was counterbalanced across participants, resulting in six possible permutations.

For each condition, the participant first performed a few demo trials in which the stimuli were presented until the participant's keyboard response. This allowed the experimenter to explain the task. This was followed by more practice trials in which the trial sequence was as in the actual experiment, with 100 ms stimulus presentations. Once the participant confirmed they understood the task, the actual experiment began.

Each trial started with a brief 500 ms fixation period where only the fixation dot was presented on a black screen. This was followed by a 100 ms stimulus presentation, after which the screen turned completely black. The participant must then make a response to judge whether the left or right target stimulus was larger/longer by pressing the corresponding arrow key. The next trial then started immediately after the response.

The 160 trials were divided into four blocks of 40 trials each. There were four trial types: the two interleaved staircases (starting above or below true equality) and the reference stimulus could either appear to the left or the right of fixation. These four trial types were repeated 10 times per block in a pseudo-randomised order. Before each block, there was a rest screen reminding the participant of the task instructions and informing them of the number of blocks they had already completed. They commenced the block by pressing any key on the keyboard.

Stimulus levels were expressed in binary logarithms of the ratio between the test and reference size (diameter for discs, length for lines). This linearises stimulus values to account for Weber's Law and allows the use of parametric statistical tests, which would not be applicable in the case of raw size ratios.

Test stimulus levels were controlled by two interleaved 1-down, 1-up staircase procedures that converged onto the point of subjective equality (PSE). One staircase started at a higher stimulus level (0.4 log units, wider disc or longer line) than true equality (0 log units), while the other staircase started at a lower level (-0.4 log units, narrower disc or shorter line). The staircase increased or decreased the stimulus level following the response in each trial. Initially, the increment/decrement was at a larger step size of 0.1 log units. Once five reversals had occurred in a given staircase, the step size was reduced to 0.05 log units. Moreover, from this point on, there was also a 10% probability on each trial that the stimulus level was increased or decreased (depending on the starting point of the given staircase) by 0.15 log units. These "jitter trials" prevented the stimulus level from continuously hovering around the PSE, as this could result in disengagement from the task.

*Data analysis*

Observers completed 160 trials per experimental condition. Data were then analysed by
calculating the proportion of trials in which the test stimulus was chosen at each stimulus level
used (collapsing across both staircases in each condition and irrespective of which side the
test was presented). We then fit a cumulative Gaussian function to these data points using
the summed squared residuals approach, taking into account the robustness of each data
point by weighting them by the number of trials at that stimulus level. We inferred the PSE
from the stimulus level where the curve crossed the 50% response level (centroid of the
psychometric curve). We also estimated discrimination uncertainty (conceptually a threshold
or just noticeable difference) by determining the bandwidth of the psychometric curve.

Our preregistered plan was to exclude observers who failed to complete all experimental
conditions; however, all observers completed all conditions. Further, we excluded participants
for whom the bandwidth (technically, the σ parameter) of any psychometric curve fit
exceeded 0.78 log units. Beyond this bandwidth, assuming the PSE is zero an observer would
only score 10% or 90%, respectively, for stimulus levels of +/- 1 log units (i.e., test stimuli half
or twice as large as the reference). We reasoned that this constitutes an implausibly poor
performance for these simple visual discrimination tasks in observers with supposedly normal,
healthy vision. All excluded participants were to be replaced to ensure correct
counterbalancing of conditions (but see below for deviation from these criteria). One observer
was excluded based on this criterion.

The PSE (centroids of psychometric curves) for the illusion conditions and the discrimination
abilities (bandwidths of psychometric curves) for the two objective discrimination tasks (disc
width and line length) were compared using Pearson's r correlation. We tested all hypotheses
using directional (one-tailed) tests, using a Bayes Factor with a default stretched beta prior
with width 1, as implemented in JASP (0.17.2.1).

*Sampling plan*

Our minimum sample size was set at N=6 because that is required to include all sequences for
disc judgement illusions. We then incremented the sample size in batches of 6, accessing the
stopping criterion after each batch. This criterion was based on hypothesis $I_1$, that is, whether
the one-tailed Pearson's correlation between the Ebbinghaus and Delboeuf illusion
magnitudes yielded a Bayes Factor of 10 or 0.1, thus supporting either the alternative or null
hypothesis, respectively. We chose this hypothesis for the stopping criterion because we
expected these two illusions to reflect the same underlying neural process (Roberts et al.,
2005; Schwarzkopf & Rees, 2013; Urale & Schwarzkopf, 2023), yet a previous study failed to
find any strong correlation between them (Jastrzębowska et al., 2023). Our maximum sample
size in case the stopping criterion was not reached was N=90. However, the criterion was
reached after two batches were collected, that is, at N=12.

*Preregistered hypotheses*

We tested the following hypotheses of between-subject correlations between psychometric
parameters estimated for individual participants:

*Hypothesis I₁:* Ebbinghaus and Delboeuf illusion magnitudes (PSEs) correlate positively,
425 because they are effectively the same process.

*Hypothesis I₂:* Ponzo illusion magnitude does not correlate (or only weakly) with magnitudes
427 of the other illusions, because it reflects higher-level processing of three-dimensional scenes.
428 (Note that this hypothesis contradicts exploratory hypothesis M₃ below as we had no specific
429 *a priori* predictions for the Mueller-Lyer illusion. If M₃ had been confirmed, we would have
430 expected I₂ to be only partially confirmed)

*Hypothesis S₁:* Ebbinghaus and Delboeuf illusion magnitudes each correlate positively with
432 bandwidth for objective disc width discrimination.

*Hypothesis S₂:* Ponzo illusion magnitude does not correlate positively (or only weakly) with
434 bandwidths for objective disc width discrimination.

*Hypothesis S₃:* Bandwidths for the objective discrimination tasks (disc width and line length)
436 correlate positively because they both rely on local V1 processing (such as receptive field size).

*Hypothesis M₁:* Mueller-Lyer illusion magnitude correlates positively with bandwidth for line
438 length discrimination.

*Exploratory Hypothesis M₂:* Mueller-Lyer illusion magnitude correlates positively with the
440 Ebbinghaus and Delboeuf illusion magnitudes if they share similar early neural mechanisms.
441 Some previous findings suggest that this illusion could be mediated by local interactions in V1
442 (Carrasco et al., 1986; Ho & Schwarzkopf, 2022; Morgan & Glennerster, 1991), but this has not
443 been tested using classic arrow-type Mueller-Lyer stimuli. Conversely, the correlation is only
444 weak if this illusion reflects higher-order scene processing.

*Exploratory Hypothesis M₃:* Mueller-Lyer illusion magnitude correlates positively with the
446 Ponzo illusion, if they both reflect the same process of interpreting three-dimensional scenes.

447 A summary of the outcome of these hypotheses is shown in Table 1.

### Experiment 2

*Stimuli*

One of the Ebbinghaus stimulus designs (*Grey&Large*) was identical to that used in Experiment
1 and was a direct replication of our earlier study (Schwarzkopf & Rees, 2013). The other
stimulus design (*White&Small*) was based on a replication study (Jastrzębowska et al., 2023),
which was based on the incorrect method description in our original study (although we note
that correct example stimuli were included in that study). Specifically, the spatial separation
and eccentricities were identical to those in the other condition, but the filled disc stimuli
were half width, because the methods description stated "diameter" where it should have

said "radius." For unexplained reasons, this replication study also presented stimuli in white instead of dark grey. Previous research suggests that white target stimuli are generally perceived as larger, which could impact the illusion estimate (Jaeger et al., 2014). Stimulus contrast also modulates contextual interactions in early visual cortex (Nauhaus et al., 2009), and could therefore affect the circuits we hypothesize mediate the Ebbinghaus illusion. To test this, we therefore also used white discs (luminance: 140 cd/m$^2$) in this condition. Compared to the dark grey stimuli, many observers found these stimuli abrasive, especially when they were flashed briefly (see issues with reliability for this condition below). For Grey&Large disc stimuli, the fixation dot was a white dot as described in the preceding experiment; but for White&Small disc stimuli, the fixation dot was black as it fell inside an inducer.

*Procedure*

There were no demo or practice trials in this experiment, because participants had already done a very similar Ebbinghaus measurement in Experiment 1 immediately prior to this experiment. Instead, participants were shown images of the two possible stimulus parameters (Figure 2A-B) on the screen, and the experimenter explained again what the perceptual judgement entailed.

Participants then performed three tasks, with the order of tasks counterbalanced across participants, resulting in six permutations:

1. *Fast-2AFC:* This was essentially identical to the Ebbinghaus measurement in the preceding experiment, except for the inclusion of the White&Small disc condition. Participants were explicitly instructed to fixate on the white dot in the centre of the screen. There were 320 trials, divided into blocks of 80 (160 trials per stimulus condition, i.e. the same number as in the preceding experiment where there was only one condition).

2. *Unconstrained-2AFC:* This was the same as the Fast-2AFC task, but there was no fixation dot shown and the illusion stimuli remained on screen until the participant responded with the keyboard. They were explicitly told that they are free to move their eyes and inspect the stimuli as they please.

3. *Adjustment:* Stimuli appeared on the screen without a fixation dot. Participants then used the up and down arrow key on the keyboard to change the width of the test disc (surrounded by large inducers) until it was a perceptual match for the width of the reference disc. Once they were satisfied with the match, they pressed the space bar to move to the next trial. There were 20 trials in this test, 10 for each of the stimulus conditions. The side where the reference and test stimuli were presented was pseudo-randomised in each trial.

Stimulus values in the two 2AFC tasks were controlled by interleaved staircase procedures, just as in Experiment 1, except that there were independent staircases for the two stimulus conditions. In the Adjustment task, the initial radius of the test stimulus in each trial was drawn from a Gaussian distribution with a standard deviation of 0.4 log units.

 *Data analysis*

Data from the two 2AFC tasks were analysed as in Experiment 1. For the Adjustment task, the
illusion magnitude in this experiment for each stimulus condition was calculated as the
arithmetic mean across the final adjusted size of the test stimulus from all trials. The paradigm
also collected information about the time course of the Adjustment task in each trial, but this
is irrelevant for the present study and will not be discussed further.

Mean illusion magnitudes were compared using 2x2 repeated-measures analysis of variance
using stimulus condition and task as factors. Post-hoc contrasts and paired two-tailed t-tests
compared individual conditions/tasks. The statistical evidence was based on default Bayes
Factors. Correlations were calculated based on Pearson's r, evaluating statistical evidence
based on default Bayes Factors. All these analyses were conducted in JASP 0.17.2.1 (JASP Team,
2023), using a default Cauchy prior with scale $r=\sqrt{2}/2$ for comparisons of means, and a
stretched beta prior with width 1 for testing correlations.

We also estimated the test-retest reliability of the illusion magnitudes in Experiment 2, using
split-half correlations. For the Adjustment task, we calculated the illusion magnitude
separately for odd and even numbered trials in each observer. For the 2AFC tasks, we refitted
the psychometric curves separately for the two independent, pseudo-randomly interleaved
staircases in each condition. We calculated the Pearson correlation, *r,* between the two halves
across observers, and then extrapolated the reliability of the full dataset, $R_{rel}$, using the
Spearman-Brown prophecy formula (Spearman, 1910):

$$R_{rel} = 2r / (1+r)$$

The magnitude of correlations between the Adjustment and 2AFC tasks was compared using
the Williams/Steiger equation for comparing two correlations with a shared variable (Howell,
1997). Statistical inference was based on the 95% confidence limit of the Williams/Steiger t-
statistic derived by resampling pairs of observers 10,000 times with replacement. A difference
in this directional test was deemed significant if the 95[th] percentile of this bootstrap
distribution was below zero. To compare correlations between stimulus conditions for the
separate tasks, we bootstrapped the difference in correlation coefficients by resampling pairs
of observers 10,000 times and calculating the 95% confidence level of this distribution. This
directional test was deemed significant if the 5[th] percentile was above zero.

Because we reused the same participants as in Experiment 1, the sampling plan for
Experiment 2 was determined by the first experiment. We also applied the same exclusion
criteria for Experiment 2, that is, if the bandwidth (technically, σ) of the psychometric curve
fit for any of the experimental conditions exceeded 0.78 log units, observers would be
excluded (without replacement, because recruitment was via Experiment 1). Based on this
criterion, one observer was excluded, leaving only N=11 for Experiment 2.

*Preregistered hypotheses*

*Hypothesis H₁:* Illusion magnitude differs between the Adjustment task and the 2AFC tasks, specifically:

- *H₁ₐ:* Illusion magnitude differs between Fast-2AFC and Adjustment.
- *H₁ᵦ:* Illusion magnitude differs between Fast-2AFC and Unconstrained-2AFC.

*Hypothesis H₂:* Illusion magnitudes correlate between tasks as follows:

- *H₂ₐ:* Illusion magnitudes correlate between Adjustment and Unconstrained-2AFC.
- *H₂ᵦ:* Illusion magnitudes correlate less strongly between Adjustment and Fast-2AFC.

*Hypothesis H₃:* Overall, illusion magnitude differs between the stimulus parameters irrespective of task. Qualitatively similar effects are found comparing stimulus parameters in each task separately.

A summary of the outcome of these hypotheses is shown in Table 1.

*Deviations from preregistration*

Our preregistered protocol also included a fourth hypothesis about the correlations between stimulus parameters. However, after data collection had commenced, we noticed that one sub-hypothesis (H₄ᵦ) in the preregistration was phrased incorrectly, effectively repeating hypothesis H₂ᵦ. The methods also did not contain the necessary description for the statistical comparison of these independent correlations. Even though this was intended to be a preplanned analysis, we therefore decided to label this as exploratory to avoid the impression of altering the preregistration after the results were known.

We also adjusted our exclusion criteria slightly. It turns out that the Fast-2AFC task with the White&Small stimulus condition was very challenging for most observers, with several reporting that this condition was only a momentary flash that made perceptual judgements difficult. One observer (P5) exceeded the exclusion criterion of bandwidth σ>0.78 for this condition (σ=0.85). However, this larger bandwidth was consistent with the values seen in the rest of the sample, and this observer's other results accorded well with the overall data. We therefore decided to retain this observer in the study. A control analysis confirmed that exclusion of this observer did not substantially alter the conclusions.

**Results**

Across two experiments, we aimed to better understand the discrepancy between previous studies in terms of the factors governing visual contextual size illusions. We quantified individual differences in illusion magnitudes and tested the effects of task and stimulus parameters on the Ebbinghaus illusion.

*Illusions with disc targets are related*

Our first experiment used a battery of tasks to compare the magnitudes of our range of illusions. Averaged across observers, the illusion magnitudes varied considerably, with by far the strongest effect for the Mueller-Lyer and the weakest for the Ponzo illusion (Figure 1G), with the Ebbinghaus and Delboeuf magnitude at similar intermediate levels. We also quantified the mean bandwidths of psychometric curves as a reflection of objective discrimination thresholds. This was largest (worst) for the Delboeuf and slightly elevated for the Ebbinghaus, but at similar levels for the other illusions and the objective discrimination tasks (Figure 1G).

Next, we calculated the pairwise correlations between the different illusions and the bandwidths for objective disc and line discrimination (Figure 1H, see also Supplementary Figure S1A for scatter plots of each correlation). This revealed a strong correlation between the Ebbinghaus and Delboeuf illusion (r=0.716, $BF_{+0}$=15.11), confirming our primary hypothesis $I_1$. Surprisingly, the Ponzo illusion also correlated positively with both the Ebbinghaus and Delboeuf illusions (Ebbinghaus: r=0.582, $BF_{+0}$=4.027; Delboeuf: r=0.597, $BF_{+0}$=4.533), contradicting earlier findings by ourselves and others (Jastrzębowska et al., 2023; Schwarzkopf et al., 2011), but in line with other recent studies (Mazuz et al., 2023). This rejected our preregistered hypothesis $I_2$.

Delboeuf illusion magnitude also correlated strongly with objective thresholds (bandwidths) for discriminating disc stimuli (r=0.648, $BF_{+0}$=7.136). We also observed positive correlations between disc discrimination and both the Ebbinghaus and Ponzo illusions, the other illusions using disc targets – but statistical evidence for these correlations was inconclusive ($BF_{+0}$<3). Our hypothesis $S_1$ about disc discrimination abilities was therefore only partially supported, and evidence for $S_2$ was inconclusive. Interestingly, we found moderate evidence for the null hypothesis of no relationship between thresholds for line discrimination and all illusion magnitudes ($BF_{+0}$<⅓), except for the Delboeuf (r=-0.006, $BF_{+0}$=0.349). This means not even the Mueller-Lyer illusion correlated with this objective task, despite using the same line targets, which rejects our preregistered hypothesis $M_1$.

There was also evidence against any relationship between the Mueller-Lyer and the Ebbinghaus illusions (r=-0.334, $BF_{+0}$=0.192). As such, our results do not adjudicate between the two exploratory hypotheses $M_2$ and $M_3$ we specified for the Mueller-Lyer in that it was neither linked with the assumed low-level effects (Ebbinghaus, Delboeuf) nor the presumably higher-level Ponzo illusion. Finally, contrary to hypothesis $S_3$ discrimination thresholds for lines and discs were effectively uncorrelated, although statistical evidence for the alternative or null hypothesis was inconclusive (r=0.101, $BF_{+0}$=0.455). Like many psychophysical studies on fundamental visual processing, our sample included the authors of this study. The pattern of results remained qualitatively unchanged when restricting the analysis to only the naïve observers (Figure S1B).

**Figure 1.** Stimuli (A-F) and results (G-H) of Experiment 1. In separate experimental runs, we estimated the illusion
magnitude/perceptual bias and discrimination sensitivity for the Ebbinghaus (A), Delboeuf (B), Ponzo (C), Mueller-Lyer (D)
illusions, or the discrimination of disc width (E) or line length (F). Red lines and numbers indicate dimensions in degrees of

visual angle. These were not part of the presented stimuli. G. Average illusion magnitude (dark discs, solid lines) and
discrimination bandwidths (light diamonds, dotted lines) for each condition. Both measures are the ratio of the test over
reference stimulus diameter, expressed as binary logarithm (see Materials and Methods). Error bars denote ±1 standard
error of the mean across observers. H. Correlation matrix comparing the magnitudes for the four illusions, Ebbinghaus
(Ebb), Delboeuf (Del), Mueller-Lyer (Mue), Ponzo (Pon), and the discrimination bandwidths for judging disc width (Dis) or
line length (Lin), respectively. Colours indicate strength of correlation. In each cell, symbols denote Bayes Factors: asterisk:
$BF_{+0}>3$, pentagram: $BF_{+0}>10$, circles $BF_{+0}<⅓$.

### *Task modulates illusion magnitude*

We next ran a battery of Ebbinghaus illusion measurements, using an Adjustment task, an Unconstrained-2AFC task (unlimited stimulus duration and free eye gaze), and a Fast-2AFC task (brief stimulus duration and instruction to maintain fixation). In this experiment, we also compared two different stimulus designs for the illusion (Figure 2A-B). Averaged across observers, illusion magnitudes varied substantially between the three tasks (Figure 2C; effect of task: $BF_{incl}=7.143$), confirming our preregistered hypothesis $H_1$. Specifically, illusion magnitudes were lower when measured with the Adjustment task than either the Unconstrained-2AFC ($BF_{10}=155.604$) or the Fast-2AFC task ($BF_{10}=16.919$). However, average illusion magnitude was similar for the two 2AFC tasks although evidence for this null effect was inconclusive ($BF_{10}=0.414$), leaving hypothesis $H_{1B}$ undetermined.

To further explore these data, we also compared discrimination bandwidths for the 2AFC tasks (Figure 2D). Bandwidths were consistently smaller for the Unconstrained-2AFC than the Fast-2AFC task ($BF_{10}=67.82$), meaning the latter was more difficult.

### *Different stimulus parameters do not affect illusion magnitude*

While there was a hint that the stimuli with White&Small discs yielded the weaker illusions, at least for the two tasks with unlimited stimulus duration and free eye gaze, statistical support for any differences between stimulus parameters was inconclusive ($BF_{incl}=0.398$), which is inconsistent with our preregistered hypothesis $H_3$. Importantly, this applies to the relative illusion effect, expressed as the ratio between test and reference stimulus diameters on a logarithmic scale. In terms of absolute size differences in degrees of visual angle, the illusion magnitude for the White&Small stimuli would be approximately half that of the Grey&Large stimuli.

The lack of any difference in illusion magnitude stands in contrast to a pronounced difference in discrimination ability. Bandwidths were consistently worse (larger) for White&Small than Grey&Large discs (Figure 2D; $BF_{10}=18.448$). Especially for the Fast-2AFC task, judgements became difficult for White&Small discs, to the extent that we decided to deviate from our preregistered exclusion criteria: the discrimination bandwidth for one observer exceeded our predefined criterion for this condition but we decided to include them, as this pattern was consistent with the whole sample and their other data were of high quality.

Illusion magnitudes for the two stimulus conditions were also strongly correlated, both averaged across tasks ($r=0.873$, $BF_{+0}=158.861$) as well as separately for the Adjustment ($r=0.918$, $BF_{+0}=697.718$) and the Unconstrained-2AFC task ($r=0.926$, $BF_{+0}=996.191$). For the Fast-2AFC task, illusion magnitudes were also positively correlated but statistical evidence for this relationship was not conclusive ($r=0.528$, $BF_{+0}=2.416$). Exploring this further we found that the correlation between stimulus conditions was significantly lower for the Fast-2AFC task than both the Adjustment ($\delta_R=0.389$, 5th percentile: 0.037) and the Unconstrained-2AFC task ($\delta_R=0.397$, 5th percentile: 0.128).

*Brief stimulation and fixation affect illusion measurements*

We then calculated the correlations between the illusion magnitudes from different tasks and stimulus conditions. This revealed that the Adjustment and Unconstrained-2AFC tasks were strongly correlated with one another (Figure 2E; all $BF_{+0}>18.394$). In contrast, while the Fast-2AFC task also correlated positively with the Adjustment task, these correlations were significantly attenuated (Grey&Large: $\delta_T=-2.591$, 95th percentile: -1.287; White&Small: $\delta_T=-1.966$, 95th percentile: -0.488). This fully supports our preregistered hypotheses $H_2$ about the individual differences between tasks. We observed the weakest correlation between the Fast-2AFC task with White&Small stimuli and the other conditions. While positive and evidence tended to be in favour of supporting the presence of correlations (all $BF_{+0}>1$), only some of the correlations for Grey&Large stimuli (with Adjustment: $r=0.662$, $BF_{+0}=6.443$; with Unconstrained-2AFC: $r=0.734$, $BF_{+0}=13.581$) and one for White&Small stimuli (with Unconstrained-2AFC with grey and small stimuli: $r=0.626$, $BF_{+0}=4.75$) conclusively supported this relationship with the other tasks. Scatter plots for these correlations are shown in Supplementary Figure S2A. Moreover, to rule out any undue bias due to inclusion of non-naïve (author) observers, we repeated this analysis restricted to naïve observers. This yielded qualitatively similar results (Supplementary Figure S2B).

*Delboeuf illusion shares most of its variance with Ebbinghaus*

We next estimated the test-retest reliability of illusion measurements from split-half data (Supplementary Figure S3). This showed that the measurement for most tasks and stimulus conditions were highly reliable ($R_{rel}\geq0.88$). Interestingly, the Fast-2AFC task with Grey&Large stimuli was most reliable, almost reaching ceiling ($R_{rel}=0.99$). The only exception was the Fast-2AFC task with White&Small stimuli, which was far less reliable ($R_{rel}=0.67$). The lower correlations and reduced reliability for the Fast-2AFC task with White&Small stimuli were presumably due to overall poorer performance for this condition. However, the Fast-2AFC measurements for the Grey&Large stimuli were extremely reliable, even though this task was more challenging than the Adjustment and Unconstrained-2AFC (Figure 2D). Since this condition was effectively the same as the Ebbinghaus task in the first experiment and used the same observers (except for one excluded based on preregistered criteria in the Experiment 2), we were able to obtain a separate reliability estimate. Comparing these measurements also revealed a strong correlation (albeit weaker than the internal reliability in Experiment 2)

between illusion magnitudes (Figure 2F, black curve; r=0.788, $BF_{+0}$=28.569) and a slightly weaker correlation between discrimination bandwidths (Figure 2F, blue curve; r=0.704, $BF_{+0}$=9.718). Of note, this test-retest reliability for illusion magnitude is only slightly greater than the correlation between the Ebbinghaus and Delboeuf illusions we observed in Experiment 1 (Figure 1H; r=0.716).

**Table 1.** Preregistered hypotheses in this study. *Ebb:* Ebbinghaus illusion magnitude. *Del:* Delboeuf illusion magnitude. *Pon:* Ponzo illusion magnitude. *Mue:* Mueller-Lyer illusion magnitude. *Dis:* Disc width discrimination bandwidth. *Lin:* Line lentgh discrimination bandwidth. *Adjst:* Adjustment task. *Unco-2AFC:* Unconstrained-2AFC task.

| **Experiment 1 – Comparison of several size illusions and related discrimination tasks** | | |
|---|---|---|
| *Hypothesis* | | *Outcome* |
| $I_1$ | Ebb and Del correlate positively | **Supported** |
| $I_2$ | Pon does not correlate with Ebb, Del, or Mue | **Rejected** (except for Mue) |
| $S_1$ | Ebb and Del correlate positively with Dis | Inconclusive and **Supported** |
| $S_2$ | Pon does not correlate positively with Dis | Inconclusive |
| $S_3$ | Dis and Lin correlate positively | Inconclusive |
| $M_1$ | Mue correlates positively with Lin | **Rejected** |
| $M_2$ | Mue correlates positively with Ebb and Del | **Rejected** and Inconclusive |
| $M_3$ | Mue correlates positively with Pon | Inconclusive |
| **Experiment 2 – Effect of task and stimulus conditions on Ebbinghaus illusion** | | |
| *Hypothesis* | | *Outcome* |
| $H_{1A}$ | Ebb differs between Fast-2AFC and Adjst | **Supported** |
| $H_{1B}$ | Ebb differs between Fast-2AFC and Unco-2AFC | Inconclusive |
| $H_{2A}$ | Ebb correlates positively between Adjst and Unco-2AFC | **Supported** |
| $H_{2B}$ | Ebb correlates less strongly between Adst and Fast-2AFC | **Supported** |
| $H_3$ | Ebb differs between Grey&Large and White&Small | Inconclusive |

**Discussion**

Our findings demonstrate a strong correlation between the magnitudes of the Ebbinghaus and Delboeuf illusions when using a classic two-alternative forced choice (2AFC) procedure, as we used in our earlier work investigating the links between perception and neural processing. This correlation is almost as strong as the test-retest correlation for the Ebbinghaus task itself. These findings confirm previous suggestions that the Ebbinghaus and Delboeuf are essentially manifestations of the same underlying effect (Roberts et al., 2005; Sherman & Chouinard, 2016; Todorović & Jovanović, 2018; Urale & Schwarzkopf, 2023). Even though Ebbinghaus stimuli are considerably more complex - with multiple contextual stimuli

defined by numerous parameters that likely interact with one another as well as with the
target – the Delboeuf illusion magnitude remarkably explains most of the variance in the
Ebbinghaus.

This finding however contrasts with other recent reports that these two illusions are not
substantially related (Jastrzębowska et al., 2023). From our present findings, we deduce that
this discrepancy could be due to the differences in methodology: previous research mostly
used adjustment procedures. This approach not only fails to control stimulus duration and
retinotopic location but could also be skewed by cognitive and/or decision-making biases,
given the extended time to make a perceptual judgement (Morgan et al., 2013). Even in the
few cases where previous studies employed 2AFC procedures, experiments did not seem to
control stimulus characteristics adequately and were also likely susceptible to decision
confounds. In fact, the correlation between tasks in that earlier study (Cretenoud et al., 2021)
was considerably weaker than the correlation we found between Adjustment and
Unconstrained-2AFC tasks in our second experiment. This discrepancy could call the reliability
of these earlier experiments into question. One potential explanation is that we used more
trials for our Adjustment procedure and therefore could estimate the illusion magnitude more
precisely.

Our present study took care to match the stimulus conditions between the illusions and
discrimination tasks. We note that previous research taking a similar approach to closely
match stimulus conditions also found strongly correlated estimates for the Ebbinghaus and
Delboeuf illusions, even when using an adjustment task (Sherman & Chouinard, 2016).
Matching stimuli is essential for drawing conclusions about how these perceptual effects
relate to neural representations in visual cortex and how they may depend on local cortical
magnification. Importantly, this also maximises the chance of observing any correlation
between the illusions; all else being equal, the only remaining difference is the type of illusion
used. Assuming these effects depend on visual cortex representations, any relationship
between these effects must be diminished when the conditions differ in terms of geometric
dimensions and where in the visual field the stimuli are presented.

Therefore, it is perhaps not unsurprising that we also found some less pronounced
correlations between the Ponzo illusion and the other two using disc targets (Ebbinghaus,
Delboeuf) and that all these illusions correlated positively (albeit only relatively weakly and
mostly without being statistically conclusive) with the objective ability to discriminate disc
widths. While there have been reports of an association between the Ebbinghaus and Ponzo
(Grzeczkowski et al., 2017; Mazuz et al., 2023), other research has found weak association
between these illusions at best (Jastrzębowska et al., 2023; Schwarzkopf et al., 2011),
consistent with the idea that the Ebbinghaus is largely mediated by local interactions between
V1 neurons (Jaeger et al., 2014; Jaeger & Grasso, 1993; Jaeger & Klahs, 2015; Roberts et al.,
2005; Schwarzkopf & Rees, 2013; Song et al., 2011; Todorović & Jovanović, 2018; Urale &
Schwarzkopf, 2023), while the Ponzo involves interpretations of the three-dimensional scene
– a process that highly likely recruits higher-order processes (Chen et al., 2018; Song et al.,

2011). Our findings now add further nuance to this interpretation. As with most illusions, the
Ponzo likely represents a mixture of both low- and high-level processes. The correlation
captures the former variance, which is shared with the Ebbinghaus and Delboeuf, while
unexplained variance still remains that could stem from higher-order mechanisms.

Given our matching stimulus characteristics, the most striking aspect of our results is in fact
the *absence* of positive correlations, supported by statistical evidence, most notably when
comparing the Mueller-Lyer illusion with the others. The neural mechanisms mediating the
Mueller-Lyer remain unclear, as it could involve both high-level interpretation of three-
dimensional perspective (Gregory, 1966) and low-level image filtering (Carrasco et al., 1986;
Ho & Schwarzkopf, 2022; Morgan & Glennerster, 1991). Our results suggest that if the Ponzo
and the Mueller-Lyer both involve higher-level processes, these are unlikely to be the same
mechanism. Curiously, the Mueller-Lyer also did not correlate with the objective ability to
discriminate line lengths, and the two objective discrimination tasks (line length and disc
width) also appear unrelated. This indicates that the perceptual judgement underlying the
Mueller-Lyer is rather different from that mediating the other tasks. The Mueller-Lyer is also
by far the strongest and most consistent of the illusions we tested. In future work, it could be
interesting to design a Mueller-Lyer illusion with disc targets (e.g., surrounded by a crown of
arrowheads) to ascertain if the absence of correlation observed here is due to the different
target stimuli.

**Figure 2.** Stimuli (A-B) and results (D-F) of Experiment 2. We tested two configurations of the Ebbinghaus illusion: our standard stimulus from Experiment 1 (A) or another version with all discs half the diameter and in white (B). In separate experimental runs, we used an Adjustment, Unconstrained-2AFC, or a Fast-2AFC task to estimate the illusion magnitude and discrimination sensitivity (except for Adjustment task). Average estimates for illusion magnitude (C) and discrimination bandwidths (D) for each task (see Materials and Methods and Figure 1 for details about these measures). Large grey discs denote the Grey&Large stimulus condition and small unfilled discs the White&Small condition. Error bars denote ±1 standard error of the mean across observers. E. Correlation matrix comparing the magnitudes estimated for each condition (task and stimulus). Adjust: Adjustment task, Unco-2AFC: Unconstrained-2AFC task. G&L: Grey&Large, W&S: White&Small. Colours indicate strength of correlation. In each cell, symbols denote Bayes Factors: asterisk: $BF_{+0}>3$, pentagram: $BF_{+0}>10$, hexagram: $BF_{+0}>100$. F. Illusion

776  magnitude (black) and discrimination bandwidth (blue) estimated for the Fast-2AFC Grey&Large condition in Experiment 2
777  against the corresponding estimates in Experiment 1. Each dot denotes an individual observer. Solid lines show the linear
778  regression fit and the shaded regions its bootstrapped 95% confidence interval using 10,000 iterations and the percentile
779  method to derive confidence limits.

In our second experiment, we compared Ebbinghaus illusion measurements between different tasks. We found that estimates from Adjustment tasks were very similar to those obtained with the Unconstrained-2AFC task, where observers could inspect stimuli for as long as they needed and were free to move their eyes. In contrast, estimates were far less consistent between these tasks and the Fast-2AFC task, where eye gaze and stimulus duration were tightly controlled. It is tempting to argue that these diminished correlations are a trivial consequence of poorer discrimination performance for the Fast-2AFC task. Indeed, discrimination thresholds for this task, especially with the stimuli comprising White&Small discs, were consistently worse. However, Ebbinghaus illusion measurements, at least for Grey&Large stimuli, were highly reliable and revealed strong correlations with the Delboeuf and Ponzo illusions in Experiment 1. This again stands in contrast to a previous study using Adjustment tasks, which failed to find any compelling associations between these illusions (Jastrzębowska et al., 2023). Therefore, we surmise that the Fast-2AFC task can yield robust measurements. The lower correlation between the Fast-2AFC task and Adjustment estimates, compared to the Unconstrained-2AFC task, likely results from the latter's shared critical characteristics with Adjustment: unlimited stimulus presentation and unconstrained eye movements. Cognitive factors may also play a role. By giving observers long time windows to consider the stimuli, both Adjustment and Unconstrained-2AFC arguably involve more cognition. However, it could also be argued that the Fast-2AFC task entails a memory component because the stimuli have disappeared by the time the observer can respond, potentially involving an additional factor not involved in the unconstrained tasks. Whatever the reason, our results demonstrate that the tasks differ in terms of the underlying processes.

Indeed, by fixing eye position and limiting stimulus durations the Fast-2AFC design likely engages different neuronal populations in visual cortex than the Adjustment and Unconstrained-2AFC tasks. Only the Fast-2AFC task allows relatively tight control over this aspect. It is difficult to ascertain which regions of the visual field – and therefore which corresponding neuronal populations – are involved during perceptual judgements in the Adjustment and Unconstrained-2AFC tasks. Whenever an observer foveates one of the two target stimuli, it is represented with the highest spatial resolution in foveal cortex, while the other stimulus is relegated to a peripheral location with reduced resolution. Consequently, the eccentricity of this fellow stimulus would then be double that of both, the test and reference stimuli, in the Fast-2AFC task. If the cortical representation matters to the perceptual effect, the illusion estimate must inevitably differ. Notably, stimuli are typically perceived as smaller in peripheral vision (Anstis, 1998; Moutsiana et al., 2016; Newsome, 1972; von Helmholtz, 1924). By directing fixation to the midpoint between the two stimuli, the Fast-2AFC controls for this aspect, while the illusion estimates in the other two tasks may include this general

perceptual bias. The situation is complicated by the fact that observers can alternately foveate
the two targets, and they may also occasionally fixate in between them. It is difficult to
determine which of these states contribute most to the illusion estimate. Either way, it also
seems likely that these unconstrained tasks leave more room for decision factors to influence
the response. While not directly comparable, a study comparing the vertical-horizontal
(bisection) illusion, reported a profound effect of stable eye fixation (Chouinard et al., 2017).

In that light, it is puzzling that several previous studies using adjustment tasks found so few
correlations between different illusions (e.g., Axelrod et al., 2017; Jastrzębowska et al., 2023).
One might reasonably expect that if these tasks were dominated by decision factors, they
would show greater consistency with each other. For example, if these results reflected the
conservativism of their decision criterion to reproduce a perceptual match, why would this
criterion differ so wildly between these illusions? Such a decision criterion could be
considered a common factor to these illusions, possibly related to personality factors
(Makowski et al., 2023) – the accumulated data across the literature speak for themselves that
such a factor either does not exist or only explains such a small amount of variance that could
not be reliably detected even in many large-scale studies.

Certainly, such a decision factor should not be reflected by the functional architecture in V1.
Rather, local properties of V1, which vary with visual field location and geometric stimulus
dimensions, exert specific effects on perception. This does not mean illusion estimates do not
generalise up to a point across stimulus characteristics and visual field locations – some
generalisation is expected, as neural properties also generalise. For instance, the cortical
magnification factor or population receptive field sizes at 3° eccentricity must at least
correlate somewhat with those at 1° eccentricity. In our study, the relative illusion estimates
for the two stimulus conditions in the second experiment were indeed very similar, despite
substantial differences in absolute change in degrees of visual angle. More generally, although
task-related differences were significant, illusion estimates remained relatively well correlated
across tasks. Up to a point it is feasible to use comparable stimuli and experimental
parameters in perceptual studies and how it relates to the brain. However, the more
differences are introduced into the design, the greater the variability becomes and the less
likely it is that two measures correlate.

*Limitations*

We acknowledge that the sample size in our study (Experiment 1: N=12, Experiment 2: N=11)
was very small for studying individual differences. Our preregistered sampling plan prescribed
stopping data collection when we obtained strong evidence either for ($BF_{+0}>10$) or against
($BF_{+0}<0.1$) a correlation between the Ebbinghaus and Delboeuf illusion magnitude. Given
similar previous research (Jastrzębowska et al., 2023; Sherman & Chouinard, 2016), we
anticipated that we would require a larger sample, setting our maximum at N=90. However,
we already obtained compelling evidence that these two illusions are correlated after the
second testing milestone. As explained in the Introduction, we deliberately designed our

illusion stimuli to closely match fundamental attributes between conditions, thereby controlling for stimulus parameters known to affect these illusions (Jaeger et al., 2014; Jaeger & Grasso, 1993; Jaeger & Klahs, 2015; Roberts et al., 2005; Todorović & Jovanović, 2018; Urale & Schwarzkopf, 2023). This could have contributed to high similarity in these illusion estimates. Either way, our sample size was evidently sufficient to obtain conclusive results.

It could also be argued that our sample was demographically too homogenous, especially considering the small number of observers, leading to potential biases. As is common in psychophysics studies, the three non-naïve authors also participated in the experiment. Our control analyses confirm that their inclusion did not qualitatively affect the results. While the age of our observers ranged from young adults to middle-aged and ethnically comparably diverse, the majority were female. Most observers were either present or recent students/staff of the University of Auckland. The Ebbinghaus illusion reportedly varies with age, with weaker illusions in children (Doherty et al., 2010; Weintraub, 1979) and older age compared to young adults (Mazuz et al., 2024). It reportedly also varies across cultural contexts (Bremner et al., 2016; de Fockert et al., 2007). Thus, it is certainly possible that our limited sample restricted the range of illusion magnitudes one would find in a broad global sample.

However, this would also be accompanied by additional confounds. A broader age range, for example, could introduce variance related to changes in general visual abilities, while a global population likely varies in terms of their practice, concentration, and skill in performing such computerized tasks. This would inevitably reduce the reliability of these measures and the strength of associations between illusions. In turn, this would run contrary to the aims of our study, to isolate the existence of shared factors in low-level visual processing under well-controlled conditions. Just as we designed our stimuli to be closely matched, our comparably homogenous sample enabled us to reveal the role of stimulus parameters and task design without contamination by other confounding factors like age, gender, cultural context, or visual diet. This is a prerequisite step before larger global studies can establish which other factors contribute to illusion magnitudes.

Another potential limitation of our experiment is that the task instructions for the 2AFC tasks inadvertently differed from those in the Adjustment tasks. In the former, observers were instructed to choose the target disc that "was larger." In the latter, the instruction was to adjust the test disc size until both discs "appeared equal." It is possible that the exact instructions can affect how observers respond, especially when studying neurodivergent observers. However, we consider it unlikely that this played a significant role in our experiments. Illusion estimates for the Adjustment and Unconstrained-2AFC tasks were strongly correlated despite this discrepancy in task instructions. By contrast, correlations between these two tasks and the Fast-2AFC were significantly weaker, even though the Fast-2AFC task with Grey&Larg stimuli was the most reliable of all our conditions. This suggests that this task probed separate processes not tapped by the other two tasks.

*Conclusions*

Any illusion estimate must reflect multiple separate mechanisms (some of which may interact), ranging from basic perceptual processing to complex decision-making. Therefore, any notion of a single common factor that could be shared by all illusions, even all spatial contextual illusions of size as studied here, seems fundamentally misguided. Even for potential common factors like general intelligence or personality traits, extraneous variables can influence the outcome of specific tasks in individuals – for example, a colour-blind person would find it difficult to pick a correct target defined by colours even if intelligence is not the limiting factor. More importantly, a hypothetical common factor, equivalent to general intelligence, that determines a person's susceptibility to visual illusions is based on circular logic. To know that your percept disagrees with the physical reality of the stimulus requires the ability to separate the perceived stimulus from the retinal image. It seems inconceivable that such a general process would have evolved, as these illusions presumably reflect stimulus processing necessary for functional vision. Our hypothesis linking the Ebbinghaus, Delboeuf, and similar contextual effects like the tilt illusion, to cortical distance is based on the idea that they result from feedforward filtering of the visual input by receptive fields. This suggests that the information about the true stimulus has been filtered out at higher stages of processing. Any measure that reflects this neural processing of visual input, such as measuring cortical magnification and/or population receptive field size, should therefore explain some of the variance in these specific illusions.

However, this also means that the opposite implication suggested by some studies (Cretenoud et al., 2021; Grzeczkowski et al., 2017; Jastrzębowska et al., 2023) is even more implausible, namely that there are no factors shared at least by some illusions at all. Taken to its logical conclusion, this notion implies that each illusion is mediated by entirely separate neural mechanisms. This is akin to the concept of a "grandmother neuron" in memory. Just as it seems unlikely that there is only one cell (or a small population of cells) that encodes the identify of a specific person, it seems highly implausible that there are wholly separate neural mechanisms for the Ebbinghaus, Delboeuf, Ponzo, or Mueller-Lyer illusions. Many visual illusions are discovered by accident or deliberately designed based on prior knowledge of how visual processing works. They tap into mechanisms. Therefore, illusions which partly tap into the same mechanisms must also be statistically related to each other. More research is required to identify these factors.

For example, it remains possible that the Ebbinghaus illusion – despite being profoundly determined by basic stimulus attributes (Jaeger et al., 2014; Jaeger & Grasso, 1993; Jaeger & Klahs, 2015; Roberts et al., 2005; Sherman & Chouinard, 2016; Todorović & Jovanović, 2018; Urale & Schwarzkopf, 2023) – also includes a genuine size contrast factor. This could underlie at least some similarity effects reported for this illusion (Choplin & Medin, 1999; Coren & Enns, 1993; Deni & Brigner, 1997). The contribution of component mechanisms presumably also varies between individuals and with demographic factors. While the Ebbinghaus varies between populations (Bremner et al., 2016; de Fockert et al., 2007) and in children (Doherty

et al., 2010), such differences could result from changes in the size contrast factor or decision processes (Weintraub, 1979), rather than being explained by changes in cortical magnification. In fact, at least the macroscopic and mesoscopic functional architecture of early visual cortex is close to adult levels already at a young age (Dekker et al., 2019), although more subtle changes still occur over the course of development (Himmelberg et al., 2023). This underscores the importance of experimental paradigms that can disentangle these factors. Classical or novel forced choice procedures (e.g., Finlayson et al., 2017; Jogan & Stocker, 2014; Morgan et al., 2013; Patten & Clifford, 2015) with closely matched stimulus properties as used in our study, are a prerequisite for testing these low-level geometric factors of subjective perceptual experience. However, this does not mean that procedures like adjustment cannot yield important insights. Indeed, for determining the role of separate factors, a combination of different methods (Manning et al., 2017) may be the most fruitful approach.

**Data availability**

All stimulus presentation materials, data, and analysis code are publicly available: https://osf.io/5zs7y and https://osf.io/b3h4x

**Acknowledgements**

We thank Adina Giurgiu and Arash Chaparian for their help in setting up our new testing room.

**References**

Altan, E., Boyaci, H., & Schwarzkopf, D. S. (2023). Evidence for high-level processing in a Ponzo-like size illusion. *Journal of Vision*, *23*, 4715. https://doi.org/10.1167/jov.23.9.4715

Anstis, S. (1998). Picturing peripheral acuity. *Perception*, *27*(7), 817–825.

Axelrod, V., Schwarzkopf, D. S., Gilaie-Dotan, S., & Rees, G. (2017). Perceptual similarity and the neural correlates of geometrical illusions in human brain structure. *Scientific Reports*, *7*(1), Article 1. https://doi.org/10.1038/srep39968

Bosten, J. M., & Mollon, J. D. (2010). Is there a general trait of susceptibility to simultaneous contrast? *Vision Research*, *50*(17), 1656–1664. https://doi.org/10.1016/j.visres.2010.05.012

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

Bremner, A. J., Doherty, M. J., Caparos, S., de Fockert, J., Linnell, K. J., & Davidoff, J. (2016). Effects of Culture and the Urban Environment on the Development of the Ebbinghaus Illusion. *Child Development*, *87*(3), 962–981. https://doi.org/10.1111/cdev.12511

Brislin, R. W. (1974). The Ponzo Illusion: Additional Cues, Age, Orientation, and Culture. *Journal of Cross-Cultural Psychology*, *5*(2), 139–161. https://doi.org/10.1177/002202217400500201

Cappe, C., Clarke, A., Mohr, C., & Herzog, M. H. (2014). Is there a common factor for vision? *Journal of Vision*, *14*(8), 4. https://doi.org/10.1167/14.8.4

Carrasco, M., Figueroa, J. G., & Willen, J. D. (1986). A test of the spatial-frequency explanation of the Müller-Lyer illusion. *Perception*, *15*(5), 553–562. https://doi.org/10.1068/p150553

Chen, L., Qiao, C., Wang, Y., & Jiang, Y. (2018). Subconscious processing reveals dissociable contextual modulations of visual size perception. *Cognition*, *180*, 259–267. https://doi.org/10.1016/j.cognition.2018.07.014

Choplin, J. M., & Medin, D. L. (1999). Similarity of the perimeters in the Ebbinghaus illusion. *Perception & Psychophysics*, *61*(1), 3–12.

Chouinard, P. A., Peel, H. J., & Landry, O. (2017). Eye-Tracking Reveals that the Strength of the Vertical-Horizontal Illusion Increases as the Retinal Image Becomes More Stable with Fixation. *Frontiers in Human Neuroscience*, *11*. https://doi.org/10.3389/fnhum.2017.00143

Coren, S. (1970). Lateral inhibition and geometric illusions. *The Quarterly Journal of Experimental Psychology*, *22*(2), 274–278. https://doi.org/10.1080/00335557043000212

Coren, S., & Enns, J. T. (1993). Size contrast as a function of conceptual similarity between test and inducers. *Perception & Psychophysics*, *54*(5), 579–588.

Coren, S., & Girgus, J. S. (1972). A comparison of five methods of illusion measurement. *Behavior Research Methods & Instrumentation*, *4*(5), 240–244. https://doi.org/10.3758/BF03210006

Coren, S., & Porac, C. (1979). Heritability in visual-geometric illusions: A family study. *Perception*, *8*(3), 303–309.

Coren, S., & Porac, C. (1987). Individual differences in visual-geometric illusions: Predictions from measures of spatial cognitive abilities. *Perception & Psychophysics*, *41*(3), 211–219. https://doi.org/10.3758/BF03208220

Cretenoud, A. F., Francis, G., & Herzog, M. H. (2020). When illusions merge. *Journal of Vision*, *20*(8), 12–12. https://doi.org/10.1167/jov.20.8.12

Cretenoud, A. F., Grzeczkowski, L., Bertamini, M., & Herzog, M. H. (2020). Individual differences in the Müller-Lyer and Ponzo illusions are stable across different contexts. *Journal of Vision*, *20*(6), 4. https://doi.org/10.1167/jov.20.6.4

Cretenoud, A. F., Grzeczkowski, L., Kunchulia, M., & Herzog, M. H. (2021). Individual differences in the perception of visual illusions are stable across eyes, time, and measurement methods. *Journal of Vision*, *21*(5), 26. https://doi.org/10.1167/jov.21.5.26

de Fockert, J., Davidoff, J., Fagot, J., Parron, C., & Goldstein, J. (2007). More accurate size contrast judgments in the Ebbinghaus Illusion by a remote culture. *Journal of Experimental Psychology. Human Perception and Performance*, *33*(3), 738–742. https://doi.org/10.1037/0096-1523.33.3.738

Dekker, T. M., Schwarzkopf, D. S., de Haas, B., Nardini, M., & Sereno, M. I. (2019). Population receptive field tuning properties of visual cortex during childhood. *Developmental Cognitive Neuroscience*, 100614. https://doi.org/10.1016/j.dcn.2019.01.001

Deni, J. R., & Brigner, W. L. (1997). Ebbinghaus illusion: Effect of figural similarity upon magnitude of illusion when context elements are equal in perceived size. *Perceptual and Motor Skills*, *84*(3 Pt 2), 1171–1175.

Doherty, M. J., Campbell, N. M., Tsuji, H., & Phillips, W. A. (2010). The Ebbinghaus
illusion deceives adults but not young children. *Developmental Science*, *13*(5),
714–721. https://doi.org/10.1111/j.1467-7687.2009.00931.x

Fang, F., Boyaci, H., Kersten, D., & Murray, S. O. (2008). Attention-dependent
representation of a size illusion in human V1. *Current Biology*, *18*(21), 1707–
1712. https://doi.org/10.1016/j.cub.2008.09.025

Finlayson, N. J., Papageorgiou, A., & Schwarzkopf, D. S. (2017). A new method for
mapping perceptual biases across visual space. *Journal of Vision*, *17*(9), 5.
https://doi.org/10.1167/17.9.5

Gallagher, R. M., Suddendorf, T., & Arnold, D. H. (2019). Confidence as a diagnostic tool
for perceptual aftereffects. *Scientific Reports*, *9*(1), 7124.
https://doi.org/10.1038/s41598-019-43170-1

Green, D. M., & Swets, J. A. (1966). *Signal detection theory and psychophysics* (pp. xi,
455). John Wiley.

Gregory, R. L. (1966). *Eye and brain: The psychology of seeing*. McGraw-Hill.

Gregory, R. L. (2008). Emmert's Law and the moon illusion. *Spatial Vision*, *21*(3–5), 407–
420. https://doi.org/10.1163/156856808784532509

Grzeczkowski, L., Clarke, A. M., Francis, G., Mast, F. W., & Herzog, M. H. (2017). About
individual differences in vision. *Vision Research*, *141*, 282–292.
https://doi.org/10.1016/j.visres.2016.10.006

Grzeczkowski, L., Roinishvili, M., Chkonia, E., Brand, A., Mast, F. W., Herzog, M. H., &
Shaqiri, A. (2018). Is the perception of illusions abnormal in schizophrenia?
*Psychiatry Research*, *270*, 929–939.
https://doi.org/10.1016/j.psychres.2018.10.063

Halpern, S. D., Andrews, T. J., & Purves, D. (1999). Interindividual variation in human
visual performance. *Journal of Cognitive Neuroscience*, *11*(5), 521–534.

He, D., Mo, C., Wang, Y., & Fang, F. (2015). Position shifts of fMRI-based population
receptive fields in human visual cortex induced by Ponzo illusion. *Experimental
Brain Research*. https://doi.org/10.1007/s00221-015-4425-3

Himmelberg, M. M., Tünçok, E., Gomez, J., Grill-Spector, K., Carrasco, M., & Winawer, J.
(2023). Comparing retinotopic maps of children and adults reveals a late-stage
change in how V1 samples the visual field. *Nature Communications*, *14*(1), 1561.
https://doi.org/10.1038/s41467-023-37280-8

Ho, M.-L., & Schwarzkopf, D. S. (2022). The human primary visual cortex (V1) encodes
the perceived position of static but not moving objects. *Communications
Biology*, *5*, Article 1. https://doi.org/10.1038/s42003-022-03136-y

Howell, D. C. (1997). *Statistical Methods for Psychology*. Duxbury Press.

Jaeger, T., & Grasso, K. (1993). Contour lightness and separation effects in the
Ebbinghaus illusion. *Perceptual and Motor Skills*, *76*(1), 255–258.
https://doi.org/10.2466/pms.1993.76.1.255

Jaeger, T., & Klahs, K. (2015). The Ebbinghaus illusion: New contextual effects and
theoretical considerations. *Perceptual and Motor Skills*, *120*(1), 177–182.
https://doi.org/10.2466/24.27.PMS.120v13x4

Jaeger, T., Klahs, K., & Newton, D. (2014). Ebbinghaus illusions with disc figures: Effects
of contextual size, separation, and lightness. *Perceptual and Motor Skills*, *118*(3),
805–817. https://doi.org/10.2466/24.27.PMS.118k29w0

JASP Team. (2023). *JASP (Version 0.18.3)* [Computer software]. https://jasp-stats.org

Jastrzębowska, M. A., Ozkirli, A., Cretenoud, A. F., Draganski, B., & Herzog, M. H. (2023).
*Is there a neural common factor for visual illusions?* (p. 2023.12.27.573437).
bioRxiv. https://doi.org/10.1101/2023.12.27.573437

Jogan, M., & Stocker, A. A. (2014). A new two-alternative forced choice method for the
unbiased characterization of perceptual bias and discriminability. *Journal of*
*Vision*, *14*(3), 20. https://doi.org/10.1167/14.3.20

Makowski, D., Te, A. S., Kirk, S., Liang, N. Z., & Chen, S. H. A. (2023). A novel visual
illusion paradigm provides evidence for a general factor of illusion sensitivity and
personality correlates. *Scientific Reports*, *13*(1), 6594.
https://doi.org/10.1038/s41598-023-33148-5

Manning, C., Morgan, M. J., Allen, C. T. W., & Pellicano, E. (2017). Susceptibility to
Ebbinghaus and Müller-Lyer illusions in autistic children: A comparison of three
different methods. *Molecular Autism*, *8*(1), 16. https://doi.org/10.1186/s13229-
017-0127-y

Mareschal, I., Morgan, M. J., & Solomon, J. A. (2010). Cortical distance determines
whether flankers cause crowding or the tilt illusion. *Journal of Vision*, *10*(8), 13.
https://doi.org/10.1167/10.8.13

Massaro, D. W., & Anderson, N. H. (1971). Judgmental model of the Ebbinghaus illusion.
*Journal of Experimental Psychology*, *89*(1), 147–151.

Mazuz, Y., Kessler, Y., & Ganel, T. (2023). The BTPI: An online battery for measuring
susceptibility to visual illusions. *Journal of Vision*, *23*(10), 2.
https://doi.org/10.1167/jov.23.10.2

Mazuz, Y., Kessler, Y., & Ganel, T. (2024). Age-related changes in the susceptibility to
visual illusions of size. *Scientific Reports*, *14*(1), 14583.
https://doi.org/10.1038/s41598-024-65405-6

Morgan, M., Dillenburger, B., Raphael, S., & Solomon, J. A. (2012). Observers can
voluntarily shift their psychometric functions without losing sensitivity. *Attention,*
*Perception & Psychophysics*, *74*(1), 185–193. https://doi.org/10.3758/s13414-
011-0222-7

Morgan, M. J., & Glennerster, A. (1991). Efficiency of locating centres of dot-clusters by
human observers. *Vision Research*, *31*(12), 2075–2083.
https://doi.org/10.1016/0042-6989(91)90165-2

Morgan, M. J., Melmoth, D., & Solomon, J. A. (2013). Linking hypotheses underlying
Class A and Class B methods. *Visual Neuroscience*, *30*(5–6), 197–206.
https://doi.org/10.1017/S095252381300045X

Moutsiana, C., de Haas, B., Papageorgiou, A., van Dijk, J. A., Balraj, A., Greenwood, J. A.,
& Schwarzkopf, D. S. (2016). Cortical idiosyncrasies predict the perception of
object size. *Nature Communications*, *7*, 12110.
https://doi.org/10.1038/ncomms12110

Murray, S. O., Boyaci, H., & Kersten, D. (2006). The representation of perceived angular
size in human primary visual cortex. *Nature Neuroscience*, *9*(3), 429–434.
https://doi.org/10.1038/nn1641

Nauhaus, I., Busse, L., Carandini, M., & Ringach, D. L. (2009). Stimulus contrast
modulates functional connectivity in visual cortex. *Nature Neuroscience*, *12*(1),
70–76. https://doi.org/10.1038/nn.2232

Newsome, L. R. (1972). Visual angle and apparent size of objects in peripheral vision.
*Perception & Psychophysics*, *12*(3), 300–304.
https://doi.org/10.3758/BF03207209
Ni, A. M., Murray, S. O., & Horwitz, G. D. (2014). Object-Centered Shifts of Receptive
Field Positions in Monkey Primary Visual Cortex. *Current Biology: CB*.
https://doi.org/10.1016/j.cub.2014.06.003
Patten, M. L., & Clifford, C. W. G. (2015). A bias-free measure of the tilt illusion. *Journal*
*of Vision*, *15*(15), 8. https://doi.org/10.1167/15.15.8
Poom, L. (2020). Influences of orientation on the Ponzo, contrast, and Craik-O'Brien-
Cornsweet illusions. *Attention, Perception, & Psychophysics*, *82*(4), 1896–1911.
https://doi.org/10.3758/s13414-019-01953-8
Pooresmaeili, A., Arrighi, R., Biagi, L., & Morrone, M. C. (2013). Blood Oxygen Level-
Dependent Activation of the Primary Visual Cortex Predicts Size Adaptation
Illusion. *Journal of Neuroscience*, *33*(40), 15999–16008.
https://doi.org/10.1523/JNEUROSCI.1770-13.2013
Roberts, B., Harris, M. G., & Yates, T. A. (2005). The roles of inducer size and distance in
the Ebbinghaus illusion (Titchener circles). *Perception*, *34*(7), 847–856.
Sánchez-Fuenzalida, N., van Gaal, S., Fleming, S. M., Haaf, J. M., & Fahrenfort, J. J.
(2023). Predictions and rewards affect decision-making but not subjective
experience. *Proceedings of the National Academy of Sciences of the United*
*States of America*, *120*(44), e2220749120.
https://doi.org/10.1073/pnas.2220749120
Schwarzkopf, D. S., & Rees, G. (2013). Subjective size perception depends on central
visual cortical magnification in human v1. *PLoS One*, *8*(3), e60550.
https://doi.org/10.1371/journal.pone.0060550
Schwarzkopf, D. S., Song, C., & Rees, G. (2011). The surface area of human V1 predicts
the subjective experience of object size. *Nature Neuroscience*, *14*(1), 28–30.
https://doi.org/10.1038/nn.2706
Sherman, J. A., & Chouinard, P. A. (2016). Attractive Contours of the Ebbinghaus
Illusion. *Perceptual and Motor Skills*, *122*(1), 88–95.
https://doi.org/10.1177/0031512515626632
Song, C., Schwarzkopf, D. S., Kanai, R., & Rees, G. (2015). Neural Population Tuning
Links Visual Cortical Anatomy to Human Visual Perception. *Neuron*, *85*(3), 641–
656. https://doi.org/10.1016/j.neuron.2014.12.041
Song, C., Schwarzkopf, D. S., Lutti, A., Li, B., Kanai, R., & Rees, G. (2013). Effective
connectivity within human primary visual cortex predicts interindividual diversity
in illusory perception. *The Journal of Neuroscience: The Official Journal of the*
*Society for Neuroscience*, *33*(48), 18781–18791.
https://doi.org/10.1523/JNEUROSCI.4201-12.2013
Song, C., Schwarzkopf, D. S., & Rees, G. (2011). Interocular induction of illusory size
perception. *BMC Neuroscience*, *12*, 27. https://doi.org/10.1186/1471-2202-12-
27
Song, C., Schwarzkopf, D. S., & Rees, G. (2013). Variability in visual cortex size reflects
tradeoff between local orientation sensitivity and global orientation modulation.
*Nature Communications*, *4*, 2201. https://doi.org/10.1038/ncomms3201

Spearman, C. (1910). Correlation Calculated from Faulty Data. *British Journal of*
*Psychology, 1904-1920, 3*(3), 271–295. https://doi.org/10.1111/j.2044-
8295.1910.tb00206.x
Sperandio, I., Chouinard, P. A., & Goodale, M. A. (2012). Retinotopic activity in V1
reflects the perceived and not the retinal size of an afterimage. *Nature*
*Neuroscience, 15*(4), 540–542. https://doi.org/10.1038/nn.3069
Todorović, D., & Jovanović, L. (2018). Is the Ebbinghaus illusion a size contrast illusion?
*Acta Psychologica, 185*, 180–187. https://doi.org/10.1016/j.actpsy.2018.02.011
Urale, P. W. B., & Schwarzkopf, D. S. (2023). Effects of cortical distance on the
Ebbinghaus and Delboeuf illusions. *Perception, 52*(7), 459–483.
https://doi.org/10.1177/03010066231175014
von Helmholtz, H. (1924). *Helmholtz's treatise on physiological optics, Vol. 1, Trans.*
*From the 3rd German ed* (pp. xxi, 482). Optical Society of America.
https://doi.org/10.1037/13536-000
Weintraub, D. J. (1979). Ebbinghaus illusion: Context, contour, and age influence the
judged size of a circle amidst circles. *Journal of Experimental Psychology.*
*Human Perception and Performance, 5*(2), 353–364.
https://doi.org/10.1037//0096-1523.5.2.353
Yildiz, G. Y., Sperandio, I., Kettle, C., & Chouinard, P. A. (2021). The conceptual
understanding of depth rather than the low-level processing of spatial
frequencies drives the corridor illusion. *Vision Research, 181*, 21–31.
https://doi.org/10.1016/j.visres.2020.11.012
1180
1181
1182

**Figure S1.** A. Scatter plots underlying the correlation matrix of data from Experiment 1 shown in Figure 1H. Each dot plots illusion magnitude and/or discrimination bandwidth for an individual observer from a given condition. Pearson correlation statistics are shown at the top of each panel (p-values rounded to three decimal points). Black line and shaded area denote linear regression and its 95% confidence interval. Black dots: Naïve observers. Red dots: Non-naïve (author) observers. B. Correlation matrix from Experiment 1 but excluding non-naïve (author) observers. Correlations compared the magnitudes for the four illusions, and discrimination abilities. All conventions as in Figure 1H. Ebb: Ebbinghaus. Del: Delboeuf. Mue: Mueller-Lyer. Pon: Ponzo. Dis: disc width. Lin: line length.

**Figure S2.** A. Scatter plots underlying the correlation matrix of data from Experiment 2 shown in Figure 2E. Each dot plots illusion magnitude for an individual observer from a given task and stimulus condition. Black line and shaded area denote linear regression and its 95% confidence interval. Pearson correlation statistics are shown at the top of each panel (p-values rounded to three decimal points). Black dots: Naïve observers. Red dots: Non-naïve (author) observers. Adj: Adjustment task. Unc: Unconstrained-2AFC task. F&F: Fast-2AFC task with fixation. B. Correlation matrix from Experiment 2 but excluding non-naïve (author) observers. All conventions as in Figure 2E.

**Figure S3.** Scatter plots of the test-retest reliability for illusion magnitudes in each task and stimulus condition from Experiment 2. Each black dots denotes the illusion magnitude (point of subjective equality, PSE) estimated for a given observer from the second half of trials against the first half. In the Adjustment tasks, data were split into odd and even numbered trials. In the 2AFC tasks, separate psychometric curves were fit to the two independent, interleaved staircases. $R_{rel}$ shows the test-retest correlation extrapolated for the full data set using Spearman-Brown prophecy formula (see Materials and Methods). Black line and shaded area denote linear regression and its 95% confidence interval.