Title

Pseudomonas taxonomic and functional microdiversity in the wheat rhizosphere is cultivar-

dependent and links to disease resistance profile and root diameter

Authors

Courtney Horn Herms[1], Rosanna Catherine Hennessy[1], Frederik Bak[1,2], Ying Guan[1], Patrick Denis

Browne [1], Tue Kjærgaard Nielsen [1], Lars Hestbjerg Hansen [1], Dorte Bodin Dresbøll[3], Mette Haubjerg

Nicolaisen[1*]

[1]Section for Microbial Ecology and Biotechnology, Department of Plant and Environmental Sciences,

University of Copenhagen, Thorvaldsensvej 40, Frederiksberg C, Denmark

[2]Bioresources Unit, AIT Austrian Institute of Technology, Konrad-Lorenz-Straße 24, Tulln, Austria

[3]Section for Crop Sciences, Department of Plant and Environmental Sciences, University of

Copenhagen, Højbakkegård Allé 13, Taastrup, Denmark

* Corresponding Author. E-mail: meni@plen.ku.dk. Telephone: +4535332649. Address:

Thorvaldsensvej 40, 1871 Frederiksberg C, Denmark

Abbreviations

ASV: amplicon sequence variant

BGCs: Biosynthetic gene clusters

CLPs: cyclic lipopeptides

CSV: culture sequence variant

NRPS: non-ribosomal peptide synthetase

RiPP: ribosomally synthesized and post-translationally modified peptide

Abstract

Diversity within lower taxonomic units in microbial communities is a key trait, giving rise to

important ecological functions. In the rhizosphere, these functions include disease suppression and

pathogen inhibition. However, limited effort has been given to exploring intragenus microdiversity in

an increasingly homogenous agricultural system. Through an integrative approach combining

culture-dependent and -independent methods, we explore the rhizosphere *Pseudomonas*

pangenome and demonstrate cultivar-dependent taxonomic and functional microdiversity between

two closely related modern winter wheat cultivars. A *Fusarium*-resistant cultivar demonstrated

increased *Pseudomonas* taxonomic diversity but not biosynthetic diversity when compared to the

susceptible cultivar, coinciding with a thinner root diameter of the resistant cultivar. We found

enrichment of *Pseudomonas* isolates capable of antagonizing *Fusarium* as well as chitinase-encoding

genes and pyoverdine gene clusters in the resistant cultivar. Across closely related *Pseudomonas*

isolates from the two cultures, there were differences in genomic content and biosynthetic gene

clusters. Ultimately, we highlight the need for fine-scale analysis to uncover the hidden

microdiversity within rhizosphere *Pseudomonas*.

1.   Introduction

Plants and soil microorganisms interact through such a tight collaboration that the plant and its

rhizosphere microorganisms are considered a holobiont (Vandenkoornhuyse et al., 2015). The

microbiome in the rhizosphere generally assembles into a stable community comprised primarily of

the phyla Actinobacteria, Bacteroidetes, Firmicutes, and Proteobacteria, which harbor key functions

important for plant health (Bai et al., 2022). But, at the order and genus level, plant microbiome

communities can differ because of plant type, root exudation, and age (Qu et al., 2020), leading to

functional community differences between closely related crops (Mendes et al., 2018).

Diversity within genera and species, i.e., microdiversity (Moore et al., 1998), plays an important role

in microbial ecology (García-García et al., 2019). Bacterial species arise when populations split into

ecologically distinct fractions allowing for specific niche adaptations (Koeppel et al., 2013). In this

case, genes involved in functions such as iron acquisition and biofilm formation become part of

lineage-specific gene pools (Shalev et al., 2021). Thus, increased diversity within genera and species

contributes to ecological interactions through resource and niche competition as well as direct

antagonism (Jousset et al., 2011; Hu et al., 2016; Yang et al., 2017; Li et al., 2023; Spragge et al.,

2023).

Despite this, only few studies to date have addressed microdiversity of rhizosphere microbiomes

(Mauchline et al., 2015; Oni et al., 2019; Pacheco-Moreno et al., 2021, 2024). These studies

demonstrate the potential genomic and function diversity yet to be discovered in the rhizosphere

environment. The lack of fine taxonomic resolution in microbiome studies of the plant holobiont

may be a consequence of short-read amplicon-based analyses that conceal diversity and functional

differences between closely related microbes (Chiniquy et al., 2021; Edgar, 2018; Wang et al., 2022).

Advances in sequencing technologies are increasing amplicon read lengths, allowing for better

culture-independent identification of soil microbes (Stevens et al., 2023), but isolation efforts, whole

genome sequencing, and pangenome analysis remain the gold standard for the identification of

microdiversity (Huang et al., 2023; Zhang et al., 2024).

The small body of research on rhizosphere microdiversity has focused on comparing cultivars distinct

by their breeding history or cropping strategies, but there is limited attention to closely related

modern cultivars that demonstrate their own microdiversity in terms of root exudation and root

morphology (Iannucci et al., 2021). Root exudation is a plant trait that is key to contributing to

microbial community diversity (Feng et al., 2023; Sasse et al., 2018; Yang et al., 2023), but it alone

cannot fully explain rhizosphere microbiome assembly (Takamatsu et al., 2023). The physical

landscape is also suggested as a driver of microbial community diversity (Dubey et al., 2021). The

physical and consequent chemical landscapes implicated by different root phenotypes have the

potential to modulate root-microbe interactions (Herms et al., 2022). Specifically, there is growing

evidence for a link between thinner root diameter and bacterial diversity in the rhizosphere for both

woody and herbaceous species (Fleishman et al., 2023; King et al., 2023; Luo et al., 2021; Pervaiz et

al., 2020; Saleem et al., 2018; Zai et al., 2021). However, to date, this relationship has not been

examined in cereals, nor have microbial microdiversity and root diameter been related.

A genus of particular interest for the study of microdiversity in the rhizosphere is *Pseudomonas*. This

ubiquitous soil genus boasts high intrinsic intragenus genomic diversity (Girard et al., 2021; Loper et

al., 2012; Lopes et al., 2018) and a wide assortment of secondary metabolites (Gavriilidou et al.,

2022; Gross and Loper, 2009; Gu et al., 2020). Additionally, *Pseudomonas* is a key member of

disease-resistant microbiomes (Hong et al., 2023; Lv et al., 2023; Qiu et al., 2022) thanks to their

production of a wide spectrum of bioactive secondary metabolites such as siderophores,

phenazines, cyclic lipopeptides (CLPs), and chitinases (Mauchline and Malone, 2017; Pacheco-

Moreno et al., 2021).

In the present work, we explored the root-associated *Pseudomonas* taxonomic and functional

microdiversity of closely related and commercially available modern cultivars of winter wheat

(*Triticum aestivum* L.) grown in agricultural soil. By choosing two cultivars with contrasting *Fusarium*

culmorum resistance profiles: the resistant Sheriff cultivar and the susceptible Heerup cultivar

(Jørgensen et al., 2023, 2020), we further aimed at investigating whether a link between

microdiversity and cultivar resistance to soil-borne fungal infections could be observed. We

undertook an integrative approach combining culture-dependent and culture-independent

pangenome analyses coupled to root morphological analysis. We expected the taxonomic and

functional microdiversity to be different between the two cultivars. Specifically, we hypothesized

that the *Fusarium*-resistant cultivar, Sheriff, harbors a more taxonomically and functionally diverse

soil *Pseudomonas* community. Second, we hypothesized that pseudomonads from the Sheriff

cultivar inhibit *F. culmorum* growth and harbor a greater abundance of genes that may be

responsible for antagonizing *F. culmorum*. Lastly, we hypothesized that the expected increase in

Sheriff *Pseudomonas* microdiversity would coincide with a smaller root diameter of the Sheriff root

system.

2. Materials and Methods

2.1 Wheat cultivation and sampling

Two winter wheat (*Triticum aestivum* L.) cultivars, Heerup and Sheriff, were chosen based on their

documented resistance to *F. culmorum* in the field and were supplied by Sejet Plant Breeding

(Horsens, Denmark). One seed of each cultivar was sown in a 24 x 7 cm PVC pot filled with soil mixed

with sand (DANSAND, Denmark; filter sand no. 2) in a 2:1 ratio. The soil was a sandy loam collected

from the plough layer (0-25 cm) in the University of Copenhagen's Long Term Nutrient Depletion

Experiment located at Højbakkegård in Høje Taastrup, Denmark (van der Bom et al., 2017). The soil

had received mineral NPK fertilizer at a rate of 120 kg nitrogen, 20 kg phosphorus and 120 kg

potassium $ha^{-1}y^{-1}$ for the past 25 years and contained 170 g kg^{-1} clay, 174 g kg^{-1} silt, 336 g kg^{-1} fine

sand, 255 g kg^{-1} coarse sand, and 40 g kg^{-1} organic matter. The soil had a pH(CaCl$_2$) of 5.4, an Olsen-P

content of 11.4 mg kg^{-1}. The water holding capacity (WHC) was 33%. The soil was air-dried and

sieved to 8mm prior to the experiment.

The plants were grown under controlled greenhouse conditions in three stages: an 18-day

germination period (19°C day/15°C night; 16h day/8h night; light intensity 300 µE), followed by a 13-

week vernalization period (6°C day/4°C night; 8h day/16h night; light intensity 150 µE), and finally a

17-week growth period to maturity (19°C day/15°C night; 16h day/8h night; light intensity 500 µE).

Pots were regularly rotated and watered to 70% WHC by weighing.

Three different time points were sampled from three independent setups following the above

method (Methods Fig 1). After four weeks, a set of plants were sampled for root morphology

scanning. A second set of plants were sampled for *Pseudomonas* isolation at first node emergence

(T1). A third set of plants were used for two rounds of long-read 16S rRNA amplicon sequencing, one

at first node emergence (T1) and one at flag leaf emergence (T2).

2.2 *Pseudomonas* library building

Three replicate plants were sampled at first node emergence (T1) to obtain *Pseudomonas* colonies

(Methods Fig 1). To obtain rhizoplane samples, one gram (corresponding to approximately 4 cm) of

root was washed in 25 mL sterile PBST (10% 10x PBS, 0.1% Tween 80) and subsequently placed in 25

mL of fresh, sterile PBST. The roots were sonicated in a water bath for two minutes to release

rhizoplane soil and microorganisms. To isolate *Pseudomonas* species, 50 μL of undiluted and a 1:10

dilution in PBST (PBS, 0.1% Tween 80) of the rhizoplane sample was plated onto Gould's S1 media

(Gould et al., 1985) to select for *Pseudomonas* isolates. After incubation at room temperature for

two to four days, single colonies were streaked for purification first on Gould's S1 media and a

second time on Luria-Bertani (LB) agar (Miller, 1972). Pure isolates were maintained as frozen stocks

(15% glycerol w/w).

2.3 16S rRNA gene sequencing of cultured isolates

A modified version of colony PCR (von Stein et al., 1997) was used to amplify the full-length 16S

rRNA gene of the isolates in the strain library. A 30 μL PCR reaction mixture was prepared: 15 μL

Master Mix (Supreme NZYTaq II 2x Master Mix, NZYTech, Portugal), 0.6 μL of 10 μM forward and

reverse primer (27F, 1492R; Weisburg et al., 1991), 11.8 μL Sigma water, and 2 μL 1:1000 diluted

overnight culture. The cycling conditions were as follows: preliminary boiling step of 96°C for 5 min

followed by 25 cycles of denaturing at 98°C for 10 s, annealing at 57°C for 30 s, extension at 72°C for

45 s, and a final extension of 10 min at 72°C. Successful amplification of the 16S rRNA gene was

visualized on 1% agarose gels. PCR purification and Sanger sequencing of the PCR product was

completed by Eurofins (Germany). The ends of the reads were trimmed by Eurofins based on quality

scores. Reads of short length (<900 base pairs (bp)) were manually discarded using CLC Genomics

Workbench (v. 12.0.03) (QIAGEN).

Sequences were oriented to the same strand using USEARCH (Edgar, 2010) and the RDP training set

(v. 18) as reference and then aligned using MUSCLE (Edgar, 2004). Gaps at the end of the aligned

sequences were removed using Gblocks (Castresana, 2000). Sequences were clustered into

sequence variant cluster of 100% identity using the USEARCH *cluster_fast* command (Edgar, 2010).

The 16S rRNA sequences from each sequence variant cluster were submitted to NBCI BLASTn for

alignment to the 16S rRNA database (Altschul et al., 1997).

The 16S rRNA genes of the strain library isolates are available in Genbank under accession numbers

PP894319-PP894691.

2.4 Phenotypic assays

The wheat fungal pathogen *F. culmorum* was sporulated in Mung bean media (Ilmi et al., 2019) (20

grams boiled mung beans, strained, in 1 L distilled water). A high-throughput, agar-based

confrontation assay was developed to screen the strain library for antifungal activity. Briefly, the

Pseudomonas isolates were inoculated onto a pre-germinated lawn of 1.45×10^3 *F. culmorum* spores

on 1/5 potato dextrose agar (BD DIFCO, Denmark). After three days of co-inoculation at room

temperature, the isolates were scored for their ability or inability to inhibit *F. culmorum* growth. A

positive result required a clearance zone of at least 1 mm around a colony. *Serratia inhibens* S40

(Hennessy et al., 2020) was used as the positive control.

A high throughput drop collapse assay (Bodour and Miller-Maier, 1998) was used to screen the strain

library for biosurfactant production. The isolates were grown in 1.5 mL liquid LB media for four days

in a deep-well 96-well plate at 28°C while shaking (200 rpm). To test for biosurfactant production, 40

µL of culture was pipetted onto Parafilm, and at least two minutes passed before the droplet size was

measured. Water was used as the negative control, and *P. fluorescens* SBW25, which produces the

surfactant viscosin (Alsohim et al., 2014), was used as the positive control.

2.5 Genome sequencing and bioinformatics pipeline

A representative isolate from each unique culture sequence variant (CSV) clusters identified from

the strain library via 16S rRNA gene sequencing were subject to full genome sequencing. CSV

clusters containing many isolates were sequenced deeper, proportional to the size of the CSV cluster

and how many isolates in the cluster originated from each cultivar. Genomic DNA was prepared from

overnight cultures using the QIAGEN Genomic DNA Handbook using the QIAGEN Buffer Kit

(Germany) or the Genomic Mini Ax Bacteria 96-Well kit (A&A Biotechnology, Poland) according to

the manufacturer's instructions. Quality was checked using a NanoDrop ND-1000

spectrophotometer (Thermo Fisher Scientific, Carlsbad, CA, USA) prior to library building. Genomic

DNA was prepared for sequencing using the Rapid Barcoding Sequencing Kit (SQK-RBK004) or the

Rapid Sequencing gDNA barcoding kit (SQK-RBK110.96) and sequenced on MinION and PromethION

flow cells (v. 9.4.1) (Oxford Nanopore Technologies, United Kingdom). Reads were basecalled with

Guppy (v. 6.2.1) using the "super accuracy" dna_r9.4.1_450bps_sup.cfg basecalling model. Adapter

sequence and barcodes were trimmed from basecalled reads with Porechop (v. 0.2.4). Genomes

were assembled using Flye (v. 2.9-b1774) (Lin et al., 2016; Kolmogorov et al., 2019), polished with

Medaka (v. 1.7.1) (https://nanoporetech.github.io/medaka), and annotated using Prokka (v. 1.14.6)

(Seemann, 2014) using default parameters. The quality of the genomes was assessed using CheckM

(v1.1.10) (Parks et al., 2015). Genomes not passing thresholds for completeness (> 90.0%) and

contamination (< 2.0%) were removed from future analyses. Dereplication with dRep (v. 3.4.5) (Olm

et al., 2017) and the Genome Taxonomy Database Toolkit (v. 2.3.0) (Chaumeil et al., 2020; Parks et

al., 2020) were used to delineate and identify species and strains. In dRep, 95% average nucleotide

identity (ANI) was chosen to delineate species, and 99.55% ANI was used to delineate strains.

The annotated genomes were screened for secondary metabolite prediction with antiSMASH (v. 7.0)

(Medema et al., 2011) using the standard cutoff of 0.3, *glocal* mode, and the *include_singletons* flag.

BiG-SCAPE (v. 1.1.7) (Navarro-Muñoz et al., 2020) and the MiBIG database (v. 3.1) (Terlouw et al.,

2023) were used for biosynthetic gene cluster family analysis. Anvi'o (v. 7.0) (Eren et al., 2021) was

used to assign functions using NCBI's Clusters of Orthologous Groups (COGs) database (Tatusov et

al., 2000), generate a pangenome, and quantify functional enrichment between sets of genomes

(Shaiber et al., 2020).

The fully assembled genomes are available under BioProject accession number PRJNA1108081.

2.6 Long-read 16S rRNA gene amplicon sequencing

For long-read 16S rRNA amplicon sequencing, rhizoplane samples were obtained as described above

at first node (T1) (n = 5 for Heerup and Sheriff) and flag leaf emergence (T2) (n = 10 for Heerup and n

= 9 for Sheriff) (Methods Fig 1). After the sonication step, the rhizoplane samples were flash frozen

in liquid nitrogen and freeze-dried. DNA was extracted using the FastPrep-24 5G bead-beating

system (MP Biomedicals, Irvine, CA, USA) at 6.0 m/s for 40 s and the FastDNA SPIN Kit for soil (MP

Biomedicals) according to the manufacturer's instructions.

PCR reactions were prepared with 32.5 µL Sigma water, 10 µL SuperFi Buffer (Thermo Fischer), 0.5

µL Platinum SuperFi DNA Polymerase (Thermo Fischer), 1 µL 10 mM dNTPs, 2.5 µL each of 10 nM

forward and reverse primer (TAG Copenhagen, Denmark), and 1 µL DNA template diluted to 5 ng/µL.

Pseudomonas-specific primers for amplifying 969 bp of the 16S rRNA gene (position 289-1258) were

used (forward primer 5'-GGTCTGAGAGGATGATCAGT-3' and reverse primer 5'-

TTAGCTCCACCTCGCGGC-3') (Widmer et al., 1998). ZymoBIOMICS Microbial Community DNA was

used as a sequencing control (Zymo Research, Irvine, CA, USA). The cycling conditions were as

follows: preliminary denaturation step of 98°C for 30 s followed by 30 cycles of denaturing at 98°C

for 10 s, annealing at 63°C for 30 s, extension at 72°C for 45 s, and a final extension of 10 min at

72°C. Successful amplification was visualized on 1% agarose gels. PCR products were purified using

AMPure XP beads at a 0.6 ratio (Beckman Coulter Inc., Brea, CA, USA). Amplicons were prepared for

sequencing using a native barcoding kit (Oxford Nanopore SQK-NBK114.96) and sequenced on a

Minion (Oxford Nanopore) on an R10.4.1 flow cell (Oxford Nanopore FLO-MIN114). Raw nanopore

data was basecalled with Guppy (v. 6.3.9) using the "super accuracy" basecalling model

dna_r10.4.1_e8.2_260bps_sup-v4.1.0.

Adapters were trimmed from Nanopore reads using porechop (v. 0.2.4)

(https://github.com/rrwick/Porechop) with the following options: adapter_threshold 95,

extra_end_trim 0, end_threshold 95, middle_threshold 95, extra_middle_trim_good_side 0,

extra_middle_trim_bad_side 0, min_split_read_size 800. The options were used to filter out overly

short inserts and to leave the primer sequences in the trimmed reads. Following this, primer

sequences were identified, and the intervening sequences (inserts) were extracted using

strip_degen_primer_deep (github.com/padbr/asat) and inserts between 880 and 970 nt were

retained. High quality inserts (Phred quality ≥ 30) were identified using Nanofilt (De Coster et al.,

2018). High quality inserts were then dereplicated and unique sequences counted using the

fastx_uniques utility of usearch (v. 11.0.667) (Edgar, 2010), and zero-width amplicon sequencing

variants (ASVs) were inferred using unoise3 (Edgar, 2016). All read inserts trimmed of adapters and

primers, within the 880 to 970 nt range, but not quality filtered, were used to create a ASV table

using the otutab utility of usearch.

The raw reads are available under BioProject accession number PRJNA1108081.

2.7 Root morphology scanning

To measure root morphology of the two cultivars, 4-week old plants were removed from the pots

and all soil was rinsed from the roots using tap water (n = 9 for Heerup and n = 10 for Sheriff).

Samples of 1 cm length were taken 1 cm (top), 11 cm (middle), and 18 cm (bottom) from the top of

each root (Methods Fig. 1). Images of each section of root were photo-scanned using the Epson

Perfection V700 scanner (8-bit gray scale, 600 dpi), and root diameter was measured using

RhizoVision (v. 2.0.3) using "broken roots" analysis mode (Seethepalli et al., 2021). To remove debris

and artifacts from the analysis, a maximum object size was set at 10 mm ^{2}, and edge smoothing was

set at a threshold of three. After segmentation, root pruning was performed with a threshold of 50.

Root diameter ranged were manually set at 0-0.2 mm, 0.2-0.3 mm, 0.3-0.4 mm, 0.4-0.5 mm, 0.5-1.0

mm, and 1.0-1.5 mm. After RhizoVision analysis, the percentage of the root system in each of

diameter class of the three root sections were averaged to obtain a single percentage for each plant

replicate.

2.8 Statistical analysis

All statistics were performed in R (v. 4.3.1). The rarefaction plot was generated using vegan (v. 2.6-4)

(Oksanen et al., 2022). Calculating the effect of cultivar on antifungal activity, biosurfactant

production, and unique isolates was performed by Fischer's Exact test. When comparing means, all

data were tested for normality using the Shapiro-Wilk test. The two-sided Wilcoxon Rank Sum test

was used in place of a two-sided unpaired Student T test when normality could not be assumed. P

values were adjusted for multiple testing as appropriate using the Benjamini & Hochberg method

(Benjamini and Hochberg, 1995). Plots were made using ggplot2 (v. 3.4.2).

Long-read amplicon sequencing data were analyzed using Qiime2 (Caporaso et al., 2010).

Normalization of the ASV table was done by rarefying with all ASVs. Following this, the ASV table was

filtered to keep only features classified in the *Pseudomonas* genus. Qiime2 was then used to derive

values for α-diversity (richness, Chao1, and Shannon) and β-diversity (Bray-Curtis dissimilarity). The

Shannon diversity values were converted to effective Shannon diversity by raising two to the power

of the Shannon diversity. Composition data and α-diversity metrics were compared between groups

of samples using the Mann-Whitney U-test to test for significance. ASVs were omitted from

composition tests if the ASV is not seen in at least 50% of the samples in one of the groups. Anosim

(Clarke, 1993), with 999 permutations, was used to test if two or more groups of samples differed

from each other.

3. Results

3.1 Culture-based screening indicates a more diverse and antifungal Sheriff *Pseudomonas*

microbiome

A strain library of 373 pseudomonads comprising 195 isolates from Heerup and 178 isolates from

Sheriff was successfully isolated. Sanger sequencing of the 16S rRNA gene confirmed all 373 isolates

as *Pseudomonas*. Clustering at 100% identity across a minimum of 900 bp and an average of 1114 bp

of the 16S rRNA gene resulted in 65 unique culture sequence variant (CSV) clusters, and BLASTn

alignment to the NBCI 16S rRNA gene database revealed 17 potential *Pseudomonas* species (Table

S1). The most abundant species in the library was identified as *P. kilonensis* with 143 isolates,

followed by *P. extremaustralis* and *P. baetica* with 86 and 36 isolates, respectively. Three and seven

species and 17 and 36 CSVs were uniquely recruited to the Heerup and Sheriff cultivar, respectively

(Figure 1). Additionally, the Sheriff cultivar harbored significantly more CSVs than the Heerup

cultivar; 33 CSVs were found in Heerup and 59 CSVs in Sheriff (Fisher's exact test, p = 0.0052).

However, the rarefaction curve demonstrated that the full *Pseudomonas* diversity was not sampled

via isolation (Figure S1).

To assess functional microdiversity within the isolated *Pseudomonas*, a high-throughput antifungal

screen and drop-collapse assay were performed. Across both cultivars, 33% (n = 129) of strains

inhibited growth of the fungal pathogen *F. culmorum*, whereas 17% (n = 67) of strains were positive

for biosurfactant activity. The high-throughput phenotypic screens showed that antifungal isolates

were enriched in the Sheriff strain library (40%) compared to Heerup strain library (27%) (Fisher's

exact test, p = 0.0030), while the proportion of biosurfactant producers did not differ between

Sheriff (20%) and Heerup (14%) (Fisher's exact test, p = 0.0603) (Fig. 2).

3.2 Targeted long-read 16S rRNA amplicon sequencing supports *Pseudomonas* microdiversity

and antifungal potential in Sheriff cultivar.

To confirm Sheriff's enrichment of *Pseudomonas* microdiversity in a culture-independent manner,

long-read 16S rRNA amplicon sequencing of the rhizoplane sample was performed on plants from a

separate experiment from the culturing. Sampling was performed at the same time point as the

culturing (at first node emergence, T1) (Heerup and Sheriff, n = 5) and one month after the culturing

time point (at flag leaf emergence, T2) (Heerup n = 10, Sheriff n = 9).

Prior to rarefying, 373,093 reads were assigned to 707 ASVs, with a range of 2,014 to 37,612 reads

per sample. Seventy of the 707 ASVs were classified within the *Pseudomonas* genus. Exact matches

to the 16S rRNA genes of the 65 unique CSVs were found in the amplicon data, but due to the length

of the amplicon at 880-970 bp, many 16S rRNA gene sequences from the CSVs were grouped

together into one ASV. Twenty-two additional, uncultured *Pseudomonas* ASVs were identified.

However, the full *Pseudomonas* diversity and richness was still not captured with amplicon

sequencing, especially at T2 (Fig S2A-B).

At T1, 67% to 91% of reads per sample were classified as *Pseudomonas* with a median of 81% (Fig

S2C). One uncultured pseudomonad, ASV7, dominated all samples at this time point and contributed

to a low percentage of cultured *Pseudomonas* reads captured by sequencing (Fig S2D). There was

also no significant difference in *Pseudomonas* diversity nor any ASVs differentially abundant

between the two cultivars at T1 (data not shown).

At T2, ASV7 decreased 95-fold in relative abundance, coinciding with a drop in *Pseudomonas* reads

per sample, with a range of 2% to 43% with a median of 5% (Fig S2C) and an increased percentage of

cultured *Pseudomonas* reads (Fig S2D).

At T2, Sheriff demonstrated a higher diversity of *Pseudomonas* than Heerup (Mann-Whitney U-test,

p = 0.00013), which aligned with the conclusion from culturing. One uncultured ASV and three ASVs

that were exact sequence matches to the 16S rRNA sequence of five isolates were significantly

enriched in the Sheriff cultivar (Table 1A). Within the ASVs matching to cultured isolates that were

enriched in the Sheriff cultivar at T2, two of the isolates showed antifungal activity against *F.*

culmorum in vitro.

Across T1 and T2, three ASVs that were exact sequence matches to the 16S rRNA sequence of 12

isolates were significantly enriched in the Sheriff cultivar compared to the Heerup cultivar (Table 1B).

Within the ASVs matching to cultured isolates that were enriched in the Sheriff cultivar across T1

and T2, seven of the isolates were antifungal against *F. culmorum in vitro.* At no time point were

there any ASVs enriched in the Heerup cultivar.

3.3 Genome sequencing highlights *Pseudomonas* microdiversity and identifies chitinase as

potential driver for enriched *Pseudomonas* antifungal activity in Sheriff cultivar

For a more detailed analysis on functional microdiversity, isolates were chosen for full-genome

sequencing based on 16S rRNA gene phylogeny. Briefly, a representative isolate of each of the 65

unique CSVs was sequenced, and large CSV clusters were sequenced deeper, proportional to the size

of the CSV cluster and how many isolates in the CSV cluster originated from each cultivar. A total of

112 isolates were fully genome sequenced of which 53 were isolated from Heerup and 59 were

isolated from Sheriff.

Dereplication (Olm et al., 2017) and the Genome Taxonomy Database (Chaumeil et al., 2020)

identified 21 species and 48 strains in the library (Table S1). Six species and 14 strains were unique to

Heerup while 10 species and 31 strains were unique to Sheriff. Hence, only five species were isolates

from both cultivars. Gene clusters generated based on amino acid sequence similarity were

generated using Anvi'o (Eren et al., 2021). In total, 14,769 gene clusters present in at least two

genomes were identified. The core genome, defined as genes present in all 112 genomes, consisted

of 2,322 gene clusters.

Gene cluster analysis revealed differences in genomic content among isolates within the same

Pseudomonas species (Fig 3A). This genomic mirodiversity was, in some instances, influenced by

cultivar of origin. In the case of the six isolates identified as *P. brassicacearum* strain R, three loci

composed of 65 gene clusters, 6 gene clusters, and 9 gene clusters, respectively, were present only

in the four isolates from the Heerup cultivar while absent in the two isolates from the Sheriff cultivar

(Fig 3B). In this same taxon, a locus of 24 gene clusters was present only in the Sheriff-isolated

strains (Fig 3B); this locus was predicted to be associated with the synthesis of flagella (Table S2).

Additionally, there is a locus of 76 gene clusters in *P. simiae* that were only present in isolates

cultured from the Heerup cultivar (Fig 3C).

In order to link genomic microdiversity to functional potential, enrichment scores for COG20

pathways, categories, and functions were calculated between isolates cultured from Heerup and

Sheriff using Anvi'o (Eren et al., 2021). There were no differentially enriched COG20 pathways.

Twenty-nine COG20 categories were differentially enriched between strains originating from the two

cultivars (adjusted *q*-value < 0.05): 15 in Heerup and 14 in Sheriff (Table S3). The most common

differentially abundant COG20 categories between the two cultivars were signal transduction

mechanisms (n = 10), secondary metabolites biosynthesis, transport and catabolism (n = 7), and cell

membrane biogenesis (n = 7). A total of 213 COG20 functions were differentially enriched between

strains originating from the two cultivars (adjusted *q*-value < 0.05): 104 in Heerup and 109 in Sheriff

(Table S4). Notably, chitinase family GH19 (accession: COG3179) was enriched in isolates originating

from the Sheriff cultivar (adjusted *q*-value = 9.2E-6). Of the remaining differentially enriched COG20

functions, no others were known to be directly implicated in antifungal activity (Table S3).

3.4 Genome mining reveals *Pseudomonas* functional partitioning between Heerup and Sheriff via cultivar-specific species enrichment

We mined the genomes for biosynthetic gene clusters (BGCs) to examine the biosynthetic potential of the *Pseudomonas* strain library. Screening with antiSMASH (Medema et al., 2011) and analysis via BiG-SCAPE (Navarro-Muñoz et al., 2020) revealed 1315 BGCs in the strain library that grouped into 137 BGC families of 21 product classes. The most common class of BGC in the strain library was ribosomally synthesized and post-translationally modified peptides (RiPP, n = 450) followed by NRPS (n = 378). The most common product class was RiPP-like (n = 287), followed by redox co-factors (n = 118), NRPS-like (n = 117), and NRPS (n = 111). The BGC families identified in the strain library were mapped to the MiBIG database, and this revealed six known products (Table 2).

To test the hypothesis of increased functional diversity and enrichment of *Fusarium* antagonism in Sheriff pseudomonads, we determined BGC families that were differentially abundant in Heerup and Sheriff. Overall, there were not a difference in average number of BGCs in isolates from the two cultivars (two-sided Wilcoxon Rank Sum test, $p > 0.05$) (Fig 4A), but 28 of the 137 BGC families identified in the strain library were differentially abundant between the two cultivars (two-sided Wilcoxon Rank Sum test, fdr-adjusted $p < 0.05$) (Fig 4B). Nineteen and nine BGC families were enriched in Heerup and Sheriff, respectively, spanning 11 different product classes (Fig 4B). The only known BGC family differentially enriched between the two cultivars was BGC family 3049, an NRPS encoding a Pf-5 pyoverdine enriched in Sheriff (Table 2). Two CLPs were identified in the strain library, viscosin and lokisin, but they were not found to be differentially abundant between Heerup and Sheriff (Table 2).

Our next objective was to uncover the factors underpinning the differing abundance of BGC families within the Heerup and Sheriff *Pseudomonas* communities. First, we determined the effect of cultivar of origin on the number of BGCs per genome in each species (Figure 5A). Five species were found in both cultivars (*P. asgharzadehiana*, *P. simiae*, *P. neuropathica, P. siliginis*, and *P. brassicacearum*);

however, only three of these species were found more than once on both cultivars (*P.*

asgharzadehiana, *P. neuropathica*, and *P. brassicacearum*). Thus, statistical analysis was only done

on these three species. The number of BGCs per genome was only affected by cultivar in the species

P. brassicacearum (two-sided Wilcoxon Rank Sum test, *p* = 0.0140) (Figure 5A). Due to this

unconvincing result of the number of BGCs per genome being dependent on cultivar of origin in only

one species out of 22, we next analyzed the distribution of each differentially abundant BGC family

across the strain collection (Figure 5B). This revealed that the variation in BGC family abundance was

primarily driven by the enrichment or uniqueness of specific *Pseudomonas* species that were

harboring many different BGC families to a single wheat cultivar.

3.5 Root scanning relates Sheriff *Pseudomonas* community to thinner root diameter

Culture-dependent and culture-independent 16S rRNA gene sequencing indicated that the Sheriff

cultivar harbors a more diverse *Pseudomonas* population. To test the hypothesis that a more diverse

Pseudomonas community would coincide with a thinner root diameter, we scanned the roots of

four-week old Heerup and Sheriff plants and measured the root diameter of the system. Sheriff had

a higher percentage of roots under 0.4 mm in diameter, while Heerup had a higher percentage of

roots above 0.5 mm (Figure 6). This contributed to the thinner average root diameter of Sheriff (0.35

± 0.03 mm) compared to Heerup (0.40 ± 0.03 mm) (two-sided Student T test, *p* = 0.0026).

4.   Discussion

Our study provides an in-depth insight into the microdiversity of *Pseudomonas* communities in two

commercially available modern cultivars of winter wheat with contrasting susceptibility to the fungal

pathogen *F. culmorum.* With 373 isolates and 112 genomes, this is to our knowledge the largest

study to characterize *Pseudomonas* microdiversity. Additionally, we provide a first glance into the

Pseudomonas pangenome of two closely related modern cultivars, demonstrating that microbial

microdiversity remains despite increasing homogeneity of agricultural systems (Elouafi, 2024).

In field-grown wheat rhizospheres, *Pseudomonas* are present in low abundances (Mauchline et al.,

2015; Simonin et al., 2020), but their ability to dictate microbiome assembly (Garrido-Sanz et al.,

2023; Getzke et al., 2023) and function (Hong et al., 2023; Lv et al., 2023) suggests their importance

regardless. The results of our work indicate that the microdiversity of this genus leaves even more to

be discovered. The taxonomic breadth in a single niche can be extensive, with our study revealing 22

species and 48 strains from 373 isolates as well as 70 ASVs from long-read 16S rRNA amplicon

sequencing. Previous work in cocoyam and potato both identified seven phylogenetic groups from

134 and 69 *Pseudomonas* isolates, respectively (Oni et al., 2019; Pacheco-Moreno et al., 2021).

Interestingly, our low-abundance ASVs were different from our low-abundance isolates, indicating

unique biases in both culturing and sequencing. Thus, we suggest that both approaches are needed

to capture the full picture of *Pseudomonas* microdiversity in the rhizoplane.

In our first hypothesis, we predicted that the *F. culmorum* resistant cultivar, Sheriff, would have

increased taxonomic diversity compared to the susceptible Heerup cultivar due to the link between

microbial diversity and disease resistance (Hu et al., 2016; Li et al., 2023). Both culture-based and

sequencing approaches indicated the preference of *Pseudomonas* species and strains to one cultivar

or the other. These results agree with previous work that reports the ability of the plant to

differentiate between phylogenetically similar bacteria (Thoms et al., 2023; Zhang and Kong, 2022).

Culturable and *in silico* community analysis established that the Sheriff cultivar harbors a more

taxonomically diverse *Pseudomonas* community, in support of our hypothesis. While not fully

explored in this work, it may not be coincidental that the *Fusarium*-resistant cultivar Sheriff harbors

two times the number of unique *Pseudomonas* strains on its roots compared to its susceptible

relative, given the link between microbial diversity and plant health (Berg et al., 2017). Testing the

protective ability of synthetic microbial communities of increasing intragenus complexity could

support the idea of the beneficial effect of taxonomic microdiversity on plant health (Hu et al., 2016;

Li et al., 2023).

Additionally in our first hypothesis, we predicted that increased taxonomic microdiversity in Sheriff

pseudomonads would coincide with increased functional diversity, as *Pseudomonas* taxonomy is

often related to functional abilities of the microbe (Lyng et al., 2024; Oni et al., 2019). While many

COG20 functions were differentially abundant between the Heerup and Sheriff pseudomonads, they

were equally distributed between the cultivars, giving no indication of increased functional diversity

in one cultivar or the other. However, secondary metabolites biosynthesis, transport and catabolism

was a COG20 category affected by cultivar. Thus, we analyzed the secondary metabolite biosynthetic

potential of our strain collection in more detail. BGC families are one way to group architecturally

similar BGCs that can be linked to a natural product chemotype (Navarro-Muñoz et al., 2020), and

thus, may be used as a proxy for different potential functions based on secondary metabolites.

Interestingly, the less taxonomically diverse Heerup pseudomonads demonstrated increased

biosynthetic functional diversity, boasting 19 enriched BGC families of 11 different product types

while Sheriff only had nine families of five product types. The preferential colonization of the highly

biosynthetic species *P. asgharzadehiana* on Heerup and the lack of biosynthetic genes in the many

Sheriff-enriched species underpins this result. This goes against our preconceived hypothesis that

increased taxonomic microdiversity results in increased functional microdiversity. Instead, few taxa

were responsible for boosting biosynthetic diversity within the Heerup cultivar. This may be a

conserved characteristic in soil, as bacterial taxonomic and functional diversity have previously been

shown to be negatively correlated in this niche (Wang et al., 2023). Additional work in this area

would elucidate the causes and consequences of such an inverse relationship and what this means

for microdiversity as a beneficial trait in plant microbiomes.

Functional diversity within a microbial genus is only rarely examined between crop genotypes, and

not yet between closely related modern cultivars (Oni et al., 2019; Pacheco-Moreno et al., 2024). We

demonstrated the prevalence of microbial microdiversity in the absence of crop diversity by

determining cultivar-dependent functional traits within the culturable *Pseudomonas* community. We

found several instances of genomic loci enriched in one cultivar or the other. The distribution of BGC

families also revealed cultivar-specific biosynthetic potential within the pseudomonads. In one case,

it stemmed from cultivar-dependent interspecies microdiversity. The species *P. brassicacearum*

carried BGC family 3677 only when it was isolated from the Sheriff cultivar. Interestingly, *P.*

brassicacearum was also the only species where the average number of BGCs per genome was

affected by cultivar, with Sheriff *P. brassicacearum* strains harboring more BGCs per genome in

comparison to their Heerup counterparts. This may reflect an exciting ability of the plant to rely on

bacterial secondary metabolites to structure its microbiome at the fine taxonomic scale. Previously,

secondary metabolites have been implicated in driving plant distinction between closely-related

microbes (Thoms et al., 2023) and used by *Streptomycetes* to carve a niche on the plant root (Nicolle

et al., 2024). Deeper genome sequencing rhizosphere pseudomonads from additional wheat

cultivars would confirm this interesting trend of cultivar-dependent interspecies microdiversity. It is

also still up for debate why pseudomonads carrying these BGC families are found on their respective

cultivar, especially since most of the BGCs in the strain library are unclassified beyond their product

class. Further work in this area could resolve how crops recruit *Pseudomonas* producing certain

metabolites to their microbiome, a mechanism previously suggested but poorly understood (Liu et

al., 2021; Feng et al., 2023; Yang et al., 2023).

Our second hypothesis predicted that pseudomonads from the *F. culmorum*-resistant Sheriff cultivar

would inhibit *F. culmorum* growth and harbor a greater abundance of genes capable of antagonizing

F. culmorum. *Pseudomonas* are often implicated in disease suppression, even when grouped

together at the genus level (Hong et al., 2023; Lv et al., 2023; Qiu et al., 2022). Thus, we examined

antifungal activity as a microdiverse trait within the *Pseudomonas* genus. In support of our

hypothesis, we found more *in vitro* antifungal activity against *F. culmorum* in the Sheriff isolates.

Isolates that were positive for fungal inhibition also mapped to ASVs that were enriched in the

Sheriff microbiome. In the pangenome, the BGC for the antagonistic metabolite 2,4-DAPG was found

equally distributed between the two cultivars. Neither of the two CLPs found in the strain library,

viscosin and lokisin, were enriched in either Heerup or Sheriff cultivar. In line with our hypothesis,

we found a chitinase-encoding gene of the GH19 family enriched in Sheriff isolates. The GH19 family,

originating from plants but also found in some bacteria including *Pseudomonas* (Ren et al., 2022),

are notable for their activity against *Fusarium* (García-Fraga et al., 2015). We also found a predicted

P. fluorescens-type pyoverdine BGC that was enriched in Sheriff pseudomonads. Bacterial

siderophores empower phytopathogen control via competition for iron (Gu et al., 2020) and have

been long documented as a *Pseudomonas* bioactive molecule against *Fusarium* (Kloepper et al.,

1980). In agreement with our observations, siderophore-mediated iron uptake was previously shown

to be a lineage-specific trait in *Pseudomonas* (Shalev et al., 2021). Thus, the specific *Pseudomonas*

traits present in a microbiome may be more important than the abundance of the *Pseudomonas*

genus as a whole in the context of plant health. Future efforts to analyze the plant holobiont for

improved health should therefore be focused on function, rather than taxonomy.

Our last hypothesis predicted that the wheat cultivar with the higher taxonomically diverse

microbiome would have a thinner average root diameter, and in support of this hypothesis, we saw

that the more diverse Sheriff cultivar had a thinner average root diameter and a higher proportion of

thinner roots compared to Heerup. This contributes to the growing body of literature that describes

a more diverse microbiome in both woody and herbaceous plant species which have thinner roots or

a root system dominated by fine roots (Saleem et al., 2018; Pervaiz et al., 2020; Luo et al., 2021; Zai

et al., 2021; Fleishman et al., 2023). Nevertheless, it is not possible at this time to attribute the

differences in *Pseudomonas* communities to the root morphology of the cultivars, since the cultivars

are genetically different. Other factors including root exudation and other plant genes are certainly

at play (Sasse et al., 2018; Song et al., 2021). Still, root diameter and root morphology as a whole

might play a previously unexplored role in microbial community composition, as we have suggested

previously (Herms et al., 2022), and deserve further evaluation.

5. Conclusion

The rhizosphere microbiome is a key determinant of plant health, and ongoing research aims to

unravel the secrets to utilizing beneficial plant-microbe interactions to promote sustainable

agriculture. As agricultural systems become increasingly homogenous, it is important to understand

how closely related modern cultivars can still differ in their microbiomes. In this work, we

demonstrate that the status quo of grouping together highly diverse genera, such as *Pseudomonas*,

in 16s rRNA analyses overlooks crucial differences in microbial functional activity (Chiniquy et al.,

2021; Jaspers and Overmann, 2004). Failing to consider the importance of microdiversity in shaping

the composition and function of the rhizosphere microbiome conceals the true potential of root-

associated microbial communities and should be a focus area in future plant holobiont research. We

suggest that microdiversity should be explored as a potential metric for the health and productivity

of plant microbiomes, but the relationship between taxonomy and function remains a confounding

factor on the mechanism underpinning taxonomic diversity and pathogen suppression. In addition to

providing novel insight into the microbial microdiversity of elite crop cultivars, our unique collection

of cultivar-specific *Pseudomonas* isolates can act as a tool to further dissect high-resolution plant-

microbe interactions. Exploration of these cultivar-dependent genomic loci may reveal insights

regarding the specificity of plant-microbe interactions at the species level. Ultimately, unraveling this

interaction can allow for designer crops that use microbial metabolites for increased resilience in the

face of challenging growing conditions.

CREDIT authorship statement

Courtney Horn Herms: Conceptualization, Data Curation, Formal analysis, Investigation,

Methodology, Project administration, Software, Visualization, Writing - Original Draft, Writing -

Review & Editing. Rosanna Catherine Hennessy: Conceptualization, Investigation, Methodology,

Supervision, Writing - Review & Editing. Frederik Bak: Conceptualization, Data curation, Formal

analysis, Methodology, Software, Visualization, Writing - Review & Editing. Ying Guan: Investigation,

Methodology, Writing - Review & Editing. Patrick Denis Browne: Data curation, Formal analysis,

Methodology, Software, Visualization, Writing - Original Draft, Writing - Review & Editing. Tue

Kjærgaard Nielsen: Data curation, Formal analysis, Methodology, Software, Writing - Review &

Editing. Lars Hestbjerg Hansen: Methodology, Resources, Writing - Review & Editing. Dorte Bodin

Dresbøll: Methodology, Resources, Writing - Review & Editing. Mette Haubjerg Nicolaisen:

Conceptualization, Funding acquisition, Methodology, Project administration, Resources,

Supervision, Writing - Review & Editing.

Conflict of Interests

The authors report no conflicts of interest.

Acknowledgements

We thank Dorette Müller-Stöver and Marie Louise Bornø, University of Copenhagen, for supplying

experimental soils. The *Fusarium culmorum* strain was kindly provided by Birgit Jensen, University of

Copenhagen. Thanks to our group members Kitzia Yashvelt Molina Zamudio and Dorthe Thybo

Ganzhorn for their support with the sampling.

Funding

This work was supported by the Novo Nordisk Foundation [grant number NNF19SA0059360].

Tables

ASV	Antifungal isolates within ASV	median number of reads in Sheriff	median number of reads in Heerup	Pval
S1E05	0/1	1.0	0.0	0.0472
S2B12_S2C03	1/2	3.0	0.0	0.0322
ASV67	n/a	1.0	0.0	0.0098
H3E07_S3A09	1/2	1.0	0.0	0.0398

562   **Table 1A. Enriched ASVs in Heerup and Sheriff at T2, with notice of the isolates that were**

563   **antifungal in the *in vitro* screen. P values were determined using the Mann-Whitney U-test. ASVs**

564   **were omitted from composition tests if the ASV is not seen in at least 50% of the samples in one of**

565   **the groups.**

ASV	Antifungal isolates within ASV	median number of reads in Sheriff	median number of reads in Heerup	Pval
H1D09_H2B05_H2D02_S1A03_S2D10_S3C01_S3E11_S3F07_S3H10	6/9	14.0	6.0	0.0305
H3E07_S3A09	1/2	1.0	0.0	0.0036
H3H08	0/1	1.0	0.0	0.0079

566   **Table 1B. Enriched ASVs in Heerup and Sheriff across T1 and T2, with notice of the isolates that**

567   **were antifungal in the *in vitro* screen. P values were determined using the Mann-Whitney U-test.**

568   **ASVs were omitted from composition tests if the ASV is not seen in at least 50% of the samples in**

569   **one of the groups.**

BGC family	Known BGC	Product Class	Enriched in	Adjusted p-value
2428	3-thiaglutamate	RiPP-like	n/a	0.22
2549	Viscosin	NRPS	n/a	0.09
2653	Arthrofactin A, anikasin, lokisin, stechlisin B2	NRPS	n/a	0.84
3049	Pf-5 pyoverdine	NRPS	Sheriff	**0.0297**
3453	2,4-diacetylphloroglucinol	T3PKS	n/a	0.37
3764	Tabtoxin	blactam	n/a	0.36

571 **Table 2. BGC families in the strain library that were identified as known products and the cultivar**

572 **they were enriched in, if applicable. Significant differences in the abundance of the BGC family**

573 **between cultivars was calculated using the two-sided Wilcoxon Rank Sum test with fdr-**

574 **adjustment using the Benjamini & Hochberg method. Significant p-values ($p < 0.05$) are bolded.**

References

Alsohim, A.S., Taylor, T.B., Barrett, G.A., Gallie, J., Zhang, X.X., Altamirano-Junqueira, A.E., Johnson,

L.J., Rainey, P.B., Jackson, R.W., 2014. The biosurfactant viscosin produced by Pseudomonas

fluorescensSBW25 aids spreading motility and plant growth promotion. Environmental

Microbiology 16, 2267–2281. https://doi.org/10.1111/1462-2920.12469

Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W., Lipman, D.J., 1997. Gapped

BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids

Research 25, 3389–3402. https://doi.org/10.1093/nar/25.17.3389

Bai, B., Liu, W., Qiu, X., Zhang, Jie, Zhang, Jingying, Bai, Y., 2022. The root microbiome: Community

assembly and its contributions to plant fitness. Journal of Integrative Plant Biology 64, 230–

243. https://doi.org/10.1111/jipb.13226

Benjamini, Y., Hochberg, Y., 1995. Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B

(Methodological) 57, 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Berg, G., Köberl, M., Rybakova, D., Müller, H., Grosch, R., Smalla, K., 2017. Plant microbial diversity is

suggested as the key to future biocontrol and health trends. FEMS Microbiology Ecology 93,

1–9. https://doi.org/10.1093/femsec/fix050

Bodour, A.A., Miller-Maier, R.M., 1998. Application of a modified drop-collapse technique for

surfactant quantitation and screening of biosurfactant-producing microorganisms. Journal of

Microbiological Methods 32, 273–280. https://doi.org/10.1016/S0167-7012(98)00031-1

Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Fierer, N.,

Peña, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley, S.T., Knights, D., Koenig, J.E.,

Ley, R.E., Lozupone, C.A., McDonald, D., Muegge, B.D., Pirrung, M., Reeder, J., Sevinsky, J.R.,

Turnbaugh, P.J., Walters, W.A., Widmann, J., Yatsunenko, T., Zaneveld, J., Knight, R., 2010.

QIIME allows analysis of high-throughput community sequencing data. Nat Methods 7, 335–

336. https://doi.org/10.1038/nmeth.f.303

Castresana, J., 2000. Selection of Conserved Blocks from Multiple Alignments for Their Use in

603   Phylogenetic Analysis. Molecular Biology and Evolution 17, 540–552.

604   https://doi.org/10.1093/oxfordjournals.molbev.a026334

605   Chao, A., 1984. Nonparametric Estimation of the Number of Classes in a Population. Scandinavian

606   Journal of Statistics 11, 265–270.

607   Chaumeil, P.-A., Mussig, A.J., Hugenholtz, P., Parks, D.H., 2020. GTDB-Tk: a toolkit to classify

608   genomes with the Genome Taxonomy Database. Bioinformatics 36, 1925–1927.

609   https://doi.org/10.1093/bioinformatics/btz848

610   Chiniquy, D., Barnes, E.M., Zhou, J., Hartman, K., Li, X., Sheflin, A., Pella, A., Marsh, E., Prenni, J.,

611   Deutschbauer, A.M., Schachtman, D.P., Tringe, S.G., 2021. Microbial Community Field

612   Surveys Reveal Abundant Pseudomonas Population in Sorghum Rhizosphere Composed of

613   Many Closely Related Phylotypes. Front. Microbiol. 12.

614   https://doi.org/10.3389/fmicb.2021.598180

615   Clarke, K.R., 1993. Non-parametric multivariate analyses of changes in community structure.

616   Australian Journal of Ecology 18, 117–143. https://doi.org/10.1111/j.1442-

617   9993.1993.tb00438.x

618   De Coster, W., D'Hert, S., Schultz, D.T., Cruts, M., Van Broeckhoven, C., 2018. NanoPack: visualizing

619   and processing long-read sequencing data. Bioinformatics 34, 2666–2669.

620   https://doi.org/10.1093/bioinformatics/bty149

621   Dubey, M., Hadadi, N., Pelet, S., Carraro, N., Johnson, D.R., van der Meer, J.R., 2021. Environmental

622   connectivity controls diversity in soil microbial communities. Commun Biol 4, 1–15.

623   https://doi.org/10.1038/s42003-021-02023-2

624   Edgar, R.C., 2018. Updating the 97% identity threshold for 16S ribosomal RNA OTUs. Bioinformatics

625   34, 2371–2375. https://doi.org/10.1093/bioinformatics/bty113

626   Edgar, R.C., 2016. UNOISE2: improved error-correction for Illumina 16S and ITS amplicon

627   sequencing. https://doi.org/10.1101/081257

Edgar, R.C., 2010. Search and clustering orders of magnitude faster than BLAST. Bioinformatics 26,

2460–2461. https://doi.org/10.1093/bioinformatics/btq461

Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput.

Nucleic Acids Research 32, 1792–1797. https://doi.org/10.1093/nar/gkh340

Elouafi, I., 2024. Why biodiversity matters in agriculture and food systems. Science 386, eads8197.

https://doi.org/10.1126/science.ads8197

Eren, A.M., Kiefl, E., Shaiber, A., Veseli, I., Miller, S.E., Schechter, M.S., Fink, I., Pan, J.N., Yousef, M.,

Fogarty, E.C., Trigodet, F., Watson, A.R., Esen, Ö.C., Moore, R.M., Clayssen, Q., Lee, M.D.,

Kivenson, V., Graham, E.D., Merrill, B.D., Karkman, A., Blankenberg, D., Eppley, J.M., Sjödin,

A., Scott, J.J., Vázquez-Campos, X., McKay, L.J., McDaniel, E.A., Stevens, S.L.R., Anderson,

R.E., Fuessel, J., Fernandez-Guerra, A., Maignien, L., Delmont, T.O., Willis, A.D., 2021.

Community-led, integrated, reproducible multi-omics with anvi'o. Nat Microbiol 6, 3–6.

https://doi.org/10.1038/s41564-020-00834-3

Feng, H., Fu, R., Luo, J., Hou, X., Gao, K., Su, L., Xu, Y., Miao, Y., Liu, Y., Xu, Z., Zhang, N., Shen, Q., Xun,

W., Zhang, R., 2023. Listening to plant's Esperanto via root exudates: reprogramming the

functional expression of plant growth-promoting rhizobacteria. New Phytologist 239, 2307–

2319. https://doi.org/10.1111/nph.19086

Fleishman, S.M., Centinari, M., Bell, T.H., Eissenstat, D.M., 2023. Assessing microbial communities

across the fine root landscape. Journal of Experimental Botany erad019.

https://doi.org/10.1093/jxb/erad019

García-Fraga, B., da Silva, A.F., López-Seijas, J., Sieiro, C., 2015. A novel family 19 chitinase from the

marine-derived *Pseudoalteromonas tunicata* CCUG 44952T: Heterologous expression,

characterization and antifungal activity. Biochemical Engineering Journal 93, 84–93.

https://doi.org/10.1016/j.bej.2014.09.014

García-García, N., Tamames, J., Linz, A.M., Pedrós-Alió, C., Puente-Sánchez, F., 2019. Microdiversity

ensures the maintenance of functional microbial communities under changing

654 environmental conditions. ISME J 13, 2969–2983. https://doi.org/10.1038/s41396-019-0487-

655 8

656 Garrido-Sanz, D., Čaušević, S., Vacheron, J., Heiman, C.M., Sentchilo, V., van der Meer, J.R., Keel, C.,

657 2023. Changes in structure and assembly of a species-rich soil natural community with

658 contrasting nutrient availability upon establishment of a plant-beneficial Pseudomonas in

659 the wheat rhizosphere. Microbiome 11, 214. https://doi.org/10.1186/s40168-023-01660-5

660 Gavriilidou, A., Kautsar, S.A., Zaburannyi, N., Krug, D., Müller, R., Medema, M.H., Ziemert, N., 2022.

661 Compendium of specialized metabolite biosynthetic diversity encoded in bacterial genomes.

662 Nat Microbiol 7, 726–735. https://doi.org/10.1038/s41564-022-01110-2

663 Getzke, F., Hassani, M.A., Crüsemann, M., Malisic, M., Zhang, P., Ishigaki, Y., Böhringer, N.,

664 Jiménez Fernández, A., Wang, L., Ordon, J., Ma, K.-W., Thiergart, T., Harbort, C.J., Wesseler,

665 H., Miyauchi, S., Garrido-Oter, R., Shirasu, K., Schäberle, T.F., Hacquard, S., Schulze-Lefert, P.,

666 2023. Cofunctioning of bacterial exometabolites drives root microbiota establishment.

667 Proceedings of the National Academy of Sciences 120, e2221508120.

668 https://doi.org/10.1073/pnas.2221508120

669 Girard, L., Lood, C., Höfte, M., Vandamme, P., Rokni-Zadeh, H., van Noort, V., Lavigne, R., De Mot, R.,

670 2021. The Ever-Expanding Pseudomonas Genus: Description of 43 New Species and Partition

671 of the Pseudomonas putida Group. Microorganisms 9, 1766.

672 https://doi.org/10.3390/microorganisms9081766

673 Gould, W.D., Hagedorn, C., Bardinelli, T.R., Zablotowicz, R.M., 1985. New Selective Media for

674 Enumeration and Recovery of Fluorescent Pseudomonads from Various Habitats. Applied

675 and Environmental Microbiology 49, 28–32. https://doi.org/10.1128/aem.49.1.28-32.1985

676 Gross, H., Loper, J.E., 2009. Genomics of secondary metabolite production by Pseudomonas spp.

677 Natural Product Reports 26, 1408–1446. https://doi.org/10.1039/b817075b

678 Gu, S., Wei, Z., Shao, Z., Friman, V.-P., Cao, K., Yang, T., Kramer, J., Wang, X., Li, M., Mei, X., Xu, Y.,

679 Shen, Q., Kümmerli, R., Jousset, A., 2020. Competition for iron drives phytopathogen control

by natural rhizosphere microbiomes. Nat Microbiol 5, 1002–1010.

https://doi.org/10.1038/s41564-020-0719-8

Hennessy, R.C., Dichmann, S.I., Martens, H.J., Zervas, A., Stougaard, P., 2020. Serratia inhibens sp.

nov., a new antifungal species isolated from potato (Solanum tuberosum). Int J Syst Evol

Microbiol 70, 4204–4211. https://doi.org/10.1099/ijsem.0.004270

Herms, C.H., Hennessy, R.C., Bak, F., Dresbøll, D.B., Nicolaisen, M.H., 2022. Back to our roots:

exploring the role of root morphology as a mediator of beneficial plant–microbe

interactions. Environmental Microbiology 00. https://doi.org/10.1111/1462-2920.15926

Hong, S., Yuan, X., Yang, J., Yang, Y., Jv, H., Li, R., Jia, Z., Ruan, Y., 2023. Selection of rhizosphere

communities of diverse rotation crops reveals unique core microbiome associated with

reduced banana Fusarium wilt disease. New Phytologist 238, 2194–2209.

https://doi.org/10.1111/nph.18816

Hu, J., Wei, Z., Friman, V.P., Gu, S.H., Wang, X.F., Eisenhauer, N., Yang, T.J., Ma, J., Shen, Q.R., Xu,

Y.C., Jousset, A., 2016. Probiotic diversity enhances rhizosphere microbiome function and

plant disease suppression. mBio 7, 1–8. https://doi.org/10.1128/mBio.01790-16

Huang, Y., Sheth, R.U., Zhao, S., Cohen, L.A., Dabaghi, K., Moody, T., Sun, Y., Ricaurte, D., Richardson,

M., Velez-Cortes, F., Blazejewski, T., Kaufman, A., Ronda, C., Wang, H.H., 2023. High-

throughput microbial culturomics using automation and machine learning. Nat Biotechnol 1–

10. https://doi.org/10.1038/s41587-023-01674-2

Iannucci, A., Canfora, L., Nigro, F., De Vita, P., Beleggia, R., 2021. Relationships between root

morphology, root exudate compounds and rhizosphere microbial community in durum

wheat. Applied Soil Ecology 158, 103781. https://doi.org/10.1016/j.apsoil.2020.103781

Ilmi, M., Putri, L.K., Muhamad, A.A.K., Cholishoh, A., Ardiansyah, S.A., 2019. Use of Mung Bean

Sprout (Tauge) as Alternative Fungal Growth Medium. J. Phys.: Conf. Ser. 1241, 012015.

https://doi.org/10.1088/1742-6596/1241/1/012015

Jaspers, E., Overmann, J., 2004. Ecological Significance of Microdiversity: Identical 16S rRNA Gene

Sequences Can Be Found in Bacteria with Highly Divergent Genomes and Ecophysiologies.

Applied and Environmental Microbiology 70, 4831–4839.

https://doi.org/10.1128/AEM.70.8.4831-4839.2004

Jørgensen, L.N., Heick, T.M., Abuley, I.K., Kudsk, P., Hansen Kemezys, A., 2023. Applied Crop

Protection 2022 (Research Dissemination No. 216). Aarhus University, Danish Centre for

Food and Agriculture.

Jørgensen, L.N., Heick, T.M., Abuley, I.K., Mathiassen, S., Jensen, P.K., Kristjansen, H.S., Hartvig, P.,

2020. Applied Crop Protection 2019 (Research Dissemination No. 167). Aarhus University,

Danish Centre for Food and Agriculture.

Jousset, A., Schulz, W., Scheu, S., Eisenhauer, N., 2011. Intraspecific genotypic richness and

relatedness predict the invasibility of microbial communities. The ISME Journal 5, 1108–

1114. https://doi.org/10.1038/ismej.2011.9

King, W.L., Yates, C.F., Cao, L., O'Rourke-Ibach, S., Fleishman, S.M., Richards, S.C., Centinari, M.,

Hafner, B.D., Goebel, M., Bauerle, T., Kim, Y.-M., Nicora, C.D., Anderton, C.R., Eissenstat,

D.M., Bell, T.H., 2023. Functionally discrete fine roots differ in microbial assembly, microbial

functional potential, and produced metabolites. Plant, Cell & Environment 46, 3919–3932.

https://doi.org/10.1111/pce.14705

Kloepper, J.W., Leong, J., Teintze, M., Schroth, M.N., 1980. Pseudomonas siderophores: A

mechanism explaining disease-suppressive soils. Current Microbiology 4, 317–320.

https://doi.org/10.1007/BF02602840

Koeppel, A.F., Wertheim, J.O., Barone, L., Gentile, N., Krizanc, D., Cohan, F.M., 2013. Speedy

speciation in a bacterial microcosm: new species can arise as frequently as adaptations

within a species. The ISME Journal 7, 1080–1091. https://doi.org/10.1038/ismej.2013.3

Kolmogorov, M., Yuan, J., Lin, Y., Pevzner, P.A., 2019. Assembly of long, error-prone reads using

repeat graphs. Nat Biotechnol 37, 540–546. https://doi.org/10.1038/s41587-019-0072-8

Li, J., Yang, C., Jousset, A., Yang, K., Wang, X., Xu, Z., Yang, T., Mei, X., Zhong, Z., Xu, Y., Shen, Q.,

Friman, V.-P., Wei, Z., 2023. Engineering multifunctional rhizosphere probiotics using

consortia of Bacillus amyloliquefaciens transposon insertion mutants. eLife 12, e90726.

https://doi.org/10.7554/eLife.90726

Lin, Y., Yuan, J., Kolmogorov, M., Shen, M.W., Chaisson, M., Pevzner, P.A., 2016. Assembly of long

error-prone reads using de Bruijn graphs. Proceedings of the National Academy of Sciences

113, E8396–E8405. https://doi.org/10.1073/pnas.1604560113

Liu, H., Li, J., Carvalhais, L.C., Percy, C.D., Prakash Verma, J., Schenk, P.M., Singh, B.K., 2021. Evidence

for the plant recruitment of beneficial microbes to suppress soil-borne pathogens. New

Phytologist 229, 2873–2885. https://doi.org/10.1111/nph.17057

Loper, J.E., Hassan, K.A., Mavrodi, D.V., Davis, E.W., Lim, C.K., Shaffer, B.T., Elbourne, L.D.H.,

Stockwell, V.O., Hartney, S.L., Breakwell, K., Henkels, M.D., Tetu, S.G., Rangel, L.I., Kidarsa,

T.A., Wilson, N.L., van de Mortel, J.E., Song, C., Blumhagen, R., Radune, D., Hostetler, J.B.,

Brinkac, L.M., Durkin, A.S., Kluepfel, D.A., Wechter, W.P., Anderson, A.J., Kim, Y.C., Pierson,

L.S., Pierson, E.A., Lindow, S.E., Kobayashi, D.Y., Raaijmakers, J.M., Weller, D.M.,

Thomashow, L.S., Allen, A.E., Paulsen, I.T., 2012. Comparative genomics of plant-associated

pseudomonas spp.: Insights into diversity and inheritance of traits involved in multitrophic

interactions. PLoS Genetics 8. https://doi.org/10.1371/journal.pgen.1002784

Lopes, L.D., Davis, E.W., Pereira e Silva, M. de C., Weisberg, A.J., Bresciani, L., Chang, J.H., Loper, J.E.,

Andreote, F.D., 2018. Tropical soils are a reservoir for fluorescent Pseudomonas spp.

biodiversity. Environmental Microbiology 20, 62–74. https://doi.org/10.1111/1462-

2920.13957

Luo, W., Zai, X., Sun, J., Li, D., Li, Y., Li, G., Wei, G., Chen, W., 2021. Coupling Root Diameter With

Rooting Depth to Reveal the Heterogeneous Assembly of Root-Associated Bacterial

Communities in Soybean. Frontiers in Microbiology 12.

https://doi.org/10.3389/fmicb.2021.783563

Lv, N., Tao, C., Ou, Y., Wang, J., Deng, X., Liu, H., Shen, Z., Li, R., Shen, Q., 2023. Root-Associated

758   Antagonistic Pseudomonas spp. Contribute to Soil Suppressiveness against Banana Fusarium

759   Wilt Disease of Banana. Microbiology Spectrum 0, e03525-22.

760   https://doi.org/10.1128/spectrum.03525-22

761   Lyng, M., Þórisdóttir, B., Sveinsdóttir, S.H., Hansen, M.L., Jelsbak, L., Maróti, G., Kovács, Á.T., 2024.

762   Taxonomy of Pseudomonas spp. determines interactions with Bacillus subtilis. mSystems 0,

763   e00212-24. https://doi.org/10.1128/msystems.00212-24

764   Mauchline, T.H., Chedom-Fotso, D., Chandra, G., Samuels, T., Greenaway, N., Backhaus, A.,

765   Mcmillan, V., Canning, G., Powers, S.J., Hammond-Kosack, K.E., Hirsch, P.R., Clark, I.M.,

766   Mehrabi, Z., Roworth, J., Burnell, J., Malone, J.G., 2015. An analysis of Pseudomonas

767   genomic diversity in take-all infected wheat fields reveals the lasting impact of wheat

768   cultivars on the soil microbiota. Environmental Microbiology 17, 4764–4778.

769   https://doi.org/10.1111/1462-2920.13038

770   Mauchline, T.H., Malone, J.G., 2017. Life in earth – the root microbiome to the rescue? Current

771   Opinion in Microbiology, Environmental microbiology * CRISPRcas9 37, 23–28.

772   https://doi.org/10.1016/j.mib.2017.03.005

773   Medema, M.H., Blin, K., Cimermancic, P., de Jager, V., Zakrzewski, P., Fischbach, M.A., Weber, T.,

774   Takano, E., Breitling, R., 2011. antiSMASH: rapid identification, annotation and analysis of

775   secondary metabolite biosynthesis gene clusters in bacterial and fungal genome sequences.

776   Nucleic Acids Research 39, W339–W346. https://doi.org/10.1093/nar/gkr466

777   Mendes, L.W., Raaijmakers, J.M., De Hollander, M., Mendes, R., Tsai, S.M., 2018. Influence of

778   resistance breeding in common bean on rhizosphere microbiome composition and function.

779   ISME Journal 12, 212–224. https://doi.org/10.1038/ismej.2017.158

780   Miller, J.H., 1972. Experiments in molecular genetics. CSH NY.

781   Moore, L.R., Rocap, G., Chisholm, S.W., 1998. Physiology and molecular phylogeny of coexisting

782   Prochlorococcus ecotypes. Nature 393, 464–467. https://doi.org/10.1038/30965

Navarro-Muñoz, J.C., Selem-Mojica, N., Mullowney, M.W., Kautsar, S.A., Tryon, J.H., Parkinson, E.I.,

De Los Santos, E.L.C., Yeong, M., Cruz-Morales, P., Abubucker, S., Roeters, A., Lokhorst, W.,

Fernandez-Guerra, A., Cappelini, L.T.D., Goering, A.W., Thomson, R.J., Metcalf, W.W.,

Kelleher, N.L., Barona-Gomez, F., Medema, M.H., 2020. A computational framework to

explore large-scale biosynthetic diversity. Nat Chem Biol 16, 60–68.

https://doi.org/10.1038/s41589-019-0400-9

Nicolle, C., Gayrard, D., Noël, A., Hortala, M., Amiel, A., Grat, S., Le Ru, A., Marti, G., Pernodet, J.-L.,

Lautru, S., Dumas, B., Rey, T., 2024. Root-associated Streptomyces produce galbonolides to

modulate plant immunity and promote rhizosphere colonization. The ISME Journal wrae112.

https://doi.org/10.1093/ismejo/wrae112

Oksanen, J., Simpson, G.L., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'Hara, R.B.,

Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B.,

Borcard, D., Carvalho, G., Chirico, M., Caceres, M.D., Durand, S., Evangelista, H.B.A., FitzJohn,

R., Friendly, M., Furneaux, B., Hannigan, G., Hill, M.O., Lahti, L., McGlinn, D., Ouellette, M.-

H., Cunha, E.R., Smith, T., Stier, A., Braak, C.J.F.T., Weedon, J., 2022. vegan: Community

Ecology Package.

Olm, M.R., Brown, C.T., Brooks, B., Banfield, J.F., 2017. dRep: a tool for fast and accurate genomic

comparisons that enables improved genome recovery from metagenomes through de-

replication. ISME J 11, 2864–2868. https://doi.org/10.1038/ismej.2017.126

Oni, F.E., Geudens, N., Omoboye, O.O., Bertier, L., Hua, H.G.K., Adiobo, A., Sinnaeve, D., Martins, J.C.,

Höfte, M., 2019. Fluorescent Pseudomonas and cyclic lipopeptide diversity in the

rhizosphere of cocoyam (Xanthosoma sagittifolium). Environmental Microbiology 21, 1019–

1034. https://doi.org/10.1111/1462-2920.14520

Pacheco-Moreno, A., Bollmann-Giolai, A., Chandra, G., Brett, P., Davies, J., Thornton, O., Poole, P.,

Ramachandran, V., Brown, J.K.M., Nicholson, P., Ridout, C., DeVos, S., Malone, J.G., 2024.

The genotype of barley cultivars influences multiple aspects of their associated microbiota

via differential root exudate secretion. PLOS Biology 22, e3002232.

https://doi.org/10.1371/journal.pbio.3002232

Pacheco-Moreno, A., Stefanato, F.L., Ford, J.J., Trippel, C., Uszkoreit, S., Ferrafiat, L., Grenga, L.,

Dickens, R., Kelly, N., Kingdon, A.D.H., Ambrosetti, L., Nepogodiev, S.A., Findlay, K.C.,

Cheema, J., Trick, M., Chandra, G., Tomalin, G., Malone, J.G., Truman, A.W., 2021. Pan-

genome analysis identifies intersecting roles for pseudomonas specialized metabolites in

potato pathogen inhibition. eLife 10, 1–46. https://doi.org/10.7554/eLife.71900

Parks, D.H., Chuvochina, M., Chaumeil, P.-A., Rinke, C., Mussig, A.J., Hugenholtz, P., 2020. A

complete domain-to-species taxonomy for Bacteria and Archaea. Nat Biotechnol 38, 1079–

1086. https://doi.org/10.1038/s41587-020-0501-8

Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P., Tyson, G.W., 2015. CheckM: assessing the

quality of microbial genomes recovered from isolates, single cells, and metagenomes.

Genome Res. 25, 1043–1055. https://doi.org/10.1101/gr.186072.114

Pervaiz, Z.H., Contreras, J., Hupp, B.M., Lindenberger, J.H., Chen, D., Zhang, Q., Wang, C., Twigg, P.,

Saleem, M., 2020. Root microbiome changes with root branching order and root chemistry

in peach rhizosphere soil. Rhizosphere 16, 100249.

https://doi.org/10.1016/j.rhisph.2020.100249

Qiu, Z., Verma, J.P., Liu, H., Wang, J., Batista, B.D., Kaur, S., de Araujo Pereira, A.P., Macdonald, C.A.,

Trivedi, P., Weaver, T., Conaty, W.C., Tissue, D.T., Singh, B.K., 2022. Response of the plant

core microbiome to Fusarium oxysporum infection and identification of the pathobiome.

Environmental Microbiology 24, 4652–4669. https://doi.org/10.1111/1462-2920.16194

Qu, Q., Zhang, Z., Peijnenburg, W.J.G.M., Liu, W., Lu, T., Hu, B., Chen, Jianmeng, Chen, Jun, Lin, Z.,

Qian, H., 2020. Rhizosphere Microbiome Assembly and Its Impact on Plant Growth. Journal

of Agricultural and Food Chemistry 68, 5024–5038. https://doi.org/10.1021/acs.jafc.0c00073

Ren, X.-B., Dang, Y.-R., Liu, S.-S., Huang, K.-X., Qin, Q.-L., Chen, X.-L., Zhang, Y.-Z., Wang, Y.-J., Li, P.-Y.,

2022. Identification and Characterization of Three Chitinases with Potential in Direct

Conversion of Crystalline Chitin into N,N'-diacetylchitobiose. Mar Drugs 20, 165.

https://doi.org/10.3390/md20030165

Saleem, M., Law, A.D., Sahib, M.R., Pervaiz, Z.H., Zhang, Q., 2018. Impact of root system architecture

on rhizosphere and root microbiome. Rhizosphere 6, 47–51.

https://doi.org/10.1016/j.rhisph.2018.02.003

Sasse, J., Martinoia, E., Northen, T., 2018. Feed Your Friends: Do Plant Exudates Shape the Root

Microbiome? Trends in Plant Science 23, 25–41.

https://doi.org/10.1016/j.tplants.2017.09.003

Seemann, T., 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30, 2068–2069.

https://doi.org/10.1093/bioinformatics/btu153

Seethepalli, A., Dhakal, K., Griffiths, M., Guo, H., Freschet, G.T., York, L.M., 2021. RhizoVision

Explorer: open-source software for root image analysis and measurement standardization.

AoB PLANTS 13, plab056. https://doi.org/10.1093/aobpla/plab056

Shaiber, A., Willis, A.D., Delmont, T.O., Roux, S., Chen, L.-X., Schmid, A.C., Yousef, M., Watson, A.R.,

Lolans, K., Esen, Ö.C., Lee, S.T.M., Downey, N., Morrison, H.G., Dewhirst, F.E., Mark Welch,

J.L., Eren, A.M., 2020. Functional and genetic markers of niche partitioning among enigmatic

members of the human oral microbiome. Genome Biology 21, 292.

https://doi.org/10.1186/s13059-020-02195-w

Shalev, O., Ashkenazy, H., Neumann, M., Weigel, D., 2021. Commensal Pseudomonas protect

Arabidopsis thaliana from a coexisting pathogen via multiple lineage-dependent

mechanisms. ISME Journal 1–10. https://doi.org/10.1038/s41396-021-01168-6

Simonin, M., Dasilva, C., Terzi, V., Ngonkeu, E.L.M., Diouf, Di., Kane, A., Béna, G., Moulin, L., 2020.

Influence of plant genotype and soil on the wheat rhizosphere microbiome: Evidences for a

core microbiome across eight African and European soils. FEMS Microbiology Ecology 96, 1–

18. https://doi.org/10.1093/femsec/fiaa067

Song, Y., Wilson, A.J., Zhang, X.C., Thoms, D., Sohrabi, R., Song, S., Geissmann, Q., Liu, Y., Walgren, L.,

He, S.Y., Haney, C.H., 2021. FERONIA restricts Pseudomonas in the rhizosphere microbiome

via regulation of reactive oxygen species. Nature Plants 7, 644–654.

https://doi.org/10.1038/s41477-021-00914-0

Spragge, F., Bakkeren, E., Jahn, M.T., B. N. Araujo, E., Pearson, C.F., Wang, X., Pankhurst, L., Cunrath,

O., Foster, K.R., 2023. Microbiome diversity protects against pathogens by nutrient blocking.

Science 382, eadj3502. https://doi.org/10.1126/science.adj3502

Stevens, B.M., Creed, T.B., Reardon, C.L., Manter, D.K., 2023. Comparison of Oxford Nanopore

Technologies and Illumina MiSeq sequencing with mock communities and agricultural soil.

Sci Rep 13, 9323. https://doi.org/10.1038/s41598-023-36101-8

Takamatsu, K., Toyofuku, M., Okutani, F., Yamazaki, S., Nakayasu, M., Aoki, Y., Kobayashi, M., Ifuku,

K., Yazaki, K., Sugiyama, A., 2023. α-Tomatine gradient across artificial roots recreates the

recruitment of tomato root-associated Sphingobium. Plant Direct 7, e550.

https://doi.org/10.1002/pld3.550

Tatusov, R.L., Galperin, M.Y., Natale, D.A., Koonin, E.V., 2000. The COG database: a tool for genome-

scale analysis of protein functions and evolution. Nucleic Acids Res 28, 33–36.

Terlouw, B.R., Blin, K., Navarro-Muñoz, J.C., Avalon, N.E., Chevrette, M.G., Egbert, S., Lee, S., Meijer,

D., Recchia, M.J.J., Reitz, Z.L., van Santen, J.A., Selem-Mojica, N., Tørring, T., Zaroubi, L.,

Alanjary, M., Aleti, G., Aguilar, C., Al-Salihi, S.A.A., Augustijn, H.E., Avelar-Rivas, J.A., Avitia-

Domínguez, L.A., Barona-Gómez, F., Bernaldo-Agüero, J., Bielinski, V.A., Biermann, F., Booth,

T.J., Carrion Bravo, V.J., Castelo-Branco, R., Chagas, F.O., Cruz-Morales, P., Du, C., Duncan,

K.R., Gavriilidou, A., Gayrard, D., Gutiérrez-García, K., Haslinger, K., Helfrich, E.J.N.,

van der Hooft, J.J.J., Jati, A.P., Kalkreuter, E., Kalyvas, N., Kang, K.B., Kautsar, S., Kim, W.,

Kunjapur, A.M., Li, Y.-X., Lin, G.-M., Loureiro, C., Louwen, J.J.R., Louwen, N.L.L., Lund, G.,

Parra, J., Philmus, B., Pourmohsenin, B., Pronk, L.J.U., Rego, A., Rex, D.A.B., Robinson, S.,

Rosas-Becerra, L.R., Roxborough, E.T., Schorn, M.A., Scobie, D.J., Singh, K.S., Sokolova, N.,

Tang, X., Udwary, D., Vigneshwari, A., Vind, K., Vromans, S.P.J.M., Waschulin, V., Williams,

S.E., Winter, J.M., Witte, T.E., Xie, H., Yang, D., Yu, J., Zdouc, M., Zhong, Z., Collemare, J.,

Linington, R.G., Weber, T., Medema, M.H., 2023. MIBiG 3.0: a community-driven effort to

annotate experimentally validated biosynthetic gene clusters. Nucleic Acids Research 51,

D603–D610. https://doi.org/10.1093/nar/gkac1049

Thoms, D., Chen, M.Y., Liu, Y., Moreira, Z.M., Luo, Y., Song, S., Wang, N.R., Haney, C.H., 2023. Innate

immunity can distinguish beneficial from pathogenic rhizosphere microbiota. bioRxiv

2023.01.07.523123. https://doi.org/10.1101/2023.01.07.523123

van der Bom, F., Magid, J., Jensen, L.S., 2017. Long-term P and K fertilisation strategies and balances

affect soil availability indices, crop yield depression risk and N use. European Journal of

Agronomy 86, 12–23. https://doi.org/10.1016/j.eja.2017.02.006

Vandenkoornhuyse, P., Quaiser, A., Duhamel, M., Le Van, A., Dufresne, A., 2015. The importance of

the microbiome of the plant holobiont. New Phytologist 206, 1196–1206.

https://doi.org/10.1111/nph.13312

von Stein, O.D., Thies, W.G., Hofmann, M., 1997. A high throughput screening for rarely transcribed

differentially expressed genes. Nucleic Acids Res 25, 2598–2602.

https://doi.org/10.1093/nar/25.13.2598

Wang, C., Yu, Q.-Y., Ji, N.-N., Zheng, Y., Taylor, J.W., Guo, L.-D., Gao, C., 2023. Bacterial genome size

and gene functional diversity negatively correlate with taxonomic diversity along a pH

gradient. Nat Commun 14, 7437. https://doi.org/10.1038/s41467-023-43297-w

Wang, N.R., Wiesmann, C.L., Melnyk, R.A., Hossain, S.S., Chi, M.-H., Martens, K., Craven, K., Haney,

C.H., 2022. Commensal Pseudomonas fluorescens Strains Protect Arabidopsis from Closely

Related Pseudomonas Pathogens in a Colonization-Dependent Manner. mBio 13.

https://doi.org/10.1128/mbio.02892-21

Weisburg, W.G., Barns, S.M., Pelletier, D.A., Lane, D.J., 1991. 16S ribosomal DNA amplification for

phylogenetic study. J Bacteriol 173, 697–703.

Widmer, F., Seidler, R.J., Gillevet, P.M., Watrud, L.S., Di Giovanni, G.D., 1998. A Highly Selective PCR

Protocol for Detecting 16S rRNA Genes of the Genus Pseudomonas (Sensu Stricto) in

Environmental Samples. Appl Environ Microbiol 64, 2545–2553.

Yang, K., Fu, R., Feng, H., Jiang, G., Finkel, O., Sun, T., Liu, M., Huang, B., Li, S., Wang, X., Yang, T.,

Wang, Y., Wang, S., Xu, Y., Shen, Q., Friman, V.-P., Jousset, A., Wei, Z., 2023. RIN enhances

plant disease resistance via root exudate-mediated assembly of disease-suppressive

rhizosphere microbiota. Molecular Plant. https://doi.org/10.1016/j.molp.2023.08.004

Yang, T., Wei, Z., Friman, V., Xu, Y., Shen, Q., Kowalchuk, G.A., Jousset, A., 2017. Resource availability

modulates biodiversity-invasion relationships by altering competitive interactions.

Environmental Microbiology 19, 2984–2991. https://doi.org/10.1111/1462-2920.13708

Zai, X., Luo, W., Bai, W., Li, Y., Xiao, X., Gao, X., Wang, E., Wei, G., Chen, W., 2021. Effect of Root

Diameter on the Selection and Network Interactions of Root-Associated Bacterial

Microbiomes in Robinia pseudoacacia L. Microbial Ecology 82, 391–402.

https://doi.org/10.1007/s00248-020-01678-4

Zhang, M., Kong, X., 2022. How plants discern friends from foes. Trends in Plant Science 27, 107–

109. https://doi.org/10.1016/j.tplants.2021.11.001

Zhang, Z.J., Cole, C.G., Coyne, M.J., Lin, H., Dylla, N., Smith, R.C., Pappas, T.E., Townson, S.A.,

Laliwala, N., Waligurski, E., Ramaswamy, R., Woodson, C., Burgo, V., Little, J.C., Moran, D.,

Rose, A., McMillin, M., McSpadden, E., Sundararajan, A., Sidebottom, A.M., Pamer, E.G.,

Comstock, L.E., 2024. Comprehensive analyses of a large human gut Bacteroidales culture

collection reveal species- and strain-level diversity and evolution. Cell Host & Microbe 0.

https://doi.org/10.1016/j.chom.2024.08.016

Species

Species
P. baetica
P. extremaustralis
P. kilonensis
P. koreensis
P. migulae

Core Genome (2,322 gene clusters)

Cultivar isolated from   ■ Heerup (53)   ■ Sheriff (59)

Species   ■ tritici (1)   ■ brassicacearum (33)   ■ sivasensis (5)   ■ neuropathica (5)   ■ siliginis (10)   ■ asgharzadehiana (18)   ■ alvandae (1)   ■ sp900581005 (9)   ■ mandelii (1)
■ simiae (3)   ■ sp004011755 (3)   ■ mucoides (1)   ■ sp001429045 (4)   ■ sp002000165 (1)   ■ abietaniphila (1)   ■ sp009764265 (1)   ■ atacamensis (2)   ■ sp.1 (3)   ■ sp.2 (1)
■ sp.4 (5)   ■ sp.3 (4)

Strain   ■ C (1)   ■ N (2)   ■ D (5)   ■ I (3)   ■ LL (2)   ■ P (2)   ■ FF (1)   ■ G (16)   ■ J (1)   ■ Q (5)   ■ X (1)   ■ S (1)   ■ H (9)   ■ R (6)   ■ NN (1)   ■ TT (1)   ■ BB (3)   ■ A (2)   ■ O (1)
■ KK (1)   ■ E (4)   ■ W (1)   ■ Y (2)   ■ UU (1)   ■ DD (4)   ■ HH (1)   ■ EE (2)   ■ RR (1)   ■ GG (1)   ■ Z (2)   ■ AA (1)   ■ PP (4)   ■ T (1)   ■ MM (1)   ■ SS (1)   ■ VV (1)
■ OO (1)   ■ QQ (1)   ■ F (2)   ■ B (1)   ■ CC (2)   ■ II (1)   ■ U (6)   ■ JJ (2)   ■ V (1)   ■ M (1)   ■ L (1)   ■ K (1)

Root scanning

1cm sample between 1cm-2cm from top of the root

Isolation

4cm sample from middle of the root

T1: First node

1cm sample between 11cm-12cm from top of the root

4 weeks

1cm sample between 18cm-19cm from top of the root

1/4 root system

T1: First node

T2: Flag leaf

Long-read 16S rRNA sequencing

Methods Figure 1. Sampling for Pseudomonas isolation, long-read 16S rRNA amplicon sequencing,

and root scanning occurred along different root axes as well as across different time points.

Created with Biorender.com.

Figure 1. Composition of the Heerup and Sheriff _Pseudomonas_ strain libraries identified by Sanger

sequencing of minimum 900 bp of the 16S rRNA gene from 195 purified isolates from Heerup and

178 purified isolates from Sheriff. Each color represents a _Pseudomonas_ species or CSV. A)

Composition at the species level identified by sequence alignment to the NBCI 16S rRNA gene

database. A legend identifying the top five most abundant species is shown. B) Composition at the

CSV level, defined as 100% identity across a minimum of 900 bp of the 16S rRNA gene.

Figure 2. _In vitro F. culmorum_ inhibition and biosurfactant production of the Heerup and Sheriff

strain libraries. P-values of Fisher's exact test are given above the bars. Strains _Serratia inhibens_ S40

and _Pseudomonas fluorescens_ SBW25 were used as the positive control for the antifungal assay and

the biosurfactant assay, respectively.

Figure 3. Pangenome of the 112 genomes. A) Blue highlight represents an isolate cultured from

Heerup, and an orange highlight represents an isolate cultured from Sheriff. Shaded bars indicate

the presence of gene clusters in the genome. The core genome of 2,322 gene clusters is collapsed

for increased clarity. Species and strain identification is indicated. B) Highlight boxes and arrows

indicates the missing gene cluster locus in the Heerup-isolated strains of _P. brassicacearum_ strain R,

and the missing three gene cluster loci in Sheriff-isolated strains, despite its presence in closely

related (same species and strain) isolates. C) Highlight box and arrow indicates the missing one gene

cluster locus in Sheriff-isolated strains of _P. simiae_ despite its presence in closely related (same

species) isolates from Heerup.

Figure 4. Biosynthetic potential in the Heerup and Sheriff strain libraries. A) Number of BGCs per

genome of isolates cultured from Heerup and Sheriff. B) Differentially abundant BGC families found

in Heerup- or Sheriff-associated pseudomonads. BGC families are color-coded along the x-axis by

their predicted product class. The x-axis highlight box indicates if the BGC family is enriched in

Heerup (blue) or Sheriff (orange).

Figure 5. Distribution of BGCs in the species found in the _Pseudomonas_ strain library. A) Number of

BGCs per genome. True data points are overlaid the violin plot as filled circles. Significant differences

in average BGCs per genome of a species between cultivars was calculated using the two-sided

Wilcoxon Rank Sum test for the species _P. brassicacearum_, _P. siliginis_, and _P. asgharzadehiana,_ and

only _p_ < 0.05 are displayed. B) Proportion of isolates of each species harboring the BGC family that is

differentially enriched between Heerup and Sheriff. The x-axis highlight box represents if the

Pseudomonas species is unique to Heerup (blue), Sheriff (orange), or found on both cultivars (grey).

The y-axis highlight box represents if the BGC family is enriched in Heerup (blue) or Sheriff (orange).

Figure 6. Root scanning on four-week old plants to measure root morphology parameters. A) Root

system composition of Heerup and Sheriff divided into six root diameter ranges. The bar plot

displays the mean percentage of root diameter in the root system of each cultivar. True data points

are overlaid the bar plot as filled circles. B) Representative images of washed Heerup and Sheriff

root systems after four weeks of growth.