Reconfiguration of the visual code and retinal cell type complement in closely related diurnal and nocturnal mice

Authors

Annette E Allen*[1,2], Joshua Hahn[3], Rose Richardson[1,2], Andreea Pantiru[1,2], Josh Mouland[1,2,4], Beatriz Baño-Otalora[2,4], Aboozar Monavarfeshani[5], Wenjun Yan[5], Christopher Williams[1,2], Jonathan Wynne[1,2], Jessica Rodgers[1,2], Nina Milosavljevic[1,2], Patrycja Orlowska-Feuer[1,2], Riccardo Storchi[1,2], Joshua R. Sanes[5], Karthik Shekhar[3,6], Robert J Lucas[1,2]

*Corresponding author
[1]Division of Neuroscience, School of Biological Sciences, Faculty of Biology, Medicine and Health, University of Manchester, UK
[2]Centre for Biological Timing, Faculty of Biology Medicine and Health, University of Manchester, Manchester M13 9PT, UK
[3]Department of Chemical and Biomolecular Engineering, University of California, Berkeley, Berkeley, CA, USA
[4]Division of Diabetes, Endocrinology and Gastroenterology, School of Medical Sciences, Faculty of Biology, Medicine and Health, University of Manchester, UK.
[5]Department of Cellular and Molecular Biology, Center for Brain Science, Harvard University, Cambridge, MA, USA, 02138
[6]Helen Wills Neuroscience Institute; Vision Science Graduate Group; Center for Computational Biology; Biophysics Graduate Group; California Institute of Quantitative Biosciences (QB3), University of California, Berkeley, Berkeley, CA, USA

Abstract

How does evolution act on neuronal populations to match computational characteristics to functional demands? We address this problem by comparing visual code and retinal cell composition in closely related murid species with different behaviours. *Rhabdomys pumilio* are diurnal and have substantially thicker inner retina and larger visual thalamus than nocturnal *Mus musculus*. High-density electrophysiological recordings of visual response features in the dorsal lateral geniculate nucleus (dLGN) reveals that *Rhabdomys* attains higher spatiotemporal acuity both by denser coverage of the visual scene and a selective expansion of elements of the code characterised by non-linear spatiotemporal summation. Comparative analysis of single cell transcriptomic cell atlases reveals that realignment of the visual code is associated with increased relative abundance of bipolar and ganglion cell types supporting OFF and ON-OFF responses. These findings demonstrate how changes in retinal cell complement can reconfigure the coding of visual information to match changes in visual needs.

Introduction

Vertebrate central nervous systems share common structural, developmental and genetic blueprints. This standard architecture must accommodate species-specific differences in dominant sense(s), modes of locomotion, and strategies for food seeking, predator avoidance, and reproduction. Such profound differences in functional requirements can exist for species that are closely related by phylogeny, highlighting the need for plasticity in the common blueprint of the brain.[1] Recent technological developments allow exploration of how such adaptations arise with unprecedented resolution. On the one hand, large-scale recordings of neuronal populations facilitate an unbiased comparison of information channels across species[2,3], linking coding strategies to ecological niches. On the other, high-throughput single cell transcriptomics allows comprehensive identification of cell types and cross-species comparisons[4-6]. With these techniques in hand, it becomes possible to describe the process of central nervous system evolution by tracing changes in cell type complement and adaptations in computational characteristics.

Here we apply this approach to a comparison of the early visual system in two closely related murid
rodents. The laboratory mouse (*Mus musculus*) is a popular model for understanding mammalian
vision in health and disease and for establishing general principles of neural function. Like most murid
species, *Mus* are predominantly nocturnal, and this is reflected in key features of the visual system,
including a rod dominant retina and a UV transmitting lens[7]. However, some Muridae are diurnal,
including the four striped mouse, *Rhabdomys pumilio*, of sub-saharan Africa[8,9]. The switch to
diurnality in *Rhabdomys* has been associated with substantial changes in visual system anatomy[10,11].
The *Rhabdomys* retina is cone-dominated, its lens absorbs UV light, and both its retina and visual
centres in the brain are expanded in volume compared to *Mus*[10-12]. Moreover, an unbiased analysis of
the *Rhabdomys* genome reveals that genes involved in vision exhibit accelerated evolution compared
to related murids[13].

Rhabdomys and Mus represent a case study in how differences in temporal niche are reflected in
visual system anatomy[14]. However, an important unanswered question is how such anatomical
expansion impacts the visual code. In principle, the enhanced capacity to process visual information
in *Rhabdomys* could allow: the same computations to be performed with higher precision; a wider
array of visual features to be extracted, describing the scene with greater granularity; or
reconfigurations of the visual code towards features of particular ecological importance. Similarly,
contributors to the anatomical expansion remain unclear. For example, both inner nuclear and
ganglion cell layers are thicker in the *Rhabdomys* retina[10-12], but the increased cell number might be
accompanied by altered frequencies of cell types shared with *Mus,* emergence of new types, or both.
Addressing these questions will help reveal how the blueprint of neural circuits can be adjusted to
align computation to changing ecology.

To address these issues, we combined high-density electrophysiological recordings of visual
response properties in the dorsal lateral geniculate nucleus (dLGN; primary visual thalamus) of *Mus*
and *Rhabdomys* with comparative analyses of retinal cell atlases from these species. We find that the
enhanced capacity of the *Rhabdomys* visual system is applied not only to sample the scene with
higher density but also to facilitate a realignment towards nonlinear spatiotemporal summation. This
functional realignment can be traced to adjustments in the relative abundance of multiple bipolar and
retinal ganglion cell types. Our data thus show how evolution can shape computation by selective
expansion of cell types within the framework of a common neural circuit blueprint.

Results

Realignment in thalamocortical visual code between *Mus* and *Rhabdomys*
Based on the assumption that thalamocortical vision may more reliably reflect inter-species
differences in visual ecology than parts of the visual system responsible for reflexive behaviours such
as optomotor coordination, pupil regulation and circadian entrainment, we set out to compare the
visual code in *Rhabdomys* and *Mus* dorsal lateral geniculate nucleus (dLGN; principal relay of visual
information to the cortex)*. Using a 256-channel recording electrode across multiple placements in
both sagittal and coronal orientations we recorded visually evoked activity for neurons across the
dLGN of both species (Figure 1a; Supplementary Figure 1). To broadly survey the diversity of visual
responses in the two species, we applied a full field temporal modulation stimulus (Figure 1b)
previously applied in both the retina and dLGN of *Mus* to identify functional response types[15]. dLGN
neurons in both species showed strong yet diverse responses to this stimulus (n=611 and n=422 units
in *Rhabdomys* and *Mus,* respectively; Figure 1).

To explore the characteristics and numbers of response types within each species, we pooled light
responsive units from *Rhabdomys* and *Mus,* and extracted isolated response features from the
peristimulus histograms (PSTH) of light-responsive neurons using a sparse principal component
analysis (sPCA; Supplementary Figure 2; see Methods). We then clustered the data by applying a
Gaussian mixture model to the feature set and optimised the cluster number by minimising the

Bayesian Information Criteria (Supplementary Figure 2b). This approach produced 15 functional
clusters, which varied in their response sign, temporal frequency tuning and/or contrast sensitivity
function (Figure 1b-c; Supplementary Figure 3 and 4). Mapping the location of all recorded units
across the dLGN ruled out any systematic sampling bias across clusters (Supplementary Figure 1c).

As expected from earlier work, we found response types corresponding to transient ON (clusters 4, 5,
9, 13) and transient OFF (10, 11), sustained ON (12, 14, 15), and ON-OFF responses (1, 2, 3, 7, 8).
Within those categories, there was further diversity in temporal and contrast sensitivities. Clusters 1,
3, 5, 9 and 13 showed band-pass temporal frequency tuning, with the remainder showing low-pass
tuning. Clusters 5, 13 and 14 showed the highest contrast sensitivity (sensitivity defined by half-
maximum of Naka-Rushton curve, see Methods), while 4 and 6 had lowest sensitivity, and clusters 2
and 5 showed saturating contrast sensitivity curves. Suppressed-by-contrast responses were also
identified (clusters 6 and 8).

We next compared the proportion of neurons from *Rhabdomys* or *Mus* that were assigned to each
functional cluster (Figure 1d). Most clusters (12/15) contained units from both *Mus* and *Rhabdomys*,
and where this was the case, the mean response profile remained consistent between species
(Supplementary Figure 2H). However, three types appeared to be unique to one species (1, 3 and 15)
and frequencies differed by >5-fold for others (2, 5, 9, 12, 13 and 14; Figure 1d). Accordingly, the
overall frequency distribution of units across clusters was highly significantly different between
species (chi square test, p<0.001). This pattern was consistent across experimental animals and
robust to controls for cluster number, batch effects and critical analysis parameters (Supplementary
Figure 2C-G).

The 3 most common clusters in *Rhabdomys* (transient ON-OFF; clusters 1-3) were represented by
only 4 units in *Mus.* Conversely the clusters with highest *Mus* representation (sustained ON; 12-15)
were sparsely represented in *Rhabdomys* (21 units across these 4 clusters). To independently verify
these prominent differences between the two species, we separately classified dLGN response units
according to the kinetics and polarity of their response to a 2s step (2s 97% contrast step every 20s)
(Methods; Figure 1e-f; Supplementary Figure 3). The distribution of units across the resultant
categories was again significantly different between species (chi square test p<0.001), confirming that
the most common response type was ON-OFF in *Rhabdomys* and sustained ON in *Mus.*

Among the responses to the chirp element of the stimulus >50% of *Rhabdomys* units but only ~33%
of *Mus* units were from clusters displaying band-pass temporal frequency tuning (1, 3, 5, 9, 13).
Accordingly, calculations across the whole dLGN population confirmed a preference for higher
frequencies in *Rhabdomys* (Supplementary Figure 4). There was a slight but significant shift towards
increased contrast sensitivity in *Rhabdomys* compared to *Mus* (Supplementary Figure 4c, d, g and h).
Mus neurons were overrepresented in clusters with the lowest contrast sensitivity (5, 13, 14), while
Rhabdomys neurons predominated clusters with high contrast sensitivity (4, and 6). Clusters showing
saturating responses (2 and 5) were also predominant in *Rhabdomys*. Suppressed-by-contrast
profiles were detected in both species, albeit at low frequency (clusters 6 and 8).

Rhabdomys dLGN provides higher temporal fidelity and denser coverage of the visual
field
To define the spatial information capacity of *Mus* and *Rhabdomys*, we next mapped receptive fields
using a binary dense noise stimulus (5Hz). Using spike triggered averages (STAs), we were able to
reconstruct receptive fields in 41% and 64% of light-responsive neurons in *Mus* and *Rhabdomys*,
respectively. Spatial receptive fields (RFs) had robust centre responses, which were delineated as
ON or OFF polarity (Figure 1g-i). Robust opposing surround responses were rare in both species,
consistent with previous work in *Mus,* and are not further quantified here. In *Mus,* the mean RF
diameter was consistent with prior estimates (median: 5.54;[16,17]; Figure 1i). In *Rhabdomys*, this value
was only 6% smaller (median: 5.20; Kolmogorov-Smirnov P=0.036) although the distribution of RF

sizes in each species covered a similar range. The *Rhabdomys* dLGN displayed higher temporal
fidelity, with the temporal filter in the receptive field centre having significantly shorter latency in this
species (median = 134ms and 186ms in *Rhabdomys* and *Mus,* respectively; Kolmogorov-Smirnov
test: P<0.001; Figure 1j). Although individual RFs were similar in the two species, the expanded
volume of the dLGN allowed greater overlap of RFs in *Rhabdomys* and therefore denser coverage at
the population level (Figure 1k).

Enhanced non-linear spatial integration in *Rhabdomys* dLGN
The higher frequency of ON-OFF units in *Rhabdomys* than *Mus* suggests that visual responses in this
species may involve increased non-linear spatiotemporal integration[18]. To assess this directly, we
recorded dLGN responses to inverting gratings at 6 spatial frequencies (each at two phases and at 4
orientations; 1Hz; Michelson contrast = 97%) and identified the optimum phase/orientation for each
neuron offline (see Methods). Quantifying the change in firing rate as a function of spatial frequency
revealed a significant increase in the sensitivity of *Rhabdomys* dLGN neurons to higher spatial
frequencies compared to *Mus.* Thus, at their preferred phase/orientation, most *Mus* neurons showed
low-pass tuning across the range of spatial frequencies tested, responding up to around 0.6 cpd
(Figure 2a,c,d), while most *Rhabdomys* neurons remained responsive to inverting gratings at the
highest spatial frequency tested (1.2 cpd; Figure 2b-e), much smaller than the predicted RF size (~3-
8°). Taken together, these results support the preponderance of non-linear spatial summation in
Rhabdomys neurons, i.e. combining visual signals over space in a non-linear fashion. To further
examine this, we quantified the response amplitude to inverting gratings that were larger than the
calculated receptive field size of each neuron (identifying the optimal phase and stimulus orientation).
We then calculated a linearity index (LI), a ratio of the response at the fundamental frequency of the
stimulus (F1, 1Hz), to the second harmonic (F2; 2Hz)) in response to this stimulus in each species
(Figure 2f-o). Around 25% of neurons were 'non-linear' in *Mus* dLGN (similar to[17]), in contrast with at
least 60% in *Rhabdomys*. In both species, non-linear units responded to inverting gratings at spatial
frequencies that were smaller than their receptive field sizes (Figure 2 I,j,n,o) whereas linear units did
not (Figure 2g, h, l, m).

Analysis of the inverting grating responses enabled us to determine the degree of orientation
selectivity by calculating an orientation selectivity index (OSI; methods) for each neuron at its
preferred spatial frequency and phase. The range of OSI values in the *Mus* dLGN was consistent with
previous reports[16,17], and a similar distribution was found in *Rhabdomys* (Figure 2p-r). At a
conservative threshold of OSI=0.5, 10% and 11% of light-responsive neurons in *Mus* and
Rhabdomys, respectively, were classed as orientation selective. In OSI neurons, we found a strong
preference for horizontally orientated bars in *Mus,* whereas *Rhabdomys* showed a more even
distribution around the tested orientations (Figure 2s); however, the low numbers of OS neurons
make it hard to attribute significance to those differences.

Direction selectivity and responses to motion
We applied a moving bar stimulus to explore motion-sensitivity and direction-selectivity. ~90% of light-
responsive neurons in each species responded to this stimulus with a phasic change in firing.
Responsive neurons showed changes in firing at the leading and/or trailing edge of the bar when
moving over its receptive field (see representative neurons in Figure 3a,b). Consistent with the
evidence of enhanced non-linear spatiotemporal summation in *Rhabdomys*, there was a bias towards
responses to both the leading and trailing edge in this species. This could be seen most clearly in the
spike-triggered average within an individual neuron's receptive field (Figure 3c,d). We calculated a
direction selectivity index (DSI) for neurons that responded to the moving bar. In a subset of neurons,
the strength of responses was dependent on the direction of movement (Figure 3b), consistent with
previous reports in *Mus* dLGN[16,17], however there was a similar distribution of DSIs in both species
(Figure 3e). We did not find any statistical difference in the preferred direction of motion between

species (Figure 3f; chi square test p<0.001), though *Rhabdomys* showed a bias towards motion
sensitivity in the dorso-temporal and ventro-nasal directions.

Electroretinogram recordings in *Mus* and *Rhabdomys*

While the visual code is sculpted in the dLGN[19], its fundamental properties are inherited from the
retina. Given the marked increase in appearance of OFF excitation in *Rhabdomys* dLGN we therefore
turned to *in vivo* electroretinography to determine whether this was also apparent in the retinal light
response. Three components of the electroretinogram (ERG) are relevant: the a-wave is derived from
photoreceptors, the b wave from ON bipolars and the d wave from OFF bipolars. In response to brief
flashes of light under dark-adapted conditions, the a-wave was larger in *Rhabdomys* at high
intensities and higher in *Mus* at low intensities, consistent with the relative paucity of rods, which are
more sensitive than cones, in *Rhabdomys* (Figure 4A-C). Similarly, the b-wave implicit time (latency to
peak), which reflects ON bipolar cell activity, was reduced in *Rhabdomys*, which would be consistent
with a reduced contribution of the more sluggish rod pathway to ON responses. Turning to the
question of OFF excitation, we recorded ERGs in response to an extended (250ms) step under light
adapted (cone isolating) conditions. This stimulus can reveal the activity of ON and OFF cone BCs as
separate b- and d-waves associated with the appearance and disappearance of the light step
respectively. In accordance with the literature, we found that the light pulse ERG is dominated by the
b-wave in *Mus* (Figure 4E-G). Conversely, in *Rhabdomys*, d-waves were at least as prominent (Figure
4E-G) consistent with a strong contribution of OFF bipolar cells and a realignment towards OFF
excitation from the earliest step of visual signal transduction in this species.

Comparison of *Mus* and *Rhabdomys* retinal cell classes and types

To understand the distinct coding properties in the primary visual pathway of *Rhabdomys* and *Mus* we
compared retinal neuronal types of the two species, using atlases derived from single-cell and single-
nucleus RNA-seq[20-23]. The *Rhabdomys* atlas[24] (Figure 5) contained 65,930 nuclei from 2 *Rhabdomys*,
which could be classified via standard computational procedures into the five retinal neuronal classes
(photoreceptors [PR], horizontal cells [HC], bipolar cells [BC], amacrine cells [AC] and retinal ganglion
cells [RGC]) as well as multiple glial types (Muller, Astrocyte, and Microglia), endothelial and retinal
pigment epithelial cells (Figure 5a). Each neuronal classes could be further divided into
transcriptomically distinct clusters: 3 PR, 1 HC, 18 BC, 33 AC and 33 RGC clusters (Figure 5b).
Altogether, the atlas included over 100 retinal clusters, representing putative cell types. Within each
class, nearly all *Rhabdomys* clusters mapped in a specific fashion to *Mus* types (see below). This
correspondence allowed us to transfer cell type labels of the better-studied *Mus* to *Rhabdomys*,
facilitating comparison between the two species.

Rhabdomys retina is cone-dominated

Analysis of photoreceptor transcriptomes confirmed and quantified the known shift in rod:cone ratio
between *Mus* and *Rhabdomys*: ~3% of *Mus* photoreceptors but >65% *Rhabdomys* photoreceptors
are cones, a >20-fold difference (Figure 5c). Also anticipated was the degree of cone opsin co-
expression. ~40% of *Mus* M cones express S-opsins at low levels[25], whereas this was true for only
~1.3% of *Rhabdomys* M cones (112 out of 8622 cones), with the majority (92.7%) expressing only M-
cone opsin.

A reduction in rod BCs and increased proportion of OFF BCs in *Rhabdomys* inner retina

BCs can be subdivided into those that depolarize (ON) or hyperpolarize (OFF) to illumination. Cones
innervate both ON and OFF BCs whereas all rod BCs are ON BCs. Clustering the ~11,400
Rhabdomys BC transcriptomes generated 18 putative types (Figure 6a). Based on known markers,
these types comprised 7 OFF cone BC types, 9 ON cone BC types, and 1 rod BC type. The final type,
BC1B, receives no direct photoreceptor input[20,26] so cannot be confidently classified as either ON or
OFF. A supervised classification analysis based on transcriptomic signatures indicated a

predominantly 1:1 correspondence between the *Rhabdomys* BC types and *Mus* BC types (Figure 6b).
The patterns of correspondence were consistent with the results of our recent comparative analysis of
BC types across 13 mammals using an integrative clustering approach[24].

Three notable differences in BC composition between species were evident. First, the 18 BC types in
Rhabdomys exceeded the 15 that have been identified and validated in *Mus*. *Mus* types BC1A, 3B
and 5A each mapped to two *Rhabdomys* types (Figure 6b). Members of each pair (C3 and C18 for
BC1B, C2 and C8 for BC3B, C6 and C9 for BC5A) differentially expressed multiple genes
(Supplementary Figure 5), supporting their identity as distinct types. Second, rod BCs comprised
around 3.5% of the total BCs in *Rhabdomys*, in contrast to ~43% in *Mus*, correlating with the
difference in rod:cone ratio between species (Figure 6c; Supplementary Table 1). Immunostaining
with the RBC marker PKCα validated this difference (Figure 6d). Third, among cone BCs (that is,
separate from the difference in rod BC frequency), there are more OFF BCs and fewer ON BCs in
Rhabdomys than in *Mus*. The four most abundant cone BC types in *Rhabdomys* retina (C1-4; clusters
are numbered in order of their abundance) all corresponded to OFF types in *Mus*. Conversely, all but
one of the ON cone BC types were more abundant in *Mus* (Figure 6e; Supplementary Table 2). This
redistribution of CBC types would predict an enhanced OFF type response at the first retinal synapse,
consistent with the enhanced photopic d-wave observed in the ERG (Figure 4).

Increased abundance of OFF and ON-OFF RGCs in *Rhabdomys*
Reclustering the 26,500 *Rhabdomys* RGCs transcriptomes yielded 33 clusters (Figure 7a), which is
substantially lower than the 45 molecularly distinct RGC types in *Mus*. However, supervised classifiers
trained on either *Mus* or *Rhabdomys* data indicated that all known 45 *Mus* RGC types were
represented among the *Rhabdomys* clusters (Figure 7b). Thus, in several instances a *Rhabdomys*
cluster was composed of a group of closely related *Mus* types. The difference in resolution was likely
due differences in cell number (26539 in *Rhabdomys* vs 35699 in *Mus*), sequencing depth and/or
modality (single-cell vs. single-nucleus).

RGCs are conventionally classified as ON, OFF or ON-OFF types depending on whether they are
excited by increases or decreases in light intensity or both. Previous studies have combined
morphological, physiological, and transcriptomic data to characterize *Mus* RGCs[21,27-29]. Based on
those results, we were able to assign 43 *Mus* types to one of these three categories (Figure 7c).
Lacking physiological data from *Rhabdomys* RGCs, we used our supervised classification model to
provisionally categorize *Rhabdomys* RGCs as ON, OFF or ON-OFF based on their assigned *Mus*
label. The 8 most abundant putative ON types in *Mus* were all under-represented in *Rhabdomys*, and
most OFF or ON-OFF types were more abundant in *Rhabdomys* than in *Mus*, including all four of the
known ON-OFF direction-selective types (ooDSGCs; Figure 7c,d; Supplementary Table 3). This bias
is consistent with the preponderance of OFF BCs in retina and OFF and ON-OFF responses in dLGN
noted above. Interestingly, orthologues of a subclass of *Mus* RGCs defined by expression in the
"THWY3" transgenic line[21,30,31] were enriched in *Rhabdomys* (Figure 7d); 2 of these types are OFF, 2
ON, 1 ON-OFF, and 1 unknown.

Depletion of ipRGCs in *Rhabdomys*
Several types of melanopsin- (*Opn4*-) expressing ipRGCs have been characterized in *Mus* [32-35]. Our
Mus RGC atlas identifies 5 ipRGC types, which we have provisionally called M1a, M1b, Mx, M2 and
M4. Two additional *Mus* RGC types, C7 and C8, are closely related to the ipRGCs; both express the
subclass-defining transcription factor *Eome*s, and both express *Opn4* at low levels[21]. *Mus* ipRGC
types co-mapped to *Rhabdomys* clusters C16 and C27 (Figure 7b). However, for all these types, the
relative abundance was substantially lower in *Rhabdomys* than in *Mus* (Figure 8a), which is the likely
reason that they did not form separate clusters.

To validate these results, we used fluorescent *in situ* hybridisation chain reaction (HCR-FISH) to
compare *Opn4*+ RGCs in retinal wholemounts from the two species (Figure 8b). Using an automated
cell detection method (Supplementary Figure 6), we found that the number of *Opn4*+ cells per unit
area was substantially lower in *Rhabdomys* than *Mus* (Figure 8c,d) indicating that both the absolute
density of ipRGCs (5x lower) and their representation as a function of the total RGC population were
greatly reduced in *Rhabdomys* compared to *Mus*. Quantification of soma size and FISH intensity in
the two species (Figure 8e-g) revealed a bias towards cells with smaller soma sizes in *Rhabdomys,*
and an increased proportion of cells with high *Opn4* expression.

Discussion

We applied physiological and transcriptomic techniques to understand how the visual systems of
closely related murid rodents are adapted to divergent visual ecology. Diurnal *Rhabdomys* have
thicker inner retinas and larger dLGNs than nocturnal *Mus*. Our data reveal that although they apply
this extra information capacity to provide denser coverage of the visual scene, this does not amount
to simply 'more of the same' visual code seen in *Mus*. Rather, we find a marked realignment towards
nonlinear spatiotemporal summation in the *Rhabdomys* dLGN. This is apparent in a shift from
predominantly ON responses in *Mus* towards biphasic, ON-OFF, responses and increased
abundance of units able to track gratings sufficiently fine to fall entirely within their receptive field
centre in *Rhabdomys*. These inter-species differences in visual code are accompanied by selective
expansions in the abundance of OFF BCs and RGCs with transcriptional fingerprints of ON-OFF
response types in *Rhabdomys* compared to *Mus*.

In principle, the enhanced information capacity of the *Rhabdomys* early visual system could have
been employed in numerous ways. One option would have been to transmit a higher fidelity, less
filtered representation of the scene to the cortex. That would facilitate richer computations and
interpretation at higher levels and is the strategy often employed by other species with enhanced
visual systems such as primates, where a relatively unprocessed visual code is coupled with an
expansion of the visual cortex[36-38]. However, we find that such simple 'pixel encoders' (characterised
by sustained responses and linear spatial summation) are substantially less common in *Rhabdomys*
than *Mus* (Supplementary Figure 3c). At the other extreme, *Rhabdomys*' extra capacity could
plausibly be applied to allow more diverse computations in the early visual system, parsing the visual
scene in more complex ways [39,40]. Our cell atlas analysis provides some support for that proposition,
as there is evidence of increased diversity in the BC population with *Mus* subtypes BC1A, 3B and 5A
each represented by two transcriptionally distinct clusters in *Rhabdomys*. However, we did not
observe a corresponding increase in diversity in the *Rhabdomys* visual code. Thus, unbiased
clustering returned around 14 functionally distinct channels in the dLGN of each species (with the
proviso that more complexity is expected with a wider array of visual stimuli and/or in distinct RGC
targets).

Given the alternatives, what advantage might *Rhabdomys* attain by selectively increasing
representation of units with non-linear spatiotemporal summation, especially as such non-linear
feature detectors are often more abundant in the retinas of animals that invest less neural resources
in (form) vision (e.g.[30]). One possibility is that it represents an efficient approach to enhance
spatiotemporal resolution. We find that RF sizes are only marginally finer in *Rhabdomys* compared to
Mus. With this constraint in mind, natural selection can take two routes to exploit the additional
information of *Rhabdomys* to increase spatiotemporal acuity. On the one hand, it can increase the
degree of overlap in RF between units to achieve denser coverage of the scene at neuronal
population level. We find that this is indeed a feature of the *Rhabdomys* visual code. Alternatively, it
can employ non-linear spatiotemporal summation to allow individual neurons to respond to patterns
within their RF, achieving greater acuity at the single unit level. *Rhabdomys* appear to exploit this
opportunity too, as we find that realignment of the visual code towards non-linear summation allows
many units to respond to high frequency gratings.

The primary features of the thalamic visual code are inherited from the retina, and species differences
in the visual code that we observe are accompanied by some substantial differences in retinal cellular
composition. Most obviously, the *Rhabdomys* retina has many more cone photoreceptors and this is
achieved by selective expansion of the M-cone opsin expressing cones (the fraction of photoreceptors
expressing S-cone opsin hardly differs between species). The shift towards cones is expected for a
more day active species. That this is restricted to M-cone opsin expressing cones is perhaps
associated with *Rhabdomys*' combination of UV sensitive S-cone opsin with a lens that filters out UV-
wavelength light[11].

Second order neurons are also substantially different in *Rhabdomys* with no overlap in the 3 most
abundant BC types in each species. The switch to a cone-rich retina in *Rhabdomys* is accompanied
by a large decrease in the relative proportion of rod BCs. This may be predicted given the cone bias
of the *Rhabdomys* retina, but recent analysis reveals that rod BCs can be numerous even in species
with high cone density[24]. Evidence that cones can signal via rod BCs supports the contention that rod
BCs do not necessarily become redundant when rod numbers fall. Nevertheless, both anatomical and
physiological characteristics of cone BCs facilitate information transfer at higher spatiotemporal
resolution.

The decrease in rod BC frequency in *Rhabdomys* compared to *Mus* is accompanied by an increase in
cone BC frequency. However, the increase is not distributed equally across cone BC types. Instead, it
appears disproportionately in OFF BCs (in order of fractional abundance BC3B, BC4, BC1A, BC3),
while cone ON BCs occupy equivalent fractions of the BC population in the two species. The
enrichment of OFF BCs in *Rhabdomys* is consistent with the enhanced ERG d-wave amplitude
(measuring OFF responses) and the preponderance of OFF excitation responses in the dLGN of this
species. A plausible function of the increase in OFF BC frequency facilitates non-linear
spatiotemporal summation and, indeed, the two most expanded BC classes in *Rhabdomys* (BC3B
and BC4) have characteristics well suited to this purposes (small dendritic and axonal fields and
transient responses[41,42]). But realignment towards OFF BCs may bring additional benefits. OFF BCs
respond faster than ON BCs, allowing the possibility not only for faster visual reflexes but also higher
acuity[43]. As a day active species, *Rhabdomys* may place a higher premium on detecting shadows
(attributed to OFF pathways[44]) and in a more general sense the negative spatial contrast to which
natural scenes are biased[45] than high sensitivity vision (attributed to ON pathways[46]). Finally, the
enhanced capacity of OFF pathways could support better ability to detect overhead (dark) looming
stimuli indicative of aerial predators[47].

Species differences in the RGC population are more nuanced than those in photoreceptor or BC
makeup but are in accordance with the changes in visual code that we observe. Thus, in the
Rhabdomys atlas RGC types characterised by ON responses in mice are relatively under-represented
in favour of those with OFF and ON-OFF responses. The population of ON type, melanopsin-
expressing, ipRGCs provides a noteworthy example of this realignment. Not only is the density of
Opn4-positive RGCs in the *Rhabdomys* retina ~5-fold less than *Mus*, but their anatomical features
suggest a particular reduction in those ipRGCs contributing to thalamocortical vision. Compared to
Mus, *Rhabdomys* Opn4+ cells are biased towards smaller soma and slightly higher melanopsin
expression. These are characteristics of the M1-3 ipRGCs which subserve reflex responses of
circadian entrainment and pupil constriction in Mus. It follows that cells with the morphological
characteristics of the *Mus* M4 population[34,48] that contribute to sustained-ON responses in the dLGN[49-
51] are especially rare in *Rhabdomys*.

The *Mus* vs *Rhabdomys* comparison represents a case study in how sensory systems can adapt to
different ecology. Despite their similarity in phylogeny, size and diet these species have evolved
markedly different visual systems. *Rhabdomys* is adapted to exploit the higher signal:noise of visual
signals in the day with its cone-rich retina and expansion in the thickness of the inner retina and
dLGN. We find that it applies this additional information capacity in both predictable and more

surprising ways. Thus, in addition to simply achieving greater density of coverage of the visual scene,
changes in composition of the retinal cell population drive a realignment in favour of OFF responses
and non-linear spatiotemporal summation, which increase spatial resolution of the visual code and
may provide other benefits. Our data thus show how evolutionarily advantageous changes in
computational outcome can be produced by selective expansion/contraction of cell types comprising a
neural circuit.

Methods

Animals

396
397 Animal care was in accordance with the UK Animals, Scientific Procedures Act of 1986, and the study
398 was approved by the University of Manchester ethics committee. Animals were housed on a 12h:12h
399 light:dark cycle at 22°C with food and water available ad libitum. All experiments were performed in
400 adult *Rhabdomys pumilio* or C57BL6J mice (aged 3–8 months).

In vivo electrophysiology

In vivo electrophysiological recordings were performed in 6 *Rhabdomys* and 5 *Mus* (male), using
methods described previously[11]. Anaesthesia was induced with 2% isofluorane in oxygen, and
maintained with an intraperitoneal injection of urethane (1.6 g kg–1, 30% w/v; Sigma-Aldrich). A
topical mydriatic (tropicamide 1%; Bausch and Lomb) and mineral oil (Sigma-Aldrich) were applied to
the left eye prior to recording. After placement into a stereotaxic frame, the skull was exposed and a
small hole drilled ~2.5 mm posterior and ~2.5 mm lateral to bregma (*Rhabdomys*); or ~2.3 mm
posterior and ~2.3 mm lateral to bregma (Mus). A 256-channel recording probe (A4x64-Poly2-5mm-
23s-250-177-S256, NeuroNexus Technologies, Inc., Ann Arbor, MI, USA) consisting of 4 shanks
spaced 200 µm apart, each with 64 recording sites, was lowered a depth of ~3–3.5 mm into the
Rhabdomys brain, or 2.5-3mm into the *Mus* brain, targeting the dLGN in each species. Broadband
neural signals were then acquired using a SmartBox recording system (NeuroNexus Technologies,
Inc.), sampling at 20 kHz. Following recordings, data from each of the four electrode shanks were pre-
processed by common median referencing, high-pass filtered at 250 Hz and then passed to an
automated template-matching-based algorithm for single unit isolation (Kilosort; [52]). Isolated units
were then extracted as virtual tetrode waveforms for validation in Offline Sorter (V3, Plexon, Dallas,
TX, USA). Here, unit isolation was confirmed by reference to MANOVA F statistics, J3 and Davies-
Bouldin validity metrics and the presence of a distinct refractory period (greater than 1.5 ms) in the
interspike interval distribution. Spike sorted data were further analysed in MATLAB R2018a (The
MathWorks).

Visual stimuli

Responses were recorded to a standardised set of temporally and spatially patterned monochromatic
stimuli, displayed using an LCD display (width: 26.8cm height: 47.4cm; Hanns-G HE225DPB ; Taipei,
Taiwan) angled at ~45° from vertical and placed at a distance of ~21cm from the contralateral eye to
occupy ~96° x ~63° visual angle. The temporal stimulus set consisted of a 2s step from minimum to
maximum light intensity (98% contrast), followed by 2s of dark, 2s at half maximum light intensity, an
8s temporal chirp (sinusoidal modulation between dark and maximum intensity at 1-8Hz accelerating
at rate of 1Hz/s), 2s at half maximum light intensity, and an 8s contrast chirp (sinusoidal modulation at
2Hz increasing from 3% to 97% contrast), as in[15]. Spatial stimuli comprised of a sparse binary noise
stimulus (5Hz, square size = 4.2°); inverting grating stimuli (1Hz) at spatial frequencies of 0.03 to 1.2
cpd, presented at two phases and four orientations (0°, 45°, 90° and 135°); and a single bar (4.2°)
moving in one of 8 directions (0°, 45°, 90°, 135°, 180°, 225°, 270° and 315°) in a pseudorandom
sequence. Stimulus spectra were designed to approximate the activation of each photoreceptor by
natural daylight for each species (14.8 MWS effective photons/cm^2/s; 12.8/12.0 SWS effective
photons/cm^2/s for *Mus* and *Rhabdomys*, respectively; 14.8 rod effective photons/cm^2/s and 14.7
melanopsin effective photons/cm^2/s).

Analysis of dLGN responses to visual stimuli

Full field stimuli

Perievent spike histograms (PSTH) were generated with bin size of 30ms. Light responsive units were identified using confidence limits test based on responses to the initial 2s step of the chirp stimulus: units were classified as significant if spike firing rate during the response window was greater than 2 standard deviations above (excitation) or below (inhibition) mean firing rate during baseline window – equivalent to 95% confidence limit. ON:OFF Bias index and Sustainedness index were calculated using previously described methods (Farrow & Masland, 2011; Lindner et al., 2021).

To analyse temporal chirps, the mean response amplitude (maximum – minimum normalised firing rate) for each temporal frequency was fit with a half-gaussian model ([16]) using least-squares minimisation to identify 5 best-fit parameters (low baseline, high baseline, gaussian spread, peak response and peak frequency).

Equation for Half Gaussians is:

$$Response = b_1 + (a - b_1) * e^{-\left[\frac{p-w}{s}\right]^{\wedge}2} \text{ for } w < p$$

$$Response = b_2 + (a - b_2) * e^{-\left[\frac{p-w}{s}\right]^{\wedge}2} \text{ for } w > p$$

Where w is the temporal frequency (Hz), p is the temporal frequency (TF) that produces peak response, a is the maximum response amplitude at optimum TF, s is the Gaussian spread, $b1$ is the baseline for frequencies lower than peak TF, $b2$ is the baseline for frequencies greater than the peak TF. Peak temporal frequency was rounded to nearest integer to address the limited resolution of temporal frequency analysis (sample every 1Hz).

For contrast chirps, the response amplitude (maximum – minimum normalised firing rate during each period of the contrast chirp stimulus) was normalised to baseline activity (1s before contrast chirp onset) for each unit. This was plotted against Michaelson contrast and then fit using a Naka-Rushton curve using least-squares minimisation to identify 4 best-fit parameters (top, bottom, C50 and slope).

Equation for Naka-Rushton curve is

$$Response = Bottom + \left(Top * \frac{C^n}{C^n + C50^n}\right)$$

Where n = slope, C = Michelson contrast and C50 is contrast that produces half maximum response. C50 was constrained between 0 and 1, and slope was constrained between 0 and 10.

Functional clustering

Sparse Principal components were generated for the full-field temporal stimulus using the SPaSM toolbox[53], as described in [15]. This allows the extraction of response features that are localised in time. We pooled mean PSTH (25ms bins) for all light responsive units from both groups and extracted up to 30 features with 5 non-zero time bins. We then discarded those that accounted for < 1% of the variance. Response features that met these criteria for each window were then combined to produce a total of 30 features for dLGN. sPCs from integrated *Mus* and *Rhabdomys* data were then clustered with a mixture of Gaussian models, a probabilistic model using random initialisation. The optimum number of clusters was determined based on the lowest Bayesian information criteria, which rewards fit but penalises complexity, and a Bayes factor below 6 as a threshold for when there was no longer evidence for further splitting.

To compare distribution of units across communities between two groups, we calculated the distance (Euclidian norm of the difference in the mean relative proportion of neurons in each cluster) between

477 *Mus* and *Rhabdomys* data and compared this with a null distribution obtained by randomly shuffling
478 retinal recordings between two groups 10,000 times. To compare proportion of units from each group
479 within each cluster, we first calculated the % of total units from each group in a given community for
480 each recording, and then found the difference between mean of *Mus* and *Rhabdomys*. This was
481 compared with a null distribution generated as above.

Receptive Field mapping

The spatio-temporal receptive field was derived for each unit by generating the spike triggered
average (STA) of responses to a sparse binary noise stimulus (5Hz, square size = 4.2°). The
separable spatial and temporal components where then extracted from the raw STA matrix by finding
the signal peak. RF locations and sizes were then generated by fitting spatial receptive fields with 2D
Gaussian function (using lsqcurvefit function, MATLAB). The receptive field size for individual cells
was the average of the standard deviation of Gaussians fitted to each dimension. Temporal receptive
fields were generated by plotting the temporal response of the RF centre. Receptive field overlap was
calculated as a percentage overlap of the 2D Gaussian for pairs of receptive fields recorded in the
same animal.

Spatial frequency tuning and linearity of spatial summation

To assay changes in spatial frequency tuning, inverting gratings (Michelson contrast between dark
and light bars = 98%) were presented in 4 orientations at two phases (phase shifted 90°), at 5
different spatial frequencies (0.03 to 1.2 cpd) at 1Hz. For each unit, response amplitudes were
quantified (relative to pre-stimulus firing) for each phase/orientation combination for each spatial
frequency to determine the optimal stimulus (R_{pref}). R_{orth} was the response to stimuli presented at 90°
to the preferred orientation. The orientation selectivity index (OSI) was calculated as the ratio of (R_{pref}-
R_{orth})/(R_{pref}+R_{orth}) at the preferred spatial frequency. Cells exceeding an OSI of 0.5 were classed as
'orientation selective'.

Response linearity was evaluated by quantifying the firing rate of units in response to stimuli that were
larger than the calculated receptive field size for an individual unit. Continuous firing rates during the
stimulus presentation were Fourier analysed to extract amplitudes of the first and second harmonic
components (F1 and F2, at 1Hz and 2Hz), at the preferred and null (90° phase shifted) stimulus
($F1_{pref}$, $F1_{null}$, $F2_{pref}$, $F2_{null}$). A linearity index (LI) of the response was then calculated as F1/F2 for both
preferred and null phases, whereby a LI<1 describes a dominant F2 amplitude, indicating nonlinear
spatial summation.

Motion selectivity

A single drifting bar (4.2°; Michelson contrast= 98%) moving in one of 8 directions (0°, 45°, 90°, 135°,
180°, 225°, 270° and 315°) in a pseudorandom at a speed of . For each unit, response amplitudes
were quantified during the presentation of movement (relative to pre-stimulus baseline) for each
direction of motion. Since the location of the stimulus relative to the location of a unit's receptive field
was not precisely known, to explore the kinetics of responses, a STA was generated (as described
above) for the preferred direction of motion. For presentation purposes, responses were clustered
using a Kmeans cluster (MATLAB) finding 3 response clusters.

Two methods were used to explore direction selectivity. First, the mean and variance of circular data
were computed using the CircStat, a MATLAB toolbox[54], to describe the angle and magnitude of
directional selectivity. A direction selectivity index was also calculated as the ratio of (Rpref-
Rnull)/(Rpref+Rnull), where Rpref was the direction of motion at which the maximum evoked
response occurred, and Rnull was response to movement in the opposite direction to this. Cells
exceeding a direction selectivity index of 0.33 were classed as 'direction selective'.

Electroretinography

ERGs were recorded in adult *Mus* (aged 4-5 months) and adult *Rhabdomys* (aged 7-8 months). All
ERGs were recorded at subjective midday following dark adaptation for a period of 6 hours. Animals
were anaesthetised under isoflurane in a 95/5% Oxygen/CO2 mix at a flow rate of 0.5 – 1.0L/min.
Isoflurane concentrations of 5% and 1.5-3.5% were used for induction and maintenance of
anaesthesia, respectively. A topical mydriatic (tropicamide 1%; Bausch and Lomb) and hypromellose
eye drops were applied to the recording eye before placement of a corneal contact-lens–type
electrode. A needle reference electrode (Ambu, Neuroline) was inserted approximately 5mm from the
base of the contralateral eye, and a second subcutaneous needle in the scruff acted as a ground.
Electrodes were connected to a Windows PC via a signal conditioner (Model 1902 Mark III, CED) that
differentially amplified (X3000) and filtered (band-pass filter cut off 0.5 to 200Hz) the signal, and a
digitizer (Model 1401, CED). Core body temperature was maintained at 37°C throughout recording
with a homeothermic heat mat (Harvard Apparatus).

Visual stimuli were generated with a combination of violet, blue and cyan elements of a multispectral
LED light source (Lumencor). Intensities were modulated via pulse width modulations via an Arduino
Uno. Light from the light engine passed through a filter-wheel containing neutral-density filters
(reducing the light by between 10^1 and 10^5) and focused onto opal diffusing glass (5mm diameter;
Edmund Optics Inc.) positioned <5mm from the eye. All LED intensities were controlled dynamically
with a PC. Stimuli were measured at the corneal plane using a spectroradiometer (SpectroCAL II,
Cambridge Research Systems, UK) between 350-700nm. Dark-adapted stimuli were presented as a
10ms flash of stimulus spectra from background across a range of 9.1 to 15.8 photons/cm^2/s
(interstimulus intervals ranging from 1-6s with increasing intensities). Light-adapted stimuli were
presented as square-wave modulations from background at 80.5% Michelson contrast at 2Hz at a
background of 14.6 log photons/cm^2/s, after 15 minutes background adaptation. ERG responses were
analysed in MATLAB. For flash responses, a-wave amplitude was calculated relative to baseline prior
to stimulus onset, with b-wave with reference to the a-wave trough. For step responses, b- and d-
waves were calculated relative to preceding a-wave.

Immunohistochemistry
Immunostaining of *Mus* and *Rhabdomys* retinal sections and wholemounts was performed as
described previously[11]. *Rhabdomys* retinal wholemounts were labelled using rabbit anti-melanopsin
antibody (UF006, Advance Targeting Systems, 1:2000) to label ipRGCs. Retinal sections were
labelled using rabbit anti-PKCα (ab32376, Abcam, 1:1000) to label rod bipolar cells. Sections and low
magnification wholemount retinas were imaged with an Axio Imager D2 upright microscope and
captured using a Coolsnap Hq2 camera (Photometrics) through Micromanager software v1.4.23. High
magnification images were collected using an inverted LSM 710 laser scanning confocal microscope
(Zeiss) and Zen 2009 image acquisition software (Zeiss).

Fluorescent in situ hybridisation chain reaction
Mus (n=3) and *Rhabdomys* (n=3) retinas were dissected and used for HCR[TM] RNA-FISH (Molecular
Instruments) according to the manufacturer protocol for fixed wholemount tissue. Briefly, retinas
underwent a series of dehydration and rehydration steps (75%, 50%, 25% methanol solution) and
then treated with proteinase K (10 µg/mL). Retinas were pre-hybridized with probe hybridization
solution and incubated overnight in probe solution containing custom ordered probes *Mus* or
Rhabdomys Opn4 (2 pmol). Retinas were pre-amplified in amplification buffer and incubated
overnight in amplification solution containing hairpins H1 and H2 (30 pmol each; amplification
fluorophore 594). Retinas were washed in sodium chloride sodium citrate tween 20 solution and
stored at 4 ∘C before imaging. Cells were detected using QuPath[55] cell detection feature.

Analysis of Transcriptomic Datasets

Alignment and quantification of gene expression

Preprocessing of raw sequencing data was performed using Cellranger (v6.2, 10X Genomics). Sequencing reads were demultiplexed using "cellranger mkfastq" to obtain a separate set of fastq.gz files for each of the 7 samples. These files were then aligned to a reference genome [13] using "cellranger count" with the --include-introns flag to include both exonic and intronic reads, resulting in a gene expression matrix (GEM) summarizing transcript counts within each sample. GEMs from each of the 7 samples were combined (column-wise concatenated) to yield a total GEM.

Segregation of major retinal cell classes

Analysis of the total GEM was performed in R, with the workflow based on Seurat v4.3.0 for single-cell analysis developed and maintained by the Satija laboratory [56] (https://satijalab.org/seurat/). Transcript counts in each cell were normalized to a total library size of 10,000 and log-transformed $(X \to \log(X + 1))$. We identified the top 2,000 highly variable genes and applied principal component analysis (PCA) to obtain a linear factorization of the submatrix corresponding to these highly variable genes. Using the top 20 principal components for each cell, we built a k-nearest neighbor graph on the data, and then clustered with a resolution parameter of 0.5 using Seurat's FindClusters function. Each cluster of cells was assigned to a retinal cell class based on canonical markers characterized in mice[22]; for example, *Vsx2*, *Otx2* and *Grik1* were used to identify bipolar cells, and *Rbpms*, *Nefl* and *Nefm* were used to identify RGCs.

Supervised classification analysis of transcriptional correspondence between Rhabdomys and Mouse types

RGCs were separated from the *Rhabdomys* total GEM and, as the *Rhabdomys* genome was annotated with mammalian orthologs across murid genome assemblies, were merged with a reference *Mus* RGC atlas[21] using genes that were present in both the *Rhabdomys* GEM and *Mus* reference GEM to create a joint GEM. We used the Canonical Correlation Analysis-based framework in Seurat, integrating by species of origin, to produce an integrated version of the joint GEM adjusted to account for species specific differences in gene expression. The top 2000 variable features of the integrated GEM were used to train a gradient boosted decision tree using the reference *Mus* RGC cells, implemented in R using the xgboost package[57]. The *Mus* RGC-trained classifier was used to assign a *Mus* identity to each *Rhabdomys* RGC based on its expression of the 2000 training features. To identify how rare *Mus* RGC types mapped to *Rhabdomys* types, a similar decision tree model was trained using the *Rhabdomys* RGC cells, and applied to *Mus* RGCs to assign a *Rhabdomys* identity to each *Mus* RGC. Similarly, this was done with *Rhabdomys* bipolar cells with a relevant *Mus* bipolar cell atlas to assign *Mus* bipolar labels to each *Rhabdomys* bipolar cell, and vice versa. These reciprocal mappings helped us validated the robustness of the label transfer procedure.

To summarize the results of the supervised classification mapping, we calculated a modified Jaccard index for each pair of *Rhabdomys* and assigned *Mus* types. For a given *Rhabdomys* type A and assigned *Mus* type B, we calculated

$$J(A, B) = \frac{|A \cap B|}{\min(|A|, |B|)}$$

Data and Code Availability

Analysis scripts for the *Rhabdomys* snRNA-seq data is available via Zenodo (https://zenodo.org/record/8067826) and on our Github (https://github.com/shekharlab/RetinaEvolution). The raw and processed sequencing data produced

611 in this work are available via the Gene Expression Omnibus (GEO) under accession number
612 GSE237210.

Acknowledgements
614 This work was supported by a Sir Henry Dale Fellowship 218556/Z/19/Z, jointly funded by the
615 Wellcome Trust and the Royal Society (AEA); a Wellcome Investigator Award 210684/Z/18/Z (RJL);
616 NIH grant U01MH105960 (JRS); and the McKnight Foundation (KS).

Author contributions
AEA, JS, KS and RJL supervised the project; AEA performed electrophysiological data collection and
analysis, with contributions from JM, JR, PO and RS; AEA performed ERG data collection and
analysis with contributions from BBO; RR, AP and AEA performed HCR-FISH experiments and
analysis; CW and JW performed retinal immunohistochemistry, with contributions from NM; JH
performed computational analysis of scRNA-seq data with contribution from WY. AM performed
scRNA-seq experiments. AEA, JH, JS, KS and RJL wrote manuscript with input and approval of all
authors.

Declaration of interests
626 The authors declare no competing interests

 Figures and legends

 <u>**Figure 1**</u>

dLGN response diversity in _Mus_ and _Rhabdomys_

a) Above: coronal sections from _Mus_ and _Rhabdomys_ with dLGN highlighted with orange edging. Below: Histological reconstruction of recording sites in _Rhabdomys_ dLGN. Fluorescence shows diI markings remaining after electrode insertion. Peristimulus time histograms (PSTHs) representing multiunit light-evoked activity recorded at each recording site overlayed (2s light step with 1s pre and post stimulus). All data are scaled according to their maximum firing rate. **b**) Heat map showing normalised PSTHs of dLGN neurons as a function of response cluster (labelled to right), organised as a function of number of _Rhabdomys_ responses. Stimulus profile denoted above. **c**) Mean response profile of normalised responses from each cluster (combined across species). Stimulus profile denoted above. **d**) Bar graphs showing proportion of responses within each cluster, for each species (_Mus_: cyan, _Rhabdomys_: red). **e**) Histogram of ON:OFF bias in _Mus_ (cyan) and _Rhabdomys_ (red). **f**) Histogram of sustainedness index in _Mus_ (cyan) and _Rhabdomys_ (red). **g&h**) Example spatial **(g)** and temporal (**h**) spike triggered averages recorded in response to binary noise stimulus. **i**) Histogram of receptive field sizes in _Mus_ (cyan) and _Rhabdomys_ (red); Kolmogorov-Smirnov test P=0.036. **j**) Histogram of receptive field latency (time of peak absolute response) in _Mus_ (cyan) and _Rhabdomys_ (red); Kolmogorov-Smirnov test: P<0.001. **k**) Histogram of receptive field overlap between pairs of neurons recorded in _Mus_ (cyan) and _Rhabdomys_ (red); Kolmogorov-Smirnov test: P<0.001.

647
<u>Figure 2</u>
Spatial frequency tuning and linearity in *Mus* and *Rhabdomys*.
a) Schematic of inverting grating stimuli of increasing spatial frequency. **b)** Mean+/-SEM PSTH for
1Hz inverting grating stimuli at 6 spatial frequencies (0.0375, 0.075, 0.15, 0.3, 0.6, 1.2cpd) for *Mus*
(left) and *Rhabdomys* (right) **c)** Amplitude of response to increasing frequency inverting gratings at
preferred orientation/phase, shown for *Mus* (cyan) and *Rhabdomys* (red); data show mean+/- SEM.
d&e) Distribution of % of neurons with peak response at a particular frequency (**d**); and with a
threshold to response at a particular spatial frequency (**e**), shown for *Mus* (cyan) and *Rhabdomys*
(red). **f,** Distribution of linearity index (LI) in *Mus*. **g,** Mean normalised response of linear neurons in
preferred (top panel) or null (lower panel) phase, in response to 2 grating inversions. Shaded regions
indicate inversions. **h,** fft amplitude in preferred and null phase of the response, for spatial frequencies
greater than (left) or less than (right) the receptive field size, for F1 (black) and F2 (green) frequency
components. **i&j,** as in **g&h** but for non-linear responses in *Mus*. **k-o,** as in **f-j**, but for *Rhabdomys*. **p)**
Distribution of orientation selectivity index (OSI) for *Mus*. **q&r)** mean response of *Mus* (**q**) and
Rhabdomys (**r**) neurons with OSI<0.5 (left) and OSI>0.5 (right) in response to inverting grating at
optimal spatial frequency, in preferred (black) and null (red) phase. **s,** distribution of preferred
orientation of bars for OS neurons (note double plot from 180-360▢) for *Mus* and *Rhabdomys* (top
and bottom panels, respectively).

666
<u>Figure 3</u>
Direction selectivity and responses to motion
a&b) Representative neurons from *Rhabdomys* with low (DSI = 0.12) or high (DSI = 0.44) DSI values.
Polar plot in centre plots change in firing rate in each direction of motion, and rasters plots to side
showing the response to repeated stimuli at each direction. Dorsal, ventral, nasal and temporal
locations indicated by D, V, N and T. **c&d)** Heat maps show spike triggered average of luminance
changes in the RF centre, as a function of time (-80ms – 0), for all neurons in which we were able to
map receptive fields, in *Mus* (c) and *Rhabdomys* (d). **e)** histogram of DSI values for *Rhabdomys* (red)
and *Mus* (cyan) dLGN neurons (Kolmogorov-Smirnov test: P=0.193) **f),** distribution of tuning
preference for DS neurons in *Mus* (top) and *Rhabdomys* (bottom) with DSI > 0.33.
677

Figure 4

Retinal sensitivity to light in *Mus* and *Rhabdomys*

a) Representative traces of dark adapted flash electroretinograms in *Mus* and *Rhabdomys* (left and right, respectively). Flash intensities range from 9.1 to 15.8 log photons/cm^2/s (shown to left of ERG traces). Arrow indicates flash onset. **b&c)** Amplitude (**b**) and implicit time (**c**) of a and b-waves recorded in *Mus* (cyan) and *Rhabdomys* (red). Data shown mean ± SEM, n=5. **d)** Representative ERG traces recorded from *Mus* and *Rhabdomys* in response to 250ms light step (timecourse indicated below; light intensity 14e- log photons/cm^2/s). **e)** Quantification of b-wave (light onset) and d-wave amplitudes (light offset) in response to 250ms light step. **f)** Quantification of b-wave and d-wave implicit time in response to 250ms light step. **g)** Quantification of b-wave and d-wave implicit times in response to 250ms light step. d-wave amplitude expressed as % of b-wave amplitude in *Mus* (cyan) and *Rhabdomys* (red). Data shown mean ± SEM, n=5.

691
Figure 5
A retinal cell atlas for *Rhabdomys*
a) Transcriptionally distinct clusters of *Rhabdomys* retinal cells visualized using UMAP[58]. Cells are
coloured by class identity (RPE, retinal pigment epithelial cells; vEndo, vascular endothelial cells;
Lens Epi, lens epithelial cells; Fibro, fibroblasts). **b)** Dendrogram showing transcriptional relationships
of *Rhabdomys* cell clusters, with major clades corresponding to cell classes. **c)** Bar chart indicating
proportion of photoreceptor types in *Rhabdomys* (red) and *Mus* (cyan).

699
Figure 6
Proportions of bipolar cell types in *Rhabdomys* and *Mus*
a) *Rhabdomys* BCs clustered separately and displayed using UMAP. Cells coloured by cluster identity
b) Confusion matrix indicating the transcriptional correspondence between *Rhabdomys* BC cluster
identity (rows) and classifier-assigned *Mus* BC type identity (columns). Cells are coloured based on
the modified Jaccard Index (colourbar, right), which ranges from 0 (no-correspondence) to 1 (perfect
correspondence; see Methods for details). Pairs of *Rhabdomys* clusters and *Mus* types that belong to
the same BC "orthotype" in[24] are indicated by a star. The preponderance of stars along the diagonal
indicates a high concordance between the correspondence analysis presented here and the orthotype
analysis in [24] **c**) Bar graph showing percentage of BCs assigned rod BC (grey), cone ON BC (red),
cone OFF BC (blue) or BC1B (white) cell types in *Mus* and *Rhabdomys*. Dark horizontal lines within
each subclass demarcate distinct types. Percentages for biological replicates are shown in
Supplementary Table 1. **d**) Transverse sections of *Mus* and *Rhabdomys* stained with antibodies to
PKCalpha. INL, Inner nuclear layer; IPL, inner plexiform layer; GCL, ganglion cell layer. RBC axon
terminals are labelled intensely; arrowhead indicates an RBC cell body. **e**) Scatter plot comparing the
relative frequency of ON (red) and OFF (blue) cone BC types (labelled according to *Mus*) between the
two species. BC1B, which cannot be classified as either ON or OFF, is labelled grey.

717
<u>**Figure 7**</u>
Proportions of RGC types in *Rhabdomys* and *Mus*
a) *Rhabdomys* RGCs were clustered separately and displayed using UMAP. Cells are coloured by
cluster identity. **b**) Confusion matrix indicating the transcriptional correspondence between
Rhabdomys RGC cluster identity (columns) and *Mus* RGC type identity (rows). Cells are coloured
based on the modified Jaccard Index (colourbar, right), as in panel 6b. Stars indicate pairs of
Rhabdomys clusters and *Mus* types that belong to the same orthotype in [24]. This figure shows results
of a classifier trained on *Rhabdomys* data. Correspondence was similar when the classifier was
trained on *Mus* data. **c**) Bar graph showing the relative frequency of each *Mus* RGC type in both the
species. Types are grouped by response polarity – ON, OFF or ON-OFF – based on results in [21,27-29].
Percentages for biological replicates are shown in Supplementary Table 3. **d)** Scatter plot comparing
relative frequency of RGC types between *Rhabdomys* and *Mus*. Response polarity is as shown in **c**.
All 6 types of the W3 subclass[21,31] are more abundant in *Rhabdomys* than *Mus*.

Figure 8

Quantifying ipRGCs in *Rhabdomys* and *Mus* retinae using HCR

a) Scatterplot showing the reduced frequency of putative ipRGC types in *Rhabdomys* (y-axis)
compared to *Mus* (x-axis). **b**) Representative HCR-FISH of *Mus* and *Rhabdomys* retinal wholemounts,
with probe for *Opn4* (species specific). Scale bar = 50μm. **c**) As in b, but showing location of *Opn4*
+ve RGCs across whole retina. Scale bar = 1mm. **d**) Number of *Opn4* +ve cells/mm^2 in *Mus* and
Rhabdomys retinal wholemounts (n=3; lines show mean±SEM). **e**) Histogram showing diameter of
Opn4 +ve cells/mm^2 in *Mus* (cyan) and *Rhabdomys* (red) retinal wholemounts (n=3; data show
mean±SEM). **f**) Scatter plot showing diameter of *Opn4* +ve cells/mm^2 vs normalised signal intensity in
Mus (cyan) and *Rhabdomys* (red) retinal wholemounts (n=3). **g**) Histogram showing normalised
intensity of *Opn4* +ve cells in *Mus* (cyan) and *Rhabdomys* (red) retinal wholemounts (n=3 retinae;
Kolmogorov-Smirnov test: P<0.001)

References

1   Butler, A. B., Hodos, W., ProQuest & ProQuest Ebook, C. *Comparative vertebrate*
neuroanatomy : evolution and adaptation. (2005).

2   Buzsaki, G. *et al.* Tools for probing local circuits: high-density silicon probes combined with
optogenetics. *Neuron* **86**, 92-105, doi:10.1016/j.neuron.2015.01.028 (2015).

3   Jun, J. J. *et al.* Fully integrated silicon probes for high-density recording of neural activity.
Nature **551**, 232-236, doi:10.1038/nature24636 (2017).

4   Roberts, R. J. V., Pop, S. & Prieto-Godino, L. L. Evolution of central neural circuits: state of
the art and perspectives. *Nat Rev Neurosci* **23**, 725-743, doi:10.1038/s41583-022-00644-y
(2022).

5   Tanay, A. & Sebe-Pedros, A. Evolutionary cell type mapping with single-cell genomics.
Trends Genet **37**, 919-932, doi:10.1016/j.tig.2021.04.008 (2021).

6   Zeng, H. & Sanes, J. R. Neuronal cell-type classification: challenges, opportunities and the
path forward. *Nat Rev Neurosci* **18**, 530-546, doi:10.1038/nrn.2017.85 (2017).

7   Chalupa, L. M. *et al. Eye, Retina, and Visual System of the Mouse.* (2008).

8   Dewsbury, D. A. & Dawson, W. W. African 4-Striped Grass Mice (Rhabdomys-Pumilio), a
Diurnal-Crepuscular Muroid Rodent, in the Behavioral Laboratory. *Behav Res Meth Instr* **11**,
329-333, doi:Doi 10.3758/Bf03205671 (1979).

9   Schumann, D. M., Cooper, H. M., Hofmeyr, M. D. & Bennett, N. C. Circadian rhythm of
locomotor activity in the four-striped field mouse, Rhabdomys pumilio: A diurnal African
rodent. *Physiol Behav* **85**, 231-239, doi:10.1016/j.physbeh.2005.03.024 (2005).

10   van der Merwe, I. *et al.* The topography of rods, cones and intrinsically photosensitive retinal
ganglion cells in the retinas of a nocturnal (Micaelamys namaquensis) and a diurnal
(Rhabdomys pumilio) rodent. *PLoS One* **13**, e0202106, doi:10.1371/journal.pone.0202106
(2018).

11   Allen, A. E. *et al.* Spectral sensitivity of cone vision in the diurnal murid Rhabdomys pumilio. *J*
Exp Biol **223**, doi:10.1242/jeb.215368 (2020).

12   Schumann, D. M., Cooper, H. M., Hofmeyr, M. D. & Bennett, N. C. Light-induced Fos
expression in the suprachiasmatic nucleus of the four-striped field mouse, Rhabdomys
pumilio: A southern African diurnal rodent. *Brain Res Bull* **70**, 270-277,
doi:10.1016/j.brainresbull.2006.04.009 (2006).

13   Richardson, R. *et al.* The genomic basis of temporal niche evolution in a diurnal rodent. *Curr*
Biol **33**, 3289-3298 e3286, doi:10.1016/j.cub.2023.06.068 (2023).

14   Morrow, A., Smale, L., Meek, P. D. & Lundrigan, B. Tradeoffs in the sensory brain between
diurnal and nocturnal rodents. *Brain Behav Evol*, doi:10.1159/000538090 (2024).

15   Baden, T. *et al.* The functional diversity of retinal ganglion cells in the mouse. *Nature* **529**,
345-350, doi:10.1038/nature16468 (2016).

16   Grubb, M. S. & Thompson, I. D. Quantitative characterization of visual response properties in
the mouse dorsal lateral geniculate nucleus. *J Neurophysiol* **90**, 3594-3607,
doi:10.1152/jn.00699.2003 (2003).

17   Piscopo, D. M., El-Danaf, R. N., Huberman, A. D. & Niell, C. M. Diverse visual features
encoded in mouse lateral geniculate nucleus. *J Neurosci* **33**, 4642-4656,
doi:10.1523/JNEUROSCI.5187-12.2013 (2013).

18   Enroth-Cugell, C. & Robson, J. G. The contrast sensitivity of retinal ganglion cells of the cat. *J*
Physiol **187**, 517-552, doi:10.1113/jphysiol.1966.sp008107 (1966).

19   Roman Roson, M. *et al.* Mouse dLGN Receives Functional Input from a Diverse Population of
Retinal Ganglion Cells with Limited Convergence. *Neuron* **102**, 462-476 e468,
doi:10.1016/j.neuron.2019.01.040 (2019).

20   Shekhar, K. *et al.* Comprehensive Classification of Retinal Bipolar Neurons by Single-Cell
Transcriptomics. *Cell* **166**, 1308-1323 e1330, doi:10.1016/j.cell.2016.07.054 (2016).

21   Tran, N. M. *et al.* Single-Cell Profiles of Retinal Ganglion Cells Differing in Resilience to Injury
Reveal Neuroprotective Genes. *Neuron* **104**, 1039-1055 e1012,
doi:10.1016/j.neuron.2019.11.006 (2019).

22   Macosko, E. Z. *et al.* Highly Parallel Genome-wide Expression Profiling of Individual Cells
Using Nanoliter Droplets. *Cell* **161**, 1202-1214, doi:10.1016/j.cell.2015.05.002 (2015).

23   Yan, W. *et al.* Mouse Retinal Cell Atlas: Molecular Identification of over Sixty Amacrine Cell
Types. *J Neurosci* **40**, 5177-5195, doi:10.1523/JNEUROSCI.0471-20.2020 (2020).

24   Hahn, J. *et al.* Evolution of neuronal cell classes and types in the vertebrate retina. *bioRxiv*,
doi:10.1101/2023.04.07.536039 (2023).

25 Applebury, M. L. *et al.* The murine cone photoreceptor: a single cone type expresses both S
and M opsins with retinal spatial patterning. *Neuron* **27**, 513-523, doi:10.1016/s0896-
6273(00)00062-3 (2000).
26 Della Santina, L. *et al.* Glutamatergic Monopolar Interneurons Provide a Novel Pathway of
Excitation in the Mouse Retina. *Curr Biol* **26**, 2070-2077, doi:10.1016/j.cub.2016.06.016
(2016).
27 Goetz, J. *et al.* Unified classification of mouse retinal ganglion cells using function,
morphology, and gene expression. *Cell Rep* **40**, 111040, doi:10.1016/j.celrep.2022.111040
(2022).
28 Rousso, D. L. *et al.* Two Pairs of ON and OFF Retinal Ganglion Cells Are Defined by
Intersectional Patterns of Transcription Factor Expression. *Cell Rep* **15**, 1930-1944,
doi:10.1016/j.celrep.2016.04.069 (2016).
29 Huang, W. *et al.* Linking transcriptomes with morphological and functional phenotypes in
mammalian retinal ganglion cells. *Cell Rep* **40**, 111322, doi:10.1016/j.celrep.2022.111322
(2022).
30 Zhang, Y., Kim, I. J., Sanes, J. R. & Meister, M. The most numerous ganglion cell type of the
mouse retina is a selective feature detector. *Proc Natl Acad Sci U S A* **109**, E2391-2398,
doi:10.1073/pnas.1211547109 (2012).
31 Laboulaye, M. A., Duan, X., Qiao, M., Whitney, I. E. & Sanes, J. R. Mapping Transgene
Insertion Sites Reveals Complex Interactions Between Mouse Transgenes and Neighboring
Endogenous Genes. *Front Mol Neurosci* **11**, 385, doi:10.3389/fnmol.2018.00385 (2018).
32 Hattar, S. *et al.* Central projections of melanopsin-expressing retinal ganglion cells in the
mouse. *J Comp Neurol* **497**, 326-349, doi:10.1002/cne.20970 (2006).
33 Ecker, J. L. *et al.* Melanopsin-expressing retinal ganglion-cell photoreceptors: cellular diversity
and role in pattern vision. *Neuron* **67**, 49-60, doi:10.1016/j.neuron.2010.05.023 (2010).
34 Berson, D. M., Castrucci, A. M. & Provencio, I. Morphology and mosaics of melanopsin-
expressing retinal ganglion cell types in mice. *J Comp Neurol* **518**, 2405-2422,
doi:10.1002/cne.22381 (2010).
35 Estevez, M. E. *et al.* Form and function of the M4 cell, an intrinsically photosensitive retinal
ganglion cell type contributing to geniculocortical vision. *J Neurosci* **32**, 13608-13620,
doi:10.1523/JNEUROSCI.1422-12.2012 (2012).
36 Dacey, D. M., Peterson, B. B., Robinson, F. R. & Gamlin, P. D. Fireworks in the primate
retina: in vitro photodynamics reveals diverse LGN-projecting ganglion cell types. *Neuron* **37**,
15-27, doi:10.1016/s0896-6273(02)01143-1 (2003).
37 Van Essen, D. C., Newsome, W. T. & Maunsell, J. H. The visual field representation in striate
cortex of the macaque monkey: asymmetries, anisotropies, and individual variability. *Vision*
Res **24**, 429-448, doi:10.1016/0042-6989(84)90041-5 (1984).
38 Van Essen, D. C., Anderson, C. H. & Felleman, D. J. Information processing in the primate
visual system: an integrated systems perspective. *Science* **255**, 419-423,
doi:10.1126/science.1734518 (1992).
39 Jun, N. Y., Field, G. D. & Pearson, J. M. Efficient coding, channel capacity, and the
emergence of retinal mosaics. *Adv Neural Inf Process Syst* **35**, 32311-32324 (2022).
40 Ocko, S., Lindsey, J., Ganguli, S. & Deny, S. The emergence of multiple retinal cell types
through efficient coding of natural movies. *Advances in Neural Information Processing*
Systems, 9389-9400 (2018).
41 Vielma, A. H. & Schmachtenberg, O. Electrophysiological fingerprints of OFF bipolar cells in
rat retina. *Sci Rep* **6**, 30259, doi:10.1038/srep30259 (2016).
42 Euler, T., Haverkamp, S., Schubert, T. & Baden, T. Retinal bipolar cells: elementary building
blocks of vision. *Nat Rev Neurosci* **15**, 507-519, doi:10.1038/nrn3783 (2014).
43 Mazade, R., Jin, J., Pons, C. & Alonso, J. M. Functional Specialization of ON and OFF
Cortical Pathways for Global-Slow and Local-Fast Vision. *Cell Rep* **27**, 2881-2894 e2885,
doi:10.1016/j.celrep.2019.05.007 (2019).
44 Westo, J. *et al.* Retinal OFF ganglion cells allow detection of quantal shadows at starlight.
Curr Biol **32**, 2848-2857 e2846, doi:10.1016/j.cub.2022.04.092 (2022).
45 Tadmor, Y. & Tolhurst, D. J. Calculating the contrasts that retinal ganglion cells and LGN
neurones encounter in natural scenes. *Vision Res* **40**, 3145-3157, doi:10.1016/s0042-
6989(00)00166-8 (2000).
46 Smeds, L. *et al.* Paradoxical Rules of Spike Train Decoding Revealed at the Sensitivity Limit
of Vision. *Neuron* **104**, 576-587 e511, doi:10.1016/j.neuron.2019.08.005 (2019).

47   Yilmaz, M. & Meister, M. Rapid innate defensive responses of mice to looming visual stimuli.
Curr Biol **23**, 2011-2015, doi:10.1016/j.cub.2013.08.015 (2013).
48   Schmidt, T. M. *et al.* A role for melanopsin in alpha retinal ganglion cells and contrast
detection. *Neuron* **82**, 781-788, doi:10.1016/j.neuron.2014.03.022 (2014).
49   Brown, T. M. *et al.* Melanopsin contributions to irradiance coding in the thalamo-cortical visual
system. *PLoS Biol* **8**, e1000558, doi:10.1371/journal.pbio.1000558 (2010).
50   Allen, A. E., Storchi, R., Martial, F. P., Bedford, R. A. & Lucas, R. J. Melanopsin Contributions
to the Representation of Images in the Early Visual System. *Curr Biol* **27**, 1623-1632 e1624,
doi:10.1016/j.cub.2017.04.046 (2017).
51   Brown, T. M. *et al.* Melanopsin-based brightness discrimination in mice and humans. *Curr Biol*
22, 1134-1141, doi:10.1016/j.cub.2012.04.039 (2012).
52   Pachitariu, M., Steinmetz, N., Kadir, S., Carandini, M. & Harris, K. D. Kilosort:realtime spike-
sorting for extracellular electrophysiology with hundreds of channels. *BioRxiv* (2016).
53   Sjöstrand, K., Clemmensen, L. H., Larsen, R., Einarsson, G. & Ersbøll, B. SpaSM: A MATLAB
Toolbox for Sparse Statistical Modeling. *Journal of Statistical Software* **84**, 1 - 37,
doi:10.18637/jss.v084.i10 (2018).
54   Berens, P. CircStat: A MATLAB Toolbox for Circular Statistics. *Journal of Statistical Software*
31, 1 - 21, doi:10.18637/jss.v031.i10 (2009).
55   Bankhead, P. *et al.* QuPath: Open source software for digital pathology image analysis. *Sci*
Rep **7**, 16878, doi:10.1038/s41598-017-17204-5 (2017).
56   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902 e1821,
doi:10.1016/j.cell.2019.05.031 (2019).
57   Chen, T. & Guestrin, C. XGBoost: A Scalable Tree Boosting System. *Proceedings of the 22nd*
ACM SIGKDD International Conference on Knowledge Discovery and Data Mining 785-794
(2016).
58   McInnes, L., Healy, J. & Melville, J. Umap: Uniform manifold approximation and projection for
dimension reduction. *arXiv preprint* arXiv:1802.03426 (2018).

889