Taking the perspective of an embodied avatar modulates the temporal dynamics of vicarious pain and pleasure: a combined Immersive Virtual Reality and EEG study

Nicolardi V.[1,2], Tieri G.[2,3*], Lisi M.P. [1,2], Mello M.[1,2], Fusaro M.[1,2], Aglioti S.M. [1,2,*]

[1]Department of Psychology, Sapienza University of Rome and CLN[2]S@Sapienza, Istituto Italiano di Tecnologia, 00161, Rome, Italy.

[2] IRCCS Santa Lucia Foundation, 00142, Rome, Italy

[3]Virtual Reality & Digital Neuroscience Lab, Department of Law and Digital Society, University of Rome Unitelma Sapienza, 00161, Rome, Italy.

[*] **Corresponding authors:**

Salvatore Maria Aglioti, E-mail: salvatoremaria.aglioti@uniroma1.it

Gaetano Tieri, E-mail: gaetano.tieri@unitelmasapienza.it

Keywords: Immersive Virtual Reality, Vicarious Pain and Pleasure, EEG, Time and Time-frequency domains.

Abstract

Observing negative and positive valence virtual stimuli can influence the onlookers' subjective and brain reactivity. However, information about the complex link between vicarious pain and pleasure, observer's perspective taking, and cerebral activity is scarce. To address this knowledge gap, we asked twenty-four, VR-immersed healthy participants to report about pleasant, painful, and neutral stimuli delivered to a virtual hand seen from either a first-person perspective (1PP) or third-person perspective (3PP) while undergoing time and time-frequency EEG recording. Participants experienced a stronger sense of ownership over a virtual hand when they viewed it from a 1PP compared to a 3PP. Furthermore, participants rated pain-inducing and pleasure-inducing stimuli as most unpleasant and pleasant, and as more intense. We observed distinct EEG patterns in early (N2, early posterior negativity- EPN) and late (late positive potential-LPP) event-related potentials, as well as in EEG power. The N2 and EPN components showed higher amplitudes for pain and pleasure stimuli compared to neutral stimuli particularly when seen from a 1PP. Conversely, the LPP component exhibited a smaller amplitude for pleasure stimuli compared to both pain and neutral stimuli. We also found that theta-band power increased and alpha power decreased for pain and pleasure stimuli viewed from a 1PP vs a 3PP perspective. Also, in the ultra-late time-window, we observed a decrease in theta, alpha, and beta-band power specifically associated with pleasure stimuli. Our study provides novel evidence on how perspective taking influences the temporal dynamics of vicarious sensations and on distinct electrocortical markers of observed pain and pleasure.

Introduction

Affective and cognitive variables influence the way in which we experience first-hand and vicarious sensations (Lisi et al, 2024). While the vicarious experience of others' feelings and their neural underpinnings has been studied extensively in the domain of pain (Lockwood, 2016; Singer & Lamm, 2009; Betti and Aglioti, 2016) much less attention has been paid to pleasurable touch (Morrison et al., 2011; Lamm et al., 2015; Chiesa et al., 2015, 2017). Interestingly, increasing attention is given to the vicarious experience of pain and pleasure (Pamplona et al., 2022; Mello et al., 2022; Seinfeld et al., 2022) also thanks to the ecological validity of Immersive Virtual Reality (IVR; Tieri et al., 2018). Indeed, IVR typically elicits the sensation that the virtual body observed from 1PP is one's own body, i.e. illusory ownership (Kilteni et al., 2015). Moreover, seeing virtual pain or pleasure stimuli (e.g. a syringe penetrating or a virtual caress) delivered to an embodied virtual hand elicits in the onlooker unpleasant and pleasurable sensations, respectively (Fusaro et al., 2016, 2019, 2021). However, despite the call for research concerning empathy for positive events (Morelli et al., 2015; Mello et al., 2024) research about the brain correlates of vicarious mapping of pleasure is generally limited, and almost absent in IVR. A similar pain-pleasure asymmetry is found also for EEG studies. For example, vicarious experience of pain is reflected in a modulation of components like N1/N2/P3 (Coll, 2018), Early Posterior Negativity (EPN; Fabi and Leuthold, 2017), and Late Positive Potential (LPP; Fan and Han, 2008), as well as modulation of theta and alpha (Mu et al., 2008; Fabi and Leuthold, 2017) and beta responses (Riečanský et al., 2014). Much less is known about whether vicarious social touch modulates EEG activity. Interestingly, Peled-Avron et al. (2016) demonstrate that behavioural and electrocortical reactivity to observed social touch is modulated by levels of empathy. Schirmer & McGlone (2019) found observing affectionate touch elicited larger LPPs (socio-affective marker), but the overlap in brain responses to vicarious pain and pleasure remains unclear. To fill this gap, we combined an adapted version of our previous IVR paradigm (Fusaro et al., 2016, 2019) with time and time-frequency domain EEG recordings in healthy participants who observed virtual painful, pleasurable, or neutral stimuli, delivered to the hand of a virtual avatar seen from a 1PP ("own" hand) or 3PP (other's hand). We focused on cortical responses related to salience, arousal, attentional engagement, and affective salience. In particular, the N2-Visual Evoked Potential, indexed selective attention and stimulus intrinsic relevance (Olofsson, 2008; Luck and Kappenman, 2013; Valentini et al., 2017a); the LPP, reflected the arousal and the motivational salience of the stimulus (Cuthbert et al., 2000; Luck and Kappenman, 2011; Hajcak et al., 2009; Valentini et al., 2015), and the EPN reflected automatic emotion-related influences on information processing (e.g., Olofsson et al., 2008; Schupp et al., 2004). We also focused on: fronto-central theta and centro-lateral alpha power that seem modulated by vicarious experience (Sarlo et al., 2005; González-Franco et al.,

2014; Fabi and Leuthold, 2017); fronto-central theta rhythm (Moreau et al., 2020; Valentini et al., 2017a; Cavanagh et al., 2012) that is enhanced by observation of pain and affective touch compared to control condition (Mu et al., 2008). Finally, we computed central alpha/mu power that appears to decrease when observing painful situations (Mu et al., 2008; Riečanský et al., 2014).

We expected that participants would exhibit greater ownership and valence (pleasant/unpleasant) ratings for pain and pleasure stimuli (compared to neutral) in 1PP vs. 3PP. Furthermore, we expected N2-VEP, LPP and EPN greater amplitude associated to vicarious pain and pleasure, compared to neutral, and in 1PP compared to 3PP. We also expected that stimulus valence and perspective affect fronto-central theta and central alpha power; theta power increases for pain and pleasure compared to neutral and in 1PP vs 3PP; alpha power decreases for pain and pleasure compared to neutral, and for 1PP vs 3PP.

Materials and methods

Participants

Twenty-four healthy volunteers took part in the study (nine males; age mean ± SD, 27.25 ± 3.95). All participants were right-handed with normal visual acuity and were naive as to the purposes of the experiment. The experimental protocol was approved by the ethics committee of the IRCCS Santa Lucia Foundation and was carried out in accordance with the ethical standards of the 2013 Declaration of Helsinki. All participants gave their written informed consent to take part in the study. The number of participants was determined by a priori power analysis performed with MorePower 6.0 (Campbell, 2012) which indicated that 24 participants would be required to detect an effect size of $\eta 2p = 0.19$ with a power of 0.80 and an α level of 0.05. This estimation was based on previous studies by our group using the same experimental paradigm (Fusaro et al., 2016, 2019).

Experimental stimuli and setup

The virtual environment was designed using 3DS Max 2017 (Autodesk, Inc.) and implemented in Unity 2017 game software environment (https://unity.com/). It consisted of a real-size living room equipped with some furniture, lamps and a table where two virtual avatars were sitting facing each other. A virtual gray panel was placed between the avatars in order to avoid their reciprocal face view and to leave visible only the bodies (see Fig 1. Panel A). Avatars were created by using the Morph Character System (MCS) Male avatar asset and were presented in Unity. Participants observed the virtual scenario and the virtual upper limb from a first-person perspective (1PP) by means of an Oculus Rift head-mounted display (HMD; www.meta.com), having 110° field of view (diagonal FOV) with a resolution of 2.160 × 1.200. (see Fig 1.A Right Panel). In order to reach the best spatial

match between real and virtual environment, the virtual table was designed with the same height of the real one and virtual body observed in 1PP was designed and programmed with same posture and position of the participant's real body. This method, widely used in our previous studies (Tieri et al., 2015a, 2015b, 2017; Fusaro et al., 2016, 2019. 2021; Monti et al., 2020; Fusco et al., 2020), allowed us to implement the basic condition (based on visuo-proprioceptive congruence between the hidden real and observed virtual body) to induce illusory ownership over the virtual body. Experimental stimuli consisted of about three different animations implemented in the virtual environment (all stimuli were matched in order to have same animation length): i.e. (1) a virtual arm holding a needle on the avatar's right hand and penetrating it, (2) a virtual hand approaching the avatar's right hand and caressing it and (3) a virtual green ball approaching and touching the avatar's right hand (Fig 1, Panel B). The virtual caress was realized in Unity by readapting the real actor's kinematics used in our previous study (Fusaro et al., 2019). Each stimulus was delivered on the virtual hand observed from 1PP (and thus perceived as 'own') or observed from 3PP (and thus perceived as 'other'). Subjective ratings, concerning the valence of the stimuli ((Un)pleasantness ratings), their intensity and the illusory ownership over the observed virtual hand were collected by means of a PC keyboard placed in front of the participant (not displayed in the virtual environment) and interfaced with Unity with a customized C# script (see paragraph *Subjective ratings* for more details). A Neuroscan SynAmps RT amplifiers system (Compumedics) with a 64-channel EEG cap (Electro-Cap International) has been used to record EEG signals (see paragraph *EEG recording* for more details). The EEG system was interfaced with VR by means of a TriggerStation TS832U (Braintrends ltd, www.braintrends.it).

-------------------------------- Please add Fig 1 near here --------------------------------

General procedure

Participants were seated comfortably on a chair in front of a table and wore a 64-channel EEG cap and the HMD (through which they passively observed the virtual body and the scenario from 1PP). Before the experiment, participants underwent the Calibration and Familiarization phases. In the Calibration phase, the point-of-view of each participant were adjusted to pair the 1PP avatar with individual positioning in order to obtain the best spatial-match between the participant's real and virtual body. In the Familiarization phase, participants (1) were invited to look both at their virtual body and at the environment and to verbally describe what they were seeing (60 sec; Tieri et al., 2015b; Slater et al., 2010) and (2) familiarized with each stimulus (pain, neutral and pleasure in both 1PP and 3PP) and requested to rate the stimuli-related sensations on the Visual Analogue Scale

(VAS). The experiment was composed of two separate blocks (one for each perspective 1PP-3PP) presented in a counterbalanced order across subjects. Each block consisted of 90 trials including each stimulus valence (30 Pain, 30 Pleasure and 30 Neutral presented in pseudo-randomized order). Thus, participants performed a total of 180 trials (90 1PP, 90 3PP). Each trial started with the observation of a right virtual hand (in 1PP or 3PP). After 5500 ± 500 msec, the stimulus was delivered to the virtual right hand on a region overlying the first dorsal interosseous. Each stimulus lasted ~2.500 msec. Upon its disappearance, participants continued to observe the virtual hand until a sequence of two VAS was presented. Participants were requested to provide ratings about stimulus-related sensations concerning the perceived valence and intensity (see Fig. 2 for the sequence of events per trial). Moreover, three additional VAS were presented in order to allow participants to rate the illusory feeling of ownership over the observed virtual hand (see Table 1, Items 3-5). In particular, in six trials of each block, the Ownership ratings' VASs appeared in counterbalanced order across trials. Thus, while VAS items 1-2 (Table 1) appeared after each stimulus, VAS items 3-5 were presented in counterbalanced order, 6 times per block, i.e. after 15 trials (two times for pain, pleasure and neutral stimuli in each block).

-------------------------------- Please add Fig 2 near here --------------------------------

Subjective Ratings

After each stimulus, a black panel with a horizontal green line appeared in the virtual space in front of the participants (see Fig. 2). The horizontal green line was 60 cm long with left and right extremities marked as "−−" and "++", respectively. Participants used their left index and middle finger to control a green cursor along the horizontal line to provide the evaluations listed in Table 1 (items selected and readapted from previous studies; Fusaro et al., 2016, 2019). In particular, item 1 concerned the valence ("−−" = strongly unpleasant, middle=neutral, "++" strongly pleasant) and item 2 the intensity ("−−" = very weak, "++" = very strong;) of the observed stimuli (Bufalari et al., 2007). Item 3-5 concerned the illusory feeling of body ownership over the virtual hand ("−−"=no illusion; "++"= maximal illusion).

-------------------------------- Please add Tab 1 near here --------------------------------

EEG recording and processing

EEG signals were recorded and amplified using a Neuroscan SynAmps RT amplifiers system (Compumedics Limited, Melbourne, Australia). These signals were acquired from 60 tin scalp electrodes embedded in a fabric cap (Electro-Cap International, Eaton, OH), arranged according to

the 10-10 system. The EEG was recorded from the following channels: Fp1, Fpz, Fp2, AF3, AF4, F7, F5, F3, F1, Fz, F2, F4, F6, F8, FC5, FC3, FC1, FCz, FC2, FC4, FC6, T7, C5, C3, C1, Cz, C2, C4, C6, T8, TP7, CP5, CP3, CP1, CPz, CP2, CP4, CP6, TP8, P7, P5, P3, P1, Pz, P2, P4, P6, P8, PO7, PO3, AF7, POz, AF8, PO4, PO8, O1, Oz, O2, FT7, and FT8. All electrodes were physically referenced to an electrode placed on the right earlobe and were re-referenced off-line to the average of all electrodes. Impedance was kept below 5 KΩ for all electrodes for the whole duration of the experiment, amplifier hardware band-pass filter was 0.01 to 200 Hz and sampling rate was 1000 Hz. The continuous EEG data were pre-processed with EEGLAB (Delorme and Makeig, 2004) and Letswave 6 (www.letswave.org). Data were band-pass filtered from 0.1 to 30 Hz (filter order 4) and re-sampled at 250 Hz. Bad channels have been interpolated when needed, based on the signal of the three closest electrodes. Data were segmented into epochs lasting from 1 s before to 1.5 s after the stimulus onset. Eye movements or eye blinks were removed using an Independent Component Analysis (ICA; Jung et al., 2000). The resultant signal was then baseline corrected to a reference interval from -1 to -0.5 s before the stimulus onset and re-referenced to the average reference.

EEG analyses in the time-domain

In the time domain, middle and late latency visual evoked potentials (VEPs) and Early posterior negativity (EPN) were analyzed separately. Specifically, we focused on: the N2-P2 waveforms, occurring between 200-300 ms which have been proposed to reflect selective attention processes in a complex visual scene (Olofsson et al., 2008); the Late Positive Potential (LPP), a positive deflection occurring after 300 ms (Hajcak et al., 2009), the magnitude of which has been associated with motivational salience, subjective arousal and more in general, processing of emotional contents (Cuthbert et al., 2000; Luck and Kappenman 2012, Valentini et al., 2017a) and the EPN, a negative deflection occurring between 200-300 ms (Farkas et al., 2020), that has been proposed to reflect "natural selective attention", and to be more sensitive to affective arousing contents (Schupp et al., 2004; Olofsson et al., 2008). VEPs amplitude was measured at occipital ROI (POz-Oz) (Olofsson et al., 2008; Luck and Kappenman, 2012) while EPN amplitude was measured at left parietal ROI contralaterally to stimulus presentation side(P5-P7-PO7) (Frank & Sabatinelli, 2019; Farkas et al., 2020). In order to investigate any differences between amplitudes across the experimental conditions, single subject average ERPs amplitude for each condition was entered in whole-waveform 2 (Perspective, 1PP vs 3PP) X 3 (stimulus valence, neutral vs pain vs pleasure) ANOVAs. Post-hoc comparisons were performed using the Letswave 6 software, t-tests with correction for multiple comparisons (Maris and Oostenveld, 2007). This method of correction, based on a cluster-level randomisation, allowed us to control the Type-1 error rate involving multiple comparisons. It consists

of the following steps: firstly, selecting the points represented by F/t values lower than P < 0.05 to identify the groups (clusters) of contiguous points that show a significant effect. Then, estimating the magnitude of each cluster by calculating the sum of the F/t values comprising each cluster. Afterwards, random permutation (500 times) of the differences between data-points/pixels at the cluster level within each individual was used to obtain a reference distribution. This distribution was obtained by randomly swapping the conditions within participants and calculating the maximum cluster-level test statistic. For each cluster, a threshold of significance was found around the value Z > 2 standard deviations from the mean. Then, for each cluster, a value corresponding to F/t and P (two-tailed) was obtained and the statistical significance ascribed only to differences lower than 0.05 at the end of the permutation process.

EEG analysis in the time-frequency domain

A short-term Fast-Fourier transform, as implemented in Letswave 6, has been computed to measure responses power in the time-frequency domain. The following parameters have been used: frequency between 1-30 Hz, step of 25 for 30 lines, to compute power for each trial, then averaged. The resultant signals, were displayed as an event-related change of average power (ER%), relative to a reference interval prior to the onset of the stimulus (-1 to −0.5 s). We focused on the theta and mu power measured at a central ROI (FCz, Cz, C3) that are consided possible indices in perspective taking, conflict monitoring and action mirroring (Pavone et al, 2016; Petereit et al, 2023; Mu et al., 2008; Riečanský et al., 2014).

In order to investigate any differences in time-frequency domain between experimental conditions (Neutral-1PP, Neutral-3PP; Pleasure-1PP, Pleasure 3PP; Pain-1PP, Pain-3PP), single subject average power of each experimental condition underwent ANOVA and t-test in Letswave 6, with correction for multiple comparisons (Maris and Oostenveld, 2007, see previous paragraph for details).

Results

Subjective Ratings

Valence ratings (Table 1, Item 1) were not normally distributed (Shapiro–Wilk test for one out of six conditions p< 0.05). Thus, nonparametric analysis including Friedman ANOVA and Wilcoxon test for the within-group effects of valence (pain, neutral, pleasure) and perspective (1PP, 3PP) was used. Since Friedman's ANOVA only allows a One-Way within-subject analysis, we collapsed perspective and valence into a unique 6-level Condition within-subject factor, to test if there is a general effect of Condition. The Friedman ANOVA revealed a significant effect of Condition ($\chi 2(5) = 89.31$, p < 0.0001; Fig. 3A). Bonferroni-corrected Wilcoxon tests for nine planned comparisons (α level =

0.0055) showed that pain (mean ± SD: 27.67 ± 17.24 for 1PP; 26.17 ± 16.25 for 3PP) elicited stronger unpleasantness than neutral (50.75 ± 6.26 for 1PP; 50.75 ± 6.05 for 3PP) and pleasure (66.97 ± 14.92 for 1PP; 64.18 ± 16.50 for 3PP; note that lower and higher ratings represent unpleasant and pleasant sensations, respectively); moreover, pleasure elicited higher pleasantness than neutral and pain (see Table 2-3 for all comparisons). No effect of perspective was found (all ps > 0.05). These results suggest that pain and pleasure stimuli elicited unpleasant and pleasant vicarious sensations, regardless of the perspective from which the stimuli were observed.

Intensity ratings (Table 1, Item 2) were not normally distributed (Shapiro–Wilk test for two out of six conditions p < 0.05). We conducted a Friedman ANOVA with the perspective and valence factors collapsed into a unique 6-level Condition within-subject factor. The Friedman ANOVA revealed a significant effect of Condition ($\chi2(5) = 45.27$, p < 0.0001; Fig. 3B). Bonferroni-corrected Wilcoxon tests for nine planned comparisons (α level = 0.0055) showed significant differences between pain (mean ± SD: 52.92 ± 25.62 for 1PP; 48.06 ± 25.14 for 3PP) vs neutral in 1PP (35.26 ± 21.15 for 1PP; 33.65 ± 21.71 for 3PP) and pleasure (55.97 ± 23.25 for 1PP; 49.49 ± 24.35 for 3PP) vs neutral in both perspectives (see Table 2-3), but not pain vs pleasure (ps > 0.05). No effect of perspective was found (all ps > 0.05). These results suggest that participants perceived as more intense pain than neutral stimuli in 1PP, and pleasure as more intense than the neutral stimuli in both perspectives.

------------------------------ Please insert Table 2 near here ------------------------------
------------------------------ Please insert Table 3 near here ------------------------------

Ownership mean scores for experimental (Table 1, Items 3-4) and the control items (Table 1, Item 5) were not normally distributed (Shapiro–Wilk test for six out of twelve conditions p < 0.05). Thus, we conducted a Friedman ANOVA with the item (experimental, control), perspective and valence factors collapsed into a unique 12-level Condition within-subject factor. The Friedman ANOVA revealed a significant effect of Condition ($\chi2(11) = 104.38$, p < 0.0001; Fig. 3C). Firstly, the mean scores of experimental items and the control one were compared for each condition (1PP-Pain, 1PP-Pleasure, 1PP-Neutral, 3PP-Pain, 3PP-Pleasure, 3PP-Neutral) using Bonferroni correction for six planned comparisons (α level = 0.0083). Wilcoxon tests showed that the scores on the experimental items were higher than the control in 1PP (ps < 0.001), but not in 3PP (ps > 0.05; see Table 2-3 for descriptive statistics and planned comparisons). Secondly, comparisons showed that this effect was not modulated by valence (α level= 0.0055), as no significant differences were found on the experimental items between 1PP pain (69.57 ± 17.44), 1PP neutral (64.10 ± 22.25) and 1PP pleasure

(66.78 ± 17.79; all ps > 0.05, see Table 2-3). These results suggest that illusory ownership was experienced only when the virtual hand was observed from 1PP and regardless of the stimulus valence.

------------------------------ Please insert Fig 3 near here ------------------------------

EEG in the time domain: ERPs results

The ANOVA on VEPs revealed a main effect of valence (pain, neutral, pleasure; F=32.83, p<0.001) between 160-400 ms, and perspective (1PP, 3PP; F=22.17, p<0.001) between 190-310 ms, on the N2-P2 complex (see Fig. 4). In line with our hypothesis, post-hoc comparisons showed greater amplitude for pain and pleasure as compared to neutral (pain vs neutral: t=-6.53, p<0.001; pleasure vs neutral: t=-6.55, p<0.001; Fig.4 panel A) as well as for 1PP compared to 3PP (Fig. 4 panel B), for the N2-P2 waveforms. Analysis of LPP showed a main effect of valence (F=13.59, p<0.001) between 400-600 ms. Surprisingly, this effect was accounted for by greater amplitude for neutral and pain compared to pleasure (pain vs pleasure: t=-4.66, p<0.001; pleasure vs neutral: t=-4.40, p<0.001; Fig.4 panel A). Figure 4 shows the effect of valence (panel A), and of perspective (panel B); significant post-hoc comparisons are highlighted in the yellow boxes of each panel.

------------------------------ Please add Fig 4 near here ------------------------------

The ANOVA on EPN revealed a main effect of valence (F=32.83, p<0.001) between 200-535 ms. Post-hoc comparisons indicate greater amplitude for pain and pleasure compared to neutral (pain vs neutral: t=-4.98, p<0.001; pleasure vs neutral: t=-6.81, p=<0.001; fig.5, panel A). The main effect of perspective was also significant (F=32.83, p<0.001, fig. 5, panel B) between 260-485ms, an effect that was accounted for by greater amplitude for 1PP compared to 3PP. No significant interaction was found.

------------------------------ Please add Fig 5 near here------------------------------

EEG in the Time-frequency domain: power changes in different frequency bands

ANOVA on EEG power revealed a main effect of valence (F=2.9, p<0.001) and perspective (F=3.86, p<0.001) (see Fig. 6), and a significant interaction between the two (see Fig. 7) (F=4.83, p=0.008 for the theta range; F=10.63, p<0.001 for the alpha range). Concerning the main effect of valence (Fig.

6, upper row), specific comparisons between the levels of valence (neutral, pain and pleasure), did not show significant differences after correcting p-value for multiple comparisons (500 permutations, see methods). The main effect of perspective (Fig. 6, bottom row), was accounted for by increased theta power in 1PP compared to 3PP in the 300-500ms window, and decreased alpha power in 1PP compared to 3PP in the 500-1000 ms window. The significant interaction showed increased theta power in the 200-350ms window, for pain (t=6.08, p<0.001) and pleasure (t=5.29, p<0.001) compared to neutral, in 1PP (Fig. 7, panel A) but not in 3PP (Fig. 7, panel B). Interestingly, in the ultra late time-window (after 800ms), a specific effect of pleasure was found, indicating a power decrease for pleasure compared to neutral. In 1PP, a beta power decrease was found for pleasure compared to neutral (t=-3.01, p=0.006; Fig. 7, panel A). Moreover, in 3PP, a power decrease for pleasure compared to neutral was found in the theta (t=-3.83, p<0.001), alpha (t=-3.56, p=0.003) and beta (t=-3.72, p=0.001) band (Fig. 7, panel B, second row) in the ultra late time-window.

-------------------------------- Please add Fig 6 and Fig 7 near here --------------------------------

Correlations between Subjective Ratings and EEG in the time-frequency domain

We used Pearson R correlation to test any possible association between the EEG power and the unpleasantness and intensity ratings. In specific, the analysis concerned subjective indices (computed subtracting the Neutral condition from either Pleasure or Pain) and time-frequency values that were significantly modulated in the different stimulus observation conditions. Ultralate Beta (1PP) and Theta (3PP) for the contrast between Pleasure and Neutral showed a moderate association with the Intensity ratings (r=0.38 and r=0.35, respectively). However, no association turned out to be statistically significant. All the ratings and across conditions comparisons are reported in Table 4.

-------------------------------- Please insert Table 4 near here --------------------------------

Discussion

In the present study we explored, for the first time using a within-subject design, the behavioural and brain reactivity to the observation of pain and pleasure stimuli delivered to an embodied vs. a non-embodied virtual avatar. To achieve our aim, we combined immersive virtual reality with time- and time-frequency domain EEG recording in healthy participants observing from a 1PP or 3PP: i) a virtual arm holding a needle that approached and then penetrated the right hand of an avatar (pain condition); ii) a virtual hand approaching the avatar's right hand and caressing it (pleasure) and; iii)

a virtual green ball approaching and touching the avatar's right hand (neutral). Subjective and neurophysiological data indicate that cues to virtual embodiment like body ownership, may operate at both explicit (Fusco et al., 2020; Monti et al., 2020; Tieri et al., 2015a) and implicit levels (Tieri et al., 2015b, 2017; Fusaro et al., 2016, 2021; Pavone et al., 2016; Casula et al., 2022; Pezzetta et al., 2023). Expanding on previous evidence (Tieri et al., 2015a; Pavone et al. 2016; Fusaro et al. 2016, 2019), subjective reports indicate that our experimental approach is adept to induce vicarious sensations and embodiment over artificial agents. In specific, participants reported stronger illusory feelings of ownership over the virtual hand when observing it from a 1PP compared to a 3PP. Subjective ratings also indicate that our painful and pleasant virtual stimuli were effective in eliciting most unpleasant and most pleasant sensations, respectively, and were rated as more intense compared to neutral stimuli. These data confirm that IVR paradigms are effective in reproducing veridical real-life like vicarious experiences. Tellingly, additional novel results of the present study concern the effect of stimulus valence (painful, pleasurable, neutral) and observational perspective (1PP vs 3PP) on time- and time-frequency domain EEG findings.

EEG results in the time domain.
Valence and perspective differentially modulated early and late stimulus-evoked EEG components. In specific, early N2-VEPs and late EPN showed greater amplitude for pain and pleasure (vs. neutral), and for 1PP (vs 3PP). This pattern of results likely reflects attentional engagement, that turns out to be higher for emotionally valued stimuli compared with the neutral ones, and for 1PP compared with the 3PP. N2-VEPs components are thought to reflect processes involved in selective attention (Olofsson et al., 2008; Luck and Kappenman 2011). Here, the early component did not distinguish between positive or negative value of the stimuli, but coded its salience, either attentional or emotional (Todd et al., 2012). A greater EPN amplitude was found for 1PP compared to 3PP, and for Pain and Pleasure compared to Neutral conditions suggesting that this EEG component is linked to emotionally salient scenarios with no differences between positive and negative valence. Indeed, the EPN reflects automatic emotion-related influences on information (e.g., Olofsson et al., 2008; Schupp et al., 2004), processing of body parts (Hietanen et al., 2014), and processing of salient affective stimuli, with no category-specific sensitivity (Fabi & Leuthold, 2017).
Interestingly, vicarious pleasant touch elicited smaller LPP (400-600 ms) amplitude than painful and neutral stimuli. LPP has been proposed to reflect the arousal and the motivational salience of the stimulus (Cuthbert et al., 2000; Luck and Kappenman 2012; Hajcak et al., 2009), with greater amplitude for more relevant and arousing stimuli. However, studies indicate that central-LPP can be modulated by the pleasantness of the received touch (Haggarty et al., 2020), with smaller LPP for

more pleasant touch, in those people who preferred slow over fast touch (Schirmer et al., 2023). Thus, our result that LPP specifically responded to the pleasant scenario is in line with previous studies concerning real touch. Given the adopted paradigm, we cannot disambiguate if our LPP modulation was due to the pleasantness of the vicarious touch, or to its positive valence with respect to the other scenarios. Yet, our results expand current knowledge by showing that the effect of virtual pleasant touch is similar to what reported for real stimuli.

EEG results in the time-frequency domain.

EEG power in theta and alpha band was modulated by stimulus valence and perspective between 200-500 ms. Within this time-interval, for the theta band, the effect of valence led to an increase in power for pain and pleasure compared to neutral, while an increased power was found for 1PP scenarios compared to 3PP. The significant interaction scenarios X perspective interaction is explained by the modulation of theta power for pain and pleasure only in 1PP. The modulation of the theta-band frequency activity observed for affective stimuli is thought to be related to the emotional processing and affective valence discrimination, with enhanced theta power for affective stimuli, particularly in central and frontal areas (Aftanas et al., 2001; Mu et al., 2008; Schirmer et al., 2018). However, theta power increase has also been associated with cognitive effort (Gregory et al., 2022), sensorimotor integration (Kober & Neuper, 2011) and conflict/error-processing (Cavanagh et al., 2012; Pezzetta et al., 2018, 2023). Thus, the enhanced theta power for pain and pleasure vs neutral scenarios found in our study may be explained by the high cognitive load related to affective content. Moreover, finding enhanced theta-power only in 1PP suggests that different mechanisms may underpin vicarious experiences depending on the adopted perspective. Alpha band power was modulated by perspective only, showing power decrement in 1PP compared to 3PP, with no differences for the stimulus valence, nor an interaction between valence and perspective. Alpha band responses to emotionally arousing stimuli led to conflicting results, with pleasant and unpleasant stimuli associated with both alpha power increase or decrease respectively (Schubring & Schupp, 2019; Uusberg et al., 2013). However, less is known about the self-other distinction when stimuli are presented in 'own' vs. 'other' perspective. Alpha bands responses have been related to the allocation of visuospatial attention (Mathewson et al., 2011), the activation of sensory (Van Ede et al., 2011) and/or motor cortex (Perry et al., 2011), with alpha/mu suppression linked to social information processing (Pineda & Hecht, 2009) and perception of others' pain (Cheng et al., 2008). Moreover, alpha suppression has been related to participants' engagement in a task, with larger mu suppression when participants are engaged in a social game compared to passively looking at the other's action (Oberman et al., 2007; Perry et al., 2011). In this vein, greater alpha suppression is associated with higher entrainment in

time during a musical joint action paradigm (Novembre et al., 2016). Here, we found that the cognitive perspective rather than the stimulus valence, modulated alpha power in response to virtual scenarios, likely indicating different attentional engagement elicited by 1PP or 3PP respectively.

Finally, the significant interaction between valence and perspective also showed a power decrease after 800 ms, in the theta, alpha and beta band, an effect that was specific for pleasure scenarios. Interestingly, while theta power increase over midfrontal areas has been associated with conflict/error processing and valence discrimination (Aftanas et al., 2001; Cavanagh et al., 2012), theta power decrease in virtual reality has been associated with calmness (Eswaran et al., 2018). Beta desynchronization has been associated with motor readiness and sensorimotor activation (Riečanský et al., 2014; Fabi and Leuthold, 2017), the observation of a moving hand (Pfurtscheller et al., 2007), as well as cognitive aspects of pain processing (Valentini et al., 2017b). Moreover, while the power decrease for theta and alpha concerns only the 3PP, the power decrease for beta band is found for both perspectives. Overall, this power decrease seems specifically associated to pleasant stimuli and occurs in the ultra late time-window. This effect may be associated to the attentional engagement and sensorimotor integration contingent upon vicarious pleasant touch processing. Thus, although somewhat speculatively, we submit that, our decreased beta power may reflect a reduction of alert or negative state. A similar meaning may be attributed to the the late time-window LPP amplitude reduction we observed for pleasure compared with neutral scenarios. Our findings of late and ultralate time- and time-frequency EEG signs of vicarious processing, is in keeping with a recent multivariate pattern analysis whole-brain EEG study showing that seeing touch evokes overlapping representations at later stages of information processing (Smit et al., 2023).

All in all, our study indicates that stimulus valence and perspective differentially affect ERPs related to attentional and motivational/affective processing, as well as EEG responses in theta, alpha and beta band. In particular, the affective scenarios (pain and pleasure) elicited similar EEG activity related to attentional engagement (N2, alpha and theta before 500 ms), while exhibiting distinct activity patterns at late and ultra-late time-window. Although these novel effects need further characterized, our patterns of results is compatible with the idea that vicarious pleasant experience, and its calming/consoling effect (Peled-Avron et al., 2018; Morrison, 2016) could be related to a specific cortical reactivity pattern, occurring after 500 ms in the neural processing. These results have potential implications for clinical research, for example on patients with abnormal perception or emotional reactivity to painful or pleasant stimuli (Boheme et al., 2020; Habig et al., 2021; Fusaro et al., 2023; Nicolardi et al., 2023).

Acknowledgements

This work was supported by BIAL Foundation (n° 218/2016) and by Italian Ministry of Health (Ricerca Finalizzata, Giovani Ricercatori 2019, n°GR-2019-12369761). S.M.A. gratefully acknowledges the support of the Institut d'études avancées de Paris.

Conflict of interest

The authors declare no competing financial interests.

Data availability statement

Raw data and scripts for the analyses will be available on a public data repository at the moment of publication of the manuscript in a peer-reviewed journal.

Author contributions

G.T., M.F., V.N. and S.M.A. designed research; G.T. created the 3D models, software and implemented the VR setup; V.N., M.P.L. and M.M. collected the data; V.N. analyzed EEG data; G.T. and M.P.L analyzed subjective data; V.N. provided the first draft of the manuscript. S.M.A supervised the research. All authors approved the final version of the manuscript.

References

Aftanas, L., Varlamov, A., Pavlov, S., Makhnev, V., & Reva, N. (2001). Event-related synchronization and desynchronization during affective processing: emergence of valence-related time-dependent hemispheric asymmetries in theta and upper alpha band. International journal of Neuroscience, 110(3-4), 197-219.

Betti, V., & Aglioti, S. M. (2016). Dynamic construction of the neural networks underpinning empathy for pain. Neuroscience & Biobehavioral Reviews, 63, 191-206.

Bucchioni, G., Fossataro, C., Cavallo, A., Mouras, H., Neppi-Modona, M., & Garbarini, F. (2016). Empathy or ownership? Evidence from corticospinal excitability modulation during pain observation. Journal of Cognitive Neuroscience, 28(11), 1760-1771.

Bufalari, I., Aprile, T., Avenanti, A., Di Russo, F., & Aglioti, S. M. (2007). Empathy for pain and touch in the human somatosensory cortex. Cerebral Cortex, 17(11), 2553–2561. https://doi.org/10.1093/cercor/bhl161

Campbell, J. I. D. & Thompson, V. A. MorePower 6.0 for ANOVA with relational confidence intervals and Bayesian analysis. *Behav.Res. Methods* 44, 1255–1265 (2012).

Casula, E. P., Tieri, G., Rocchi, L., Pezzetta, R., Maiella, M., Pavone, E. F., … Koch, G. (2022). Feeling of Ownership over an Embodied Avatar's Hand Brings About Fast Changes of Fronto-Parietal Cortical Dynamics. *The Journal of Neuroscience*, *42*(4), 692–701. https://doi.org/10.1523/JNEUROSCI.0636-21.2021

Cavanagh, J. F., Figueroa, C. M., Cohen, M. X., & Frank, M. J. (2012). Frontal theta reflects uncertainty and unexpectedness during exploration and exploitation. Cerebral cortex, 22(11), 2575-2586.

Cavanagh, J. F., Zambrano-Vazquez, L., & Allen, J. J. (2012). Theta lingua franca: A common mid-frontal substrate for action monitoring processes. Psychophysiology, 49(2), 220-238.

Cheng, Y., Yang, C. Y., Lin, C. P., Lee, P. L., & Decety, J. (2008). The perception of pain in others suppresses somatosensory oscillations: a magnetoencephalography study. Neuroimage, 40(4), 1833-1840.

Chiesa, P. A., Liuzza, M. T., Acciarino, A., & Aglioti, S. M. (2015). Subliminal perception of others' physical pain and pleasure. Experimental Brain Research, 233, 2373-2382.

Chiesa, P. A., Liuzza, M. T., Macaluso, E., & Aglioti, S. M. (2017). Brain activity induced by implicit processing of others' pain and pleasure. Human Brain Mapping, 38(11), 5562-5576.

Coll, M. P. (2018). Meta-analysis of ERP investigations of pain empathy underlines methodological issues in ERP research. Social cognitive and affective neuroscience, 13(10), 1003-1017.

Coll, M. P., Bird, G., Catmur, C., & Press, C. (2015). Cross-modal repetition effects in the mu rhythm indicate tactile mirroring during action observation. Cortex, 63, 121-131.

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. Journal of neuroscience methods, 134(1), 9-21.

ESwaran, V. S. B., VEEzhinathan, M., Balasubramanian, G., & Taneja, A. (2018). Virtual Reality Therapy for Mental Stress Reduction. Journal of Clinical & Diagnostic Research, 12(10).

Fabi, S., & Leuthold, H. (2017). Empathy for pain influences perceptual and motor processing: Evidence from response force, ERPs, and EEG oscillations. Social neuroscience, 12(6), 701-716.

Fan, Y., & Han, S. (2008). Temporal dynamic of neural mechanisms involved in empathy for pain: an event-related brain potential study. Neuropsychologia, 46(1), 160-173.

Farkas, A. H., Oliver, K. I., & Sabatinelli, D. (2020). Emotional and feature-based modulation of the early posterior negativity. Psychophysiology, 57(2), e13484.

Frank, D. W., & Sabatinelli, D. (2019). Hemodynamic and electrocortical reactivity to specific scene contents in emotional perception. Psychophysiology, 56(6), e13340. https://doi.org/10.1111/psyp.13340

Fusaro, M., Bufacchi, R. J., Nicolardi, V., & Provenzano, L. (2022). The analgesic power of pleasant touch in individuals with chronic pain: Recent findings and new insights. Frontiers in Integrative Neuroscience, 16, 956510.

Fusaro, M., Lisi, M. P., Tieri, G., & Aglioti, S. M. (2021). Heterosexual, gay, and lesbian people's reactivity to virtual caresses on their embodied avatars' taboo zones. *Scientific Reports*, *11*(1). https://doi.org/10.1038/s41598-021-81168-w

Fusaro, M., Tieri, G., & Aglioti, S. M. (2016). Seeing pain and pleasure on self and others: behavioral and psychophysiological reactivity in immersive virtual reality. *Journal of Neurophysiology*, *116*(6), 2656–2662. https://doi.org/10.1152/jn.00489.2016

Fusaro, M., Tieri, G., & Aglioti, S. M. (2019). Influence of cognitive stance and physical perspective on subjective and autonomic reactivity to observed pain and pleasure: An immersive virtual reality study. *Consciousness and Cognition*, *67*(November 2018), 86–97. https://doi.org/10.1016/j.concog.2018.11.010

Fusco, G., Tieri, G., & Aglioti, S. M. (2020). Visual feedback from a virtual body modulates motor illusion induced by tendon vibration. *Psychological Research*, (0123456789). https://doi.org/10.1007/s00426-020-01366-5

Gaetz, W., & Cheyne, D. (2006). Localization of sensorimotor cortical rhythms induced by tactile stimulation using spatially filtered MEG. Neuroimage, 30(3), 899-908.

González-Franco, M., Peck, T. C., Rodríguez-Fornells, A., & Slater, M. (2014). A threat to a virtual hand elicits motor cortex activation. Experimental brain research, 232(3), 875-887.

Haggarty, C. J., Malinowski, P., McGlone, F. P., & Walker, S. C. (2020). Autistic traits modulate cortical responses to affective but not discriminative touch. European Journal of Neuroscience, 51(8), 1844-1855.

Hajcak, G., Dunning, J. P., & Foti, D. (2009). Motivated and controlled attention to emotion: time-course of the late positive potential. Clinical neurophysiology, 120(3), 505-510.

Hietanen, J. K., Kirjavainen, I., & Nummenmaa, L. (2014). Additive effects of affective arousal and top-down attention on the event-related brain responses to human bodies. Biological psychology, 103, 167-175.

Ionta, S., Costantini, M., Ferretti, A., Galati, G., Romani, G. L., & Aglioti, S. M. (2020). Visual similarity and psychological closeness are neurally dissociable in the brain response to vicarious pain. cortex, 133, 295-308.

Jung, T. P., Makeig, S., Humphries, C., Lee, T. W., Mckeown, M. J., Iragui, V., & Sejnowski, T. J. (2000). Removing electroencephalographic artifacts by blind source separation. Psychophysiology, 37(2), 163-178.

Kilteni, K., Maselli, A., Kording, P. K., & Slater, M. (2015). Over my fake body: body ownership illusions for studying the multisensory basis of own-body perception. *Frontiers in Human Neuroscience*, *9*(March). https://doi.org/10.3389/fnhum.2015.00141

Kober, S. E., & Neuper, C. (2011). Sex differences in human EEG theta oscillations during spatial navigation in virtual reality. International Journal of Psychophysiology, 79(3), 347-355.

Lamm, C., Silani, G., & Singer, T. (2015). Distinct neural networks underlying empathy for pleasant and unpleasant touch. Cortex, 70, 79-89.

Lisi, M. P, Fusaro, M., & Aglioti, S. M. (2024). Visual perspective and body ownership modulate vicarious somatosensory processing: A systematic review. *Psychon Bull Rev* (2024). https://doi.org/10.3758/s13423-024-02477-5

Lockwood, P. L. (2016). The anatomy of empathy: Vicarious experience and disorders of social cognition. Behavioural brain research, 311, 255-266.

Luck, S. J., & Kappenman, E. S. (Eds.). (2013). The Oxford handbook of event-related potential components. Oxford university press.

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG-and MEG-data. Journal of neuroscience methods, 164(1), 177-190.

Mathewson, K. E., Lleras, A., Beck, D. M., Fabiani, M., Ro, T., & Gratton, G. (2011). Pulsed out of awareness: EEG alpha oscillations represent a pulsed-inhibition of ongoing cortical processing. Frontiers in psychology, 2, 99.

Mello, M., Fusaro, M., & Aglioti, S. M. (2024). The neuroscience of human empathy for pleasure: protocol for a scoping review. Systematic Reviews, 13(1), 82.

Mello, M., Fusaro, M., Tieri, G., & Aglioti, S. M. (2022). Wearing same- and opposite-sex virtual bodies and seeing them caressed in intimate areas. *Quarterly Journal of Experimental Psychology*, *75*(3), 461–474. https://doi.org/10.1177/17470218211031557

Monti, A., Porciello, G., Tieri, G., & Aglioti, S. M. (2020). The "embreathment" illusion highlights the role of breathing in corporeal awareness. *Journal of Neurophysiology*, *123*(1), 420–427. https://doi.org/10.1152/jn.00617.2019

Moreau, Q., Candidi, M., Era, V., Tieri, G., & Aglioti, S. M. (2020). Midline frontal and occipito-temporal activity during error monitoring in dyadic motor interactions. *Cortex*, *127*, 131–149. https://doi.org/10.1016/j.cortex.2020.01.020

Morelli, S. A., Sacchet, M. D., & Zaki, J. (2015). Common and distinct neural correlates of personal and vicarious reward: A quantitative meta-analysis. NeuroImage, 112, 244-253.

Morrison, I. (2016). Keep calm and cuddle on: social touch as a stress buffer. Adaptive Human Behavior and Physiology, 2, 344-362.

Mu, Y., Fan, Y., Mao, L., & Han, S. (2008). Event-related theta and alpha oscillations mediate empathy for pain. Brain research, 1234, 128-136.

Nicolardi, V., Fanizza, I., Accogli, G., Macchitella, L., Scoditti, S., & Trabacca, A. (2023). Pain assessment in autism: updating the ethical and methodological challenges through a state-of-the-art review. Neurological Sciences, 44(11), 3853-3861.

Nicolardi, V., Panasiti, M. S., D'Ippolito, M., Pecimo, G. L., & Aglioti, S. M. (2020). Pain perception during social interactions is modulated by self-related and moral contextual cues. Scientific Reports, 10(1), 1-12.

Novembre, G., Sammler, D., & Keller, P. E. (2016). Neural alpha oscillations index the balance between self-other integration and segregation in real-time joint action. Neuropsychologia, 89, 414-425.

Oberman, L. M., McCleery, J. P., Ramachandran, V. S., & Pineda, J. A. (2007). EEG evidence for mirror neuron activity during the observation of human and robot actions: Toward an analysis of the human qualities of interactive robots. Neurocomputing, 70(13-15), 2194-2203.

Olofsson, J. K., Nordin, S., Sequeira, H., & Polich, J. (2008). Affective picture processing: an integrative review of ERP findings. Biological psychology, 77(3), 247-265.

Pamplona, G. S., Salgado, J. A., Staempfli, P., Seifritz, E., Gassert, R., & Ionta, S. (2022). Illusory Body Ownership Affects the Cortical Response to Vicarious Somatosensation. Cerebral Cortex, 32(2), 312-328.

Pavone, E. F., Tieri, G., Rizza, G., Tidoni, E., Grisoni, L., & Aglioti, S. M. (2016). Embodying Others in Immersive Virtual Reality: Electro-Cortical Signatures of Monitoring the Errors in the Actions of an Avatar Seen from a First-Person Perspective. *Journal of Neuroscience*, *36*(2), 268–279. https://doi.org/10.1523/JNEUROSCI.0494-15.2016

Peled-Avron, L., Goldstein, P., Yellinek, S., Weissman-Fogel, I., & Shamay-Tsoory, S. G. (2018). Empathy during consoling touch is modulated by mu-rhythm: An EEG study. Neuropsychologia, 116, 68-74.

Peled-Avron, L., Levy-Gigi, E., Richter-Levin, G., Korem, N., & Shamay-Tsoory, S. G. (2016). The role of empathy in the neural responses to observed human social touch. Cognitive, Affective, & Behavioral Neuroscience, 16(5), 802-813.

Perry, A., Stein, L., & Bentin, S. (2011). Motor and attentional mechanisms involved in social interaction—Evidence from mu and alpha EEG suppression. Neuroimage, 58(3), 895-904.

Pezzetta, R., Nicolardi, V., Tidoni, E., & Aglioti, S. M. (2018). Error, rather than its probability, elicits specific electrocortical signatures: a combined EEG-immersive virtual reality study of action observation. Journal of neurophysiology, 120(3), 1107-1118.

Pezzetta, R., Ozkan, D. G., Era, V., Tieri, G., Zabberoni, S., Taglieri, S., ... & Aglioti, S. M. (2023). Combined EEG and immersive virtual reality unveil dopaminergic modulation of error monitoring in Parkinson's Disease. npj Parkinson's Disease, 9(1), 3.

Pfurtscheller, G., Scherer, R., Leeb, R., Keinrath, C., Neuper, C., Lee, F., & Bischof, H. (2007). Viewing moving objects in virtual reality can change the dynamics of sensorimotor EEG rhythms. Presence, 16(1), 111-118.

Pineda, J. A., & Hecht, E. (2009). Mirroring and mu rhythm involvement in social cognition: are there dissociable subcomponents of theory of mind?. Biological psychology, 80(3), 306-314.

Riečanský, I., Lengersdorff, L. L., Pfabigan, D. M., & Lamm, C. (2020). Increasing self-other bodily overlap increases sensorimotor resonance to others' pain. Cognitive, Affective, & Behavioral Neuroscience, 20(1), 19-33.

Riečanský, I., Paul, N., Kölble, S., Stieger, S., & Lamm, C. (2014). Beta oscillations reveal ethnicity ingroup bias in sensorimotor resonance to pain of others. Social cognitive and affective neuroscience, 10(7), 893-901.

Sarlo, M., Buodo, G., Poli, S., & Palomba, D. (2005). Changes in EEG alpha power to different disgust elicitors: the specificity of mutilations. Neuroscience letters, 382(3), 291-296.

Schirmer, A., & McGlone, F. (2019). A touching sight: EEG/ERP correlates for the vicarious processing of affectionate touch. Cortex, 111, 1-15.

Schirmer, A., Lai, O., Cham, C., & Lo, C. (2023). Velocity-tuning of somatosensory EEG predicts the pleasantness of gentle caress. NeuroImage, 119811.doi: 10.1016/j.neuroimage.2022.119811. Epub 2022 Dec 13.

Schupp, H. T., Junghöfer, M., Weike, A. I., & Hamm, A. O. (2004). The selective processing of briefly presented affective pictures: an ERP analysis. Psychophysiology, 41(3), 441-449.

Seinfeld, S., Schmidt, I., & Müller, J. (2022). Evoking realistic affective touch experiences in virtual reality. arXiv preprint arXiv:2202.13389.

Singer, T., & Lamm, C. (2009). The social neuroscience of empathy. Annals of the New York Academy of Sciences, 1156(1), 81-96.

Slater, M., Spanlang, B., Sanchez-Vives, M. V, & Blanke, O. (2010). First person experience of body transfer in virtual reality. *PloS One*, *5*(5), e10564. https://doi.org/10.1371/journal.pone.0010564

Smit S, Moerel D, Zopf R, Rich AN (2023) Vicarious touch: Overlapping neural patterns between seeing and feeling touch Neuroimage Sep:278:120269. doi: 10.1016/j.neuroimage.2023.120269.

Tieri, G., Gioia, A., Scandola, M., Pavone, E. F., & Aglioti, S. M. (2017). Visual appearance of a virtual upper limb modulates the temperature of the real hand: a thermal imaging study in Immersive Virtual Reality. *European Journal of Neuroscience*, 1–11. https://doi.org/10.1111/ejn.13545

Tieri, G., Morone, G., Paolucci, S., & Iosa, M. (2018). Virtual reality in cognitive and motor rehabilitation: facts, fiction and fallacies. Expert Review of Medical Devices, 15(2), 107–117. https://doi.org/10.1080/17434440.2018.1425613

Tieri, G., Tidoni, E., Pavone, E. F., & Aglioti, S. M. (2015a). Mere observation of body discontinuity affects perceived ownership and vicarious agency over a virtual hand. *Experimental Brain Research*, *233*(4), 1247–1259. https://doi.org/10.1007/s00221-015-4202-3

Tieri, G., Tidoni, E., Pavone, E. F., & Aglioti, S. M. (2015b). Body visual discontinuity affects feeling of ownership and skin conductance responses. *Scientific Reports*, *5*, 17139. https://doi.org/10.1038/srep17139

Todd, R. M., Cunningham, W. A., Anderson, A. K., & Thompson, E. (2012). Affect-biased attention as emotion regulation. Trends in cognitive sciences, 16(7), 365-372.

Uusberg, A., Uibo, H., Kreegipuu, K., & Allik, J. (2013). EEG alpha and cortical inhibition in affective attention. International Journal of Psychophysiology, 89(1), 26-36.

Valentini, E., Koch, K., Nicolardi, V., & Aglioti, S. M. (2015). Mortality salience modulates cortical responses to painful somatosensory stimulation: Evidence from slow wave and delta band activity. NeuroImage, 120, 12-24.

Valentini, E., Nicolardi, V., & Aglioti, S. M. (2017a). Visual reminders of death enhance
    nociceptive-related cortical responses and event-related alpha desynchronisation. Biological
    Psychology, 129, 121-130.

Valentini, E., Nicolardi, V., & Aglioti, S. M. (2017b). Painful engrams: Oscillatory correlates of
    working memory for phasic nociceptive laser stimuli. Brain and cognition, 115, 21-32.

Valeriani, M., Betti, V., Le Pera, D., De Armas, L., Miliucci, R., Restuccia, D., ... Aglioti, S. M.
    (2008). Seeing the pain of others while being in pain: A laser-evoked potentials study.
    NeuroImage, 40(3), 1419–1428. https://doi.org/10.1016/j.neuroimage.2007.12.056.

Van Ede, F., De Lange, F., Jensen, O., & Maris, E. (2011). Orienting attention to an upcoming
    tactile event involves a spatially and temporally specific modulation of sensorimotor alpha-and
    beta-band oscillations. Journal of Neuroscience, 31(6), 2016-2024.

Figures

Figure 1 | A: representation of virtual environment and the two avatars sat in front of the table facing each other with a grey panel placed between them. B: experimental stimuli observed from 1PP and 3PP (pain, neutral and pleasure).

Figure 2 | Sequence of events for each trial. VAS 1 and 2, refers to items 1 and 2 of the Table 1.

Figure 3 | Visual Analog Scale (VAS) ratings (means and SE) of valence (**A**), intensity (**B**), and ownership (**C**) elicited by virtual pain, pleasure and neutral stimuli in 1PP and 3PP.

Figure 4 | VEPs results. Grand average waveform and scalp topographies of VEPs observed at POz-Oz electrodes. The ANOVA with correction for multiple comparison revealed a main effect of both, valence between 160-400 ms (left, indicated by the first yellow rectangle), and perspective between 190-310 ms (right, in the yellow rectangle), on the N2-P2 complex while a main effect of valence only, on the LPP between 400-600 ms (left, second yellow rectangle), with no significant interaction. Post-hoc comparisons showed Greater amplitude for pain and pleasure as compared to neutral, as well as for 1PP compared to 3PP, was found for the N2-P2 waveforms. The main effect of valence on the LPP was accounted by greater amplitude for neutral and pain compared to pleasure.

Figure 5 | EPN results. Grand average waveforms and topography for EPN at the electrodes P5-P7-PO7. The ANOVA with correction for multiple comparison on EPN revealed a main effect of valence between 200-500ms (left, yellow rectangles) and perspective between 260-500ms (right, yellow rectangle), with no significant interaction. The main effect of perspective was accounted by greater amplitude for 1PP compared to 3PP. Post-hoc comparisons between the levels of valence indicate greater amplitude for pain and pleasure compared to neutral

Figure 6 | Time-frequency results. Main effects of Valence and Perspective Grand average of EEG total power for the two main effects of valence and perspective at the Central ROI (FCz, Cz, C3) and significant comparisons with corrected p-values within the green-background boxes. For the main effect of valence, single comparisons between the three levels (neutral, pain and pleasure), did not showed significant differences after correcting p-value for multiple comparisons (500 permutation, see methods). The main effect of perspective shows increased theta power in 1PP compared to 3PP between 300-500ms, and decreased alpha power in 1PP compared to 3PP between 500-1000ms.

Figure 7 | Time-frequency results. (A): Grand average of EEG total power at the central ROI and significant comparison within the green-background boxes for the significant interaction Valence x Perspective. (B) and (C): t-test for the significant interaction corrected for multiple comparisons. B is for 1PP and C for 3PP. Red-background boxes indicate significant results and corrected p-values

Tables

Item	Index	stimulus-related sensation
1	*Valence*	was the stimulus pleasant or unpleasant for you?
2	*Intensity*	how intense was the stimulus for you?

	Index	body ownership
3	*Ownership*	I felt as if I were looking at my own right hand
4	*Ownership*	I felt as if the virtual hand were part of my body
5	*control*	It felt as if I had more than one right hand

Table 1 | VAS questions concerning the stimulus's valence (Item 1: "−−" = strongly unpleasant, middle=neutral, "++" strongly pleasant) and intensity (Item 2: "−−" = very weak, "++" = very strong) and VAS statements assessing the illusory feeling of body ownership (Item 3-5: "−−"=no illusion; "++"= maximal illusion).

	First-person perspective (1PP)			Third-person perspective (3PP)		
	Pleasure (mean ± SD)	Pain (mean ± SD)	Neutral (mean ± SD)	Pleasure (mean ± SD)	Pain (mean ± SD)	Neutral (mean ± SD)
Valence	66.97 ± 14.92	27.67 ± 17.24	50.75 ± 6.26	64.18 ± 16.50	26.17 ± 16.25	50.75 ± 6.05
Intensity	55.97 ± 23.25	52.92 ± 25.62	35.26 ± 21.15	49.49 ± 24.35	48.06 ± 25.14	33.65 ± 21.71
Ownership (experimental items)	66.78 ± 17.79	69.57 ± 17.44	64.10 ± 22.25	32.30 ± 23.29	32.67 ± 24.13	29.26 ± 24.28
Ownership (control item)	23.88 ± 23.66	20.83 ± 22.66	26.67 ± 26.02	24.05 ± 26.17	25.88 ± 26.20	19.24 ± 23.99

Table 2 | Means and SDs of each VAS rating (UnPleasantness, Intensity and Ownership) of the type of observed stimulus (pain-pleasure-neutral) in the different Perspectives (1PP and 3PP). High and low UnPleasantness values indicate pleasant and unpleasant sensations, respectively.

Rating	Post-hoc comparisons (mean ± SD) (* = p < 0.05; ** = p < 0.001)
Valence	Pain 1PP (27.67 ± 17.24) < Pleasure 1PP (66.97 ± 14.92); **
	Pain 1PP (27.67 ± 17.24) < Neutral 1PP (50.75 ± 6.26); **
	Pleasure 1PP (66.97 ± 14.92) > Neutral 1PP (50.75 ± 6.26); **
	Pain 3PP (26.17 ± 16.25) < Pleasure 3PP (64.18 ± 16.50); **
	Pain 3PP (26.17 ± 16.25) < Neutral 3PP (50.75 ± 6.05); **
	Pleasure 3PP (64.18 ± 16.50) > Neutral 3PP (50.75 ± 6.05); **
Intensity	Pain 1PP (52.92 ± 25.62) > Neutral 1PP (35.26 ± 21.15);*
	Pleasure 1PP (55.97 ± 23.25) < Neutral 1PP (35.26 ± 21.15);**
	Pleasure 3PP (49.49 ± 24.35) < Neutral 3PP (33.65 ± 21.71);**
Ownership	Experimental, Pain 1PP (69.57 ± 17.44) > Control, Pain 1PP (20.83 ± 22.66);**
	Experimental, Pleasure 1PP (66.78 ± 17.79) > Control, Pleasure 1PP (23.88 ± 23.66);**
	Experimental, Neutral 1PP (64.10 ± 22.25) > Control, Neutral 1PP (26.67 ± 26.02);**
	Experimental, Pain 1PP (69.57 ± 17.44) > Experimental, Pain 3PP (32.67 ± 24.13);**
	Experimental, Pleasure 1PP (66.78 ± 17.79) > Experimental, Pleasure 3PP (32.30 ± 23.29);**
	Experimental, Neutral 1PP (64.10 ± 22.25) > Experimental, Neutral 3PP (29.26 ± 24.28);**

Table 3 | Summary of the main results regarding the subjective UnPleasantness, Intensity and Ownership.

Index	Frequency Band	Subjective Rating	Pearson R	p-value
Pleasure 1PP - Neutral 1PP	Beta 800-1500ms	Pleasantness	0.15	0.47
Pleasure 1PP - Neutral 1PP	Beta 800-1500ms	Intensity	0.38	0.07
Pleasure 3PP - Neutral 3PP	Beta 800-1500ms	Pleasantness	0.07	0.75
Pleasure 3PP - Neutral 3PP	Beta 800-1500ms	Intensity	-0.18	0.40
Pleasure 3PP - Neutral 3PP	Theta 800-1500ms	Pleasantness	-0.07	0.75
Pleasure 3PP - Neutral 3PP	Theta 800-1500ms	Intensity	0.36	0.08
Pleasure 3PP - Neutral 3PP	Alpha 800-1500ms	Pleasantness	-0.08	0.70
Pleasure 3PP - Neutral 3PP	Alpha 800-1500ms	Intensity	0.12	0.57
Pleasure 1PP - Neutral 1PP	Theta 200-350ms	Pleasantness	0.09	0.66
Pleasure 1PP - Neutral 1PP	Theta 200-350ms	Intensity	0.01	0.97
Pain 1PP - Neutral 1PP	Theta 200-350ms	Pleasantness	0.26	0.22
Pain 1PP - Neutral 1PP	Theta 200-350ms	Intensity	0.15	0.50

Table 4 | Correlations between Subjective Ratings and EEG in the time-frequency domain. We used Pearson correlation to test the associations between the EEG power and the Unpleasantness and Intensity ratings. Indexes were calculated (separately for each perspective) by subtracting the Neutral condition from Pleasure and Pain.