The expression patterns of different cell types and their interactions in the tumor microenvironment are predictive of breast cancer patient response to neoadjuvant chemotherapy

Saugato Rahman Dhruba[1], Sahil Sahni[1], Binbin Wang[1], Di Wu[2], Padma Sheila Rajagopal[1,3], Yael Schmidt[1], Eldad D. Shulman[1], Sanju Sinha[1,4], Stephen-John Sammut[5,6], Carlos Caldas[7,8], Kun Wang[1,*], Eytan Ruppin[1,*]

[1] Cancer Data Science Laboratory, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA.

[2] Laboratory of Pathology, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA.

[3] Women's Malignancies Branch, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA.

[4] NCI-Designated Cancer Center, Sanford Burnham Prebys Medical Discovery Institute, San Diego, CA, USA.

[5] Breast Cancer Now Toby Robins Research Centre, The Institute of Cancer Research, London, UK.

[6] The Royal Marsden Hospital NHS Foundation Trust, London, UK.

[7] Institute of Metabolic Science, School of Clinical Medicine, University of Cambridge, Cambridge, UK.

[8] Department of Clinical Biochemistry, University of Cambridge, Cambridge, UK.

[*] Co-corresponding authors (eytan.ruppin@nih.gov, kun.wang@nih.gov)

ABSTRACT

The tumor microenvironment (TME) is a complex ecosystem of diverse cell types whose interactions govern tumor growth and clinical outcome. While the TME's impact on immunotherapy has been extensively studied, its role in chemotherapy response remains less explored. To address this, we developed DECODEM (DEcoupling Cell-type-specific Outcomes

using <u>DE</u>convolution and <u>M</u>achine learning), a generic computational framework leveraging

cellular deconvolution of *bulk transcriptomics* to associate the gene expression of individual cell

types in the TME with clinical response. Employing DECODEM to analyze the gene expression of

breast cancer (BC) patients treated with neoadjuvant chemotherapy, we find that the gene

expression of specific immune cells (*myeloid*, *plasmablasts*, *B-cells*) and stromal cells (*endothelial*,

normal epithelial, *CAFs*) are highly predictive of chemotherapy response, going beyond that of

the malignant cells. These findings are further tested and validated in a single-cell cohort of triple

negative breast cancer. To investigate the possible role of immune cell-cell interactions (CCIs) in

mediating chemotherapy response, we extended DECODEM to DECODEMi to identify such CCIs,

validated in single-cell data. Our findings highlight the importance of active pre-treatment

immune infiltration for chemotherapy success. The tools developed here are made publicly

available and are applicable for studying the role of the TME in mediating response from readily

available bulk tumor expression in a wide range of cancer treatments and indications.

INTRODUCTION

The tumor microenvironment (TME) is a complex and multifaceted ecosystem that encompasses

a plethora of cell types including immune cells (such as T- and B-lymphocytes, macrophages,

dendritic cells *etc.*), stromal cells (such as fibroblasts, endothelial cells *etc.*), and the extracellular

matrix. It plays crucial roles in regulating tumor progression and response to therapy. Emerging

evidence suggest the context-specific behavior of the TME as either tumor-suppressive or

supportive, presenting an attractive target for therapeutic intervention [1][2]. Numerous studies

have elucidated the critical role of TME in mediating response to immunotherapy, however,

insights into how the TME is relevant to chemotherapy response remains limited. Here we focus

on studying the influence of the TME on chemotherapy response in breast cancer (BC), where

neoadjuvant chemotherapy (NAC) is used prior to primary surgery to enable tumor downstaging

and increase the likelihood of breast-conserving surgery [3][4][5].

Recent studies have underscored the relevance of the BC TME in determining patient

response to chemotherapy. For instance, chemotherapy can induce dynamic changes in the TME

that lead to an increase in regulatory T-cells and myeloid-derived suppressor-like cells, calling for

a balance in immune population for positive treatment outcomes [6]. Lymphocyte density and

tumor infiltrating lymphocytes have consistently been associated with BC patients achieving

pathological complete response (pCR) following NAC and anti-HER2 therapy [7][8][9][10][11].

Additionally, multiple studies have reported the active involvement of TME in therapy resistance

[12][13][14], with macrophages, endothelial, and BC stem cells promoting chemoresistance

[15][16][17], and B-cells and plasma cells displaying varying roles in NAC response [18][19][20].

However, to our knowledge, no study has yet attempted to quantify the contribution and

predictive power of each individual cell type in the TME to chemotherapy response in a systematic

and rigorous manner. We, therefore, aim to investigate three fundamental research questions:

(1) Can we apply a rigorous computational approach to delineate the cell-type-specific

transcriptome predictive power in the BC TME from bulk transcriptomics data? (2) Which are the

cell types in the BC TME and the corresponding pathways whose gene expression are most

predictive of chemotherapy response? (3) Does aggregating the inferred expression of multiple

predictive cell types in the BC TME lead to a more accurate predictor?

Current advances in precision oncology have highlighted the utility of artificial intelligence

[1][2][21][22][23][24] in guiding treatment using diverse data types such as genomics [1][25][26],

transcriptomics [29][30][31][32][33], histopathology [34][35][36][37][38][39], or through

multimodal approaches [11][27][28][40][41][42]. However, investigations into neoadjuvant

therapy response often encounter limitations like small sample size, treatment heterogeneity, or

inadequate capturing of the TME's complexity, as highlighted by Sammut et al. [11]. In their

pioneering work, Sammut et al. presented a novel multiomic predictor of neoadjuvant therapy

response in BC [11]. They have leveraged machine learning (ML) to incorporate 34 features

extracted from clinical, DNA, RNA, pathology, and treatment information to develop a single

model to predict patient response across all BC subtypes and different treatment regimens with

high accuracy. Expanding on their groundwork and addressing the limitations, our study aims to

decouple the cell-type-specific effects to NAC response in the BC TME from transcriptomics alone

by focusing on the HER2-negative patients, where NAC is recommended to increase the likelihood

of achieving pCR following surgery, especially in triple-negative breast cancer (TNBC) [49].

To systematically explore the association between the TME and chemotherapy response,

we developed a computational framework, **DECODEM (DEcoupling Cell-type-specific Outcomes**

using DEconvolution and Machine learning). This framework leverages cellular deconvolution

with machine learning to elucidate the association of gene expression of diverse cell types in the

TME to patient response to a given treatment. Recognizing that these phenotype effects are

intricately regulated by the interactions among relevant cell types within the TME, we further

extended DECODEM to **DECODEMi**, where 'i' stands for interaction. DECODEMi incorporates the

inferred cell-to-cell communications to pinpoint the key cell-cell interactions (CCIs) influencing

treatment response. As mentioned, we demonstrate the utility of our tools in an example

scenario of investigating the role of the BC TME in modulating response to NAC. A recent study

by Kester et al. investigated the association of cellular compositions of the TME with patient

survival in BC [43], but to our knowledge, the current study is the first to analyze the *cell-type-*

specific transcriptomic profiles in the BC TME to objectively quantify their association with

treatment response.

Through the application of DECODEM, we illustrate the markedly improved predictive

capabilities of cell-type-specific transcriptome over bulk transcriptome, untangling the individual

contributions of different cell types in treatment response. Through the application of DECODEMi,

we further pinpoint the notable immune interactions among these cell types. Together all these

findings point to the necessity of an active pre-treatment immune system to facilitate positive

treatment outcome, as supported by Sammut et al. [11]. Therefore, from a methodological

perspective, our study provides tools to systematically assess the role of individual cellular

components within the TME in anticancer therapy response (DECODEM), and to identify the

prominent cell-to-cell communications associated with this response (DECODEMi). These tools

will be made publicly available upon publication.

RESULTS

Overview of the analysis

DECODEM and DECODEMi developed and presented here are new computational frameworks designed to explore the predictive powers of individual cell types and relevant cell-cell interactions, respectively, within the TME to response to therapy. They harness the prowess of CODEFACS and LIRICS, two recent tools developed in our laboratory [44]. CODEFACS facilitates the extraction of the cell-type-specific expression profiles from bulk gene expression data for each patient, while LIRICS discerns the activity of cell-type-specific ligand-receptor interactions within the patient's TME. In essence, DECODEM takes the pre-treatment bulk gene expression as input and outputs the quantified predictive power (in terms of AUC and AP *i.e.*, the area under the receiver operating characteristics (ROC) curve and average precision, equivalent to the area under the precision-recall curve, respectively) of each cell type to treatment response. DECODEMi then follows through to extract the most predictive cell-cell interactions among these cell types.

Specifically, DECODEM comprises of two sequential steps: *First*, for each bulk expression cohort, it generates nine cell-type-specific expression profiles (as annotated by Wu et al. [45] which were used for molecular signature generation; see details in **METHODS**: *B-cells*, *Cancer-associated Fibroblasts (CAFs)*, *Cancer Epithelial (malignant)*, *Endothelial*, *Myeloid*, *Normal Epithelial*, *Plasmablasts*, *Perivascular-like Cells (PVL)*, and *T-cells*) in each sample using CODEFACS; *Second*, by leveraging these nine deconvolved expression profiles, it generates nine cell-type-specific predictors of clinical response (responders vs. non-responders *e.g.*, pCR vs. residual disease) using a multi-stage ML pipeline (**Figure 1**). Within this pipeline, we performed rigorous feature selection to keep the top 'm' genes per cell type ($2 \leq m \leq 25$; the upper limit is chosen to avoid overfitting and facilitate faster runtime) before feeding their expression into an unweighted ensemble classifier consisting of four ML algorithms: regularized logistic regression, random forest, support vector machine, and XGBoost (see details in **METHODS**). The pipeline was trained with repeated three-fold cross-validation (CV) to optimize model hyperparameters. Additionally, we built a baseline predictor with bulk expression using the same ML pipeline to serve as a control, comparative rod.

To assess the performance of DECODEM and evaluate the contribution of each cell type,

we analyzed bulk transcriptomics data from three breast cancer cohorts with different ratios of

responders to non-responders (R:NR) to NAC (**Supplementary Figure 1**) in two scenarios: (a)

cross-validation with an additional five-fold CV on the TransNEO cohort (n = 94, R:NR = 22:72),

recently published by Sammut et al. accompanied by a multiomic clinical response predictor [11],

and (b) external validation on the ARTemis + PBCP (n = 55, R:NR = 8:47) and BrighTNess (n = 106,

R:NR = 65:41) cohorts [46][47]. TransNEO and ARTemis + PBCP included all major BC subtypes,

among which we focused on the HER2-negative patients [11][46] (**Supplementary Figure 1B-C**),

whereas the BrighTNess cohort included only TNBC patients from arm B of the trial, treated with

a combination of paclitaxel and carboplatin with a placebo for veliparib [47]. Moreover, we

established DECODEM's generalizability to single-cell transcriptomics through external validation

on an independent cohort from Zhang et al, which consisted of 15 TNBC patients who received

either paclitaxel alone (R:NR = 3:3) or in combination with atezolizumab, an anti-PD-L1 agent

(R:NR = 4:5) [48]. Notably, we considered the Sammut et al. predictor built with 18 extracted

features from clinical and RNA-seq data (e.g. not including the DNA-seq data) as a comparative

rod, which achieved an AUC of 0.88 in CV and 0.89 in validation with ARTemis + PBCP [11]. This

predictor outperformed a baseline predictor built with seven clinical features including ER status

and HER2 status, which are routinely used to guide clinical decisions [49].

To investigate the effect of incorporating the expression profiles of multiple predictive cell

types together, as indicated by our third question, we applied two extensions to DECODEM: (i)

append the cell-type-specific expression profiles for two and three most predictive cell types

together to train a multi-cell-ensemble ML pipeline, and (ii) extend to DECODEMi to incorporate

the inferred CCIs to train a CCI-based ML pipeline (**Figure 1**; see details in **METHODS**). The former

model maximizes the predictive performance by leveraging the complementary information from

different cell types while the latter identifies the most predictive cell-to-cell communications for

clinical response, thereby providing deeper insights to the underlying mechanism.

Figure 1: The analysis pipeline, DECODEM and DECODEMi. *First*, we apply CODEFACS on bulk expression to generate nine cell-type-specific expression profiles and further apply LIRICS to infer the cell-cell interactions (CCIs) present in the tumor microenvironment. *Second*, we train a multi-stage machine learning pipeline with either the nine cell-type-specific expression profiles to build nine cell-type-specific clinical response predictors (DECODEM) or the CCI profile to build a CCI-based predictor (DECODEMi). In validation, we deploy the abovementioned first stage to procure the cell-type-specific expression or CCIs and directly feed those as inputs to the trained predictors to evaluate model performance.

Gene expression of immune and stromal cells predicts response to neoadjuvant chemotherapy as accurately as the expression of malignant cells

To assess the impact of individual cell types on chemotherapy response, we applied DECODEM to TransNEO in a CV analysis. Our analysis demonstrated significant differences (one-tailed Wilcoxon rank-sum test p-value ≤ 0.001) in the resulting prediction scores between responders and non-responders across 91 patients (the patients with available predictions from Sammut et al.), with the identified cell types outperforming bulk and Sammut et al. predictors (**Figure 2A**). Seven out of nine cell types exhibited markedly enhanced predictive capabilities in identifying responders.

Notably, immune cells (*myeloid*, *plasmablasts*, *B-cells*) contribute as prominently as the malignant

(*cancer epithelial*) and stromal cells (*endothelial*, *normal epithelial*, *CAFs*) (**Figure 2B**). Our

subsequent analyses, focused on these seven 'prominent' predictive cell types, whose abilities to

accurately identify true responders to chemotherapy in terms of the odds ratio and sensitivity

values are shown in **Supplementary Figure 2A,C**.

These cell types have frequently been reported to influence clinical response in BC,

particularly in the context of chemotherapy-induced immunogenic cell death as the proposed

mechanism [51][11]. Changes in the immune population, such as an increase in regulatory T-cells

and myeloid-derived suppressor-like cells, and a decrease in CD8$^+$ T-cells, have been associated

with achieving pCR following NAC [6]. Myeloid cells have been implicated in therapy response

through interactions with the malignant and endothelial cells, inducing chemoresistance *via* the

CXCL1/2 − S100A8/9 loop, whereas tumor-associated macrophages have been linked to adverse

outcomes and chemoresistance through survival factor secretion, anti-apoptotic pathway

activation, and modulation of signaling pathways [6][15][16]. *VEGF*-mediated upregulation of

survivin in Endothelial cells was further implicated in playing central roles in chemoresistance

[15]. Additionally, plasmablasts, as precursors of plasma cells, have been associated with better

prognosis [19][20], while B-cells have shown both pro- and anti-tumor roles [18][19][20].

We further tested and validated the predictive power of these cell types independently in

external validation using the ARTemis + PBCP (*n* = 55) and BrighTNess (*n* = 106) cohorts, observing

consistent results (**Figure 2C-D, Supplementary Figure 2A,C**). In the ARTemis + PBCP cohort

(dominated by ER-positive, HER2-negative BC), the prominent contributors to chemotherapy

response were CAFs, normal epithelial, and cancer epithelial (**Figure 2C**), whereas in the TNBC-

specific BrighTNess cohort, the key players were myeloid, plasmablasts, and cancer epithelial

(**Figure 2D**). When stratifying TransNEO and ARTemis + PBCP cohorts by the available BC subtypes,

our cell-type-specific models retain good performance across both subtypes (**Supplementary**

Figure 3A-D). The models for ER-positive, HER2-negative BC were more predictive than TNBC,

possibly due to sample size limitations as well as increased heterogeneity in TNBC. These findings

support the prominent involvement of immune and stromal cells alongside the malignant cells in

mediating patient response to chemotherapy in the TME, with variations observed likely due to

tumor subtype composition and ER/HER2 status differences across cohorts.

Our analysis did not identify T-cells as a prominent predictive cell type, as their expression

provide marginal to no enhancement in predictive power over bulk expression (**Figure 2A-B**). This

lack of signal may stem from aggregating different T-cell subtypes together (due to limits on the

resolution of the deconvolution, which breaks down for low-abundance cell types). To study this

further, we performed an enrichment analysis to delineate the contributions of $CD4^+$ and $CD8^+$ T-

cells to chemotherapy response. However, the expression of each of these cell types still exhibited

limited stratification power in distinguishing responders from non-responders across the three

bulk cohorts (**Supplementary Figure 4A-D**). Additionally, we complemented the expression-based

analysis with an analysis of the association between cell abundances and clinical response. This

revealed that the overall T-cell abundance exhibited predictive power similar to or even higher

than $CD4^+$ and $CD8^+$ subtypes (**Supplementary Figure 4E-G**), albeit still significantly lower than

that of bulk transcriptomics.

Figure 2: The prominent cell types mediating chemotherapy response in breast cancer tumor microenvironment.

A-B. Comparison of prediction scores (**A**) and model performance (**B**) in cross-validation with TransNEO (*n* = 91) for nine cell-type-specific, bulk, and Sammut et al. predictors. R and NR stand for responders and non-responders. The differences between the prediction scores were computed by using a one-tailed Wilcoxon rank-sum test. AUC and AP stand for the area under the receiver operating characteristics curve and average precision (equivalent to the area under precision-recall curve), respectively. Cell types are ranked by their AUC values in a descending order.

C. Comparison of model performance in external validation with ARTemis + PBCP (*n* = 55) for the seven prominent cell types, bulk, and Sammut et al. predictors. Cell types are ranked by their AUC values in a descending order.

D. Comparison of model performance in external validation with BrighTNess (*n* = 106) for the seven prominent cell types and bulk predictors. Cell types are ranked by their AUC values in a descending order.

An ensemble model incorporating the expression of immune and stromal cells significantly boosts the predictive power of patient response

To further enhance DECODEM's performance, we aggregated multiple individual cell types together by appending the corresponding expression profiles as the input features. We explored 35 combinations of two to three cell types exhaustively, This comprehensive analysis yielded a boost in predictive performance in both ARTemis + PBCP and BrighTNess cohorts, with significant overlaps observed in the top multi-cell-type combinations across cohorts (**Figure 3A-D, Supplementary Figure 2B,D**). Remarkably, the ensemble of immune and stromal cells, specifically the combination of the expressions of endothelial, myeloid, and plasmablast cells exhibited the strongest stratification capability in identifying chemotherapy responders (*ARTemis + PBCP*: AUC = 0.94, AP = 0.78, **Figure 3B**; *BrighTNess*: AUC = 0.75, AP = 0.81, **Figure 3D**). Additionally, the immune-cell-ensemble comprising myeloid and plasmablasts displayed high stratification (*ARTemis + PBCP*: AUC = 0.92, AP = 0.66, **Figure 3A**; *BrighTNess*: AUC = 0.73, AP = 0.80, **Figure 3C**). These findings testify to the complementary contributions of different cell types within the TME to chemotherapy response. However, the top ensemble for the BrighTNess cohort provided a similar performance to the top cell type, myeloid (AUC = 0.76, AP = 0.83, **Figure 2D**), suggesting that the interactions among different cell types are less prominent in this cohort compared to the other cohorts, illustrated further in **Figure 4D**. The subtype-specific stratification of ARTemis + PBCP reinforced the superior predictive capabilities of these ensembles across both available BC subtypes while recapitulating their diminished impact in TNBC (**Supplementary Figure 3E-F**).

We conducted gene set enrichment analysis (GSEA) with Reactome pathways [52] to explore the functional contributions of these prominent cell types in chemotherapy response prediction. This analysis revealed the enrichment of cell cycle checkpoint and DNA replication pathways across different cell types (false discovery rate (FDR)-adjusted *p*-value ≤ 0.2; **Figure 3E**)

as the most common functional contribution. Dysregulation of these pathways are frequent in BC

and promotes tumor proliferation [53][54][55], making them valuable prognostic markers and

therapeutic targets [56][57][58][59]. Immune cells exhibited enrichment in signal transduction

pathways (*e.g.*, *VEGFA − VEGFR2* and *VEGF* signaling pathways in plasmablasts and multiple

NOTCH1 signaling pathways in myeloid), implicated in malignant progression and therapy

resistance, particularly in TNBC [60][61][62][63][64][65][66]. Stromal endothelial cells showed

enrichment in development- and immune-related pathways (*e.g.*, *L1CAM* interactions and

antigen processing & presentation), contributing to aggressive BC progression and immune

evasion with implications for both chemotherapy and immunotherapy

[67][68][69][70][71][72][73].

To delve deeper into the underlying mechanism, we assessed DECODEM's performance in

light of the prevalent chemotherapy regimen within each cohort. TransNEO and ARTemis + PBCP

predominantly employed T-FEC (*n* = 61 and 40, respectively; including docetaxel/taxotere, 5-

flurouracil, epirubicin, and cyclophosphamide), while BrighTNess used a uniform regimen of

paclitaxel and carboplatin (**Supplementary Figure 1C**). All six drugs have known clinical activities

stimulating anticancer immunity *via* releasing immunostimulatory molecules from cancer cells

(*on-target* effects) or promoting activation of effector cells and diminishing immunosuppressive

cells (*off-target* effects) [73]. Remarkably, across all three cohorts spanning two BC subtypes and

two regimens, DECODEM identified myeloid and plasmablasts to consistently stratify responders

to *immunogenic* drug regimens, alongside malignant cells (**Supplementary Figure 5A,B,D**). For

ARTemis + PBCP (dominated by ER-positive, HER2-negative BC), although the bulk mixture

displayed comparable or superior predictive power than each of these two individual immune

cells, their ensemble markedly boosted the performance, supporting the possibility that there

may be prominent CCIs between these cell types that may be associated with the likelihood of

response (**Supplementary Figure 5C**). Among the three-cell-ensemble models, the combination

of endothelial, myeloid, and plasmablasts retains its highest performance, further underscoring

the critical roles of the stromal − immune interactions. Additionally, for BrighTNess, the individual

immune cell types showed notably strong predictive abilities, with less pronounced CCI effects

(**Supplementary Figure 5E**, **Figure 4D**). Collectively, these findings corroborate the immunogenic

effects mediated by chemotherapy, suggestive of *immunogenic cell death* [73] as suggested by

Sammut et al. [11].

Figure 3: The prominent multi-cell-type ensembles mediating chemotherapy response in breast cancer tumor microenvironment.

A-B. Validation performance with ARTemis + PBCP (n = 55) for the five most prominent two-cell-ensembles (**A**) and three-cell-ensembles (**B**) along with the bulk. AUC and AP stand for the area under the receiver operating characteristics curve and average precision (equivalent to the area under the precision-recall curve), respectively. Ensembles are ranked by their AUC values in a descending order.

C-D. Validation performance with BrighTNess (n = 106) for the five most prominent two-cell-ensembles (**C**) and three-cell-ensembles (**D**) along with the bulk. Ensembles are ranked by their AUC values in a descending order.

E. The enriched Reactome pathways (FDR-adjusted p-value ≤ 0.2) across the prominent cell types. For each cell type, only the top 10 relevant pathways are displayed. NES and log$P$ stand for the normalized enrichment score from GSEA and log-scaled p-value, respectively.

Identifying key cell-cell interactions associated with chemotherapy response

To investigate the impact of the presence of interactions among multiple cell types in modulating chemotherapy response, we use DECODEMi to analyze cell-cell interactions (inferred using LIRICS [44], starting with 2,422 ligand – receptor (L-R) pairs from Ramilowski et al. [50]; see details in **METHODS**) that are predictive of patient response. As before, we conducted two analyses: (a) cross-validation on TransNEO (n = 94), and (b) external validation on ARTemis + PBCP (n = 55) and BrighTNess (n = 106). We found that CCIs encompassing only the four most prominent cell types in the TME (*cancer epithelial*, *endothelial*, *myeloid*, and *plasmablasts*) exhibited a comparable classification power to all available CCIs across all three cohorts (one-tailed Wilcoxon rank-sum test p-value ≤ 0.05), further testifying to the influential roles of these specific cell types in patient response (**Figure 4A-D**, **Supplementary Figure 6A-D**). The CCIs involving these four cell types markedly improved the predictive power of CCI-based predictors in both TransNEO and ARTemis + PBCP (**Figure 4A-C**). However, the performance of CCIs was generally lower than the top predictive myeloid cells in BrighTNess cohort (**Figure 4D**). This discrepancy can again be attributed to cohort composition and treatment-specific variations. It is worth noting that this cohort

included only TNBC patients treated with paclitaxel and carboplatin, for which very few samples

were available in training (**Supplementary Figure 1B-C**). These results reinforced that leveraging

information from multiple cell types can notably improve clinical response prediction and the

inter-cellular crosstalk may involve mediating treatment response in the TME. To further assess

the generalizability of DECODEMi beyond bulk cohorts, we assessed its effectiveness on an

independent single-cell (SC) TNBC cohort without additional training. To overcome the limitation

of small sample size (n = 6), we constructed a SC-TNBC cohort of 200 'pseudopatients' treated

with chemotherapy (R:NR = 100:100) by downsampling 28,209 cells across the six TNBC patients

who received paclitaxel in Zhang et al.'s study [48]. We focused on the CCIs among the three cell

types (encompassing B-cells, myeloid, and T-cells) present in both SC and deconvolved bulk data,

and identified the 170 top predictive CCIs (covering 134 L-R pairs) in bulk TransNEO cohort. The

prediction scores for the SC-TNBC cohort were computed by counting the number of activated

CCIs within each pseudopatient (as inferred by Kumar et al. [103]; see details in **METHODS**). Our

analysis demonstrated the significant and robust predictive capabilities of these top CCIs (one-

tailed Wilcoxon rank-sum test p-value ≤ 0.001, **Figure 4G-H**), affirming DECODEMi's ability to

identify the CCIs predictive of patient response.

By computing feature importance from DECODEMi, we identified the most predictive cell-

to-cell communications across the ARTemis + PBCP and BrighTNess cohorts, finding a significant

overlap in top CCIs (**Figure 4E-F**). Remarkably, the interaction between '*GDF9*' (growth

differentiation factor 9) and '*BMPR1B*' (bone morphogenetic protein receptor 1B), both members

of the TGF-β superfamily (the corresponding protein-protein interaction (PPI) network, obtained

from STRING [100], is provided in **Supplementary Figure 6E**) with roles in reproductive biology

and signal transduction [74][75][76][77], emerged as a key CCI with the same directionality across

multiple cell types. These genes have been implicated in BC prognosis and therapeutic outcomes

[78][79][80], with *BMPR1B* exhibiting dual roles in chemotherapy response [81][82][83]. Another

top CCI involved the renowned *GDNF − RET* signaling pathway interactions between '*NRTN*'

(neurturin), or its paralog '*ARTN*' (artemin), and '*GFRA1*' (GDNF family receptor alpha 1) in

multiple cell types (**Figure 4E-F**; see PPI network in **Supplementary Figure 6F**), which are all

members of the glial cell line-derived neurotrophic factor (GDNF) family within the TGF-β

superfamily with roles in neuronal survival, growth, and differentiation [84]. The GDNF family

exhibits diverse roles in cancer [85][86], including an association with therapy resistance in ER-

positive BC, highlighting *GFRA1* as a potential target [87][88][89][90]. Additionally, the well-

studied chemokine interaction between '*CX3CL1*' (C-X3-C chemokine ligand 1) and '*CX3CR1*' (C-

X3-C chemokine receptor 1) [91] is prominent in BrighTNess (**Figure 4F**; see PPI network in

Supplementary Figure 6G), which is known to be involved in progression and metastasis across

multiple cancers including BC, presenting a promising pan-cancer target [92][93][94][95][96][97].

Overall, DECODEMi application results in: (a) a simpler predictor leveraging CCIs that

improves patient stratification compared to bulk transcriptomics, and (b) the identification of cell-

to-cell communications predictive of treatment response, whose activation is associated with

pro-cancerous roles, thus offering potential novel therapeutic targets.

Figure 4: The prominent cell-cell interactions (CCIs) mediating chemotherapy response in breast cancer tumor microenvironment.

A. Prediction scores for DECODEMi in cross-validation with TransNEO (*n* = 94) and in external validation with ARTemis + PBCP (*n* = 55) and BrighTNess (*n* = 106). R and NR stand for responders

and non-responders, respectively. The differences between the prediction scores were computed

by using a one-tailed Wilcoxon rank-sum test.

B. Model performance in cross-validation with TransNEO for CCI-based, the top cell-type-specific,

and bulk predictors. AUC and AP stand for the area under the receiver operating characteristics

curve and average precision (equivalent to the area under precision-recall curve), respectively.

C-D. Model performance in external validation with ARTemis + PBCP (**C**) and BrighTNess (**D**) for

CCI-based, the top cell-type-specific, and bulk predictors.

E-F. The 10 most predictive CCIs in external validation with ARTemis + PBCP (**E**) and BrighTNess

(**F**). Each CCI is displayed as a quadruplet where the first pair containing the ligand and receptor

cell types (separated by '–') is separated by '::' from the second pair containing the corresponding

ligand and receptor genes (separated by '–'). Mean decrease in Gini impurity is the built-in feature

importance measure in random forest where a higher value indicates a higher importance and

vice versa [98]. Feature directionalities were computed using Fisher's enrichment analysis.

G-H. Prediction scores (**G**) and AUC value (**H**) for chemotherapy response prediction in external

validation with the SC-TNBC cohort (n = 200, sourced from Zhang et al. [48]) using the top 170

response-relevant CCIs (among B-cells, myeloid, and T-cells) identified in bulk by DECODEMi.

DECODEM identifies responders to chemotherapy and anti-PD-L1 therapy from

single-cell transcriptomics

To further assess the generalizability of DECODEM, we evaluated the predictive capabilities of the

cell-type-specific models with single-cell transcriptomics. We analyzed a SC cohort containing the

expression of B-cells, myeloid, and T-cells for 15 patients with triple negative breast cancer from

Zhang et al., who were either treated with chemotherapy (paclitaxel) alone or combined with the

anti-PD-L1 agent, atezolizumab [48]. We first stratified the patient cohort into those who received

chemotherapy alone (n = 6) and those who underwent chemotherapy + immunotherapy (n = 9).

For each subset, we applied our established DECODEM models for these three cell types which

were trained with deconvolved TransNEO data alongside a baseline 'pseudobulk' model

(analogous to bulk in previous scenarios; see details in **METHODS**). Our findings revealed that the

expression of B-cells exhibited the highest predictive potential within this cohort (**Figure 5A-B**,

Supplementary Figure 7A-D). Notably, this B-cells-specific predictor from DECODEM effectively

distinguishes the responders from non-responders for both intervention types (one-tailed

Wilcoxon rank-sum test *p*-value ≤ 0.05, **Figure 5A-B**). These findings further attest to the

generalizability and robustness of DECODEM, highlighting its potential in capturing treatment-

invariant properties of the BC TME to predict response to the combination of chemotherapy and

immunotherapy through SC transcriptomics.

DECODEM effectively stratifies TCGA patients' survival

We finally explored whether DECODEM can stratify the survival of 705 breast cancer patients in

The Cancer Genome Atlas (TCGA-BRCA) from their deconvolved expression, obtained from Wang

et al. [44]. Focusing on the expression of the prominent cell type of cancer epithelial (and

endothelial) cells, DECODEM effectively stratified these patients into low-risk, likely responders

(scores > median score, *n* = 353) and high-risk, unlikely responders (scores ≤ median score, *n* =

352), as displayed by their overall survival and progression-free interval (Log-rank test *p*-value ≤

0.05, **Figure 5C-D**, **Supplementary Figure 7E-F**). Remarkably, this stratification was achieved

regardless of treatment regimen, attesting that DECODEM captured the treatment-invariant

properties of the BC TME that are predictive of patient survival.

Figure 5: Generalizability of DECODEM to single-cell (SC) transcriptomics and for TCGA patient survival stratification.

A-B. Prediction scores (**A**) and the area under the receiver operating characteristics curve (AUC) values (**B**) in external validation with SC expression of TNBC patients (SC-TNBC) treated with chemotherapy alone (n = 6) or combined with immunotherapy (n = 9) from Zhang et al. [48]. R and NR stand for responders and non-responders, respectively. The differences between the prediction scores were computed by using a one-tailed Wilcoxon rank-sum test.

C-D. Kaplan-Meier plots depicting the overall survival (OS; **C**) and progression-free interval (PFI; **D**) of 705 BC patients from TCGA (TCGA-BRCA). Patients were stratified into low-risk, likely responder (n = 353) and high-risk, unlikely responder (n = 352) groups by their median DECODEM score computed using the expression of cancer epithelial cells, whereby the low-risk and high-risk groups comprised individuals with scores above and below the median, respectively. The differences between the two curves were computed by using a Log-rank test.

DISCUSSION

Our study introduces DECODEM, a transcriptomics-based modeling framework employing CODEFACS, a cutting-edge deconvolution tool, with a multi-stage machine learning pipeline to systematically assess the cell-type-specific contributions to predicting clinical response to treatment. As a representative scenario, we apply it to build predictors of patient response to neoadjuvant chemotherapy in breast cancer. We further present DECODEMi, a companion tool that identifies the key cell-cell interactions associated with clinical response. A recent study by Kester et al. investigated the association of cellular compositions of the BC TME with patient survival [43], but to our knowledge, the current study is the first-of-its-kind to analyze the cell-type-specific transcriptomics profiles in the BC TME to quantify their respective contributions to treatment response prediction.

We applied DECODEM to investigate the predictive power of the expression of various cell types in the BC TME to patient response to NAC. Our findings highlight the prominence of immune cells (*myeloid*, *plasmablasts*, *B-cells*) in chemotherapy response, consistent with Kester et al. [43], elucidating the complementary roles of diverse cell types that allows for the incorporation of top predictive cell types to build accurate clinical response predictors, and showcase DECODEMi's ability to identify key CCIs impacting response across HER2-negative BC subtypes. Furthermore, we illustrated DECODEM's generalizability to single-cell transcriptomics where it captures treatment-invariant properties of the TME, enabling accurate classification of responders for chemotherapy and immunotherapy, and its application for patient survival stratification in TCGA.

Immune cells have a clear association with prognosis in BC, both without chemotherapy and after chemotherapy in the early-stage setting [104][30][31]. The International TILs Working Group was established for the express purpose of standardizing incorporation of tumor infiltrating lymphocyte pathological review for clinical use, and established an online prognosis calculator for patients with TNBC (https://www.tilsinbreastcancer.org/prognosis-tool/). Plasma cells and naïve B-cells have been associated with better pCR rates in ER-positive tumors, whereas regulatory T-cells have been associated with better pCR rates in TNBC [105]. In our analysis, immune cells such

as plasmablasts and myeloid cells exhibit involvement in signal transduction, particularly through

VEGF signaling and *NOTCH* signaling pathways, which are associated with tumor-associated

angiogenesis, enrichment of BC stem cells and epithelial-to-mesenchymal transition, leading to

poor prognosis and therapy resistance [60][61][64][66]. Stromal cells such as endothelial cells

show enrichments in development- and immune-related pathways including *L1CAM* interactions

and antigen processing & presentation, with *L1CAM*'s oncogenic role and defects in antigen

processing leading to poor outcomes and therapy resistance [67][68][69][70][71]. Overall, our

results paired with the existing clinical evidence reinforce the notion that the presence of an

active immune infiltration within the TME prior to treatment, as indicated by the prominent

contributions of immune cells, leads to favorable response to chemotherapy, as suggested by

Sammut et al. [11].

It is also important to note that DECODEM and DECODEMi offer to extend upon the

existing application of gene expression in the early-stage breast cancer setting. Currently, gene

expression-based tests such as Oncotype DX and MammaPrint are used for prediction and

prognostication in ER-positive BC patients, reporting if a subset should receive chemotherapy due

to worse predicted outcomes [106]. These tests use quantitative RT-PCR or cDNA microarray to

assess a known subset of genes, and largely apply only to a subset of early-stage ER-positive BC

patients. DECODEM and DECODEMi provide complimentary context to these tools, expanding the

applicable subtypes for chemotherapy response prediction as well as greater biological context.

Although we presented a comprehensive framework to assess the TME in a quantitative

manner, certain limitations should be acknowledged and addressed in future studies. *First*, our

predictors rely on data-driven algorithms involving deconvolution and ML, necessitating a large

sample size to attain statistical robustness and avoid overfitting. Accordingly, the available

datasets are limited in size, limiting our ability to perform robust subtype-specific analyses.

Second, any limitations inherent to CODEFACS, such as discrepancies in RNA-seq processing and

reliance on appropriate molecular signatures [44], will affect DECODEM in the downstream

analysis. The same goes for DECODEMi and its dependence on LIRICS using an appropriate ligand

– receptor database [43]. *Third*, our deconvolution analysis could not be performed separately

for the CD4$^+$ and CD8$^+$ T-cell subsets. Alternatively, we provided an enrichment analysis assessing

the predictive powers of CD4$^+$ and CD8$^+$ T-cells which corroborated our findings from using the

overall T-cells.

Our work focused on examining the association of the BC TME, specifically in HER2-

negative subtypes, to clinical response to NAC. However, both DECODEM and DECODEMi can be

applied to investigate diverse treatment regimens in different cancer indications. Although both

TransNEO and ARTemis + PBCP cohorts include different chemotherapy combinations, limited

sample size prevented us from exploring further. Therefore, an intriguing direction could be to

investigate the distinct role of the TME in different chemotherapy families with different

mechanisms of actions (MOAs) *e.g.*, taxanes, antimetabolites, platinum agents and so on. With

DECODEMi, one could further pinpoint the family-specific significant cell-cell interactions to

understand the underlying MOA and uncover novel CCIs with promising therapeutic implications.

Additionally, expanding the investigation beyond the scope of cancer and examining the impact

of the microenvironment in various indications such as aging-related disorders and autoimmune

diseases could provide further valuable insights.

In summary, our study introduces two data-driven computational frameworks that enable

systematic investigation of the roles of the microenvironment in various processes, including

treatment response. These frameworks can utilize bulk transcriptomics from healthy and

diseased conditions when single-cell information is not easily accessible, providing a valuable

approach to study the TME in a comprehensive manner. Their application in breast cancer

uncovers the considerable predictive power of diverse cell types within the TME and highlights

key stromal – immune cell-cell interactions that are strongly predictive of patient response.

METHODS

Overview of the datasets

We analyzed three bulk transcriptomics datasets, encompassing a total of 255 breast cancer patients who received neoadjuvant chemotherapy (**Supplementary Figure 1**). The first dataset, recently published by Sammut et al., was the TransNEO cohort which included 94 HER2-negative breast tumors patients undergoing various NAC regimens [11]. Notably, Sammut et al. presented the complete TransNEO cohort, covering 168 patients with early and locally advanced BC from all major subtypes, and providing DNA-seq, RNA-seq, digital pathology, and treatment information for various subsets of patients [11]. The second dataset was the ARTemis + PBCP cohort, validated externally by Sammut et al., combining the control arm of the ARTemis trial (n = 38) received the combination therapy of FEC-T [46] and the HER2-negative cases from the Personalised Breast Cancer Programme (PBCP; n = 17) undergoing various NAC regimens [11] (**Supplementary Figure 1B-C**). The third dataset, the BrighTNess cohort, included 106 patients from arm B of the BrighTNess trial, a randomized, double-blind, placebo-controlled three-arm clinical study for veliparib in stage II/III TNBC [47]. Arm B included only patients who received the NAC combination of paclitaxel, carboplatin, and a placebo for veliparib (**Supplementary Figure 1B-C**). In all three cohorts, patient response was assessed at surgery using the residual cancer burden (RCB) index where patients achieving pathological complete response (pCR; RCB = 0) were considered as responders (R) and patients with residual diseases (RCB-I, II, III) as non-responders (NR) with different R:NR ratios (**Supplementary Figure 1A**).

Furthermore, we analyzed a recent single-cell cohort presented by Zhang et al. comprising 22 patients with advanced triple negative breast cancer. Half of the patients received paclitaxel alone, while the other half received paclitaxel combined with the anti-PD-L1 agent, atezolizumab [48]. This study included 78 tumor biopsies and blood samples collected at three time-points, including baseline. Zhang et al. provided the single-cell expression for four immune cell types (B-cells, T-cells, myeloid, and innate lymphoid cells) and patient response assessed using the Response Evaluation Criteria in Solid Tumors (RECIST) criteria [48]. We categorized the patients achieving complete or partial response as responders (R) and those with stable or progressive

diseases as non-responders (NR). Our analysis focused solely on pre-treatment tissue samples,

resulting in two treatment groups: (a) chemotherapy (n = 6, R:NR = 3:3) and (b) chemotherapy +

immunotherapy (n = 9, R:NR = 4:5). To align with the output of CODEFACS, which provides the

mean cell-type-specific expression profile per patient [44], we harmonized the cell-type-specific

SC expression by averaging across all corresponding cells for each patient, while filtering for low-

variance genes (threshold = 0.8). Additionally, we calculated the 'pseudobulk' expression as the

mean expression across all cells encompassing the three cell types for each patient (post low-

variance filtering) to generate a baseline expression profile similar to the bulk mixture.

Deconvolution with CODEFACS

The first step of DECODEM involves employing CODEFACS, a cellular deconvolution tool recently

developed in our lab [44]. CODEFACS characterizes the TME by reconstructing the cell-type-

specific expression in each sample from the input bulk expression, utilizing the corresponding cell

type abundance or a cell-type-specific signature profile as another input. The output includes the

cell-type-specific expression profiles, the corresponding cell type abundances, and a confidence

score matrix that quantifies the confidence level (as a value between 0 and 1) of the estimated

expression values for each gene in each cell type across all samples.

We derived a cell type signature matrix of 1,400 genes representing nine cell types from

a single-cell RNA-seq (scRNA-seq) cohort from Wu et al. [45] using an in-house signature

derivation tool based on the CIBERSORTx algorithm [101]. We then deconvolved the three bulk

expression profiles with this signature into cell-type-specific expression profiles, encompassing

the nine cell types: *B-cells*, *Cancer-associated Fibroblasts (CAFs)*, *Cancer Epithelial (malignant)*,

Endothelial, *Myeloid*, *Normal Epithelial*, *Plasmablasts*, *Perivascular-like Cells (PVL)*, and *T-cells*.

Extraction of cell-cell interactions with LIRICS

592   We extended DECODEM to DECODEMi by incorporating multi-cell-type information through cell-

593   cell interactions. To this end, DECODEMi employs LIRICS [44] on the output of CODEFACS to infer

594   the activity of the corresponding CCIs. LIRICS applies a two-step approach to infer the cell-type-

specific ligand – receptor (L-R) pairs that are likely to be 'active' in each sample from deconvolved expression. The first step is to query a database of 49,397 plausible L-R interactions between each pair of cell types in BC TME (see below for details). The second step infers an L-R interaction to be active (indicated by '1') in a pair of cell types in a given sample if the expressions of both ligand and receptor genes exceed their median cell-type-specific expression values or inactive (indicated by '0') otherwise. We used this binary CCI activity matrix output from LIRICS to assess the predictive potential of cell-to-cell communications.

Curation of cell-type-specific ligand – receptor database for BC TME from single-cell transcriptomics

To infer the cell-cell interaction activity using LIRICS, we applied a two-step approach to construct a database of cell-type-specific ligand – receptor interactions (LRIs) to characterize all cell-to-cell communications that are likely to occur in the TME of BC patients. We first obtained a list of 2,422 potential L-R pairs, as annotated by Ramilowski et al. [50]. We next evaluated the potential of the L-R pairs occurring between the nine cell-types-of-interest using scRNA-seq data from Wu et al [45]. To discern the activity of the curated L-R list, we pooled together all 130,246 cells for the nine cell types from Wu et al. and adopted an empirical p-value based CCI inference approach following Kumar et al. [103], where an LRI with p-value ≤ 0.05 is categorized as 'likely' to occur in the TME of BC patients and 'unlikely' otherwise. We constructed a database of 49,397 cell-type-specific L-R pairs (*i.e.*, CCI quadruplets) that are likely to occur in BC TME. This database composed the initial feature space for LIRICS to yield different numbers of CCIs across three bulk cohorts: TransNEO = 46,755, ARTemis + PBCP = 43,887, and BrighTNess = 38,285.

Machine learning pipeline with deconvolved expression and CCIs

In both DECODEM and DECODEMi, the second step involves building ML predictors of clinical response using cell-type-specific expression profiles and CCIs, respectively. For DECODEM, we built nine cell-type-specific predictors to classify responders and non-responders following treatment using a four-stage ML pipeline (**Figure 1**): (a) prefiltering to remove genes with CODEFACS confidence score < 0.99, (b) low-variance filtering to remove genes with variance < 0.1,

(c) feature selection to select the top 'm' genes maximizing performance, where $2 \leq m \leq 25$ with

genes ranked by their ANOVA F-scores for association with response followed by standardization

to scale each gene to zero mean and unit variance, and (d) classification with an unweighted

ensemble classifier comprised of regularized logistic regression, random forest, support vector

machine, and XGBoost (extreme gradient boosting) to predict the probabilities of achieving pCR.

Here, the upper limit on the feature size is chosen to avoid model overfitting and longer runtimes

(and evaluated to provide similar performance across different feature sizes in cross-validation).

Furthermore, each prediction vector was rescaled between 0 and 1 to improve generalization for

external validation. We trained this ML pipeline with three-fold cross-validation (CV) for

hyperparameter tuning to maximize the AUC for each classifier separately. To avoid overfitting,

we repeated the training procedure five times with five different CV splits and took the mean of

the five prediction vectors as our final prediction. We analyzed the three bulk expression cohorts

with this pipeline in two scenarios: (a) cross-validation with TransNEO (n = 94) where we applied

an additional five-fold CV to estimate the performance, and (b) external validation on ARTemis +

PBCP (n = 55) and BrighTNess (n = 106) using predictors trained on TransNEO. We further built a

baseline predictor using bulk expression in ML pipeline directly and identified the 'prominent' cell

types in CV that outperformed the bulk, subsequently validated in external validation.

To accommodate the sparse nature of SC transcriptomics for external validation with

Zhang et al. [48], we slightly modified the pipeline to enable modeling for all available genes post

filtering in step (c) *i.e.*, $2 \leq m \leq$ 'all'. Additionally, for CCI-based analysis with DECODEMi, we further

updated the low-variance cut-off in step (b) to 0.08 and the classifier in step (d) to a random forest

that is better suited for the binary nature of the CCI matrix and includes a built-in measure of

feature importance, mean decrease in Gini impurity (similar performance to the permutation-

based approaches) [98]. The corresponding CCI directionalities were computed using a two-tailed

Fisher's test (followed by false discovery rate (FDR) adjustment for multiple hypothesis correction

i.e., FDR-adjusted *p*-value $\leq$ 0.05), where an odds ratio (OR) > 1 denotes an enrichment in

responders and OR < 1 indicates enrichment in non-responders. Moreover, to filter CCIs for a

subset of cell types, we only include the interactions involving the confident L-R genes (CODEFACS

confidence level ≥ 0.99) in the corresponding cell type.

CD4$^+$ / CD8$^+$ T-cells enrichment analysis

To analyze the contribution of the individual T-cell subtypes to chemotherapy response, we

performed a gene set variation analysis (GSVA) to summarize pathway activity per sample for

each T-cell subtype. We employed two single-cell expression signatures derived from Wu et al.

[45], collected *via* the Tumor Immune Single Cell Hub 2 (TISCH2) [102]. These signatures consisted

of 207 and 424 differentially expressed genes (log-fold-change ≥ 0.5) for CD4$^+$ and CD8$^+$ T-cells,

respectively (with an overlap of 151 genes). Subsequently, we evaluated the association of the

GSVA enrichment scores (rescaled between 0 and 1) with clinical response across the three bulk

cohorts through a one-tailed Wilcoxon rank-sum test (FDR-adjusted *p*-value ≤ 0.05), and also

assessed the AUC and AP values independently.

Cell-type-specific enrichment analysis

We conducted gene set enrichment analysis (GSEA) to investigate the association of each cell type

with different biological processes. For each cell type, we first ranked the 17,680 protein-coding

genes available in TransNEO by their association with positive outcome (*i.e.*, pCR), measured by

the *p*-values from a one-tailed Wilcoxon rank-sum test, adjusted by CODEFACS confidence scores.

We next performed a separate GSEA analysis using the Reactome pathways [52] for each

prominent cell type (except CAFs). This resulted in six different sets of enriched pathways (FDR-

adjusted *p*-value ≤ 0.2) across six prominent cell types. From these sets, we reported the top 10

relevant pathways for each cell type.

Single-cell validation of top cell-cell interactions

We validated the most predictive cell-cell interactions from DECODEMi using a single-cell cohort

from Zhang et al. that includes six TNBC patients treated with chemotherapy [48]. We devised a

three-step validation procedure: *First,* for consistency, we retrained DECODEMi using only the

interactions among cell types available in both bulk and SC (namely *B-cells*, *myeloid*, and *T-cells*)

in TransNEO. This resulted in the extraction of the top predictive CCIs (m = 170) involving a total

of 134 unique L-R pairs. *Second*, to enhance the statistical power, we pooled the 28,209 cells

encompassing these three cell types from the six chemotherapy patients in Zhang et al., and

randomly downsampled 30% of the cells 100 times within each treatment group for each cell

type. This procedure generated a profile of 200 'pseudopatients' (R:NR = 100:100) with empirical

p-values encompassing the significance levels of the 804 possible CCIs involving the 134 L-R pairs

in three cell types (= 134 × 6) *i.e.*, the 'SC-TNBC cohort'. To identify the activation status of the

170 CCIs-of-interest in this cohort, we again adopted the CCI inference approach from Kumar et

al. [103], where a CCI is denoted as 'active' if *p*-value ≤ 0.05 and 'inactive' otherwise. *Third*, for

each pseudopatient, we predicted the chemotherapy response as the sum of the active CCIs,

weighted by their directionalities (*i.e.*, +1 or -1 when OR > 1 or < 1, FDR-adjusted *p*-value ≤ 0.05;

and 0 otherwise), followed by a rescaling between 0 and 1 across all pesudopatients. We used

these prediction scores to evaluate the predictive performance of these top CCIs learned from

bulk on the SC-TNBC cohort through using a one-tailed Wilcoxon rank-sum test (FDR-adjusted *p*-

value ≤ 0.05) and receiver operating characteristics (ROC) analysis.

DATA AVAILABILITY

For TransNEO cohort, RNA-seq counts and clinical data were acquired from Sammut et al.

(https://doi.org/10.1038/s41586-021-04278-5). RNA-seq counts were converted to TPM by using

Ensembl gene annotations (GRCh37.87, http://ftp.ensembl.org/pub/grch37/release-

87/gtf/homo_sapiens/). For ARTemis + PBCP cohort, clinical data were collected from Sammut et

al. (https://doi.org/10.1038/s41586-021-04278-5) and RNA-seq TPM values were obtained

directly from the authors. For BrighTNess cohort, RNA-seq FPKM values and clinical data were

acquired from GEO (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE164458). The

FPKM values were converted into TPM. For Zhang et al. cohort, scRNA-seq counts and clinical

data were obtained from Zhang et al. (https://doi.org/10.1016/j.ccell.2021.09.010). For the

single-cell signature for deconvolution, scRNA-seq counts and cell annotations were collected

from Wu et al. (https://doi.org/10.1038/s41588-021-00911-1). The ligand-receptor list for LIRICS

were obtained from Ramilowski et al. via GitHub (https://github.com/LewisLabUCSD/Ligand-

Receptor-Pairs). The deconvolved expression and cell fraction data using CODEFACS on TCGA-BRCA patients were collected from Wang et al. (https://doi.org/10.1158%2F2159-8290.CD-21-0887). The corresponding TCGA-BRCA patients' overall survival and progression-free interval (synonymous to progression-free survival) data were obtained from the UCSC Xena browser (https://xenabrowser.net). All data used in this study are either publicly available or available upon request to the original authors publishing the data. All deconvolved expression data generated in this study will be made publicly available upon publication.

CODE AVAILABILITY

All relevant codes for reproducing the figures in this study will be made publicly available upon publication at https://github.com/ruppinlab/DECODEM/.

ACKNOWLEDGEMENTS

This research is supported in part by the Intramural Research Program of the National Institutes of Health, National Cancer Institute, Center for Cancer Research. This work has utilized the computational resources of the NIH HPC Biowulf cluster (http://hpc.nih.gov). The results presented are in part based on the data generated by The Cancer Genome Atlas Research network (https://www.cancer.gov/tcga/). We would additionally like to acknowledge Dr. E. Michael Gertz and other members of the Cancer Data Science Laboratory for their help with data collection and constructive feedback on this work. Figure 1 has been created with http://biorender.com/.

AUTHOR CONTRIBUTIONS

S.R.D, K.W., and E.R. conceived and designed the study. S.R.D., S.S., and K.W. performed the model development and validation analyses. S.R.D., K.W., S.S., P.S.R., B.W., Y.S., E.S., S-J.S., C.C., and E.R. acquired, analyzed, or interpreted the data. S.R.D., S.S., K.W., D.W., and B.W. worked on data visualization. E.R. and K.W. supervised the study. All authors critically revised the manuscript for important intelligent content.

COMPETING INTERESTS

738 **COMPETING INTERESTS**

739 E.R. is a co-founder of Medaware Ltd. (https://www.medaware.com/), Metabomed
740 (https://www.metabomed.com/), and Pangea Biomed (https://pangeamedicine.com/). He has
741 divested and serves as an unpaid scientific consultant to the latter company. The rest of the
742 authors declare no conflicts of interest.
743

REFERENCES

[1] Tsimberidou, A. M., Fountzilas, E., Nikanjam, M., & Kurzrock, R. (2020). Review of precision cancer medicine: Evolution of the treatment paradigm. Cancer treatment reviews, 86, 102019.

[2] Huang K, Xiao C, Glass LM, Critchlow CM,. Machine learning applications for therapeutic tasks with genomics data. Patterns 2021; 2(10):100328.

[3] Fisher, B., Brown, A., Mamounas, E., Wieand, S., Robidoux, A., Margolese, R. G., … & Dimitrov, N. V. (1997). Effect of preoperative chemotherapy on local-regional disease in women with operable breast cancer: findings from National Surgical Adjuvant Breast and Bowel Project B-18. Journal of clinical oncology, 15(7), 2483-2493.

[4] Bear, H. D., Anderson, S., Brown, A., Smith, R., Mamounas, E. P., Fisher, B., ... & Wolmark, N. (2003). The effect on tumor response of adding sequential preoperative docetaxel to preoperative doxorubicin and cyclophosphamide: preliminary results from National Surgical Adjuvant Breast and Bowel Project Protocol B-27. Journal of Clinical Oncology, 21(22), 4165-4174.

[5] Golshan, M., Loibl, S., Wong, S. M., Huober, J. B., O'Shaughnessy, J., Rugo, H. S., ... & Untch, M. (2020). Breast conservation after neoadjuvant chemotherapy for triple-negative breast cancer: surgical results from the BrighTNess randomized clinical trial. JAMA surgery, 155(3), e195410-e195410.

[6] Urueña, C., Lasso, P., Bernal-Estevez, D. et al. The breast cancer immune microenvironment is modified by neoadjuvant chemotherapy. Sci Rep 12, 7981 (2022). https://doi.org/10.1038/s41598-022-12108-5

[7] Ali, H. R., Dariush, A., Provenzano, E., Bardwell, H., Abraham, J. E., Iddawela, M., ... & Caldas, C. (2016). Computational pathology of pre-treatment biopsies identifies lymphocyte density as a predictor of response to neoadjuvant chemotherapy in breast cancer. Breast Cancer Research, 18(1), 1-11.

[8] Ali, H. R., Dariush, A., Thomas, J., Provenzano, E., Dunn, J., Hiller, L., ... & Caldas, C. (2017). Lymphocyte density determined by computational pathology validated as a predictor of

response to neoadjuvant chemotherapy in breast cancer: secondary analysis of the ARTemis

trial. Annals of Oncology, 28(8), 1832-1835.

[9] Dieci, M. V., Prat, A., Tagliafico, E., Paré, L., Ficarra, G., Bisagni, G., ... & Guarneri, V. (2016).

Integrated evaluation of PAM50 subtypes and immune modulation of pCR in HER2-positive

breast cancer patients treated with chemotherapy and HER2-targeted agents in the CherLOB

trial. Annals of Oncology, 27(10), 1867-1873.

[10] Griguolo, G., Serna, G., Pascual, T. et al. Immune microenvironment characterisation and

dynamics during anti-HER2-based neoadjuvant treatment in HER2-positive breast cancer. npj

Precis. Onc. 5, 23 (2021). HYPERLINK "https://doi.org/10.1038/s41698-021-00163-6.

https://doi.org/10.1038/s41698-021-00163-63434

[11] Sammut, SJ., Crispin-Ortuzar, M., Chin, Sf. et al. Multi-omic machine learning predictor of

breast cancer therapy response. Nature 601, 623–629 (2022).

https://doi.org/10.1038/s41586-021-04278-5

[12] Klemm F, Joyce JA. Microenvironmental regulation of therapeutic response in cancer. Trends

Cell Biol. 2015 Apr;25(4):198-213. doi: 10.1016/j.tcb.2014.11.006. Epub 2014 Dec 22. PMID:

25540894; PMCID: PMC5424264.

[13] Son B, Lee S, Youn H, Kim E, Kim W, Youn B. The role of tumor microenvironment in

therapeutic resistance. Oncotarget. 2017 Jan 17;8(3):3933-3945. doi:

10.18632/oncotarget.13907. PMID: 27965469; PMCID: PMC5354804.

[14] Wu, P., Gao, W., Su, M., Nice, E. C., Zhang, W., Lin, J., & Xie, N. (2021). Adaptive mechanisms

of tumor therapy resistance driven by tumor microenvironment. Frontiers in cell and

developmental biology, 9, 641469.

[15] Mehraj, U., Dar, A.H., Wani, N.A. et al. Tumor microenvironment promotes breast cancer

chemoresistance. Cancer Chemother Pharmacol 87, 147–158 (2021).

https://doi.org/10.1007/s00280-020-04222-w

[16] Ruffell, B., & Coussens, L. M. (2015). Macrophages and therapeutic resistance in cancer.

Cancer cell, 27(4), 462-472.

[17] Acharyya, S., Oskarsson, T., Vanharanta, S., Malladi, S., Kim, J., Morris, P. G., ... & Massagué,

J. (2012). A CXCL1 paracrine network links cancer chemoresistance and metastasis. Cell,

150(1), 165-178.

[18] Li, M., Quintana, A., Alberts, E., Hung, M. S., Boulat, V., Ripoll, M. M., & Grigoriadis, A. (2023).

B Cells in Breast Cancer Pathology. Cancers, 15(5), 1517.

[19] Sakaguchi, A., Horimoto, Y., Onagi, H. et al. Plasma cell infiltration and treatment effect in

breast cancer patients treated with neoadjuvant chemotherapy. Breast Cancer Res 23, 99

(2021). https://doi.org/10.1186/s13058-021-01477-w

[20] Yeong, J., Lim, J. C. T., Lee, B., Li, H., Chia, N., Ong, C. C. H., ... & Iqbal, J. (2018). High densities

of tumor-associated plasma cells predict improved prognosis in triple negative breast cancer.

Frontiers in immunology, 9, 1209.

[21] Bhinder, B., Gilvary, C., Madhukar, N. S., & Elemento, O. (2021). Artificial intelligence in

cancer research and precision medicine. Cancer Discovery, 11(4), 900-915.

[22] Singla, N., & Singla, S. (2020). Harnessing big data with machine learning in precision

oncology. Kidney cancer journal: official journal of the Kidney Cancer Association, 18(3), 83.

[23] Senft, D., Leiserson, M. D., Ruppin, E., & Ze'ev, A. R. (2017). Precision oncology: the road

ahead. *Trends in molecular medicine*, *23*(10), 874-898.

[24] Tsimberidou, A. M., Fountzilas, E., Bleris, L., & Kurzrock, R. (2020, September).

Transcriptomics and solid tumors: The next frontier in precision cancer medicine. In Seminars

in cancer biology. Academic Press.

[25] Siravegna, G., Marsoni, S., Siena, S., & Bardelli, A. (2017). Integrating liquid biopsies into the

management of cancer. Nature reviews Clinical oncology, 14(9), 531-548.

[26] Heitzer, E., Haque, I. S., Roberts, C. E., & Speicher, M. R. (2019). Current and future

perspectives of liquid biopsies in genomics-driven oncology. Nature Reviews Genetics, 20(2),

71-88.

[27] Sawabata, N. (2020). Circulating tumor cells: From the laboratory to the cancer clinic.

Cancers, 12(10), 3065.

[28] Beaubier, N., Bontrager, M., Huether, R., Igartua, C., Lau, D., Tell, R., ... & White, K. P. (2019).
Integrated genomic profiling expands clinical options for patients with cancer. *Nature*
biotechnology, *37*(11), 1351-1360.

[29] Hayashi, H., Takiguchi, Y., Minami, H., Akiyoshi, K., Segawa, Y., Ueda, H., ... & Nakagawa, K.
(2020). Site-specific and targeted therapy based on molecular profiling by next-generation
sequencing for cancer of unknown primary site: a nonrandomized phase 2 clinical trial. JAMA
oncology, 6(12), 1931-1938.

[30] Vaske, O. M., Bjork, I., Salama, S. R., Beale, H., Shah, A. T., Sanders, L., ... & Haussler, D. (2019).
Comparative tumor RNA sequencing analysis for difficult-to-treat pediatric and young adult
patients with cancer. *JAMA network open*, *2*(10), e1913968-e1913968.

[31] Lee, J. S., Nair, N. U., Dinstag, G., Chapman, L., Chung, Y., Wang, K., ... & Ruppin, E. (2021).
Synthetic lethality-mediated precision oncology via the tumor transcriptome. Cell, 184(9),
2487-2502.

[32] Sinha, S., Dhruba, S. R., Wu, W., Kerr, D. L., Stroganov, O. V., Grishagin, I., ... & Ruppin, E.
(2022). Predicting patient treatment response and resistance via single-cell transcriptomics
of their tumors. bioRxiv.

[33] Dinstag, G., Shulman, E. D., Elis, E., Ben-Zvi, D. S., Tirosh, O., Maimon, E., ... & Aharonov, R.
(2022). Clinically oriented prediction of patient response to targeted and immunotherapies
from the tumor transcriptome. bioRxiv.

[34] Fu, Y., Jung, A. W., Torne, R. V., Gonzalez, S., Vöhringer, H., Shmatko, A., ... & Gerstung, M.
(2020). Pan-cancer computational histopathology reveals mutations, tumor composition and
prognosis. Nature Cancer, 1(8), 800-810.

[35] Noorbakhsh, J., Farahmand, S., Namburi, S., Caruana, D., Rimm, D., Soltanieh-ha, M., ... &
Chuang, J. H. (2020). Deep learning-based cross-classifications reveal conserved spatial
behaviors within tumor histological images. Nature communications, 11(1), 1-14.

[36] Yu, K. H., Wang, F., Berry, G. J., Re, C., Altman, R. B., Snyder, M., & Kohane, I. S. (2020).
Classifying non-small cell lung cancer types and transcriptomic subtypes using convolutional
neural networks. Journal of the American Medical Informatics Association, 27(5), 757-769.

[37] Couture, H. D., Williams, L. A., Geradts, J., Nyante, S. J., Butler, E. N., Marron, J. S., ... &
Niethammer, M. (2018). Image analysis with deep learning to predict breast cancer grade,
ER status, histologic subtype, and intrinsic subtype. NPJ breast cancer, 4(1), 1-8.

[38] Qu, H., Zhou, M., Yan, Z., Wang, H., Rustgi, V. K., Zhang, S., ... & Metaxas, D. N. (2021). Genetic
mutation and biological pathway prediction based on whole slide images in breast carcinoma
using deep learning. NPJ precision oncology, 5(1), 1-11.

[39] Hoang, D. T., Dinstag, G., Hermida, L. C., Ben-Zvi, D. S., Elis, E., Caley, K., ... & Ruppin, E. (2022).
Synthetic lethality-based prediction of cancer treatment response from histopathology
images. bioRxiv.

[40] Tanioka, M., Fan, C., Parker, J. S., Hoadley, K. A., Hu, Z., Li, Y., ... & Perou, C. M. (2018).
Integrated Analysis of RNA and DNA from the Phase III Trial CALGB 40601 Identifies
Predictors of Response to Trastuzumab-Based Neoadjuvant Chemotherapy in HER2-Positive
Breast CancerIntegrated Response Models in HER2-Positive Breast Cancer. Clinical Cancer
Research, 24(21), 5292-5304.

[41] Wong, M., Mayoh, C., Lau, L., Khuong-Quang, D. A., Pinese, M., Kumar, A., ... & Cowley, M. J.
(2020). Whole genome, transcriptome and methylome profiling enhances actionable target
discovery in high-risk pediatric cancer. Nature medicine, 26(11), 1742-1753.

[42] Rodon, J., Soria, J. C., Berger, R., Miller, W. H., Rubin, E., Kugel, A., ... & Kurzrock, R. (2019).
Genomic and transcriptomic profiling expands precision cancer medicine: the WINTHER trial.
Nature medicine, 25(5), 751-758.

[43] Kester, L., Seinstra, D., van Rossum, A. G., Vennin, C., Hoogstraat, M., van der Velden, D., ...
& van Rheenen, J. (2022). Differential Survival and Therapy Benefit of Patients with Breast
Cancer Are Characterized by Distinct Epithelial and Immune Cell Microenvironments Tumor
Cellular Composition Predicts Benefit to Therapies. *Clinical Cancer Research*, OF1-OF12.

[44] Wang, K., Patkar, S., Lee, J.S., Gertz, E.M., Robinson, W., Schischlik, F., Crawford, D.R.,
Schäffer, A.A. and Ruppin, E. (2022). Deconvolving Clinically Relevant Cellular Immune Cross-
talk from Bulk Gene Expression Using CODEFACS and LIRICS Stratifies Patients with
Melanoma to Anti–PD-1 Therapy. Cancer discovery, 12(4), 1088-1105.

[45] Wu, S. Z., Al-Eryani, G., Roden, D. L., Junankar, S., Harvey, K., Andersson, A., ... & Swarbrick,
A. (2021). A single-cell and spatially resolved atlas of human breast cancers. *Nature*
genetics, *53*(9), 1334-1347.

[46] Earl, H. M., Hiller, L., Dunn, J. A., Blenkinsop, C., Grybowicz, L., Vallier, A. L., ... & Hayward, L.
(2015). Efficacy of neoadjuvant bevacizumab added to docetaxel followed by fluorouracil,
epirubicin, and cyclophosphamide, for women with HER2-negative early breast cancer
(ARTemis): an open-label, randomised, phase 3 trial. The lancet oncology, 16(6), 656-666.

[47] Loibl, S., O'Shaughnessy, J., Untch, M., Sikov, W. M., Rugo, H. S., McKee, M. D., ... & Geyer Jr,
C. E. (2018). Addition of the PARP inhibitor veliparib plus carboplatin or carboplatin alone to
standard neoadjuvant chemotherapy in triple-negative breast cancer (BrighTNess): a
randomised, phase 3 trial. The Lancet Oncology, 19(4), 497-509.

[48] Zhang, Y., Chen, H., Mo, H., Hu, X., Gao, R., Zhao, Y., ... & Liu, Z. (2021). Single-cell analyses
reveal key immune cell subsets associated with response to PD-L1 blockade in triple-negative
breast cancer. Cancer Cell, 39(12), 1578-1593.

[49] Korde, L. A., Somerfield, M. R., Carey, L. A., Crews, J. R., Denduluri, N., Hwang, E. S., ... &
Hershman, D. L. (2021). Neoadjuvant chemotherapy, endocrine therapy, and targeted
therapy for breast cancer: ASCO guideline. Journal of Clinical Oncology, 39(13), 1485-1505.

[50] Ramilowski, J., Goldberg, T., Harshbarger, J. *et al.* A draft network of ligand–receptor-
mediated multicellular signalling in human. *Nat Commun* **6**, 7866 (2015).
https://doi.org/10.1038/ncomms8866

[51] Green, D. R., Ferguson, T., Zitvogel, L., & Kroemer, G. (2009). Immunogenic and tolerogenic
cell death. Nature Reviews Immunology, 9(5), 353-363.

[52] Fabregat, A. et al. The Reactome Pathway Knowledgebase. Nucleic Acids Res. 46, D649–D655
(2018).

[53] Molinari, M. (2000). Cell cycle checkpoints and their inactivation in human cancer. Cell
proliferation, 33(5), 261-274.

[54] Bower, J.J., Vance, L.D., Psioda, M. et al. Patterns of cell cycle checkpoint deregulation
associated with intrinsic molecular subtypes of human breast cancer cells. npj Breast Cancer
3, 9 (2017). https://doi.org/10.1038/s41523-017-0009-7

[55] Zhou, K., Sun, Y., Dong, D. et al. EMP3 negatively modulates breast cancer cell DNA
replication, DNA damage repair, and stem-like properties. Cell Death Dis 12, 844 (2021).
https://doi.org/10.1038/s41419-021-04140-6

[56] Repo, H., Löyttyniemi, E., Kurki, S. et al. A prognostic model based on cell-cycle control
predicts outcome of breast cancer patients. BMC Cancer 20, 558 (2020).
https://doi.org/10.1186/s12885-020-07045-3

[57] Vihervuori, H., Korpinen, K., Autere, T. A., Repo, H., Talvinen, K., & Kronqvist, P. (2022).
Varying outcomes of triple-negative breast cancer in different age groups–prognostic value
of clinical features and proliferation. Breast Cancer Research and Treatment, 1-12.

[58] Thu, K. L., Soria-Bretones, I., Mak, T. W., & Cescon, D. W. (2018). Targeting the cell cycle in
breast cancer: towards the next phase. Cell Cycle, 17(15), 1871-1885.

[59] Zhang, J., Chan, D. W., & Lin, S. Y. (2022). Exploiting DNA Replication Stress as a Therapeutic
Strategy for Breast Cancer. Biomedicines, 10(11), 2775.

[60] Zhu, X., & Zhou, W. (2015). The emerging regulation of VEGFR-2 in triple-negative breast
cancer. Frontiers in Endocrinology, 6, 159.

[61] Ceci, C., Atzori, M. G., Lacal, P. M., & Graziani, G. (2020). Role of VEGFs/VEGFR-1 signaling
and its inhibition in modulating tumor invasion: Experimental evidence in different
metastatic cancer models. International journal of molecular sciences, 21(4), 1388.

[62] Zhang, Q., Lu, S., Li, T., Yu, L., Zhang, Y., Zeng, H., ... & Lin, Y. (2019). ACE2 inhibits breast
cancer angiogenesis via suppressing the VEGFa/VEGFR2/ERK pathway. Journal of
Experimental & Clinical Cancer Research, 38(1), 1-12.

[63] Edwards, A., & Brennan, K. (2021). Notch signalling in breast development and cancer.
Frontiers in Cell and Developmental Biology, 9, 692173.

[64] Zhou, B., Lin, W., Long, Y. *et al.* Notch signaling pathway: architecture, disease, and
therapeutics. *Sig Transduct Target Ther* **7**, 95 (2022). https://doi.org/10.1038/s41392-022-
00934-y.

[65] Miao, K., Lei, J.H., Valecha, M.V. et al. NOTCH1 activation compensates BRCA1 deficiency and
promotes triple-negative breast cancer formation. Nat Commun 11, 3256 (2020).
https://doi.org/10.1038/s41467-020-16936-9

[66] BeLow, M., & Osipo, C. (2020). Notch signaling in breast cancer: a role in drug resistance.
Cells, 9(10), 2204.

[67] Li, L., Wang, X., Hu, K. et al. ZNF133 is a potent suppressor in breast carcinogenesis through
dampening L1CAM, a driver for tumor progression. Oncogene (2023).
https://doi.org/10.1038/s41388-023-02731-5

[68] van der Maten, M., Reijnen, C., Pijnenborg, J. M., & Zegers, M. M. (2019). L1 cell adhesion
molecule in cancer, a systematic review on domain-specific functions. International journal
of molecular sciences, 20(17), 4180.

[69] Zhang, J., Yang, F., Ding, Y., Zhen, L., Han, X., Jiao, F., & Tang, J. (2015). Overexpression of L1
cell adhesion molecule correlates with aggressive tumor progression of patients with breast
cancer and promotes motility of breast cancer cells. International journal of clinical and
experimental pathology, 8(8), 9240.

[70] Bandola-Simon, J., & Roche, P. A. (2019). Dysfunction of antigen processing and presentation
by dendritic cells in cancer. Molecular immunology, 113, 31-37.

[71] Dhatchinamoorthy, K., Colbert, J. D., & Rock, K. L. (2021). Cancer immune evasion through
loss of MHC class I antigen presentation. Frontiers in immunology, 12, 636568.

[72] McDonnell, A. M., Lesterhuis, W. J., Khong, A., Nowak, A. K., Lake, R. A., Currie, A. J., &
Robinson, B. W. (2015). Tumor-infiltrating dendritic cells exhibit defective cross-presentation
of tumor antigens, but is reversed by chemotherapy. European journal of immunology, 45(1),
49-59.

[73] Galluzzi, L., Humeau, J., Buqué, A., Zitvogel, L., & Kroemer, G. (2020). Immunostimulation
with chemotherapy in the era of immune checkpoint inhibitors. Nature reviews Clinical
oncology, 17(12), 725-741.

[74] Li, J., Li, C., Liu, X. et al. GDF9 concentration in embryo culture medium is linked to human
embryo quality and viability. J Assist Reprod Genet 39, 117–125 (2022).
https://doi.org/10.1007/s10815-021-02368-x

[75] Stocker, W. A., Walton, K. L., Richani, D., Chan, K. L., Beilby, K. H., Finger, B. J., ... & Harrison,
C. A. (2020). A variant of human growth differentiation factor-9 that improves oocyte
developmental competence. Journal of Biological Chemistry, 295(23), 7981-7991.

[76] Gomez-Puerto, M. C., Iyengar, P. V., García de Vinuesa, A., Ten Dijke, P., & Sanchez-Duffhues,
G. (2019). Bone morphogenetic protein receptor signal transduction in human disease. The
journal of pathology, 247(1), 9-20.

[77] Feary, E. S., Juengel, J. L., Smith, P., French, M. C., O'Connell, A. R., Lawrence, S. B., ... &
McNatty, K. P. (2007). Patterns of expression of messenger RNAs encoding GDF9, BMP15,
TGFBR1, BMPR1B, and BMPR2 during follicular development and characterization of ovarian
follicular populations in ewes carrying the Woodlands FecX2W mutation. Biology of
reproduction, 77(6), 990-998.

[78] Hanavadi, S., Martin, T. A., Watkins, G., Mansel, R. E., & Jiang, W. G. (2007). The role of
growth differentiation factor-9 (GDF-9) and its analog, GDF-9b/BMP-15, in human breast
cancer. Annals of surgical oncology, 14, 2159-2166.

[79] Harrath, A. H., Jalouli, M., Oueslati, M. H., Farah, M. A., Feriani, A., Aldahmash, W., ... &
Alwasel, S. (2021). The flavonoid, kaempferol-3-O-apiofuranosyl-7-O-rhamnopyranosyl, as a
potential therapeutic agent for breast cancer with a promoting effect on ovarian function.
Phytotherapy Research, 35(11), 6170-6180.

[80] Bach, D. H., Park, H. J., & Lee, S. K. (2018). The dual role of bone morphogenetic proteins in
cancer. Molecular Therapy-Oncolytics, 8, 1-13.

[81] Bokobza, S. M., Ye, L., Kynaston, H. E., Mansel, R. E., & Jiang, W. G. (2009). Reduced
expression of BMPR-IB correlates with poor prognosis and increased proliferation of breast
cancer cells. Cancer genomics & proteomics, 6(2), 101-108.

[82] Zheng, Y., Jiang, X., Wang, M., Yang, S., Deng, Y., Li, Y., ... & Chen, L. (2022). BMPR1B
Polymorphisms (rs1434536 and rs1970801) are Associated With Breast Cancer Susceptibility
in Northwest Chinese Han Females: A Case-Control Study. Clinical Breast Cancer, 22(5), e641-
e646.

[83] Dai, K., Qin, F., Zhang, H., Liu, X., Guo, C., Zhang, M., ... & Ma, Y. (2016). Low expression of
BMPRIB indicates poor prognosis of breast cancer and is insensitive to taxane-anthracycline
chemotherapy. Oncotarget, 7(4), 4770.

[84] Jing, S., Wen, D., Yu, Y., Holst, P. L., Luo, Y., Fang, M., ... & Fox, G. M. (1996). GDNF–induced
activation of the ret protein tyrosine kinase is mediated by GDNFR-α, a novel receptor for
GDNF. Cell, 85(7), 1113-1124.

[85] Fielder, G. C., Yang, T. W. S., Razdan, M., Li, Y., Lu, J., Perry, J. K., ... & Liu, D. X. (2018). The
GDNF family: a role in cancer?. Neoplasia, 20(1), 99-117.

[86] Li, Q., Cao, Z., & Zhao, S. (2022). The Emerging Portrait of Glial Cell Line-derived Neurotrophic
Factor Family Receptor Alpha (GFRα) in Cancers. International Journal of Medical Sciences,
19(4), 659.

[87] Pecar, G., Liu, S., Hooda, J. et al. RET signaling in breast cancer therapeutic resistance and
metastasis. Breast Cancer Res 25, 26 (2023). https://doi.org/10.1186/s13058-023-01622-7

[88] Horibata, S., Rice, E. J., Mukai, C., Marks, B. A., Sams, K., Zheng, H., ... & Danko, C. G. (2018).
ER-positive breast cancer cells are poised for RET-mediated endocrine resistance. PLoS One,
13(4), e0194023.

[89] Bhakta, S., Crocker, L. M., Chen, Y., Hazen, M., Schutten, M. M., Li, D., ... & Junutula, J. R.
(2018). An Anti-GDNF Family Receptor Alpha 1 (GFRA1) Antibody–Drug Conjugate for the
Treatment of Hormone Receptor–Positive Breast Cancer Targeting Hormone-Positive Breast
Cancer with a GFRA1 ADC. Molecular cancer therapeutics, 17(3), 638-649.

[90] Kim, M., & Kim, D. J. (2018). GFRA1: a novel molecular target for the prevention of
osteosarcoma chemoresistance. International journal of molecular sciences, 19(4), 1078.

[91] Liu, H., Yang, Z., Lu, W., Chen, Z., Chen, L., Han, S., ... & Cai, Y. (2020). Chemokines and
chemokine receptors: A new strategy for breast cancer therapy. Cancer medicine, 9(11),
3786-3799.

[92] Liang, Y., Yi, L., Liu, P., Jiang, L., Wang, H., Hu, A., ... & Dong, J. (2018). CX3CL1 involves in
breast cancer metastasizing to the spine via the Src/FAK signaling pathway. Journal of
Cancer, 9(19), 3603.

[93] Tardáguila, M., Mira, E., García-Cabezas, M. A., Feijoo, A. M., Quintela-Fandino, M., Azcoitia,
I., ... & Manes, S. (2013). CX3CL1 Promotes Breast Cancer via Transactivation of the EGF
PathwayCX3CL1 Promotes Breast Cancer. Cancer research, 73(14), 4461-4473.

[94] Jamieson-Gladney, W.L., Zhang, Y., Fong, A.M. et al. The chemokine receptor CX3CR1 is

directly involved in the arrest of breast cancer cells to the skeleton. Breast Cancer Res 13,

R91 (2011). https://doi.org/10.1186/bcr3016

[95] Rivas-Fuentes, S., Salgado-Aguayo, A., Arratia-Quijada, J., & Gorocica-Rosete, P. (2021).

Regulation and biological functions of the CX3CL1-CX3CR1 axis and its relevance in solid

cancer: A mini-review. Journal of Cancer, 12(2), 571.

[96] Yue, Y., Zhang, Q., & Sun, Z. (2022). CX3CR1 acts as a protective biomarker in the tumor

microenvironment of colorectal Cancer. Frontiers in Immunology, 12, 5972.

[97] Ishida, Y., Kuninaka, Y., Yamamoto, Y., Nosaka, M., Kimura, A., Furukawa, F., ... & Kondo, T.

(2020). Pivotal involvement of the CX3CL1-CX3CR1 axis for the recruitment of M2 tumor-

associated macrophages in skin carcinogenesis. *Journal of Investigative*

Dermatology, *140*(10), 1951-1961.

[98] Breiman, L. (2001). Random forests. Machine learning, 45, 5-32.

[99] Michaud, M., Martins, I., Sukkurwala, A. Q., Adjemian, S., Ma, Y., Pellegatti, P., ... & Kroemer,

G. (2011). Autophagy-dependent anticancer immune responses induced by

chemotherapeutic agents in mice. Science, 334(6062), 1573-1577.

[100] Szklarczyk, D., Gable, A. L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., ... & Mering, C.

V. (2019). STRING v11: protein–protein association networks with increased coverage,

supporting functional discovery in genome-wide experimental datasets. Nucleic acids

research, 47(D1), D607-D613.

[101] Newman, A. M., Steen, C. B., Liu, C. L., Gentles, A. J., Chaudhuri, A. A., Scherer, F., ... &

Alizadeh, A. A. (2019). Determining cell type abundance and expression from bulk tissues

with digital cytometry. Nature biotechnology, 37(7), 773-782.

[102] Han, Y., Wang, Y., Dong, X., Sun, D., Liu, Z., Yue, J., ... & Wang, C. (2023). TISCH2: expanded

datasets and new tools for single-cell transcriptome analyses of the tumor

microenvironment. Nucleic acids research, 51(D1), D1425-D1431.

[103] Kumar MP, Du J, Lagoudas G, Jiao Y, Sawyer A, Drummond DC, et al. Analysis of Single-

Cell RNA-Seq Identifies Cell-Cell Communication Associated with Tumor Characteristics. Cell

Rep   [Internet].   2018;25(6):1458-1468.e4.   Available   from:
https://www.sciencedirect.com/science/article/pii/S221112471831636X

[104] Leon-Ferre, R. A., Jonas, S. F., Salgado, R., Loi, S., De Jong, V., Carter, J. M., ... &
International Immuno-Oncology Biomarker Working Group. (2024). Tumor-Infiltrating
Lymphocytes in Triple-Negative Breast Cancer. JAMA, 331(13), 1135-1144.

[105] Ali, H. R., Chlon, L., Pharoah, P. D., Markowetz, F., & Caldas, C. (2016). Patterns of immune
infiltration in breast cancer and their clinical implications: a gene-expression-based
retrospective study. PLoS medicine, 13(12), e1002194.

[106] Chowdhury, A., Pharoah, P. D., & Rueda, O. M. (2023). Evaluation and comparison of
different breast cancer prognosis scores based on gene expression data. Breast Cancer
Research, 25(1), 17.

1063

SUPPLEMENTARY FIGURES

1065

Supplementary Figure 1: Summary of the three bulk transcriptomics cohorts used.

A. Distribution of responders (R) vs. non-responders (NR) across the three cohorts. R and NR indicate the patients who achieved pCR following neoadjuvant treatment and those who did not, respectively.

B. Distribution of breast cancer subtypes across the three cohorts. ER+ / HER2- and TN stand for ER-positive, HER2-negative tumor subtype and triple negative tumor subtype, respectively.

C. Fractions of chemotherapy regimens used for neoadjuvant treatment across the three cohorts. E = Epirubicin, C = Cyclophosphamide, F = 5-Flurouracil, T = Docetaxel (Taxotere), P = Paclitaxel (Taxol), Cb = Carboplatin.

1075

1076

Supplementary Figure 2: The prominent cell types and ensembles mediating chemotherapy

response in breast cancer tumor microenvironment.

A-B. Odds ratios for the prominent cell types (**A**) and multi-cell-type ensembles (**B**) as compared

to the bulk across three cohorts. Cell types are ranked by their area under the receiver operating

characteristics curve (AUC) values in cross-validation in a descending order (see **Figure 2B**).

Ensembles are ranked by their AUC values in external validation in a descending order (see **Figure**

3A-D).

C-D. Sensitivities for the prominent cell types (**C**) and multi-cell-type ensembles (**D**) as compared

to the bulk across the three cohorts. Cell types are ranked by their AUC values in cross-validation

in a descending order (see **Figure 2B**). Cell ensembles are ranked by their AUC values in external

validation in a descending order (see **Figure 3A-D**).

1088

1089

Supplementary Figure 3: The prominent cell types and ensembles mediating chemotherapy response in breast cancer tumor microenvironment in a subtype-specific manner.

A-B. Comparison of model performances in cross-validation with TransNEO for the ER-positive, HER2-negative (ER+ / HER2-; **A**) and triple negative (TN; **B**) subtypes across nine cell-type-specific, bulk, and Sammut et al. predictors. AUC and AP stand for the area under the receiver operating characteristics curve and average precision (equivalent to the area under precision-recall curve), respectively. Cell types are ranked by their AUC values in a descending order.

C-D. Comparison of model performances in external validation with ARTemis + PBCP for the ER+ / HER2- (**C**) and TN (**D**) subtypes across nine cell-type-specific, bulk, and Sammut et al. predictors. Cell types are ranked by their AUC values in a descending order.

E-F. Comparison of model performances in external validation with ARTemis + PBCP for the ER+ / HER2- (**E**) and TN (**F**) subtypes across five most prominent two-cell-ensembles, five most prominent three-cell-ensembles, bulk, and Sammut et al. predictors. Ensembles are ranked by their AUC values in a descending order.

1104

1105

Supplementary Figure 4: The CD4⁺ / CD8⁺ T-cells enrichment and cell abundance − clinical

response association analyses reaffirm the poor chemotherapy response stratification by T-cells

at the current resolution.

A-B. GSVA enrichment scores (**A**) and predictive performance (**B**) of CD4⁺ T-cells for chemotherapy

response stratification across TransNEO (n = 94), ARTemis + PBCP (n = 55), and BrighTNess (n =

106) cohorts. R and NR stand for responders and non-responders, respectively. The differences

between the prediction scores were computed by using a one-tailed Wilcoxon rank-sum test. AUC

and AP stand for the area under the receiver operating characteristics curve and average precision

(equivalent to the area under precision-recall curve), respectively.

C-D. GSVA enrichment scores (**C**) and predictive performance (**D**) of CD8+ T-cells for chemotherapy

response stratification across the three cohorts.

E-G. Predictive performance of abundances of the nine cell types for TransNEO (**E**), ARTemis +

PBCP (**F**), and BrighTNess (**G**) cohorts. Cell types are ranked by their AUC values for TransNEO in a

descending order.

1120

1121

Supplementary Figure 5: The prominent immune cell types and ensembles mediating response

to the most prevalent immunogenic chemotherapy regimen in each breast cancer cohort.

A. Comparison of model performances in cross-validation with TransNEO for the T-FEC regimen

(*n* = 61) across nine cell-type-specific and bulk predictors. AUC and AP stand for the area under

the receiver operating characteristics curve and average precision (equivalent to the area under

precision-recall curve), respectively. Cell types are ranked by their AUC values in a descending

order. Here, T = docetaxel/taxotere, F = 5-flurouracil, E = epirubicin, and C = cyclophosphamide.

B-C. Comparison of model performances in external validation with ARTemis + PBCP for the T-FEC

regimen (*n* = 40) across the top cell-type-specific models (**B**) and top multi-cell-type ensembles

(**C**). Cell types and ensembles are ranked by their AUC values in a descending order.

D-E. Comparison of model performances in external validation with BrighTNess for paclitaxel +

carboplatin regimen (*n* = 106) across the top cell-type-specific models (**D**) and top multi-cell-type

ensembles (**E**). Cell types and ensembles are ranked by their AUC values in a descending order.

1135

1136

Supplementary Figure 6: The cell-cell interactions (CCIs) encompassing prominent cell types

capture the performance of CCIs using all nine cell types in mediating chemotherapy response

and are validated in single-cell (SC) breast cancer tumor microenvironment.

A-B. Prediction scores (**A**) and model performance (**B**) for DECODEMi using all available CCIs in

cross-validation with TransNEO (n = 94) and in external validation with ARTemis + PBCP (n = 55)

and BrighTNess (n = 106). R and NR stand for responders and non-responders, respectively. The

differences between the prediction scores were computed by using a one-tailed Wilcoxon rank-

sum test. AUC and AP stand for the area under the receiver operating characteristics curve and

average precision (equivalent to the area under precision-recall curve), respectively.

C-D. Odds ratios (**C**) and sensitivities (**D**) for DECODEMi using different sets of CCIs in cross-

validation with TransNEO (n = 94) and in external validation with ARTemis + PBCP (n = 55) and

BrighTNess (n = 106).

E-G. Protein-protein interaction networks obtained from STRING [48] displaying the known

interaction networks for three prominent CCIs *i.e.*, *GDF9 − BMPR1B* (**E**), *NRTN − GFRA1* and *ARTN*

− GFRA1 (**F**), and *CX3CL1 − CX3CR1* (**G**).

1152

1153

Supplementary Figure 7: The generalizability of DECODEM to single-cell (SC) transcriptomics and TCGA patient survival stratification.

A-B. Prediction scores (**A**) and the area under the receiver operating characteristics curve (AUC) values (**B**) for all available cell types in external validation with SC expression of TNBC patients (SC-TNBC) treated with chemotherapy (*n* = 6) from Zhang et al. [48]. Pseudobulk is shown as a baseline and computed as the mean expression across all available cells per patient after filtering for low-variance genes (threshold = 0.8). R and NR stand for responders and non-responders, respectively. The differences between the prediction scores were computed by using a one-tailed Wilcoxon rank-sum test.

C-D. Prediction scores (**C**) and the AUC values (**D**) for four cell types in external validation with SC expression of TNBC patients (SC-TNBC) treated with chemotherapy and immunotherapy (*n* = 9) from Zhang et al. [48].

E-F. Kaplan-Meier curves depicting the overall survival (OS; **E**) and progression-free interval (PFI;

F) of the 705 BC patients from TCGA (TCGA-BRCA). Patients were stratified into low-risk, likely

responder (n = 353) and high-risk, unlikely responder (n = 352) groups by their median DECODEM

score computed using the expression of endothelial cells, whereby the low-risk and high-risk

groups comprised individuals with scores above and below the median, respectively. The

differences between the two curves were computed by using a Log-rank test.

1172