Style deflection is determined by the handedness of phyllotaxis and differential cell

elongation in a species with mirror-image flowers

Caroline Robertson[1,*], Haoran Xue[2,*], Marco Saltini[3,*], Alice L. M. Fairnie[4], Dirk Lang[5], Merijn

H. L. Kerstens[6], Viola Willemsen[6], Robert A. Ingle[1], Spencer C. H. Barrett[4], Eva E. Deinum[3],

Nicola Illing[1], Michael Lenhard[2,6]

[1] University of Cape Town, Department of Molecular and Cell Biology, Rondebosch 7701,

South Africa

[2] University of Potsdam, Institute for Biochemistry and Biology, Karl-Liebknecht-Str. 24-25, D-

14476 Potsdam-Golm, Germany

[3] Mathematical and Statistical Methods (Biometris), Plant Science Group, 6708 PB

Wageningen, Wageningen, The Netherlands

[4] Department of Ecology and Evolutionary Biology, University of Toronto, 25 Willcocks Street,

Toronto, Ontario, Canada M5S 3B2

[5] University of Cape Town, Department of Human Biology, Observatory, 7925, South Africa

[6] Laboratory of Cell and Developmental Biology, Wageningen University, Droevendaalsesteeg

1, 6708 PB Wageningen, the Netherlands

* These authors contributed equally.

[6] author for correspondence: michael.lenhard@uni-potsdam.de

ABSTRACT

Many animals and plants show left-right (LR) asymmetry. In some animal systems, handedness has a simple genetic basis, which has allowed identifying how handedness is determined at the molecular level, even if its functional relevance remains unclear. Mirror-image flowers represent an example of LR asymmetry of clear functional significance, with the reciprocal placement of male and female organs in left- versus right-handed flowers promoting cross-pollination. Here, we use the South African geophyte *Cyanella alba* to study how handedness of its mirror-image flowers is determined and elaborated during development. Inflorescences of *C. alba* produce flowers with a largely consistent handedness. However, we find that this handedness has no simple genetic basis, and individual plants can switch handedness between years. Rather, it is the direction of the phyllotactic spiral that determines floral handedness. Cellular analysis combined with biophysical modelling demonstrates that style deflection is driven by increased cell expansion in the adaxial carpel facing the next oldest flower compared to the other adaxial carpel. The carpel with greater expansion shows transcriptional signatures of increased auxin signaling and cell-wall modifications compared to the less expanding one. We propose that a recently described inherent LR auxin asymmetry in the initiating organs of spiral phyllotaxis determines handedness in *C. alba*, representing a conserved non-genetic mechanism for creating a stable floral polymorphism. This mechanism links chirality across different levels of plant development and exploits a developmental constraint in a core patterning process to produce morphological variation of ecological relevance.

SIGNIFICANCE STATEMENT

Several plant species show left-right asymmetry in their flowers, with the female reproductive organ displaced to the left or to the right of the midline, and at least one of the male organs positioned reciprocally. This increases reproductive success by promoting cross-pollination between individuals, yet how the handedness of flowers is established is unknown. Here, we show that most *Cyanella alba* plants form flowers of only one handedness, but this is not genetically controlled. Rather, left-right determination in flowers builds on a patterning process at the shoot tip that forms leaf and flower primordia in a clockwise or counter-clockwise spiral. Thus, an inbuilt handedness at one level of development is used to establish left-right asymmetry of ecological relevance.

INTRODUCTION

Left-right (LR) asymmetry is a fascinating feature of many plants and animals. Striking examples include the asymmetric placement of internal organs in vertebrates and the left versus right coiling of snail shells (1, 2). Such LR asymmetries raise three fundamental questions. First, how is symmetry broken in a consistent manner to determine left and right? Second, how is this translated, from genes to cellular and tissue level processes, to produce asymmetric morphologies? Third, what is the biological function of LR asymmetry? The first two questions are being answered for the LR asymmetry of internal organs in vertebrates and shell coiling direction in snails (3–8). In both systems the inherent chirality of different cytoskeletal elements appears to be exploited for the initial symmetry breaking event, which is then translated into an asymmetric morphology via a partially conserved pathway. In mouse embryos, motile cilia on the cells in the node cause a left-ward flow of extracellular fluid that triggers the upregulation of *Nodal* expression on the left side (5, 6). This left-ward flow results from the inherent chirality of the microtubule arrangement in motile cilia, which causes them to rotate in a clock-wise direction, combined with their tilt towards the posterior side (9). In *Lymnaea* snails a genetic polymorphism determines coiling direction, with the dominant *D* allele causing right-handed coiling (10). The *D* locus is a duplicated copy of a formin gene (3, 7) that determines direction via actin-cytoskeleton based symmetry breaking (4, 10). In both cases LR-asymmetric upregulation of *Nodal* and of *Pitx2* transcription-factor expression translates the initial symmetry breaking event into an asymmetric morphology. Despite this progress in understanding the molecular basis of handedness, the functional relevance of LR asymmetry in these examples remains unclear.

Several examples of LR asymmetry and left- or right-handed helical growth also exist in plants (2). These include the left-right asymmetry of leaves in several species (11), left- or right-handed helical arrangement of leaves and flowers around the stem (i.e. phyllotaxis) (12), the twining of tendrils (13), and the left- or right-handed helical growth of mutants (14). Several important components that underlie helical growth *per se* have been identified, such as a polar auxin transport-based mechanism that determines the initiation of organs at the shoot meristem in a phyllotactic spiral and indirectly seems to cause LR asymmetries in leaves (15, 16), or gelatinous G-fibres in twisting tendrils (17). But how the handedness of the resulting helices is determined remains unknown. The important exception are mutants with consistently left- or right-handed helical growth (14). Most of these result from mutations in tubulin or microtubule-associated proteins that cause a spiral arrangement of cortical microtubules, with the handedness of this spiral determining the handedness of helical growth at the cell or organ level (18, 19). These mutants underscore the value of genetic

polymorphisms causing helical growth with a consistent orientation for identifying the
molecular determinants of handedness.

Mirror-image flowers (enantiostyly) represent a case of LR asymmetry in plants for which a
clear functional relevance has been demonstrated and where - at least in some taxa - the
handedness of the flowers is genetically determined (20–22). This provides an opportunity to
identify both the molecular and developmental control of handedness and its adaptive
significance (23). In enantiostylous flowers the style is deflected to the left- or right-side of the
midline that runs along the dorsoventral axis of the flower (24), and in most species a
pollinating anther is deflected in the opposite direction (Fig. 1). Enantiostylous species are
classified based on the distribution of left- and right-styled flowers on individuals (24). In
monomorphic enantiostyly, both forms of flower can be found on a single individual, whereas
in dimorphic enantiostyly a plant has only left- or right-styled flowers, called L-morph or R-
morph plants, respectively.

Enantiostyly is reliably reported from 11 unrelated angiosperm families and represents a
striking case of convergent evolution in form and function. Most enantiostylous species are of
the monomorphic type (23, 24), with dimorphic enantiostyly reliably reported from just two
monocotyledonous lineages: Pontederiaceae (*Heteranthera*) and Haemodoraceae
(*Wachendorfia*, *Barberetta*). A third instance in Tecophilaeaceae (*Cyanella*) has been reported
(20), but this has not been fully confirmed. Experimental studies have demonstrated that
enantiostyly represents a floral adaptation that promotes cross-pollination and reduces sexual
interference between male and female function in hermaphroditic flowers (25, 26). It does so
by optimizing the placement of pollen on the bodies and wings of pollinating insects for efficient
disassortative pollen transfer to styles on the opposite morph. Studies using genetic markers
indicate that dimorphic enantiostyly promotes significantly higher levels of outcrossing than
the monomorphic condition (21, 27). The efficiency of dimorphic enantiostyly in promoting
disassortative pollination has also been demonstrated using fluorescently labelled pollen
grains in *Wachendorfia* (28), although similar studies in *Barberetta aurea* failed to observe this
effect (29). Thus, at least for some species/pollinator combinations the pollen that left-styled
(L) flowers deposit on the right side of visiting insects is transferred efficiently to the stigmas
of right-styled (R) flowers, and vice versa. The mode of inheritance of dimorphic enantiostyly
has only been determined in *Heteranthera multiflora* (now *H. missouriensis*), where
handedness is determined by segregation at a single Mendelian locus (22), with the dominant
allele controlling R flowers and the homozygous recessive genotype having L flowers.

Here, we investigate the molecular and development basis of enantiostyly in *Cyanella alba*
subsp. flavescens (hereafter *C. alba),* a long-lived, deciduous geophyte restricted to the

Western and Northern Cape regions of South Africa (30). Plants form deep-seated corms that

produce a basal rosette of narrow erect leaves after winter rains with flowering occurring from

early August to early October. The inflorescence is a compact raceme with very little stem

elongation, and flowers are borne individually on long pedicels (10-20 cm), giving the

appearance of solitary flowers emerging from the rosette. Each plant produces a small number

(generally not more than 15) of long-lived flowers during the flowering season, but daily display

size is most commonly one to two flowers (31). *C. alba* is a candidate for showing dimorphic

enantiostyly based on previous work (20, 24, 32), as the majority of plants were reported to

form flowers of only one handedness and only a few plants produced both types of flowers

(32). Thus, floral handedness in *C. alba* may have a genetic basis.

Our investigation used a sequencing-based approach to search for the presumed genetic

basis of floral handedness in *C. alba*. By analogy to other similar floral polymorphisms and to

enantiostyly in *H. missouriensis*, we hypothesized that floral handedness in *C. alba* was under

the control of a single *ENANTIOSTYLY* (*E*) locus with a dominant and recessive allele. Under

such a scenario, disassortative (intermorph) mating between L- and R-morphs would produce

an equal ratio of L- and R-morphs in the progeny, with one morph homozygous recessive and

the other heterozygous at the *E* locus. Our study aimed to answer three questions. First, is

there indeed a genetic basis to floral handedness in *C. alba*? Second, what is the symmetry-

breaking event that determines the direction of style deflection in flowers? Third, which cellular

and developmental processes cause consistent style deflection to one side?

RESULTS

Sequencing-based analysis and transcriptomics do not support a simple genetic basis for floral handedness

To determine whether there is a genetic basis for handedness, we established a reference genome assembly from an individual that had formed seven consistently R-flowers in 2021 (see Supporting Information). We also performed Illumina-sequencing on 25 *C. alba* individuals that had four or more flowers of consistent handedness in 2021 (15 L and 10 R), and 20 with six or more flowers of consistent handedness in 2022 (11 L and 9 R; see Supporting Information).

Assuming a similar system as in *H. missouriensis* (22) (see above), we tested two hypotheses regarding a possible simple, single-locus genetic basis of floral handedness in *C. alba*. The first was that one of the morphs is heterozygous and the other homozygous recessive. To test this, we performed a genome wide association study (GWAS) to identify single nucleotide polymorphisms (SNPs) associated with each phenotype. We aligned all 45 Illumina samples to the reference assembly and phenotype-associated SNPs were identified (see Supporting Information Text). The list of SNPs was filtered for sites where more than 80% of the individuals of one phenotype were heterozygous for the alternate alleles of the SNP, whereas in the other only <20% were heterozygous. We chose a relaxed threshold of 80% to account for the possibility that some individuals may have been mis-categorised, especially in the 2021 sample with its lower flower number per individual. In other words, we considered it possible that a genotypically L-plant might still have produced four R-flowers by a series of developmental errors or vice versa. The Illumina2021 and Illumina2022 datasets were first analysed separately, before combining all 45 samples (Table 1). In the Illumina2021 dataset, more heterozygous SNP sites were identified in R- than L-plants (Table 1), but the opposite pattern occurred in the Illumina2022 samples (Table 1). The combined dataset identified more SNPs that were heterozygous in L- than in R-plants. However, none of the identified SNPs were supported across more than one analysis. Therefore, we conclude that the identified SNPs are most likely unrelated to the putative *E* locus and that enantiostyly in *C. alba* does not appear to be governed by a bi-allelic Mendelian locus.

Our second hypothesis was that the putative *E* locus is hemizygous, as demonstrated in several heterostylous species (33–35), and present only in either L- or R-plants. As our reference genome was from a presumed R-plant, we first searched for an R-exclusive region by performing a coverage analysis. We calculated the number of Illumina reads mapping to each region of the haploid *C. alba* genome and compared left and right pools to identify regions where there was a significant difference in coverage. We used the two haploid assemblies

generated by Hifiasm separately for this analysis. As above, we performed the analysis first
with the Illumina2021 and Illumina2022 datasets separately, and then with the combined
dataset of all 45 samples. Again, none of the windows identified as having significantly
different coverage in L- compared to R-samples were shared between the Illumina2021 and
Illumina2022 datasets (Table S1), and no plausible candidate for a hemizygous region in the
right-handed *C. alba* assembly was identified.

We next asked whether the putative *E* locus may be exclusively present in L-plants and thus
undetectable with our R-plant reference assembly. Such a hemizygous region should be
detectable based on L-plant exclusive *k*-mers in the L-pool (see Supporting Information). In
the Illumina2021 data set there were around 2.5 times more R- than L-associated *k*-mers
(Figure 1A). By contrast, analysis of the Illumina2022 samples identified over five times more
L- than R-associated *k*-mers. Surprisingly, all individuals, regardless of handedness,
possessed more L- than R-associated *k*-mers. The much higher number of phenotype-
associated *k*-mers in the Illumina2022 than the Illumina2021 dataset may reflect the smaller
sample size of the former (20 as opposed to 25 individuals), and thus a possibly higher rate
of false-positive phenotype-associated *k*-mers. As there was no agreement between the
Illumina2021 and Illumina2022 datasets in our analysis, there is no strong support for a
hemizygous region exclusive to L-plants.

As a complement to the coverage- or *k*-mer based search for a hemizygous region, we used
RNA-seq of dissected styles (including the upper half of the ovary) to search for morph-specific
transcripts. We harvested triplicate samples of styles from very early-, early- and mid-stage
buds of plants with exclusively R- or L-flowers (see Supporting Information Text for definition
of stages). No style deflection is visible in very early-stage buds, but this becomes detectable
in early buds and very obvious in mid-stage buds (see Figure 3A below). Thus, our samples
should cover the critical time window when the orientation of style deflection is determined.
We identified 137 differentially expressed genes between L- and R-samples (FDR: $\alpha < 0.05$),
of which 52 were upregulated and 85 downregulated in R- versus L-buds (Table S2). None of
the genes were exclusively expressed in L or R plants. Thus, the transcriptomic analysis does
not provide evidence for genes only present or only expressed in one of the phenotypes.
Taken together, our results show no convincing evidence for a simple genetic basis to floral
handedness in *C. alba*.

Figure 1: Floral handedness has no simple genetic basis in *C. alba*

(**A**) Image of L- and R-flowers of *Cyanella alba* subsp*. flavescens.* Feeding anthers (FA) and pollinating anther (PA) are indicated. L and R indicate left- and right-deflected styles.

(**B,C**) Phenotype-associated *k*-mer composition of *C. alba* individuals in the Illumina2021 (**B**) and Illumina2022 (**C**) datasets. *K*-mers associated with either the L- or R-pools were identified using KmerGO2. The total number of L- and R-associated 31-mers in each individual was calculated and plotted.

(**D**) Distribution of floral phenotypes of the same cohort of 135 plants in 2022 and 2023. The colour gradient indicates the consistency in floral orientation between years (red: fully consistent, yellow: fully inconsistent). Black bars on the outside show marginal distributions.

Switching floral handedness on individual plants between flowering seasons indicates that the direction of style deflection is not genetically determined

The above conclusion that there is unlikely to be a genetic basis to handedness received further support from our demographic studies of individually marked plants during the flowering seasons of 2022 and 2023. In 2022, 188 plants in three populations were permanently marked with metal pins, and in 2023 we recovered 135 of these plants. We followed the plants throughout the duration of both flowering seasons and recorded the number of L- and R-flowers produced by each individual (Figure 1D). Strikingly, of the 44 plants recovered in 2023 that had formed exclusively L-flowers in 2022, 22 retained their handedness and 22 switched handedness in 2023. Of the latter, 16 produced exclusively R-flowers and 6 produced both L- and R- flowers (proportion of R flowers within a plant ranged from 40-75%). Conversely, of the 36 plants recovered in 2023 that had exclusively R-flowers in 2022, 19 switched their handedness in 2023. Again, the same trend was seen as above with the majority, 15 plants, producing exclusively L-flowers in 2023, and the remaining four plants (21%) producing both flower types (proportion of L flowers within a plant ranged from 33-60%). Thus, these results demonstrate that the handedness of flowers produced by an individual plant during one flowering season is highly consistent, but that this consistent handedness can switch within an individual from one year to the next. This argues strongly against the genetic determination of stylar handedness in *C. alba* and suggests that populations of the species exhibit monomorphic enantiostyly and not dimorphic enantiostyly as earlier assumed.

The handedness of the phyllotactic spiral determines the consistent bias in floral handedness within plants

The lack of direct genetic control of handedness raises an intriguing question: How can flowers
in an inflorescence consistently develop an L- or R morphology without genetic input, and how
can they switch their annually consistent bias for one side to the other in the next year? One
developmental process with a highly consistent handedness over the growing period of an
inflorescence is the direction of the phyllotactic spiral. *C. alba* shows such spiral phyllotaxis
(Figure S1), with organs initiated at the shoot apical meristem in either a clockwise or counter-
clockwise direction. Changes in the handedness of the phyllotactic spiral produced by one
shoot meristem appear to be exceedingly rare (36), providing stability in any one year. This
aslo seems to hold true in *C. alba*, as none of 133 plants we sampled showed a consistent
handedness reversal of the phyllotactic spiral. Moreover, studies of many angiosperm species
indicate that the handedness of the phyllotactic spiral is not under genetic control, and that
initiating shoot meristems randomly establish either a clockwise or counter-clockwise spiral
with equal probability (12). This pattern is also true for the individual lateral meristems on a
single plant. Thus, it is likely that the shoot meristems initiated in successive years by *C. alba*
will also randomly adopt a clockwise or counter-clockwise orientation. We therefore asked
whether the direction of style deflection in *C. alba* flowers is predicted by the handedness of
the phyllotactic spiral.

To address this question, we determined the handedness of the phyllotactic spiral on 132 *C.*
alba plants in the 2023 flowering season and the phenotype of between 1 and 7 open flowers
and sufficiently mature buds for recognizing the style orientation per plant (average 2.74,
median 3 flowers per plant; a small number of flowers with unclear style orientation were
excluded). This revealed a strong correlation between the direction of the phyllotactic spiral
and floral handedness, with a correlation coefficient of 0.82 (0.76 – 0.87). Most plants with a
counter-clockwise or right-handed phyllotactic spiral (defined from older to younger organs)
formed only R flowers, and *vice versa* for plants with a clockwise or left-handed phyllotactic
spiral. Twenty plants produced both R and L flowers, and there were very few plants (5/132)
where floral handedness consistently contradicted that predicted by the phyllotaxis direction.
Thus, the handedness of the phyllotactic spiral in *C. alba* determines floral handedness,
suggesting that it provides the consistently biased input into flower development to ensure the
non-random distribution of style deflection of individual plants through time.

We next asked whether deviations from the floral handedness predicted by phyllotaxis were
randomly distributed in an inflorescence. For this we plotted the fraction of such deviations per
absolute flower position in an inflorescence and compared it to the overall average of all
flowers. The distribution of deviations was non-random across the different positions
($p<0.0001$, X^2-test), with higher deviation rates at the very first and at higher positions beyond
position 5, and the lowest deviation rates at positions 2 to 4, i.e., the very first and the later

flowers of the season were more likely to have the deviating handedness. The reason for this non-random distribution of errors is unclear at present, but it may suggest an additional flower-flower interaction that cannot operate on the first flower and becomes weaker over time.

Figure 2: Direction of the phyllotactic spiral determines floral handedness in *Cyanella alba* subsp*. flavescens*

(**A**) Correlation of phyllotactic direction (CW: clockwise, CCW: counter-clockwise) and floral handedness. Numbers in the diagram indicate the number of individual plants. 'Unclear' phyllotaxis indicates plants with individual divergence angles that are inconsistent with their overall phyllotaxis orientation. See Materials and Methods in Supporting Information for more details and our interpretation of these cases.

300 (**B**) Distribution of deviations from main pattern (flowers with the opposite style deflection compared to
301 the orientation predicted based on phyllotaxis) across flower positions in an inflorescence. Numbers in
302 diagram indicate the number of flowers scored at each position. Horizontal red line shows average
303 fraction of such errors over all flower positions. Dots are observed values and vertical lines indicate
304 95% confidence intervals for the fraction of deviations per position calculated based on a binomial test.
305

Style deflection results from differential elongation of the two adaxial carpels

The styles of *C. alba* are initially straight with no discernible difference between L- and R-
flowers (Figure 3A). When styles of early buds reach 70% or more of the length of the
pollinating anther, they begin to deflect slightly to the left or right of the midline. The stigma tip
can also be slanted at this stage, with the slant facing the direction to which the style will be
deflected. In mid-stage buds, one to two days before anthesis, the style is clearly deflected
from the midline, and in open flowers the style and pollinating anther are fully deflected to
opposite sides of the flower. The style is relatively straight, with the exception of the stigma tip
curving sharply upwards. The style movement to one side can be observed using time-lapse
microscopy in dissected flowers cultivated *in vitro* (Video S1). The hinge-like motion of the
style is coupled with expansion and elongation of the ovary, as confirmed by measuring the
distances between the three marked spots on the ovary, which increased by 14% and 10% in
the length and width direction over the course of the video, respectively (Video S1). These
observations suggested that style deflection might be driven by differential elongation of the
carpels. The *C. alba* gynoecium consists of three carpels, with the two adaxial carpels lying
on either side of the dorsoventral midline of the flower and the third carpel below, such that
the midline bisects it (Figure 3B). If one of the two adaxial carpels elongated more than the
other, this could push the style away from the more strongly elongating carpel.

To test the feasibility of this mechanism and investigate the contribution of differential
elongation of different carpel walls, we generated a biophysical model of the *C. alba* pistil. We
used a three-dimensional polyhedral model to represent the ovary, which is composed of a
triangular prism topped by a truncated pyramid (Supporting Information Text; Figures S2, S3;
Tables S3 to S8). Deflection of the truncated face of the pyramid such that it is no longer
parallel to the base of the triangular prism indicates style deflection. We applied this model to
images of six ovaries and identified for each the polyhedron whose two-dimensional side view
most accurately represented the ovary in its later stages of development (see Figure 3B-E,
and Figures S1, S2). The resulting solid figure is skewed, reflecting the differential carpel
elongation responsible for stylar deflection. The best fitting polyhedron was then recreated
with a static bead-spring system, with the edges of such a polyhedron replaced by springs
with different elongation and stiffness (Figure 3F). The elongation and stiffness of each spring

was determined by random search optimisation algorithms (37), to identify the parameters of

our mechanistic model that enable a reproduction of the best fitting polyhedron and, hence, of

the ovary of *C. alba* during the late developmental stage (Figures S4, S5).

The results of this optimization indicated that both the inner and outer walls of carpel 1 (the

adaxial carpel opposite the direction of style deflection) undergo more significant elongation

than those of carpels 2 (the adaxial carpel on the side to which the style deflects) and 3 (the

abaxial carpel) (Figure 3G, Figure 4A). The slight elongation seen in carpels 2 and 3, as

predicted by our model, arises from the tensile forces generated by links between different

carpels at the base of the style. This effect is driven by the greater relative stiffness of the base

compared to other system walls (Figure 3H), contributing to guiding the shape and

morphogenesis of the style (38, 39). Our analysis revealed that both inner and outer walls of

carpel 1 end up being compressed, whereas those in carpels 2 and 3 are stretched. Based on

the analyses of six *C. alba* ovaries through our models, the primary driver behind the

elongation of carpel 1 is the extension of the outer wall, with an average detected elongation

relative to the initial length of the early bud stage of 2.47 (SD = 0.41) times, as opposed to the

inner wall mean relative elongation of 1.28 (SD = 0.18) times.

Figure 3: Schematic of the data-driven and mechanistic models and analysis of ovaries of *Cyanella alba* subsp*. flavescens*.

(**A**) Stereomicroscope images of *C. alba* buds at four stages of development reveal the developmental sequence of stylar bending. Blue line indicates the slant of the stigma in early-stage buds. Scale bars are 1 mm.

(**B**) Cross-section of a *C. alba* ovary indicating the trilocular architecture. Orange lines represent a cross-section of the modelling polyhedron. Red arrows indicate the edges of the polyhedron representing inner and outer wall of a carpel. The outer-wall positions overlie the carpel midveins.

(**C**) The ovaries are modelled as polyhedra composed of triangular prisms topped with a truncated
pyramid. Red edges denote the inner and outer walls of the same carpel highlighted in (**B**).
(**D**) Two-dimensional projection of the lateral view of a collection of polyhedra.
(**E**) The selection process of the best fitting polyhedron for modelling the ovaries of *C. alba*, consists of
choosing among all possible polyhedra based on area, perimeter, and length and width of the ovary.
(**F**) Biophysical parameters of our bead-spring models are optimised to achieve an accurate fit of the
polyhedron to the ovary structure.
(**G**, **H**) Elongation of the inner and outer walls of the *C. alba* ovaries (**G**) and stiffness of the organ walls
(**H**) as predicted by our mechanistic model. Error bars represent standard deviation. n=6
371
The above model makes two key predictions. First, the outer wall of carpel 1 should elongate
more than the outer wall of carpel 2. Thus, in an R-flower we would predict the top left carpel
(when viewed from the front) would elongate more than the top right carpel (Figure 4A). The
second prediction is that the outside wall of carpel 1 should elongate more than the inner wall
corresponding to the placenta. To test the first prediction, we measured the lengths of the
three carpels in the ovary along their prominently visible midveins at four different stages of
development and combined data for L- and R-flowers. Carpel 1 was significantly longer than
carpel 2 and the abaxial carpel 3 in both open flowers and mid-stage buds (Figure 4B). Carpel
2 was on average 89.9% ± 4.3% and 88.9% ± 5.42% of the length of carpel 1 in open flowers
and mid-stage buds, respectively. By contrast, carpel lengths were indistinguishable in early
or very early-stage buds (Figure 4B). This mirrors the observation that clear deflection of the
style to the left or right is only visible in mid-stage buds and open flowers and supports the
hypothesis that differential elongation between the two adaxial carpels causes style deflection
in *C. alba*.
To further support the link between differential carpel elongation and style deflection, we
repeated this experiment on *C. hyacinthoides*, a close relative of *C. alba* that does not exhibit
enantiostyly (Figure 4C). Styles of *C. hyacinthoides* are not deflected to the left or right, but
instead towards the abaxial side. The structure of the trilocular *C. hyacinthoides* ovary
matches that of *C. alba*. Carpels 1 and 2 of *C. hyacinthoides* were indistinguishable in length,
but carpel 3 was significantly shorter than carpel 1 (Figure 4D). This suggests that the reduced
expansion of carpel 3 underlies the downward deflection of the style in *C. hyacinthoides*,
whereas the differential elongation of carpels 1 and 2 determines the LR deflection of the style
in *C. alba*.

Figure 4: Differential carpel elongation underlies style deflection in *Cyanella alba* subsp*. flavescens*

(**A**) An open *C. alba* flower. The arrangement of the three carpels in the trilocular ovary is indicated by the subdivided circle.

(**B**) Measurements of carpel lengths from ovaries of flowers at the indicated stages of development. Lengths were normalised to the average value of the three carpels per ovary. Values are means ± SD here and throughout the other panels. Sample sizes are indicated and show the number of ovaries from which the paired measurements were taken. *: significantly different from carpel 1 (p<0.05, paired t-test).

(**C**) *C. hyacinthoides* is a closely related spec*ies* without enantiostyly. The style is deflected downwards and placed centrally between the six anthers.

(**D**) Measurements of carpel lengths from *C. hyacinthoides* ovaries. *: significantly different from carpel 1 (p<0.05, paired t-test).

(**E**) Fluorescence micrograph of a calcofluor white-stained pistil. Blue and orange boxes indicate the
regions where the measurements of cell sizes shown in (**F**, **G**) and (**H**) were taken, respectively. V2
indicates midvein of carpel 2.
(**F**, **G**) Cell lengths (**F**) and cell widths (**G**) of epidermal cells overlying the midveins of carpels 1 and 2
in the upper third of the ovary are shown. Cell length is the extension of the cell along the long axis of
the ovary, and cell width the extension perpendicular to the long axis of the ovary. *: significantly
different from carpel 1 ($p<0.0001$, paired t-test).
(**H**) Cell lengths of epidermal cells overlying the midveins of carpels 1 and 2 at the tip of the ovary are
shown. *: significantly different from carpel 1 ($p<0.0001$, paired t-test).
(**I**) Lengths of subepidermal cells in the outer and inner walls of carpels 1 and 2. The number of carpels
from which the measurements were taken is indicated inside the graph. Letters indicate mean cell
lengths that are significantly different as determined by a two-way repeated measures ANOVA followed
by Tukey's HSD post-hoc test ($\alpha=0.05$). In total 14 ovaries were dissected and imaged, but for some of
them individual walls could not be imaged in sufficient quality to allow cell-length measurements.
Scale bars are 1 mm.
424
We next asked whether the differential elongation of carpels 1 and 2 in *C. alba* was due to
differences in cell elongation or cell division. The length of the epidermal cells overlying the
carpel midveins was measured from confocal laser-scanning microscopy images of the distal
third of ovaries (Figure 4E, Figure S6). Cell length (i.e. the size of the cell along the ovary's
longitudinal axis) was significantly greater along the midvein of carpel 1 than along carpel 2,
with mean cell length in carpel 2 only 81.8% of that observed in carpel 1 (Figure 4F). This
difference is more than sufficient to explain the observation that carpel 2 length is on average
89.9% of carpel 1 length (Figure 4B). In addition, carpel 1 cells are significantly wider than
carpel 2 cells (Figure 4G). To confirm the observed differences in cell length, we repeated the
measurement at the tip of the ovary where it transitions into the style. Again carpel 1 cells
were significantly longer than carpel 2 (and also carpel 3) cells, with the difference between
carpels 1 and 2 even greater (25%) than in the upper third of the ovary (Figure 4H), suggesting
a proximo-distal gradient in differential cell elongation.

To test the second prediction from our model, we compared cell lengths between the outer
carpel walls and the inner walls at the placenta for carpels 1 and 2 (Figure S7). Subepidermal
cells were used to test this prediction, as they could be clearly and reproducibly measured,
and the epidermis was often damaged due to the need to remove the ovules before imaging.
As expected, outer-wall cells from carpel 1 were longer than outer-wall cells from carpel 2,
confirming the above measurements. Outer-wall cells from carpel 1 were also longer than
inner-wall cells from carpel 1, while those of carpel 2 were not. Thus, these measurements
fully support the predictions from the model.

Collectively our empirical results and biophysical model indicate that stronger isotropic cell
expansion in the outer wall of carpel 1 than carpel 2 drives style deflection to the opposite side
of the midline from carpel 1.

450 Differential carpel growth associated with increased gene expression of functional
451 categories linked to cell elongation in carpel 1

Given the differential elongation of the two adaxial carpels, we compared their gene
expression patterns using RNA-seq data. Outer walls of carpel 1 and 2 were dissected from
early- and mid-stage buds with their individual directions of style deflection clearly discernible.
We identified 53 differentially expressed transcripts (FDR<0.05, absolute log2 fold-
change(carpel 1/carpel2) >1). These included four *Auxin/Indole-3-Acetic Acid* (*Aux/IAA*) and
one *SMALL AUXIN UP-REGULATED* (*SAUR*) transcripts, all of which were upregulated in
carpel 1 (Table S9). To complement this individual-transcript based analysis, we functionally
annotated transcripts using Mercator4 (40) and tested for bins that differed significantly from
the remaining transcriptome in terms of their distribution of log2 fold-change values. In total,
106 bins showed a significantly different distribution compared to the background (Table S10).
The empirical cumulative-distribution function (ECDF) plots indicated that for many of these
bins gene-expression differences between the two carpels were less variable than the
transcriptomic background, consistent with these being large bins of house-keeping genes
(Figure S8). We identified three bins with a possible link to cell elongation as significantly
different from the background. These were "Cell wall organisation", "Phytohormone action"
and "Cytoskeleton organisation". The most specific significant subcategories within these
were "pectin methylesterase" and "Fasciclin-type arabinogalactan protein"; "RALF/RALFL-
precursor polypeptide", "auxin-responsive genes *(SAUR)" and "transcriptional repressor
*(Aux/IAA)"; and "alpha-beta-tubulin heterodimer", respectively. For each of these, transcript
levels were higher in the carpel 1 than carpel 2 samples, visible by a consistent shift in the
ECDF plot to positive values (Figure 5A). This higher expression in carpel 1 was particularly
evident when considering those members of each gene family with higher absolute expression
levels (Figure 5B). As discussed below, the elevated expression of transcripts encoding pectin
methylesterases, Fasciclin-type arabinogalactan proteins, SAUR and Aux/IAA proteins in
carpel 1 is consistent with increased cell elongation driven by higher auxin levels in carpel 1
versus carpel 2. Our transcriptome analysis therefore suggests a molecular basis for the
differential growth of the two adaxial carpels driving style deflection.

480
481

Figure 5: Transcriptome comparison of the two adaxial carpels in *Cyanella alba* subsp*.*
flavescens

(**A**) Empirical cumulative distribution functions (ECDFs) of the log2 fold-changes (log2FC) in transcript
abundance between carpels 1 and 2 are shown for selected Mercator bins with significantly different
log2FC distributions compared to the transcriptome-wide background ('other bins'). See Table S10 for
results for all bins.

(**B**) Log2 fold-changes (log2FC) in transcript abundance between carpels 1 and 2 are shown relative to
the average base expression across all samples for all genes in the Mercator bins shown in (**A**).

(**C**) Schematic model for the development of a consistent floral handedness based on an initial
asymmetry in auxin concentration in the initiating organs (gradients in circles, P2 to P0) at the shoot
meristem (green). The direction of this initial asymmetry depends on the direction of the phyllotactic
spiral (curved arrow, counter-clockwise in this illustration) and is later translated into style deflection (to
the right in this illustration), by stronger elongation of carpel 1 than carpel 2.

DISCUSSION

Our results provide multiple lines of evidence indicating that *C. alba* possesses monomorphic rather than dimorphic enantiostyly. Molecular analyses revealed that the consistent handedness of the mirror-image flowers of individual plants within any single flowering season is not determined by their genotype. Rather, style deflection depends on the apparently random choice that each annually initiated shoot meristem makes concerning the handedness of the phyllotactic spiral. Once determined, the handedness of the phyllotactic spiral is stable throughout one growing season, causing the flowers formed during that season to have a predictable and consistent direction of style and anther deflection, with only a few flowers per individual not conforming to this pattern. We also show that this left-right asymmetric morphology has a simple cellular and developmental basis. Developmentally, style deflection is caused by differential cell elongation between the two adaxial carpels, coupled with a very stiff base to the ovary. The stronger elongation of one of the carpels is associated with, and likely caused by, increased auxin signaling and higher expression of cell-expansion promoting genes. Thus, our work identifies a non-genetic mechanism for ensuring consistent left-right handedness that links chirality across different levels of plant development to produce adaptive morphological variation.

The mechanism for determining stylar deflection based on the handedness of the phyllotactic spiral mirrors recent findings about leaf asymmetries in other plants. In tomato, *Arabidopsis thaliana* and several other species, the direction of more or less subtle LR asymmetries in leaves can be predicted from the handedness of the phyllotactic spiral (11, 16, 41). These asymmetries have been traced back to an unequal distribution of auxin and auxin signalling in initiating leaf primordia at the shoot apical meristem (16). In particular, the side of the primordium facing the next oldest (the so-called descending side) has higher auxin levels than the ascending side, resulting from the pattern of polar auxin transport. We therefore consider it highly likely that floral asymmetry in *C. alba* is ultimately caused by such an initial auxin asymmetry in initiating bract/floral primordia (Figure 5C) for two reasons. First, auxin maxima seem to determine the sites of organ initiation at the shoot apical meristem across all mono- and dicotyledonous plants studied, so also most likely in *C. alba*. Second, the asymmetry of auxin levels in initiating primordia in spiral phyllotaxis appears to result from the pattern of polar auxin transport itself that generates these auxin maxima, leading to what has been interpreted as an inherent mechanistic constraint on the starting conditions in an initiating primordium (16), which most likely applies to all species with spiral phyllotaxis.

This model raises three questions. How is the presumed initial auxin asymmetry translated into floral handedness? Is a similar mechanism operating in other plants with mirror-image

flowers? What could be the functional significance of linking floral variation to the working of a core shoot-patterning mechanism? In a *C. alba* individual with counter-clockwise phyllotaxis and style deflection to the right, the left adaxial carpel (when viewed facing the flower) corresponds to the descending side of the initial primordium. Thus, according to our model the stronger auxin signalling on the descending side of the primordium ultimately triggers stronger cell expansion in the corresponding adaxial carpel (Figure 5C). When combined with a very stiff base of the ovary, such differential carpel elongation is predicted by our biophysical model to cause style deflection away from the descending side of the flower. Strikingly, our transcriptomic comparison between the more- and less-expanding adaxial carpels provides evidence for stronger auxin signalling in the carpel on the descending side of the initial primordium. This carpel has higher expression of Aux/IAA and SAUR encoding genes, both of which are induced by auxin signalling (42, 43). The latter also have a well-established role in mediating acidification of the extracellular space, resulting in increased cell expansion as described by the acid-growth hypothesis (42, 44, 45).

In addition, the more strongly expanding adaxial carpel also shows higher expression of genes encoding extracellular pectin methylesterases and Fasciclin-like arabinogalactan proteins (FLAs). Overexpression of a pectin methylesterase in *A. thaliana* shoot meristems resulted in less rigid cell walls (46), presumably facilitating cell expansion, while overexpression of a FLA in cotton-fibre cells increased their elongation (47), suggesting that both activities also contribute to stronger expansion of the adaxial carpel on the descending side of the flower. Auxin and the auxin-response factor ARF3/ETTIN have been shown to induce pectin methylesterase activity or expression in tobacco and *A. thaliana*, respectively, suggesting a similar link in the descending carpel (48–50). At the same time, this carpel also shows higher expression of potential negative regulators of cell expansion, in particular Rapid Alkalinisation Factor (RALF) peptides and three pectin-methylesterase inhibitors, which limit cell expansion in *A. thaliana* roots and shoot meristems, respectively (46, 51). RALF signalling has been shown to increase auxin biosynthesis in *A. thaliana* roots (51), providing another possible link between the differentially expressed gene categories in the *C. alba* carpels.

Thus, even though the mechanistic details remain to be resolved, the more strongly expanding adaxial carpel shows higher expression of both auxin-induced genes and genes linked to the control of cell elongation than the less expanding one. These findings suggest a molecular basis for its stronger growth and ultimately style deflection. At the same time, it remains an open question how the information from the initial auxin asymmetry in the primordium could be maintained for the long time across many cell divisions between floral-meristem initiation and late stages of pistil morphogenesis. An epigenetic process may be involved, but this remains speculative at present. However, this long duration between the presumed initial

trigger and the later elaboration of the morphological asymmetry may explain the small fraction of 'developmental errors' where individual flowers do not follow the pattern of style deflection predicted from the handedness of the phyllotactic spiral.

We propose that our model for *C. alba* may be more widely applicable to other species with floral asymmetry. An interaction between processes at the inflorescence meristem and flower development predicting the handedness of mirror-image flowers is also seen in other species with monomorphic enantiostyly, such as *Solanum rostratum* (52). The inflorescence of this species is a scorpioid monochasial cyme, i.e. each meristem terminates in a flower and inflorescence growth is continued by a single lateral meristem, whose position alternates relative to the respective main axis. Styles are almost always deflected towards the lateral branch, resulting in pendulum asymmetry (53), and suggesting that positional information from the primordium stage translates into floral handedness. A more directly comparable case is found in *Monochoria australasica* (also referred to as *Pontederia australasica*). This species of Australian perennial marsh plant shows inflorescence-level monomorphic enantiostyly, i.e. the flowers within one inflorescence have a consistent handedness, yet different inflorescences produced by the same individual can have a different floral handedness (24). Inspection of publicly available photographs of *M. australasica* inflorescences shows that the species has spiral phyllotaxis (Figure S9, Table S11). Across all available images where the handedness of both phyllotaxis and style deflection could be reliably assessed, we found that 7 of 9 inflorescences with a clockwise phyllotactic spiral had styles deflected to the right (one inflorescence mixed, one with L-flowers), and all 12 inflorescences with counter-clockwise phyllotaxis had styles deflected to the left. Thus, the handedness of the phyllotactic spiral again appears to determine floral handedness, although with an inverted relation compared to *C. alba*.

In plants with monomorphic enantiostyly where several flowers with different handedness are open on a given day, there is a risk of between-flower selfing (geitonogamy), as approximately half of the flowers should donate pollen efficiently to the other half (21, 27). This could result in gamete wastage due to inbreeding depression and pollen discounting. Linking the handedness of individual flowers to phyllotaxis as a stable pattern-generating process at the shoot apical meristem can unify floral handedness across an inflorescence, thus limiting the mating costs of geitonogamy (54). At the same time, the random establishment of clockwise versus counter-clockwise phyllotaxis across individual shoot meristems results in an equal distribution of L- and R-biased inflorescences in a population, maximizing mating opportunities and effectively functioning like dimorphic enantiostyly. These beneficial effects would be maximal in a situation where the relation between phyllotactic and floral handedness was absolute, individuals only formed a single inflorescence, and there was a large floral display

with many open flowers on a given day. The latter seems to be the case in *M. australasica*,
yet its clonal growth with the possibility of forming more than one inflorescence per individual
reintroduces some risk of within-plant selfing owing to between-inflorescence geitonogamy. In
C. alba, each plant appears to form only a single inflorescence per growing season and daily
floral display size is small, but often still larger than one. This suggests that inflorescence-level
uniformity of floral handedness can limit within-plant selfing and promote disassortative mating
between individuals of opposite phyllotactic and floral handedness.

Thus, in conclusion we describe a conserved mechanism whereby plants exploit an inherent
developmental constraint leading to chirality in a core developmental process to generate
ecologically relevant morphological variation in the form of LR asymmetry.

ACKNOWLEDGEMENTS

We are grateful to Marrietta and Barry Lubbe of Mertenhof farm and Barend Salomo and all
members of the Wupperthal Original Rooibos Cooperative for permission and support in
sampling *Cyanella alba*. We thank Sinéad Watchorn and Lia Hemerick for support in the field
sampling. We thank Christian Kappel and Mathias Scharmann for suggestions on genome
assembly and transcriptome analysis. CR held a Harry Crossley Research Fellowship. M.K
was funded by the Nederlandse Organisatie voor Wetenschappelijk Onderzoek
(GSGT.2019.019). This work was supported by a grant from the Human Frontiers Science
Program to E.D., N.I., S.C.H.B. and M.L. (grant number RGP0036/2021).

DATA AVAILABILITY

Whole genome and transcriptome datasets generated in this work will be available in NCBI
under under BioProject PRJNA1123744.

 TABLES

631 **Table 1: Numbers of morph-associated SNPs in *Cyanella alba* subsp. *flavescens***

632 The number of morph-associated SNPs in each dataset where 80% or more individuals of a
633 given handedness are heterozygous at the SNP site is given. The analysis was restricted to
634 the protein-coding regions of the *C. alba flavescens* genome.

	SNPs where L-plants are heterozygous	SNPs where R-plants are heterozygous
Illumina2021 dataset (25 individuals)	10	12
Illumina2022 dataset (20 individuals)	12	17
Combined dataset (45 individuals)	0	1

REFERENCES

1. M. Blum, K. Feistel, T. Thumberger, A. Schweickert, The evolution and conservation of left-right patterning mechanisms. *Development* **141**, 1603–1613 (2014).

2. D. R. Smyth, Helical growth in plant organs: mechanisms and significance. *Development* **143**, 3272–3282 (2016).

3. A. Davison, *et al.*, Formin Is Associated with Left-right asymmetry in the pond snail and the frog. *Curr Biol* **26**, 654–660 (2016).

4. R. Kuroda, B. Endo, M. Abe, M. Shimizu, Chiral blastomere arrangement dictates zygotic left-right asymmetry pathway in snails. *Nature* **462**, 790-U112 (2009).

5. S. Nonaka, H. Shiratori, Y. Saijoh, H. Hamada, Determination of left-right patterning of the mouse embryo by artificial nodal flow. *Nature* **418**, 96–99 (2002).

6. Y. Okada, S. Takeda, Y. Tanaka, J. C. I. Belmonte, N. Hirokawa, Mechanism of nodal flow: A conserved symmetry breaking event in left-right axis determination. *Cell* **121**, 633–644 (2005).

7. M. Abe, R. Kuroda, The development of CRISPR for a mollusc establishes the formin Lsdia1 as the long-sought gene for snail dextral/sinistral coiling. *Development* **146** (2019).

8. D. T. Grimes, R. D. Burdine, Left-right patterning: Breaking symmetry to asymmetric morphogenesis. *Trends Genet* **33**, 616–628 (2017).

9. N. Hirokawa, Y. Tanaka, Y. Okada, Left–right determination: Involvement of molecular motor KIF3, cilia, and nodal flow. *Cold Spring Harb. Perspect. Biol.* **1**, a000802 (2009).

10. R. Kuroda, A twisting story: how a single gene twists a snail? Mechanogenetics. *Q. Rev. Biophys.* **48**, 445–452 (2015).

11. R. W. Korn, Anodic asymmetry of leaves and flowers and its relationship to phyllotaxis. *Ann. Bot.* **97**, 1011–1015 (2006).

12. H. A. Allard, The ratios of clockwise and counterclockwise spirality observed in the phyllotaxy of some wild plants. *Castanea* **16**, 1–6 (1951).

13. M. S. Sousa-Baena, L. G. Lohmann, J. Hernandes-Lopes, N. R. Sinha, The molecular control of tendril development in angiosperms. *New Phytol.* **218**, 944–958 (2018).

14. H. Buschmann, A. Borchers, Handedness in plant cell expansion: a mutant perspective on helical growth. *New Phytol* (2019). https://doi.org/10.1111/nph.16034.

15. D. Reinhardt, *et al.*, Regulation of phyllotaxis by polar auxin transport. *Nature* **426**, 255–60 (2003).

16. D. H. Chitwood, *et al.*, Leaf asymmetry as a developmental constraint imposed by auxin-dependent phyllotactic patterning. *Plant Cell* **24**, 2318–2327 (2012).

17. S. J. Gerbode, J. R. Puzey, A. G. McCormick, L. Mahadevan, How the cucumber
tendril coils and overwinds. *Science* **337**, 1087–1091 (2012).

18. S. Thitamadee, K. Tuchihara, T. Hashimoto, Microtubule basis for left-handed helical
growth in Arabidopsis. *Nature* **417**, 193–196 (2002).

19. T. Ishida, Y. Kaneko, M. Iwano, T. Hashimoto, Helical microtubule arrays in a collection
of twisting tubulin mutants of Arabidopsis thaliana. *Proc. Natl. Acad. Sci. U. S. A.* **104**,
8544–8549 (2007).

20. S. C. H. Barrett, The evolution of plant sexual diversity. *Nat Rev Genet* **3**, 274–84
(2002).

21. L. K. Jesson, S. C. H. Barrett, Experimental tests of the function of mirror-image
flowers. *Biol. J. Linn. Soc.* **85**, 167–179 (2005).

22. L. K. Jesson, S. C. H. Barrett, The genetics of mirror-image flowers. *Proc. R. Soc. B-*
Biol. Sci. **269**, 1835–1839 (2002).

23. S. C. H. Barrett, A. L. M. Fairnie, The neglected floral polymorphism: mirror-image
flowers emerge from the shadow of heterostyly. *Evol. J. Linn. Soc.* **3**, kzae004 (2024).

24. L. K. Jesson, S. C. H. Barrett, The comparative biology of mirror-image flowers. *Int J*
Plant Sci **164**, S237–S249 (2003).

25. S. C. H. Barrett, Darwin's legacy: the forms, function and sexual diversity of flowers.
Philos Trans R Soc Lond B Biol Sci **365**, 351–68 (2010).

26. S. C. H. Barrett, Sexual interference of the floral kind. *Heredity* **88**, 154–159 (2002).

27. L. K. Jesson, S. C. H. Barrett, Enantiostyly: Solving the puzzle of mirror-image flowers.
Nature **417**, 707–707 (2002).

28. C. Minnaar, B. Anderson, A combination of pollen mosaics on pollinators and floral
handedness facilitates the increase of outcross pollen movement. *Curr. Biol. CB* **31**,
3180-3184.e3 (2021).

29. S. D. Johnson, J. J. Midgley, N. Illing, The enantiostylous floral polymorphism of
Barberetta aurea (Haemodoraceae) facilitates wing pollination by syrphid flies. *Ann.*
Bot. **132**, 1107–1118 (2023).

30. J. C. Manning, P. Goldblatt, A revision of Tecophilaeaceae subfam. Tecophilaeoideae
in Africa. *Bothalia Afr. Biodivers. Conserv.* **42** (2012).

31. B. R. Paudel, *et al.*, Loss of buzz pollination results in chronic pollen limitation in an
enantiostylous plant. *South Afr. J. Bot.* (in press).

32. R. Dulberger, R. Ornduff, Floral morphology and reproductive-biology of 4 species of
Cyanella (Tecophilaeaceae). *New Phytol* **86**, 45- (1980).

33. C. Kappel, C. N. Huu, M. Lenhard, A short story gets longer: recent insights into the
molecular basis of heterostyly. *J Exp Bot* **68**, 5719–5730 (2017).

34. C. N. Huu, *et al.*, Presence versus absence of CYP734A50 underlies the style-length
dimorphism in primroses. *Elife* **5**, e17956 (2016).

35. J. S. Shore, *et al.*, The long and short of the S-locus in Turnera (Passifloraceae). *New*
Phytol. **224**, 1316–1329 (2019).

36. V. Mirabet, F. Besnard, T. Vernoux, A. Boudaoud, Noise and robustness in phyllotaxis.
PLOS Comput. Biol. **8**, e1002389 (2012).

37. J. A. Nelder, R. Mead, A simplex method for function minimization. *Comput. J.* **7**, 308–
313 (1965).

38. M. A. Holland, T. Kosmata, A. Goriely, E. Kuhl, On the mechanics of thin films and
growing surfaces. *Math. Mech. Solids MMS* **18**, 561–575 (2013).

39. O. Hamant, *et al.*, Developmental patterning by mechanical signals in Arabidopsis.
Science **322**, 1650–5 (2008).

40. M. Bolger, R. Schwacke, B. Usadel, "MapMan Visualization of RNA-Seq Data Using
Mercator4 Functional Annotations" in *Solanum Tuberosum: Methods and Protocols*,
Methods in Molecular Biology., D. Dobnik, K. Gruden, Ž. Ramšak, A. Coll, Eds.
(Springer US, 2021), pp. 195–212.

41. C. C. Martinez, D. H. Chitwood, R. S. Smith, N. R. Sinha, Left-right leaf asymmetry in
decussate and distichous phyllotactic systems. *Philos. Trans. R. Soc. Lond. B. Biol.*
Sci. **371**, 20150412 (2016).

42. M. Du, E. P. Spalding, W. M. Gray, Rapid auxin-mediated cell expansion. *Annu. Rev.*
Plant Biol. **71**, 379–402 (2020).

43. N. T. Krogan, X. Yin, W. Ckurshumova, T. Berleth, Distinct subclades of Aux/IAA genes
are direct targets of ARF5/MP transcriptional regulation. *New Phytol.* **204**, 474–483
(2014).

44. A. K. Spartz, *et al.*, The SAUR19 subfamily of SMALL AUXIN UP RNA genes promote
cell expansion. *Plant J* **70**, 978–990 (2012).

45. A. K. Spartz, *et al.*, SAUR Inhibition of PP2C-D phosphatases activates plasma
membrane H+-ATPases to promote cell expansion in Arabidopsis. *Plant Cell* **26**, 2129–
2142 (2014).

46. A. Peaucelle, *et al.*, Pectin-induced changes in cell wall mechanics underlie organ
initiation in Arabidopsis. *Curr Biol* **21**, 1720–6 (2011).

47. G.-Q. Huang, *et al.*, A fasciclin-like arabinogalactan protein, GhFLA1, is involved in
fiber initiation and elongation of cotton. *Plant Physiol.* **161**, 1278–1290 (2013).

48.   W. H. Bryan, K. H. Newcomb, Stimulation of pectin methylesterase activity of cultured
tobacco pith by indoleacetic acid. *Physiol. Plant.* **7**, 290–297 (1954).

49.   A. Andres-Robin, *et al.*, Evidence for the regulation of gynoecium morphogenesis by
ETTIN via cell wall dynamics. *Plant Physiol.* **178**, 1222–1232 (2018).

50.   A. Andres-Robin, *et al.*, Immediate targets of ETTIN suggest a key role for pectin
methylesterase inhibitors in the control of Arabidopsis gynecium development. *Plant*
Signal. Behav. **15**, 1771937.

51.   L. Li, *et al.*, RALF1 peptide triggers biphasic root growth inhibition upstream of auxin
biosynthesis. *Proc. Natl. Acad. Sci. U. S. A.* **119**, e2121058119 (2022).

52.   L. K. Jesson, J. Kang, S. L. Wagner, S. C. H. Barrett, N. G. Dengler, The development
of enantiostyly. *Am J Bot* **90**, 183–195 (2003).

53.   W. A. Charlton, "Pendulum symmetry" in *Symmetry in Plants*, Series in Mathematical
Biology and Medicine., (WORLD SCIENTIFIC, 1998), pp. 61–87.

54.   L. D. Harder, S. C. H. Barrett, Mating cost of large floral displays in hermaphrodite
plants. *Nature* **373**, 512–515 (1995).