Development of capsaicin derivatives as prohibitin ligands to modulate the Aurora kinase A/PHB2 interaction in cancer cells

Amel Djehal[a,b,‡], Claire Caron[c,‡], Deborah Giordano[d], Valentina Pizza[c,d], Kimberley Farin[b], Angelo Facchiano[d], Laurent Désaubry[b,*] and Giulia Bertolin[c,*].

[a]Higher National School of Biotechnology of Constantine, Constantine 25100, Algeria

[b]Laboratory of Regenerative Nanomedicine, Center of Research and Biomedicine, University of Strasbourg, 67000 Strasbourg, France

[c]CNRS, Univ Rennes, IGDR [(Institut de Génétique et Développement de Rennes)] - UMR 6290, F-35000 Rennes, France

[d]National Research Council, Institute of Food Science, via Roma 64, 83100 Avellino, Italy

‡ Contributed equally to this work

*To whom correspondence should be addressed: desaubry@unistra.fr (L.D.), giulia.bertolin@univ-rennes.fr (G.B.).

Keywords: PHB2; Aurora kinase A/AURKA; capsaicin; ligands; mitochondria; FRET/FLIM microscopy; molecular docking; cancer.

Abstract

Aurora kinase A/AURKA is a serine/threonine kinase overexpressed in a variety of solid and hematological malignancies. In the last decades, clinical trials aiming to counteract the overexpression of AURKA turned out to be largely unsuccessful. Meanwhile, recent discoveries pointed at new functions of AURKA at the subcellular level, including at mitochondria. At this location, AURKA induces organelle clearance by mitophagy by acting in complex with the mitophagy mediator LC3, and its inner mitochondrial membrane receptor PHB2. The natural polyphenol xanthohumol was shown to act as a PHB2 ligand, altering the interaction between AURKA and PHB2 and restoring mitochondrial functions in cancer cells. However, its chemical nature prevents its broader use as an anti-cancer agent.

Using Förster's Resonance Energy Transfer/Fluorescence Lifetime Imaging Microscopy (FRET/FLIM) in live breast cancer cells, we here explore the effects of alternative PHB ligands in altering the proximity between AURKA and PHB2. Among the already-available compounds, we found that the pungent natural product capsaicin partially alters the AURKA/PHB2 protein-protein proximity. We then synthesized 16 novel capsaicin analogues to enhance the effects of capsaicin. We found that replacing the long hydrophobic acyl moiety by a butyryl one increases the AURKA/PHB2 interaction. Among the capsaicin derivatives carrying this modification, we uncover that compound 12 and 13 enhance the AURKA/PHB2 proximity. Molecular docking approaches corroborate FRET/FLIM data, and we visualize compound 12 and 13 in complex with AURKA, PHB2 and LC3. We show that compound 12 and 13 stabilize the AURKA/PHB2 interaction, and that they can bind to the inhibitory pocket of PHB2 and to the AURKA active site.

Together, our data indicate that compound 12 and 13 are promising PHB ligands acting on the AURKA/PHB2 interaction. Through their molecular action on either AURKA or PHB2, they may provide the basis for the development of new anticancer drugs targeting the mitochondrial functions of AURKA.

Highlights

- The cancer-related kinase Aurora A/AURKA interacts with the mitochondrial protein PHB2 to induce mitophagy in cancer cells.

- Among the available PHB ligands, capsaicin mildly lowers the AURKA/PHB2 interaction.

- Screening of 16 newly-synthesized capsaicin derivatives show differential effects on the AURKA/PHB2 interaction.

- Compound 12 and 13 stabilize the AURKA/PHB2 interaction.

- Compound 12 and 13 bind to the AURKA active site and to the PHB2 inhibitory pocket.

Introduction

Mitochondria are in constant equilibrium between turnover and biogenesis. To be able to metabolically sustain an uncontrolled cell proliferation in cancer cells, mitochondria are frequently hijacked to produce significant amounts of ATP[1]. In parallel, damaged mitochondria can be eliminated through a process called mitophagy[2,3]. Mitophagy is controlled by several proteins, including the scaffold protein Prohibitin-2 (PHB2) which binds to the autophagy mediator LC3[4]. PHB2 and its homolog Prohibitin-1 (PHB1) form a heterodimeric complex located on the Inner Mitochondrial Membrane (IMM), and constitute a hub for many signaling pathways beyond mitophagy in cancer cells[5].

Recently, the overexpression of the cancer-related protein Aurora kinase A/AURKA was shown to induce mitophagy by functionally interacting with PHB2[6]. AURKA regulates a wide variety of mitotic and non-mitotic functions at different subcellular compartments, including mitochondrial dynamics and ATP production[7–11]. We previously showed that the AURKA-dependent phosphorylation of PHB2 on Ser39 induces the formation of an AURKA-LC3-PHB2 tripartite complex[6]. We also showed that this tripartite complex triggers mitophagy in a PARK2/Parkin-independent manner[6]. AURKA-dependent mitophagy is used to eliminate defective mitochondria in cancer cells, while selecting a population of hyperfunctional organelles with high ATP production levels. We also found that the natural polyphenol xanthohumol acts as a PHB2 ligand on the inner membrane of mitochondria to destabilize the AURKA-LC3-PHB2 tripartite complex. By blocking the interaction between AURKA and LC3, treatment with xanthohumol results in mitophagy inhibition. Under these conditions, the AURKA-dependent mitochondrial ATP overproduction is also abolished, thereby linking an effective mitophagy and the capacity of mitochondria to increase their ATP production rates[6]. However, the polyphenolic structure of xanthohumol hampers its development as an anticancer agent.

As part of our ongoing programs on the development of new PHB ligands[12–16], we here screen already-available alternative PHB ligands in their capacity to alter the protein-protein interaction between AURKA and PHB2. We then optimize the activity of the most promising hit – capsaicin - to obtain 16 novel analogues targeting the AURKA/PHB2 interaction. Last, we show that these ligands differentially affect the capacity of AURKA to interact with PHB2. Using a combination of quantitative fluorescence microscopy and molecular docking, we identify two new PHB ligands dramatically reinforcing the interaction between AURKA and PHB2. By binding both to the AURKA active site and to the inhibitory pocket of PHB2, compound 12 and 13 could become relevant candidates for follow-up studies.

Results

Screening of available PHB ligands identifies capsaicin as a compound lowering the interaction between AURKA and PHB2.

As an alternative to xanthohumol[6], we selected three PHB ligands with different chemical structures and with different functional consequences on PHB2 signaling pathways such as FL3, fluorizoline and capsaicin[5] (Figure 1).

Figure 1: Structure of the PHB ligands xanthohumol, FL3, fluorizoline and capsaicin examined for their effects on the AURKA/PHB2 interaction in breast cancer cells.

We then compared the capacity of FL3, capsaicin and fluorizoline to alter the protein-protein proximity between AURKA and PHB2 using Förster's Resonance Energy Transfer coupled to Fluorescence Lifetime Imaging Microscopy (FRET/FLIM) in live cells[6,17,18]. In these experiments, AURKA was fused to GFP and was used as a FRET donor, while PHB2 was fused to mCherry and was used as a FRET acceptor. FRET was represented using ΔLifetime, which corresponds to the net difference between the lifetime of AURKA-GFP in the absence (AURKA-GFP + Empty vector, hereby donor-only condition) and in the presence of the acceptor (AURKA-GFP + PHB2-mCherry) (Figure 2). MCF7 breast cancer cells expressing AURKA-GFP in the presence or absence of PHB2-mCherry were incubated for 6h with each of the three compounds, or with the vehicle (DMSO) prior to FRET measurements.

Treatment with FL3 did not significantly impact the interaction between AURKA and PHB2 in FRET/FLIM analyses. In contrast, treatment with capsaicin or fluorizoline brought the ΔLifetime of AURKA-GFP in the presence of PHB2-mCherry closer to that of their respective donor-only counterparts (Figure 2). This indicates that both compounds significantly lower the interaction between AURKA and PHB2. However, fluorizoline also displayed a significant green autofluorescence that made image-based analyses difficult. Thus, we selected capsaicin as a promising compound for the development of novel PHB ligands that could differentially modulate the AURKA/PHB2 interaction. Capsaicin did not promote a nuclear localization of PHB2 in MCF7 cells, in contrast with a previous report in HeLa cells[19]. Instead, our FRET/FLIM analyses corroborate previous evidence, showing that PHB2 could only be located at mitochondria in MCF7 cells[6].

Our data show that capsaicin mildly lowers the protein-protein interaction between AURKA and PHB2 at mitochondria. We next synthesized a series of new capsaicin derivatives, and we evaluated their effects on the AURKA/PHB2 interaction.

Figure 2: FRET/FLIM analyses on MCF7 cells expressing AURKA-GFP with an empty vector (donor-only condition), or with PHB2-mCherry. Cells were treated with vehicle (DMSO), capsaicin, FL3 or fluorizoline. Pseudocolour scale: pixel-by-pixel ΔLifetime. Graph: corresponding ΔLifetime quantifications performed in the dotted area. *n* = 10 cells per condition of one representative experiment (of three). Data extend from the min to the max. Dots indicate individual cells. Scale bar: 10 μm. *P<0.01 against each corresponding "AURKA-GFP + Empty vector" condition.

Chemical synthesis of novel capsaicin derivatives

After comparing several of methods of acylation, we found that Schotten-Baumann acylation allows the amine of vanillylamine **1a** to be selectively acylated without affecting the phenol as described by Short and coll. (Figure 3A)[20]. With this process, we synthesized 16 new capsaicin derivatives potentially behaving as PHB ligands (Figure 3B). After the synthesis was completed, we then screened these compounds to explore whether they could alter the AURKA/PHB2 interaction at mitochondria.

Figure 3: Synthesis reactions (**A**) and structures (**B**) of capsaicin analogues used in the study. Bold numbers below each structure in (B) indicate the number of each compound tested.

Capsaicin derivatives differentially affect the interaction between AURKA and PHB2.

To evaluate the effects of the 16 newly-synthesized capsaicin derivatives (Figure 3B) towards the AURKA/PHB2 interaction, MCF7 cells co-expressing AURKA-GFP and PHB2-mCherry were treated for 6h with each capsaicin derivative or with the DMSO control compound. Their effects on the interaction between AURKA and PHB2 were then evaluated using FRET/FLIM. The dynamic range of a GFP-mCherry chimeric tandem construct shows a maximum of 250 psec of ΔLifetime when compared to a GFP donor-only condition[17]. Given that the AURKA-GFP/PHB2-mCherry pair in the DMSO condition shows a mean ΔLifetime of 120-150 psec *per se*, we explored whether the 16 capsaicin derivatives could induce variations to these mean ΔLifetime values in cells co-expressing AURKA-GFP and PHB2-mCherry.

To this end, we calculated the mean ΔLifetime difference from the AURKA-GFP/PHB2-mCherry pair in the DMSO condition for each novel capsaicin derivative (Table 1). Variations were considered mild when a given compound induced changes in the AURKA-GFP ΔLifetime of +/- 30-50 additional psec compared to what observed in the AURKA-GFP/PHB2-mCherry pair in the DMSO condition. This corresponds to 12-25% of the maximum ΔLifetime response

of a GFP-mCherry tandem[21]. Alternatively, variations were considered significant when a given compound induced changes to the AURKA-GFP ΔLifetime which were superior to +/- 50 additional psec compared to what observed in the DMSO condition. This is more than 25% of the maximum ΔLifetime response of a GFP-mCherry tandem[21].

The replacement of the acyl moiety of capsaicin by various linear and unsaturated fatty acids in compounds 1-7 (6 to 18 carbons) was detrimental to the FRET effect. Indeed, compound 1, 4, 6 and 7 showed no effect on AURKA-GFP ΔLifetime when AURKA-GFP and PHB2-mcherry were co-expressed. Compound 2, 3 and 5 mildly lowered the AURKA/PHB2 interaction, but with a standard deviation superior to that of capsaicin (Table 1).

Compound	Mean Δlifetime difference from DMSO (psec)	S.D. (psec)	Effect on the AURKA/PHB2 interaction
Capsaicin	-41.92	14.1	Mildly lowered
1	-23.17	131.18	No significant variation
2	-48.26	85.25	Mildly lowered
3	-40.44	145.43	Mildly lowered
4	-2.93	109.07	No significant variation
5	-67.92	94.85	Mildly lowered
6	21.12	68.6	No significant variation
7	8.76	44.84	No significant variation
8	47.02	47.38	Mildly increased
9	48.96	63.07	Mildly increased
10	44.06	49.07	Mildly increased
11	74.89	77.81	Increased
12	77.09	79.45	Increased
13	66.35	13.97	Increased
14	12.87	55.99	No significant variation
15	0.77	24.93	No significant variation
16	51.92	63.9	Increased

Table 1. Effect of capsaicin derivatives on the AURKA/PHB2 interaction. Mean ΔLifetime differences and standard deviation (S.D.) of each capsaicin derivative when compared to the vehicle compound, in MCF7 cells co-expressing AURKA-GFP and PHB2-mCherry. Mild variations are between 30 and 50 psec, while variations are considered significant when they are more than 50 psec from the DMSO condition. Results are means from three independent experiments, with n = 10 cells per condition in each replicate.

Unexpectedly, the introduction of 5 or 6-membered ring in the hydrophobic acyl chain resulted in an opposite pharmacological activity. Indeed, the constrained analogues 8, 9 and 10 mildly increased the interaction between AURKA and PHB2. Of note, these compounds also show a lower degree of experimental variability than the previously-tested compounds in our FRET/FLIM analyses (Table 1). Similarly to compounds 8-10, *N*-vanillylgeranamide/compound 11 displayed some rigidification as well. However, this compound promoted AURKA/PHB2 interaction to a greater extent than the previous analogues (Table 1). Shortening the hydrophobic chain into a crotonic group as in compound 12 also induced significant changes of more than 50 psec in the AURKA-GFP ΔLifetime, thus increasing the interaction between AURKA and PHB2 compared to that observed in the DMSO condition (Figure 4A, C and Table 1). Nevertheless, both compound 11 and 12 displayed a high standard deviation. Switching to the more flexible butyryl group as in compound 13 was another suitable strategy to significantly increase the AURKA/PHB2 interaction. Furthermore, compound 13 displayed a lower standard deviation than compound 11 or 12 (Figure 4B, C and Table 1). Thus, we chose to modify this analogue to explore the vanillyl moiety.

Figure 4: FRET/FLIM analyses on MCF7 cells co-expressing AURKA-GFP (donor) and PHB2-mCherry (acceptor). Cells were treated with vehicle (DMSO), 12 (**A**) or 13 (**B**). Pseudocolour scale: pixel-by-pixel ΔLifetime. Dotted area: PHB2-rich mitochondrial area.

The methylation of the phenolic moiety and leading to compound 14 and 15 left the interaction between AURKA and PHB2 unaltered (Table 1). On the contrary, its replacement by an aniline as in compound 16 increased the interaction between the two proteins (Table 1). This indicates the requirement for a H-bond donor in this position.

To exclude that the capsaicin derivatives could significantly perturb the lifetime of AURKA-GFP *per se,* each compound was evaluated in the donor-only condition where AURKA-GFP was co-expressed with an empty vector (Supplementary Table 1). Previous reports demonstrated that a ΔLifetime difference below 50 psec corresponds to the standard deviation of our FLIM setup[17,18]. Therefore, a difference in the AURKA-GFP ΔLifetime above 50 psec may indicate that the tested ligand intrinsically affects the lifetime of AURKA-GFP, and this beyond the FRET effect observed when AURKA-GFP is co-expressed with PHB2-mCherry. Globally, the majority of the compounds tested showed a non-significant variation of ΔLifetime in the donor-only condition. Nevertheless, we observed that compound 16 shows a slightly higher variability than the other derivatives under these conditions. This suggest that the differences in ΔLifetime observed upon the co-expression of AURKA-GFP and PHB2-mCherry (Table 1) in cells treated with compound 16 could be partially independent of the FRET effect, and could be explained with an intrinsic effect of the compound 16 on the donor lifetime. Therefore, we decided not to use compound 16 for follow-up analyses in living cells.

Overall, our FRET/FLIM results show that the newly-generated capsaicin derivatives differentially modulate the interaction between AURKA and PHB2 at mitochondria in live cells. While the majority had either no effect or a mild effect on this interaction, we determined that compound 11, 12 and 13 significantly increase the AURKA/PHB2 protein-protein proximity. Among these, compound 12 and 13 increase the AURKA/PHB2 interaction with a low standard deviation. This makes of these capsaicin derivatives the most promising hits among the 16 compounds here generated. However, insights at the molecular level are needed to explain why compound 12 and 13 increase the AURKA/PHB2 interaction.

Capsaicin and compound 5 induce a molecular clash lowering the AURKA/PHB2 interaction.

After observing that compound 12 and 13 robustly increase the AURKA/PHB2 interaction, we sought to uncover their binding mechanisms to AURKA and PHB2. To this end, we used molecular docking approaches and we compared a total of four molecules with opposite FRET/FLIM efficiencies. On one side, we selected capsaicin and complex 5, which mildly lower the AURKA/PHB2 interaction. On the other side, we selected compound 12 and 13, which show the highest FRET/FLIM efficiencies.

First, a reverse docking approach was used to discover possible interesting protein targets for all the selected compounds. Reverse docking results performed by GalaxySagittarius-

AlphaFold (Supplementary File 1) highlighted that the four compounds tested were predicted to bind AURKA, even if with different scores. Capsaicin and compound 5 could bind AURKA with a Galaxy best final score of 0.573 and 0.498, respectively. Conversely, compound 12 and 13 could bind AURKA with higher Galaxy best final scores, which were of 0.765 and 0.846, respectively. Interesting, the key capsaicin target Transient Receptor Potential cation channel subfamily V member 1 (TRPV1)[22,23] could be retrieved only in the presence of capsaicin and compound 5 for the "Binding compatibility prediction" search mode (Table 2). TRPV1 could be retrieved in the list of targets of compound 12, but only with the "Similarity combined prediction" search mode. TRPV1 also showed a score significantly lower than the score of AURKA, and comparable to that of MAP1LC3 which occupies the lowest position on the list (Table 2). This may suggest a higher specificity of compound 12 and 13, that could result in a superior affinity and/or selectivity for AURKA.

| Compound | TARGET | | | | | | | |
| | GALAXY SIMILARITY AlphaFold | | | | GALAXY COMPATIBILITY AlphaFold | | | |
	rank	uniprot_id	name	score	rank	uniprot_id	name	score
Capsaicin	20	O14965	AURKA	0,498	63	Q8NER1	TRPV1	0,243
	321	Q8NER1	TRPV1	0,35	175	O14965	AURKA	0,222
Compound 5	24	O14965	AURKA	0,573	457	Q8NER1	TRPV1	0,238
	552	Q8NER1	TRPV1	0,363	780	O14965	AURKA	0,211
Compound 12	7	O14965	AURKA	0,765	10	O14965	AURKA	0,273
	277	Q8NER1	TRPV1	0,275				
	834	Q9H492	MAP1LC3A	0,211				
Compound 13	4	O14965	AURKA	0,846	7	O14965	AURKA	0,273

Table 2. Extract of GalaxySagittarius-AF reverse docking results. For each compound, the position (rank) and the score of the proteins of the tripartite complex or of the capsaicin receptor TRPV1 are reported if the target is present among all the hits retrieved.

Second, we analyzed the affinity of capsaicin and of compound 5, 12 and 13 for AURKA and PHB2. To this end, we performed rigid and flexible docking analyses to detect possible relevant binding sites on the activated (pThr288) AURKA structure bound to ADP and Mg^{2+}, on the PHB2 model, and on the tripartite AURKA/PHB2/LC3 complex model (Supplementary Table 2). The tripartite AURKA/PHB2/LC3 complex model was built by exploiting rigid protein-protein docking and homology modelling procedures. The best model obtained presents the Ser39 of PHB2[6] oriented towards the AURKA active site (Supplementary Figure 1). This model also shows the region of LC3 involved in recognition of the LC3-Interacting Region (LIR) binding motif on PHB2 facing the LC3 binding motif YQRL (residues 121-124), and towards the putative LC3 binding motif FNRI (residues 52-55) (Supplementary Figure 1). Binding energy is equal to -12.6 kCal/mol for the AURKA/PHB2 interaction, -10.8 kCal/mol for PHB2/ LC3 interaction, and -1.4 kCal/mol for the AURKA/LC3 interaction. This last value could be due to the very small interface area between AURKA and LC3, which is substantiated by low FRET/FLIM efficiencies in a previous report[6]. Of note, the present model lacks the first 125

residues of AURKA, which are a well-known disordered region. AURKA, LC3 and PHB2 present Z-scores equal to -7.1, -6.6, and -5.91, respectively, in line with the original structures having Z-score values of -7.06, and -6.54 for the crystallographic structure of AURKA and LC3, respectively, and -5.75 for the PHB2 model. The Ramachandran plot of the trimeric complex model obtained presents the following values: 97.8% of dihedral angles in permitted regions of the plot, 4.3% in regions allowed and 0% in generously allowed and disallowed regions.

After generating the model of the tripartite complex (Supplementary Figure 1), we then evaluated the effects of capsaicin and compound 5 on this model. Docking results for compound 5 and capsaicin highlighted a similar behavior of these two compounds (Table 3).

Protein	Ligand	Docking type[a]	Lowest Binding Energy (Kcal/mol)	Mean Binding Energy (Kcal/mol)	N. of poses in cluster[b]	Ki	Pocket of Interaction[c]
AURKA	Capsaicin	Focused and flexible	-7.88	-7.88	1	1.66 µM	K143-F144-H176-Q177-**D256-**K258-E260-**N261-**D274-W277-Y288-L289-**ADP** (active site)
			-7.10	-6.74	5	6.21 µM	K141-G142-K143-Q177-L178-E181-E260-D274-G276-W277-**ADP** (active site)
	Compound 5	Focused and flexible	-8.83	-8.28	4	336.01 nM	K143-F144-K162-L164-V174-Q177-L178-E181-D274-G276-W277-L289-A290-G291-**ADP** (active site)
			-8.37	-7.51	5	738.07 nM	K143-F144-K162-L164-L178-Q177-E181-D274-G276-W277-L289-A290-G291-L293-L296-I301-**ADP** (active site)
Tripartite AURKA/PHB2/LC3 complex	Capsaicin	Focused and rigid	-7.62	-6.77	3	2.60 µM	AURKA: K141-T217-Y219-K258-E260-G291-T292-**ADP-Mg²⁺**; PHB2: V36-S39-F51-F70-R71-Q76
			-7.01	-6.55	4	7.30 µM	AURKA: K143-F144-K171-A172-Q177-**D256**-D274-

							G276-W277 **ADP-Mg²⁺**; PHB2: A31-V32-G35-W74
	Compound 5	Focused and rigid	-7.78	-7.78	1 (+1)	6.28 µM	AURKA: K141-K143-Y219-K258-E260-W277-G291-T292-**ADP-Mg²⁺**; PHB2: V32-G35-V36-S39-Q59-F70-I72-Q76
			-7.10	-5.96	6 (+10)	1.98 µM	AURKA: F144-E168-K171-A172; PHB2: A31-Y34-E38-R71-W74-P73-Y77; LC3: R14-T54
PHB2	Capsaicin	Focused and rigid	-5.91	-5.05	23	46.93 µM	P78-I79-I80-**D82**-L126-**D127**-Y128-E129 (inhibitor site)
	Compound 5	Focused and rigid	-5.29	-4.39	5	132.16 µM	H47-P78-I79-I80-Y81-**D82**-R86-L126-**D127**-Y128-E129-E130 (inhibitor site)

Table 3: Docking simulation results of capsaicin and compound 5.

[a]: All docking simulations were performed allowing flexibility to the ligand. The annotation "flexible" refers to a docking focused on the AURKA active site, and allowing flexibility not only to the ligand but also to the D256-N261-D274-W277 residues.

[b]: Number of different conformations detected for each ligand in the pocket of interaction, and it is calculated on 100 total poses.

[c]: Residues underlined are involved in H-bond with the ligand, and residues relevant for binding or catalysis are bolded.

Blind and focused, rigid and flexible docking analyses highlighted that compound 5 and capsaicin bind a similar area in the active site of AURKA both with two different possible conformations (Supplementary Table 2). Compound 5 shows interaction with ADP and Mg²⁺ in its best conformation, and binding the AURKA active form. However, it should be noted that only few poses are present in the two clusters, potentially indicating a low selectivity for this area. Both conformations are comparable to the best energetic pose of capsaicin (Supplementary figure 2). While compound 5 binds exclusively to the cofactors, capsaicin appears to interact with the cofactor and with the active site residue D256. Interestingly, capsaicin binds the AURKA active site with a very low binding energy equal to -7.88 kCal/mol and a constant of inhibition (Ki) of 1.66 µM, and compound 5 does so with an energy value of -8.83 kCal/mol and a Ki of 336.01 nM. All the poses detected suggest a possible inhibitory role of these molecules on the AURKA active site. With this information, we sought to visualize possible sterical clashes due to the presence of the ligands. Clashes are conformations

Figure 5: Molecular docking of capsaicin and of compound 5, 12 and 13 in complex with the tripartite AURKA/PHB2/LC3 complex model. In all panels, AURKA is represented in grey,

PHB2 in cyan and LC3 in orange. Sticks highlight the N-terminal region of PHB2, D256 and ADP in AURKA. (**A**) Binding poses of capsaicin (green sticks and surface) and of compound 5 (pink sticks and surface) to the monomeric active form of AURKA. The overlap shows a clash among the ligands and the N-terminal region of PHB2. (**B**) Results of the docking analysis performed on the tripartite complex with compound 5. In purple, the best energy binding pose detected is shown, while in pink the representative pose of the most populated cluster is shown. These two poses indicate that compound 5 is forced to move from the pose detected on AURKA alone. (**C**) Results of the docking analysis performed on the tripartite complex with capsaicin. Light green represents the best energy binding pose detected, while the representative pose for the most populated cluster detected is shown in dark green. Capsaicin is forced either to move from the pose detected on AURKA alone, or to change its orientation in the pocket. (**D**) Superposition of the conformations of compound 12 and 13 bound to AURKA alone, with the results of the docking analysis performed on the tripartite complex. Yellow sticks and surface show compound 12 in its AURKA-binding conformation, and in orange the ones detected for the tripartite complex. Cyan sticks and surface show compound 13 in its AURKA-binding conformation. Violet and blue show the binding pose of compound 13 with the best binding energy detected for the tripartite complex and the one representative of the most populated cluster, respectively.

occurring when two atoms, which are not covalently bonded to each other, are found to be impossibly close to each other. All the structures of the monomeric proteins bound to the ligands were superimposed on the model of the tripartite complex. The overlap highlighted a possible clash between the ligand – either capsaicin or compound 5 – and PHB2 (Figure 5A), suggesting that the binding of either ligand to AURKA could weaken the AURKA/PHB2 interaction.

To verify this hypothesis, additional docking simulations were performed on the tripartite complex model. Docking was focused at the interface areas of the three proteins. As expected, compound 5 was found to change its localization on AURKA, with a further 1 kCal/mol loss of its binding energy and reaching values of -7.78 or of -7.10 kCal/mol. Two binding sites were observed to reach the best binding energy with compound 5, and involve the AURKA K141-K143-Y219-K258-E260-W277-G291-T292 residues, the cofactors ADP-Mg2+ and PHB2 V32-G35-V36-S39-Q59-F70-I72-Q76 residues. They also involve AURKA F144-E168-K171-A172, PHB2 A31-Y34-E38-R71-W74-P73-Y77 and LC3 R14-T54 residues (Figure 5B). Similarly to compound 5, capsaicin is able to interact with cofactors in the active site (Figure 5C). However, capsaicin is forced to move into a more dislocated binding site with the best binding energy of -7.62 kCal/mol. Alternatively, capsaicin is forced to adopt an alternative conformation similar to the second one already observed when binding to AURKA, and showing the lowest binding energy equal to -7.10 kCal/mol (Figure 5C).

We then analyzed the effects of capsaicin and of compound 5 on PHB2. Docking analyses performed on the PHB2 model were focused on the most organized tertiary structure area of this protein. Results reveal a preference of capsaicin and compound 5 for the inhibitory site of the protein, originally detected when studying the PHB2 inhibitor YL-939[24]. Both capsaicin and compound 5 appear to bind to PHB2 by interacting with the same residues, P78-I79-I80-D82-L126-D127-Y128-E129. The binding energies of these compounds with PHB2 are 5.91KCal/mol and -5.29 kCal/mol for capsaicin and compound 5. They are higher than the binding energies measured for the two compounds in complex with AURKA, indicating that

capsaicin and compound 5 could bind PHB2 with less affinity than for AURKA. However, it should also be noted that the number of poses in the cluster are higher for capsaicin in complex with PHB2 than with AURKA. Despite its lower binding affinity towards PHB2, the number of poses still indicates a good selectivity of capsaicin for the inhibitory site of PHB2.

Overall, capsaicin and compound 5 appear to move from the best energy binding pose to a second pose on AURKA alone. These compounds have a steric hindrance that induces a steric clash at the interface between AURKA and PHB2, and potentially lowering or destabilizing the interaction between these two proteins. The molecular docking data thus substantiate the mild effect of capsaicin and of compound 5 in lowering the AURKA/PHB2 interaction observed with FRET/FLIM experiments.

Compound 12 and 13 create a molecular bridge between AURKA and PHB2, and they bind to the AURKA active site and the PHB2 inhibitory pocket.

Given the previous results on capsaicin and compound 5, we performed similar binding and docking analyses for compound 12 and 13.

Compound 12 and 13 reach the best binding energy with the active conformation of AURKA, and they appear to specifically bind the substrate-binding site of the kinase (Table 4). Differently from compound 5, both molecules appear to interact with the D256 residue of the active site and with the cofactors. When observing the structure of the two compounds bound to the tripartite complex, no sterical clash between AURKA and PHB2 is observed (Figure 5D). On the contrary, this superposition rather suggests a possible action of compound 12 and 13 as a bridge between AURKA and PHB2. The docking performed on the complex also revealed that compound 12 and 13 show a preference for the AURKA binding site (Figure 5D). The binding site is preserved and no clashes occur to alter the binding. The two compounds appear to adapt their orientation in the pocket to preserve the binding and stabilize the complex. Compound 12 preserves its interaction with AURKA Q177, D274, W277 residues, ADP and Mg^{2+}. Compound 13 preserves its interaction with AURKA Q177, E181, D256, K258, N261, D274, W277 residues, and with ADP and the two Mg^{2+} ions. Furthermore, the two compounds show an interaction with PHB2. Compound 12 shows interaction with V32, and compound 13 shows interaction with S39, F51, F70, R71, I72, and Q76 in one of the two alternative binding conformations. Binding energies for the compounds binding to the complex or to AURKA alone are comparable, with -6.67 kCal/mol and -6.73 kCal/mol for compound 12 bound to the tripartite complex or AURKA alone, and with -6.74 kCal/mol and -6.77 kCal/mol for compound 13 bound to the complex or to AURKA alone. This is additional evidence that the binding site does not undergo changes with these two compounds. In addition, the higher number of poses in the cluster suggests that the selectivity of these compounds for the AURKA active is higher than what measured for capsaicin and compound 5.

Protein	Ligand	Docking type[a]	Lowest Binding	Mean Binding	N. of poses in cluster[b]	Ki	Pocket of Interaction[c]

			Energy (kCal/mol)	Energy (kCal/mol)			
AURKA	Compound 12	Blind and rigid	-6.73	-6.30	6 (+11)	11.58 µM	F144-L164-L169-Q177-L178-E181-**D256**-K258-N261-D274-W277-**ADP-Mg²⁺-Mg²⁺**(active site)
	Compound 13	Blind and Rigid	-6.77	-5.84	9 (+9)	10.86 µM	V74-L169-Q177- L178-E181-**D256**-K258-N261-D274-W277-**ADP-Mg²⁺ -Mg²⁺** (active site)
Tripartite AURKA/PHB2/LC3 complex	Compound 12	Focused and Rigid	-6.67	-5.85	3 (+10)	12.83 µM	AURKA: K143-V174-Q177-D274-W277; **ADP-Mg²⁺**; PHB2: V32
	Compound 13	Focused and Rigid	-6.74	-6.45	6	11.46 µM	AURKA: K143-F144-K162-Q177-E181-**D256**-K258-N261-D274-G276-W277-**ADP-Mg²⁺ -Mg²⁺**
			-6.51	-6.19	7(+9)	16.96 µM	AURKA: K141-Y219-K258-N261-D274-E260-**ADP-Mg²⁺;** PHB2: S39-F51-F70-R71-I72-Q76
PHB2	Compound 12	Focused and Rigid	-5.55	-4.81	20	85.40 µM	I80-Y81-**D82**-R86-L126-**D127**-Y128-E129-E130 (inhibitor site)
	Compound 13	Focused and Rigid	-4.92	-4.68	17 (+5 +10)	248.14 µM	I80-Y81-**D82**-R86-R88-L126-**D127**-Y128-E129-E130 (inhibitor site)

Table 4. Docking simulation results of compound 12 and 13.
[a]: All docking simulations were performed allowing flexibility to the ligand. The annotation "rigid" refers to protein residues.
[b]: The number of poses in cluster indicates the number of different conformations detected for each ligand in the pocket of interaction, and it is calculated on a sampling of 100 total poses.
[c]: Residues underlined are involved in H-bond with the ligand, and residues relevant for binding or catalysis are bolded.

As performed for capsaicin or compound 5, we explored the effects of compound 12 and 13 on PHB2 (Table 4). Both compounds appear to bind the inhibitory site of PHB2, as they establish an interaction with the D82-I80-D127 binding site of the PHB2 inhibitor YL-939[24]. The two compounds bind to this site with comparable mean binding energy (-4.81 kCal/mol, and -4.68 kCal/mol for compound 12 and 13, respectively). The number of poses in each docking cluster is higher for compound 12 and 13 than for compound 5. This supports a higher selectivity of the former compounds for the inhibitory pocket than of the latter. The binding energies detectable are lower than those obtained for the AURKA active site. Despite this, the selectivity for the inhibitory pocket of PHB2 is higher, with almost 20 poses out of 100 detected in the same binding area.

Overall, the conformation of compound 12 and 13 bridges AURKA and PHB2 and brings them close to each other. These compounds potentially strengthen and/or stabilize the interaction between these two proteins. In addition, docking analyses show that these two compounds can not only bind to the AURKA active site, but also to the PHB2 inhibitory pocket. Therefore, molecular docking data thus substantiate the effect of compound 12 and 13 in strengthening the AURKA/PHB2 interaction observed with FRET/FLIM experiments.

In conclusion, our results determined that capsaicin is a pharmacological component altering the AURKA/PHB2 protein-protein proximity. Newly-generated capsaicin derivatives showed differential effects on the capacity of AURKA to interact with PHB2, and they either lower or strengthen the interaction in FRET/FLIM experiments. Molecular docking analyses revealed that capsaicin derivatives stabilizing the AURKA/PHB2 proximity can bind to the AURKA/PHB2/LC3 tripartite complex, such as compound 12 or 13. These compounds target the AURKA active site and the PHB2 inhibitory pocket, resulting in enhanced binding features compared to molecules lowering the AURKA/PHB2 interaction. We propose that capsaicin derivatives stabilizing the AURKA/PHB2 interaction are promising novel inhibitors targeting the mitochondrial roles of AURKA.

Discussion

Aurora kinase A/AURKA is a hallmark of epithelial and hematological cancers, where it is frequently overexpressed both at the mRNA or at the protein levels[8]. It is known that AURKA has multiple subcellular locations, including mitochondria[9,10].

We here identified capsaicin as a PHB2 ligand that lowers the AURKA/PHB2 interaction at mitochondria. This compound was previously shown to induce a dissociation between PHB and the Adenine Nucleotide Translocator 2 (ANT2, SLC25A5). Proteomics studies suggest that the auto/mitophagy modulator LC3 may interact with ANT2 and in our previous study, we identified ANT2 as an interactor of AURKA using MS/MS-based approaches[9]. In addition, we previously observed that AURKA interacts with LC3 in a PHB2-dependent manner[6]. This raises the interesting possibility that PHB2, AURKA ANT2 and LC3 may form a quaternary complex in mitochondria.

Capsaicin is known to be cytotoxic in a wide range of cancer cell lines. Beyond PHB2[19], these effects have been attributed to the action of capsaicin on several targets. These include the neuronal vanilloid receptor TRPV1[22,23], the Tumor-Associated NADH Oxidase class (tNOX/ENOX2) protein, which is ubiquitously expressed in cancer but not in non-transformed cells[25], the nuclear lysine specific demethylase 1A (KDM1A/LSD1)[26], and the nuclear transcription factor 2 (POU3F2)[27]. Last, capsaicin was shown to bind to PHB2, and to induce its translocation from the inner mitochondrial membrane to the nucleus in HeLa cells[19]. This also contributes to the induction of apoptosis in human myeloid leukemia cells. Given these multiple targets of capsaicin, it is interesting to note that compound 12 and 13 are unable to significantly bind to TRPV1[28]. We also observed that capsaicin and compound 5 are the only compounds capable to induce a cytostatic effect in MCF7 cells, while no cytotoxicity was measured with any compound (Supplementary Table 3). The lack of overall cytotoxicity exerted by capsaicin and its ligands may suggest low expression levels of canonical capsaicin targets in MCF7 cells. The binding activity of compound 12 and 13 is shown when the tripartite AURKA/PHB2/LC3 complex is formed, a condition observable only upon AURKA overexpression[6]. While this makes of compound 13 and 12 two more selective PHB ligands than capsaicin, it also suggests that AURKA should perhaps be overexpressed to detect the cytotoxic effects of these compounds.

Alternatively, the lack of cytotoxicity could also be explained by the reverse docking results obtained with GalaxySagittarius-AlphaFold predictions. Predictions uncovered that TRPV1 is a preferential target of capsaicin and of compound 5, while it was not a preferential target of compound 12 or 13 (Table 2, Supplementary File 1). This leads to the hypothesis that the cytostatic effect of capsaicin and of compound 5 might be related to their activity on TRPV1. With compound 12 and 13 instead, AURKA stood out as a preferential hit (Table 2, Supplementary File 1). The results on TRPV1 highlight how compound 12 and 13, despite being derived from capsaicin, have developed their own selectivity and show a high specificity for other targets such as AURKA. The presence of TRPV1 in the capsaicin list with scores comparable to the best results further corroborates the predictions for the other compounds analyzed. It is important to underline that the absence of PHB2 from the GalaxySagittarius list is not related to a potential inability of the compounds to interact with it. Despite the use of GalaxySagittarius in AlphaFold mode, the PHB2 protein model is not present in the reference database. Therefore, it was impossible to evaluate the interaction score between PHB2 and the compounds studied.

Despite the multiple targets of capsaicin, it is important to note that they are not localized at mitochondria and that the effect of capsaicin at this subcellular compartment was previously unknown. We here provide the first evidence for the capacity of capsaicin and its derivatives to modulate the protein/protein proximity between PHB2 and a cancer-related kinase such as AURKA at this compartment. At mitochondria, AURKA was shown to play a variety of roles including the modulation of mitochondrial morphology, of ATP production and of mitochondrial clearance by mitophagy[6,9]. We previously showed that the PHB2 ligand xanthohumol globally alters protein-protein interactions within the AURKA-PHB2-LC3 tripartite complex[6]. Indeed, xanthohumol lowers the AURKA/LC3 interaction while concomitantly increasing the AURKA/PHB2 interaction. Altering these interactions abolishes AURKA-dependent mitophagy and restores normal ATP levels in cancer cells with AURKA overexpression[6]. When looking at the newly-generated capsaicin derivatives, it is interesting to observe that compound 11, 12 and 13 significantly increase the AURKA/PHB2 interaction as xanthohumol does[6]. Thanks to their low standard deviation compared to that of compound 11 in FRET/FLIM experiments, our data also suggest that compound 12 and 13 may be promising capsaicin derivatives for functional studies.

Molecular docking data further corroborate the notion that compounds reinforcing protein-protein interactions by FRET have binding mechanisms compatible with the inhibition of PHB2. All the compounds tested show a preference for a binding site on PHB2 already known to be the target of the YL-939 PHB2 inhibitor[24]. YL-939 appears to bind PHB2 with two H-bonds involving the D82 and D127 residues, while its distal phenyl group forms hydrophobic interactions with V43 and I80. Mutagenesis experiments performed on D82 and D127 highlighted that these residues are fundamental for the inhibitor binding and reduce the sensitivity to erastin in ferroptosis[24]. Similarly, the binding of capsaicin and of compound 5, 12 and 13 to this pocket could exert an inhibitory effect. Moreover, the proximity of the PHB2 inhibition site to the LC3 binding site - YQRL motif, residues 121-124 - may also indicate the involvement of these compounds in modulating the binding of LC3 to PHB2.

Interestingly, capsaicin and compound 5, 12 and 13 all display an interaction with AURKA and show binding energy levels which are better than the ones obtained when interacting with PHB2. However, the different size of the molecules and their different hindrance may explain their differential effects on AURKA. First, they could compete for the binding to Ser39 of PHB2 a residue targeted by AURKA and required for AURKA-dependent mitophagy[6]. As a consequence, the functions of PHB2 within or outside the AURKA/PHB2/LC3 complex could also be compromised. In addition, the modelling of the tripartite complex shows that compound 5 represent a steric hindrance for PHB2 accommodation, and this in the two conformations detected. Compound 12 and 13 instead accommodate at the interface of AURKA and PHB2. This may give rise to a stabilized AURKA/PHB2 interaction, which is corroborated by FRET/FLIM experiments in living cells. Docking simulations performed directly on the tripartite complex also underline that compound 12 and 13 can reach the AURKA active site. On the contrary, capsaicin and compound 5 are forced to move away from the active site of the kinase and to change their orientation when the complex is formed.

Our results show that capsaicin derivatives are pertinent chemical tools to alter the formation of the tripartite AURKA/PHB2/LC3 complex both in live cells and at the molecular level. Future reports will elucidate whether the AURKA/PHB2 interaction could be considered as a proxy to predict the efficacy of novel pharmacological compounds towards the mitochondrial functions

of AURKA and ultimately, the proliferation rates of cancer cells with AURKA overexpression. Therefore, our study paves the way to the use of capsaicin derivatives to be used as lead compounds towards the development of a new class of anticancer drugs targeting the mitochondrial activation of AURKA.

Materials and methods

Cell culture procedures

MCF7 (HTB-22) cells were purchased from the American Type Culture Collection and were kept free from mycoplasma throughout experiments. They were grown in Dulbecco's Modified Eagle's Medium (DMEM) already containing 1% L-glutamine (Thermo Fisher Scientific), and supplemented with 10% FBS (Thermo Fisher Scientific) and 1% penicillin-streptomycin (Thermo Fisher Scientific). For all live microscopy experiments, cells were grown at 37°C either in Nunc Lab-Tek Chamber slides (Thermo Fisher Scientific), or in polymer-coated μ-Slide 8 Well high slides (Ibidi). Before proceeding with FLIM imaging, standard growth media was replaced with phenol red-free Leibovitz's L-15 medium (Thermo Fisher Scientific) supplemented with 20% FBS and 1% penicillin–streptomycin. Plasmid transfections were performed with Lipofectamine 2000 (Thermo Fisher Scientific), according to the manufacturer's instructions and incubated for 48h before FLIM analyses. pcDNA3 AURKA-GFP, a pcDNA 3.1 empty vector and pmCherry N1-PHB2 were previously described[6].

Fluorizoline, FL3, capsaicin and all capsaicin derivatives were resuspended in DMSO and stored at -20°C for long-term storage. Each compound was used at a final concentration of 10 μM, and incubated for 6h before proceeding with microscopy experiments.

FRET/FLIM microscopy

Images were acquired with a time-gated custom-made setup based on a spinning disk microscope as described in [18]. GFP was used as a FRET donor in all experiments, and it was excited at 480 ± 10 nm with a supercontinuum picosecond pulsed laser source. Emission was selected using a band pass filter of 483/35 nm. The FLIM setup was controlled by the Inscoper Suite solution (Inscoper, France), and GFP lifetime was measured in real-time during acquisition with the Inscoper software. In all experiments, AURKA-GFP lifetime was calculated by the Inscoper software only when the pixel- by-pixel fluorescence intensity in the first gate was above 2000 gray levels. To calculate the ΔLifetime, the mean lifetime of the cells in the donor-only condition (AURKA-GFP + Empty vector) was calculated, and then used to normalize data in all the analyzed conditions and for each independent experiment as previously reported[29].

Cell proliferation and viability

MCF7 cells were cultured in Eagle's Minimal Essential Medium (EMEM, Sigma-Aldrich) supplemented with 10% FBS (Thermo Fisher Scientific), 2 mM L-Glutamine (Sigma-Aldrich), 1mM sodium pyruvate (Sigma), 100 U/ml penicillin (Sigma-Aldrich) and 100 μg/ml streptomycin (Sigma-Aldrich). Cells were grown at 37°C in NUNC Edge96 Nunclon Delta Surface 96-well plates. All compounds were incubated for 48h prior to cell proliferation and viability experiments, at the following final concentrations: 0.1 μM; 0.5 μM; 1 μM; 5 μM; 10 μM, 50 μM with three independent points measured for each concentration. Only data for 50 μM were reported in Supplementary Table 3. Cellular proliferation was calculated with an IncuCyte S3, driven by the IncuCyte 2023A software and by measuring cellular confluence. Cellular viability was measured with the WST-1 test (Takara). 450 nm absorbance was recorded using an Envision (Perkin Elmer) plate reader.

Molecular modelling of capsaicin and capsaicin derivatives, structural models of the AURKA-PHB2-LC3 tripartite complex and molecular docking simulations.

The molecular structures of compounds 5, 12 and 13 were built by Discovery Studio 4.5 (Dassault Systèmes BIOVIA), and minimized by Chimera 1.14[30]. The molecular structure of capsaicin was instead downloaded from PubChem[31] (CID: 1548943). The putative protein targets of capsaicin and compounds 5-12-13 were estimated with GalaxySagittarius-AF[32], exploiting both similarity and compatibility mode, and re-ranked using docking. The structures of Aurora kinase A (AURKA) in its active conformation, and of Microtubule-associated protein 1A/1B light chain 3B (LC3) were retrieved from PDB[33] (PDB code: 5G1X, and 7ELG respectively). For the 3D-structure of Prohibitin-2 (PHB2), the AlphaFold complete model (identifier: AF-Q99623-F1) was used[34]. The H-dock software[35] was used to create a model of the AURKA-PHB2-LC3 tripartite complex. The complexes obtained were analyzed in terms of compatibility of the interface orientation with the information on the binding site from the literature. Once the most suitable complex was selected, it was used as a template for Modeller9.22[36] to refine the interactions among the chains of the complex. PDBePISA[37] was used to calculate the binding energy and the interactions established at the interfaces. ProSA-web[38] was used to analyze the quality of the model obtained.

Blind and focused docking simulations were performed by AutoDock 4.2.5.1 and AutoDockTools 1.5.6[39]. For blind docking with AURKA active conformation, a 126x126x126 grid box with grid center coordinates equal to (37.908; -15.701; 37.348) was set; spacing was set at 0.397 Å. Focused flexible docking was performed on the AURKA active site to refine the binding energy and the conformation of the compounds in the pocket. A 60x60x60 grid box with center coordinates (39.621; -19.626; 37.357) and a spacing of 0.375 Å was set as default. Four residues - D256-N261-D274-W277 - were set as flexible. This is because D256 is the active site, and the other three residues occupy positions with a relevant role for the access to the active site.

For PHB2, docking was focused on the portion of the protein having a more organized tertiary structure. The greed box was equal to 126x126x126 with center coordinates equal to (-15.757; -2.72; 16.28) and a spacing value of 0.386 Å. Docking simulations performed on the complex with capsaicin and compounds 5, 12 or 13 were performed focusing the box at the interface of the three molecules with a greed box equal to 126x126x126, a greed center with coordinates (-31.084; -6.4; 22.525), and spacing value of 0.386 Å.

All the docking simulations allowed the flexibility of the ligand. The results obtained were analyzed by AutoDockTools and LigPlot v2.2[40].

Statistical analyses

Two-way ANOVA was used to compare the effect of the transfection conditions and of the pharmacological treatments (Figure 2), and was performed using GraphPad Prism v. 10. Statistics and summary statistics for biological replicates comparing the effect of the transfection conditions and the pharmacological treatments were performed with SuperPlotsofData[41] (Table 1, Supplementary Table 1, Figure 4C).

Author contributions

CRediT taxonomy: Conceptualization (G.B., L.D.), Data Curation (A.D., C.C., D.G., V.P., K.F., A.F., G.B., L.D.), Formal Analysis (A.D., C.C., D.G, V.P., K.F., A.F., G.B., L.D.), Funding Acquisition (G.B., L.D.), Investigation (A.D., C.C., D.G., V.P., K.F., A.F., G.B., L.D.), Methodology (A.F., G.B., L.D.), Project Administration (G.B., L.D.), Resources (G.B., L.D.), Software (C.C., D.G., V.P., A.F., G.B.), Supervision (A.F, G.B., L.D.), Validation (G.B., L.D.), Visualization (C.C., D.G., V.P., A.F., G.B., L.D.), Writing – original, review and editing (C.C., D.G, A.F., G.B., L.D.).

Acknowledgements

We thank S. Dutertre, X. Pinson and G. Le Marchand at the Microscopy Rennes Imaging Center (MRic, *Biologie, Santé, Innovation Technologique* - BIOSIT, Rennes, France) for assistance with FLIM experiments. MRic is member of the national infrastructure France-BioImaging, supported by the French National Research Agency (ANR-10-INBS-04). We thank the *Plateforme de chimie biologique integrative* (PCBIS, University of Strasbourg) for assistance with cell proliferation and viability experiments. We are grateful to N. Jolivet for critical reading of the manuscript. This work was supported by the *Centre National de la Recherche Scientifique* (CNRS), the University of Rennes, the *Ligue Contre le Cancer, Comités d'Ille et Vilaine et du Finistère* to G.B. C.C was supported by a PhD fellowship from the *Ligue Nationale Contre le Cancer* (grant no. IP/SC – 17653).

Competing interests

The authors declare no competing or financial interests.

References

1.   Wang, T., Ma, F., and Qian, H. (2021). Defueling the cancer: ATP synthase as an emerging target in cancer therapy. Molecular Therapy - Oncolytics *23*, 82–95. https://doi.org/10.1016/j.omto.2021.08.015.

2.   Pickrell, A.M., and Youle, R.J. (2015). The Roles of PINK1, Parkin and Mitochondrial Fidelity in Parkinson's Disease. Neuron *85*, 257–273. https://doi.org/10.1016/j.neuron.2014.12.007.

3.   Wang, S., Long, H., Hou, L., Feng, B., Ma, Z., Wu, Y., Zeng, Y., Cai, J., Zhang, D., and Zhao, G. (2023). The mitophagy pathway and its implications in human diseases. Sig Transduct Target Ther *8*, 1–28. https://doi.org/10.1038/s41392-023-01503-7.

4.   Wei, Y., Chiang, W.-C., Sumpter, R., Mishra, P., and Levine, B. (2017). Prohibitin 2 Is an Inner Mitochondrial Membrane Mitophagy Receptor. Cell *168*, 224-238.e10. https://doi.org/10.1016/j.cell.2016.11.042.

5.   Wang, D., Tabti, R., Elderwish, S., Abou-Hamdan, H., Djehal, A., Yu, P., Yurugi, H., Rajalingam, K., Nebigil, C.G., and Désaubry, L. (2020). Prohibitin ligands: a growing armamentarium to tackle cancers, osteoporosis, inflammatory, cardiac and neurological diseases. Cell. Mol. Life Sci. *77*, 3525–3546. https://doi.org/10.1007/s00018-020-03475-1.

6.   Bertolin, G., Alves-Guerra, M.-C., Cheron, A., Burel, A., Prigent, C., Borgne, R.L., and Tramier, M. (2021). Mitochondrial Aurora kinase A induces mitophagy by interacting with MAP1LC3 and Prohibitin 2. Life Science Alliance *4*. https://doi.org/10.26508/lsa.202000806.

7.   Kashatus, D.F., Lim, K.-H., Brady, D.C., Pershing, N.L.K., Cox, A.D., and Counter, C.M. (2011). RALA and RALBP1 regulate mitochondrial fission at mitosis. Nature Cell Biology *13*, 1108–1115. https://doi.org/10.1038/ncb2310.

8.   Nikonova, A.S., Astsaturov, I., Serebriiskii, I.G., Dunbrack, R.L., and Golemis, E.A. (2013). Aurora A kinase (AURKA) in normal and pathological cell division. Cellular and Molecular Life Sciences *70*, 661–687. https://doi.org/10.1007/s00018-012-1073-7.

9.   Bertolin, G., Bulteau, A.-L., Alves-Guerra, M.-C., Burel, A., Lavault, M.-T., Gavard, O., Le Bras, S., Gagné, J.-P., Poirier, G.G., Le Borgne, R., et al. (2018). Aurora kinase A localises to mitochondria to control organelle dynamics and energy production. eLife *7*. https://doi.org/10.7554/eLife.38111.

10.   Grant, R., Abdelbaki, A., Bertoldi, A., Gavilan, M.P., Mansfeld, J., Glover, D.M., and Lindon, C. (2018). Constitutive regulation of mitochondrial morphology by Aurora A kinase depends on a predicted cryptic targeting sequence at the N-terminus. Open Biology *8*, 170272. https://doi.org/10.1098/rsob.170272.

11.   Bertolin, G., and Tramier, M. (2020). Insights into the non-mitotic functions of Aurora kinase A: more than just cell division. Cell. Mol. Life Sci. *77*, 1031–1047. https://doi.org/10.1007/s00018-019-03310-2.

12.   Chouha, N., Abou-Hamdan, H., Yurugi, H., Yoshii, R., Ii, H., Najem, A., Ghanem, G.E., Nakata, S., Rajalingam, K., Peng, Y., et al. (2022). Development of fluorizoline analogues as prohibitin ligands that modulate C-RAF signaling, p21 expression and melanogenesis. European Journal of Medicinal Chemistry *242*, 114635. https://doi.org/10.1016/j.ejmech.2022.114635.

13.   Tabti, R., Lamoureux, F., Charrier, C., Ory, B., Heymann, D., Bentouhami, E., and

Désaubry, L. (2021). Development of prohibitin ligands against osteoporosis. European Journal of Medicinal Chemistry *210*, 112961. https://doi.org/10.1016/j.ejmech.2020.112961.

14.   Yurugi, H., Zhuang, Y., Siddiqui, F.A., Liang, H., Rosigkeit, S., Zeng, Y., Abou-Hamdan, H., Bockamp, E., Zhou, Y., Abankwa, D., et al. (2020). A subset of flavaglines inhibits KRAS nanoclustering and activation. Journal of Cell Science *133*, jcs244111. https://doi.org/10.1242/jcs.244111.

15.   Elderwish, S., Audebrand, A., Nebigil, C.G., and Désaubry, L. (2020). Discovery of 3,3′-pyrrolidinyl-spirooxindoles as cardioprotectant prohibitin ligands. European Journal of Medicinal Chemistry *186*, 111859. https://doi.org/10.1016/j.ejmech.2019.111859.

16.   Djehal, A., Krayem, M., Najem, A., Hammoud, H., Cresteil, T., Nebigil, C.G., Wang, D., Yu, P., Bentouhami, E., Ghanem, G.E., et al. (2018). Targeting prohibitin with small molecules to promote melanogenesis and apoptosis in melanoma cells. European Journal of Medicinal Chemistry *155*, 880–888. https://doi.org/10.1016/j.ejmech.2018.06.052.

17.   Padilla-Parra, S., Audugé, N., Coppey-Moisan, M., and Tramier, M. (2008). Quantitative FRET analysis by fast acquisition time domain FLIM at high spatial resolution in living cells. Biophys. J. *95*, 2976–2988. https://doi.org/10.1529/biophysj.108.131276.

18.   Leray, A., Padilla-Parra, S., Roul, J., Héliot, L., and Tramier, M. (2013). Spatio-Temporal Quantification of FRET in living cells by fast time-domain FLIM: a comparative study of non-fitting methods [corrected]. PLoS ONE *8*, e69335. https://doi.org/10.1371/journal.pone.0069335.

19.   Kuramori, C., Azuma, M., Kume, K., Kaneko, Y., Inoue, A., Yamaguchi, Y., Kabe, Y., Hosoya, T., Kizaki, M., Suematsu, M., et al. (2009). Capsaicin binds to prohibitin 2 and displaces it from the mitochondria to the nucleus. Biochemical and Biophysical Research Communications *379*, 519–525. https://doi.org/10.1016/j.bbrc.2008.12.103.

20.   Janusz, J.M., Buckwalter, B.L., Young, P.A., LaHann, T.R., Farmer, R.W., Kasting, G.B., Loomans, M.E., Kerckaert, G.A., Maddin, C.S., and Berman, E.F. (1993). Vanilloids. 1. Analogs of capsaicin with antinociceptive and antiinflammatory activity. J Med Chem *36*, 2595–2604. https://doi.org/10.1021/jm00070a002.

21.   Padilla-Parra, S., Audugé, N., Lalucque, H., Mevel, J.-C., Coppey-Moisan, M., and Tramier, M. (2009). Quantitative Comparison of Different Fluorescent Protein Couples for Fast FRET-FLIM Acquisition. Biophys J *97*, 2368–2376. https://doi.org/10.1016/j.bpj.2009.07.044.

22.   Amantini, C., Mosca, M., Nabissi, M., Lucciarini, R., Caprodossi, S., Arcella, A., Giangaspero, F., and Santoni, G. (2007). Capsaicin-induced apoptosis of glioma cells is mediated by TRPV1 vanilloid receptor and requires p38 MAPK activation. J Neurochem *102*, 977–990. https://doi.org/10.1111/j.1471-4159.2007.04582.x.

23.   Xu, S., Cheng, X., Wu, L., Zheng, J., Wang, X., Wu, J., Yu, H., Bao, J., and Zhang, L. (2020). Capsaicin induces mitochondrial dysfunction and apoptosis in anaplastic thyroid carcinoma cells *via* TRPV1-mediated mitochondrial calcium overload. Cellular Signalling *75*, 109733. https://doi.org/10.1016/j.cellsig.2020.109733.

24.   Yang, W., Mu, B., You, J., Tian, C., Bin, H., Xu, Z., Zhang, L., Ma, R., Wu, M., Zhang, G., et al. (2022). Non-classical ferroptosis inhibition by a small molecule targeting PHB2. Nat Commun *13*, 7473. https://doi.org/10.1038/s41467-022-35294-2.

25.   Islam, A., Su, A.J., Zeng, Z.-M., Chueh, P.J., and Lin, M.-H. (2019). Capsaicin

Targets tNOX (ENOX2) to Inhibit G1 Cyclin/CDK Complex, as Assessed by the Cellular Thermal Shift Assay (CETSA). Cells *8*, 1275. https://doi.org/10.3390/cells8101275.

26.   Jia, G., Cang, S., Ma, P., and Song, Z. (2020). Capsaicin: A "hot" KDM1A/LSD1 inhibitor from peppers. Bioorg Chem *103*, 104161. https://doi.org/10.1016/j.bioorg.2020.104161.

27.   Chen, H.Y., Lee, Y.H., Chen, H.Y., Yeh, C.A., Chueh, P.J., and Lin, Y.-M.J. (2016). Capsaicin Inhibited Aggressive Phenotypes through Downregulation of Tumor-Associated NADH Oxidase (tNOX) by POU Domain Transcription Factor POU3F2. Molecules *21*, 733. https://doi.org/10.3390/molecules21060733.

28.   Yang, F., Xiao, X., Cheng, W., Yang, W., Yu, P., Song, Z., Yarov-Yarovoy, V., and Zheng, J. (2015). Structural mechanism underlying capsaicin binding and activation of the TRPV1 ion channel. Nat Chem Biol *11*, 518–524. https://doi.org/10.1038/nchembio.1835.

29.   Bertolin, G., Marchand, G.L., and Tramier, M. (2020). Real-Time Monitoring of Aurora kinase A Activation using Conformational FRET Biosensors in Live Cells. JoVE (Journal of Visualized Experiments), e61611. https://doi.org/10.3791/61611.

30.   Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., and Ferrin, T.E. (2004). UCSF Chimera--a visualization system for exploratory research and analysis. J Comput Chem *25*, 1605–1612. https://doi.org/10.1002/jcc.20084.

31.   Kim, S., Chen, J., Cheng, T., Gindulyte, A., He, J., He, S., Li, Q., Shoemaker, B.A., Thiessen, P.A., Yu, B., et al. (2023). PubChem 2023 update. Nucleic Acids Research *51*, D1373–D1380. https://doi.org/10.1093/nar/gkac956.

32.   Kwon, S., Jung, N., Yang, J., and Seok, C. (2024). GalaxySagittarius-AF: Predicting Targets for Drug-Like Compounds in the Extended Human 3D Proteome. Journal of Molecular Biology, 168617. https://doi.org/10.1016/j.jmb.2024.168617.

33.   Berman, H.M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T.N., Weissig, H., Shindyalov, I.N., and Bourne, P.E. (2000). The Protein Data Bank. Nucleic Acids Research *28*, 235–242. https://doi.org/10.1093/nar/28.1.235.

34.   Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly accurate protein structure prediction with AlphaFold. Nature *596*, 583–589. https://doi.org/10.1038/s41586-021-03819-2.

35.   Yan, Y., Tao, H., He, J., and Huang, S.-Y. (2020). The HDOCK server for integrated protein–protein docking. Nat Protoc *15*, 1829–1852. https://doi.org/10.1038/s41596-020-0312-x.

36.   Eswar, N., Webb, B., Marti-Renom, M.A., Madhusudhan, M.S., Eramian, D., Shen, M.-Y., Pieper, U., and Sali, A. (2006). Comparative protein structure modeling using Modeller. Curr Protoc Bioinformatics *Chapter 5*, Unit-5.6. https://doi.org/10.1002/0471250953.bi0506s15.

37.   Krissinel, E., and Henrick, K. (2007). Inference of Macromolecular Assemblies from Crystalline State. Journal of Molecular Biology *372*, 774–797. https://doi.org/10.1016/j.jmb.2007.05.022.

38.   Wiederstein, M., and Sippl, M.J. (2007). ProSA-web: interactive web service for the recognition of errors in three-dimensional structures of proteins. Nucleic Acids Res *35*,

W407-410. https://doi.org/10.1093/nar/gkm290.

39.   Morris, G.M., Huey, R., Lindstrom, W., Sanner, M.F., Belew, R.K., Goodsell, D.S., and Olson, A.J. (2009). AutoDock4 and AutoDockTools4: Automated Docking with Selective Receptor Flexibility. J Comput Chem *30*, 2785–2791. https://doi.org/10.1002/jcc.21256.

40.   Wallace, A.C., Laskowski, R.A., and Thornton, J.M. (1995). LIGPLOT: a program to generate schematic diagrams of protein-ligand interactions. Protein Eng *8*, 127–134. https://doi.org/10.1093/protein/8.2.127.

41.   Goedhart, J. (2021). SuperPlotsOfData – a web app for the transparent display and quantitative comparison of continuous data from different conditions. MBoC, mbc.E20-09-0583. https://doi.org/10.1091/mbc.E20-09-0583.