Human transcription factor combinations mapped by footprinting with deaminase

Runsheng He[1,2]†, Wenyang Dong[1,2,4]†, Wenping Ma[2]†, Zhi Wang[1,2,4]†, Long Gao[2]†, Chen

Xie[1,2]†, Dubai Li[1,2,3], Ke Shen[1,2,4], Fanchong Jian[1,2,5], Jiankun Zhang[1,2,4], Yuan Yuan[1,2,5], Xinyao

Wang[1,3], Yuxuan Pang[1,2,3], Zhen Zhang[2], Yinghui Zheng[1], Shuang Liu[2], Cheng Luo[3], Xiaoran

Chai[1], Jun Ren[2], Zhanxing Zhu[2], Xiaoliang Sunney Xie[1,2] *

Affiliations

[1] Beijing Advanced Innovation Center for Genomics (ICG) & Biomedical Pioneering Innovation

Center (BIOPIC), Peking University; Beijing 100871, China.

[2] Changping Laboratory; Beijing, 102206, China.

[3] Academy for Advanced Interdisciplinary Studies, Peking University; Beijing, 100871, China.

[4] School of Life Sciences, Peking University; Beijing, 100871, China.

[5] College of Chemistry and Molecular Engineering, Peking University; Beijing, 100871, China.

† Authors contributed equally.

* Corresponding author. Email: sunneyxie@biopic.pku.edu.cn

Abstract

An individual's somatic cells have the same genome but exhibit cell-type-specific

transcriptome regulated by a combination of transcription factors (TFs) for each gene. Mapping of

TF sites on the human genome is critically important for understanding functional genomics. Here

we report a novel technique to measure human TFs' binding sites genome-wide with single-base

resolution by footprinting with deaminase (FOODIE). Single-molecule sequencing reads from

thousands of cells after *in situ* deamination yielded site-specific TF binding fractions and the

cooperativity among adjacent TFs. In a human lymphoblastoid cell line, we found that, genes in a

correlated gene module (CGM) share TF(s) in their *cis*-regulatory elements to facilitate a particular

biological function. Finally, single-cell resolved experiments (scFOODIE) allow cell-type-specific

TF footprinting in heterogeneous brain tissues.

Introduction

Every somatic cell in the human body has essentially the same genome but carries out distinct functions in specific tissues with different gene expression. Transcription factors (TFs) bind to *cis*-regulatory elements, including promoters and enhancers of genes on the genome, to control gene expression and regulation, cell differentiation, and development, in response to environmental signals through regulatory networks (*1*). Our knowledge about gene expression and regulation first came from the study of bacteria, where one TF, acting as a key to turn on and off an operon, contains multiple protein-coding genes that carry out a particular biological function (*2, 3*). Such TF activities have been directly monitored and quantitatively described on a single-cell and single-molecule basis (*4, 5*).

In contrast, TFs in eukaryotes, ever since their first isolation (*6*), seem to operate differently in that each binds to many sites on the genome with lower binding affinities (*7*), and gene regulation is achieved through the binding of a combination of several TFs at promoters and enhancers (*8*). The fact that several TFs, acting together as a set of keys, control one protein-coding gene makes the total number of ~1600 TFs sufficient to control ~22000 genes in human cells. More importantly, we hypothesize, that this provides a way to coordinate multiple genes in a gene module to accomplish a particular biological function, such as protein synthesis or glucose metabolism (*9-12*). However, due to the lack of experimental techniques, little is known about how TF combinations are organized to coordinate regulatory networks, which remains the central question in eukaryotic biology.

Most TF binding sites in eukaryotes have been identified by chromatin immunoprecipitation ChIP-on-chip (*13*) and ChIP-seq (*14, 15*), analyzing one TF at a time, through antibody pulling down followed by DNA sequencing, with a low spatial resolution of a few hundred base pairs. A common method for determining cellular binding motifs is footprinting, i.e. detecting DNA sequences that are protected from enzymatic treatment, most notably by DNase-seq, which detects the endpoints of DNA fragments cleaved by DNase I (*16*). However, both ChIP-seq and DNase-seq require bulk samples with millions of cells, usually of cell lines, and are incapable of detection on a single DNA.

To overcome these limitations, another footprinting method by cytosine methylation had been previously attempted with Single Molecule Footprinting (SMF) (*17*). However, it is not appliable

to normal mammalian cells, as the extraneous methyltransferase used was mostly indistinguishable

from endogenous methylation.

Here we report a novel approach based on double-stranded DNA cytosine deamination, named

FOOtprinting with DeamInasE (FOODIE), to detect TFs' binding sequences at single-base

resolution along the genome, as the TFs' steric hindrance prevents the conversion of cytosine to

uracil. Cost-effective and scalable,　FOODIE requires sequencing only thousands of cells,

significantly fewer than those required for ChIP-seq or DNase-seq.

Genome-wide mapping of TF footprints

FOODIE relies on double-stranded DNA deaminase with high efficiency (Fig. 1A). We

initially tested DddA, the first double-stranded DNA deaminase reported (*18*), which only

deaminates cytosine preceded by thymine, not capable of footprinting all TFs bound to the genome

(Fig. S1A). To overcome this limitation, we characterized a new dsDNA cytosine deaminase from

a large group of reported toxins (*19*), which we named DddB (also known as BadTF3 (*20*)) after

DddA. Unlike DddA, DddB converts cytosine to uracil largely independent of adjacent bases (Fig.

1B), allowing us to footprint almost all TF motif sequences (Fig. S1A). We determined the crystal

structure of DddB in complex with its immunity protein at 2.5 Å resolution (Fig. 1C). Despite the

overall structural similarity between DddB and DddA, DddB features an intervening helix near the

active site, whereas DddA has a loop in this region (Fig. S1B).

Similar but one step more than ATAC-seq (*21*), FOODIE uses transposase Tn5 in

permeabilized cells to enrich the open chromatin regions, where most TFs bind and exert their

regulatory roles. DddB is then added after Tn5 tagmentation to detect protein binding on DNA,

whose underlying cytosines are protected from enzymatic deamination. As such, FOODIE is

capable of probing the regulatory sites in open chromatin regions in a cost-effective and high-

throughput way.

TF binding profiles detected by ChIP-seq, FOODIE, and DNase-seq are shown in Fig. 1D.

ChIP-seq had a low spatial resolution (200-500bp), not capable of telling whether both or just one

of the detected CTCF sequences were occupied. In contrast, FOODIE accurately identified CTCF

that the footprint on the right is bound, which was consistent with DNase-seq footprinting. We

further compared the number of genome-wide binding sites of CTCF detected by FOODIE with

those obtained from ChIP-seq and DNase-seq. The CTCF binding sites detected by FOODIE show

strong concordance with those detected by ChIP-seq and largely overlap with those detected by

DNase-seq footprinting (Fig. 1E).

Next we employed a heat shock experiment to validate the reliability of FOODIE. Upon a

temperature increase, the transcription factor HSF is activated to bind to the promoters of target

genes to induce expression (*22*). At HSP90AA1 promoter, we observed a decrease in the

conversion ratio within HSF motif sequences, which corresponded to the gain of HSF binding

upon heat shock (Fig. 1F). The above data suggested that FOODIE can identify precise binding of

TFs with high accuracy.

To support the interactive analyses of genome-wide footprints identified by FOODIE, we

developed a web server (http://foodie.sunneyxielab.org) with a user-friendly interface. This

platform includes a genome browser for visualization and a data portal with our FOODIE data.

Fig. 2A shows a browser display of the genomic region around the TRNAU1AP promoter in

chromosome 1, a chromatin open region between nucleosomes -1 (and -2) and +1 (and +2), with

three distinct footprints. The first row is the ATAC track seen by Tn5 tagmentation. The second

row is the C to U conversion efficiency showing the footprints. The raw reads of single-molecule

sequences below, with the converted cytosines on the Watson strand (red) and the Crick strand

(green), show the binding fractions along the genome.

We found that the deamination efficiency of DddB is dependent on the neighboring sequences

of cytosines (especially from -4 to +2) (Fig. S2), which needs to be normalized when calculating

the binding fraction. We measured deamination efficiency along the naked human genome as a

positive control for the conversion ratio correction. The binding fraction is one minus conversion

ratio.

Fig. 2B shows the mean number of footprints present in *cis*-regulatory elements of genes and

their relative arrangement (see Material and Methods). The median number of footprints per gene

promoter was found to be 5 (median number), compared to 2 in enhancer regions. Notably, the

binding fractions of TFs in the promoter regions were higher than those of TFs in enhancer regions.

The FOODIE footprints need to be assigned to particular TFs, commonly through motifs

tabulated on websites, such as JASPAR (*23*) and HOCOMOCO (*24*), which collected vast amounts

of information from *in vitro* data of SELEX (*25*) and PBM's (*26*) as well as *in vivo* data of ChIP-

seq (*27*). It turned out that a footprint has many different possible assignments of TFs, which we

can not pin down because of the lack of binding affinities from databases and the intracellular

concentrations. While we are working on the methods to fully assign the footprints, in this work

we assigned footprints to certain TFs only when they have strong ChIP-seq signal in GM12878

cell line. This resulted in reliable assignments to only seventy-nine known TFs.

As such, the abundance of the assigned TFs is shown in promoter and enhancer regions, we

found promoters were particularly enriched for the NRF1, KLF/SP, ZNF143, ETS, and YY1,

footprints, which were rarely detected in enhancers (Fig. 2C). Taken together, TF bound to

promoters and enhancers differ in average number within open regions, binding fractions and TF

types.

Single-molecule sequencing reads revealing binding cooperativity among TFs

While the binding fraction of a particular binding site can be determined by the proportion of

single molecules of sequenced DNA with unconverted C to that of the total of converted and

unconverted C. The fact that sequencing occurs on individual molecules allows us to determine

the cooperativity between TFs bound to two adjacent binding sites.

Single-molecule reads at a CTCF binding site in the genome are shown in Fig. 2D, exhibiting

two separate footprints of the bridging structure of the zinc finger hinge (*28*), which are either

simultaneously occupied or not. In Fig. 2D, two adjacent TFs RFX and CREB have two distinct

footprints (*29*). To evaluate their binding cooperativity, we performed Fisher's exact test for the

significance of independence on a pair of cytosines, each selected from the two binding sites. The

Φ correlation coefficient is between 0 and 1 for positive cooperativity and between -1 and 0 for

negative cooperativity. A red dot indicates significant positive cooperativity ($P<0.001$). A pair of

cytosines within the RFX (or CREB) binding sequence are positively cooperative because the

binding of RFX to them is simultaneous. Interestingly, cytosine pairs across CREB and RFX

binding sequences are also positively cooperative, i.e. RFX and CREB's binding is not

independent but facilitates each other (Fig. 2D). We attribute this positive coopertavity to protein-

protein interactions between two TFs, which has been previously reported (*29*).

Conversely, negative cooperativity was seen between two NRF1 binding sites within the

ACTN4 promoter (Fig. 2D). While the Φ of the two cytosines within each NRF1 binding site is

positive, the pairs across different sites are negative (blue: $P<0.001$), meaning their binding is

exclusive. Such single-molecule measurements provide unprecedented information about how TFs

work together in the genome.

Shared TFs among genes within the same CGM

Many protein-coding genes form a module to work together to accomplish a particular

biological function. People have been looking for gene modules with the bulk (*12*) and single cells

(*9, 10, 30*) by perturbation measurements. By improving the detectability of single-cell

transcriptome, we demonstrated the identification of correlated gene modules (CGMs) without

perturbation under the steady state condition (*9, 10*). The bursty nature of gene expression has

been long established and attributed to the single-molecule nature of DNA in single cells (*31-35*).

The coordination of a particular biological function by multiple genes requires their

synchronization. Experimentally, such synchronization can be probed by measuring the covariance

matrix of any pair of RNA in a single-cell transcriptome experiment (MALBAC-DT) (*9, 10*).

Recently we have developed a new method for more accurate identification of CGMs, Triangle

Screening of Hierarchical Accumulated-Edges (TriSHA) (*36*). Over Representation Analysis

(ORA) reveals that nearly 80% of the modules are functionally annotated with a False Discovery

Rate (FDR) below 0.05. Through module connection analyses, we have identified three distinct

types of CGMs within the GM12878 cell line based on functional annotation, associated with

specific cellular functions: housekeeping, tissue-specific, and cell cycle (Fig. 3A). Supplementary

Table S1 lists the CGMs with size larger than 10 that we identified. Gene t-SNE plot of GM12878

clearly shows the cluster of genes in the identified CGMs (Fig. 3B).

How are the genes in the same module distributed within the 3D genome? Are they clustered

together, or spatially dispersed? Our group determined the 3D genome structure of a somatic

human cell (*37*). To answer the above question, we employed our latest high-resolution single-cell

3D genome structures of GM12878 cells using scMicro-C (*38*). Taking the DNA replication CGM

(MD20, cell cycle clade) of 317 genes as an example, Fig. 3C shows that they are not clustered

but relatively randomly distributed within the nucleus, with the more quantitative distribution of

pair-wise distance shown in Fig. 3D. No spatial clustering of genes in other gene modules was

observed either. This observation does not support the idea that the correlated genes are transcribed

within certain locations in the nucleus. We hypothesize that the coordination of different genes in

the module is achieved through TF presented in the contiguous euchromatin space.

With FOODIE, we are in a position to determine how TFs organize the CGMs. We looked at

the enrichment of TFs shared among both promoters and enhancers of genes within a module,

which is characterized by the hypergeometric test. Out of sixty-five CGMs identified in GM12878

(Fig. 3A and Supplementary Table S1), we successfully identified that many TFs are statistically

enriched in different gene modules, including for example, E2F factors in the promoters of genes

in MD16 cell cycle-related CGM (*39*) and RELB in the enhancers of genes in MD47 regulation of

immune system CGM (*40*) (Fig. 4A). As a control, when we randomly selected tens of genes (size

from 30 to 150) out of the ~4000 expressed genes, no TFs were significantly enriched.

Significantly we found enrichments of specific TFs are clearly different for the three types of

CGMs. For cell cycle and housekeeping CGMs, the TF enrichment is primarily in promoters. For

example, the NF-Y factor is enriched in the genes of MD14 cholesterol metabolism CGM (Fig.

4A). The number of enriched TFs ($P<0.05$) were 31 and 59 for cell cycle modules and house

keeping modules, respectively. As a control, if the cell cycle and house keeping modules were

grouped together two large groups, the number enriched TFs ($P<0.05$) were reduced to 9 and 30,

respectively. This lends further support on module-specific TF enrichment.

For tissue-specific CGMs, the TF enrichment is primarily in enhancers (Fig. 4A). For example,

STAT3 enrichment is present in MD 48 immune response CGM. This suggests that there are two

distinct modes of CGM regulation: genes with essential cellular functions rely on TFs' binding at

promoters for robustness, while tissue-specific genes depend more on enhancers for larger

tunability.

We further observed certain TF pairs are shared among genes in both promoters and enhancers

within the same CGM (Fig. 4B) and evaluated the enrichment of TF using the hypergeometric test.

TF pairs are highly abundant in GCMs, and there are more TF pairs in enhancer regions (Fig. S2A).

Around one-third of the CGMs have enriched TF pairs, either at the promoters (Fig. 4B), or at the

enhancers (Fig. S2B). A higher fraction is expected when TF footprints are fully assigned. This

suggests the coordination of different genes in a module for a particular biology function through

TF combinations is ubiquitous and by no means random. Concrete examples of TF pairs shared

by multiple genes in a CGM are shown in Fig. 4C and 4D for promoters and enhancers.

As such, we are able to use FOODIE to probe how regulatory networks are orchestrated by a

combination of TFs. Although we do not yet have all the TF footprints assigned, even an

incomplete set (seventy-nine assigned TFs), already uncovered TFs pairs previously unknown in

order to coordinate gene modules.

Footprinting after cell typing by scFOODIE in heterogeneous tissues

Our goal is to characterize different human tissues with FOODIE and determine their genome-wide TF footprints. However, human tissues are heterogeneous with many different cell types, so it is necessary to determine each cell's identity. We note that each gene only has one maternal allele and one paternal allele in an individual cell, which is not sufficient to provide reliable statistics for TF footprints. However, thousands of single cells identified for a particular cell type would be sufficient to carry out footprinting.

Similar to cell typing by single-cell transcriptome or single-cell ATAC-seq, we use FOODIE signal to do unsupervised clustering of a population of cells. We evaluated the "non-conversion ratio", i. e. the ratio of unconverted cytosines to the total number of detected cytosines within the gene body ± 1000 bp regions for each gene. Using four well-characterized human cell lines (GM12878, K562, HEK293T, and HeLa) to generate a mix of single-cell FOODIE data, different cell types were clearly identified (Fig. 5A). After merging single-cell data of a particular cell type (GM12878 for example) as a pure bulk, we observed precise TF footprints of CTCF and NRF1 at the specific genome locations (Fig. 5A).

We further conducted scFOODIE for ~1500 cells from the mouse hippocampus. Four main cell types were identified, including oligodendrocytes, microglia, astrocytes, and neurons (both inhibitory and excitatory), which were annotated by marker genes (Fig. 5B). Using merged scFOODIE data of neurons, we observed footprints of JUN and MEF2B, two TFs critical for neural cellular functions (Fig. 5B). Experiments like this are expected to provide new information of neuronal gene regulation in response to either physical or chemical stimulates in the brain.

Outlook

In conclusion, we have shown that FOODIE is a high-precision method for genome-wide *in situ* mapping of TF footprints inside the nucleus with single-base resolution. FOODIE is experimentally simple, high-throughput, and cost-effective. ScFOODIE allows for cell typing in a heterogeneous tissue sample and TF footprinting in any cell type within a complex tissue. A systematic FOODIE database of different human tissues is underway. We anticipate that such information about TF combinations of different cell types and correlated gene modules will lead to the fundamental understanding of transcriptional regulation and cellular functions in eukaryotic

247 biology, especially for human biology, which might hold the potential for understanding many
248 human diseases.
249

Acknowledgments
We thank Prof. Ge Gao for the critical reading of the manuscript. We thank Qiuwan Liu for
helping with the figures, and Wenzhong Zhang for helping with the web server. **Funding**: This
project is financially supported by Changping Laboratory (2021C040201) and the Ministry of
Science and Technology of China. The initiation work was made possible by the support from
Beijing Advanced Innovation Center for Genomics at Peking University. W. D. was supported in
part by the Postdoctoral Fellowship of Peking-Tsinghua Center for Life Sciences. X. W. was
supported in part by Peking University Boya Postdoctoral Fellowship. **Author contributions**: X.
S. X., R. H., W. D., Z. W., and W. M. designed the experiments. R. H., W. D., W. M., Z. W., Y.
Y., Y. Z., S. L., C. L., and J. R. collected the experimental data. L. G., C. X., D. L., K. S., F. J., J.
Z., X. W., Y. P., Z. Zhang, X. C. and Z. Zhu performed the data analyses. X. S. X. supervised the
project and wrote the manuscript with the help of all authors. **Competing interests**: W. D., R. H.,
Z. W., and X. S. X. are coinventors on patent application WO/2024/065721 that includes
discoveries described in this manuscript. The remaining authors declare no competing interests.
Data and materials availability: Raw sequencing data and processed data have been deposited
at http://foodie.sunneyxielab.org.

Fig. 1. Genome-wide mapping of TF binding sites by FOODIE. (A) Workflow of FOODIE.
Cells were permeabilized for Tn5 tagmentation and DddB deamination. Amplicons of DNA
undergone *in situ* enzymatic deamination were sequenced to identify TF footprints. (B) (Top)
DddB C to U conversion efficiency of naked DNA *in vitro*. The x-axis is the base following, and

the y-axis base proceeding the cytidine. (Bottom) Sequence logo of the above data. (C) The crystal structure of DddB (red) complexed with its immunity protein (green). (D) Human genome tracks of DNase-seq footprint probability, FOODIE, and ChIP-seq in GM12878 cell line. The horizontal green bars represent the two possible CTCF motif sequences. Only the right one is bound by CTCF. (E) Venn diagrams of FOODIE with ChIP-seq or DNase-seq for the numbers of CTCF binding sites in open chromatin regions. (F) Heat shock factor (HSF) responding to heat shock. Upon heat shock, HSF monomers trimerize, bind to the Hsp90AA1 promoter, and leave a TF footprint on DNA.

Fig. 2. TF binding fractions and cooperativity among TFs detected by FOODIE. (A) FOODIE browser display at the promoter region of TRNAU1AP, with two nucleosomes at each side flanking the open chromatin region. The FOODIE sequencing depth is similar to ATAC-seq, whereas the FOODIE conversion ratio shows footprints of nucleosomes -1 & -2 and +1 & +2, and three distinct TFs in between. Below are single-molecule sequencing raw reads with the converted cytosines on the Watson strand (red) and the Crick strand (green). (B) (Left) Distributions for numbers of TF footprints in the promoters (median=5) and enhancers (median=2). (Right) Distributions for binding fractions of TF footprints in the promoters (median=0.348) and enhancers (median=0.317). (C) Abundance of those TF footprints with ChIP-seq-confirmed assignments on the promoters and enhancers. (D) (Left) Structure of CTCF (ZFs 3-11) and the

single-molecule sequencing reads of FOODIE footprints on a CTCF binding site. (Middle) Single-molecule sequencing reads showing RFX and CREB footprints at the HLA-DRA promoter. The positive binding cooperativity between two TFs is evident from the positive Φ correlation coefficient (dots colored in red) between a pair of cytosines, each from both TF binding sites. (Right) Single-molecule sequencing reads with two NRF1 binding sites at the ACTN4 promoter have negative binding cooperativity, which is evident from the negative Φ correlation coefficient (dots colored in blue) between a pair of cytosines, each from both TF binding sites.

Fig. 3. Genes in a CGM for a biological function not spatially clustered in nucleus. (A) Pair-wise Pearson correlation coefficients for ~ 4000 detected mRNA levels of GM12878 single cells under steady-state conditions. CGMs were identified by the improved algorithm TriSHA (*36*), and indicated by enclosed squares along with the names of major ones. Shown on the left was the

hierarchical clustering of CGMs with three major clades: cell cycle, housekeeping, and tissue-specific functions, respectively. (B) Gene t-SNE plot of the CGMs along with names of the representative ones. Each dot is an expressed gene, and genes in one CGM are labeled with a particular color. (C) 3D genome structure of a GM12878 cell experimentally determined by scMicro-C (*38*). Yellow points represent the promoter positions of the genes in the DNA replication CGM. (D) Pairwise spatial distance matrix for the genes in the DNA replication CGM, averaged over all scMicro-C structures, suggesting the genes in a CGM are far apart from each other rather than clustered.

Fig. 4. Genes in a CGM sharing TF(s) in their promoters and enhancers. (A) Heatmap

showing the TF enrichment (-log *p*-value) within the major CGMs. Each row indicates a CGM

and each column indicates TFs assigned by ChIP-seq to the measured footprints. Color density

denotes the -log *p*-values calculated by the hypergeometric test. (B) Heatmap showing the TF pairs

enrichment (-log *p*-value) within the major CGMs. Each row indicates a CGM and each column

indicates TF pairs. Similarly, the -log *p*-values were calculated by the hypergeometric test. (C)

NRF1-KLF5-NR2C1 TF combination at promoters regulating the genes in the M4 RNA binding

and mitochondrial translation CGM. The solids lines are between a gene that connects more than

one TFs, whereas the dash lines are between a gene with only one TF specified (and other un-

assigned TFs). (D) STAT3-MEF2C-NFATC3 TF combination at enhancers regulating the genes

in the MD48 regulation of immune response CGM.

Fig. 5. Cell-type-specific TF footprinting in heterogeneous tissues by scFOODIE. (A) Cells from a mixture of four human cell lines are classified into four different cell types based on their barcoded scFOODIE data. Unsupervised clustering was performed using UMAP. CTCF and NRF1 footprints at two genome locations from pooled cells of the GM12878 cell type are shown in the inset. (B) Cell typing of ~1500 mouse hippocampus cells based on their scFOODIE data. Different colors indicate distinct cell types, astrocytes, microglia, oligodendrocytes, and neurons (including inhibitory and excitatory neurons), with annotations based on the marker genes. Footprints of JUN and MEF2B in neurons at merged genome-wide binding sites are shown in the inset.

Materials and Methods:

<u>Plasmid construction</u>

Expression constructs for the DddA toxin domain (DddAtox) and DddB were obtained from

Genescript through a gene synthesis service. To generate a pETDuet-1-based expression construct

for DddAtox, the DddAtox encoding sequence:

ATGGGCAGCAGCCATCACCATCATCACCACAGCCAGGATCCGGGTAGCTATGCGCT

GGGTCCGTATCAGATCTCTGCTCCGCAGCTGCCGGCATATAACGGTCAGACTGTTGG

TACTTTCTATTATGTTAACGATGCTGGCGGTTTAGAAAGCAAAGTTTTCAGCTCTGGT

GGTCCGACCCCGTATCCGAACTATGCTAACGCTGGTCACGTTGAAGGTCAGTCTGCT

CTGTTCATGCGTGATAACGGTATCTCTGAAGGTCTGGTTTTCCATAACAACCCGGAA

GGTACCTGTGGTTTTTGTGTTAACATGACCGAAACCCTGCTGCCGGAAAACGCTAAA

ATGACCGTTGTTCCGCCGGAAGGTGCGATTCCGGTTAAACGTGGTGCTACCGGTGAA

ACCAAAGTTTTCACCGGTAACTCTAACTCTCCGAAATCTCCGACCAAAGGTGGTTGC

TAA (of which the translated protein sequence is:

MGSSHHHHHHSQDPGSYALGPYQISAPQLPAYNGQTVGTFYYVNDAGGLESKVFSSGG

PTPYPNYANAGHVEGQSALFMRDNGISEGLVFHNNPEGTCGFCVNMTETLLPENAKMT

VVPPEGAIPVKRGATGETKVFTGNSNSPKSPTKGGC.), is synthesized and cloned into the

MCS-1 (NcoI and HindIII sites, retaining an N-terminal hexahistidine tag), and the immunity

protein DddAI encoding sequence:

ATGTATGCGGATGACTTTGACGGGGAAATTGAGATTGATGAAGTTGATAGCCTAGTT

GAGTTTCTGAGCCGTCGTCCGGCGTTCGATGCGAACAACTTCGTTCTGACCTTCGAA

GAAAGCGGCTTCCCGCAGCTGAACATCTTCGCGAAAAACGATATCGCGGTTGTTTAC

TACATGGATATCGGCGAAAACTTCGTTAGCAAAGGCAACAGCGCGAGCGGCGGCAC

CGAAAAATTCTACGAAAACAAACTGGGCGGCGAAGTTGATCTGAGCAAAGATTGCG

TTGTTAGCAAAGAACAGATGATCGAAGCGGCGAAACAGTTCTTCGCGACCAAACAG

CGTCCGGAACAGCTGACCTGGAGCGAACTGTAA (of which the translated protein

sequence is:

MYADDFDGEIEIDEVDSLVEFLSRRPAFDANNFVLTFEESGFPQLNIFAKNDIAVVYYMD

IGENFVSKGNSASGGTEKFYENKLGGEVDLSKDCVVSKEQMIEAAKQFFATKQRPEQLT

WSEL.), is synthesized and cloned into the MCS-2 (NdeI and XhoI sites, removing an C-terminal

S-tag). To generate a pETDuet-1-based expression construct for DddB, the DddB encoding

sequence:

ATGGGCAGCAGCCATCACCATCATCACCACAGCCAGGATCCTGGTTGGAAATTTTCT

AACGGTAAACGCCGTCCGCCGCACAAAGCAACGGTAACTGTGACCGATAAAAACGG

TGTCGTTAAACACAAAAGCAACCTGGTTTCTGGCAACATGACTGAAGCCGAAAAGA

AACTGGGCTTCCCGAACAACTCCCTGGCGACCCACACCGAAAACCGTGCTACCCGC

CTGATCGATCTGAACCAAGGTGATACTATGCTGATCGAGGGCCAATACCGTCCGTGT

CCACGTTGTAAAGGTGCAATGCGCGTGAAAGCGGAGGAATCCGGTGCGAAAGTGAT

CTACACCTGGCCAGAAGATGGTGACCTGAAAAAACGTGAATGGGAAGGCACTCCGT

GCGACAAAAAATAA   (of   which   the   translated   protein   sequence   is:

MGSSHHHHHHSQDPGWKFSNGKRRPPHKATVTVTDKNGVVKHKSNLVSGNMTEAEK

KLGFPNNSLATHTENRATRLIDLNQGDTMLIEGQYRPCPRCKGAMRVKAEESGAKVIYT

WPEDGDLKKREWEGTPCDKK.), is synthesized and cloned into the MCS-1 (NcoI and HindIII

sites, retaining an N-terminal hexahistidine tag), and the immunity protein DddBI encoding

sequence:

ATGACCAAATCTAAAATGCTGAGCAACATCGTCATCCAGGAGGTCAAATTTGCGATC

GAAGATTACTGCGCTATTCTGAGCTTCGCTTCTGACTCTTATGAAGTGCCGGAGCAG

TATTTTATCATTACCCGTTCTACCACCGAACGTTCTGGCGGTATTCCGGAGGGCGAC

ATCTACCTGGAATCTAACCTGTTTCTGGATTTTAACCCGTACGGCCTGAGCGGTTACC

TGCTGTCTGAGCCGAACTGCGTAGATCTGCTGATCGAACCGAACAACTACGTTCGTC

TGCGTCTGATCGAAAAAATCGATATCCTGGAAGTGGAAAACCACCTGAAATTTCTGT

TCGACAACTAA   (of   which   the   translated   protein   sequence   is:

MTKSKMLSNIVIQEVKFAIEDYCAILSFASDSYEVPEQYFIITRSTTERSGGIPEGDIYLESN

LFLDFNPYGLSGYLLSEPNCVDLLIEPNNYVRLRLIEKIDILEVENHLKFLFDN.),   is

synthesized and cloned into the MCS-2 (NdeI and KpnI sites, removing an C-terminal S-tag). The

pETDuet-1::dddAtox + dddAI vector and pETDuet-1::DddBtox + DddBI were transformed into

E. coli strains DH5α BL21 and stored at -20 degrees.

<u>Bacterial strains and culture conditions</u>

Unless otherwise noted, all Escherichia coli (E. coli) strains used in this study were grown in

Lysogeny Broth (LB) at 37 °C or on LB medium solidified with agar (LBA, 1.5% w/v). The media

was supplemented with ampicillin (100 µg per ml) or IPTG (0.5 mM) if it is required. E. coli

strains DH5α and BL21 were used for plasmid maintenance and protein expression, respectively.

<u>Purification of DddA enzyme</u>

The purification of the DddA toxin domain (DddAtox) protein has been reported previously

(Beverly et al. 2020, Nature). Briefly, to purify the his-tagged DddAtox in complex with DddAI,

E. coli BL21 (pETDuet-1::dddAtox + dddAI) was used to inoculate 2 L of LB broth in a 1:100

dilution and cultured overnight. After the culture reached an OD600 of approximately 0.6, 0.5 mM

IPTG was added, and the culture was incubated with shaking for 16 hours at 18˚C. Bacterial cell

pellets were collected by centrifugation at 4000 g for 30 minutes and then resuspended in 50 ml of

lysis buffer (50 mM Tris-HCl pH 8.0, 500 mM NaCl, 10 mM imidazole, 1 mg/mL lysozyme, and

a protease inhibitor cocktail). The cells were lysed by sonication (five pulses, 10 seconds each),

and the supernatant was separated from the debris by centrifugation at 25,000 g for 30 minutes.

The his-tagged DddAtox–DddAI complex was purified from the supernatant using a Nickel

column and eluted with elution buffer (50 mM Tris-HCl pH 7.5, 300 mM imidazole, 500 mM

NaCl, 30 mM imidazole, and 1 mM DTT). The eluted DddAtox–DddAI complex was denatured

by adding 50 ml of 8 M urea denaturing buffer (50 mM Tris-HCl pH 7.5, 300 mM NaCl, and 1

mM DTT) and incubated for 16 hours at 4˚C. The denatured proteins in the 8 M urea denaturing

buffer were reloaded onto the Nickel column, which was then washed with 50 ml of the same

buffer to remove any remaining DddAI. Sequential washes were performed with denaturing buffer

containing decreasing concentrations of urea (6 M, 4 M, 2 M, 1 M), followed by a final wash with

buffer without urea. DddAtox bound to the column was eluted with 5 ml of elution buffer. The

eluted DddAtox was subjected to size-exclusion chromatography using fast protein liquid

chromatography (FPLC) on a Superdex200 column (GE Healthcare) with sizing buffer (20 mM

Tris-HCl pH 7.5, 200 mM NaCl, 1 mM DTT, and 5% (w/v) glycerol). The purity of the eluted

DddAtox was assessed by SDS-PAGE gel stained with Coomassie blue, and the protein was stored

at -80˚C.

<u>Purification of DddB enzyme</u>

To purify the his-tagged DddB toxin domain (DddBtox) in complex with DddBI, *E. coli* BL21

(pETDuet-1::dddBtox + dddBI) was used to inoculate 2 L of LB broth in a 1:100 dilution and

cultured overnight. After the culture reached an OD600 of approximately 0.6, 0.5 mM IPTG was

added, and the culture was incubated with shaking for 16 hours at 18˚C. Bacterial cell pellets were

collected by centrifugation at 4000 g for 30 minutes and then resuspended in 50 ml of lysis buffer

(50 mM Tris-HCl pH 8.0, 500 mM NaCl, 10 mM imidazole, 1 mg/ml lysozyme, and a protease

inhibitor cocktail). The cells were lysed by sonication (five pulses, 10 seconds each), and the

supernatant was separated from the debris by centrifugation at 25,000 g for 30 minutes. The his-

tagged DddBtox–DddBI complex was purified from the supernatant using a Nickel column and

eluted with elution buffer (50 mM Tris-HCl pH 7.5, 300 mM NaCl, 30 mM imidazole, and 1 mM

DTT). The eluted DddBtox–DddBI complex was denatured by adding 50 ml of 8 M urea

denaturing buffer (50 mM Tris-HCl pH 7.5, 300 mM NaCl, and 1 mM DTT) and incubated for 16

hours at 4˚C. The denatured proteins in the 8 M urea denaturing buffer were reloaded onto the

Nickel column, which was then washed with 50 ml of the same buffer to remove any remaining

DddBI. Sequential washes were performed with denaturing buffer containing decreasing

concentrations of urea (6 M, 4 M, 2 M, 1 M), followed by a final wash with buffer without urea.

DddBtox bound to the column was eluted with 5 ml of elution buffer. The eluted DddBtox was

subjected to size-exclusion chromatography using fast protein liquid chromatography (FPLC) on

a Superdex200 column (GE Healthcare) with sizing buffer (20 mM Tris-HCl pH 7.5, 200 mM

NaCl, 1 mM DTT, and 5% (w/v) glycerol). The purity of the eluted DddBtox was assessed by

SDS-PAGE gel stained with Coomassie blue, and the protein was stored at -80˚C.

Crystallization and structure determination

Crystals of the DddB toxin-immunity protein complex were obtained at 5~12 mg/ml in

crystallization buffer (25 mM Tris pH 7.5, 200 mM NaCl, 1.0 mM DTT), mixed 1:1 with

crystallization solution containing 10% (w/v) PEG 8000, 0.1 M Potassium phosphate monobasic/

Sodium phosphate dibasic pH 6.2, 200 mM NaCl2. Rectangular crystals grew to $400 \times 200 \times 100$

μm over 5 days. DddB toxin-immunity protein crystals displayed the symmetry of space group

C2221 (a = 84.1 Å, b = 130.6 Å, c = 61.6 Å, α = β = γ = 90°). Before data collection, crystals were

cryoprotected in a crystallization solution of 20% glycerol, 10% (w/v) PEG 8000, 0.1 M Potassium

phosphate monobasic/ Sodium phosphate dibasic pH 6.2, 200 mM NaCl2. The X-ray diffraction

data were collected using a Rigaku XTALAB SYNERGY CUSTOM diffractometer with a Cu

target and were processed with CrysAlis(Pro) software.

The DddB toxin-Immunity protein complex structure was solved by molecular replacement using Phaser in Phenix (*41*), with the Alphafold2 predicted structure as the search model (*42*). Phenix (Adams et al., 2010) and Coot (*43*) were used for the standard refinement. The structure validation was completed with MolProbity (*44*). The data processing and refinement statistics are summarized in Table S2.

DNA deamination activity assays

Lambda DNA or genomic DNA extracted from K562 or GM12878 cell lines was used to evaluate the double-stranded DNA deamination activity. Reactions were performed in 10 μl of deamination buffer consisting of 20 mM Tris-HCl pH 7.4, 100 mM NaCl, 1 mM DTT, 50ng DNA substrate, and deaminase. Due to batch effects in enzyme purification, it is recommended to perform titration tests to reach the saturation of deamination. Reactions were incubated for 1 hr or the indicated time course at 37˚C, followed by DNA purification using Zymo DNA Clean & Concentrator-5 kit. Purified DNA was subjected to library preparation using TruePrep DNA Library Prep Kit V2 for Illumina (Vazyme) following the manual, except replacing the polymerase mix with 1× Q5U PCR master mix plus Bst 3.0 polymerase (0.08 U per μl). The uracil conversion ratio at each cytosine site was calculated and averaged to evaluate the deamination activity in each possible sequence context.

Cell preparation

Human cell line K562 and GM12878 were grown with RPMI 1640 medium (cat. no. 11875093, Gibco) containing 10% FBS (cat. no. 10099141C, Gibco) in the 5% CO2 incubator at 37°C. If necessary, cells were cultured in the 5% CO2 incubator at 42°C for 30min for heat shock treatment. When needed, 0.2 million cells were crosslinked in 0.2 ml of freshly prepared 1% PFA fixation buffer at room temperature (RT) for 10 min. To quench the fixation, the cells were rinsed once with 1 ml of DPBS (cat. no. 14040117, Thermofisher)-1% BSA (cat. no. A1933, Sigma), and further incubated with DPBS-1% BSA for 5 min.

FOODIE for bulk samples

1. Nuclei preparation. To prepare nuclei, fresh or fixed cells were spun down at 450g for 5 min, followed by washing using an equal volume of cold 1× PBS and centrifugation at 500g for 5

min at 4 °C. A total of 30,000 cells were permeabilized using cold permeabilization buffer (10 mM

Tris-HCl, pH 7.4, 10 mM NaCl, 3 mM MgCl2, 0.1% IGEPAL CA-630, 0.1% Tween-20, 0.1%

digitonin). Immediately after permeabilization, nuclei were spun down at 550g for 5 min at 4 °C.

Supernatant was carefully removed from the pellet after centrifugation.

2.   Tagmentation reaction. Prepare 2X TD buffer (20mM TAPS ph8.5, 10mM MgCl2, 20%

DMF). The nuclei pellet was immediately resuspended in the transposase reaction mix (25 μL 2×

TD buffer, 1 μL transposase (Vazyme, S111), and 24 μL PBS with 0.1% digitonin). The

transposition reaction was carried out for 30 min at 37 °C on a thermomixer at 800 rpm. For fixed

cells, the transposition reaction can be extended to 1.5 h. After that, 100 μl ice-cold RSB was added

to the mixer, which was followed by centrifugation at 550g for 5 min at 4 °C. Supernatant was

carefully removed from the pellet after centrifugation.

3.   Deamination reaction. Prepare 2X DRB buffer (20 mM Tris-HCl pH 7.5, 20 mM NaCl, 2

mM DTT). The nuclei pellet was immediately resuspended in the deaminase reaction mix. For

DddAtox, the deaminase reaction mix contains 12 μl DRB and 18 μl DddA enzyme (50 uM, stored

in sizing buffer). For DddB, the deaminase reaction mix contains 5 μl DRB and 5 μl DddB enzyme

(50 uM, stored in sizing buffer). Due to batch effects in enzyme purification, it is recommended to

perform titration tests to achieve optimal deamination performance. The nuclei mix was incubated

at 37°C for 10min. For fixed cells, the deamination reaction can be extended to 45 min, followed

by a proteinase K overnight digestion for de-crosslinking. The DNA was purified using a Zymo

DNA Clean & Concentrator-5 kit.

4.   Library amplification. After DNA purification, library fragments were amplified using 1×

Q5U PCR master mix, Bst 3.0 polymerase (0.08 U per μl) and 1.25 μM of Nextera PCR primers

(forward and reverse), using the following PCR conditions: 65 °C for 5 min; 80 °C for 12 min;

98 °C for 2 min; and thermocycling at 98 °C for 15 s, 60 °C for 30 s and 72 °C for 1 min. It is

recommended to monitor the PCR reaction using qPCR to stop amplification before saturation and

avoid GC and size bias in PCR. After a pre-amplification of full libraries for five cycles, an aliquot

of the PCR reaction was collected for qPCR. A total of 20 cycles of qPCR was performed to

determine the additional number of cycles needed for the remaining PCR reaction. Generally, a

total of 10–12 cycles of amplification yields high-quality libraries. The libraries were purified

using a Zymo Select-a-Size DNA Clean & Concentrator Kit to collect DNA with fragment size

above 200bp. Size-selected libraries are ready for sequencing.

FOODIE for single cells

Single-cell FOODIE in four cell lines was performed separately in the same experiment on the same day. Cultured cells with more than 95% viability were collected freshly washed with ice-cold PBS twice and counted with cell counters (Invitrogen). 50, 000 cells of each cell line were used following the above bulk FOODIE procedure, except the incubation temperature was reduced to 24°C. After the STOP step, the nuclei pellet was diluted into 1 mL ice-cold PBS. DAPI (Sigma) was incubated in 1:1000 with the nuclei for 10 minutes in the dark on ice. Then the stained single nuclei were sorted into single wells of a 96-well plate containing 2ul lysis buffer (30mM Tris-HCL pH7.5, 20mM NaCl, 0.5% SDS, 1mM DTT, 0.4ug/ml QP) by FACS machine (BD Aria II). After the lysis step at 50C for 30min, distinct cell barcodes were added to different wells through primers, and a 10 μL volume of PCR reaction was conducted using 22 cycles. More cycles should be considered when dealing with other cell types. 3μL of the PCR product of each well in one plate was pooled together and purified using 1.1X Ampure beads twice for size selection.

The C57BL/6N mouse strain is utilized in the scFOODIE project. The procedure for isolating mouse hippocampal nuclei follows that outlined in reference (*45*), while the beginning steps for dissociated mouse brain cells were performed as previously described in bulk FOODIE. Post deaminase incubation, add 0.2 μl of 1 mg/ml DAPI per 100 μl reaction for 5 minutes for staining on ice. Subsequently, sorting single cell to a 96-well plate containing 4 μl lysis buffer (20 mM Tris-HCl pH 7.5, 20 mM NaCl, 0.4% SDS) using flow cytometry. Incubate the 96-well plate at 65°C for 15 minutes for nuclear lysis. Finally, add 8 μl of PCR mix (2×Q5U master mix 6μl, 10% Tween20 2μl), and amplify as follows: 72 °C for 5 min; 98 °C for 1 min; and cycling for 22 cycles at 98 °C for 10 s, 63 °C for 30 s and 72 °C for 1 min. After PCR, wells from the same plate are pooled and purified using 1×Ampure beads for sequencing.

FOODIE data preprocessing

Raw paired-end reads were adapter-trimmed using Trim Galore (version 0.6.10) (*46*) and then mapped to the hg38 genome and deduplicated using Bismark Bisulfite Mapper (version 0.24.2) (*47*). We then used custom Python scripts to extract C-to-T conversion states for deduplicated reads and generate a conversion ratio for every genome position. For correction of the binding fraction, we measured deamination efficiency along the naked human genome as a positive control,

from which we obtained conversion probability for all 4^6 (-4 to +2 bases adjacent to cytosine)

possible sequences. For *in vivo* FOODIE data, conversion ratios of any cytosine sites were

corrected by normalization with the positive control.

Single-cell FOODIE data processing

For single-cell FOODIE data analysis, adapter sequences are first trimmed using Trimmomatic

(version 0.39) (*48*), followed by mapping to the mouse reference genome mm10 with Bismark

(version 0.24.0) (*47*). Duplicated reads are subsequently removed using the "deduplicate_bismark"

method within Bismark. For the single-cell data, the cytosines on the deduplicated reads are then

extracted for calculating the "non-conversion ratio" of detected genes. The resulting non-

conversion ratio matrix undergoes filtering for cell clustering analysis using Scanpy (*49*), requiring

a minimum of 600 genes per cell with a non-zero non-conversion ratio and each gene is detected

in at least 5 cells. Further analysis includes Principal Component Analysis using "scanpy.tl.pca,"

construction of a neighborhood graph with "scanpy.pp.neighbors," and dimension reduction via

"scanpy.tl.umap" for UMAP visualization. Subsequently, clusters are identified using the

"scanpy.tl.leiden" algorithm. Cluster-specific marker genes are analyzed using

"scanpy.tl.rank_genes_groups," with the top 20 ranked genes selected for each cluster for cell type

annotation. Cell type annotation is performed by matching the marker genes of each cluster against

the CellMarker 2.0 database (*50*). The cell type with the highest number of matched genes is

assigned to the respective cluster.

De novo footprinting

We used a custom-modified method to de novo call TF binding footprints (original code from

https://noble.gs.washington.edu/proj/footprinting). In brief, we used a set of sliding windows with

different sizes to determine footprint positions. Windows with a conversion ratio significantly

lower than flanking regions are considered footprints.

Assignment of TF recognition sequences to FOODIE footprints

To assign TF recognition sequences to FOODIE footprints, *de novo* called footprints were

overlapped with TF recognition sequences. TF recognition sequences were downloaded from

JASPAR (*23*) website (https://frigg.uio.no/JASPAR/JASPAR_TFBSs/2024). A motif that

overlaps with a footprint of more than 50% length of the motif was reported for further analysis.

When needed, ChIP-seq data was downloaded from the ENCODE Project (*27*) for TF assignments,

using a principle that TF recognition sequences fully covered by the corresponding ChIP-seq peaks

were used for further analysis.

<u>Statistic analyses</u>

To investigate the potential cooperativity among transcription factors (TFs), we analyzed

patterns of cytosine conversion at both sites. For each cytosine pair, we constructed 2x2

contingency tables to capture the bivariate frequency distribution of their conversion events. We

then performed Fisher's exact test on each table, testing the null hypothesis that the conversion

events between cytosines are independent. The type of cooperativity (positive or negative) was

determined based on the Φ correlation coefficients derived from these tables. Specifically, a

positive Φ coefficient indicates positive cooperativity, while a negative Φ coefficient suggests

negative cooperativity. We retained cytosine pairs with significant *P*-values ($P < 0.001$) for

constructing a pairwise correlation map. In these maps, individual TFs exhibited strong self-

correlation, depicted by a densely populated triangular area with positive Φ values. The

cooperativity between two TFs was represented by the cross-section of two individual binding

sites, displaying either positive or negative patterns.

Given the binding sites of a TF, we can identify the most frequently occurring modules. Since

some modules are larger than others, they tend to show up more frequently for any TF. Thus, it is

valuable to identify the statistically significantly enriched modules. For a specific TF and a specific

module, if all analyzed modules have N total genes, among which M are involved in the module,

and the TF binds to the promoters (or enhancers) of n genes, among which m are involved in the

same module, the *p* value is calculated as follows:

688
$$p = 1 - \sum_{i=0}^{m-1} \frac{\binom{M}{i}\binom{N-M}{n-i}}{\binom{N}{n}}$$

689   Then we performed Benjamini-Hochberg FDR correction with default cutoff of 0.05.

References:

1. S. A. Lambert *et al.*, The Human Transcription Factors. *Cell* **172**, 650–665 (2018).
2. F. Jacob, J. Monod, Genetic regulatory mechanisms in the synthesis of proteins. *Journal of molecular biology* **3**, 318-356 (1961).
3. M. Ptashne, ISOLATION OF THE lambda PHAGE REPRESSOR. *Proc Natl Acad Sci U S A* **57**, 306-313 (1967).
4. J. Elf, G.-W. Li, X. S. Xie, Probing transcription factor dynamics at the single-molecule level in a living cell. *Science* **316**, 1191-1194 (2007).
5. J. Yu, J. Xiao, X. Ren, K. Lao, X. S. Xie, Probing gene expression in live cells, one protein molecule at a time. *Science* **311**, 1600-1603 (2006).
6. P. J. Mitchell, R. Tjian, Transcriptional regulation in mammalian cells by sequence-specific DNA binding proteins. *Science* **245**, 371-378 (1989).
7. J. C. M. Gebhardt *et al.*, Single-molecule imaging of transcription factor binding to DNA in live mammalian cells. *Nature methods* **10**, 421-426 (2013).
8. C. Nüsslein-Volhard, E. Wieschaus, Mutations affecting segment number and polarity in Drosophila. *Nature* **287**, 795-801 (1980).
9. A. R. Chapman *et al.*, Correlated gene modules uncovered by high-precision single-cell transcriptomics. *Proceedings of the National Academy of Sciences* **119**, e2206938119 (2022).
10. M. Hu *et al.*, Correlated Protein Modules Revealing Functional Coordination of Interacting Proteins Are Detected by Single-Cell Proteomics. *The Journal of Physical Chemistry B* **127**, 6006-6014 (2023).
11. J. M. Replogle *et al.*, Mapping information-rich genotype-phenotype landscapes with genome-scale Perturb-seq. *Cell* **185**, 2559-2575. e2528 (2022).
12. P. Langfelder, S. Horvath, WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics* **9**, 559 (2008).
13. B. Ren *et al.*, Genome-Wide Location and Function of DNA Binding Proteins. *Science* **290**, 2306-2309 (2000).
14. G. Robertson *et al.*, Genome-wide profiles of STAT1 DNA association using chromatin immunoprecipitation and massively parallel sequencing. *Nature Methods* **4**, 651-657 (2007).
15. D. S. Johnson, A. Mortazavi, R. M. Myers, B. Wold, Genome-Wide Mapping of in Vivo Protein-DNA Interactions. *Science* **316**, 1497-1502 (2007).
16. J. R. Hesselberth *et al.*, Global mapping of protein-DNA interactions in vivo by digital genomic footprinting. *Nature Methods* **6**, 283-289 (2009).
17. C. Sönmezer *et al.*, Molecular Co-occupancy Identifies Transcription Factor Binding Cooperativity In Vivo. *Molecular Cell*, (2020).
18. B. Y. Mok *et al.*, A bacterial cytidine deaminase toxin enables CRISPR-free mitochondrial base editing. *Nature* **583**, 631-637 (2020).
19. L. M. Iyer, D. Zhang, I. B. Rogozin, L. Aravind, Evolution of the deaminase fold and multiple origins of eukaryotic editing and mutagenic nucleic acid deaminases from bacterial toxin systems. *Nucleic acids research* **39**, 9473-9497 (2011).
20. M. H. de Moraes *et al.*, An interbacterial DNA deaminase toxin directly mutagenizes surviving target populations. *eLife* **10**, e62967 (2021).

21. J. D. Buenrostro, P. G. Giresi, L. C. Zaba, H. Y. Chang, W. J. Greenleaf, Transposition of
native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-
binding proteins and nucleosome position. *Nature methods* **10**, 1213-1218 (2013).
22. M. Åkerfelt, R. I. Morimoto, L. Sistonen, Heat shock factors: integrators of cell stress,
development and lifespan. *Nature Reviews Molecular Cell Biology* **11**, 545-555 (2010).
23. I. Rauluseviciute *et al.*, JASPAR 2024: 20th anniversary of the open-access database of
transcription factor binding profiles. *Nucleic Acids Research* **52**, D174-D182 (2024).
24. I. E. Vorontsov *et al.*, HOCOMOCO in 2024: a rebuild of the curated collection of
binding models for human and mouse transcription factors. *Nucleic Acids Research* **52**,
D154-D163 (2024).
25. A. Jolma *et al.*, Multiplexed massively parallel SELEX for characterization of human
transcription factor binding specificities. *Genome Research* **20**, 861-873 (2010).
26. S. Mukherjee *et al.*, Rapid analysis of the DNA-binding specificities of transcription
factors with DNA microarrays. *Nature Genetics* **36**, 1331-1339 (2004).
27. The ENCODE (ENCyclopedia Of DNA Elements) Project. *Science* **306**, 636-640 (2004).
28. M. J. MacPherson, P. D. Sadowski, The CTCF insulator protein forms an unusual DNA
structure. *BMC Molecular Biology* **11**, 101 (2010).
29. W. Reith, S. LeibundGut-Landmann, J.-M. Waldburger, Regulation of MHC class II gene
expression by the class II transactivator. *Nature Reviews Immunology* **5**, 793-806 (2005).
30. A. Dixit *et al.*, Perturb-Seq: Dissecting Molecular Circuits with Scalable Single-Cell
RNA Profiling of Pooled Genetic Screens. *Cell* **167**, 1853-1866.e1817 (2016).
31. D. M. Suter *et al.*, Mammalian Genes Are Transcribed with Widely Different Bursting
Kinetics. *Science* **332**, 472–474 (2011).
32. A. Raj, C. S. Peskin, D. Tranchina, D. Y. Vargas, S. Tyagi, Stochastic mRNA Synthesis
in Mammalian Cells. *PLOS Biology* **4**, e309 (2006).
33. J. R. Chubb, T. Trcek, S. M. Shenoy, R. H. Singer, Transcriptional Pulsing of a
Developmental Gene. *Current Biology* **16**, 1018–1025 (2006).
34. S. Chong, C. Chen, H. Ge, X. S. Xie, Mechanism of Transcriptional Bursting in Bacteria.
Cell **158**, 314–326 (2014).
35. D. E. Nelson *et al.*, Oscillations in NF-kappaB signaling control the dynamics of gene
expression. *Science* **306**, 704-708 (2004).
36. W. Ma, X. S. Xie, TriSHA: investigating modularity in high-precision single-cell
transcriptomic data. *In preparation*, (2024).
37. L. Tan, D. Xing, C.-H. Chang, H. Li, X. S. Xie, Three-dimensional genome structures of
single diploid human cells. *Science* **361**, 924-928 (2018).
38. H. Wu, J. Zhang, L. Tan, X. S. Xie, Extruding transcription elongation loops observed in
high-resolution single-cell 3D genomes. *bioRxiv*, 2023.2002. 2018.529096 (2023).
39. W. Zhu, P. H. Giangrande, J. R. Nevins, E2Fs link the control of G1/S and G2/M
transcription. *The EMBO Journal* **23**, 4615-4626-4626 (2004).
40. K. Taniguchi, M. Karin, NF-κB, inflammation, immunity and cancer: coming of age.
Nature Reviews Immunology **18**, 309-324 (2018).
41. A. J. McCoy *et al.*, Phaser crystallographic software. *J Appl Crystallogr* **40**, 658-674
(2007).
42. J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583–589 (2021).

43.   P. D. Adams *et al.*, PHENIX: a comprehensive Python-based system for macromolecular
structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221 (2010).
44.   C. J. Williams *et al.*, MolProbity: More and better reference data for improved all-atom
structure validation. *Protein Science* **27**, 293-315 (2018).
45.   L. Tan *et al.*, Changes in genome architecture and transcriptional dynamics progress
independently of sensory experience during post-natal brain development. *Cell* **184**, 741-
758 e717 (2021).
46.   F. Krueger, T. Galore, A wrapper tool around Cutadapt and FastQC to consistently apply
quality and adapter trimming to FastQ files. *Cambridge, UK: Babraham Institute*,
(2015).
47.   F. Krueger, S. R. Andrews, Bismark: a flexible aligner and methylation caller for
Bisulfite-Seq applications. *bioinformatics* **27**, 1571-1572 (2011).
48.   A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120 (2014).
49.   F. A. Wolf, P. Angerer, F. J. Theis, SCANPY: large-scale single-cell gene expression
data analysis. *Genome Biology* **19**, 15 (2018).
50.   C. Hu *et al.*, CellMarker 2.0: an updated database of manually curated cell markers in
human/mouse and web tools based on scRNA-seq data. *Nucleic Acids Res* **51**, D870-
D876 (2023).

 Supplementary Materials

822   Figure S1. (A) Percentage of TF motifs on the human genome that can be assessed by DddA

823   (probing every TC) and DddB (probing every C) enzymes. (B) Comparsion of cystal sturcutes of

824   DddA and DddB.

843 Figure S2. (A) Sequence logo of DddB conversion yields for a cytosine on naked DNA *in vitro*.

844 (B) Bases from -4 to +2 positions of cytosine show relatively large bias on conversion.

860 Figure S3. Enhancers harbor more enriched TF pairs than promoters. (A) Heatmap showing the

861 number of modules, in which the TF pairs in a promoter are enriched. Color bar on the top denotes

862 the number of enriched modules. (B) Heatmap showing the number of modules, in which the TF

863 pairs in a enhancer are enriched. Color bar on the top denotes the number of enriched modules.