The effects of 17α-estradiol treatment on endocrine system revealed by single-nucleus

transcriptomic sequencing of hypothalamus

Lei Li[#,*1], Guanghao Wu[#2], Xiaolei Xu[3], Junling Yang[4], Lirong Yi[5], Ziqing Yang[6], Zheng Mo[3], Li

Xing[3], Ying Shan[*1], Zhuo Yu[*3], Yinchuan Li[*5]

[1]Department of Obstetrics and Gynecology, National Clinical Research Center for Obstetric &

Gynecologic Diseases, Peking Union Medical College Hospital, Chinese Academy of Medical

Sciences and Peking Union Medical College, 100730 Beijing, China.

[2]School of Medical Technology , Beijing Institute of Technology, 100081, Beijing, China.

[3]Department of Medical Oncology , Beijing Tsinghua Changgung Hospital, School of Clinical

Medicine, Tsinghua University, 102218, Beijing, China.

[4]Department of Cancer Biology and Pharmacology, University of Illinois College of Medicine,

Peoria, Illinois, United states of America.

[5]Institute of Reproductive Medicine, Medical School of Nantong University, Nantong, Jiangsu

226001, China.

[6]School of Basic Medical Sciences, Shandong University, Jinan, Shandong 250012, China.

# These authors contributed equally: Lei Li, Guanghao Wu

* Corresponding author. Email: shypumch@163.com; lilei64@pumch.cn; yza02214@btch.edu.cn;

18622397604@163.com

Abstract

In this study, we investigated the role of 17α-estradiol in lifespan extension and its potential side

effects from long-term administration. Pooled hypothalami from aged male Norway brown rats

treated with 17α-estradiol (O.T), aged male controls (O), and young male controls (Y) were

subjected to single-nucleus transcriptomic sequencing (snRNA-seq). To evaluate the effects of

17α-estradiol on aging neurons, supervised clustering of neurons based on neuropeptides and their

receptors were used to evaluate the responses of each neuron subtype during aging and after

17α-estradiol treatment. The elevated cellular metabolism, stress and decreased expression levels

of pathways involved in synapse formation in neurons initiated by aging were significantly

attenuated by 17α-estradiol. Assessment of changes in neuron populations showed that neurons

related to food intake, reproduction, blood pressure, stress response, and electrolyte balance were

sensitive to 17α-estradiol treatment. 17α-estradiol treatment not only increased serum Oxytocin

(Oxt), but also heightened the activity of hypothalamic-pituitary-gonadal (HPG) axis, as

evidenced by significantly elevated levels of plasma Gnrh, total testosterone, and decreased

estradiol. Elevated Gnrh1 was confirmed to be one of the causal effects mediating the role of

17α-estradiol in energy homeostasis, neural synapse, and stress response. Notably, *Crh* neurons

exhibited prominent stressed phenotype among all the checked neuron subtypes in O.T, which

may indicate a potential side effect of 17α-estradiol treatment. Therefore, the HPG axis and energy

metabolism may be key targets of 17α-estradiol in male hypothalamus. Additionally, supervised

clustering of neurons was shown to be a useful method for assessing treatment responses among

different neuron subtypes in the hypothalamus.

Key words snRNA-seq, hypothalamus, 17α-estradiol, aging, Crh, Oxt, Gnrh

Background

The hypothalamus serves as the central hub for controlling energy homeostasis, stress response,

temperature, learning, feeding, sleep, social behavior, sexual behavior, hormone secretion,

reproduction, osmoregulation, blood pressure, visceral activities, emotion, and circadian rhythms

[1]. The hypothalamic energy-sensing system, particularly the circuits that regulate food intake,

plays a crucial role in life span extension [2]. Elevated metabolic activity in the aged

hypothalamus has been reported in aged hypothalamus, including increased mTor signaling [3, 4].

Additionally, decreases in gonadotropin-releasing hormone (GnRH), Ghrh, Trh, monoamine

neurotransmitters, and blood supply are hallmarks of aging hypothalamus [5].

Previous studies have demonstrated that 17α-estradiol extends the lifespan of male mice and has

beneficial effects on metabolism and inflammation, similar to those of rapamycin and acarbose

[6-8]. Recent study indicated that 17α-estradiol also extends the lifespan of male rats [9].Further

investigations revealed certain unique features of 17α-estradiol in life extension distinct to

rapamycin and acarbose [10, 11]. Moreover, it has been shown that 17α-estradiol targets

hypothalamic *POMC* neurons to reduce metabolism by decreasing feeding behavior through

anorexigenic pathways [12]. Interestingly, the lifespan extension effect has only been observed in

male animals [13]. The safety of 17α-estradiol is key for translation into clinical treatment, and the

potential side effects on reproduction and feminization by 17α-estradiol treatment must be

considered. However, contradictory results have been reported regarding its side effects on

reproduction and feminization [6, 14, 15]. Therefore, further investigation and verification are

needed to understand the underlying mechanisms of lifespan extension and the safety of

17α-estradiol.

In this report, we utilized single-nucleus transcriptomic sequencing and performed supervised

clustering of neurons based on neuropeptides, hormones, and their receptors. Supervised

clustering offers better resolution in cell cluster screening compared to traditional unsupervised

clustering. We assessed the effects of 17α-estradiol on metabolism, stress responses, ferroptosis,

senescence, inflammation, and pathways involved in synaptic activity in each neuron subtype,

ranking the most sensitive neurons. The effects of 17α-estradiol on reversing aging-related cellular

processes were evaluated by two opposing regulatory networks involved in hypermetabolism,

stress, inflammation, and synaptic activity. Several key endocrine factors from serum were

examined, and the potential side effects of 17α-estradiol on specific neurons were also evaluated.

Materials and methods

Animals, treatment and tissues

Twelve Norway brown male rats (12 months old) were acquired from Charles River, including 8

12-months old and 4 1-month old (Beijing). Aged rats were randomly allocated into control and

17α-estradiol-treated groups. Four Aged rats treated with 17α-estradiol (Catalog #: E834897,

Macklin Biochemical, Shanghai, China) were fed freely with regular diet mixed with

17α-estradiol at a dose of 14.4 mg/kg (14.4 ppm), starting at 24 months of age for 6 months

according to prior reports [16, 17]. The young rats were fed a regular diet without 17α-estradiol

continuously for 3 months until 4 months old. All rats had ad libitum access to food and water

throughout the experiments. The rats were then euthanized via CO_2, hypothalami, testes and blood

serum were collected for subsequent experimental procedures. All blood samples were collected at

9:00-9:30 a.m to minimize hormone fluctuation between animals. All animal procedures were

reviewed and approved by the Institutional Animal Care and Use Committee at Nantong

University.

Enzyme immunoassays

Enzyme immunoassays kits for rat Oxt (Catalog #: EIAR-OXT), Corticotropin Releasing Factor (Catalog #: EIAR-CRF), and gonadoliberin-1 (Catalog #: EIAR-GNRH) were obtained from Raybiotech (GA, USA). Enzyme immunoassay kits for rat serum total testosterone (Catalog #: ml002868), estradiol (Catalog #: ml002891), aldosterone (Catalog #: ml002876), and cortisol (Catalog #: ml002874) were obtained from Enzyme-linked Biotechnology (Shanghai, China). Sera from 3 animals per group were used and each was diluted 10 or 20 times for immunoassays.

Seminiferous tubule inflammation test

8 testes were obtained from each sample group and then subjected to fixation in 4% formalin for at least 1 week. Formalin-fixed paraffin-embedded mouse testis sections of 5 µm thickness were used for HE Staining. At least 30 seminiferous tubules in each slide were examined for inflammation test. Testis with at least 1 inflammatory seminiferous tubule was set as 1, and normal testis was set as 0 for inflammation index calculation.

snRNA-seq data processing, batch effect correction, and cell subset annotation

Intact hypothalami were cryopreserved in liquid nitrogen from sacrificed rats. Two (O) or three (Y and O.T) hypothalami were pooled within each group and homogenized in 500 µL ice-cold homogenization buffer (0.25 M sucrose, 5 mM $CaCl_2$, 3 mM $MgAc_2$, 10 mM Tris-HCl [pH 8.0], 1 mM DTT, 0.1 mM EDTA, 1× protease inhibitor, and 1 U/µL RiboLock RNase inhibitor) with Dounce homogenizer. Then, the homogenizer was washed with 700 µL ice-cold nuclei washing buffer (0.04% bovine serum albumin, 0.2 U/µL RiboLock RNase Inhibitor, 500 mM mannitol, 0.1 mM phenylmethanesulfonyl fluoride protease inhibitor in 1× phosphate buffer saline). Next, the homogenates were filtered through a 70-µm cell strainer to collect the nuclear fraction. The nuclear fraction was mixed with an equal volume of 50% iodixanol and added on top of a 30% and 33% iodixanol gradient. This solution was then centrifuged for 20 min at 10 000 ×g at 4 °C. After the myelin layer was removed from the top of the gradient, the nuclei were collected from the 30% and 33% iodixanol interface. The nuclei were resuspended in nuclear wash buffer and resuspension buffer and pelleted for 5 min at 500 ×g at 4 °C. The nuclei were filtered through a 40-µm cell strainer to remove cell debris and large clumps, and the nuclear concentration was manually assessed using trypan blue counterstaining and a hemocytometer. Finally, the nuclei were adjusted to 700–1200 nuclei/µL, and examined with a 10X Chromium platform.

Reverse transcription, cDNA amplification and library preparation were performed according to

the protocol from 10X Genomics and Chromium Next GEM Single Cell 3′ Reagent Kits v3.1.

Library sequencing was performed on the Illumina HiSeq™ 4000 by Gene Denovo Biotechnology

Co., Ltd (Guangzhou, China).

10X Genomics Cell Ranger software (version 3.1.0) was used to convert raw BCL files to FASTQ

files, and for alignment and counts quantification. Reads with low-quality barcodes and UMIs

were filtered out and then mapped to the reference genome. Reads uniquely mapped to the

transcriptome and intersecting an exon at least 50% were considered for UMI counting. Before

quantification, the UMI sequences were corrected for sequencing errors, and valid barcodes were

identified using the EmptyDrops method. The cell $\times$ gene matrices were produced via UMI

counting and cell barcodes calling. Cells with an unusually high number of UMIs ($\geq$8000) or

mitochondrial gene percent ($\geq$15%) were filtered out. Batch effect correction was performed by

SCTransform function built in Seurat V4.4.0.

Pathways, gene signatures, TFs and TF cofactors, cell communication

Gene sets and pathways were derived from Hallmark gene sets of MSigDB collections, the KEGG

pathway database, Reactome pathway database, and WikiPathways database, and some ontology

terms derived from the Gene Ontology (GO) resource. Mitochondrial pathways were derived from

MitoCarta3.0 [18]. Pathways, gene sets, and gene signatures were evaluated with the

PercentageFeatureSet function built into R package Seurat. TFs and TF cofactors were obtained

from AnimalTFDB 3.0 [19]. TFs and TF cofactors were further filtered with mean counts >0.1.

The ligand–receptor pairs were calculated via R package CommPath [20].

Correlation analysis and ROC analysis

Pearson correlation coefficient was calculated with the linkET package ($p < 0.05$). Fast Wilcoxon

rank sum test and auROC analysis was performed with the wilcoxauc function in R package

presto. The minimal cell number in either one of the comparing pairs should be no less than 15.

Ranks of area under the curve (AUC) values were in descending order. A total of 431 pathways

from Hallmark, KEGG and PID databases were used for correlation analysis with MitoCarta

OXPHOS subunits in neurons and non-neural cells (**Figure 2—figure supplement 1**). A total of

97 pathways related to synapse activity were derived from GO, including GO cellular components,

GO biological processes and GO molecular functions (**Table S1**).

The division of expression level-dependent clusters in each pathway and their gene signatures

The quarters of the mixed cell populations from O, O.T, and Y hypothalamic neurons were equally divided using the R function fivenum from the R package stats, based on pathway expression levels. Thus, the total number of neurons was evenly divided into four clusters (c1-c4) in terms of cell number. The cell proportions from O.T, O, and Y neurons in each cluster were weighted against the total number of neurons in the three groups. The unique markers of each cluster were calculated using the FindAllMarkers function from the Seurat package. The intersection of the unique markers from the six pathways was obtained for heatmap plotting. Nineteen genes that were highly expressed in c1 were identified as c1.up.signature via the PercentageFeatureSet function in the Seurat package. Twelve genes that were highly expressed in c4 were identified as c4.up.signature. There were no intersecting unique markers in clusters c2 and c3 among the six selected pathways.

TF and pathway activities

The TF resources were derived from CollecTRI, the pathway resource was from PROGENy, and the enrichment scores of TFs and pathways were performed with the Univariate Linear Model (ulm) method according to the pipeline in R package decoupleR [21].

Subtypes of neurons generated by supervised clustering

Vast majority of these subtypes were clustered by neuropeptides, hormones, and their receptors within all the neurons with the subset function from R package Seurat (the target gene expression level > 0). A total of 121 neuron subtypes were obtained, comprising 80 neuropeptide-secreting neurons and 41 neurons expressing a unique neuropeptide receptor or hormone receptor. Further groupings may exist within the identified neuron subtypes, and the category of excitatory or inhibitory neurons was not discriminated further. The cell proportion of each neuron subtype was weighted according to the total number of neurons in O.T, O, and Y samples. The mean values ± standard deviation of pathways and gene signatures were performed for each subtype. The top 10 and the bottom 10 items were calculated, and the top 10 subtypes ranked within each of 16 pathways or gene signatures were collected for word clouds annotation via R package wordcloud2.

Differential expression and pathway enrichment analysis

DEGs between groups were identified via FindMarkers (test.use = bimod, min.pct = 0.1,

logfc.Threshold = 0.25, avg_diff > 0.1 or < −0.1). DEGs were then enriched in redundant GO

terms via WebGestalt and filtered with false discovery rate < 0.05 [22].

Bidirectional Mendelian randomization (MR) study

The protein quantitative trait locus (pQTL) GWAS summary data of 204 human endocrine-related

GWAS summary data with European ancestry were obtained from open-access MRC Integrative

Epidemiology Unit (IEU) (**Table S4**) [23, 24]. Independent genome-wide significant SNPs for

exposure OXT (id:prot-a-2159) or GNRH1 (id: prot-a-1233) were used as instrumental variables

with genome-wide significance ($P < 1 \times 10^{-5}$), independence inheritance (r2 < 0.001) without

linkage disequilibrium (LD) with each other for MR. For the reverse MR, independent

genome-wide significant SNPs from 203 endocrine-related GWAS summary data ($P < 1 \times 10^{-5}$, r2

< 0.001) without LD with each other were obtained as exposures and human GWAS summary

data of OXT (id:prot-a-2159) or GNRH1 (id:prot-a-1233) were used as outcomes. Weak

instruments less than 10 were discarded via F-statistics.

MR and reverse MR analysis were conducted with method inverse-variance weighting (IVW),

MR Egger, Weighted median, Simple mode, and Weighted mode. The screening criteria: all of the

odds ratio (OR) values of the 5 methods should be simultaneously either >1 or <1 and the

significant p value of IVW was <0.05. The heterogeneity via IVW method and the horizontal

pleiotropy were also evaluated with R package TwoSampleMR [25].

Results

The overall changes in aged hypothalamus with or without long-term 17α-estradiol

treatment via snRNA-seq profiling

To investigate the hypothalamus as a potential key target of 17α-estradiol's effects on life

extension, we performed snRNA-seq on the entire hypothalamus of aged and 17α-estradiol-treated

aged Norway brown rats, using the hypothalamus from young adult male rats as a control. We

identified 10 major cell types based on specific cell markers of the hypothalamus (**Figure 1A-B**).

Notably, the proportions of all non-neural cells changed in O versus Y (**Figure 1C**). For instance,

Oligo, OPC, and Micro were found to be increased, while Astro, Tany, Fibro, PTC, and Endo were

decreased in O compared to Y. The proportions of Oligo, OPC, and Micro were also increased in

17α-estradiol-treated aged group (O.T) compared to those in Y. Furthermore, Endo was increased

in O.T compared to both Y and O. The proportions of Astro, Tany, Epen, and PTC decreased more

in O.T than those in O when compared to Y. These results indicated that 17α-estradiol treatment

had extensive effects on the proportions of non-neural cells in hypothalamus.

Cell communication analysis revealed significant changes in the ligand-receptor pairs between

neurons and other cell types, particularly in Endo, Fibro, Tany, and Astro (**Figure 1D**). Significant

ligand–receptor interactions among neurons also changed in O.T and O groups, especially in O

(**Figure 1E**). Notably, among the significant ligand–receptor pairs in neurons,

Bmp2–Acvr1/Acvr2a/Acvr2b/Bmpr, *Gdf11–Acvr2a/Acvr2b*, *Inhba–Acvr1/Acvr2a/Acvr2b*,

Nrg1/Nrg2/Nrg4–Erbb4, *Rspo1–Lgr5/Lrp6*, and *Rspo3–Lgr5* were exclusively and significantly

increased in neurons of the O group compared to those in O.T and Y, suggesting enhanced TGF

superfamily-mediated signaling activity and canonical Wnt signaling during aging. The

significantly changed ligand–receptor pairs *Nlgn1–Nrxn1/Nrxn2*, *Nlgn2–Nrxn1/Nrxn2/Nrxn3*,

Nlgn3–Nrxn1/Nrxn2/Nrxn3, *Nxph1–Nrxn1/Nrxn2/Nrxn3*, *Nxph3–Nrxn1/Nrxn2/Nrxn3*,

Pomc–Oprd1/Oprk1/Oprm1, and *Vip–Adcyap1r1/Avpr1a/Vipr2* were exclusively increased in

neurons of O.T compared to O and Y (**Figure 1E**). These ligand–receptor pairs were associated

with synaptic activity, cellular adhesion, the opioid system, and vasodilation, indicating unique

roles of 17α-estradiol in restoring certain physiological functions in the aging hypothalamus. The

increased *Pomc* signal in O.T neurons aligns with previous reports suggesting that 17α-estradiol

treatment decreases food uptake in mice, potentially correlated with *Pomc* neurons (**Figure 1E**)

[12].

Gene set enrichment analysis (GSEA) based on DEGs also corroborated the expression profiles

related to stress responses and synapse-associated cellular processes in neurons (**Figure 1F**). ROC

analysis of significantly differently expressed pathways related to neural synapses, manually

selected from Gene Ontology databases, indicated that most top-ranked pathways related to

synapses, according to AUC values, were downregulated in aged neurons, while 17α-estradiol

treatment reversed this trend (**Figure 1G, Table S1**).

Overall, these findings suggest that 17α-estradiol broadly reshapes cell populations, cellular

communication, neuropeptide secretion, and synapse-related cellular processes in the aging

hypothalamus, distinguishing it from both the young hypothalamus and the untreated aged

hypothalamus.

The two opposing signaling networks in regulating metabolism and synapse activity, which

can be balanced effectively by 17α-estradiol

To monitor the metabolism and neural status affected by 17α-estradiol, we utilized the energy

metabolism pathway MitoCarta OXPHOS subunits to calculate the positively or negatively

correlated pathways in hypothalamic neurons (**Figure 2—figure supplement 1**). Our findings

revealed that energy metabolism and synapse activity represent two opposing regulatory signaling

networks in hypothalamic neurons, with 17α-estradiol strongly playing a significant role in

balancing these networks (**Figure 2A**). At the core of these opposing signaling pathways are two

categories of contrasting TFs (**Figure 2B**). For example, *Calr*, *Clu*, *Peg3*, *Prnp*, *Ndufa13*, *Actb*,

Ywhab, *Nfe2l1*, *Mtdh*, *Npm1*, *Bex2*, *Aft4*, and *Maged1* were positively correlated with pathways

involved in OXPHOS subunits, lysosome function, protein export, mTorc1 signaling, and the

unfolded protein response (UPR) in O, O.T and Y neurons, while showing negative correlations

with pathways related to ubiquitin-mediated proteolysis, endocytosis, tight junctions, focal

adhesion, axon guidance, and MAPK signaling. Additionally, TFs *Myt1l*, *Ctnnd2*, *Tenm4*, *Camta1*,

Med12l, *Rere*, *Csrnp3*, *Erbb4*, *Jazf1*, *Dscam*, *Klf12*, and *Kdm4c* exhibited opposite correlation

patterns with these selected pathways in O, O.T and Y neurons. These TFs may take conserved

roles in regulating the two opposing biological processes in hypothalamic neurons.

We then attempted to establish gene signatures to represent these two opposing signaling networks,

thereby displaying the cell status of aging and evaluating the effects exerted by 17α-estradiol. To

achieve this, we evenly divided the expression levels of each of the six selected pathways from the

two opposing signaling networks into four quarters (c1-c4) among the mixed neurons from O, O.T,

and Y, calculating the shared unique markers in each quarter (**Figure 2C, D**). From the distribution

patterns, we observed that the proportion of neurons in O decreased from c1 to c4 in metabolic

pathways (MitoCarta OXPHOS subunits and Hallmark mTorc1 signaling), while this trend was

reversed in the opposing signaling pathways (GOBP synapse organization and KEGG MAPK

signaling pathway) (**Figure 2C**). In contrast, in Y, this trend was opposite, suggesting the

expression levels from the 4 quarters (c1-c4) of the two opposing signaling networks can be used

to monitor aging status. Treatment with 17α-estradiol alleviated this trend or even reversed it in O.

We then screened the shared unique markers of each quarter from the six selected pathways in an

attempt to establish the gene signatures representing the two opposing signaling networks. Unique

markers in c1 (19 genes, c1.up.signature) and c4 (12 genes, c4.up.signature) were identified;

however, c2 and c3 lacked unique markers shared by the 6 pathways (**Figure 2D**). Consequently,

the 19 genes in c1.up.signature displayed an inversed correlation pattern with the 12 genes in

c4.up.signature, indicating the two opposing gene signatures are capable of reflecting the two

opposing signaling networks in hypothalamic neurons (**Figure 2E**). Conversely, the balance of the

two opposing signaling networks affected by 17α-estradiol in non-neural cell types was less

pronounced than in neurons, showing variable effects on non-neural cells (**Figure 2—figure**

supplement 2). GOBP pathway enrichment analysis revealed that Micro exhibited lower levels of

synapse-related cellular processes in O.T compared to O, which was distinct from the observations

in neurons (**Figure 2—figure supplement 3**). Therefore, in this report, we primarily focused on

hypothalamic neurons and their responses to aging and 17α-estradiol.

Supervised clustering revealed distinct responses of different subtypes of hypothalamic

neurons to aging and 17α-estradiol

The hypothalamus contains numerous neuron subtypes that release various neuropeptides and

hormones to regulate fundamental body functions. To differentiate the changes occurring during

aging and the effect of 17α-estradiol on each neuron subtype, we performed supervised clustering

based on neuropeptides, hormones, or their receptors (**Figure 3A**). Most of the top 10 neuron

subtypes involved in UPR, ferroptosis, mTorc1 signaling, insulin signaling pathway, immune

response pathways, OXPHOS subunits, and senescence signature during aging were attenuated by

17α-estradiol treatment (**Figure 3B, Figure 3—figure supplement 1**). For example,

Serpine1-secreting neurons exhibited very high expression levels of UPR, ferroptosis signature,

mTorc1 signaling, Tnfα signaling via NF-kB, and OXPHOS subunits during aging, all of which

were attenuated by 17α-estradiol.

To further identify the most sensitive neurons in response to aging and 17α-estradiol, we

calculated the frequency of each neuron subtype by combining the top 10 neurons across each of

the 16 pathways and gene signatures (**Figure 3C, Table S2**). Serpine1-secreting neurons were

exclusively enriched in aged hypothalamus, suggesting they are effective targets of 17α-estradiol.

Additionally, *Mlnr*, *Nmb*, *Pomc*, *Ednra*, *Serpine3*, *Gast*, and *Pcsk6* neurons were all affected by 17α-estradiol (**Figure 3D**). *Galp*, *Glp1r*, *Serping1*, *Sstr2*, *Sstr3*, and *Vip* neurons were enriched in O, O.T and Y hypothalamus, indicating a lack of effect of 17α-estradiol on these neuron types. The unique neurons in O.T (*Crh*, *Serpinb9*, *Cckbr*, *Pth2r*, *Kiss1*, *Prpr*, *Hcrtr1*, *Npb*, *Nppa*, *Nxph3*, and *Npff*) may be a consequence of compensatory effects from 17α-estradiol treatment.

We then evaluated the most and least sensitive neurons to aging by ranking the neuron subtypes according to the levels of c1.up.signature and c4.up.signature (**Figure 3E**). *Serpine1*, *Galp*, *Calca* neurons were among the most affected subtypes in the aged hypothalamus, consistent with the combined ranks of several pathways (**Figure 3—figure supplement 1**). *Npr3*, *Crh*, *Ar*, *Esr1*, and *Esr2* neurons were among the least affected subtypes in the aged hypothalamus (**Figure 3E**). In O.T, neuron subtypes ranked by c1.up.signature and c4.up.signature indicated that *Gast*, *Npb*, *Nppa*, and *Crh* were affected most by 17α-estradiol treatment (**Figure 3F**). Neurons *Oxtr*, *Glp1r*, *Gnrh1*, and *Crh* also showed the lowest levels of c4.up.signature, indicative of an aging phenotype. Meanwhile, neurons *Glp2r*, *Ar* and *Esr1* ranked among the top neurons based on c4.up.signature, suggesting higher expression levels of synapse-related processes and relatively less stress status in these types of neurons in O.T.

These results indicate that supervised clustering of each neuron subtype of neurons facilitated the visualization of the different responses from distinct subtypes of neurons in response to aging and medical treatments. Furthermore, it strongly suggests that *Crh, Oxtr, Glp1r, Gnrh1, Glp2r, Ar,* and *Esr1* neuron subtypes are particularly sensitive to long-term 17α-estradiol treatment, which is associated with appetite regulation, glucose metabolism, stress response, and, as expected, sex hormone secretion and signaling.

The potential side effects on hypothalamic–pituitary–adrenal (HPA) axis by long-term 17α-estradiol treatment in the males

To further investigate the potential side effects or compensatory effects of 17α-estradiol treatment, we performed stricter screening by intersecting the top 10 ranks in the UPR, OXPHOS subunits, and ferroptosis signature (**Figure 4A**). *Crh* and *Nppa* were the only two intersected neuron subtypes identified through this strict screening. The treatment with 17α-estradiol elevated several key metabolic pathways in *Crh* neurons compared to those in Y and O (**Figure 4B**). Additionally,

17α-estradiol treatment increased the c1-up-signature while simultaneously reducing many

pathways associated with synapse activity and the c4-up-signature in *Crh* neurons of O.T,

indicating a potent stressed phenotype in *Crh* neuron. In contrast, in *Nppa* and *Nppc* neurons, the

decreased c1-up-signature in O.T implied a lesser extent of stressed phenotype in these neurons

compared to *Crh* neurons. The aberrant changes in *Crh* neurons were also evidenced by the

increased number of DEGs related to mitochondria-expressed genes in O.T, along with a reduced

number of DEGs in the adherens junction pathway (**Figure 4C**). The status of Crh neurons in O.T

may be associated with elevated TF activities of *Esr2*, *Usf2*, *Hdac5*, *Creb3l1*, *Tfam*, *Preb*, *Pou3f2*,

and *Hoxb5* (**Figure 4D**).

Notably, the HPA axis was altered by 17α-estradiol treatment, as evidenced by the elevated

cortisol levels in O.T compared to O ($p = 0.078$) (**Figure 4E**). The correlation between elevated

cortisol production and the heightened stress in *Crh* neurons by 17α-estradiol treatment needs

further investigation. Additionally, as a crucial component of the renin-angiotensin-aldosterone

system, the significantly increased serum aldosterone in O.T and its potential role in sodium

reabsorption and cardiovascular health also warrant more detailed investigation (**Figure 4E**).

In summary, 17α-estradiol treatment altered the activity of HPA axis in male BN rats and

introduced increased potential side effects in *Crh* neurons.

17α-estradiol increased Oxt neuron proportion and secretion and its possible role in

mediating the effect of 17α-estradiol on endocrine system

Aging and 17α-estradiol treatment also altered the proportions of various neuron subtypes among

O, O.T, and Y (**Figure 5A, B, Table S3**). The proportions of *Grp*, *Pmch*, *Npb*, *Serpinb9*, *Sstr2*,

Agrp, *Sstr3*, *Mlnr*, and *Hcrt* neurons ranked in the top 10 in O, while *Oxt*, *Vip*, *Avp*, *Calca*, *Glp2r*,

Tacr1, *Trh*, *Serping1*, *Npff*, and *Npy1r* were in the top 10 in O.T. *Galp*, *Calcrl*, *Ednra*, *Oxt*,

Serpinh1, *Pomc*, *Cck*, *Crh*, *Tacr1*, and *Kiss1* neurons were among the bottom 10 in O, whereas

Oxtr, *Galp*, *Agrp*, *Serpinb9*, *Npvf*, *Serpinh1*, *Ednrb*, *Agt*, *Gipr*, and *Pomc* neurons ranked among

the bottom 10 in O.T. *Agrp*, *Pomc*, *Oxt*, *Oxtr*, *Gipr*, and *Glp2r* neurons are well-known for their

roles in regulating food intake and energy homeostasis. *Agrp* neurons are activated by hunger,

while *Pomc* neurons are activated by satiety in the ARC of the hypothalamus. 17α-estradiol

treatment effectively elevated the expression levels of the c4.up.signature and synapse-associated

processes in neuron subtypes *Agrp*, *Pomc*, *Oxt* and *Glp2r* in O.T compared to O. This may mitigate the adverse effects of reduced cell populations in Pomc and *Agrp* neurons in aging hypothalamus (**Figure 5C**). This finding indicates a potential role of 17α-estradiol in appetite control, as previously reported [12]. Additionally, 17α-estradiol treatment resulted in elevated proportions of *Vip*, *Avp*, *Npff*, *Calca*, and *Tacr1* neurons, which ranked in the top 10, while the proportions of Agt neurons decreased. These are all associated with blood pressure regulation (**Figure 5—figure supplement 1**). Notably, the proportions of *Oxt* and *Glp2r* neurons, both of which have anorexigenic effects [26, 27], increased in O.T. In addition to the increased number of *Oxt*-positive neurons, the expression level of *Oxt* also rose in O.T. The elevated expression of synapse-related pathways was supported by the increased DEGs in the enriched synaptic membrane pathway in *Oxt* neurons (**Figure 5D**).

More importantly, the serum level of Oxt was significantly elevated in O.T compared to O (p=0.04), although it remained lower than that in Y (**Figure 5E**). Notably, the top TF activities in O.T and O differed markedly from those in Y (**Figure 5F**). The elevated levels of *Hopx* and *Xbp1* may be associated with the response to 17α-estradiol treatment.

Due to the intricate regulatory networks among various endocrine factors, elucidating the causal effect of Oxt on other endocrine factors is quite complex using traditional methods. MR analysis, employing variant SNPs as genetic tools, is advantageous for such task. We performed a bidirectional MR analysis of the GWAS summary data of human plasma OXT and 203 endocrine-related and hypothalamus-related factors, most of which are protein quantitative trait loci (pQTL) data from the IEU (**Table S4**). As an exposure, OXT revealed a significant causal effect on POMC/beta-endorphin (id:prot-a-2327, id:prot-a-2325), glucagon (id:prot-a-1181), GNRH1/Progonadoliberin-1 (id:prot-a-1233), and total testosterone levels (id:ebi–a–GCST90012112, id:ieu–b–4864) (**Figure 5G**). NPW and CBLN1 were found to be negatively associated with OXT, but the significance of these associations was not found in the reverse MR analysis (**Figure 5—figure supplement 2A, B**).

In contrast, we could not identify significant associations between OXT and estradiol levels (id:ebi-a-GCST90012105, id:ebi-a-GCST90020092, id:ebi-a-GCST90020091, id:ieu-b-4872, id:ieu-b-4873, id:ukb-e-30800_AFR, id:ukb-e-30800_CSA). Interestingly, QRFP, IGF1, AGRP, TAC4, GRP, CLU, BNF, PCSK7, PACAP, ANP, TAC3, CRH, INSL6, and PRL displayed

significant associations with OXT in both MR and reverse MR analysis, indicative of their

complex causal effects (**Figure 5—figure supplement 2A, B**).

The results suggest that elevated Oxt levels induced by 17α-estradiol may have positive

associations with endocrine factors governing feeding behavior, glucose metabolism, male

reproduction, and sex hormones. Therefore, OXT may serve as a potential mediator of

17α-estradiol.

17α-estradiol activated HPG axis and the elevated Gnrh also took important roles in

mediating the effect of 17α-estradiol on other endocrine factors

Since Gnrh and sex hormone expressing neuron subtypes were sensitive to 17α-estradiol treatment

(**Figure 3F, Figure 3—figure supplement 1**), we examined their expression profiles alongside

representative pathways of two opposing signaling networks related to metabolism and synapses,

including c1-up-signature and c4-up-signature (**Figure 6A**). However, neither the c1-up-signature

nor the c4-up-signature was up-regulated in *Gnrh1* neuron in the O.T, similar to the observations

in *Esr2* neuron. Only in *Esr1* neuron was the c1-up-signature found to be up-regulated. Meanwhile,

both *Ar* and *Esr* neurons displayed high level of c4-up-signature, indicating a relatively healthy

status. The data suggest that most of the 4 types of sex hormone-related neurons did not

experience severe aging in O.T. However, based on these expression profiles, it's difficult to

define the precise physiological status of these subtypes of neurons, particularly regarding

neuroendocrine activities. Consequently, we performed enzyme immunoassays of hormones from

the serum of O, O.T and Y. The treatment with 17α-estradiol significantly increased the plasma

level of Gnrh compared to Y (p = 0.0099) and approached significance when compared to O (p =

0.096) (**Figure 6B**). More intriguingly, testosterone levels in serum were significantly increased in

O.T compared to O (p = 0.018) and approached significance when compared to Y (p = 0.052).

Additionally, the serum estradiol levels were significantly increased in O compared to Y (p =

0.011) and significantly decreased in O.T compared to O (p = 0.019), suggesting that 17α-estradiol

treatment markedly altered the homeostasis of testosterone and estradiol.

Furthermore, most testes from 30-month-old male BN rats exhibited severe age-related

inflammation and epithelial collapse of seminiferous tubules (**Figure 6C**). The testes without

inflammation in O.T displayed normal morphology. 17α-estradiol treatment slightly decreased the

testis inflammation in O.T compared to that in O (p = 0.15), indicating a potential positive role of

17α-estradiol treatment in male reproductive system. The elevated TFs such as *Sf1*, *Pparg*, *Litaf*,

Nupr1, *Rxrg*, *E2f2*, and *Zfp42* may be involved in the transcriptional regulation by 17α-estradiol in

O.T (**Figure 6D**). Importantly, the activities of androgen and estrogen pathways were decreased in

Gnrh1 neurons in O.T compared to O, and were distinct from those in *Ar*, *Esr1*, and *Esr2* neurons

(**Figure 6E**). These signaling pathways are important for the feedback control of sex hormone

secretion in *Gnrh* neurons, and these results may also reflect the strong effect of 17α-estradiol on

Gnrh neurons.

To decipher the potential effects of elevated serum Gnrh levels on endocrine system, we

performed bidirectional MR analysis of the GWAS summary data of human GNRH1 (id:

prot-a-1233) and 203 endocrine-related factors with genetic variants SNPs. We found strong

causal effects of GNRH1 on GAL/Galanin (id:prot−a−1166), POMC/Beta−endorphin

(id:prot−a−2327, id:prot−a−2325), Adrenomedullin (id:prot−a−48), BDNF (id:prot−a−242)**,** and

LPR (id:prot−a−1724), which are involved in feeding, energy homeostasis, osmotic regulation,

and neuron plasticity (**Figure 6F**). Notably, CRH/Corticotropin (id:prot−a−2326),

PRLH/Prolactin−releasing peptide (id:prot−a−2376), NPW/Neuropeptide W (id:prot−a−2082),

Glucagon (id:prot−a−1181), Chromogranin−A (id:prot−a−538) displayed bidirectional

significance, indicating close and complex causal effects between GNRH1 and these endocrine

factors (**Figure 6G, Figure 6—figure supplement 1A, B**). These results also suggest that the role

of 17α-estradiol treatment in feeding, energy homeostasis, reproduction, osmotic regulation, stress

response, and neuronal plasticity may be mediated, at least in part, by elevated Gnrh secretion.

Discussion

The most striking role of 17α-estradiol treatment revealed in this study showed that HPG axis was

substantially improved in the levels of serum Gnrh and testosterone. The underlying molecular

mechanism remains unclear; however, prior reports have indicated that 17α-estradiol can bind to

ESR1 [9]. In our findings, 17α-estradiol treatment significantly decreased serum estradiol levels

while elevating serum testosterone. Based on this evidence, we propose that 17α-estradiol may

function similarly to estrogen receptor antagonists or aromatase inhibitors, potentially preventing

the conversion of androgens to estrogens [28, 29]. These actions could alleviate the feedback

inhibition exerted by estrogen on hypothalamus and pituitary, thereby facilitating the secretion of Gnrh, FSH, and LH [30].

The testosterone levels in men gradually decline beginning in the third decade of life [31]. Age-related deterioration of the gonadotropic axis, particularly in older males with low serum testosterone, is often linked to numerous aging symptoms, including loss of skeletal muscle mass, reduced muscle strength and power, low bone mineral density, frailty, impaired physical performance, mobility limitations, increased risk of diabetes, elevated all-cause cardiovascular mortality, cognitive decline, and heightened risk of Alzheimer's disease [32]. Consequently, testosterone supplementation in older men is beneficial. Additionally, Gnrh supplementation may help mitigate age-related declines in neurogenesis and slow aging processes [33]. Importantly, treatment with 17α-estradiol did not result in feminization or adversely affect the sperm parameters and fertility in male animals [6, 14]. Thus, the observed increases in Gnrh and serum testosterone levels due to 17α-estradiol treatment are likely advantageous for older males, particularly those experiencing late-onset hypogonadism.

Postmenopausal women with low estrogen experience aging-related syndromes similar to those of older males with low serum testosterone. Those women also face increased mortality, cardiovascular disease, osteoporosis fracture, urogenital atrophy, and dementia, all of which may benefit from hormone therapy [34]. However, a prior report indicates that 17α-estradiol treatment does not provide positive life extension effects in aged females [13]. The discrepancy may stem from the inhibitory effects of estrogens associated with 17α-estradiol treatment, as evidenced by its inability to enhance female fertility [35]. Nonetheless, due to the lack of parallel data in aged female BN rats treated with 17α-estradiol, further research is needed to definitely address this question in the female subjects in the future.

Another notable effect of 17α-estradiol is its ability to reduce the overall expression levels of energy metabolism in hypothalamic neurons of aged male BN rats. The nutrient-sensing network, mediated by MTORC1 complex, is a central regulator of mRNA and ribosome biogenesis, protein synthesis, glucose metabolism, autophagy, lipid metabolism, mitochondrial biosynthesis, and proteasomal activity [36]. Downregulation of this nutrient-sensing network has been associated with increased lifespan and healthspan [37]. Notably, 17α-estradiol treatment diminished nutrient-sensing network activity in most hypothalamic neurons, which may be a contributing

factor in promoting lifespan extension.

In this report, we demonstrated significant changes in neuron populations involved in appetite

control, including *Agrp*, *Pomc*, *Oxt*, *Oxtr*, *Gipr*, and *Glp2r* neurons. Among the identified subtypes,

the proportion of Oxt neurons saw the most considerable increase due to 17α-estradiol treatment

(**Figure 5A**). Oxt plays versatile roles in social behavior, stress response, satiety, energy balance,

reproduction, and inflammation [38]. Most *Oxt* neurons originate from the paraventricular nucleus

(PVN) and supraoptic nuclei (SON) in the hypothalamus, exhibiting high plasticity during

development and intricate circuitry [39, 40]. The PVN, arcuate nucleus (ARC), and ventromedial

hypothalamic nucleus together form a neural hub in the hypothalamus that integrates peripheral,

nutritional, and metabolic signals to regulate food intake and energy balance [41]. Many effects of

Oxt are sex-specific [42]; for instance, females are less sensitive to exogenous Oxt than males

regarding social recognition [43]. Interestingly, Oxt injections, facilitated by nanoparticles that

enhance blood-brain barrier penetration, reduced body mass while increasing social investigation

and the number of Oxt-positive cells in the SON, particularly in male rats [44]. Additionally,

intracerebroventricular injections of Oxt in rats showed a reduction in food intake in both sexes,

with a more pronounced effect in males [45].Therefore, we propose that Oxt's role in systemic

aging and feeding behavior may contribute to the sex-biased effects of 17α-estradiol, warranting

further verification.

Furthermore, 17α-estradiol treatment appears to have enhanced stress in HPA axis. One evidence

was the increased levels of ferroptosis-signature and UPR in *Crh* neurons. The other evidence was

the elevated serum cortisol, which is also a potential hallmark of aging HPA axis [46, 47].

Therefore, more attentions should be paid to the potential side effects of 17α-estradiol especially

in its clinical application.

In summary, our findings suggest that 17α-estradiol treatment positively influences the HPG axis

and neurons associated with appetite and energy balance. This may be closely linked to the

life-extension effects of 17α-estradiol in aged males. Additionally, employing supervised

clustering based on neuropeptides, hormones, and their receptors proves to be a valuable strategy

for examining pharmacological, pathological, and physiological processes in different neuronal

subtypes within the hypothalamus.

Abbreviations

Ar: androgen receptor; ARC: arcuate nucleus; Astro: astrocyte; AUC: area under the curve; Cga: glycoprotein hormones, alpha Polypeptide; Crh: corticotropin releasing hormone; DEG: differentially-expressed gene; Endo: endothelial cell; Esr1: estrogen receptor 1; Esr2: estrogen receptor 2; Fibro: fibroblast; Glp2r: glucagon-like peptide 2 receptor; GnRH: gonadotropin releasing hormone; GSEA: gene set enrichment analysis; HPA: hypothalamic–pituitary–adrenal; HPG: hypothalamic-pituitary-gonadal; IEU: MRC Integrative Epidemiology Unit; IVW: inverse-variance weighting; KEGG: Kyoto Encyclopedia of Genes and Genomes; Micro: microglia; MR: Mendelian randomization; MTORC1: mechanistic target of rapamycin kinase 1; Nppa: natriuretic peptide A; Oligo: oligodendrocyte; OPC: oligodendrocyte precursor cell; OR: odds ratio; OXPHOS: oxidative phosphorylation; PID: Pathway Interaction Database; POMC: proopiomelanocortin; pQTL: protein quantitative trait loci; PTC: pars tuberalis cell; PVN: paraventricular nucleus; ROC: receiver operating characteristics; snRNA-seq: single-nucleus transcriptomic sequencing; Tany: tanycyte; TF: transcription factor; UPR: unfolded protein response.

Acknowledgments

We appreciate Dr. Qinghua Wang and Hongyun Shi from animal facility of Nantong University in helping with the animal experiments. Special thanks to Professor Ken-ichiro Fukuchi from University of Illinois College of Medicine for constructive comments and suggestions in manuscript preparation.

Funding: This work was supported by the National Natural Science Foundation of China grant 31271448 (YL), 82171621 (LL), 82172566 (ZY), 82150107 (RL) and the National High Level Hospital Clinical Research Funding (2022-PUMCH-A-231) (LL).

Author contributions: Conceptualization, YL, LL; Methodology, YL, GW, XX; Animal operation: YL, LL; EIA assays: YL, LY; Inflammation test: YL, ZY, LL, YS. Validation, LL, JY, ZY; Formal Analysis, YL, JY; Resources, GW, YS, ZY, ZM, LX; Visualization, YL; Writing – Original Draft, YL, JY; Writing – Review & Editing, LL, YS; Supervision, YS, ZY, YL; Project Administration, YL, ZY, LL; Funding Acquisition, ZY, LL. All authors have read and approved the final manuscript.

Declaration of competing interest

The authors have no conflict of interests to declare.

Availability of data and materials: All data are available at GEO accession number GSE248413.

Competing interests

All authors declare no competing interests.

References

1.   Hajdarovic, K. H.; Yu, D.; Webb, A. E., Understanding the aging hypothalamus, one cell at a
time. *Trends Neurosci* **2022,** 45, (12), 942-954.
2.   Dacks, P. A.; Moreno, C. L.; Kim, E. S.; Marcellino, B. K.; Mobbs, C. V., Role of the
hypothalamus in mediating protective effects of dietary restriction during aging. *Front*
Neuroendocrinol **2013,** 34, (2), 95-106.
3.   Masliukov, P. M., Changes of Signaling Pathways in Hypothalamic Neurons with Aging. *Curr*
Issues Mol Biol **2023,** 45, (10), 8289-8308.
4.   Yang, S. B.; Tien, A. C.; Boddupalli, G.; Xu, A. W.; Jan, Y. N.; Jan, L. Y., Rapamycin
ameliorates age-dependent obesity associated with increased mTOR signaling in hypothalamic
POMC neurons. *Neuron* **2012,** 75, (3), 425-36.
5.   Yang, H.; Fang, B.; Wang, Z.; Chen, Y.; Dong, Y., The Timing Sequence and Mechanism of
Aging in Endocrine Organs. *Cells* **2023,** 12, (7).
6.   Stout, M. B.; Steyn, F. J.; Jurczak, M. J.; Camporez, J. G.; Zhu, Y.; Hawse, J. R.; Jurk, D.;
Palmer, A. K.; Xu, M.; Pirtskhalava, T.; Evans, G. L.; de Souza Santos, R.; Frank, A. P.; White,
T. A.; Monroe, D. G.; Singh, R. J.; Casaclang-Verzosa, G.; Miller, J. D.; Clegg, D. J.;
LeBrasseur, N. K.; von Zglinicki, T.; Shulman, G. I.; Tchkonia, T.; Kirkland, J. L.,
17α-Estradiol Alleviates Age-related Metabolic and Inflammatory Dysfunction in Male Mice
Without Inducing Feminization. *J Gerontol A Biol Sci Med Sci* **2017,** 72, (1), 3-15.
7.   Shen, Z.; Hinson, A.; Miller, R. A.; Garcia, G. G., Cap-independent translation: A shared
mechanism for lifespan extension by rapamycin, acarbose, and 17α-estradiol. *Aging Cell* **2021,**
20, (5), e13345.
8.   Wink, L.; Miller, R. A.; Garcia, G. G., Rapamycin, Acarbose and 17α-estradiol share common
mechanisms regulating the MAPK pathways involved in intracellular signaling and
inflammation. *Immun Ageing* **2022,** 19, (1), 8.
9.   Mann, S. N.; Hadad, N.; Nelson Holte, M.; Rothman, A. R.; Sathiaseelan, R.; Ali Mondal, S.;
Agbaga, M. P.; Unnikrishnan, A.; Subramaniam, M.; Hawse, J.; Huffman, D. M.; Freeman, W.
M.; Stout, M. B., Health benefits attributed to 17α-estradiol, a lifespan-extending compound,
are mediated through estrogen receptor α. *Elife* **2020,** 9.
10.   Watanabe, K.; Wilmanski, T.; Baloni, P.; Robinson, M.; Garcia, G. G.; Hoopmann, M. R.;
Midha, M. K.; Baxter, D. H.; Maes, M.; Morrone, S. R.; Crebs, K. M.; Kapil, C.; Kusebauch,
U.; Wiedrick, J.; Lapidus, J.; Pflieger, L.; Lausted, C.; Roach, J. C.; Glusman, G.; Cummings,
S. R.; Schork, N. J.; Price, N. D.; Hood, L.; Miller, R. A.; Moritz, R. L.; Rappaport, N.,
Lifespan-extending interventions induce consistent patterns of fatty acid oxidation in mouse
livers. *Commun Biol* **2023,** 6, (1), 768.
11.   Burns, A. R.; Wiedrick, J.; Feryn, A.; Maes, M.; Midha, M. K.; Baxter, D. H.; Morrone, S. R.;

Prokop, T. J.; Kapil, C.; Hoopmann, M. R.; Kusebauch, U.; Deutsch, E. W.; Rappaport, N.;
Watanabe, K.; Moritz, R. L.; Miller, R. A.; Lapidus, J. A.; Orwoll, E. S., Proteomic changes
induced by longevity-promoting interventions in mice. *Geroscience* **2024,** 46, (2), 1543-1560.

12.   Steyn, F. J.; Ngo, S. T.; Chen, V. P.; Bailey-Downs, L. C.; Xie, T. Y.; Ghadami, M.; Brimijoin,
S.; Freeman, W. M.; Rubinstein, M.; Low, M. J.; Stout, M. B., 17α-estradiol acts through
hypothalamic pro-opiomelanocortin expressing neurons to reduce feeding behavior. *Aging*
Cell **2018,** 17, (1).

13.   Harrison, D. E.; Strong, R.; Allison, D. B.; Ames, B. N.; Astle, C. M.; Atamna, H.; Fernandez,
E.; Flurkey, K.; Javors, M. A.; Nadon, N. L.; Nelson, J. F.; Pletcher, S.; Simpkins, J. W.; Smith,
D.; Wilkinson, J. E.; Miller, R. A., Acarbose, 17-α-estradiol, and nordihydroguaiaretic acid
extend mouse lifespan preferentially in males. *Aging Cell* **2014,** 13, (2), 273-82.

14.   Isola, J. V. V.; Veiga, G. B.; de Brito, C. R. C.; Alvarado-Rincón, J. A.; Garcia, D. N.; Zanini,
B. M.; Hense, J. D.; Vieira, A. D.; Garratt, M.; Gasperin, B. G.; Schneider, A.; Stout, M. B.,
17α-estradiol does not adversely affect sperm parameters or fertility in male mice:
implications for reproduction-longevity trade-offs. *Geroscience* **2023,** 45, (4), 2109-2120.

15.   Stout, M. B.; Vaughan, K. L.; Isola, J. V. V.; Mann, S. N.; Wellman, B.; Hoffman, J. M.; Porter,
H. L.; Freeman, W. M.; Mattison, J. A., Assessing tolerability and physiological responses to
17α-estradiol administration in male rhesus macaques. *Geroscience* **2023,** 45, (4), 2337-2349.

16.   Harrison, D. E.; Strong, R.; Reifsnyder, P.; Kumar, N.; Fernandez, E.; Flurkey, K.; Javors, M.
A.; Lopez-Cruzan, M.; Macchiarini, F.; Nelson, J. F.; Markewych, A.; Bitto, A.; Sindler, A. L.;
Cortopassi, G.; Kavanagh, K.; Leng, L.; Bucala, R.; Rosenthal, N.; Salmon, A.; Stearns, T. M.;
Bogue, M.; Miller, R. A., 17-a-estradiol late in life extends lifespan in aging UM-HET3 male
mice; nicotinamide riboside and three other drugs do not affect lifespan in either sex. *Aging*
Cell **2021,** 20, (5), e13328.

17.   Strong, R.; Miller, R. A.; Antebi, A.; Astle, C. M.; Bogue, M.; Denzel, M. S.; Fernandez, E.;
Flurkey, K.; Hamilton, K. L.; Lamming, D. W.; Javors, M. A.; de Magalhães, J. P.; Martinez, P.
A.; McCord, J. M.; Miller, B. F.; Müller, M.; Nelson, J. F.; Ndukum, J.; Rainger, G. E.;
Richardson, A.; Sabatini, D. M.; Salmon, A. B.; Simpkins, J. W.; Steegenga, W. T.; Nadon, N.
L.; Harrison, D. E., Longer lifespan in male mice treated with a weakly estrogenic agonist, an
antioxidant, an α-glucosidase inhibitor or a Nrf2-inducer. *Aging Cell* **2016,** 15, (5), 872-84.

18.   Rath, S.; Sharma, R.; Gupta, R.; Ast, T.; Chan, C.; Durham, T. J.; Goodman, R. P.; Grabarek,
Z.; Haas, M. E.; Hung, W. H. W.; Joshi, P. R.; Jourdain, A. A.; Kim, S. H.; Kotrys, A. V.; Lam,
S. S.; McCoy, J. G.; Meisel, J. D.; Miranda, M.; Panda, A.; Patgiri, A.; Rogers, R.; Sadre, S.;
Shah, H.; Skinner, O. S.; To, T. L.; Walker, M. A.; Wang, H.; Ward, P. S.; Wengrod, J.; Yuan, C.
C.; Calvo, S. E.; Mootha, V. K., MitoCarta3.0: an updated mitochondrial proteome now with
sub-organelle localization and pathway annotations. *Nucleic Acids Res* **2021,** 49, (D1),
D1541-d1547.

19.   Hu, H.; Miao, Y. R.; Jia, L. H.; Yu, Q. Y.; Zhang, Q.; Guo, A. Y., AnimalTFDB 3.0: a
comprehensive resource for annotation and prediction of animal transcription factors. *Nucleic*
Acids Res **2019,** 47, (D1), D33-d38.

20.   Lu, H.; Ping, J.; Zhou, G.; Zhao, Z.; Gao, W.; Jiang, Y.; Quan, C.; Lu, Y.; Zhou, G.,
CommPath: An R package for inference and analysis of pathway-mediated cell-cell
communication chain from single-cell transcriptomics. *Comput Struct Biotechnol J* **2022,** 20,
5978-5983.

21.   Badia, I. M. P.; Vélez Santiago, J.; Braunger, J.; Geiss, C.; Dimitrov, D.; Müller-Dott, S.; Taus,
P.; Dugourd, A.; Holland, C. H.; Ramirez Flores, R. O.; Saez-Rodriguez, J., decoupleR:
ensemble of computational methods to infer biological activities from omics data. *Bioinform*
Adv **2022,** 2, (1), vbac016.

22.   Liao, Y.; Wang, J.; Jaehnig, E. J.; Shi, Z.; Zhang, B., WebGestalt 2019: gene set analysis
toolkit with revamped UIs and APIs. *Nucleic Acids Res* **2019,** 47, (W1), W199-w205.

23.   Hemani, G.; Zheng, J.; Elsworth, B.; Wade, K. H.; Haberland, V.; Baird, D.; Laurin, C.;
Burgess, S.; Bowden, J.; Langdon, R.; Tan, V. Y.; Yarmolinsky, J.; Shihab, H. A.; Timpson, N.
J.; Evans, D. M.; Relton, C.; Martin, R. M.; Davey Smith, G.; Gaunt, T. R.; Haycock, P. C.,
The MR-Base platform supports systematic causal inference across the human phenome. *Elife*
2018, 7.

24.   Sun, B. B.; Maranville, J. C.; Peters, J. E.; Stacey, D.; Staley, J. R.; Blackshaw, J.; Burgess, S.;
Jiang, T.; Paige, E.; Surendran, P.; Oliver-Williams, C.; Kamat, M. A.; Prins, B. P.; Wilcox, S.
K.; Zimmerman, E. S.; Chi, A.; Bansal, N.; Spain, S. L.; Wood, A. M.; Morrell, N. W.; Bradley,
J. R.; Janjic, N.; Roberts, D. J.; Ouwehand, W. H.; Todd, J. A.; Soranzo, N.; Suhre, K.; Paul, D.
S.; Fox, C. S.; Plenge, R. M.; Danesh, J.; Runz, H.; Butterworth, A. S., Genomic atlas of the
human plasma proteome. *Nature* **2018,** 558, (7708), 73-79.

25.   Hemani, G.; Tilling, K.; Davey Smith, G., Orienting the causal relationship between
imprecisely measured traits using GWAS summary data. *PLoS Genet* **2017,** 13, (11),
e1007081.

26.   Inada, K.; Tsujimoto, K.; Yoshida, M.; Nishimori, K.; Miyamichi, K., Oxytocin signaling in
the posterior hypothalamus prevents hyperphagic obesity in mice. *Elife* **2022,** 11.

27.   Dalvi, P. S.; Belsham, D. D., Glucagon-like peptide-2 directly regulates hypothalamic neurons
expressing neuropeptides linked to appetite control in vivo and in vitro. *Endocrinology* **2012,**
153, (5), 2385-97.

28.   Leder, B. Z.; Rohrer, J. L.; Rubin, S. D.; Gallo, J.; Longcope, C., Effects of aromatase
inhibition in elderly men with low or borderline-low serum testosterone levels. *J Clin*
Endocrinol Metab **2004,** 89, (3), 1174-80.

29.   Guay, A. T.; Jacobson, J.; Perez, J. B.; Hodge, M. B.; Velasquez, E., Clomiphene increases
free testosterone levels in men with both secondary hypogonadism and erectile dysfunction:
who does and does not benefit? *Int J Impot Res* **2003,** 15, (3), 156-65.

30.   Wang, C.; Swerdloff, R. S., Testosterone Replacement Therapy in Hypogonadal Men.
Endocrinol Metab Clin North Am **2022,** 51, (1), 77-98.

31.   Camacho, E. M.; Huhtaniemi, I. T.; O'Neill, T. W.; Finn, J. D.; Pye, S. R.; Lee, D. M.; Tajar,
A.; Bartfai, G.; Boonen, S.; Casanueva, F. F.; Forti, G.; Giwercman, A.; Han, T. S.; Kula, K.;
Keevil, B.; Lean, M. E.; Pendleton, N.; Punab, M.; Vanderschueren, D.; Wu, F. C.,
Age-associated changes in hypothalamic-pituitary-testicular function in middle-aged and older
men are modified by weight change and lifestyle factors: longitudinal results from the
European Male Ageing Study. *Eur J Endocrinol* **2013,** 168, (3), 445-55.

32.   Rodrigues Dos Santos, M.; Bhasin, S., Benefits and Risks of Testosterone Treatment in Men
with Age-Related Decline in Testosterone. *Annu Rev Med* **2021,** 72, 75-91.

33.   Zhang, G.; Li, J.; Purkayastha, S.; Tang, Y.; Zhang, H.; Yin, Y.; Li, B.; Liu, G.; Cai, D.,
Hypothalamic programming of systemic ageing involving IKK-β, NF-κB and GnRH. *Nature*
2013, 497, (7448), 211-6.

34.   Shoupe, D., Individualizing hormone therapy to minimize risk: accurate assessment of risks
and benefits. *Womens Health (Lond)* **2011,** 7, (4), 475-85.
35.   Isola, J. V. V.; Zanini, B. M.; Hense, J. D.; Alvarado-Rincón, J. A.; Garcia, D. N.; Pereira, G.
C.; Vieira, A. D.; Oliveira, T. L.; Collares, T.; Gasperin, B. G.; Stout, M. B.; Schneider, A.,
Mild calorie restriction, but not 17α-estradiol, extends ovarian reserve and fertility in female
mice. *Exp Gerontol* **2022,** 159, 111669.
36.   López-Otín, C.; Blasco, M. A.; Partridge, L.; Serrano, M.; Kroemer, G., Hallmarks of aging:
An expanding universe. *Cell* **2023,** 186, (2), 243-278.
37.   Singh, P. P.; Demmitt, B. A.; Nath, R. D.; Brunet, A., The Genetics of Aging: A Vertebrate
Perspective. *Cell* **2019,** 177, (1), 200-220.
38.   Kerem, L.; Lawson, E. A., The Effects of Oxytocin on Appetite Regulation, Food Intake and
Metabolism in Humans. *Int J Mol Sci* **2021,** 22, (14).
39.   Rosen, G. J.; de Vries, G. J.; Goldman, S. L.; Goldman, B. D.; Forger, N. G., Distribution of
oxytocin in the brain of a eusocial rodent. *Neuroscience* **2008,** 155, (3), 809-17.
40.   Madrigal, M. P.; Jurado, S., Specification of oxytocinergic and vasopressinergic circuits in the
developing mouse brain. *Commun Biol* **2021,** 4, (1), 586.
41.   Cornejo, M. P.; Hentges, S. T.; Maliqueo, M.; Coirini, H.; Becu-Villalobos, D.; Elias, C. F.,
Neuroendocrine Regulation of Metabolism. *J Neuroendocrinol* **2016,** 28, (7).
42.   Carvalho Silva, R.; Pisanu, C.; Maffioletti, E.; Menesello, V.; Bortolomasi, M.; Gennarelli, M.;
Baune, B. T.; Squassina, A.; Minelli, A., Biological markers of sex-based differences in major
depressive disorder and in antidepressant response. *Eur Neuropsychopharmacol* **2023,** 76,
89-107.
43.   Dumais, K. M.; Veenema, A. H., Vasopressin and oxytocin receptor systems in the brain: Sex
differences and sex-specific regulation of social behavior. *Front Neuroendocrinol* **2016,** 40,
1-23.
44.   Duarte-Guterman, P.; Lieblich, S. E.; Qiu, W.; Splinter, J. E. J.; Go, K. A.; Casanueva-Reimon,
L.; Galea, L. A. M., Oxytocin has sex-specific effects on social behaviour and hypothalamic
oxytocin immunoreactive cells but not hippocampal neurogenesis in adult rats. *Horm Behav*
2020, 122, 104734.
45.   Liu, C. M.; Davis, E. A.; Suarez, A. N.; Wood, R. I.; Noble, E. E.; Kanoski, S. E., Sex
Differences and Estrous Influences on Oxytocin Control of Food Intake. *Neuroscience* **2020,**
447, 63-73.
46.   Veldhuis, J. D., Changes in pituitary function with ageing and implications for patient care.
Nat Rev Endocrinol **2013,** 9, (4), 205-15.
47.   Warde, K. M.; Smith, L. J.; Basham, K. J., Age-related Changes in the Adrenal Cortex:
Insights and Implications. *J Endocr Soc* **2023,** 7, (9), bvad097.
705

706 **Figure legends**

707 **Figure 1. snRNA-seq profiling of the hypothalamus from O, O.T and Y samples. (A)** UMAP

708 visualization of nuclei colored by 10 cell types: neuron (Neu), astrocyte (Astro), oligodendrocyte

709 (Oligo), oligodendrocyte precursor cell (OPC), tanycyte (Tany), ependymocyte (Epen), microglia

710 (Micro), fibroblast (Fibro), pars tuberalis cell (PTC), and endothelial cell (Endo), from

hypothalamus of aged rats (O), 17α-estradiol-treated aged rats (O.T) and young rats (Y). **(B)** Heatmap showing the classic markers of 10 major cell types in hypothalamus. **(C)** Cell-type compositions by groups (left panel) or by major cell types with the total cell numbers shown above each column. **(D)** Circos plot depicting the number of ligand–receptor pairs between Neu and other cell types (color strips) for each group. **(E)** Dot plot showing significant ligand–receptor interactions between Neurons for each group. Boxes showing the unique ligand–receptor interactions between Neuron.O (black boxes) or between Neuron.O.T (blue boxes). **(F)** Dot plot of the top 6 enriched GO biological process terms across three groups of neurons via GSEA analysis. **(G)** The top 15 changed pathways/gene sets according to the ranks of AUC values in selected pathways related to neuronal synapses and axons from Gene Ontology (GO) biological process, GO molecular function and GO cellular component.

Figure 2. Two opposing regulatory signaling networks in neuron metabolism. (A) Dot plot of the selected pathways representing the prominent changes of overall expression levels across Neuron.O, Neuron.O.T and Neuron.Y in metabolism, signaling and synaptic activity. **(B)** Correlation heatmap showing transcription factors (TFs) that correlated with the two opposing regulatory signaling networks in the mixed neurons of O, O.T and Y. **(C)** The shared unique markers of each quarter (c1-c4) in 6 pathways in hypothalamic neurons (O, O.T, and Y). The markers were then collected as c1.up.signature (19 genes) and c4.up.signature (12 genes). **(D)** The aging-related cell proportions of each quarter shown by 4 pathways. **(E)** The correlation of c1.up.signature and c2.up.signature with the two opposing regulatory signaling networks.

Figure 3. Screening of neuron subtypes by supervised clustering that responded distinctly to aging and 17α-estradiol treatment. (A) Diagram outlining the features of supervised clustering of neurons in hypothalamus compared with the traditional unsupervised clustering. **(B)** The top 10 and bottom 10 neurons of 121 neurons according to the mean expression values of eight signaling pathways from stress, apoptosis, metabolism, immune response, and senescence in Neuron.O. The expression levels of these neuron subtypes from Neuron.O.T and Neuron.Y were also indicated. **(C)** Word clouds displaying the frequency of neurons that were among the top 10 in each of the 16 signaling pathways as shown in **Table S2**. The 16 pathways were from oxidative stress, apoptosis,

metabolism, immune response, and senescence. **(D)** Venn diagram showing the number of neuron

subtypes exclusively in each group or shared by the groups in (C). The neurons with frequency >3

were selected for calculation in (C). **(E)** The top 15 and bottom 15 neurons among 121 neurons

according to the mean expression levels of c1.up.signature and c4.up.signature in Neuron.O. **(F)**

The top 15 and bottom 15 neurons among 121 neurons according to the mean expression levels of

c1.up.signature and c4.up.signature in Neuron.O.T.

Figure 4. The responses of Crh neurons to long-term 17α-estradiol treatment. (A) The two

intersected neurons among the top 10 neurons according to the mean expression levels of the three

senescence-related pathways. **(B)** The expression profiles of selected pathways from the two

opposing signaling networks in Neuron *Crh*, *Nppa* and *Nppc*. Neuron *Crh* and *Nppa* were the only

two types of neurons shared by the three top 10 neurons in (A). **(C)** The downregulated and

upregulated DEGs expressed by mitochondria or in pathway adherens junction between O.T and

O in Crh neurons. **(D)** The top 25 TF activities in *Crh* and *Gnrh1* neurons. **(E)** Enzyme

immunoassay of the serum levels of Crh, cortisol, and aldosterone in Y, O and O.T. Two-tail

unpaired T-test was performed. p values were labeled.

Figure 5. The response of Oxt neurons to 17α-estradiol and the causal effects of Oxt on other

endocrine factors.

(A), (B) The top 10 (arrows) and bottom 10 (arrows) types of neurons from 121 subtypes ranked

in cell proportions among three groups. **(C)** Dot plots showing the expression profiles of the

selected pathways from the two opposing signaling pathways in four types of food uptake-related

neurons, which decreased or increased among the top 10 ranks in (A) or (B). Blue arrows:

c1.up.signature and c4.up.signature. **(D)** Volcanic plots showing the DEGs between Neuron.O.T

and Neuron.O in the pathway synaptic membrane. **(E)** Enzyme immunoassay of the plasma levels

of Oxt in three groups. **(F)** Top 25 TF activities in neuron Oxt. **(G)** Significant causal effects (p <

0.05, IVW) between exposure OXT (id: prot-a-2159) and 203 endocrine-related outcomes, which

were not significant in reverse MR analysis. Significant heterogeneity (Q_pval < 0.05). Significant

horizontal pleiotropy (pval < 0.05).

Figure 6. The response of HPG axis in the males to 17α-estradiol and the causal effects of Gnrh on other endocrine factors. (A) The expression profiles of pathways from the two opposing signaling networks in Gnrh1-, Esr2-, Esr1- or Ar-positive neurons. **(B)** Enzyme immunoassay of the serum levels of Gnrh, total testosterone (T), and estrogen (E) in Y, O and O.T samples. Two-tail unpaired T-test was performed. **(C)** Inflammation of seminiferous tubules in testes of O and O.T. Left two panels: representative HE staining of testis inflammation in O and the normal seminiferous tubules of O.T. Right panel: the mean testis inflammation index of O and O.T. **(D)** The top 25 TF activities in Gnrh1 neurons in three groups. **(E)** The activities of 14 pathways in Gnrh1-, Esr2-, Esr1- or Ar-positive neurons. **(F)** Significant causal effects (IVW, p < 0.05) between exposure GNRH1 (id: prot-a-1233) and 203 endocrine-related outcomes, which were not significant in reverse MR analysis. **(G)** Items with significant causal effects (IVW, p < 0.05) in both directions of MR analysis between GNRH1 (id: prot-a-1233) and 203 endocrine-related outcomes.