The effects of 17α-estradiol treatment on endocrine system revealed by single-nucleus

transcriptomic sequencing of hypothalamus

Lei Li[#,*1], Guanghao Wu[#2], Xiaolei Xu[3], Junling Yang[4], Lirong Yi[5], Ziqing Yang[6], Zheng Mo[3], Li

Xing[3], Ying Shan[*1], Zhuo Yu[*3], Yinchuan Li[*5]

[1]Department of Obstetrics and Gynecology, National Clinical Research Center for Obstetric &

Gynecologic Diseases, Peking Union Medical College Hospital, Chinese Academy of Medical

Sciences and Peking Union Medical College, 100730 Beijing, China.

[2]School of Medical Technology , Beijing Institute of Technology, 100081, Beijing, China.

[3]Department of Medical Oncology , Beijing Tsinghua Changgung Hospital, School of Clinical

Medicine, Tsinghua University, 102218, Beijing, China.

[4]Department of Cancer Biology and Pharmacology, University of Illinois College of Medicine,

Peoria, Illinois, United states of America.

[5]Institute of Reproductive Medicine, Medical School of Nantong University, Nantong, Jiangsu

226001, China.

[6]School of Basic Medical Sciences, Shandong University, Jinan, Shandong 250012, China.

# These authors contributed equally: Lei Li, Guanghao Wu

* Corresponding author. Email: shypumch@163.com; lilei64@pumch.cn; yza02214@btch.edu.cn;

18622397604@163.com

Abstract

In this study, we investigated 17α-estradiol's role in lifespan extension from long-term

administration. Pooled hypothalami from aged male Norway brown rats treated with 17α-estradiol

(O.T), aged male controls (O), and young male controls (Y) underwent single-nucleus

transcriptomic sequencing (snRNA-seq). To evaluate the effects of 17α-estradiol on aging neurons,

supervised clustering of neurons based on neuropeptides and their receptors were used to evaluate

the responses of each neuron subtype during aging and after 17α-estradiol treatment. The elevated

cellular metabolism, stress and decreased expression levels of pathways involved in synapse

formation in neurons initiated by aging were significantly attenuated by 17α-estradiol. Assessment

of changes in neuron populations showed that neurons related to food intake, reproduction, blood

pressure, stress response, and electrolyte balance were sensitive to 17α-estradiol treatment.

17α-estradiol treatment not only increased serum Oxytocin (Oxt), but also heightened the activity

of hypothalamic-pituitary-gonadal (HPG) axis, as evidenced by significantly elevated levels of

plasma Gnrh, total testosterone, and decreased estradiol. Elevated Gnrh1 was confirmed to be one

of the causal effects mediating the role of 17α-estradiol in energy homeostasis, neural synapse,

and stress response. Notably, *Crh* neurons exhibited prominent stressed phenotype in O.T, distinct

to appetite-stimulating neurons *Agrp* and *Ghrl*. Therefore, the HPG axis and energy metabolism

may be key targets of 17α-estradiol in male hypothalamus. Additionally, supervised clustering of

neurons was shown to be a useful method for assessing treatment responses and cellular

perturbation among different neuron subtypes in the hypothalamus.

Key words snRNA-seq, hypothalamus, 17α-estradiol, aging, Crh, Oxt, Gnrh

Background

The hypothalamus serves as the central hub for controlling energy homeostasis, stress response,

temperature, learning, feeding, sleep, social behavior, sexual behavior, hormone secretion,

reproduction, osmoregulation, blood pressure, visceral activities, emotion, and circadian rhythms

[1]. The hypothalamic energy-sensing system, particularly the circuits that regulate food intake,

plays a crucial role in life span extension [2]. Elevated metabolic activity in the aged

hypothalamus has been reported in aged hypothalamus, including increased mTor signaling [3, 4].

Additionally, decreases in gonadotropin-releasing hormone (GnRH), Ghrh, Trh, monoamine

neurotransmitters, and blood supply are hallmarks of aging hypothalamus [5].

Previous studies have demonstrated that 17α-estradiol extends the lifespan of male mice and has

beneficial effects on metabolism and inflammation, similar to those of rapamycin and acarbose

[6-8]. Recent study indicated that 17α-estradiol also extends the lifespan of male rats [9].Further

investigations revealed certain unique features of 17α-estradiol in life extension distinct to

rapamycin and acarbose [10, 11]. Moreover, it has been shown that 17α-estradiol targets

hypothalamic *POMC* neurons to reduce metabolism by decreasing feeding behavior through

anorexigenic pathways [12]. Interestingly, the lifespan extension effect has only been observed in

male animals [13]. The safety of 17α-estradiol is key for translation into clinical treatment, and the

potential side effects on reproduction and feminization by 17α-estradiol treatment must be considered. However, contradictory results have been reported regarding its side effects on reproduction and feminization [6, 14, 15]. Therefore, further investigation and verification are needed to understand the underlying mechanisms of lifespan extension and the safety of 17α-estradiol.

In this report, we utilized single-nucleus transcriptomic sequencing and performed supervised clustering of neurons based on neuropeptides, hormones, and their receptors. Supervised clustering offers better resolution in cell cluster screening compared to traditional unsupervised clustering. We assessed the effects of 17α-estradiol on metabolism, stress responses, ferroptosis, senescence, inflammation, and pathways involved in synaptic activity in each neuron subtype, ranking the most sensitive neurons. The effects of 17α-estradiol on reversing aging-related cellular processes were evaluated by two opposing regulatory networks involved in hypermetabolism, stress, inflammation, and synaptic activity. Several key endocrine factors from serum were examined, and the potential side effects of 17α-estradiol on specific neurons were also evaluated.

Materials and methods

Animals, treatment and tissues

Twelve Norway brown male rats (12 months old) were acquired from Charles River, including 8 12-months old and 4 1-month old (Beijing). Aged rats were randomly allocated into control and 17α-estradiol-treated groups. Four Aged rats treated with 17α-estradiol (Catalog #: E834897, Macklin Biochemical, Shanghai, China) were fed freely with regular diet mixed with 17α-estradiol at a dose of 14.4 mg/kg (14.4 ppm), starting at 24 months of age for 6 months according to prior reports [16, 17]. The young rats were fed a regular diet without 17α-estradiol continuously for 3 months until 4 months old. All rats had ad libitum access to food and water throughout the experiments. The rats were then euthanized via CO_2, hypothalami, testes and blood serum were collected for subsequent experimental procedures. All blood samples were collected at 9:00-9:30 a.m to minimize hormone fluctuation between animals. All animal procedures were reviewed and approved by the Institutional Animal Care and Use Committee at Nantong University.

Enzyme immunoassays

Enzyme immunoassays kits for rat Oxt (Catalog #: EIAR-OXT), Corticotropin Releasing Factor

(Catalog #: EIAR-CRF), and gonadoliberin-1 (Catalog #: EIAR-GNRH) were obtained from

Raybiotech (GA, USA). Enzyme immunoassay kits for rat serum total testosterone (Catalog #:

ml002868), estradiol (Catalog #: ml002891), aldosterone (Catalog #: ml002876), and cortisol

(Catalog #: ml002874) were obtained from Enzyme-linked Biotechnology (Shanghai, China). Sera

from 3 animals per group were used and each was diluted 10 or 20 times for immunoassays.

Seminiferous tubule inflammation test

8 testes were obtained from each sample group and then subjected to fixation in 4% formalin for

at least 1 week. Formalin-fixed paraffin-embedded rat testis sections of 5 µm thickness were used

for HE Staining. At least 30 seminiferous tubules in each slide were examined for inflammation

test. Testis with at least 1 inflammatory seminiferous tubule was set as 1, and normal testis was set

as 0 for inflammation index calculation.

snRNA-seq data processing, batch effect correction, and cell subset annotation

Intact hypothalami were cryopreserved in liquid nitrogen from sacrificed rats. Two (O) or three (Y

and O.T) hypothalami were pooled within each group and homogenized in 500 µL ice-cold

homogenization buffer (0.25 M sucrose, 5 mM $CaCl_2$, 3 mM $MgAc_2$, 10 mM Tris-HCl [pH 8.0], 1

mM DTT, 0.1 mM EDTA, 1× protease inhibitor, and 1 U/µL RiboLock RNase inhibitor) with

Dounce homogenizer. Then, the homogenizer was washed with 700 µL ice-cold nuclei washing

buffer (0.04% bovine serum albumin, 0.2 U/µL RiboLock RNase Inhibitor, 500 mM mannitol, 0.1

mM phenylmethanesulfonyl fluoride protease inhibitor in 1× phosphate buffer saline). Next, the

homogenates were filtered through a 70-µm cell strainer to collect the nuclear fraction. The

nuclear fraction was mixed with an equal volume of 50% iodixanol and added on top of a 30%

and 33% iodixanol gradient. This solution was then centrifuged for 20 min at 10 000 ×g at 4 °C.

After the myelin layer was removed from the top of the gradient, the nuclei were collected from

the 30% and 33% iodixanol interface. The nuclei were resuspended in nuclear wash buffer and

resuspension buffer and pelleted for 5 min at 500 ×g at 4 °C. The nuclei were filtered through a

40-µm cell strainer to remove cell debris and large clumps, and the nuclear concentration was

manually assessed using trypan blue counterstaining and a hemocytometer. Finally, the nuclei

were adjusted to 700–1200 nuclei/µL, and examined with a 10X Chromium platform.

Reverse transcription, cDNA amplification and library preparation were performed according to

the protocol from 10X Genomics and Chromium Next GEM Single Cell 3′ Reagent Kits v3.1.

Library sequencing was performed on the Illumina HiSeq™ 4000 by Gene Denovo Biotechnology

Co., Ltd (Guangzhou, China).

10X Genomics Cell Ranger software (version 3.1.0) was used to convert raw BCL files to FASTQ

files, and for alignment and counts quantification. Reads with low-quality barcodes and UMIs

were filtered out and then mapped to the reference genome. Reads uniquely mapped to the

transcriptome and intersecting an exon at least 50% were considered for UMI counting. Before

quantification, the UMI sequences were corrected for sequencing errors, and valid barcodes were

identified using the EmptyDrops method. The cell $\times$ gene matrices were produced via UMI

counting and cell barcodes calling. Cells with an unusually high number of UMIs ($\geq$8000) or

mitochondrial gene percent ($\geq$15%) were filtered out. Batch effect correction was performed by

harmony.

Pathways, gene signatures, TFs and TF cofactors, cell communication

Gene sets and pathways were derived from Hallmark gene sets of MSigDB collections, the KEGG

pathway database, Reactome pathway database, and WikiPathways database, and some ontology

terms derived from the Gene Ontology (GO) resource. Mitochondrial pathways were derived from

MitoCarta3.0 [18]. Pathways, gene sets, and gene signatures were evaluated with the

PercentageFeatureSet function built into R package Seurat. TFs and TF cofactors were obtained

from AnimalTFDB 3.0 [19]. TFs and TF cofactors were further filtered with mean counts >0.1.

The ligand–receptor pairs were calculated via R package CommPath [20].

Correlation analysis and ROC analysis

Pearson correlation coefficient was calculated with the linkET package ($p < 0.05$). Fast Wilcoxon

rank sum test and auROC analysis was performed with the wilcoxauc function in R package

presto. The minimal cell number in either one of the comparing pairs should be no less than 15.

Ranks of area under the curve (AUC) values were in descending order. A total of 431 pathways

from Hallmark, KEGG and PID databases were used for correlation analysis with MitoCarta

OXPHOS subunits in neurons and non-neural cells (**Figure 2—figure supplement 1**). A total of

97 pathways related to synapse activity were derived from GO, including GO cellular components,

GO biological processes and GO molecular functions (**Table S1**).

The division of expression level-dependent clusters in each pathway and their gene

signatures

The quarters of the mixed cell populations from O, O.T, and Y hypothalamic neurons were equally

divided using the R function fivenum from the R package stats, based on pathway expression

levels. Thus, the total number of neurons was evenly divided into four clusters (c1-c4) in terms of

cell number. The cell proportions from O.T, O, and Y neurons in each cluster were weighted

against the total number of neurons in the three groups. The unique markers of each cluster were

calculated using the FindAllMarkers function from the Seurat package. The intersection of the

unique markers from the six pathways was obtained for heatmap plotting. Nineteen genes that

were highly expressed in c1 were identified as c1.up.signature via the PercentageFeatureSet

function in the Seurat package. Twelve genes that were highly expressed in c4 were identified as

c4.up.signature. There were no intersecting unique markers in clusters c2 and c3 among the six

selected pathways.

TF and pathway activities

The TF resources were derived from CollecTRI, the pathway resource was from PROGENy, and

the enrichment scores of TFs and pathways were performed with the Univariate Linear Model

(ulm) method according to the pipeline in R package decoupleR [21].

Subtypes of neurons generated by supervised clustering and cell prioritization

Vast majority of these subtypes were clustered by neuropeptides, hormones, and their receptors

within all the neurons with the subset function from R package Seurat (the target gene expression

level > 0). A total of 209 neuron subtypes were obtained, comprising 104 neuropeptide-secreting

or hormone-secreting neurons and 105 neurons expressing a unique neuropeptide receptor or

hormone receptor (Table S2). Further groupings may exist within the identified neuron subtypes,

and the category of excitatory or inhibitory neurons was not discriminated further. The cell

proportion of each neuron subtype was weighted according to the total number of neurons in O.T,

O, and Y samples. The mean values ± standard deviation of pathways and gene signatures were

performed for each subtype. The top 20 and the bottom 20 items were calculated. The cell type

prioritization was performed using the R package Augur, with the subsample_size parameter of

the calculate_auc function set to 6 [22]. In each comparison pair, the minimum number of cells in

a subcluster shall not be less than 6 when performing cell prioritization with function

calculate_auc.

Differential expression and pathway enrichment analysis

DEGs between groups were identified via FindMarkers (test.use = bimod, min.pct = 0.1, logfc.Threshold = 0.25, avg_diff > 0.1 or < −0.1). DEGs were then enriched in redundant GO terms via WebGestalt and filtered with false discovery rate < 0.05 [23].

Bidirectional Mendelian randomization (MR) study

The protein quantitative trait locus (pQTL) GWAS summary data of 204 human endocrine-related GWAS summary data with European ancestry were obtained from open-access MRC Integrative Epidemiology Unit (IEU) (**Table S4**) [24, 25]. Independent genome-wide significant SNPs for exposure OXT (id:prot-a-2159) or GNRH1 (id: prot-a-1233) were used as instrumental variables with genome-wide significance (P < 1 × 10^{-5}), independence inheritance (r2 < 0.001) without linkage disequilibrium (LD) with each other for MR. For the reverse MR, independent genome-wide significant SNPs from 203 endocrine-related GWAS summary data (P < 1 × 10^{-5}, r2 < 0.001) without LD with each other were obtained as exposures and human GWAS summary data of OXT (id:prot-a-2159) or GNRH1 (id:prot-a-1233) were used as outcomes. Weak instruments less than 10 were discarded via F-statistics.

MR and reverse MR analysis were conducted with method inverse-variance weighting (IVW), MR Egger, Weighted median, Simple mode, and Weighted mode. The screening criteria: all of the odds ratio (OR) values of the 5 methods should be simultaneously either >1 or <1 and the significant p value of IVW was <0.05. The heterogeneity via IVW method and the horizontal pleiotropy were also evaluated with R package TwoSampleMR [26].

Results

The overall changes in aged hypothalamus with or without long-term 17α-estradiol treatment via snRNA-seq profiling

To investigate the hypothalamus as a potential key target of 17α-estradiol's effects on life extension, we performed snRNA-seq on the entire hypothalamus of aged and 17α-estradiol-treated aged Norway brown rats, using the hypothalamus from young adult male rats as a control. We identified 10 major cell types based on specific cell markers of the hypothalamus (**Figure 1A-B**). Notably, the proportions of all non-neural cells changed in O versus Y (**Figure 1C**). For instance, Oligo, OPC, and Micro were found to be increased, while Astro, Tany, Fibro, PTC, and Endo were

decreased in O compared to Y. The proportions of Oligo, OPC, and Micro were also increased in

17α-estradiol-treated aged group (O.T) compared to those in Y. Furthermore, Endo was increased

in O.T compared to both Y and O. The proportions of Astro, Tany, Epen, and PTC decreased more

in O.T than those in O when compared to Y. These results indicated that 17α-estradiol treatment

had extensive effects on the proportions of non-neural cells in hypothalamus.

Cell communication analysis revealed significant changes in the ligand-receptor pairs between

neurons and other cell types, particularly in Endo, Fibro, Tany, and Astro (**Figure 1D**). Significant

ligand–receptor interactions among neurons also changed in O.T and O groups, especially in O

(**Figure 1E**). Notably, among the significant ligand–receptor pairs in neurons,

Bmp2–Acvr1/Acvr2a/Acvr2b/Bmpr, *Gdf11–Acvr2a/Acvr2b*, *Inhba–Acvr1/Acvr2a/Acvr2b*,

Nrg1/Nrg2/Nrg4–Erbb4, *Rspo1–Lgr5/Lrp6*, and *Rspo3–Lgr5* were exclusively and significantly

increased in neurons of the O group compared to those in O.T and Y, suggesting enhanced TGF

superfamily-mediated signaling activity and canonical Wnt signaling during aging. The

significantly changed ligand–receptor pairs *Nlgn1–Nrxn1/Nrxn2*, *Nlgn2–Nrxn1/Nrxn2/Nrxn3*,

Nlgn3–Nrxn1/Nrxn2/Nrxn3, *Nxph1–Nrxn1/Nrxn2/Nrxn3*, *Nxph3–Nrxn1/Nrxn2/Nrxn3*,

Pomc–Oprd1/Oprk1/Oprm1, and *Vip–Adcyap1r1/Avpr1a/Vipr2* were exclusively increased in

neurons of O.T compared to O and Y (**Figure 1E**). These ligand–receptor pairs were associated

with synaptic activity, cellular adhesion, the opioid system, and vasodilation, indicating unique

roles of 17α-estradiol in restoring certain physiological functions in the aging hypothalamus. The

increased *Pomc* signal in O.T neurons aligns with previous reports suggesting that 17α-estradiol

treatment decreases food uptake in mice, potentially correlated with *Pomc* neurons (**Figure 1E**)

[12].

Gene set enrichment analysis (GSEA) based on DEGs also corroborated the expression profiles

related to stress responses and synapse-associated cellular processes in neurons (**Figure 1F**). ROC

analysis of significantly differently expressed pathways related to neural synapses, manually

selected from Gene Ontology databases, indicated that most top-ranked pathways related to

synapses, according to AUC values, were downregulated in aged neurons, while 17α-estradiol

treatment reversed this trend (**Figure 1G, Table S1**).

Overall, these findings suggest that 17α-estradiol broadly reshapes cell populations, cellular

communication, neuropeptide secretion, and synapse-related cellular processes in the aging

hypothalamus, distinguishing it from both the young hypothalamus and the untreated aged hypothalamus.

The two opposing signaling networks in regulating metabolism and synapse activity, which can be balanced effectively by 17α-estradiol

To monitor the metabolism and neural status affected by 17α-estradiol, we utilized the energy metabolism pathway MitoCarta OXPHOS subunits to calculate the positively or negatively correlated pathways in hypothalamic neurons (**Figure 2—figure supplement 1**). Our findings revealed that energy metabolism and synapse activity represent two opposing regulatory signaling networks in hypothalamic neurons, with 17α-estradiol strongly playing a significant role in balancing these networks (**Figure 2A**). At the core of these opposing signaling pathways are two categories of contrasting TFs (**Figure 2B**). For example, *Calr*, *Clu*, *Peg3*, *Prnp*, *Ndufa13*, *Actb*, *Ywhab*, *Nfe2l1*, *Mtdh*, *Npm1*, *Bex2*, *Aft4*, and *Maged1* were positively correlated with pathways involved in OXPHOS subunits, lysosome function, protein export, mTorc1 signaling, and the unfolded protein response (UPR) in O, O.T and Y neurons, while showing negative correlations with pathways related to ubiquitin-mediated proteolysis, endocytosis, tight junctions, focal adhesion, axon guidance, and MAPK signaling. Additionally, TFs *Myt1l*, *Ctnnd2*, *Tenm4*, *Camta1*, *Med12l*, *Rere*, *Csrnp3*, *Erbb4*, *Jazf1*, *Dscam*, *Klf12*, and *Kdm4c* exhibited opposite correlation patterns with these selected pathways in O, O.T and Y neurons. These TFs may take conserved roles in regulating the two opposing biological processes in hypothalamic neurons.

We then attempted to establish gene signatures to represent these two opposing signaling networks, thereby displaying the cell status of aging and evaluating the effects exerted by 17α-estradiol. To achieve this, we evenly divided the expression levels of each of the six selected pathways from the two opposing signaling networks into four quarters (c1-c4) among the mixed neurons from O, O.T, and Y, calculating the shared unique markers in each quarter (**Figure 2C, D**). From the distribution patterns, we observed that the proportion of neurons in O decreased from c1 to c4 in metabolic pathways (MitoCarta OXPHOS subunits and Hallmark mTorc1 signaling), while this trend was reversed in the opposing signaling pathways (GOBP synapse organization and KEGG MAPK signaling pathway) (**Figure 2C**). In contrast, in Y, this trend was opposite, suggesting the expression levels from the 4 quarters (c1-c4) of the two opposing signaling networks can be used

to monitor aging status. Treatment with 17α-estradiol alleviated this trend or even reversed it in O.

We then screened the shared unique markers of each quarter from the six selected pathways in an

attempt to establish the gene signatures representing the two opposing signaling networks. Unique

markers in c1 (19 genes, c1.up.signature) and c4 (12 genes, c4.up.signature) were identified;

however, c2 and c3 lacked unique markers shared by the 6 pathways (**Figure 2D**). Consequently,

the 19 genes in c1.up.signature displayed an inversed correlation pattern with the 12 genes in

c4.up.signature, indicating the two opposing gene signatures are capable of reflecting the two

opposing signaling networks in hypothalamic neurons (**Figure 2E**). Conversely, the balance of the

two opposing signaling networks affected by 17α-estradiol in non-neural cell types was less

pronounced than in neurons, showing variable effects on non-neural cells (**Figure 2—figure**

supplement 2). GOBP pathway enrichment analysis revealed that Micro exhibited lower levels of

synapse-related cellular processes in O.T compared to O, which was distinct from the observations

in neurons (**Figure 2—figure supplement 3**). Therefore, in this report, we primarily focused on

hypothalamic neurons and their responses to aging and 17α-estradiol.

Supervised clustering revealed distinct responses of different subtypes of hypothalamic

neurons to aging and 17α-estradiol

The hypothalamus contains numerous neuron subtypes that release various neuropeptides and

hormones to regulate fundamental body functions. To differentiate the changes occurring during

aging and the effect of 17α-estradiol on each neuron subtype, we performed supervised clustering

based on neuropeptides, hormones, or their receptors (**Table S2**) (**Figure 3A**). The cell counts in

each neuronal subcluster classified by neuropeptide secretion (neuropeptide-secreting subtypes)

and subclusters defined by neuropeptide or hormone receptor expression (receptor-expressing

subtypes) were quantified and compared in sample Y (**Figure 3B**). Notably, neurons expressing

Prlr, *Esr1*, and *Ar* ranked among the top 20 receptor-expressing subtypes across all analyzed

neuron populations. The similarity indices among these cell subtypes were further calculated

(**Figure 3—figure supplement 1**), revealing high positive correlations in neuron subtypes

expressing *Cartpt*, *Nxph4*, *Bdnf*, *Cck*, *Crh*, *Nppa*, *Adcyap1*, and *Penk*, as well as those expressing

Esr1, *Calcrl*, and *Pth2r*. These similarities may partially reflect cellular overlap between subtypes

(**Figure 3—figure supplement 1**).

We next calculated the prioritization of cellular perturbation induced by aging and/or 17α-estradiol treatment across these screened neuron subtypes (**Figure 3C, D**). The *Gnrh1* neuron subtype ranked among the top perturbed neuropeptide-secreting subtypes in both O vs Y and O.T vs Y comparisons (purple arrows). Notably, *Sct* and *Kiss1* neuron subtypes emerged as the top 2 perturbed populations in the O.T vs O analysis (red arrows), highlighting their heightened sensitivity to 17α-estradiol in the aged hypothalamus. Among receptor-expressing subtypes, *Insr* neurons showed the highest sensitivity to perturbation in both O vs Y and O.T vs Y comparisons (purple arrows, **Figure 3D**), while *Adipor2* and *Mlnr* neurons (blue arrows) ranked as the top 2 sensitive subtypes in the O.T vs O analysis. Intriguingly, neurons expressing *Ar* and *Esr1* ranked among the top 20 most perturbed receptor subtypes during aging (O vs Y), but were no longer ranked in this group following treatment (O.T vs Y and O.T vs O comparisons). This indicates that 17α-estradiol administration attenuated age-associated perturbation in these neuronal subtypes (**Figure 3D**).

Differential senescence or stress levels and subtype-specific susceptibility in aged hypothalamic neurons

To gain a deeper understanding of the effects of 17α-estradiol treatment on the aged hypothalamus, we selected 3 gene signatures and 2 gene sets associated with aging, apoptosis, and stress to characterize the differential responses of distinct neuronal subtypes to aging and 17α-estradiol. These neuronal subtypes were then ranked separately based on neuropeptide-secreting subtypes and receptor-expressing subtypes (**Figure 4A, B**).

Neuropeptide-secreting subtypes, such as *Prlh-*, *Sct-*, *Gast-*, *Nppa-*, *Nxph1-*, *Ucn-*, *Pnoc-*, *Galp-*, and *Ghrl*-expressing neurons, were consistently ranked among the top 20 in at least 4 out of the 5 gene signatures or gene sets. These neurons are involved in gastrointestinal function, food intake, hunger, energy homeostasis, water homeostasis, vascular regulation, and pain, suggesting that aging exacerbates senescence or stress in these physiological processes.

In contrast, neurons expressing *Igf2*, *Crh*, *Npy*, *Npw*, *Npff*, *Nmu*, *Agrp*, or *Adipoq* ranked among the bottom 20 in at least four of the five signatures or gene sets. These neuropeptides and hormones are associated with cortical excitability, stress response, food intake, circadian rhythms, fat metabolism, insulin sensitivity, heart rate, and blood pressure. Notably, although

Crh-expressing neurons exhibited high overall cellular perturbation among neuropeptide-secreting

subtypes (**Figure 3C**), the relatively lower senescence and stress burden in *Crh*-, *Npy*-, *Npw*-, and

Nmu-expressing neurons—key mediators of the stress response—compared to other neuronal

subtypes represents a defining characteristic of the aged hypothalamus.

Regarding receptor-expressing subtypes, *Mc3r*-, *Sstr1*-, *Kiss1r*-, *Ntsr2*-, *Mlnr*-, *Ntsr1*-, *Npy1r*-,

and *Avpr1a*-expressing neurons were consistently among the top 20 in at least 4 of the 5 gene

signatures or gene sets. These receptor-expressing subtypes are involved in food intake,

neurotransmission, reproduction, gut function, fat metabolism, circadian rhythm, and

vasoconstriction, thereby indicating heightened stress in the aging hypothalamus. Conversely,

Glp2r-, *Lepr*-, *Paqr8*-, and *Npr3*-expressing neurons were among the bottom 20 in at least 4 of the

5 gene signatures or gene sets, with associations to glucose regulation, fat metabolism,

progesterone signaling, blood volume, and blood pressure.

Notably, most of the 5 signatures or gene sets in top ranked neurons exhibited alleviated

senescence or stress following 17α-estradiol treatment, indicating that such treatment mitigates

senescence or stress in these specific neuronal populations (**Figure 4A, B**).

The appetite-controlling neurons and hypothalamic–pituitary–adrenal (HPA) axis were

altered by long-term 17α-estradiol treatment in the males

To further investigate the positive effects, potential side effects or compensatory effects of

17α-estradiol treatment, we performed stricter screening by intersecting the top 20 and bottom 20

ranks of the scores of c1-up-signature, ferroptosis gene signature, UPR, Mtorc1 signaling, and

OXPHOS subunits (**Figure 5A**). Neurons expressing *Calcb*, *Edn3*, *Ucn*, *Ghrl*, *Nmu*, *Npff*, *Cnp*,

and *Agrp* ranked among the bottom 20 in at least 4 out of the 5 gene signatures or gene sets. These

neurons are involved in stress responses, vascular activity, appetite regulation, and muscle

contraction. Notably, the lower levels of *Agrp*- and *Ghrl*-expressing neurons in the

Mitocarta_OXPHOS_subunits signature may also indicate reduced physiological activity of these

potent appetite-promoting neurons during 17α-estradiol treatment, which could represent a key

clue to its role in lifespan prolongation.

In contrast, neurons expressing *Gast*, *Npb*, *Nppa*, *Crh*, *Scg3*, and *Npw* consistently ranked among

the top 20 in at least 4 of the 5 gene signatures or gene sets. These neurons participate in

gastrointestinal activity, feeding behavior, stress responses, cardio-renal homeostasis, and

angiogenesis. Of note, the expression pattern of *Crh* neurons in O.T was opposite to that in O

(**Figure 4A**). Additionally, the Mitocarta_OXPHOS_subunits score in *Crh* neurons was the

highest among all examined neuropeptide-expressing subtypes (**Figure 5A**), which contrasted

sharply with those of *Agrp* and *Ghrl* neurons. Additionally, the treatment with 17α-estradiol in O.T

also elevated several key metabolic pathways in *Crh* neurons compared to those in Y and O

(**Figure 5B**). 17α-estradiol treatment increased the c1-up-signature while simultaneously reducing

many pathways associated with synapse activity and the c4-up-signature in *Crh* neurons of O.T,

indicating a potent stressed phenotype in *Crh* neuron. In contrast, in *Kiss1* and *Prlh* neurons, the

decreased c1-up-signature in O.T implied a lesser extent of stressed phenotype in these neurons

compared to *Crh* neurons. The status of Crh neurons in O.T may be associated with elevated TF

activities of *Esr2*, *Usf2*, *Hdac5*, *Creb3l1*, *Tfam*, *Preb*, *Pou3f2*, and *Hoxb5* (**Figure 5C**). The

aberrant changes in *Crh* neurons were also evidenced by the increased expression of DEGs related

to mitochondria-expressed genes and reduced expression of DEGs in the adherens junction

pathway in O.T, indicative of higher energetic activity and altered extracellular adhesion in this

type of neuron by 17α-estradiol treatment (**Figure 5D**).

Notably, the HPA axis was altered by 17α-estradiol treatment, as evidenced by the elevated

cortisol levels in O.T compared to O (p = 0.078) (**Figure 5E**). The correlation between elevated

cortisol production and the heightened stress in *Crh* neurons by 17α-estradiol treatment needs

further investigation. Additionally, as a crucial component of the renin-angiotensin-aldosterone

system, the significantly increased serum aldosterone in O.T and its potential role in sodium

reabsorption and cardiovascular health also warrant more detailed investigation (**Figure 5E**).

In summary, 17α-estradiol treatment altered the activity of appetite-promoting neurons and the

hypothalamic-pituitary-adrenal (HPA) axis in male BN rats, while also inducing enhanced stress

responses in *Crh* neurons.

17α-estradiol increased Oxt neuron proportion and secretion and its possible role in

mediating the effect of 17α-estradiol on endocrine system

Aging and 17α-estradiol treatment also altered the proportions of various neuron subtypes among

O, O.T, and Y (**Figure 6A, B**). The proportions of *Grp*, *Pmch*, *Npb*, *Serpinb9*, *Sstr2*, *Agrp*, *Sstr3*,

Mlnr, and *Hcrt* neurons ranked in the top 10 in O, while *Oxt*, *Vip*, *Avp*, *Calca*, *Glp2r*, *Tacr1*, *Trh*,

Serping1, *Npff*, and *Npy1r* were in the top 10 in O.T. *Galp*, *Calcrl*, *Ednra*, *Oxt*, *Serpinh1*, *Pomc*,

Cck, *Crh*, *Tacr1*, and *Kiss1* neurons were among the bottom 10 in O, whereas *Oxtr*, *Galp*, *Agrp*,

Serpinb9, *Npvf*, *Serpinh1*, *Ednrb*, *Agt*, *Gipr*, and *Pomc* neurons ranked among the bottom 10 in

O.T. *Agrp*, *Pomc*, *Oxt*, *Oxtr*, *Gipr*, and *Glp2r* neurons are well-known for their roles in regulating

food intake and energy homeostasis. *Agrp* neurons are activated by hunger, while *Pomc* neurons

are activated by satiety in the ARC of the hypothalamus. 17α-estradiol treatment effectively

elevated the expression levels of the c4.up.signature (blue arrows) and synapse-associated

processes in neuron subtypes *Agrp*, *Pomc*, *Oxt* and *Glp2r* in O.T compared to O. This may

mitigate the adverse effects of reduced cell populations in Pomc and *Agrp* neurons in aging

hypothalamus (**Figure 6C**). This finding indicates a potential role of 17α-estradiol in appetite

control, as previously reported [12]. Additionally, 17α-estradiol treatment resulted in elevated

proportions of *Vip*, *Avp*, *Npff*, *Calca*, and *Tacr1* neurons, which ranked in the top 10, while the

proportions of Agt neurons decreased. These are all associated with blood pressure regulation

(**Figure 6—figure supplement 1**). Notably, the proportions of *Oxt* and *Glp2r* neurons, both of

which have anorexigenic effects [27, 28], increased in O.T. In addition to the increased number of

Oxt-positive neurons, the expression level of *Oxt* also rose in O.T. The elevated expression of

synapse-related pathways was supported by the increased DEGs in the enriched synaptic

membrane pathway in *Oxt* neurons (**Figure 6D**).

More importantly, the serum level of Oxt was significantly elevated in O.T compared to O (p =

0.04), yet remained lower than those in Y (**Figure 6E**). Notably, the top TF activities in O.T and O

differed markedly from those in Y (**Figure 6F**). The elevated levels of *Hopx* and *Xbp1* may be

associated with the response to 17α-estradiol treatment.

Due to the intricate regulatory networks among various endocrine factors, elucidating the causal

effect of Oxt on other endocrine factors is quite complex using traditional methods. MR analysis,

employing variant SNPs as genetic tools, is advantageous for such task. We performed a

bidirectional MR analysis of the GWAS summary data of human plasma OXT and 203

endocrine-related and hypothalamus-related factors, most of which are protein quantitative trait

loci (pQTL) data from the IEU (**Table S3**). As an exposure, OXT revealed a significant causal

effect on POMC/beta-endorphin (id:prot-a-2327, id:prot-a-2325), glucagon (id:prot-a-1181),

GNRH1/Progonadoliberin-1   (id:prot-a-1233),   and   total   testosterone   levels

(id:ebi−a−GCST90012112, id:ieu−b−4864) (**Figure 6G**). NPW and CBLN1 were found to be

negatively associated with OXT, but the significance of these associations was not found in the

reverse MR analysis (**Figure 6—figure supplement 2A, B**).

In contrast, we could not identify significant associations between OXT and estradiol levels

(id:ebi-a-GCST90012105, id:ebi-a-GCST90020092, id:ebi-a-GCST90020091, id:ieu-b-4872,

id:ieu-b-4873, id:ukb-e-30800_AFR, id:ukb-e-30800_CSA). Interestingly, QRFP, IGF1, AGRP,

TAC4, GRP, CLU, BNF, PCSK7, PACAP, ANP, TAC3, CRH, INSL6, and PRL displayed

significant associations with OXT in both MR and reverse MR analysis, indicative of their

complex causal effects (**Figure 6—figure supplement 2A, B**).

The results suggest that elevated Oxt levels induced by 17α-estradiol may have positive

associations with endocrine factors governing feeding behavior, glucose metabolism, male

reproduction, and sex hormones. Therefore, OXT may serve as a potential mediator of

17α-estradiol.

17α-estradiol activated HPG axis and the elevated Gnrh also took important roles in

mediating the effect of 17α-estradiol on other endocrine factors

Given the sensitivity of GnRH- and sex hormone receptor-expressing neuron subtypes to

17α-estradiol treatment (**Figure 3C, D**), we analyzed their expression profiles alongside

representative pathway genes from two opposing signaling networks - those related to metabolism

and synapses - such as the c1-up-signature and c4-up-signature (**Figure 7A**). However, neither the

c1-up-signature nor the c4-up-signature was up-regulated in *Gnrh1* neuron in the O.T in

comparison with Y. *Ar* and *Esr2* neuron displayed decreased level of c1-up-signature in

comparison with O. Only in *Esr1* neuron was the c1-up-signature found to be up-regulated.

Meanwhile, both *Ar* and *Esr* neurons displayed increased level of c4-up-signature in O.T

comparing with O. *Ar*, *Pgr*, and *Esr1* were also among the top 20 of c4-up-signature, suggesting

long-term 17α-estradiol treatment did not impose significant stress on hypothalamic neurons

expressing these hormone receptors (**Figure 7—figure supplement 1**). But *Gnrh1* and *Crh*

neurons were among the bottom 20, indicative of higher cellular stress by long-term 17α-estradiol

treatment. However, based on these cellular perturbations, it's difficult to define the precise

physiological status of these subtypes of neurons, particularly regarding neuroendocrine activities.

Consequently, we performed enzyme immunoassays of hormones from the serum of O, O.T and Y.

The treatment with 17α-estradiol significantly increased the plasma level of Gnrh compared to Y

($p = 0.0099$) and approached significance when compared to O ($p = 0.096$) (**Figure 7B**). More

intriguingly, testosterone levels in serum were significantly increased in O.T compared to O ($p =$

0.018) and approached significance when compared to Y ($p = 0.052$). Additionally, the serum

estradiol levels were significantly increased in O compared to Y ($p = 0.011$) and significantly

decreased in O.T compared to O ($p = 0.019$), suggesting that 17α-estradiol treatment markedly

altered the homeostasis of testosterone and estradiol.

Furthermore, most testes from 30-month-old male BN rats exhibited severe age-related

inflammation and epithelial collapse of seminiferous tubules (**Figure 7C**). The testes without

inflammation in O.T displayed normal morphology. 17α-estradiol treatment slightly decreased the

testis inflammation in O.T compared to that in O ($p = 0.15$), indicating a potential positive role of

17α-estradiol treatment in male reproductive system. The elevated TFs such as *Sf1*, *Pparg*, *Litaf*,

Nupr1, *Rxrg*, *E2f2*, and *Zfp42* may be involved in the transcriptional regulation by 17α-estradiol in

O.T (**Figure 7D**). Importantly, the activities of androgen and estrogen pathways were decreased in

Gnrh1 neurons in O.T compared to O, and were distinct from those in *Ar*, *Esr1*, and *Esr2* neurons

(**Figure 7E**). These signaling pathways are important for the feedback control of sex hormone

secretion in *Gnrh* neurons, and these results may also reflect the strong effect of 17α-estradiol on

Gnrh neurons.

To decipher the potential effects of elevated serum Gnrh levels on endocrine system, we

performed bidirectional MR analysis of the GWAS summary data of human GNRH1 (id:

prot-a-1233) and 203 endocrine-related factors with genetic variants SNPs. We found strong

causal effects of GNRH1 on GAL/Galanin (id:prot−a−1166), POMC/Beta−endorphin

(id:prot−a−2327, id:prot−a−2325), Adrenomedullin (id:prot−a−48), BDNF (id:prot−a−242)**,** and

LPR (id:prot−a−1724), which are involved in feeding, energy homeostasis, osmotic regulation,

and neuron plasticity (**Figure 7F**). Notably, CRH/Corticotropin (id:prot−a−2326),

PRLH/Prolactin−releasing peptide (id:prot−a−2376), NPW/Neuropeptide W (id:prot−a−2082),

Glucagon (id:prot−a−1181), Chromogranin−A (id:prot−a−538) displayed bidirectional

significance, indicating close and complex causal effects between GNRH1 and these endocrine

factors (**Figure 7G, Figure 7—figure supplement 2A, B**). These results also suggest that the role

of 17α-estradiol treatment in feeding, energy homeostasis, reproduction, osmotic regulation, stress

response, and neuronal plasticity may be mediated, at least in part, by elevated Gnrh secretion.

Discussion

The most striking role of 17α-estradiol treatment revealed in this study showed that HPG axis was

substantially improved in the levels of serum Gnrh and testosterone. The underlying molecular

mechanism remains unclear; however, prior reports have indicated that 17α-estradiol can bind to

ESR1 [9]. In our findings, 17α-estradiol treatment significantly decreased serum estradiol levels

while elevating serum testosterone. Based on this evidence, we propose that 17α-estradiol may

function similarly to estrogen receptor antagonists or aromatase inhibitors, potentially preventing

the conversion of androgens to estrogens [29, 30]. These actions could alleviate the feedback

inhibition exerted by estrogen on hypothalamus and pituitary, thereby facilitating the secretion of

Gnrh, FSH, and LH [31].

The testosterone levels in men gradually decline beginning in the third decade of life [32].

Age-related deterioration of the gonadotropic axis, particularly in older males with low serum

testosterone, is often linked to numerous aging symptoms, including loss of skeletal muscle mass,

reduced muscle strength and power, low bone mineral density, frailty, impaired physical

performance, mobility limitations, increased risk of diabetes, elevated all-cause cardiovascular

mortality, cognitive decline, and heightened risk of Alzheimer's disease [33]. Consequently,

testosterone supplementation in older men is beneficial. Additionally, Gnrh supplementation may

help mitigate age-related declines in neurogenesis and slow aging processes [34]. Importantly,

treatment with 17α-estradiol did not result in feminization or adversely affect the sperm

parameters and fertility in male animals [6, 14]. Thus, the observed increases in Gnrh and serum

testosterone levels due to 17α-estradiol treatment are likely advantageous for older males,

particularly those experiencing late-onset hypogonadism.

Postmenopausal women with low estrogen experience aging-related syndromes similar to those of

older males with low serum testosterone. Those women also face increased mortality,

cardiovascular disease, osteoporosis fracture, urogenital atrophy, and dementia, all of which may

benefit from hormone therapy [35]. However, a prior report indicates that 17α-estradiol treatment

does not provide positive life extension effects in aged females [13]. The discrepancy may stem

from the inhibitory effects of estrogens associated with 17α-estradiol treatment, as evidenced by

its inability to enhance female fertility [36]. Nonetheless, due to the lack of parallel data in aged

female BN rats treated with 17α-estradiol, further research is needed to definitely address this

question in the female subjects in the future.

The stressed phenotype observed in neuronal subtypes discussed herein likely represents a

transcriptomic manifestation of heightened physiological activity. For instance, as evidenced in

this study, prolonged 17α-estradiol treatment induces a pronounced stressed phenotype in *Gnrh*

neurons alongside elevated Gnrh secretion and consequent high serum testosterone levels.

Similarly, the stressed phenotype reflected in the *Crh* neuronal transcriptome coincides with

substantially increased serum cortisol. Furthermore, long-term 17α-estradiol treatment markedly

alleviates the stressed phenotype in appetite-stimulating neurons (*Agrp* and *Ghrl*), suggesting an

appetite-suppressing effect in rats. This aligns with previously reported findings that 17α-estradiol

treatment inhibits feeding behavior in mice [12].

Another notable effect of 17α-estradiol is its ability to reduce the overall expression levels of

energy metabolism in hypothalamic neurons of aged male BN rats. The nutrient-sensing network,

mediated by MTORC1 complex, is a central regulator of mRNA and ribosome biogenesis, protein

synthesis, glucose metabolism, autophagy, lipid metabolism, mitochondrial biosynthesis, and

proteasomal activity [37]. Downregulation of this nutrient-sensing network has been associated

with increased lifespan and healthspan [38]. Notably, 17α-estradiol treatment diminished

nutrient-sensing network activity in most hypothalamic neurons, which may be a contributing

factor in promoting lifespan extension.

In this report, we demonstrated significant changes in neuron populations involved in appetite

control, including *Agrp*, *Pomc*, *Oxt*, *Oxtr*, *Gipr*, and *Glp2r* neurons. Among the identified subtypes,

the proportion of Oxt neurons saw the most considerable increase due to 17α-estradiol treatment

(**Figure 6A**). Oxt plays versatile roles in social behavior, stress response, satiety, energy balance,

reproduction, and inflammation [39]. Most *Oxt* neurons originate from the paraventricular nucleus

(PVN) and supraoptic nuclei (SON) in the hypothalamus, exhibiting high plasticity during

development and intricate circuitry [40, 41]. The PVN, arcuate nucleus (ARC), and ventromedial

hypothalamic nucleus together form a neural hub in the hypothalamus that integrates peripheral,

nutritional, and metabolic signals to regulate food intake and energy balance [42]. Many effects of

Oxt are sex-specific [43]; for instance, females are less sensitive to exogenous Oxt than males

regarding social recognition [44]. Interestingly, Oxt injections, facilitated by nanoparticles that

enhance blood-brain barrier penetration, reduced body mass while increasing social investigation

and the number of Oxt-positive cells in the SON, particularly in male rats [45]. Additionally,

intracerebroventricular injections of Oxt in rats showed a reduction in food intake in both sexes,

with a more pronounced effect in males [46].Therefore, we propose that Oxt's role in systemic

aging and feeding behavior may contribute to the sex-biased effects of 17α-estradiol, warranting

further verification.

Furthermore, 17α-estradiol treatment appears to have enhanced stress in HPA axis. One evidence

was the increased levels of ferroptosis-signature and UPR in *Crh* neurons. The other evidence was

the elevated serum cortisol, which is also a potential hallmark of aging HPA axis [47, 48].

Therefore, more attentions should be paid to the potential side effects of 17α-estradiol especially

in its clinical application.

In summary, our findings suggest that 17α-estradiol treatment positively influences the HPG axis

and neurons associated with appetite and energy balance. This may be closely linked to the

life-extension effects of 17α-estradiol in aged males.   Additionally, employing supervised

clustering based on neuropeptides, hormones, and their receptors proves to be a valuable strategy

for examining pharmacological, pathological, and physiological processes in different neuronal

subtypes within the hypothalamus.

Abbreviations

Ar: androgen receptor; Agrp, agouti related neuropeptide; ARC: arcuate nucleus; Astro: astrocyte;

AUC: area under the curve; Cga: glycoprotein hormones, alpha polypeptide; Crh: corticotropin

releasing hormone; DEG: differentially-expressed gene; Endo: endothelial cell; Esr1: estrogen

receptor 1; Esr2: estrogen receptor 2; Fibro: fibroblast; Ghrl, Ghrelin; Glp2r: glucagon-like

peptide 2 receptor; GnRH: gonadotropin releasing hormone; GSEA: gene set enrichment analysis;

HPA: hypothalamic–pituitary–adrenal**;** HPG: hypothalamic-pituitary-gonadal; IEU: MRC

Integrative Epidemiology Unit; IVW: inverse-variance weighting; KEGG: Kyoto Encyclopedia of

Genes and Genomes; Kiss1, kisspeptin, Micro: microglia; MR: Mendelian randomization;

MTORC1: mechanistic target of rapamycin kinase 1; Oligo: oligodendrocyte; OPC:

oligodendrocyte precursor cell; OR: odds ratio; OXPHOS: oxidative phosphorylation; PID:

Pathway Interaction Database; POMC: proopiomelanocortin; pQTL: protein quantitative trait loci;

Prlh, prolactin releasing hormone; PTC: pars tuberalis cell; PVN: paraventricular nucleus; ROC:

receiver operating characteristics; snRNA-seq: single-nucleus transcriptomic sequencing; Tany:

tanycyte; TF: transcription factor; UPR: unfolded protein response.

Acknowledgments

We appreciate Dr. Qinghua Wang and Hongyun Shi from animal facility of Nantong University in

helping with the animal experiments. Special thanks to Professor Ken-ichiro Fukuchi from

University of Illinois College of Medicine for constructive comments and suggestions in

manuscript preparation.

Funding: This work was supported by the National Natural Science Foundation of China grant

31271448 (YL), 82171621 (LL), 82172566 (ZY), 82150107 (RL) and the National High Level

Hospital Clinical Research Funding (2022-PUMCH-A-231) (LL).

Author contributions: Conceptualization, YL, LL; Methodology, YL, GW, XX; Animal

operation: YL, LL; EIA assays: YL, LY; Inflammation test: YL, ZY, LL, YS. Validation, LL, JY,

ZY; Formal Analysis, YL, JY; Resources, GW, YS, ZY, ZM, LX; Visualization, YL; Writing –

Original Draft, YL, JY; Writing – Review & Editing, LL, YS; Supervision, YS, ZY, YL; Project

Administration, YL, ZY, LL; Funding Acquisition, ZY, LL. All authors have read and approved

the final manuscript.

Declaration of competing interest

The authors have no conflict of interests to declare.

Availability of data and materials: All data are available at GEO accession number GSE248413.

Competing interests

All authors declare no competing interests.

References

1.   Hajdarovic, K. H.; Yu, D.; Webb, A. E., Understanding the aging hypothalamus, one

cell at a time. *Trends Neurosci* 2022, 45, (12), 942-954.

2.   Dacks, P. A.; Moreno, C. L.; Kim, E. S.; Marcellino, B. K.; Mobbs, C. V., Role of the

hypothalamus in mediating protective effects of dietary restriction during aging. *Front*

Neuroendocrinol **2013,** 34, (2), 95-106.

3.   Masliukov, P. M., Changes of Signaling Pathways in Hypothalamic Neurons with

Aging. *Curr Issues Mol Biol* **2023,** 45, (10), 8289-8308.

4.   Yang, S. B.; Tien, A. C.; Boddupalli, G.; Xu, A. W.; Jan, Y. N.; Jan, L. Y., Rapamycin

ameliorates age-dependent obesity associated with increased mTOR signaling in

hypothalamic POMC neurons. *Neuron* **2012,** 75, (3), 425-36.

5.   Yang, H.; Fang, B.; Wang, Z.; Chen, Y.; Dong, Y., The Timing Sequence and

Mechanism of Aging in Endocrine Organs. *Cells* **2023,** 12, (7).

6.   Stout, M. B.; Steyn, F. J.; Jurczak, M. J.; Camporez, J. G.; Zhu, Y.; Hawse, J. R.; Jurk,

D.; Palmer, A. K.; Xu, M.; Pirtskhalava, T.; Evans, G. L.; de Souza Santos, R.; Frank,

A. P.; White, T. A.; Monroe, D. G.; Singh, R. J.; Casaclang-Verzosa, G.; Miller, J. D.;

Clegg, D. J.; LeBrasseur, N. K.; von Zglinicki, T.; Shulman, G. I.; Tchkonia, T.;

Kirkland, J. L., 17α-Estradiol Alleviates Age-related Metabolic and Inflammatory

Dysfunction in Male Mice Without Inducing Feminization. *J Gerontol A Biol Sci Med*

Sci **2017,** 72, (1), 3-15.

7.   Shen, Z.; Hinson, A.; Miller, R. A.; Garcia, G. G., Cap-independent translation: A

shared mechanism for lifespan extension by rapamycin, acarbose, and 17α-estradiol.

Aging Cell **2021,** 20, (5), e13345.

8.   Wink, L.; Miller, R. A.; Garcia, G. G., Rapamycin, Acarbose and 17α-estradiol share

common mechanisms regulating the MAPK pathways involved in intracellular

signaling and inflammation. *Immun Ageing* **2022,** 19, (1), 8.

9.   Mann, S. N.; Hadad, N.; Nelson Holte, M.; Rothman, A. R.; Sathiaseelan, R.; Ali

Mondal, S.; Agbaga, M. P.; Unnikrishnan, A.; Subramaniam, M.; Hawse, J.; Huffman,

D. M.; Freeman, W. M.; Stout, M. B., Health benefits attributed to 17α-estradiol, a

lifespan-extending compound, are mediated through estrogen receptor α. *Elife* **2020,**

9.

10.   Watanabe, K.; Wilmanski, T.; Baloni, P.; Robinson, M.; Garcia, G. G.; Hoopmann, M.

R.; Midha, M. K.; Baxter, D. H.; Maes, M.; Morrone, S. R.; Crebs, K. M.; Kapil, C.;

Kusebauch, U.; Wiedrick, J.; Lapidus, J.; Pflieger, L.; Lausted, C.; Roach, J. C.;

Glusman, G.; Cummings, S. R.; Schork, N. J.; Price, N. D.; Hood, L.; Miller, R. A.;

Moritz, R. L.; Rappaport, N., Lifespan-extending interventions induce consistent

patterns of fatty acid oxidation in mouse livers. *Commun Biol* **2023,** 6, (1), 768.

11.   Burns, A. R.; Wiedrick, J.; Feryn, A.; Maes, M.; Midha, M. K.; Baxter, D. H.; Morrone, S.

R.; Prokop, T. J.; Kapil, C.; Hoopmann, M. R.; Kusebauch, U.; Deutsch, E. W.;

Rappaport, N.; Watanabe, K.; Moritz, R. L.; Miller, R. A.; Lapidus, J. A.; Orwoll, E. S.,

Proteomic changes induced by longevity-promoting interventions in mice.

Geroscience **2024,** 46, (2), 1543-1560.

12.   Steyn, F. J.; Ngo, S. T.; Chen, V. P.; Bailey-Downs, L. C.; Xie, T. Y.; Ghadami, M.;

Brimijoin, S.; Freeman, W. M.; Rubinstein, M.; Low, M. J.; Stout, M. B., 17α-estradiol

acts through hypothalamic pro-opiomelanocortin expressing neurons to reduce

feeding behavior. *Aging Cell* **2018,** 17, (1).

13.   Harrison, D. E.; Strong, R.; Allison, D. B.; Ames, B. N.; Astle, C. M.; Atamna, H.;

Fernandez, E.; Flurkey, K.; Javors, M. A.; Nadon, N. L.; Nelson, J. F.; Pletcher, S.;

Simpkins, J. W.; Smith, D.; Wilkinson, J. E.; Miller, R. A., Acarbose, 17-α-estradiol,

and nordihydroguaiaretic acid extend mouse lifespan preferentially in males. *Aging*

Cell **2014,** 13, (2), 273-82.

14.   Isola, J. V. V.; Veiga, G. B.; de Brito, C. R. C.; Alvarado-Rincón, J. A.; Garcia, D. N.;

Zanini, B. M.; Hense, J. D.; Vieira, A. D.; Garratt, M.; Gasperin, B. G.; Schneider, A.;

Stout, M. B., 17α-estradiol does not adversely affect sperm parameters or fertility in

male mice: implications for reproduction-longevity trade-offs. *Geroscience* **2023,** 45,

(4), 2109-2120.

15.   Stout, M. B.; Vaughan, K. L.; Isola, J. V. V.; Mann, S. N.; Wellman, B.; Hoffman, J. M.;

Porter, H. L.; Freeman, W. M.; Mattison, J. A., Assessing tolerability and physiological

responses to 17α-estradiol administration in male rhesus macaques. *Geroscience*

2023, 45, (4), 2337-2349.

16.   Harrison, D. E.; Strong, R.; Reifsnyder, P.; Kumar, N.; Fernandez, E.; Flurkey, K.;

Javors, M. A.; Lopez-Cruzan, M.; Macchiarini, F.; Nelson, J. F.; Markewych, A.; Bitto,

A.; Sindler, A. L.; Cortopassi, G.; Kavanagh, K.; Leng, L.; Bucala, R.; Rosenthal, N.;

Salmon, A.; Stearns, T. M.; Bogue, M.; Miller, R. A., 17-a-estradiol late in life extends

lifespan in aging UM-HET3 male mice; nicotinamide riboside and three other drugs do

not affect lifespan in either sex. *Aging Cell* **2021,** 20, (5), e13328.

17.   Strong, R.; Miller, R. A.; Antebi, A.; Astle, C. M.; Bogue, M.; Denzel, M. S.; Fernandez,

E.; Flurkey, K.; Hamilton, K. L.; Lamming, D. W.; Javors, M. A.; de Magalhães, J. P.;

Martinez, P. A.; McCord, J. M.; Miller, B. F.; Müller, M.; Nelson, J. F.; Ndukum, J.;

Rainger, G. E.; Richardson, A.; Sabatini, D. M.; Salmon, A. B.; Simpkins, J. W.;

Steegenga, W. T.; Nadon, N. L.; Harrison, D. E., Longer lifespan in male mice treated

with a weakly estrogenic agonist, an antioxidant, an α-glucosidase inhibitor or a

Nrf2-inducer. *Aging Cell* **2016,** 15, (5), 872-84.

18. Rath, S.; Sharma, R.; Gupta, R.; Ast, T.; Chan, C.; Durham, T. J.; Goodman, R. P.;

Grabarek, Z.; Haas, M. E.; Hung, W. H. W.; Joshi, P. R.; Jourdain, A. A.; Kim, S. H.;

Kotrys, A. V.; Lam, S. S.; McCoy, J. G.; Meisel, J. D.; Miranda, M.; Panda, A.; Patgiri,

A.; Rogers, R.; Sadre, S.; Shah, H.; Skinner, O. S.; To, T. L.; Walker, M. A.; Wang, H.;

Ward, P. S.; Wengrod, J.; Yuan, C. C.; Calvo, S. E.; Mootha, V. K., MitoCarta3.0: an

updated mitochondrial proteome now with sub-organelle localization and pathway

annotations. *Nucleic Acids Res* **2021,** 49, (D1), D1541-d1547.

19. Hu, H.; Miao, Y. R.; Jia, L. H.; Yu, Q. Y.; Zhang, Q.; Guo, A. Y., AnimalTFDB 3.0: a

comprehensive resource for annotation and prediction of animal transcription factors.

Nucleic Acids Res **2019,** 47, (D1), D33-d38.

20. Lu, H.; Ping, J.; Zhou, G.; Zhao, Z.; Gao, W.; Jiang, Y.; Quan, C.; Lu, Y.; Zhou, G.,

CommPath: An R package for inference and analysis of pathway-mediated cell-cell

communication chain from single-cell transcriptomics. *Comput Struct Biotechnol J*

2022, 20, 5978-5983.

21. Badia, I. M. P.; Vélez Santiago, J.; Braunger, J.; Geiss, C.; Dimitrov, D.; Müller-Dott, S.;

Taus, P.; Dugourd, A.; Holland, C. H.; Ramirez Flores, R. O.; Saez-Rodriguez, J.,

decoupleR: ensemble of computational methods to infer biological activities from

omics data. *Bioinform Adv* **2022,** 2, (1), vbac016.

22. Skinnider, M. A.; Squair, J. W.; Kathe, C.; Anderson, M. A.; Gautier, M.; Matson, K. J.

687  E.; Milano, M.; Hutson, T. H.; Barraud, Q.; Phillips, A. A.; Foster, L. J.; La Manno, G.;

688  Levine, A. J.; Courtine, G., Cell type prioritization in single-cell data. *Nat Biotechnol*

689  **2021,** 39, (1), 30-34.

690 23. Liao, Y.; Wang, J.; Jaehnig, E. J.; Shi, Z.; Zhang, B., WebGestalt 2019: gene set

691  analysis toolkit with revamped Uls and APls. *Nucleic Acids Res* **2019,** 47, (W1),

692  W199-w205.

693 24. Hemani, G.; Zheng, J.; Elsworth, B.; Wade, K. H.; Haberland, V.; Baird, D.; Laurin, C.;

694  Burgess, S.; Bowden, J.; Langdon, R.; Tan, V. Y.; Yarmolinsky, J.; Shihab, H. A.;

695  Timpson, N. J.; Evans, D. M.; Relton, C.; Martin, R. M.; Davey Smith, G.; Gaunt, T. R.;

696  Haycock, P. C., The MR-Base platform supports systematic causal inference across

697  the human phenome. *Elife* **2018,** 7.

698 25. Sun, B. B.; Maranville, J. C.; Peters, J. E.; Stacey, D.; Staley, J. R.; Blackshaw, J.;

699  Burgess, S.; Jiang, T.; Paige, E.; Surendran, P.; Oliver-Williams, C.; Kamat, M. A.;

700  Prins, B. P.; Wilcox, S. K.; Zimmerman, E. S.; Chi, A.; Bansal, N.; Spain, S. L.; Wood,

701  A. M.; Morrell, N. W.; Bradley, J. R.; Janjic, N.; Roberts, D. J.; Ouwehand, W. H.; Todd,

702  J. A.; Soranzo, N.; Suhre, K.; Paul, D. S.; Fox, C. S.; Plenge, R. M.; Danesh, J.; Runz,

703  H.; Butterworth, A. S., Genomic atlas of the human plasma proteome. *Nature* **2018,**

704  558, (7708), 73-79.

705 26. Hemani, G.; Tilling, K.; Davey Smith, G., Orienting the causal relationship between

706  imprecisely measured traits using GWAS summary data. *PLoS Genet* **2017,** 13, (11),

707  e1007081.

708 27. Inada, K.; Tsujimoto, K.; Yoshida, M.; Nishimori, K.; Miyamichi, K., Oxytocin signaling

in the posterior hypothalamus prevents hyperphagic obesity in mice. *Elife* **2022,** 11.

28.   Dalvi, P. S.; Belsham, D. D., Glucagon-like peptide-2 directly regulates hypothalamic

neurons expressing neuropeptides linked to appetite control in vivo and in vitro.

Endocrinology **2012,** 153, (5), 2385-97.

29.   Leder, B. Z.; Rohrer, J. L.; Rubin, S. D.; Gallo, J.; Longcope, C., Effects of aromatase

inhibition in elderly men with low or borderline-low serum testosterone levels. *J Clin*

Endocrinol Metab **2004,** 89, (3), 1174-80.

30.   Guay, A. T.; Jacobson, J.; Perez, J. B.; Hodge, M. B.; Velasquez, E., Clomiphene

increases free testosterone levels in men with both secondary hypogonadism and

erectile dysfunction: who does and does not benefit? *Int J Impot Res* **2003,** 15, (3),

156-65.

31.   Wang, C.; Swerdloff, R. S., Testosterone Replacement Therapy in Hypogonadal Men.

Endocrinol Metab Clin North Am **2022,** 51, (1), 77-98.

32.   Camacho, E. M.; Huhtaniemi, I. T.; O'Neill, T. W.; Finn, J. D.; Pye, S. R.; Lee, D. M.;

Tajar, A.; Bartfai, G.; Boonen, S.; Casanueva, F. F.; Forti, G.; Giwercman, A.; Han, T.

S.; Kula, K.; Keevil, B.; Lean, M. E.; Pendleton, N.; Punab, M.; Vanderschueren, D.;

Wu, F. C., Age-associated changes in hypothalamic-pituitary-testicular function in

middle-aged and older men are modified by weight change and lifestyle factors:

longitudinal results from the European Male Ageing Study. *Eur J Endocrinol* **2013,** 168,

(3), 445-55.

33.   Rodrigues Dos Santos, M.; Bhasin, S., Benefits and Risks of Testosterone Treatment

in Men with Age-Related Decline in Testosterone. *Annu Rev Med* **2021,** 72, 75-91.

34.   Zhang, G.; Li, J.; Purkayastha, S.; Tang, Y.; Zhang, H.; Yin, Y.; Li, B.; Liu, G.; Cai, D.,

732   Hypothalamic programming of systemic ageing involving IKK-β, NF-κB and GnRH.

733   *Nature* **2013,** 497, (7448), 211-6.

734   35.   Shoupe, D., Individualizing hormone therapy to minimize risk: accurate assessment of

735   risks and benefits. *Womens Health (Lond)* **2011,** 7, (4), 475-85.

736   36.   Isola, J. V. V.; Zanini, B. M.; Hense, J. D.; Alvarado-Rincón, J. A.; Garcia, D. N.;

737   Pereira, G. C.; Vieira, A. D.; Oliveira, T. L.; Collares, T.; Gasperin, B. G.; Stout, M. B.;

738   Schneider, A., Mild calorie restriction, but not 17α-estradiol, extends ovarian reserve

739   and fertility in female mice. *Exp Gerontol* **2022,** 159, 111669.

740   37.   López-Otín, C.; Blasco, M. A.; Partridge, L.; Serrano, M.; Kroemer, G., Hallmarks of

741   aging: An expanding universe. *Cell* **2023,** 186, (2), 243-278.

742   38.   Singh, P. P.; Demmitt, B. A.; Nath, R. D.; Brunet, A., The Genetics of Aging: A

743   Vertebrate Perspective. *Cell* **2019,** 177, (1), 200-220.

744   39.   Kerem, L.; Lawson, E. A., The Effects of Oxytocin on Appetite Regulation, Food Intake

745   and Metabolism in Humans. *Int J Mol Sci* **2021,** 22, (14).

746   40.   Rosen, G. J.; de Vries, G. J.; Goldman, S. L.; Goldman, B. D.; Forger, N. G.,

747   Distribution of oxytocin in the brain of a eusocial rodent. *Neuroscience* **2008,** 155, (3),

748   809-17.

749   41.   Madrigal, M. P.; Jurado, S., Specification of oxytocinergic and vasopressinergic

750   circuits in the developing mouse brain. *Commun Biol* **2021,** 4, (1), 586.

751   42.   Cornejo, M. P.; Hentges, S. T.; Maliqueo, M.; Coirini, H.; Becu-Villalobos, D.; Elias, C.

752   F., Neuroendocrine Regulation of Metabolism. *J Neuroendocrinol* **2016,** 28, (7).

43. Carvalho Silva, R.; Pisanu, C.; Maffioletti, E.; Menesello, V.; Bortolomasi, M.;

Gennarelli, M.; Baune, B. T.; Squassina, A.; Minelli, A., Biological markers of

sex-based differences in major depressive disorder and in antidepressant response.

Eur Neuropsychopharmacol **2023,** 76, 89-107.

44. Dumais, K. M.; Veenema, A. H., Vasopressin and oxytocin receptor systems in the

brain: Sex differences and sex-specific regulation of social behavior. *Front*

Neuroendocrinol **2016,** 40, 1-23.

45. Duarte-Guterman, P.; Lieblich, S. E.; Qiu, W.; Splinter, J. E. J.; Go, K. A.;

Casanueva-Reimon, L.; Galea, L. A. M., Oxytocin has sex-specific effects on social

behaviour and hypothalamic oxytocin immunoreactive cells but not hippocampal

neurogenesis in adult rats. *Horm Behav* **2020,** 122, 104734.

46. Liu, C. M.; Davis, E. A.; Suarez, A. N.; Wood, R. I.; Noble, E. E.; Kanoski, S. E., Sex

Differences and Estrous Influences on Oxytocin Control of Food Intake. *Neuroscience*

2020, 447, 63-73.

47. Veldhuis, J. D., Changes in pituitary function with ageing and implications for patient

care. *Nat Rev Endocrinol* **2013,** 9, (4), 205-15.

48. Warde, K. M.; Smith, L. J.; Basham, K. J., Age-related Changes in the Adrenal Cortex:

Insights and Implications. *J Endocr Soc* **2023,** 7, (9), bvad097.

Figure legends

Figure 1. snRNA-seq profiling of the hypothalamus from O, O.T and Y samples. **(A)** UMAP

visualization of nuclei colored by 10 cell types: neuron (Neu), astrocyte (Astro), oligodendrocyte

(Oligo), oligodendrocyte precursor cell (OPC), tanycyte (Tany), ependymocyte (Epen), microglia

(Micro), fibroblast (Fibro), pars tuberalis cell (PTC), and endothelial cell (Endo), from

hypothalamus of aged rats (O), 17α-estradiol-treated aged rats (O.T) and young rats (Y). **(B)** Heatmap showing the classic markers of 10 major cell types in hypothalamus. **(C)** Cell-type compositions by groups (left panel) or by major cell types with the total cell numbers shown above each column. **(D)** Circos plot depicting the number of ligand–receptor pairs between Neu and other cell types (color strips) for each group. **(E)** Dot plot showing significant ligand–receptor interactions between Neurons for each group. Boxes showing the unique ligand–receptor interactions between Neuron.O (black boxes) or between Neuron.O.T (blue boxes). **(F)** Dot plot of the top 6 enriched GO biological process terms across three groups of neurons via GSEA analysis. **(G)** The top 15 changed pathways/gene sets according to the ranks of AUC values in selected pathways related to neuronal synapses and axons from Gene Ontology (GO) biological process, GO molecular function and GO cellular component.

Figure 2. Two opposing regulatory signaling networks in neuron metabolism. (A) Dot plot of the selected pathways representing the prominent changes of overall expression levels across Neuron.O, Neuron.O.T and Neuron.Y in metabolism, signaling and synaptic activity. **(B)** Correlation heatmap showing transcription factors (TFs) that correlated with the two opposing regulatory signaling networks in the mixed neurons of O, O.T and Y. **(C)** The shared unique markers of each quarter (c1-c4) in 6 pathways in hypothalamic neurons (O, O.T, and Y). The markers were then collected as c1.up.signature (19 genes) and c4.up.signature (12 genes). **(D)** The aging-related cell proportions of each quarter shown by 4 pathways. **(E)** The correlation of c1.up.signature and c2.up.signature with the two opposing regulatory signaling networks.

Figure 3. Screening of neuron subtypes via supervised clustering, which responded distinctly to aging and 17α-estradiol treatment. (A) Diagram outlining the features of supervised clustering of neurons in the hypothalamus in comparison with traditional unsupervised clustering. **(B)** The ranks of cell counts in neuropeptide-secreting neuron subclusters (left panel) and subclusters expressing neuropeptide receptors or hormone receptors (right panel) in sample Y. The cell number (n) in each subset is ≥ 10. **(C, D)** The prioritization of the top 20 neuron subclusters across the 3 types of perturbation (O vs Y, O.T vs Y, and O.T vs O) calculated by the Augur algorithm, in neuropeptide-secreting neurons (C) and neuron subclusters expressing neuropeptide

receptors or hormone receptors (D).

Figure 4. Ranking of neuron subtypes with distinct responses to aging and 17α-estradiol

treatment. (A, B) The top 20 and bottom 20 neuron subtypes based on the mean expression

values of five signatures or gene sets, ranked by their values in sample O, in

neuropeptide-secreting subtypes (A) and in neuron subtypes expressing neuropeptide receptors or

hormone receptors (B).

Figure 5. Responses of Crh neurons to long-term 17α-estradiol treatment. (A) The top 20 and

bottom 20 neuropeptide-secreting neuron subtypes, ranked by their mean expression values of five

signatures or gene sets in sample O. **(B)** Expression profiles of selected pathways from two

opposing signaling networks in *Crh*, *Kiss1*, and *Prlh* neurons. **(C)** Downregulated and upregulated

differentially expressed genes (DEGs) associated with mitochondria or the adherens junction

pathway in *Crh* neurons, comparing O.T vs O. **(D)** Top 25 transcription factor (TF) activities in

Crh and *Gnrh1* neurons. **(E)** Serum levels of Crh, cortisol, and aldosterone in Y, O, and O.T

groups as measured by enzyme immunoassay; two-tailed unpaired t-tests were performed, with

p-values indicated.

Figure 6. The response of *Oxt* neurons to 17α-estradiol and the causal effects of Oxt on other

endocrine factors.

(A)   The relative cell proportions of peptide-expressing subclusters (upper panel) and

receptor-expressing subclusters (lower panel) across Y, O, and O.T (sorted in descending order of

proportions in Y). Only subclusters with a cell count of n ≥ 10 in sample Y were included for

calculation. **(B)** Dot plots showing the expression profiles of the selected pathways from the two

opposing signaling pathways in four types of food uptake-related neurons, which decreased or

increased among the top 10 ranks in (A) or (B). Blue arrows: c1.up.signature and c4.up.signature.

(C) Volcanic plots showing the DEGs between Neuron.O.T and Neuron.O in the pathway synaptic

membrane. **(D)** Enzyme immunoassay of the plasma levels of Oxt in three groups. **(E)** Top 25 TF

activities in neuron Oxt. **(F)** Significant causal effects ($p < 0.05$, IVW) between exposure OXT (id:

prot-a-2159) and 203 endocrine-related outcomes, which were not significant in reverse MR

analysis. Significant heterogeneity (Q_pval < 0.05). Significant horizontal pleiotropy (pval <

0.05).

Figure 7. The response of HPG axis in the males to 17α-estradiol and the causal effects of

Gnrh on other endocrine factors. (A) The expression profiles of pathways from the two

opposing signaling networks in *Gnrh1-*, *Esr2-*, *Esr1-* or *Ar*-positive neurons. **(B)** Enzyme

immunoassay of the serum levels of Gnrh, total testosterone (T), and estrogen (E) in Y, O and O.T

samples. Two-tail unpaired T-test was performed. **(C)** Inflammation of seminiferous tubules in

testes of O and O.T. Left two panels: representative HE staining of testis inflammation in O and

the normal seminiferous tubules of O.T. Right panel: the mean testis inflammation index of O and

O.T. **(D)** The top 25 TF activities in *Gnrh1* neurons in three groups. **(E)** The activities of 14

pathways in *Gnrh1-*, *Esr2-*, *Esr1-* or *Ar*-positive neurons. **(F)** Significant causal effects (IVW, p <

0.05) between exposure GNRH1 (id: prot-a-1233) and 203 endocrine-related outcomes, which

were not significant in reverse MR analysis. **(G)** Items with significant causal effects (IVW, p <

0.05) in both directions of MR analysis between GNRH1 (id: prot-a-1233) and 203

endocrine-related outcomes.