Differential effects of diphenyl diselenide (PhSe)₂ on mitochondria-related pathways depending on the cellular energy status in Bovine Vascular Endothelial Cells

Letícia Selinger Galant[1,2], Laura Doblado[2], Rafael Radi[3], João Batista Teixeira da Rocha[4], Andreza Fabro de Bem [1,5,6*], Maria Monsalve[2*]

[1] Biochemistry PhD Program, Federal University of Santa Catarina, Florianopolis, SC, Brazil.

[2] Instituto de Investigaciones Biomédicas Sols-Morreale (CSIC-UAM). Arturo Duperier 4. 28029, Madrid, Spain.

[3] Center for Free Radical and Biomedical Research (CEINBIO), Facultad de Medicina, Universidad de la República, Montevideo, Uruguay

[4] Department of Biochemistry and Molecular Biology, Federal University of Santa Maria, Santa Maria, Brazil.

[5] Laboratory of Bioenergetics and Metabolism, Institute for Biological Sciences; University of Brasília, Brasília, Brazil.

[6] National Institute of Science and Technology on Neuroimmunomodulation, Rio de Janeiro, Brazil.

Abstract

Cellular energy metabolism varies depending on tissue and cell type, as well as the availability of energy substrates and energy demands. We recently investigated the variations in cellular metabolism and antioxidant responses in primary bovine vascular endothelial cells (BAECs) under different energetic substrate conditions in vitro, specifically glucose or galactose. In this context, pharmacological agents may exert different effects on cells depending on their energy metabolism status. In this study, we aimed to characterize the effects of $(PhSe)_2$, a redox-active molecule known for its prominent cardiovascular effects, on redox-bioenergetic cellular pathways under glycolytic or oxidative conditions in BAECs. Under glucose conditions, $(PhSe)_2$ positively impacted mitochondrial oxidative capacity, as assessed by respirometry, and was associated with changes in mitochondrial cellular dynamics. However, these changes were not observed in cells cultured with galactose. Although $(PhSe)_2$ induced the nuclear translocation of NRF2 in both glucose and galactose media, NRF2 remained in the nuclei of cells cultured in galactose for a longer duration. Additionally, activation of FOXO3a was only detected in galactose media. Notably, $(PhSe)_2$ strongly induced the expression of genes controlling mitochondrial antioxidant capacity and glutathione synthesis and recycling in glucose media, whereas its effects in galactose media were primarily focused on glutathione homeostasis. In conclusion, our findings underscore the critical influence of cellular metabolic status on the antioxidant capacity of redox-active molecules such as $(PhSe)_2$.

Key words: endothelial cell; diphenyl diselenide, glucose; galactose; mitochondria; antioxidants; oxidative metabolism; REDOX; FOXO3; NRF2.

Introduction

Cellular energy metabolism may vary depending on the tissue and cell type as well as on the energy substrates available and the energy demands. Cancer cells, in particular, commonly exhibit a more avid non-oxidative glycolytic metabolism than non-tumor cells [1], although they also need to adapt their metabolism in response to substrate availability, which results in corresponding changes in cell physiology and redox-bioenergetic status [2]. Similarly, pharmacology agents may have different cellular effects, depending on the energy metabolism status of the target cells [3].

The use of different cell culture media conditions provides a suitable model for evaluating cellular metabolic performance *in vitro*. While both glucose and galactose are good sources of cellular energy, galactose enters the glycolytic pathway through the energy-consuming Leloir pathway [4]. When glucose in the culture medium is replaced by galactose, cells are forced to shift to the mitochondrial oxidative metabolism to meet their energy demands. This condition not only activates mitochondrial OXPHOS but fully reprograms mitochondrial function and dynamis, and facilitates the evaluation of mitochondrial susceptibility to various agents revealing sensitivities related to oxidative metabolism that are obscured when cells are cultured in high glucose-containing medium [5][6]. In contrast, glucose catabolism tends to be primarily non-oxidative when the levels of glucose reach a certain threshold, known as the "Crabtree effect". As a result, cells grown in media with standard of glucose concentrations (25 mM) tend to acquire a highly non-oxidative glycolytic phenotype [7].

The activation of the mitochondrial OXPHOS is controlled by the master transcriptional coactivator PGC-1α and linked to the reprograming of the cellular capacity to regulate its REDOX status [8]. Furthermore, Nrf2 and FoxO, transcription factors that control antioxidant gene expression, and respond to oxidative stress

translocating to the nucleus [9][10], are also well know to be sensitive to the cellular

metabolic status [11][12], and are regulated by PGC-1α [13][14]. However, studies that

evaluate the redox activity of molecules rarely consider the differential effect of the

cellular metabolic status. In a recent study, we observed that, endothelial cells maintained

in a galactose-containing medium, not only exhibited higher mitochondrial oxidative

capacity and intercellular metabolic coupling than were cultured in galactosa, but also

show differences in the nuclear levels of NRF22 and FOXO3 which may be indicative of

a differential sensitivity to molecules with redox activity (Galant 2024), highlighting the

relevance of redox-metabolic coupling and the need to control for metabolism in cell

assys. Accordingly, it is reasonable to expect that pharmacological drug protocols will

have different cellular effects depending on the energy metabolism status of the target

cells.

Diphenyl diselenide (PhSe)$_2$, is a promising organoselenium compound with

antiatherogenic properties [15–17]. The cardiovascular actions of (PhSe)$_2$ have been

previously tested in several *in vivo* and *in vitro* models, including hypercholesterolemic

mice and rabbits, human LDL, as well as vascular endothelial and macrophage cells

[18][19].

(PhSe)$_2$, mechanism of action has been proposed to be related to is antioxidat

activity, it has been shown to react with reactive cellular redox-sensitive thiols, such as

cysteine and the tripeptide glutathione (GSH), an important antioxidant, functioning as a

glutathione peroxidase (GPx) mimic, and facilitating the recycling of reduced glutathione.

Additionally, (PhSe)$_2$ has been show to promote the oxidation of critical cysteinyl

residues in KEA1, a redox sensing protein that maintains Nrf2 in the cytoplasm, leading

to the NRF2 nuclear translocation and activation [20][16][21][22]. Furthermore, (PhSe)$_2$

has been shown to increase the levels of two important mitochondrial antioxidant

enzymes, PRDX3 and MnSOD [21] that are transcriptional targets of FOXO3a [13] and NRF2 [23], although it has to yet been stablished with unclear which transcription factor/s are mediate these effects.

Therefore, aiming to fully characterize the molecular mechanisms involved in the cardiovascular effects of $(PhSe)_2$ and taking into consideration the relevance of metabolic alterations in cardiovascular diseases, we analyzed how $(PhSe)_2$ impacted redox-bioenergetic cellular pathways under glycolytic or oxidative *in vitro* conditions in primary bovine vascular endothelial cells (BAEC).

Materials and Methods

Cell culture and treatments –BAEC were extracted from a fresh bovine thoracic aorta, obtained in an authorized slaughterhouse, as previously described by Peluffo and colleagues [24]. BAEC were cultured in growth medium (DMEM) supplemented with 10% of fetal bovine serum (FBS; Gibco/Invitrogen), containing 2 mM glutamine, 100 units/mL penicillin, 100 μg/mL streptomycin, 10 mM Hepes, 25 mM glucose, 44 mM $NaHCO_3$ and incubated at 37 °C in a humidified atmosphere with 5% CO_2. Cells suspensions were seeded in 96, 24 or 6-well plates (100 × 20 mm), at different cell densities, depending on the experimental procedure. Cells were allowed to reach 90% confluence and then submitted to the experimental protocols: i) When BAEC were submitted to the *glycolytic protocol*, culture media was changed to DMEM without FBS. Cells were treated with 1 μM $(PhSe)_2$ or vehicle (DMSO 0.05%) for 3, 6, 12, 24 or 48h. ii) When BAEC were submitted to the *oxidative protocol,* culture media was changed to glucose-free RPMI medium with 5 mM galactose and without FBS. Cells were maintained in this medium for 16 hours and then treated with 1 μM $(PhSe)_2$ or vehicle (DMSO 0.05%) for 3, 6, 12, 24 or 48h.

High-resolution respirometry by Seahorse- To evaluate mitochondrial oxygen consumption in BAEC, cells were plated in XF24 cell culture microplates (24-well plates at a cell density of 5×10^3 cells/well), submitted to the glycolytic (glucose) or oxidative (galactose) protocols and then treated with 1 μM (PhSe)$_2$ or vehicle for 3, 12 or 48 h. A calibration cartridge (Seahorse Bioscience) was equilibrated overnight and then loaded with unbuffered DMEM (port A), 0.6 μM oligomycin (port B), 0.3 μM FCCP (port C), and 0.1 μM rotenone plus 0.1 μM antimycin A (port D), all from Sigma-Aldrich. This allowed the determination of basal respiration, maximal respiration, reserve capacity, and non- mitochondrial respiration. In all experiments, the protein concentration in each well was determined at the end of the measurements, using the Pierce BCA protein assay kit (Thermo Scientific) following cell lysis with RIPA buffer (Sigma-Aldrich) supplemented with a protease inhibitor cocktail (Complete Mini; Roche), and used to calibrate the oxygen consumption data.

MitoSOX Imaging - Mitochondrial superoxide (O_2^-) was evaluated by MitoSOX Red (Molecular Probes, Carlsbad, CA, USA) staining, essentially as described [25]. Briefly, BAEC were grown on coverslips in 24-well culture (1×10^5 cells/well) plates, submitted to the glycolytic (glucose) or oxidative (galactose) protocols and treated with 1 μM (PhSe)$_2$ or vehicle (DMSO 0.05%) for 24h. Then, BAEC were exposed to 500 μM H_2O_2 for 4h or 80 μM DMNQ for 2h, and finally incubated with 3 μM MitoSOX Red for 10 min and fixed with paraformaldehyde (2%) and analyzed by fluorescence microscopy (Leica TCS SP5, Buffalo Grove, IL).

Immunofluorescence (IF)– BAEC were grown on coverslips, 24-well culture (1×10^5 cells/well) plates and submitted to the glycolytic (glucose) or oxidative (galactose)

protocols and then treated with 1 μM (PhSe)$_2$ or vehicle for 3, 6, 12, 24 or 48h. At the end

of the incubation period, cells were then fixed with 3.7 % formaldehyde, permeabilized

with 0.1 % Triton, and incubated consecutively with a primary antibody α-TOMM22

(1:200, Santa Cruz Biotechnology), α-NRF2 (1:200, Cell signaling) or α-FOXO3 (1:100,

Cell signaling) and then a secondary antibody (αIgG rabbit ALEXA-488 conjugate,

1:2500). Finally, cells were counterstained with DAPI, mounted and examined under a

confocal microscope (Zeiss LSM 700, Obercochen, Germany), as previously described

[25].

The total TOMM22 signal was used for the evaluation of the cellular

mitochondrial content. Mitochondrial fission was determined as the standard deviation of

TOMM22 intensity signal across the cytosol. Mitochondrial subcellular distribution was

also evaluated using TOMM22 signal. The perinuclear region signal was compared to

the total cytosolic signal, and the asymmetry of the signal in the perinuclear region was

also evaluated comparing the max and min signals in opposite nuclear sides. The nuclei

were identified by DAPI staining.

Protein extraction and Western blotting - BAEC (5 x 10^5 cells/well) were grown in 6-

well culture plates, submitted to the glycolytic (glucose) or oxidative (galactose)

protocols and treated with 1 μM (PhSe)$_2$ or vehicle for 3, 6, 12, 24 or 48 h. The cells were

then washed with phosphate-buffered saline (PBS) and lysed in 150 μl lysis buffer

containing 150 mM NaCl, 0.1% sodium dodecyl sulphate (SDS), 1% sodium

deoxycholate, 1% NP40 and 25 mM Tris–HCl pH 7.6, in the presence of protease

(Complete, Roche Diagnostics, Mannheim, Germany) and phosphatase inhibitors

(Sigma-Aldrich, St. Louis, MO). Cells were harvested by scraping, samples were clarified

by centrifugation at 13.000 rpm for 15 min at 4 °C and protein concentration was

determined using the BCA assay (Thermo Scientific, Rockford, IL, USA). Equal amounts

of protein (40 – 50 μg) from whole extracts were separated by SDS-PAGE

electrophoresis, using 10/12% acrylamide/bisacrylamide (29:1) 0.1% SDS gels and

transferred to a 0.2 μm PVDF Hybond-P membranes (Amersham/GE healthcare) at 20V

for 1h in a semi-dry Trans-Blot SD cell system (Bio-Rad, Hercules, CA). Following the

blockade of the membranes using TBS-T (20 mM Tris, 150mM NaCl, 0.1% Tween 20)

with 3% BSA, they were incubated with primary antibodies and then secondary

antibodies in blocking solution, α-FOXO3 (1:1000, Cell signaling), α-pFOXO3 (1:1000,

Cell signaling), α-AKT (1:1000, Cell signaling), α-pAKT (1:1000, Cell signaling), α-

MnSOD (1:3000, ADI-SOD-110, Enzo Life Sciences), α-PRDX3 (1:1000; LabFrontier,

Daehyun-dong Seodaemun-gu Seoul, Korea), α-GCLC (1:1000; Santa Cruz

Biotechnology, CA, USA), α-GCLM (1:1000; Santa Cruz Biotechnology, CA, USA) and

β-Tubulin (1:80000; Sigma, St. Louis, M), that was used as loading control. Secondary

antibodies α-rabbit and α-mouse IgG were from LI-COR Biosciences (Lincoln, NE).

Membranes were then extensively washed with TBS-T and developed using Bio-Rad

Clarity Western ECL Substrate. Images were captured using a luminometer scanner and

subject to densitometry analysis.

Image analysis - Image J software was used for the analysis of areas, signals in an area

and cross section signals from fluorescence and confocal microscopic images for

Tomm22, Nrf2, FOXO3 and MitoSOX and also to analyze Western bloting membranes.

Gene expression analysis of genes coding for antioxidant proteins by RT-qPCR - To

evaluate mRNA levels, BAEC (5 x 10^5 cells/well) were seeded in 6-well plates and

cultivated for 24h. Following cell treatment, total RNA was isolated using TRIzol

Reagent (Invitrogen, Carlsbad, CA) following the manufacturer instructions. 1 μg of total

RNA was then reverse transcribed by extension of random primers using M-MLV

(Promega, Madison,WI). Relative expression levels were determined by real-time PCR

(qPCR) in a 7900 Sequence Detection System (Applied Biosystems, Carlsbad, CA) using

the primer sets listed below/in Table 1.

Target	Forward	Reverse
GCLC	5'-gcacggcatcctccagttcct-3'	5'-tcggatggttggggtttgtcc-3'
GCLM	5'-gtgatgccaccagatttgact-3'	5'-cccactcgtgcgcttgaat-3'
GPx1	5'-ctgctcattgagaacgtagca-3'	5'tctcgccattcacctcgca-3'
MnSOD	5'-ggaacaacaggtcttatccccct-3'	5'-ttacttgctgcaagccgtgtatc-3'
Prx3	5'-ttcgggcttcgctcatccga-3'	5'-atggtatgaggaactggtgct-3'

Table 1 Primer pairs used in RT-qPCR to quantify gene expression changes

Statistical analysis and graphics. Numerical raw data was compiled and processed in

Excell files (Microsoft) and then final data was incorporated into GraphPad PRISM®

software version 6.0 for Windows (GraphPad Software, San Diego, CA, USA), that was

used for statistical analysis and preparation of graphs. Normal (Gaussian) distribution was

evaluated with the Shapiro-Wilk normality test. Significance of differences among data

sets was evaluated using unpaired test *t* or two-way of variance analysis (ANOVA), using

Bonferroni's correction for multiple comparisons. Results are expressed as mean ± SEM.

$p < 0.05$ was considered statistically significant. n ≥ 4 in all experiments. The number of

independent experiments is indicated in the corresponding figure legends.

Results

Effect of (PhSe)$_2$ on mitochondrial respiration in BAEC

We first tested the differential effect of (PhSe)$_2$ on mitochondrial respiration in BAEC

cultured in standard glucose containing media or in galatose-containing medium to force

the cells to rely on OXPHOS for ATP production. We found that 3h treatment with

$(PhSe)_2$ decreased maximum respiration in both glucose- and galactose-containing

medium (Fig. 1A, C and D) but the oxygen consumption rate (OCR) recovered after 48h

of treatment with $(PhSe)_2$ in both glucose- and galactose-containing medium (Fig. 1B, C

and D). Furthermore, when the maximum OCR was normalized for the mitochondrial

volume, estimated by immunofluorescence analysis of the mitochondrial protein

Tomm22, we found that this ratio was significantly increased following 48h of treatment

with $(PhSe)_2$ in glucose-containing medium, suggesting a higher respiratory capacity per

mitochondria in this context (Fig. 1E). However, in galactose-containing medium the

treatment with $(PhSe)_2$ did not significantly change the maximum

respiration/mitochondrial volume unit ratio (Fig. 1F).

Fig. 1 Effect of (PhSe)₂ on mitochondrial respiration in BAEC conditioned in glucose- or galactose-containing medium. (A) Representative respirometry assay of BAEC conditioned in glucose-containing medium for 3h and (B) 48h, data corrected for protein amount. (C) Maximum respiration in BAEC conditioned in glucose-containing medium. (D) Maximum respiration in BAEC conditioned in galactose-containing medium. (E) Ratio of maximum respiration per mitochondrial volume unit in BAEC treated with (PhSe)₂ for 3, 12 or 48h in glucose-containing medium and (F) galactose-containing medium. (A and B) Plots of O₂ consumption (OCR) quantification was corrected using total μg of protein and expressed as the percentage of treatment with (PhSe)₂ (black bars) with respect to vehicle group (dashed line). (C and D) Average points recorded in samples

with (PhSe)$_2$ after addition of FCCP/average points recorded in samples with vehicle after addition of FCCP x100. (E and F) The mitochondrial volume unit was analyzed by Tomm22 immunofluorescence of 4 independent experiments (Figure 2). The average of the samples after FCCP by Seahorse/average of Tomm22 was analyzed by immunofluorescence. Data were represented as mean ± SEM (n=4). *$p < 0.05$, indicate statistical difference from vehicle group by two-way ANOVA, followed by the Bonferroni's *post hoc* test.

Effect of (PhSe)$_2$ on mitochondrial dynamic in BAEC conditioned in glucose or galactose

medium

To understand the changes elicited in the mitochondria by (PhSe)$_2$ we monitored mitochondrial dynamics by immunofluorescence analysis of Tomm22 using confocal microscopy. Total mitochondrial volume, fusion and subcellular distribution were analyzed in BAEC treated with (PhSe)$_2$ for 3, 12 and 48h in glucose- or galactose-containing medium. We found that in glucose-containing medium (Fig 2A-E), although the treatment with (PhSe)$_2$ for 48h decreased total mitochondrial content (Fig. 2B), it increased mitochondrial fussion (Fig. 2C), reduced the mitochondrial content in the perinuclear region (Fig. 2D), consistent with the observed increase in oxidative capacity per mitochondria. Furthermore, it increased the asimetrical distribution of mitochondria within the cells, that could be related to intercellular coupling/organization of the epithelium polarity (Fig. 2E). In contrast, cells maintained in galactose-containing medium (Fig. 2F-J) did not change the total mitochondrial content (Fig. 2G), fission (Fig. 2H), or mitochondrial localization in the perinuclear region (Fig. 2I) in response to the treatment with (PhSe)$_2$. However, a transient increase in the the asimetrical distribution of mitochondria was observed at 12 h, that reverted after 48h of treatment (Fig. 2J).

Fig. 2 Effect of (PhSe)$_2$ on mitochondrial dynamics in BAEC conditioned in glucose- or galactose-containing medium. BAEC were treated with (PhSe)$_2$ (1 μM) or vehicle (DMSO) for 3, 12, or 48h in medium containing glucose (A-E) or galactose (F-J) and the mitochondrial parameters were determined by immunofluorescence of Tomm22 antibody. (A and F) Representative image of mitochondrial Tomm22 (green) in BAEC treated with (PhSe)$_2$, the white bars represent 50 μm. (B and G) Quantification of mitochondrial volume, (C and H) mitochondrial fission, (D and I) intensity of the perinuclear zone, (E and J) ratio of left/right intensity. Results were expressed as the percentage of treatment with (PhSe)$_2$ (white bars) with respect to vehicle group (dashed line). Data were represented as mean $\pm$ SEM (n=4). *$p < 0.05$, indicate statistical difference from vehicle group by unpaired t-test.

(PhSe)$_2$ prevents the mitochondrial superoxide production in BAEC exposed to oxidants.

To evaluate if (PhSe)$_2$ could modity the cellular capacity to respond to an oxidative challenge, we pre-treated BAEC with 1 μM of (PhSe)$_2$ and then exposed the cells to H$_2$O$_2$ (500 μM) or DMNQ (80 μM), an inhibitor of the mitochondrial electron transport chain that drives the formation of mitochondrial superoxide (O$_2^-$).

Mitochondrial RS generation was evaluated using the MitoSOX probe. In BAEC cultured

in glucose-containing medium, DMNQ induced a significant increase in O_2^- production.

Importantly, $(PhSe)_2$ pre-treatment reduced mitochondrial O_2^- levels in DMNQ challenged

cells. H_2O_2 in glucose cultured cells actually reduced mitochondrial O_2^- levels, and

$(PhSe)_2$ pre-treatment partially rescued H_2O_2 inhibition (Fig. 3A-B), suggesting that

$(PhSe)_2$ pre-treatment improved redox homeostasis in glucose cultured BAEC. In

contrast, in BAEC cultured in galactose-containing medium DMNQ did not generate a

significant increase in O_2^- production, possibly because in these conditions the

mitochondria are more resistant to DMNQ action, while H_2O_2 did. Furthermore, pre-

treatment with $(PhSe)_2$ reduced tmitochondrial O_2^- levels following H_2O_2 exposure (Fig.

3C-D). These results consistently support the idea that the $(PhSe)_2$ reduces the

mitochondrial accumulation of O_2^- levels.

Fig. 3 Effect of $(PhSe)_2$ on mitochondrial O_2^- production in BAEC conditioned in glucose-
or galactose-containing medium. BAEC were pretreated with $(PhSe)_2$ (1 µM) or vehicle
(DMSO) for 24h in medium containing glucose (A-B) or galactose (C-D) and subjected
to oxidants as DMNQ or H_2O_2. The $O_2^{•-}$ production was determined by confocal

microscopy through the MitoSOX probe. (A and C) Representative image of mitochondrial O_2^- production in BAEC pretreated with $(PhSe)_2$ and exposed to DMNQ (80 μM) or H_2O_2 (500 μM) (for 2 or 4h, respectively), the white bars represent 50 μm. (B and D) Quantification of mitochondrial O_2^- production (values x 10^4) in BAEC pretreated with $(PhSe)_2$ and exposed to DMNQ or H_2O_2. * $p < 0.05$, indicate statistical difference between the groups exposed to oxidants and $(PhSe)_2$, # $p < 0.05$, indicate statistical difference between vehicle and DMNQ or H_2O_2 groups by two-way ANOVA, followed by the Bonferroni's *post hoc* test.

Time-course effect of (PhSe)₂ in the expression of antioxidant proteins in BAEC

conditioned in glucose or galactose medium.

Next, we decided to evaluate if the observed effect of $(PhSe)_2$ on mitochondrial $O_2^{•-}$ levels could be derived from an increased cellular detoxification capacity. To that end we used targeted gene expression and protein analysis of some mayor antioxidant coding genes regulated by the two important redox sensitive transcription factors, NRF2 and FOXO3. In BAEC cultured in glucose-containing medium, following 6h of treatment with $(PhSe)_2$ we observed a significant increase in MnSOD and *SOD2* gene expression levels (Fig. 4A-B) as well as an increase in PRDX3 protein levels (Fig. 4C-D). Both MnSOD and PRDX3 are antioxidants proteints that localize in the mitochondrial matrix. We also monitored the effect of $(PhSe)_2$ on GCL, the enzyme of the limiting step for the synthesis of the antioxidant tripeptide glutathione, and GPx1 the main enzyme involved in the recovery of reduced gluthatione following its oxidation, in glucose-containing medium. GCL is made up of two subunits. We noted that GCLC protein and *GCLC* gene expression levels increased following 24h of treatment with $(PhSe)_2$ (Fig. 4E-F) as well as *GCLM* levels at a 6h of treatment (Fig. 4 G-H). Furthermore, *GPx1* levels were increased at 12 and 24h of treatment with $(PhSe)_2$ in BAEC cultured in in glucose-containing medium (Supplementary Fig. 1A). Similar results were observed in BAEC maintained in galactose-containing medium. We observed an increase in the mRNA levels of *GCLC* after 12h and GCLC protein content at 24h (Fig. 4 M-N), as well as an

increase in GCLM protein levels after 6h of treatment with (PhSe)$_2$ (Fig. 4 O-P). (PhSe)$_2$

also increase the mRNA expression of GPx1 at 12 and 24h of treatment in galactose-

containing medium (Supplementary Fig. 1B). In contrast, *SOD2* mRNA levels and

protein of MnSOD were unaltered (Fig. 4 I-J), while the PRDX3 levels decreased after

24h of treatment with (PhSe)$_2$ in galactose-containing medium (Fig. 4 K-L). The mRNA

levels of *CAT*, *UCP2*, *CYTC*, *GSS* and *NRF2* were unaltered by the treatment with (PhSe)$_2$

of BAEC maintained in either glucose- or galactose containing medium (Supplementary

Fig. 1 C-L). Overall these results suggest that in glucose containing media (PhSe)$_2$ induces

both the main mitochondrial and cytosolic antioxidant systems, while in galactose media

(PhSe)$_2$ induces mainly the cytosolic antioxidant systems, a result consistent with the

expected pre-activation of OXPHOS and mitochondrial antioxidant capacity in galactose

media.

Fig. 4 Effect of (PhSe)$_2$ in time-course expression profiles of antioxidant proteins in glucose- or galactose-containing medium. BAEC were treated with (PhSe)$_2$ or vehicle (DMSO) for 6, 12, 24 or 48h in glucose or galactose medium. Protein expression by Western Blotting analyses and mRNA expression by qPCR. Results were expressed as protein /β-tubulin ratio and the data were presented as the percentage of treatment with (PhSe)$_2$ (bars) with respect to vehicle group (dashed line). V = vehicle, D = (PhSe)$_2$. Data were represented as mean ± SEM (n=3-4). *$p < 0.05$, indicate statistical difference from vehicle group by two-way ANOVA, followed by the Bonferroni's *post hoc* test.

Effect of (PhSe)$_2$ on Nrf2 nuclear translocation of BAEC conditioned in glucose or

galactose medium.

Since (PhSe)$_2$ increased the levels of antioxidant genes known to be targeted ty

the transcription factor NRF2, we evaluated the effect of (PhSe)$_2$ on Nrf2 nuclear

translocation, an thus activation, in BAEC cultured in glucose- or galactose-containing

medium. We found that NRF2 nuclear localization was incrased following 3h of

incubation with (PhSe)$_2$ in BAEC cultured in both glucose and galactose containing

medium (Fig. 5A-B). The increased nuclear localization was gradually reduced overtime

in both glucose and galactose cultured cells, but was maintained over a more extended

period of time in galactose cultured cells (Fig. 5C-D), suggesting a more effective nuclear

retention in galactose. The separate evaluation of nuclear and cytosolic levels of NRF2

showed a significant increase in cytosolic NRF2 levels following 3 h of treatment with

(PhSe)$_2$ in BAEC maintained in glucose- containing medium but not in galactose cultured

cells possibly suggesting that global NRF2 levels stabilized by (PhSe)$_2$ in glucose but not

in galactose media (Supplementary Fig. 2). NRF2 degradation is a tightly regulated

process, cytosolic retention of NRF2 by KEAP1 is linked to its proteasomal degradation,

and therefore, KEAP1 release is generally associated not only to NRF2 nuclear

translocation but also to its stabilization [26]. In sum, NRF2 activation is detectable in

both glucose and galactose media, but in galactose media is likely to be effective for more

extended periods of time.

Fig. 5 Effect of (PhSe)$_2$ on Nrf2 translocation in BAEC conditioned in glucose- or galactose-containing medium. BAEC were treated with (PhSe)$_2$ (1 μM) or vehicle (DMSO) for 3, 6, 12, 24 and 48h in medium containing glucose (A-B) or galactose (C-D) and the immunofluorescence the Nrf2 were analyzed. (A and C) Representative image of immunofluorescence from Nrf2 (green) and DAPI (blue) in BAEC, the white bars represent 50 μm. (B and D) Ratio of Nrf2 in the nuclei/cytosol. Data were represented as mean ± SEM (n=4). * $p < 0.05$, indicate statistical difference from vehicle group by two-way ANOVA, followed by the Bonferroni's *post hoc* test.

Effect of (PhSe)₂ on FOXO3 activation in BAEC conditioned in glucose or galactose

medium.

Finally, we also analyzed the posible involvement of FOXO3 transcriptional

factor in (PhSe)₂ activation of antioxidant gene expresion. Treatment of BAEC in

glucose-containing medium with (PhSe)₂ for 6, 12 or 24h increased the level of FOXO3

phosphorylation (pFOXO3) at Thr32, a mark of AKT phosphorylation (pAKT),

suggesting an inhibition of FOXO3 activity and nuclear extrussion (Fig. 6A). However,

no significant changes in the nuclear/cytosolic ratio of FOXO3 (Fig 7) nor in the

cytosolicy and nuclear levels when analyzed separately (Supplementary Figure 3), and,

although an increase in AKT phosphorylation was detectable in response to (PhSe)₂, it

could only be observed at 24 h of treatment in glucose media. (Fig. 6C). In contrast,

although the nuclear/cytosolic ratio of FOXO3 transiently decreased at 3h of treatmen in

galactose media, (PhSe)₂ decreased pFOXO3/FOXO3 ratio at 24h, suggesting an increase

in FOXO3 activity and nuclear localization (Fig. 6B). In these conditions although the

changes in pAKT levels did not reach statistical significant, a consistent tendency to

lower pAKT/AKT ratio was detectable at 24h (Fig. 6D) as well as detectably higher

FOXO3 levels in both the nuclei and the cytosol at 12h of treatment (Supplementary

Figure 3). All together these results suggest that FOXO3 activity is more likely to play a

 relevant role in antioxidant gene regulation in galactose than in glucose cultured cells.

Fig. 6 Effect of (PhSe)2 on pFOXO3 and pAkt levels in BAEC conditioned in glucose- or galactose-containing medium. BAEC were treated with (PhSe)2 or vehicle (DMSO) for 6, 12, 24 and 48h in glucose (A-B) or galactose (C-D) medium to analyze pFOXO3 (A-B) and pAkt (C-D) protein content by Western blotting. Results are expressed as protein/β-tubulin ratio and the data is presented as the relative % of treatment with (PhSe)2 (bars) *vs* vehicle. V = vehicle, D = (PhSe)2. Data were represented as mean ± SEM (n=4-7). * $p < 0.05$, indicate statistical difference from vehicle group by two-way ANOVA, followed by the Bonferroni's *post hoc* test.

Fig. 7 Effect of (PhSe)$_2$ on the translocation of FOXO3 in BAEC conditioned in glucose-
or galactose-containing medium. BAEC were treated with (PhSe)$_2$ (1 μM) or vehicle
(DMSO) for 3, 6, 12, 24 and 48h in medium containing glucose (A) or galactose (B) and
the immunofluorescence for FOXO3 were analyzed. Ratio of FOXO3 in the
nuclei/cytosol. Data were represented as mean ± SEM (n=4). * $p < 0.05$, indicate
statistical difference from vehicle group (dashed line) by two-way ANOVA, followed by
the Bonferroni's *post hoc* test.

Discussion

Endothelial cells are the primary target of circulating redox-active molecules. Among

these, primary bovine aortic endothelial cells (BAEC), since preserving their original

metabolic plasticity, represent a relevant *in vitro* model for investigating the effects of

pharmacological agents across diverse metabolic contexts. The choice of cell culture

media profoundly influences the metabolic performance of cells *in vitro*. Our recent

findings, particularly concerning the modulation of transcription factors NRF2 and

FOXO3—key regulators of antioxidant gene expression—in BAECs, underscore the

significance of celular metabolic state. Specifically, alterations in cell culture conditions,

leading to heightened reliance on oxidative metabolism, demonstrate a consequential

impact on the efficacy of redox-active agents on vascular cells. These observations

suggest that the effectiveness of redox-active agents on endothelial cells may be

intricately linked to their metabolic milieu.

Therefore, we investigate the effects of (PhSe)$_2$ on BAEC cultured in media containing

glucose or galactose, assessing its impact on metabolism and antioxidant responses.

(PhSe)$_2$ has been extensively studied for its pivotal role in vascular physiology and

pathology, attributed to its capacity for redox modulation. This includes the promotion of

mitochondrial respiration and the upregulation of antioxidant gene expression, partly

mediated by the translocation of NRF2 to the nucleus [21]. Our findings reveal significant

differences in the action of (PhSe)$_2$ depending on whether BAEC are culture in galactose

and glucose media. These differences are likely to be relevant to their vascular effects *in*

vivo.

In cells cultured in glucose, treatment with (PhSe)$_2$ yielded a net positive impact on

mitochondrial oxidative capacity, assessed by respirometry. This enhancement correlated

with observed changes in mitochondrial cellular dynamics, characterized by more fused

mitochondria and a uniform distribution of mitochondria throughout the cells, with

reduced concentration in the perinuclear region. Conversely, these changes were not

observed in cells cultured with galactose, where the cells were already compelled to

utilize their oxidative capacity. Therefore, the modulation of mitochondrial function by

(PhSe)$_2$ appears to be effective only in cells exhibiting limited utilization of their

mitochondrial oxidative capacity.

Regarding (PhSe)$_2$ antioxidant capacity, we noted that although (PhSe)$_2$ induces the

nuclear translocation of NRF2 in both glucose and galactose media, NRF2 remains longer

in the nuclei of cells cultured in galactose media. This observation suggests a potential

preference for nuclear retention in the galactose environment. However, any speculation

regarding the underlying mechanism at this stage would be highly conjectural. Potential

explanations could involve interactions with other transcription factors [27] or the

inhibition of its nuclear extrusion [28].

The enhanced NRF2 activation profile observed in galactose media is even more

pronounced in the case of FOXO3a, where its activation is solely detectable in galactose

media, with reduced levels of pFOXO3a and increased nuclear protein levels. This shared

pattern suggests a more efficient induction of antioxidant gene expression in galactose

compared to glucose media. However, analysis of the gene expression changes elicited

by (PhSe)$_2$ on crucial antioxidant genes controlling mitochondrial antioxidant capacity

and glutathione synthesis and recycling reveals a stronger induction in glucose media. In

galactose media, the changes induced by (PhSe)$_2$ are not only smaller but also

predominantly focused on glutathione homeostasis, suggesting a more cytosolic, rather

than mitochondrial, effect.

Therefore, the induction of antioxidants genes by (PhSe)$_2$ in glucose media likely requires

additional transcription factors that are specially active in conditions of low mitochondrial

activity. Potential candidates include NFkB [29] a redox-sensitive transcription factor

controlling antioxidant gene expression, and c-MYC [30] among others. Additionally,

since (PhSe)$_2$ also enhances mitochondrial fission and maximal oxidative capacity, other

pathways may be involved. Acute inhibition of mitochondrial respiration could drive

compensatory mechanisms, such as the activation of PGC-1 α, which can induce both

antioxidant gene expression and mitochondrial fussion [8] while supporting FOXO3 [13]

and NRF2 [14] activation.

Despite the apparent lower capacity of (PhSe)$_2$ to induce antioxidant gene expression in

galactose media, assessments of mitochondrial antioxidant capacity using the

mitochondrial inhibitor DMNQ or the exogenous addition of H_2O_2 suggest a potentially

heightened antioxidant response in galactose-treated cells. This inference arises from the

substantial reduction elicited by (PhSe)$_2$ on mitochondrial superoxide ($O_2^{\bullet-}$) levels in

galactose-cultured cells following a challenge with H_2O_2. However, drawing definitive

conclusions from these observations is challenging. In galactose media, cells exhibit greater resistance to DMNQ, while in glucose media, they are highly susceptible to H_2O_2-induced mitochondrial damage, resulting in under basal levels of mitochondrial $O_2^{\bullet-}$. Notably, $(PhSe)_2$ partially rescues these aberrant levels and restores them to normal, consistent with its observed capacity to enhance mitochondrial oxidative capacity. Therefore, our data suggest that while $(PhSe)_2$ demonstrates greater efficacy in the activation of NRF2 and FOXO3a under pro-oxidative conditions, alternative pathways contribute to its enhancement of mitochondrial oxidative capacity and total antioxidant capacity when cells are cultured in glucose media. The observation that $(PhSe)_2$ induces acute inhibition of mitochondrial respiration shortly after treatment may indicate a significant role for mitochondrial function in this context. In sum, our findings underscore the critical influence of the cellular metabolic status on antioxidant capacity of redox-active molecules such as $(PhSe)_2$.

References

[1] Z. Abbaszadeh, S. Çeşmeli, Ç. Biray Avcı, Crucial players in glycolysis: Cancer progress, Gene 726 (2020) 144158. https://doi.org/10.1016/j.gene.2019.144158.

[2] Y. Liu, K. Birsoy, Metabolic sensing and control in mitochondria, Molecular Cell 83 (2023) 877–889. https://doi.org/10.1016/j.molcel.2023.02.016.

[3] E.A. Zaal, C.R. Berkers, The Influence of Metabolism on Drug Response in Cancer, Front Oncol 8 (2018) 500. https://doi.org/10.3389/fonc.2018.00500.

[4] M.J.C. de Kok, A.F. Schaapherder, R.C.I. Wüst, M. Zuiderwijk, J.A. Bakker, J.H.N. Lindeman, S.E. Le Dévédec, Circumventing the Crabtree effect in cell culture: A systematic review, Mitochondrion 59 (2021) 83–95. https://doi.org/10.1016/j.mito.2021.03.014.

[5] L.D. Marroquin, J. Hynes, J.A. Dykens, J.D. Jamieson, Y. Will, Circumventing the crabtree effect: Replacing media glucose with galactose increases susceptibility of hepG2 cells to mitochondrial toxicants, Toxicological Sciences 97 (2007) 539–547. https://doi.org/10.1093/toxsci/kfm052.

[6] V.M. Gohil, S.A. Sheth, R. Nilsson, A.P. Wojtovich, J.H. Lee, F. Perocchi, W. Chen, C.B. Clish, C. Ayata, P.S. Brookes, V.K. Mootha, Nutrient-sensitized screening for drugs that shift energy metabolism from mitochondrial respiration to glycolysis, Nat Biotechnol 28 (2010) 249–255. https://doi.org/10.1038/nbt.1606.

[7]   M.V. Golikov, V.T. Valuev-Elliston, O.A. Smirnova, A.V. Ivanov, Physiological Media in Studies of Cell Metabolism, Mol Biol 56 (2022) 629–637. https://doi.org/10.1134/S0026893322050077.

[8]   I. Valle, A. Álvarez-Barrientos, E. Arza, S. Lamas, M. Monsalve, PGC-1α regulates the mitochondrial antioxidant defense system in vascular endothelial cells, Cardiovasc Res 66 (2005) 562–573. https://doi.org/10.1016/j.cardiores.2005.01.026.

[9]   V. Ngo, M.L. Duennwald, Nrf2 and Oxidative Stress: A General Overview of Mechanisms and Implications in Human Disease, Antioxidants (Basel) 11 (2022) 2345. https://doi.org/10.3390/antiox11122345.

[10] K.A. Lasick, E. Jose, A.M. Samayoa, L. Shanks, K.W. Pond, C.A. Thorne, A.L. Paek, FOXO nuclear shuttling dynamics are stimulus-dependent and correspond with cell fate, Mol Biol Cell 34 (2023) ar21. https://doi.org/10.1091/mbc.E22-05-0193.

[11] J.M. DeBlasi, G.M. DeNicola, Dissecting the Crosstalk between NRF2 Signaling and Metabolic Processes in Cancer, Cancers (Basel) 12 (2020) 3023. https://doi.org/10.3390/cancers12103023.

[12] M. Monsalve, Y. Olmos, The Complex Biology of FOXO, Current Drug Targets 12 (2011) 1322–1350. https://doi.org/10.2174/138945011796150307.

[13] Y. Olmos, I. Valle, S. Borniquel, A. Tierrez, E. Soria, S. Lamas, M. Monsalve, Mutual dependence of Foxo3a and PGC-1alpha in the induction of oxidative stress genes, J Biol Chem 284 (2009) 14476–14484. https://doi.org/10.1074/jbc.M807397200.

[14] E. Navarro, L. Gonzalez-Lafuente, I. Pérez-Liébana, I. Buendia, E. López-Bernardo, C. Sánchez-Ramos, I. Prieto, A. Cuadrado, J. Satrustegui, S. Cadenas, M. Monsalve, M.G. López, Heme-Oxygenase I and PCG-1α Regulate Mitochondrial Biogenesis via Microglial Activation of Alpha7 Nicotinic Acetylcholine Receptors Using PNU282987, Antioxid Redox Signal 27 (2017) 93–105. https://doi.org/10.1089/ars.2016.6698.

[15] B. Fiuza, N. Subelzú, P. Calcerrada, M.R. Straliotto, L. Piacenza, A. Cassina, J.B.T. Rocha, R. Radi, A.F. De Bem, G. Peluffo, Impact of SIN-1-derived peroxynitrite flux on endothelial cell redox homeostasis and bioenergetics: Protective role of diphenyl diselenide via induction of peroxiredoxins, Free Radical Research 49 (2015) 122–132. https://doi.org/10.3109/10715762.2014.983096.

[16] A.F. De Bem, B. Fiuza, P. Calcerrada, P.M. Brito, G. Peluffo, T.C.P. Dinis, M. Trujillo, J.B.T. Rocha, R. Radi, L.M. Almeida, Protective effect of diphenyl diselenide against peroxynitrite-mediated endothelial cell death: A comparison with ebselen, Nitric Oxide - Biology and Chemistry 31 (2013) 20–30. https://doi.org/10.1016/j.niox.2013.03.003.

[17] M.A. Hort, M.R. Straliotto, P.M. Netto, J.B.T. Da Rocha, A.F. De Bem, R.M. Ribeiro-Do-Valle, Diphenyl diselenide effectively reduces atherosclerotic lesions in LDLr -/- Mice by attenuation of oxidative stress and inflammation, Journal of Cardiovascular Pharmacology 58 (2011) 91–101. https://doi.org/10.1097/FJC.0b013e31821d1149.

[18] M.R. Straliotto, M.A. Hort, B. Fiuza, J.B.T. Rocha, M. Farina, G. Chiabrando, A.F. de Bem, Diphenyl diselenide modulates oxLDL-induced cytotoxicity in macrophage by improving the redox signaling, Biochimie 95 (2013) 1544–1551. https://doi.org/10.1016/j.biochi.2013.04.008.

[19] A.F. de Bem, M. Farina, R. de L. Portella, C.W. Nogueira, T.C.P. Dinis, J.A.N. Laranjinha, L.M. Almeida, J.B.T. Rocha, Diphenyl diselenide, a simple glutathione peroxidase mimetic, inhibits human LDL oxidation in vitro, Atherosclerosis 201 (2008) 92–100. https://doi.org/10.1016/j.atherosclerosis.2008.02.030.

[20] S.G. Müller, N.S. Jardim, C.B. Quines, C.W. Nogueira, Diphenyl diselenide regulates Nrf2/Keap-1 signaling pathway and counteracts hepatic oxidative stress induced by

bisphenol A in male mice, Environmental Research 164 (2018) 280–287. https://doi.org/10.1016/j.envres.2018.03.006.

[21] R.L. Quispe, M.L. Jaramillo, L.S. Galant, D. Engel, A.L. Dafre, J.B. Teixeira da Rocha, R. Radi, M. Farina, A.F. de Bem, Diphenyl diselenide protects neuronal cells against oxidative stress and mitochondrial dysfunction: Involvement of the glutathione-dependent antioxidant system, Redox Biology 20 (2019) 118–129. https://doi.org/10.1016/j.redox.2018.09.014.

[22] D. Bartolini, J. Commodi, M. Piroddi, L. Incipini, L. Sancineto, C. Santi, F. Galli, Glutathione S-transferase pi expression regulates the Nrf2-dependent response to hormetic diselenides, Free Radical Biology and Medicine 88 (2015) 466–480. https://doi.org/10.1016/j.freeradbiomed.2015.06.039.

[23] M. Monsalve, S. Borniquel, I. Valle, S. Lamas, Mitochondrial dysfunction in human pathologies, Front Biosci 12 (2007) 1131–1153. https://doi.org/10.2741/2132.

[24] G. Peluffo, P. Calcerrada, L. Piacenza, N. Pizzano, R. Radi, Superoxide-mediated inactivation of nitric oxide and peroxynitrite formation by tobacco smoke in vascular endothelium: studies in cultured cells and smokers, Am J Physiol Heart Circ Physiol 296 (2009) H1781–H1792. https://doi.org/10.1152/ajpheart.00930.2008.

[25] S. Borniquel, N. García-Quintáns, I. Valle, Y. Olmos, B. Wild, F. Martínez-Granero, E. Soria, S. Lamas, M. Monsalve, Inactivation of Foxo3a and subsequent downregulation of PGC-1 alpha mediate nitric oxide-induced endothelial cell migration, Mol Cell Biol 30 (2010) 4035–4044. https://doi.org/10.1128/MCB.00175-10.

[26] F. He, X. Ru, T. Wen, NRF2, a Transcription Factor for Stress Response and Beyond, International Journal of Molecular Sciences 21 (2020) 4777. https://doi.org/10.3390/ijms21134777.

[27] W. Li, S. Yu, T. Liu, J.-H. Kim, V. Blank, H. Li, A.-N.T. Kong, Heterodimerization with small Maf proteins enhances nuclear retention of Nrf2 via masking the NESzip motif, Biochimica et Biophysica Acta (BBA) - Molecular Cell Research 1783 (2008) 1847–1856. https://doi.org/10.1016/j.bbamcr.2008.05.024.

[28] J.W. Kaspar, A.K. Jaiswal, Tyrosine phosphorylation controls nuclear export of Fyn, allowing Nrf2 activation of cytoprotective gene expression, FASEB J 25 (2011) 1076–1087. https://doi.org/10.1096/fj.10-171553.

[29] M.J. Morgan, Z. Liu, Crosstalk of reactive oxygen species and NF-κB signaling, Cell Res 21 (2011) 103–115. https://doi.org/10.1038/cr.2010.178.

[30] D.R. Wonsey, K.I. Zeller, C.V. Dang, The c-Myc target gene PRDX3 is required for mitochondrial homeostasis and neoplastic transformation, Proc Natl Acad Sci U S A 99 (2002) 6649–6654. https://doi.org/10.1073/pnas.102523299.