Differential effects of diphenyl diselenide (PhSe)$_2$ on mitochondria-related pathways depending on the cellular energy status in Bovine Vascular Endothelial Cells

Letícia Selinger Galant [a,b], Laura Doblado [b], Rafael Radi [c], João Batista Teixeira da Rocha [d], Andreza Fabro de Bem [a,e,f*], Maria Monsalve [b*]

[a] Biochemistry PhD Program, Federal University of Santa Catarina, Florianópolis, SC, Brazil.

[b] Biomedicine Research Institute Sols-Morreale (CSIC-UAM). Arturo Duperier 4. 28029, Madrid, Spain.

[c] Center for Free Radical and Biomedical Research (CEINBIO), Facultad de Medicina, Universidad de la República, Montevideo, Uruguay

[d] Department of Biochemistry and Molecular Biology, Federal University of Santa Maria, Santa Maria, Brazil.

[e] Laboratory of Bioenergetics and Metabolism, Institute for Biological Sciences; University of Brasília, Brasília, Brazil.

[f] National Institute of Science and Technology on Neuroimmunomodulation, Rio de Janeiro, Brazil.

Abstract

Cellular energy metabolism varies depending on tissue and cell type, as well as the availability of energy substrates and energy demands. We recently investigated the variations in cellular metabolism and antioxidant responses in primary bovine vascular endothelial cells (BAECs) under different energetic substrate conditions *in vitro*, specifically glucose or galactose. In this context, pharmacological agents may affect cells differently depending on their energy metabolism status. In this study, we aimed to characterize the effects of diphenyl diselenide ($(PhSe)_2$), a redox-active molecule known for its prominent cardiovascular effects, on redox-bioenergetic cellular pathways under glycolytic or oxidative conditions in BAECs. Under glucose conditions, $(PhSe)_2$ positively impacted mitochondrial oxidative capacity, as assessed by respirometry, and was associated with changes in mitochondrial cellular dynamics. However, these changes were not observed in cells cultured with galactose. Although $(PhSe)_2$ induced the nuclear translocation of the redox sensitive nuclear factor erythroid 2-related factor 2 (Nrf2) in both glucose and galactose media, Nrf2 remained in the nuclei of cells cultured in galactose for a longer duration. Additionally, activation of another redox sensitive transcription factor, forkhead O3 (FOXO3a) was only detected in galactose media. Notably, $(PhSe)_2$ induced the expression of genes controlling mitochondrial antioxidant capacity and glutathione synthesis and recycling in glucose media, whereas its effects in galactose media were primarily focused on glutathione homeostasis. In conclusion, our findings underscore the critical influence of cellular metabolic status on the antioxidant capacity of redox-active molecules such as $(PhSe)_2$.

Key words*: endothelial cell; diphenyl diselenide, glucose; galactose; mitochondria; FOXO3; Nrf2.

1. Introduction

Cellular energy metabolism may vary depending on the tissue and cell type, the energy substrates available, and the energy demands. Cancer cells, in particular, commonly exhibit a more avid non-oxidative glycolytic metabolism than non-tumor cells [1]. However, they also need to adapt their metabolism in response to substrate availability, which results in corresponding changes in cell physiology and redox-bioenergetic status [2]. Similarly, pharmacology agents may have different cellular effects, depending on the energy metabolism status of the target cells [3].

The use of different cell culture media conditions provides a suitable model for evaluating cellular metabolic performance *in vitro*. While both glucose and galactose are good cellular energy sources, galactose enters the glycolytic pathway through the energy-consuming Leloir pathway [4]. When glucose in the culture medium is replaced by galactose (5-10 mM), cells are forced to shift to the mitochondrial oxidative metabolism to meet their energy demands. This condition not only activates mitochondrial oxidative phosphorylation OXPHOS but fully reprograms mitochondrial function and dynamics and facilitates the evaluation of mitochondrial susceptibility to various agents, revealing sensitivities related to oxidative metabolism that are obscured when cells are cultured in high glucose-containing medium [5,6]. In contrast, glucose catabolism tends to be primarily non-oxidative when glucose levels reach a certain threshold, known as the "Crabtree effect." As a result, cells grown in media with a standard of glucose concentrations (25 mM) tend to acquire a highly non-oxidative glycolytic phenotype [7].

The activation of the mitochondrial OXPHOS is controlled by the master transcriptional coactivator peroxisome proliferator-activated receptor gamma coactivator -1alpha (PGC-1α) and linked to the reprogramming of the cellular capacity to regulate its REDOX status [8]. Furthermore, Nrf2 and FOXO3, transcription factors that control

antioxidant gene expression, and respond to oxidative stress translocating to the nucleus [9,10], are also well known to be sensitive to the cellular metabolic status [11,12], and are regulated by PGC-1α [13,14]. However, studies that evaluate the redox activity of molecules rarely consider the differential effect of the cellular metabolic status. In a recent study, we observed that endothelial cells maintained in a galactose-containing medium, not only exhibited higher mitochondrial oxidative capacity and intercellular metabolic coupling than were cultured in galactose but also showed differences in the nuclear levels of Nrf2 and FOXO3 which may be indicative of differential sensitivity to molecules with redox activity [15], highlighting the relevance of redox-metabolic coupling and the need to control for metabolism in cell assays. Accordingly, it is reasonable to expect that pharmacological drug protocols will have different cellular effects depending on the energy metabolism status of the target cells.

(PhSe)$_2$ is a promising organoselenium compound with antiatherogenic properties [16–18]. The cardiovascular actions of (PhSe)$_2$ have been previously tested in several *in vivo* and *in vitro* models, including hypercholesterolemic mice and rabbits, human low-density lipoprotein (LDL), and vascular endothelial and macrophage cells [19,20]. (PhSe)$_2$, mechanism of action has been proposed to be related to its antioxidant activity. It has been shown to react with reactive cellular redox-sensitive thiols, such as cysteine and the tripeptide glutathione (GSH), an important antioxidant, functioning as a glutathione peroxidase (GPx) mimic, and facilitating the recycling of reduced glutathione. Additionally, (PhSe)$_2$ promotes the oxidation of critical cysteinyl residues in kelch-like ECH-associated protein 1 (Keap1), a redox-sensing protein that retains Nrf2 in the cytoplasm. This oxidation leads to the translocation of Nrf2 into the nucleus and its subsequent activation [16,21–23]. Furthermore, (PhSe)$_2$ has been shown to increase the levels of two key mitochondrial antioxidant enzymes, peroxiredoxin 3 (Prx3) and

manganese-dependent superoxide dismutase (MnSOD) [22], which are transcriptional targets of FOXO3 [13] and Nrf2 [24]. However, it must yet be established which transcription factor/s mediate these effects.

Therefore, aiming to fully characterize the molecular mechanisms involved in the cardiovascular effects of (PhSe)$_2$ and considering the relevance of metabolic alterations in cardiovascular diseases, we analyzed how (PhSe)$_2$ impacted redox-bioenergetic cellular pathways under glycolytic or oxidative *in vitro* conditions in primary bovine vascular endothelial cells (BAEC).

2. Materials and Methods

2.1. Cell culture and treatments

BAEC were extracted from a fresh bovine thoracic aorta, obtained in an authorized slaughterhouse, as previously described by Peluffo and colleagues [25]. BAEC were cultured in growth medium (DMEM) supplemented with 10% of fetal bovine serum (FBS; Gibco/Invitrogen), containing 2 mM glutamine, 100 units/mL penicillin, 100 µg/mL streptomycin, 10 mM HEPES, 25 mM glucose, 44 mM NaHCO$_3$ and incubated at 37°C in a humidified atmosphere with 5% CO$_2$. Cell suspensions were seeded in 96, 24 or 6-well plates (100 × 20 mm), at different cell densities, depending on the experimental procedure. Cells at passages P4–P8 were used to ensure they maintained sufficient mitochondrial plasticity. Cells were allowed to reach 90% confluence and under *glycolytic or oxidative conditions,* as previously described [15], treated with 1 µM (PhSe)$_2$ or vehicle (DMSO 0.05%) for 3, 6, 12, 24 or 48h. In short, when BAEC were submitted to the *glycolytic protocol*, culture media was changed to DMEM without FBS and treated with 1 µM (PhSe)$_2$ or vehicle (DMSO 0.05%) for 3, 6, 12, 24 or 48h. When BAEC were submitted to the *oxidative protocol,* culture media was changed to glucose-free RPMI medium with 5 mM galactose and without FBS. Cells were maintained in this medium

for 16 hours and then treated with 1 μM (PhSe)$_2$ or vehicle (DMSO 0.05%) for 3, 6, 12,

24 or 48h. For each independent experiment, identical cells, derived from the same

original culture dish, were split and then exposed to glucose and galactose media

separately. Therefore, all the cells within each independent experiment had the same

passage. The use of 1 μM (PhSe)$_2$ dosage is based on previous studies from our group

and others that demonstrated the protective effect of (PhSe)$_2$ on BAECs at a concentration

of 1 μM, as well as the absence of cytotoxic effects [16,17].

2.2. High-resolution respirometry by Seahorse

To evaluate mitochondrial oxygen consumption in BAEC, cells were plated in

XF24 cell culture microplates (24-well plates at a cell density of 5 x 10^3 cells/well),

submitted to the glycolytic (glucose) or oxidative (galactose) protocols and then treated

with 1 μM (PhSe)$_2$ or vehicle for 3, 12 or 48h. A calibration cartridge (Seahorse

Bioscience) was equilibrated overnight and then loaded with unbuffered DMEM (port

A), 0.6 μM oligomycin (port B), 0.3 μM carbonyl cyanide 4-

(trifluoromethoxy)phenylhydrazone (FCCP) (port C), and 0.1 μM rotenone plus 0.1 μM

antimycin A (port D), all from Sigma-Aldrich. This allowed the determination of basal

respiration, maximal respiration, reserve capacity, and non- mitochondrial respiration. In

all experiments, the protein concentration in each well was determined at the end of the

measurements, using the Pierce BCA protein assay kit (Thermo Scientific) following cell

lysis with RIPA buffer (Sigma-Aldrich) supplemented with a protease inhibitor cocktail

(Complete Mini; Roche), and used to calibrate the oxygen consumption data.

2.3. MitoSOX Imaging

Mitochondrial superoxide (O$_2^-$) was evaluated by MitoSOX Red (Molecular

Probes, Carlsbad, CA, USA) staining, essentially as described [26]. Briefly, BAEC were

grown on coverslips in 24-well culture (1 x 10^5 cells/well) plates, submitted to the

glycolytic (glucose) or oxidative (galactose) protocols and treated with 1 μM $(PhSe)_2$ or

vehicle (DMSO 0.05%) for 24h. Then, BAEC were exposed to 500 μM hydrogen

peroxide (H_2O_2) for 4h or 80 μM 1,4-naphthoquinone (DMNQ) for 2h, and finally

incubated with 3 μM MitoSOX Red for 10 min and fixed with paraformaldehyde (2%)

and analyzed by fluorescence microscopy (Nikon 90i). Under our experimental

conditions, no nuclear fluorescence was detected, indicating that the probe remained

confined to the mitochondria. Therefore, nuclear counterstaining with DAPI or equivalent

dyes was not required for image segmentation. For each slide, four images were acquired,

and the total fluorescence intensity was determined using ImageJ software. The average

fluorescence from the four images was then calculated and considered as the

representative fluorescence value for one replica sample.

2.4. Immunofluorescence (IF)

BAEC were grown on coverslips, 24-well culture (1 x 10^5 cells/well) plates and

submitted to the glycolytic (glucose) or oxidative (galactose) protocols and then treated

with 1 μM $(PhSe)_2$ or vehicle for 3, 6, 12, 24 or 48h. At the end of the incubation period,

cells were then fixed with 3.7 % formaldehyde, permeabilized with 0.1 % Triton, and

incubated consecutively with a primary antibody Tomm22 (1:200, Sigma Life Science,

cat: HPA003037), Nrf2 (1:200, Thermo Fisher, cat: PA5-27882) or FOXO3 (1:100, Cell

signaling, cat: 9467) and then a secondary antibody (IgG rabbit ALEXA-488 conjugate,

1:2500). Finally, cells were counterstained with DAPI, mounted and examined under a

confocal microscope (Zeiss LSM 700, Obercochen, Germany), as previously described

[26].

Tomm22 was used as a marker to assess mitochondrial content and subcellular

distribution using Image J software analysis of confocal microscopy images. Total

Tomm22 positive area was considered to indicate mitochondrial volume. Subcellular

distribution was analyzed by drawing a ling along the longitudinal cell axis and by

calculating the ratio between the perinuclear Tomm22 signal (15 pixels from the nuclear

edge) and the total cytosolic signal average provided by that line. Signal asymmetry in

the perinuclear region was assessed by calculating the rate of the maximum and minimum

intensity values on opposite sides of the nucleus as divided by the above-mentioned cross-

section line. Using this line, mitochondrial fission was estimated by the calculation of the

standard deviation of Tomm22 signal across the cytosol. The nucleus was identified using

the DAPI staining signal, this analysis procedure was as previously described [15,27].

Since BAECs are primary cells with significant variability in mitochondrial and

respiratory parameters depending on passage number [15], all data were normalized to

the $t = 0$ values of each replica and expressed as a percentage in the graphs.

For the analysis of nuclear and cytosolic content of Nrf2 and FOXO3, as previously

described [15], a similar procedure was used, using Imagen J Software, a line was drawn

across the long axis of each cell the cellular compartment and the nuclear region was

identified based on DAPI fluorescence signal. The Nrf2 average fluorescence signal

corresponding to the and cytosolic regions was then calculated and then ratio of both,

resulting in the nuclear-cytosolic ratio.

2.5. Protein extraction and Western blotting

BAEC (5×10^5 cells/well) were grown in 6-well culture plates, submitted to the

glycolytic (glucose) or oxidative (galactose) protocols and treated with 1 μM (PhSe)$_2$ or

vehicle for 3, 6, 12, 24 or 48h. The cells were then washed with phosphate-buffered saline

(PBS) and lysed in 150 μl lysis buffer containing 150 mM NaCl, 0.1% sodium dodecyl

sulphate (SDS), 1% sodium deoxycholate, 1% NP40 and 25 mM Tris–HCl pH 7.6, in the

presence of protease (Complete, Roche Diagnostics, Mannheim, Germany) and

phosphatase inhibitors (Sigma-Aldrich, St. Louis, MO). Cells were harvested by scraping,

samples were clarified by centrifugation at 13.000 rpm for 15 min at 4 °C and protein

concentration was determined using the BCA assay (Thermo Scientific, Rockford, IL,

USA). Equal amounts of protein (40 – 50 μg) from whole extracts were separated by

SDS-PAGE electrophoresis, using 10/12% acrylamide/bisacrylamide (29:1) 0.1% SDS

gels and transferred to a 0.2 μm PVDF Hybond-P membranes (Amersham/GE healthcare)

at 20V for 1h in a semi-dry Trans-Blot SD cell system (Bio-Rad, Hercules, CA).

Following the blockade of the membranes using TBS-T (20 mM Tris, 150mM NaCl,

0.1% Tween 20) with 3% BSA, they were incubated with primary antibodies and then

secondary antibodies in blocking solution, FOXO3 (1:1000, Cell signaling, cat: 9467),

pFOXO3 (1:1000, Cell signaling, cat: 2309), Akt (1:1000, Cell signaling, cat: 2920), pAkt

(1:1000, Cell signaling, cat: 4060), MnSOD (1:3000, ADI-SOD-110, Enzo Life Sciences,

cat: ADI-SOD-110-F), Prx3 (1:1000; LabFrontier, Daehyun-dong Seodaemun-gu Seoul,

Korea, cat: LF-PA0030), GCLC (1:1000; Cell Signaling, cat: 48005), GCLM (1:1000;

Sigma-Aldrich, cat: SAB5700737) and β-Tubulin (1:80000; Sigma, St. Louis, M, cat:

T9026), that was used as loading control. Secondary antibodies rabbit and mouse IgG

were from LI-COR Biosciences (Lincoln, NE). Membranes were then extensively washed

with TBS-T and developed using Bio-Rad Clarity Western ECL Substrate. Images were

captured using a luminometer scanner and subject to densitometry analysis.

2.6. Image analysis

Image J software was used for the analysis of areas, signals in an area and cross

section signals from fluorescence and confocal microscopic images for Tomm22, Nrf2,

FOXO3 and MitoSOX and to analyze Western blotting membranes.

2.7. Gene expression analysis of genes coding for antioxidant proteins by RT-qPCR

To evaluate mRNA levels, BAEC (5×10^5 cells/well) were seeded in 6-well plates

and cultivated for 24h. Following cell treatment, total RNA was isolated using TRIzol

Reagent (Invitrogen, Carlsbad, CA) following the manufacturer instructions. 1 μg of total

RNA was then reverse transcribed by extension of random primers using M-MLV

(Promega, Madison,WI). Relative expression levels were determined by real-time PCR

(qPCR) in a 7900 Sequence Detection System (Applied Biosystems, Carlsbad, CA) using

the primer sets listed below/in Table 1.

Table 1 Primer pairs used in RT-qPCR to quantify gene expression changes.

Target	Forward	Reverse
GCLC	5'-gcacggcatcctccagttcct-3'	5'-tcggatggttggggtttgtcc-3'
GCLM	5'-gtgatgccaccagatttgact-3'	5'-cccactcgtgcgcttgaat-3'
GPx1	5'-ctgctcattgagaacgtagca-3'	5'tctcgccattcacctcgca-3'
MnSOD	5'-ggaacaacaggtcttatccccct-3'	5'-ttacttgctgcaagccgtgtatc-3'
Prx3	5'-ttcgggcttcgctcatccga-3'	5'-atggtatgaggaactggtgct-3'

2.8. Statistical analysis and graphics

Numerical raw data was compiled and processed in Excel files (Microsoft) and

then final data was incorporated into GraphPad PRISM® software version 6.0 for

Windows (GraphPad Software, San Diego, CA, USA), that was used for statistical

analysis and preparation of graphs. Normal (Gaussian) distribution was evaluated with

the Shapiro-Wilk normality test. Significance of differences among data sets was

evaluated using unpaired test *t* or two-way of variance analysis (ANOVA), using

Bonferroni's correction for multiple comparisons. Results are expressed as mean ± SD. *p*

<0.05 was considered statistically significant. n ≥ 3 in all experiments. The number of

independent experiments is indicated in the corresponding figure legends. Each data point

incorporated in the graphs represents an independent experiment, not technical replicas.

Independent means they used an independent vial of cells, thaw on a different day and

thus tested on different days. Data from technical replicas/individual cells/individual

images were always pooled to calculate each data point. The graphical abstract was

created in BioRender De bem, L. (2025) https://BioRender.com/pu1070I under the

agreement number XK28BVWXKS.

3. Results

3.1. Effect of (PhSe)$_2$ on mitochondrial respiration in BAEC

We first tested the differential effect of (PhSe)$_2$ on mitochondrial respiration in

BAEC cultured in standard glucose-containing media or in galactose-containing medium

to force the cells to rely on OXPHOS for adenosine triphosphate (ATP) production. All

control data comparing alterations in BAECs cultured in glucose- versus galactose-

containing media have been recently published in [15]. In that study we observed that the

maximum oxidative capacity after 12–48 h of incubation of BAEC conditioned in culture

medium containing galactose was significantly higher when compared to that observed

in cells in culture medium containing glucose. Now, we found that 3h treatment with

(PhSe)$_2$ decreased maximum respiration in both glucose- ($F_{(1, 19)} = 4{,}959$, $p = 0.0382$)

and galactose- ($F_{(2, 12)} = 10{,}56$, $p = 0.0023$) containing medium (Fig. 1A, C and D) but

the oxygen consumption rate (OCR) recovered after 48h of treatment with (PhSe)$_2$ in both

glucose- and galactose-containing medium (Fig. 1B, C and D). Furthermore, when the

maximum OCR was normalized for the mitochondrial volume, estimated by

immunofluorescence analysis of the mitochondrial protein Tomm22, we found that this

ratio was significantly increased following 48h of treatment with (PhSe)$_2$ in glucose-

containing medium ($t = 3.512$, $p = 0.0246$), suggesting a higher respiratory capacity per

mitochondria in this context (Fig. 1E). However, in a galactose-containing medium the

treatment with (PhSe)$_2$ did not significantly change the maximum

respiration/mitochondrial volume unit ratio (Fig. 1F).

Fig. 1 Effect of (PhSe)$_2$ on mitochondrial respiration in BAEC conditioned in glucose- or galactose-containing medium. (A) Representative respirometry assay of BAEC conditioned in glucose-containing medium for 3h and (B) 48h, data corrected for protein amount. (C) Maximum respiration in BAEC conditioned in glucose-containing medium. (D) Maximum respiration in BAEC conditioned in a galactose-containing medium. (E) Ratio of maximum respiration per mitochondrial volume unit in BAEC treated with (PhSe)$_2$ for 3, 12 or 48h in glucose-containing medium and (F) galactose-containing medium. (A and B) Plots of O$_2$ consumption (OCR). (C and D) Quantification was corrected using total µg of protein and expressed as the percentage of treatment with (PhSe)$_2$ (white bars) with respect to vehicle group (dashed line). Average points recorded in samples with (PhSe)$_2$ after the addition of FCCP/average points recorded in samples with vehicle after the addition of FCCP x100. (E and F) The mitochondrial volume unit was analyzed by Tomm22 immunofluorescence in 5 independent experiments (Figure 2). The average of the samples after FCCP by Seahorse/average of Tomm22 was analyzed by immunofluorescence. Data were represented as mean ± SD (n=3-5). (C and D) *$p <$ 0.05, indicate statistical difference from the vehicle group by two-way ANOVA, followed

by Bonferroni's *post hoc* test, (E and F) *$p < 0.05$, indicates statistical difference from vehicle group by paired t-test.

3.2. Effect of (PhSe)$_2$ on mitochondrial dynamic in BAEC conditioned in glucose or galactose medium

To understand the changes elicited in the mitochondria by (PhSe)$_2$, we monitored mitochondrial dynamics by immunofluorescence analysis of Tomm22 using confocal microscopy. Total mitochondrial volume, fusion, and subcellular distribution were analyzed in BAEC treated with (PhSe)$_2$ for 3, 12, and 48h in glucose- or galactose-containing medium. We found that in glucose-containing medium (Fig 2A-E), although the treatment with (PhSe)$_2$ for 48h decreased total mitochondrial content ($t = 3.161$, $p = 0.0195$) (Fig. 2B), it increased mitochondrial fusion ($t = 3.022$, $p = 0.0233$) (Fig. 2C), reduced the mitochondrial content in the perinuclear region ($t = 3.218$, $p = 0.0182$) (Fig. 2D), consistent with the observed increase in oxidative capacity per mitochondria. Furthermore, it increased the asymmetrical distribution of mitochondria within the cells, which could be related to intercellular coupling/organization of the epithelium polarity ($t = 3.506$, $p = 0.0172$) (Fig. 2E). In contrast, cells maintained in galactose-containing medium (Fig. 2F-J) did not change the total mitochondrial content (Fig. 2G), fission (Fig. 2H), or mitochondrial localization in the perinuclear region (Fig. 2I) in response to the treatment with (PhSe)$_2$. However, a transient increase in the asymmetrical distribution of mitochondria was observed at 12h ($t = 3.041$, $p = 0.0287$), which reverted after 48h ($t = 5.916$, $p = 0.001$) of treatment (Fig. 2J).

Fig. 2 Effect of (PhSe)₂ on mitochondrial dynamics in BAEC conditioned in glucose- or galactose-containing medium. BAEC were treated with (PhSe)₂ (1 μM) or vehicle (DMSO) for 3, 12, or 48h in medium containing glucose (A-E) or galactose (F-J), and the mitochondrial parameters were determined by immunofluorescence of Tomm22 antibody. (A and F) Representative image of mitochondrial Tomm22 (green) in BAEC treated with (PhSe)₂. The white bars represent 50 μm. (B and G) Quantification of mitochondrial volume, (C and H) mitochondrial fission, (D and I) intensity of the perinuclear zone, (E and J) ratio of left/right intensity. Results were expressed as the percentage of treatment with (PhSe)₂ (white bars) with respect to the vehicle group (dashed line). Data were represented as mean ± SD (n=3-4). *$p < 0.05$, indicates statistical difference from vehicle group by unpaired t-test.

3.3. (PhSe)₂ prevents the mitochondrial superoxide production in BAEC exposed to oxidants.

To evaluate if (PhSe)₂ could modify the cellular capacity to respond to an oxidative challenge, we pre-treated BAEC with 1 μM of (PhSe)₂ and then exposed the cells to H_2O_2 (500 μM) or DMNQ (80 μM), an inhibitor of the mitochondrial electron transport chain that drives the formation of mitochondrial superoxide (O_2^-). Mitochondrial RS generation was evaluated using the MitoSOX probe. In BAEC cultured

in glucose-containing medium, DMNQ significantly increased O_2^- production.

Importantly, $(PhSe)_2$ pre-treatment reduced mitochondrial O_2^- levels in DMNQ

challenged cells ($F(1,8) = 7.35$, $p = 0.0266$). H_2O_2 in glucose-cultured cells also increased

mitochondrial O_2^- levels and, consistently, $(PhSe)_2$ pre-treatment reduced H_2O_2 levels (F

$(1,7) = 13.66$, $p = 0.0077$) (Fig. 3A-B), suggesting that $(PhSe)_2$ pre-treatment improved

mitochondrial homeostasis in glucose cultured BAEC. In contrast, in BAEC cultured in

galactose-containing medium, DMNQ did not generate a significant increase in O_2^-

production, possibly because in these conditions the mitochondria are more resistant to

DMNQ action, while H_2O_2 did. Furthermore, pre-treatment with $(PhSe)_2$ reduced

mitochondrial O_2^- levels following H_2O_2 exposure ($F(1,7) = 9.171$, $p = 0.0192$) (Fig. 3C-

D). These results consistently support the idea that the $(PhSe)_2$ reduces the mitochondrial

accumulation of O_2^- levels.

Fig. 3 Effect of $(PhSe)_2$ on mitochondrial O_2^- production in BAEC conditioned in glucose-
or galactose-containing medium. BAEC were pretreated with $(PhSe)_2$ (1 µM) or vehicle
(DMSO) for 24h in medium containing glucose (A-B) or galactose (C-D) and subjected
to oxidants as DMNQ or H_2O_2. The O_2^- production was determined by Nikon 90i
microscopy through the MitoSOX probe. (A and C) Representative image of
mitochondrial O_2^- production in BAEC pretreated with $(PhSe)_2$ and exposed to DMNQ
(80 µM) or H_2O_2 (500 µM) (for 2 or 4h, respectively), the white bars represent 50 µm. (B

and D) Quantification of mitochondrial O_2^- production (values x 10^4) in BAEC pretreated
with $(PhSe)_2$ and exposed to DMNQ or H_2O_2. * $p < 0.05$, indicates statistical difference
between the groups exposed to oxidants and $(PhSe)_2$, # $p < 0.05$, indicates the statistical
difference between vehicle and DMNQ or H_2O_2 groups by two-way ANOVA, followed
by Bonferroni's *post hoc* test.
355

3.4. Time-course effect of $(PhSe)_2$ in the expression of antioxidant proteins in BAEC

conditioned in glucose or galactose medium.

Next, we decided to evaluate if the observed effect of $(PhSe)_2$ on mitochondrial

O_2^- levels could be derived from an increased cellular detoxification capacity. To that

end, we used targeted gene expression and protein analysis of some major antioxidant

coding genes regulated by the two crucial redox-sensitive transcription factors, Nrf2 and

FOXO3. In BAEC cultured in glucose-containing medium, following 6h of treatment

with $(PhSe)_2$ we observed a significant increase in MnSOD and *SOD2* gene expression

levels ($F (3,47) = 5.159, p = 0.0036$ and $F (1,29) = 7.157, p = 0.0121$, respectively) (Fig.

4A-B) as well as an increase in Prx3 protein levels ($F (3,36) = 2.722, p = 0.053$) (Fig. 4C-

D). Both MnSOD and Prx3 are antioxidant proteins that localize in the mitochondrial

matrix. We also monitored the effect of $(PhSe)_2$ on GCL, the enzyme of the limiting step

for the synthesis of the antioxidant tripeptide glutathione, and GPx1, the main enzyme

involved in the recovery of reduced glutathione following its oxidation, in glucose-

containing medium. GCL is made up of two subunits. We noted that GCLC protein and

GCLC gene expression levels increased ($F (1,20) = 30.42, p<0.0001$ and $F (1,27) = 16.94,$

$p = 0.0003$, respectively) following 24h of treatment with $(PhSe)_2$ (Fig. 4E-F) as well as

GCLM levels at 6h of treatment ($F (1,30) = 8.759, p = 0.006$) (Fig. 4 G-H). Furthermore,

GPx1 levels were increased at 12 and 24h of treatment with $(PhSe)_2$ in BAEC cultured in

glucose-containing medium ($F (1,14) = 29.11, p < 0.0001$) (Supplementary Fig. 1A).

Similar results were observed in BAEC maintained in a galactose-containing medium.

We observed an increase in the mRNA levels of *GCLC* after 12h (F (1,22) = 14.07, p =

0.0011) and GCLC protein content at 24h (F (1,31) = 9.316, p = 0.0046) (Fig. 4 M-N), as

well as an increase in GCLM protein levels after 6h of treatment with $(PhSe)_2$ (F (3,14)

= 5.9, p = 0.0076) (Fig. 4 O-P). $(PhSe)_2$ also increases the mRNA expression of GPx1 at

12 and 24h of treatment in the galactose-containing medium (F (1,12) = 40.22, p <

0.0001) (Supplementary Fig. 1B).   In contrast, *SOD2* mRNA levels and protein of

MnSOD were unaltered (Fig. 4 I-J), while the Prx3 levels decreased after 24h of treatment

with $(PhSe)_2$ in the galactose-containing medium (Fig. 4 K-L). The mRNA levels of

catalase (*CAT*), mitochondrial uncoupling protein 2 (*UCP2*), cytochrome c (*CYTC*),

glutathione synthetase (*GSS)* and *Nrf2* were unaltered by the treatment with $(PhSe)_2$ of

BAEC maintained in either glucose- or galactose-containing medium (Supplementary

Fig. 1 C-L). Overall, these results suggest that in glucose containing media $(PhSe)_2$

induces both the main mitochondrial and cytosolic antioxidant systems, while in galactose

media $(PhSe)_2$ induces mainly the cytosolic antioxidant systems, a result consistent with

the expected pre-activation of OXPHOS and mitochondrial antioxidant capacity in

galactose media.

Fig. 4 Effect of (PhSe)₂ in time-course expression profiles of antioxidant proteins in glucose- or galactose-containing medium. BAEC were treated with (PhSe)₂ or vehicle (DMSO) for 6, 12, 24 or 48h in glucose or galactose medium. Protein expression by Western Blotting analyses and mRNA expression by qPCR. Results were expressed as protein /β-tubulin ratio and the data were presented as the percentage of treatment with (PhSe)₂ (bars) with respect to vehicle group (dashed line). V = vehicle, D = (PhSe)₂. Data were represented as mean ± SD (n=3-7). *$p < 0.05$, indicate statistical difference from vehicle group by two-way ANOVA, followed by Bonferroni's *post hoc* test.

3.5. Effect of (PhSe)₂ on Nrf2 nuclear translocation of BAEC conditioned in glucose or galactose medium.

Since $(PhSe)_2$ increased the levels of antioxidant genes known to be targeted by the transcription factor Nfr2, we evaluated the effect of $(PhSe)_2$ on Nrf2 nuclear translocation, a thus activation, in BAEC cultured in glucose- or galactose-containing medium. We found that Nrf2 nuclear localization was increased following 3h of incubation with $(PhSe)_2$ in BAEC cultured in both glucose and galactose containing medium ($F_{(4,14)} = 4.737$, $p = 0.0126$ and $F_{(1,30)} = 64.98$, $p < 0.0001$, respectively) (Fig. 5A-B). The increased nuclear localization was gradually reduced overtime in both glucose and galactose cultured cells but was maintained over a more extended period in galactose cultured cells (Fig. 5C-D), suggesting a more effective nuclear retention in galactose. The separate evaluation of nuclear and cytosolic levels of Nrf2 showed a significant increase in cytosolic Nrf2 levels following 3h of treatment with $(PhSe)_2$ in BAEC maintained in glucose- containing medium but not in galactose cultured cells possibly suggesting that global Nrf2 levels stabilized by $(PhSe)_2$ in glucose but not in galactose media (Supplementary Fig. 2). Nrf2 degradation is a tightly regulated process, cytosolic retention of Nrf2 by Keap1 is linked to its proteasomal degradation, and therefore, Keap1 release is generally associated not only to Nrf2 nuclear translocation but also to its stabilization [28]. In summary, Nrf2 activation is detectable in both glucose and galactose media, but in galactose media is likely to be effective for more extended periods of time.

Fig. 5 Effect of (PhSe)₂ on Nrf2 translocation in BAEC conditioned in glucose- or galactose-containing medium. BAEC were treated with (PhSe)₂ (1 μM) or vehicle (DMSO) for 3, 6, 12, 24 and 48h in medium containing glucose (A-B) or galactose (C-D) and the immunofluorescence the Nrf2 were analyzed. (A and C) Representative image of immunofluorescence from Nrf2 (green) and DAPI (blue) in BAEC, the white bars represent 50 μm. (B and D) Ratio of Nrf2 in the nuclei/cytosol. Data were represented as mean ± SD (n=3-4). * $p < 0.05$, indicate statistical difference from vehicle group by two-way ANOVA, followed by Bonferroni's *post hoc* test.

3.6. Effect of (PhSe)₂ on FOXO3 activation in BAEC conditioned in glucose or galactose

medium.

Finally, we also analyzed the possible involvement of FOXO3 transcriptional

factor in $(PhSe)_2$ activation of antioxidant gene expression. Treatment of BAEC in

glucose-containing medium with $(PhSe)_2$ for 6, 12 or 24h increased the level of FOXO3

phosphorylation (pFOXO3) at Thr32 ($F_{(1,90)} = 8.111$, $p = 0.0055$), a mark of Akt

phosphorylation (pAkt), suggesting an inhibition of FOXO3 activity and nuclear

extrusion (Fig. 6A). However, no significant changes in the nuclear/cytosolic ratio of

FOXO3 (Fig 6E) nor in the cytosolic and nuclear levels when analyzed separately

(Supplementary Figure 3), and, although an increase in Akt phosphorylation was

detectable in response to $(PhSe)_2$, it could only be observed at 24h of treatment in glucose

media ($F_{(1,76)} = 12.86$, $p = 0.0006$) (Fig. 6C). In contrast, although the nuclear/cytosolic

ratio of FOXO3 transiently decreased at 3h of treatment in galactose media (Fig. 6F),

$(PhSe)_2$ decreased pFOXO3/FOXO3 ratio at 24h ($F_{(3,57)} = 5.642$, $p = 0.0019$),

suggesting an increase in FOXO3 activity and nuclear localization (Fig. 6B). In these

conditions although the changes in pAkt levels did not reach statistical significance, a

consistent tendency to lower pAkt/Akt ratio was detectable at 24h (Fig. 6D) as well as

detectably higher FOXO3 levels in both the nuclei and the cytosol at 12h of treatment

(Supplementary Figure 3). All together these results suggest that FOXO3 activity is more

likely to play a relevant role in antioxidant gene regulation in galactose than in glucose

cultured cells.

Fig. 6 Effect of (PhSe)$_2$ on pFOXO3 and pAkt levels in BAEC conditioned in glucose-
or galactose-containing medium. BAEC were treated with (PhSe)$_2$ or vehicle (DMSO)
for 3, 6, 12, 24 and 48h in glucose (A, C and E) or galactose (B, D and F) medium to
analyze pFOXO3 (A-B) and pAkt (C-D) protein content by Western blotting. Results are
expressed as protein/β-tubulin or FOXO3 ratio and the data is presented as the relative %
of treatment with (PhSe)$_2$ (bars) *vs* vehicle. V = vehicle, D = (PhSe)$_2$. (E and F)
Immunofluorescence for FOXO3 were analyzed. Ratio of FOXO3 in the nuclei/cytosol.
Data were represented as mean ± SD (n=3-13). * $p < 0.05$, indicate statistical difference
from vehicle group (dashed line) by two-way ANOVA, followed by Bonferroni's *post
hoc* test.

4. Discussion

Endothelial cells are the primary target of circulating redox-active molecules.

Among these, primary bovine aortic endothelial cells (BAEC), since preserving their

original metabolic plasticity, represent a relevant *in vitro* model for investigating the

effects of pharmacological agents across diverse metabolic contexts. The choice of cell

culture media profoundly influences the metabolic performance of cells *in vitro.* Our

recent findings, particularly concerning the modulation of transcription factors Nrf2 and

FOXO3 key regulators of antioxidant gene expression in BAECs, underscore the

significance of cellular metabolic state. Specifically, alterations in cell culture conditions,

leading to heightened reliance on oxidative metabolism, demonstrate a consequential

impact on the efficacy of redox-active agents on vascular cells. These observations

suggest that the effectiveness of redox-active agents on endothelial cells may be

intricately linked to their metabolic milieu. Therefore, we investigate the effects of

$(PhSe)_2$ on BAEC cultured in media containing glucose or galactose, assessing its impact

on metabolism and antioxidant responses.

$(PhSe)_2$ has been extensively studied for its pivotal role in vascular physiology

and pathology, attributed to its capacity for redox modulation. This includes the

promotion of mitochondrial respiration and the upregulation of antioxidant gene

expression, partly mediated by the translocation of Nrf2 to the nucleus [22]. Our findings

reveal significant differences in the action of $(PhSe)_2$ depending on whether BAEC are

cultured in galactose and glucose media. These differences are likely to be relevant to

their vascular effects *in vivo*. In cells cultured in glucose, treatment with $(PhSe)_2$ yielded

a net positive impact on mitochondrial oxidative capacity, assessed by respirometry. This

enhancement correlated with observed changes in mitochondrial cellular dynamics,

characterized by more fused mitochondria and a uniform distribution of mitochondria

throughout the cells, with reduced concentration in the perinuclear region. Conversely,

these changes were not observed in cells cultured with galactose, where the cells were

already compelled to utilize their oxidative capacity. Therefore, the modulation of

mitochondrial function by (PhSe)$_2$ appears to be effective only in cells exhibiting limited

utilization of their mitochondrial oxidative capacity. Regarding (PhSe)$_2$ antioxidant

capacity, we noted that although (PhSe)$_2$ induces the nuclear translocation of Nrf2 in both

glucose and galactose media, Nrf2 remains longer in the nuclei of cells cultured in

galactose media. This observation suggests a potential preference for nuclear retention in

the galactose environment. However, any speculation regarding the underlying

mechanism at this stage would be highly conjectural. Potential explanations could involve

interactions with other transcription factors [29] or the inhibition of its nuclear extrusion

[30].

The enhanced Nrf2 activation profile observed in galactose media is even more

pronounced in the case of FOXO3a, where its activation is solely detectable in galactose

media, with reduced levels of pFOXO3a and increased nuclear protein levels. This shared

pattern suggests a more efficient induction of antioxidant gene expression in galactose

compared to glucose media. However, analysis of the gene expression changes elicited

by (PhSe)$_2$ on crucial antioxidant genes controlling mitochondrial antioxidant capacity

and glutathione synthesis and recycling reveals a stronger induction in glucose media. In

galactose media, the changes induced by (PhSe)$_2$ are not only smaller but also

predominantly focused on glutathione homeostasis, suggesting a more cytosolic, rather

than mitochondrial effect. Therefore, the induction of antioxidants genes by (PhSe)$_2$ in

glucose media likely requires additional transcription factors that are especially active in

conditions of low mitochondrial activity. Potential candidates include NFkB [31] a redox-

sensitive transcription factor controlling antioxidant gene expression, and c-MYC [32]

among others. Additionally, since (PhSe)$_2$ also enhances mitochondrial fission and

maximal oxidative capacity, other pathways may be involved. Acute inhibition of

mitochondrial respiration could drive compensatory mechanisms, such as the activation

of PGC-1α, which can induce both antioxidant gene expression and mitochondrial fusion

[8] while supporting FOXO3 [13] and Nrf2 [14] activation.

Despite the apparent lower capacity of (PhSe)$_2$ to induce antioxidant gene

expression in galactose media, assessments of mitochondrial antioxidant capacity using

the mitochondrial inhibitor DMNQ or the exogenous addition of H_2O_2 suggest a

potentially heightened antioxidant response in galactose-treated cells. This inference

arises from the substantial reduction elicited by (PhSe)$_2$ on mitochondrial superoxide (O_2^-

) levels in galactose-cultured cells following a challenge with H_2O_2. However, drawing

definitive conclusions from these observations is challenging. In galactose media, cells

exhibit greater resistance to DMNQ, while in glucose media, they are highly susceptible

to H_2O_2-induced mitochondrial damage, resulting in under basal levels of mitochondrial

O_2^-. Notably, (PhSe)$_2$ partially rescues these aberrant levels and restores them to normal,

consistent with its observed capacity to enhance mitochondrial oxidative capacity.

Therefore, our data suggest that while (PhSe)$_2$ demonstrates greater efficacy in the

activation of Nrf2 and FOXO3a under pro-oxidative conditions, alternative pathways

contribute to its enhancement of mitochondrial oxidative capacity and total antioxidant

capacity when cells are cultured in glucose media. The observation that (PhSe)$_2$ induces

acute inhibition of mitochondrial respiration shortly after treatment may indicate a

significant role for mitochondrial function in this context. In summary, our findings

underscore the critical influence of the cellular metabolic status on antioxidant capacity

of redox-active molecules such as (PhSe)$_2$.

Declaration of competing interest

The authors declare that they have no known competing financial interests or
personal relationships that could have appeared to influence the work reported in this
paper.

Acknowledgments

This work was supported by MCIN/AEI/10.13039/501100011033/FEDER, UE,
[grant numbers RTI2018-093864-B-I00 and PID2021-122765OB-I00]; the European
Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-
Curie [grant agreement 721236-TREATMENT]; EU Horizon Europe Program under the
EIC Pathfinder DiBaN [GA-101162517]; the Brazilian institutions: Conselho Nacional
de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de
Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Fundação de Apoio à Pesquisa
do Distrito Federal (FAPDF) [grants 00193-00000884/2021-89 and 00193-
00002348/2022-07]; and Instituto Nacional de Ciência e Tecnologia e Neuro-
ImunoModulação (INCT-NIM) [grant 485489/2014-1]. The authors are grateful to Smart
Servier Biology Medicine for the available figures.

References

[1] Z. Abbaszadeh, S. Çeşmeli, Ç. Biray Avcı, Crucial players in glycolysis: Cancer
progress, Gene, 726 (2020).
[2] Y. Liu, K. Birsoy, Metabolic sensing and control in mitochondria, Mol Cell, 83
(2023) 877–889.
[3] E.A. Zaal, C.R. Berkers, The influence of metabolism on drug response in
cancer, Front Oncol, 8 (2018).
[4] M.J.C. de Kok, A.F. Schaapherder, R.C.I. Wüst, M. Zuiderwijk, J.A. Bakker,
J.H.N. Lindeman, S.E. Le Dévédec, Circumventing the Crabtree effect in cell
culture: A systematic review, Mitochondrion, 59 (2021) 83–95.
[5] L.D. Marroquin, J. Hynes, J.A. Dykens, J.D. Jamieson, Y. Will, Circumventing
the crabtree effect: Replacing media glucose with galactose increases
susceptibility of hepG2 cells to mitochondrial toxicants, Toxicological Sciences,
97 (2007) 539–547.
[6] V.M. Gohil, S.A. Sheth, R. Nilsson, A.P. Wojtovich, J.H. Lee, F. Perocchi, W.
Chen, C.B. Clish, C. Ayata, P.S. Brookes, V.K. Mootha, Nutrient-sensitized
screening for drugs that shift energy metabolism from mitochondrial respiration
to glycolysis, Nat Biotechnol, 28 (2010) 249–255.

[7]   M. V. Golikov, V.T. Valuev-Elliston, O.A. Smirnova, A. V. Ivanov, Physiological Media in Studies of Cell Metabolism, Mol Biol, 56 (2022) 629–637.

[8]   I. Valle, A. Álvarez-Barrientos, E. Arza, S. Lamas, M. Monsalve, PGC-1α regulates the mitochondrial antioxidant defense system in vascular endothelial cells, Cardiovasc Res, 66 (2005) 562–573.

[9]   V. Ngo, M.L. Duennwald, Nrf2 and Oxidative Stress: A General Overview of Mechanisms and Implications in Human Disease, Antioxidants, 11 (2022).

[10] K.A. Lasick, E. Jose, A.M. Samayoa, L. Shanks, K.W. Pond, C.A. Thorne, A.L. Paek, FOXO nuclear shuttling dynamics are stimulus-dependent and correspond with cell fate, Mol Biol Cell, 34 (2023).

[11] J.M. Deblasi, G.M. Denicola, Dissecting the crosstalk between nrf2 signaling and metabolic processes in cancer, Cancers (Basel), 12 (2020) 1–20.

[12] M. Monsalve, Y. Olmos, The Complex Biology of FOXO, 2011.

[13] I. Valle, Y. Olmos, S. Borniquel, A. Tierrez, E. Soria, S. Lamas, M. Monsalve, Mutual dependence of Foxo3a and PGC-1α in the induction of Oxidative stress genes, Journal of Biological Chemistry, 284 (2005) 14476–14484.

[14] E. Navarro, L. Gonzalez-Lafuente, I. Pérez-Liébana, I. Buendia, E. López-Bernardo, C. Sánchez-Ramos, I. Prieto, A. Cuadrado, J. Satrustegui, S. Cadenas, M. Monsalve, M.G. López, Heme-Oxygenase i and PCG-1α Regulate Mitochondrial Biogenesis via Microglial Activation of Alpha7 Nicotinic Acetylcholine Receptors Using PNU282987, Antioxid Redox Signal, 27 (2017) 93–105.

[15] L.S. Galant, L. Doblado, R. Radi, A.F. de Bem, M. Monsalve, Culture of Bovine Aortic Endothelial Cells in Galactose Media Enhances Mitochondrial Plasticity and Changes Redox Sensing, Altering Nrf2 and FOXO3 Levels, Antioxidants, 13 (2024).

[16] B. Fiuza, N. Subelzú, P. Calcerrada, M.R. Straliotto, L. Piacenza, A. Cassina, J.B.T. Rocha, R. Radi, A.F. De Bem, G. Peluffo, Impact of SIN-1-derived peroxynitrite flux on endothelial cell redox homeostasis and bioenergetics: Protective role of diphenyl diselenide via induction of peroxiredoxins, Free Radic Res, 49 (2015) 122–132.

[17] A.F. De Bem, B. Fiuza, P. Calcerrada, P.M. Brito, G. Peluffo, T.C.P. Dinis, M. Trujillo, J.B.T. Rocha, R. Radi, L.M. Almeida, Protective effect of diphenyl diselenide against peroxynitrite-mediated endothelial cell death: A comparison with ebselen, Nitric Oxide, 31 (2013) 20–30.

[18] M.A. Hort, M.R. Straliotto, P.M. Netto, J. Batista Teixeira Da Rocha, A. Fabro De Bem, R.M. Ribeiro-Do-Valle, Diphenyl Diselenide Effectively Reduces Atherosclerotic Lesions in LDLr 2/2 Mice by Attenuation of Oxidative Stress and Inflammation, n.d.

[19] M.R. Straliotto, M.A. Hort, B. Fiuza, J.B.T. Rocha, M. Farina, G. Chiabrando, A.F. De Bem, Diphenyl diselenide modulates oxLDL-induced cytotoxicity in macrophage by improving the redox signaling, Biochimie, 95 (2013) 1544–1551.

[20] A.F. de Bem, M. Farina, R. de L. Portella, C.W. Nogueira, T.C.P. Dinis, J.A.N. Laranjinha, L.M. Almeida, J.B.T. Rocha, Diphenyl diselenide, a simple glutathione peroxidase mimetic, inhibits human LDL oxidation in vitro, Atherosclerosis, 201 (2008) 92–100.

[21] S.G. Müller, N.S. Jardim, C.B. Quines, C.W. Nogueira, Diphenyl diselenide regulates Nrf2/Keap-1 signaling pathway and counteracts hepatic oxidative stress induced by bisphenol A in male mice, Environ Res, 164 (2018) 280–287.

[22] R.L. Quispe, M.L. Jaramillo, L.S. Galant, D. Engel, A.L. Dafre, J.B. Teixeira da Rocha, R. Radi, M. Farina, A.F. de Bem, Diphenyl diselenide protects neuronal cells against oxidative stress and mitochondrial dysfunction: Involvement of the glutathione-dependent antioxidant system, Redox Biol, 20 (2019) 118–129.

[23] D. Bartolini, J. Commodi, M. Piroddi, L. Incipini, L. Sancineto, C. Santi, F. Galli, Glutathione S-transferase pi expression regulates the Nrf2-dependent response to hormetic diselenides, Free Radic Biol Med, 88 (2015) 466–480.

[24] M. Monsalve, S. Borniquel, I. Valle, S. Lamas, Mitochondrial dysfunction in human pathologies, 2007.

[25] G. Peluffo, P. Calcerrada, L. Piacenza, N. Pizzano, R. Radi, Superoxide-mediated inactivation of nitric oxide and peroxynitrite formation by tobacco smoke in vascular endothelium: studies in cultured cells and smokers, Am J Physiol Heart Circ Physiol, 296 (2009) 1781–1792.

[26] S. Borniquel, N. García-Quintáns, I. Valle, Y. Olmos, B. Wild, F. Martínez-Granero, E. Soria, S. Lamas, M. Monsalve, Inactivation of Foxo3a and Subsequent Downregulation of PGC-1α Mediate Nitric Oxide-Induced Endothelial Cell Migration, Mol Cell Biol, 30 (2010) 4035–4044.

[27] J. Bernal-Tirapo, M.T. Bayo Jiménez, P. Yuste-García, I. Cordova, A. Peñas, F.J. García-Borda, C. Quintela, I. Prieto, C. Sánchez-Ramos, E. Ferrero-Herrero, M. Monsalve, Evaluation of Mitochondrial Function in Blood Samples Shows Distinct Patterns in Subjects with Thyroid Carcinoma from Those with Hyperplasia, Int J Mol Sci, 24 (2023).

[28] F. He, X. Ru, T. Wen, NRF2, a transcription factor for stress response and beyond, Int J Mol Sci, 21 (2020) 1–23.

[29] W. Li, S. Yu, T. Liu, J.H. Kim, V. Blank, H. Li, A.N.T. Kong, Heterodimerization with small Maf proteins enhances nuclear retention of Nrf2 via masking the NESzip motif, Biochim Biophys Acta Mol Cell Res, 1783 (2008) 1847–1856.

[30] J.W. Kaspar, A.K. Jaiswal, Tyrosine phosphorylation controls nuclear export of Fyn, allowing Nrf2 activation of cytoprotective gene expression, The FASEB Journal, 25 (2011) 1076–1087.

[31] M.J. Morgan, Z.G. Liu, Crosstalk of reactive oxygen species and NF-κB signaling, Cell Res, 21 (2011) 103–115.

[32] D.R. Wonsey, K.I. Zeller, C. V Dang, The c-Myc target gene PRDX3 is required for mitochondrial homeostasis and neoplastic transformation, n.d.