Risk factors, brain structure, and neurotransmitter systems: a large-scale multivariate study

Short title: Risk factors, brain structure, and neurotransmitter systems

Eliana Nicolaisen-Sobesky[1,2]*, Somayeh Maleki Balajoo[1,2], Mostafa Mahdipour[1],
Agoston Mihalik[3,4,5], Mahnaz Olfati[6], Felix Hoffstaedter[1,2], Janaina Mourao-Miranda[3,4],
Masoud Tahmasian[1,2,7], Simon B. Eickhoff[1,2], Sarah Genon[1,2]*

[1]Institute of Neuroscience and Medicine (INM-7: Brain and Behaviour), Research Centre Jülich, Jülich, Germany.
[2]Institute of Systems Neuroscience, Heinrich Heine University Düsseldorf, Düsseldorf, Germany.
[3]Centre for Medical Image Computing, Department of Computer Science, University College London, London, United Kingdom.
[4]Max Planck University College London Centre for Computational Psychiatry and Ageing Research, University College London, London, United Kingdom.
[5]Department of Psychiatry, University of Cambridge, Cambridge, United Kingdom.
[6]Institute of Medical Science and Technology, Shahid Beheshti University, Tehran, Iran.
[7]Department of Nuclear Medicine, University Hospital and Medical Faculty, University of Cologne, Cologne, Germany.

*Corresponding authors:

Eliana Nicolaisen-Sobesky
Institute of Neuroscience and Medicine (INM-7: Brain and Behaviour), Research Centre Jülich,
Jülich, 52428, Germany
elinicolaisen@gmail.com

Sarah Genon
Institute of Neuroscience and Medicine (INM-7: Brain and Behaviour), Research Centre Jülich,
Jülich, 52428, Germany
s.genon@fz-juelich.de

Summary

The link between brain health and risk/protective factors for non-communicable diseases (such as high blood pressure, high body mass index, diet, smoking, physical activity, etc.) is increasingly acknowledged. However, the specific effects that these factors have on brain health are still poorly understood, delaying their implementation in precision brain health. Here, we studied the multivariate relationships between risk factors for non-communicable diseases and brain structure, including cortical thickness (CT) and grey matter volume (GMV). Furthermore, we adopted a systems-level perspective to understand such relationships, by characterizing the cortical patterns (yielded in association to risk factors) with regards to brain morphological and functional features, as well as with neurotransmitter systems. Similarly, we related the pattern of risk/protective factors dimensions with a peripheral marker of inflammation.

First, we identified latent dimensions linking a broad set of risk factors for non-communicable diseases to parcel-wise CT and GMV across the whole cortex. Data was obtained from the UK Biobank (n=7370, age range=46-81 years). We used regularized canonical correlation analysis (RCCA) embedded in a machine learning framework. This approach allows us to capture inter-individual variability in a multivariate association and to assess the generalizability of the model. The brain patterns (captured in association with risk/protective factors) were characterized from a multi-level perspective, by performing correlations (spin tests) between them and different brain patterns of structure, function, and neurotransmitter systems. The association between the risk/protective factors pattern and C-reactive protein (CRP, a marker of inflammation) was examined using Spearman correlation.

We found two significant and partly replicable latent dimensions. One latent dimension linked cardiometabolic health to brain patterns of CT and GMV and was consistent across sexes. The

other latent dimension linked physical robustness (including non-fat mass and strength) to

patterns of CT and GMV, with the association to GMV being consistent across sexes and the

association to CT appearing only in men. The CT and GMV patterns of both latent dimensions

were associated to the binding potentials of several neurotransmitter systems. Finally, the

cardiometabolic health dimension was correlated to CRP, while physical robustness was only

very weakly associated to it.

We observed robust, multi-level and multivariate links between both cardiometabolic health

and physical robustness with respect to CT, GMV, and neurotransmitter systems. Interestingly,

we found that cardiometabolic health and physical robustness are associated with not only

increases in CT or GMV, but also with decreases of CT or GMV in some brain regions. Our

results also suggested a role for low-grade chronic inflammation in the association between

cardiometabolic health and brain structural health. These findings support the relevance of

adopting a holistic perspective in health, by integrating neurocognitive and physical health.

Moreover, our findings contribute to the challenge to the classical conceptualization of

neuropsychiatric and physical illnesses as categorical entities. In this perspective, future studies

should further examine the effects of risk/protective factors on different brain regions in order

to deepen our understanding of the clinical significance of such increased and decreased CT

and GMV.

Keywords: Brain   structure,   physical   robustness,   cardiometabolic,   risk   factors,

neurotransmitter systems

Introduction

Non-communicable diseases, including cardiovascular, metabolic, mental, and neurological disorders, represent the predominant global public health challenge nowadays (1,2). Most non-communicable diseases share a common set of risk/protective factors (3–5) (here referred simply as risk factors), like tobacco smoking, unhealthy diet, physical inactivity (1,2), excessive alcohol consumption, hypertension (2), sleep problems (6), obesity (increased body mass index (BMI)) and air pollution (2,6). Since non-communicable diseases include mental and neurological disorders, the same set of risk factors also affects brain health (2,7).

Two of the biomarkers for brain health are cortical thickness (CT) and Grey Matter Volume (GMV). Several studies have analyzed the link between these structural markers and specific risk factors such as BMI (8–14), waist circumference (13,14), cigarette smoking (13,15,16), physical exercise (17), and diet (18). Even though these studies have contributed to our understanding of the link between risk factors and brain structure they have some caveats. One limitation is that the reported findings have been inconsistent across studies. For instance, associations between BMI and both, global or frontal CT, have been reported as negative (8–10,13,14), positive (11–13), or not significant (9,10). To address this point, studies using robust approaches that evaluate the generalizability of results are needed. In this regard, machine learning approaches use cross-validation to test for the generalizability of the implemented models.

A second limitation of studies linking brain structure to risk factors is that they mostly used univariate/bivariate approaches. In other words, so far studies have mainly linked specific risk factors with global measures of the brain, or with specific brain regions. (19). However, these factors usually do not happen in isolation (e.g. BMI, diet and physical activity are strongly interrelated in population data) and hence represent a multivariate set. Accordingly, previous

studies only provide a partial view of this association (15,16,19,20) impeding the discovery of

distributed brain networks that are associated to a range of risk factors simultaneously (21).

Identifying distributed brain networks and their interactions, as well as their relationships with

several risk factors requires "'doubly" multivariate approaches. These approaches can take full

advantage of the rich phenotyping included in large-scale datasets such as the UK Biobank

(UKB) and embrace collinearity among risk factors variables, as well as among brain regions.

In this respect, canonical correlation analysis (CCA) is a multivariate, data-driven approach

that can be used to discover large-scale distributed cortical patterns associated with several risk

factors (22,23). Of note, a regularized version of CCA (RCCA) mitigates the effect of

collinearity and yields more stable results than CCA (22), improving the interpretability of

results. CCA and RCCA have been previously used to search for robust and generalizable

multivariate associations between different data modalities, such as brain structure, brain

function, hippocampal structure, behavior, environmental variables, and psychiatric features

(20,23–28). Using these methods, numerous studies have shown that several healthy and

illness-related phenotypes are linked to axes of brain structural organization (23,24,26).

Therefore, despite the link between risk factors for non-communicable diseases and brain

health being acknowledged, there are still several important aspects of this association which

are incompletely understood (21,29). The lack of understanding of the relationship between

risk factors and brain health prevents their use as biomarkers in the clinical practice and hence

is a major research priority (21). Since structural brain changes may be long-lasting and lead

to various neuropsychiatric diseases, it is critical to pinpoint modifiable risk factors that can

reduce the risk of those conditions. Understanding how risk factors for non-communicable

diseases are related to brain health from a comprehensive and broad perspective requires

understanding how multiple risk factors simultaneously relate to structural patterns across the

whole brain. Thus, the main question of this study is how several risk factors for non-communicable diseases are associated with patterns of CT and GMV.

Another important point to consider is the multi-level nature of the interplay between risk factors and brain health. For instance, risk factors for non-communicable diseases have been associated with several brain features, including brain structure (30,31), function (32), genetics, and neurotransmitter systems (33). This needed integration of different neurobiological features (34) can now be done quantitively with neuromaps (35), which provides access to a wide set of brain maps, including for instance genetic transcription and brain molecular features. Similarly, a role for immunoinflammatory processes in the association between risk factors and brain health has been frequently discussed (36–40). Linking CCA-derived cortical pattern and risk factors pattern to known neurobiological patterns and inflammation will allow to gain a systems-level understanding of the association between risk factors and brain health.

Hence, to gain a comprehensive, generalizable, and multi-level understanding of the relationship between risk factors and brain health, it is needed to use robust machine learning approaches with generalizability testing, along with methods that integrate different data modalities (24,41). With that aim, we first searched for latent dimensions linking a wide range of risk factors for non-communicable diseases to region-wise CT and GMV across the whole brain cortex using RCCA embedded in a machine learning framework (Figure 1). In other words, we searched for those combinations of risk factors which are most relevant for CT and GMV interindividual variability (20). On a second step we aimed to characterize the captured brain patterns from a neurobiological perspective comparing them with existing brain maps spanning brain structure, function, genetic variability, and neurotransmitter systems. Finally, we analyzed the link between the risk factors dimension and a peripheral marker of inflammation: C-reactive protein (CRP).

153 **Figure 1. Overview of the analyses.** Top panel: Latent dimensions were search using RCCA,
154 linking cortical thickness and grey matter volume measures (parcel-wise across the whole brain
155 cortex) with a wide set of risk factors for non-communicable diseases. Below panel: In order
156 to interpret the latent dimension, the brain loadings were compared with several brain features.
157 Created in BioRender. Nicolaisen, E. (2025) https://BioRender.com/22pyru2

Methods

Participants

We used data from UKB (application 41655) (42,43). Inclusion criteria for this study were

having no self-reported illnesses (Data-Field ID 20002-2.*) and having complete data in all

variables utilized in this study (excluding responses like "Do not know" or "Prefer not to

answer"). To ensure that our sample represent male and female population equally, we selected

a sample (n=7370) with equal number of age-matched males and females. We hereafter refer

to it as the main or mixed sample, since it includes individuals of both sexes/genders: 3685

women (age: range 46-81 years, mean 62.3, standard deviation 7.6) and 3685 men (age: range

46-81 years, mean 62.3, standard deviation 7.6). In order to check for a potential sex-bias in

the results, we also analyzed each sex-specific subsample, separately (see Supplementary

Methods 1.1).

Risk factors data

We identified 68 risk factor variables in the UKB dataset (after removing variables which were

duplicated (i.e. same variable measured several times), variables with missing values, and

variables with skewed distribution). These 68 variables spanned categories of general health,

body size measures, diet, physical activity, residential air pollution, sleep health, alcohol

consumption, and smoking (Table S1). As done previously, we computed waist-to-hip ratio

dividing waist circumference by hip circumference (13). All variables corresponding to risk

factors were acquired on the Imaging visit, with the exception of those in the Category of

"Residential air pollution" (Table S1), which were acquired before the Imaging visit.

In order to first gain a better understanding of the interrelation between risk factors alone

(without taking into account brain data), the intercorrelation among risk factors was analyzed

with Pearson's correlation and visualized in correlation matrices.

Neuroimaging data

Neuroimaging data for the sample used in this study were collected by UKB in four sites using

identical protocols and 3T Siemens Skyra scanners with standard Siemens 32-channel receive

head coils (44). T1-weighted structural imaging (3D-MPRAGE, sagittal) was acquired with the

following parameters (42,44): voxel resolution: 1x1x1 mm, FoV: 208x256x256 matrix, TI/TR

= 880/2000 ms, in-plane acceleration iPAT=2. T1 images were processed by UKB using a

custom pipeline based on FSL (44), including gradient distortion correction, cutting down the

field of view, registration (linear and then non-linear) to the MNI152 standard-space T1

template, brain extraction, defacing, and brain segmentation (44). CT and GMV were estimated

using FreeSurfer a2009s.

We used mean raw CT and mean raw GMV of 148 cortical parcels (Destrieux atlas, a2009s)

(45). To check the effect of brain size correction on the results, we also used proportional

estimations of CT and GMV (computed subject-wise, dividing the mean raw CT or GMV of

each parcel by the mean raw CT or Total Intracranial Volume (TIV), respectively, across all

parcels) and CT and GMV corrected for brain size (regressing out mean CT or TIV,

respectively, in a cross-validation consistent manner to avoid data leakage) (46). These three

different brain structural measures (raw, proportional, or corrected) represent different

biological properties and can show different patterns of associations with risk factors. For

instance, raw CT represents the absolute CT of a brain region, while proportional CT can be

understood as the ratio between a region's CT and the whole brain's CT. Along the same line,

corrected CT can be seen as the regional value adjusted for the whole brain value, that is, the

regional value which is not explained by the total brain value. Accordingly, proportional and

corrected values are relative values, i.e. when taking into account the whole brain value. This

means that a corrected CT value for a given region expresses to which extend it has high or

low CT when taking into account the global brain structure. Accordingly, corrected values are

more likely to reveal structural patterns that pertain to specific brain regions. However, the

drawback of correction or adjustment for head/brain size is that it may discard relevant

variance, i.e. variance that pertains to the effect we are interested in. For example, if variability

in brain/head size is associated with variability in physical robustness for relevant biological

reasons (e.g. physically more robust individuals tend to be taller and more massive), adjusting

for brain/head size can remove relevant variance related to physical robustness. Therefore, row

and uncorrected/unadjusted values have pros and cons and offer different insights in the current

study.

Regularized canonical correlation analysis

Canonical Correlation Analysis (CCA) is a multivariate technique that can discover latent

dimensions linking interindividual variability in X and Y (22,23,47). Here, X included either

CT or GMV for each parcel, while Y included risk factors. CCA searches linear combinations

of variables in X (brain weights u) and of variables in Y (risk factor weights v), which

maximize the canonical correlation between the brain scores and risk factor scores (22,23). The

scores correspond to the projection of X and Y onto their respective weights (Xu and Yv) (these

correspond to subject-wise values). One limitation of CCA is that it can overfit the data or yield

unstable results, especially in high-dimensional datasets (48). Therefore, we here used a

regularized version of CCA (RCCA) which reduces the overfitting of the model by adding L2-

norm constraints to the weights (22,23,49,50).

To search for multivariate associations between risk factors and both, CT and GMV in the main

sample, we ran two RCCA models (one per brain structural measure). To check for the effect

of brain size correction on the results, we ran four more analyses for cortical models: using

either proportional or brain-size-corrected CT and GMV. In addition, to investigate potential

sex-bias in the results, we ran 6 additional RCCA models: three in the subsample of women

and three in the age-matched subsample of men (see Supplementary Materials 1.1). Age and

site effects were regressed out avoiding data leakage in the machine learning framework

(regression parameters were estimated in the training set and applied to the training, test, and

holdout sets) (46,51). In the main sample, sex was also regressed out. Confounding variables

were included in the Y matrix to check if their variance was properly removed.

To visualize and interpret the latent dimensions, loadings were computed (22) (these

correspond to variable-wise values). Brain loadings correspond to the correlation of the original

brain variables (X) with brain scores (Xu). Similarly, risk factor loadings correspond to the

correlation between the risk factors original variables (Y) with the respective scores (Yv).

Loadings indicate which variables are more strongly associated with the latent dimension. To

interpret the latent dimensions, only stable loadings were considered (loadings whose error bar

did not cross zero).

Machine learning framework

We utilized a machine learning framework that uses multiple holdouts of the data (23,52). This

framework implements two consecutive splits: the outer split divides the whole data into

optimization and hold-out sets, and is used for statistical evaluation, and the inner split divides

the optimization set into training and test sets and is used for model selection. In this study, we

used 5 inner splits and 5 outer splits. Model selection was performed based on the highest test

canonical correlation and the highest stability (similarity of weights estimated using Pearson's

correlation across the 5 inner splits).

Statistical evaluation of the latent dimensions

Statistical significance of the latent dimensions was tested with permutation tests. In each of

the 1000 iterations, the rows of the Y matrix were shuffled in the optimization and hold-out

sets. The RCCA model (hyperparameters) that was previously selected with the original data

was now fitted on the permuted optimization set, and weights were obtained. Then, the

permuted hold-out set was projected onto these weights. A canonical correlation under the null

model was hence obtained, and a p-value was computed. The permutation test was repeated for

each one of the outer splits, hence yielding 5 p-values which were corrected by multiple

comparisons (Bonferroni method over 5 comparisons). The statistical significance of the latent

dimensions was evaluated using the omnibus hypothesis (52). Here, the null hypothesis states

that there is no effect in any of the splits. Hence, if at least one split yields a p-value below

0.05, the null hypothesis is rejected, and the latent dimension is considered significant. When

a significant latent dimension was found, its variance was removed from the data using

deflation (23), and an additional latent dimension was sought.

Stability of the latent dimensions across sexes and brain structural measures

The latent dimensions were compared based on their average risk factor loadings and average

brain loadings (average across the outer 5 splits). The risk factor loadings were compared with

Pearson's correlation across models. The brain loadings were compared using spin test with

10000 permutations to account for spatial dependencies of brain data (53) using neuromaps

software (35). The p-values were corrected by multiple comparisons using Bonferroni method.

Neurobiological characterization of the brain structural patterns

The brain maps provided in neuromaps (35) (Table S3) (excluding the map 'hill2010' which

is provided only in one hemisphere) were compared to the maps of CT or GMV loadings using

spin test (53) with 10000 permutations. This was performed to assess if the latent dimensions

captured a CT or GMV pattern significantly associated with other patterns of brain features.

Multiple comparisons were corrected using the Bonferroni method.

Association of the latent dimensions with demographics

The association of the risk factor scores (subject-wise values) and demographics was analyzed

with Spearman correlation (Supplementary materials 1.2, Table S2).

Association of the latent dimensions with peripheral inflammation

We explore the association between the composite variable of risk factors (i.e. the risk factor

scores which are subject-wise values) and a peripheral marker of inflammation with Spearman

correlation. We focused on C-reactive Protein (CRP, field-id 30710-1.0 in the UKB dataset) as

the most relevant marker for chronic and systemic low-grade inflammation having a good

signal-to-noise ratio for population studies, a relative stability in time, and a relatively low

missingness in UK Biobank. Multiple comparisons were corrected with Bonferroni.

Latent dimensions linking risk factors to subcortical and cerebellar volumes

Although the current study focuses on cortical structure, we also added supplementary analyses

examining latent dimensions yielded with subcortical and cerebellar structures for readers'

information. For that, we ran one additional RCCA analysis linking the same set of risk factors

as before with a set of subcortical and cerebellar volumes in the main/mixed sample (see

Supplementary methods 1.3). Age, sex and site were regressed out.

Ethics

Analyses on the data have been approved by the University Hospital Düsseldorf ethics

committee votes 2018-317-RetroDEuA, 2018-317_1-RetroDEuA, and 2018-317_2.

Results

Risk factors collinearity

The distribution of risk factors is shown in Figures S1-S2. In the correlation matrices for risk

factors, two groups of highly intercorrelated variables were evident (Figures S3-S4). One group

shows intercorrelation among body composition measures, including BMI, body fat

percentage, body fat mass, body fat-free mass, body water mass, basal metabolic rate (which

is estimated from fat-free body mass), impedance of whole body, waist circumference, hip

circumference, and waist-to-hip ratio. Another group of highly intercorrelated variables was

characterized by air pollution.

Latent dimension linking cardiometabolic health to brain structure

A first significant latent dimension was found when analyzing the link between risk factors to

raw CT in the main/mixed sample (merging women and men) (r_{range} = 0.30-0.34, p_{range} = 0.005-

0.005) (Figure 2, Figures S5-S6, Table S4). Conceptually, this latent dimension captures

variability in cardiometabolic health, for instance with positive loadings for higher physical

activity and a negative pole for factors of metabolic conditions, such as body fat, BMI and

waist-to-hip ratio, as well as blood pressure. Of note, when analyzing the link between the same

set of risk factors and GMV in the main/mixed sample a latent dimension with a highly similar

profile of risk factors loadings was yielded (Figure 2, Figures S5 and S7, Table S7; Figure S8

for comparison across latent dimensions). Note that this latent dimension corresponded to the

first yielded latent dimension for all models except for the models for raw GMV, in which it

was yielded as second latent dimension.

Figure 2. Loadings of the latent dimension of cardiometabolic health. Risk factor loadings and brain loadings in the main/mixed sample for cortical thickness (left) and grey matter volume (right). The arrows depict the Spearman correlation of the loadings across models. The loadings represent the average over the five outer splits. Error bars depict one standard deviation. The shadowed zone marks loadings between −0.2 and 0.2.

The pattern of cardiometabolic health was linked to a specific brain pattern of CT and a specific

brain pattern of GMV (Figure 2). The CT pattern shows that better cardiometabolic health is

associated with higher CT in the insula, cingulate cortex, temporal lobe, inferior parietal,

orbitofrontal, and primary motor cortex, and to lower CT in the primary somatosensory cortex,

superior frontal areas, superior parietal areas, and occipital areas. In addition, better

cardiometabolic health is associated to higher GMV in primary motor cortex and temporal

cortex, and lower GMV in frontal, parietal, and occipital areas.

Notably, this latent dimension of cardiometabolic health was stable across sexes and across

brain size corrections (Figure S8, Supplementary results 2.1-2.3). Associations between the

latent dimensions and demographics are shown Supplementary Results 2.4.

Thus, overall, this latent dimension appears to capture a composite aspect of cardiometabolic

health that robustly relates to brain structure.

Latent dimension linking physical robustness to brain structure

Our results showed another significant latent dimension linking risk factors to raw CT (r_{range} =

0.09-0.12, p_{range} = 0.005-0.005) (Figure 3, Figures S25-S26, Table S4). Importantly, when

studying raw CT, this latent dimension was significant and stable in the sample of men but not

in the sample of women (see Supplementary results 2.5). For this reason, in the next steps we

focus on the sample of men when considering this latent dimension with respect to CT. Also,

note that this latent dimension corresponded to the second yielded latent dimension for all

models except for the models for raw GMV, in which it was yielded as first latent dimension.

This latent dimension captured variability in physical robustness (as opposed to physical

frailty), with a positive pole associated to for instance whole body fat-free mass, grip strength

and overall health rating, and a negative pole associated with smoking and air pollution.

Figure 3. Loadings of the latent dimension of physical robustness. Risk factor loadings and
brain loadings in the men sample for cortical thickness (left) and the main/mixed sample for
and grey matter volume (right). The arrows depict the Spearman correlation of the loadings
across models. The loadings represent the average over the five outer splits. Error bars depict
one standard deviation. The shadowed zone marks loadings between −0.2 and 0.2.

This latent dimension of physical robustness was associated with a specific brain pattern of CT

and a specific brain pattern of GMV (Figure 3), though the statistical comparison showed a

marginally significant Spearman correlation indicating that they do share variance (r = 0.41, p

= 0.046). However, this significant comparison was yielded only for the case of raw CT and

raw GMV, but not for corrected brain structural measures (Figure S31-f). Conceptually, our

results indicated that higher physical robustness was associated with higher CT, especially in

superior frontal areas and insula. Moreover, the physical robustness pattern was associated to

higher GMV, especially in the anterior cingulate cortex, medial superior frontal areas,

orbitofrontal cortex, and temporal lobe.

This latent dimension was stable across sexes when yielded with GMV, but it was not

significant/stable in the sample of women when yielded with CT (Supplementary results 2.5-

2.7). When yielded with GMV, and when yielded in the sample of men with CT, this latent

dimension was stable across brain size correction procedures (Figure S31, Supplementary

results 2.5-2.7). Associations between the latent dimensions and demographics are shown

Supplementary Results 2.8.

Neurobiological characterization of the pattern of brain structural loadings associated with

cardiometabolic health

To characterize the latent dimensions from a neurobiological perspective, we compared the CT

and GMV loadings with brain maps spanning brain function, structure, and neurotransmitter

systems.

For the latent dimension associated to cardiometabolic health, several associations between the

CT and GMV patterns and different brain maps were significant (Table S10, Figure 4, Figure

S41). Namely, the map of CT loadings was positively associated with the cortical distribution

of serotonin receptor 5-HT1a (54,55), dopamine transporter DAT (56), receptor GABAa (57)

and acetylcholine transporter VAChT (34,58). In addition, the CT pattern was positively

associated with cortical thickness (59), and fifth gradient of resting-state functional

connectivity (RSFC) (60), as well as negatively associated with glucose metabolism (61),

histone deacetylase (62) and with the first principal component of genes in the Allen Human

Brain Atlas (63,64). In turn, the GMV pattern of the cardiometabolic health latent dimension

was positively associated with the cortical distribution of dopamine receptor D2 (65) and

dopamine transporter DAT (66), and negatively associated with serotonin receptor 5-HT1b

(34,67), and histamine receptor H3 (68). Moreover, the GMV pattern was positively associated

with the 8th gradient of RSFC (60), and negatively associated with histone deacetylase (62) and

10th gradient of RSFC (60).

Figure 4. Association of brain structural loadings with neuromaps for the latent dimension of cardiometabolic health. Only data for neuromaps that yielded a significant association with at least one of the shown loadings maps are depicted. Colorbar represents the Spearman correlation. White tiles represent non-significant associations.

Conceptually, this indicates that the latent dimension captured a brain axis in which cortical regions whose CT covaried positively the most (increased the most, red regions in Figure 2) with the pattern of cardiometabolic health were regions that showed the highest binding

potentials for 5-HT1a, DAT, GABAa and VAChT. Similarly, the brain regions whose GMV

increased the most (red regions) with cardiometabolic health were regions with the highest

binding potential for D2 and DAT. In turn, the brain regions whose GMV covaried negatively

the most (decreased the most, blue regions) with cardiometabolic health were regions with the

highest binding potentials for 5-HT1b and H3.

Neurobiological characterization of the pattern of brain structural loadings associated with

physical robustness

The brain patterns yielded for the latent dimension of physical robustness also showed several

significant associations with brain maps (Table S11, Figure 5, Figure S42). The CT pattern

yielded in the sample of men was positively associated with glutamate receptor mGluR5

(34,69), RSFC intersubject variability (70), and developmental area scaling (71). The GMV

pattern was positively associated with the serotonin receptors 5-HT1a (54), with glutamate

receptors mGluR5 (34,69,72) and NMDA (34,73–75), with histamine receptor H3 (68), with

opioid receptors MOR (76,77) and KOR (78), with cannabinoid receptor CB1 (79,80), and with

receptor GABAa (57). In addition, the GMV pattern was positively associated with the 1st

gradient of RSFC (60), intersubject variability of RSFC (70), cortical thickness (59),

developmental area scaling (71), evolutionary expansion (81) and sensory-association axis

(82). Moreover, the GMV pattern was negatively associated with T1w/T2w (59) and with

functional homology (81).

Figure 5. Association of brain structural loadings with neuromaps for the latent dimension of physical robustness. Only data for neuromaps that yielded a significant association with at least one of the shown loadings maps are depicted. Colorbar represents the Spearman correlation. White tiles represent non-significant associations.

Conceptually, this indicates that the latent dimension captured a brain axis in which cortical

regions whose CT increased the most with the pattern of physical robustness (red regions in

Figure 3) were those with highest binding potential for mGluR5. In addition, regions whose

GMV increased the most with physical robustness (red regions) were those with highest

binding potential for 5-HT1a, mGluR5, NMDA, H3, MOR, KOR, CB1 and GABAa.

Correlation of the latent dimensions with peripheral marker of inflammation

CRP was significantly associated to the risk factors scores of cardiometabolic health when

yielded in association with raw CT ($r = -0.39$, $p < 0.001$) and with raw GMV ($r = -0.39$, $p <$

0.001) in the main/mixed sample. It was also significantly associated to the risk factors pattern

of physical robustness yielded in association with raw GMV in the main/mixed sample ($r = -$

0.08, $p = 0.04$) but with a very low effect size. CRP was not significantly associated with the

risk factors yielded in association with raw CT in the sample of men ($r = 0.038$, $p = 0.999$).

Thus, overall, the composite variable of cardiometabolic health appears moderately correlated

to a marker of low-grade systemic inflammation, while physical robustness was only very

weakly associated ($r < 0.1$) to such a marker.

Latent dimensions linking risk factors to subcortical and cerebellar volumes

Results yielded by the analysis of subcortical and cerebellar data are reported in Supplementary

results section 2.9. Overall, more significant latent dimensions appeared and only one was

partly related to the two main dimensions or risk factors found in association to cortical

structure. This suggests that subcortical regions have different patterns of interindividual

variability than cortical regions (likely higher interindividual variability and more vulnerability

to exposome factors). Accordingly, their associations with risk factors should be the focus of

specific studies. In the following discussion, we focus on cortical structure.

Discussion

We report two latent dimensions characterizing the interplay between a wide range of risk

factors for non-communicable diseases with region-wise CT and GMV across the whole cortex.

One of these latent dimensions highlighted the relevance of cardiometabolic health for inter-

individual variability of brain structure in a healthy sample. Importantly, this latent dimension

was stable across sexes, across brain structural measures (CT and GMV) and across brain size

corrections (raw, proportional, and brain-size corrected CT). Accordingly, this latent

dimension cannot be explained by a confounding effect of head size/morphology that could be

conveyed in brain structural estimates. The other latent dimension highlights the relationship

between physical robustness and brain structure. Even though this latent dimension was stable

within analyses, the stability across sexes, brain structural measures, and brain size corrections

was limited. In addition, our results underline the multi-level nature of the association between

risk factors and brain structure, linking the brain patterns of both latent dimensions with the

spatial distribution of several neurotransmitter systems.

We showed that the variability in cardiometabolic health is related to an axis of variability in

CT extending from the insula and cingulate cortex to the occipital lobes and superior parietal

regions. Overall, this latent dimension indicates that regions engaged in processing internal

information such as emotional and motivational systems (including for instance insula,

orbitofrontal cortex, and anterior cingulate cortex) show positive loadings. Accordingly, this

CT pattern was significantly related to dopamine and serotonin systems. In contrast, more

dorsal regions typically engaged in dorsal attention and executive systems (such as the lateral

superior prefrontal and parietal cortex) show negative CT loadings on the latent dimension.

Similarly, cardiometabolic health was negatively associated with GMV in the frontal regions.

However, it was positively associated with GMV primarily in sensorimotor regions (peri-

central regions), but also to a lesser extent to the ventral attention network (lateral parietal and

middle temporal regions). This suggests that cardiometabolic health may be differently related

to variability in CT and variability in GMV in the population. More concretely, our results

suggest that better cardiometabolic health is related to increased CT in emotional and

motivational systems along with increased GMV in sensorimotor cortex. Future studies should

further elucidate the mechanism behind these associations to identify whether improving

cardiometabolic health can promote brain structural health in motivation, emotion and

sensorimotor networks and hence improve mental health and motor function in aging. The

potential role of neurotransmitters system in these mechanisms is further discussed below.

In the current stage of knowledge, our results are in line with recent reports pointing to the

important role of cardiometabolic health for brain health. For instance, several studies have

linked cardiometabolic factors to brain structure, such as reduced total brain volumes

(38,83,84), reduced grey matter volumes (31,38,84), or to structural markers of brain aging

(85). Interestingly, our findings are in contrast with works that have reported only reductions,

but not increases, in brain structural measures in association with risk factors or

cardiometabolic health. Differences in the health status of the samples might explain these

differential effects. Apart from associations with brain structure, cardiometabolic health has

also been associated with brain function, such as cognition (38), dementia (83), and other

neuropsychiatric disorders and symptoms (21,29,38,86,87). In our study, a moderate

correlation between the composite variable of cardiometabolic health and a marker of

peripheral inflammation further suggests a role for low-grade chronic inflammation in the

association between cardiometabolic health and brain health.

Actually, several studies have pointed out that a causal factor or mediator in the association

between risk factors for non-communicable diseases and brain health might be inflammation

(36–40), and in particular low-grade systemic inflammation (21,36,38,39,88,89). Risk factors

for non-communicable diseases are also risk factors for low-grade systemic chronic

inflammation (6,39,89). In turn, low-grade systemic chronic inflammation has been associated

with non-communicable diseases (6,88), including neuropsychiatric illnesses (21,38,39,88–

92). For instance, chronic inflammation in the adipose tissue has been associated with the

development of neurodegenerative disorders such as Alzheimer's disease (39,93). Moreover,

inflammatory factors have been reported to mediate the association between BMI and other

risk factors with cortical structure and behavior (36,37) (however, see (94)). In fact,

inflammation has been proposed as the cause of comorbidities between non-communicable

diseases (86,90), such as depression and cardiovascular or neurodegenerative illnesses (90).

Thus, we can speculate here that cardiometabolic health, partly via inflammation pathways,

influence brain health with potential impacts on emotion and motivation behavioral systems,

as well as to motor systems.

Our results also show that interindividual variability in physical robustness (fat-free mass and

strength) is associated with interindividual variability in brain structure, in both men and

women for GMV, but only robustly in men for CT. At the brain level, this CT pattern in men

shows highest loadings mainly across the whole lateral frontal cortex and the superior medial

frontal gyrus. The pattern of GMV associated to physical robustness also shows high loadings

on medial frontal regions, along with lateral temporal regions. Accordingly, this GMV pattern

is also significantly related to cortical patterns of developmental area scaling, along with

evolutionary expansion and sensory-association axis. This may suggest common mechanisms

underlying physical health (likely musculoskeletal health) and brain health in key regions for

human high-level cognition (such as the frontal cortex). Interestingly, time spent watching

television appears on the negative pole of these dimensions suggesting that sedentary lifestyles

may threaten this aspect of human brain health.

It should be noted that the variability of physical robustness in women appears to be linked to

GMV but not to CT, while in men it is associated to both structural markers. This sex-variability

could be explained by the risk factors profile in association with CT in men capturing a negative

pole of air pollution and smoking. Previous studies have pointed towards sex-variability in the

effect of air pollution on health and particularly on brain and behavior (95–99). This could be

due to sex-specific exposures to pollutants, with more interindividual variability in exposures

in men allowing us to observe associations with health outcomes. In line with this hypothesis,

the distribution of variables related to air pollution in our sample of men include a few more

extreme values than in women (Figure S2). Alternatively, a potential sex-specific effect of air

pollutants could also be due to different biological aspects being differentially affected in men

in comparison to women, or to sex-specific pathways underpinning the association between air

pollution and brain structure. The mechanisms underpinning this difference should be explored

in future studies. Moreover, studies analyzing the association between health variables and

neuroimaging should consider this and perform sex-specific analyses. In the current stage of

knowledge, we can only speculate from our results that air pollution and smoking may

represent a specific threat for men brain health in high-level associations regions.

Interestingly, the brain patterns associated to both dimensions of risk factors were generally

significantly correlated to the spatial distribution of most neurotransmitter systems. It should

also be noted that the neurotransmitters associated to the latent dimension have in turn been

linked to phenotypes related to the risk factors captured in the latent dimension. For instance,

the serotoninergic system is associated with energy balance and feeding behavior (100,101),

obesity (100) and physical activity (102). Specifically, the 5-HT1a receptor has been associated

with food intake (101), anorexia nervosa and bulimia nervosa (103). The dopaminergic system

has been associated with physical activity (102), and specifically, the receptor D2 has been

associated with disorders related to eating behavior, such as anorexia nervosa, bulimia nervosa,

and obesity (103). Overall, the evidence indicates that these neurotransmitter systems are

associated with imbalances in energy homeostasis and feeding behavior, which are phenotypes

related to the pattern of risk factors captured in our latent dimension. This suggests that the

clinically relevant interaction between brain and body health that has been called to attention

recently (21,29) might be mediated by processes associated with neurotransmitter systems

and/or importantly alter neurotransmitter systems. For instance, cardiometabolic factors may

lead to perturbations in the serotoninergic system via brain structural alterations ultimately

leading to immune-metabolic depression (104).

Evidence of the mechanistic cause linking neurotransmitter systems with both, risk factors for

non-communicable diseases and brain structure, also points to inflammation. Several studies

have shown a crosstalk between the immune system and several neurotransmitter systems, such

as serotoninergic, noradrenergic and dopaminergic systems (92,105). For instance, certain

neurotransmitter receptors, including 5-HT1a, D1 and D2, have immunologic functions

(88,105), and can lead to the disruption of homeostasis (88) and to inflammation. Accordingly,

immune cells express neurotransmitter receptors (105). In turn, immunological factors can

regulate normal cellular functions, including neurotransmission and synaptic plasticity (88). In

sum, the crosstalk between inflammatory factors and neurotransmitter systems is bidirectional

(88,105) and is relevant for several non-communicable diseases (105). Moreover, since these

neurotransmitter systems are implicated in mental disorders and in somatic non-communicable

diseases (88,90), the mechanisms underlying the comorbidity between mental and somatic

disorders might be partly related with alterations in neurotransmitter systems (86).

Accordingly, it is likely that a vicious cycle is engaged when cardiometabolic and

musculoskeletal systems, immunoinflammatory systems, brain structure, and

neurotransmitters systems are altered complexifying treatment of neuropsychiatric disorders.

In that context, our findings contribute to the recognized urgency to characterize brain-body

interactions for its implementation in clinical practice (21,29,86). For instance, it is not

common to monitor physical illnesses in patients with neuropsychiatric disorders. However,

recently it has been pointed out that neuropsychiatric disorders are associated with symptoms

of physical illnesses and that poor physical health is a more pronounced effect than brain

phenotypes in these patients (29). Given that the interplay between brain and body health is not

well understood, the clinical practice nowadays has limited tools to exploit this interaction not

only for a comprehensive monitoring of health, but also for its use as biomarkers. Hence,

research characterizing brain and body interactions is a major priority for global health because

it will guide the discovery of new integrated disease manifestations and pave the way for the

development of new therapies and clinical interventions (21,29).

Conclusion

Our study shows two latent dimensions linking cardiometabolic health and physical robustness

to patterns of CT and GMV variability across the whole cortex. In turn, the captured CT and

GMV brain patterns were associated with the cortical distribution of several neurotransmitter

systems. Furthermore, cardiometabolic health was associated to a peripheral marker of low-

grade systemic inflammation. Hence, our study shows that body health, brain structure, and

neurotransmitter systems are interrelated, highlighting the multi-level nature of health. Also,

our work contributes to questioning the classic consideration of neuropsychiatric and somatic

illnesses as separate categories (88) and supports the view of a needed integration of brain and

physical health in clinical practice (21,29).

Declarations

Authors' contributions

ENS conceptualized the study, developed software, prepared data, performed analyses, and

contributed to discussion and interpretation of results. SMB contributed to study

conceptualization, data preparation, software development, and to interpretation and discussion

of results. MM contributed to data preparation and software development. AM contributed to

software and methodology development. FH contributed to data preparation and preprocessing.

JMM contributed to software and methodology development. MT contributed to discussion

and interpretation of results. SBE acquired funding and contributed to discussion and

interpretation of results. SG acquired funding, conceptualized the study, and contributed to

discussion and interpretation of results. The contribution of ENS has been done in partial

fulfillment of the requirements for a Ph.D. thesis. All authors contributed to the revision of the

manuscript. All authors contributed to writing and reviewing of the manuscript and approved

the final version. ENS and SMB have accessed and verified the data.

Declarations of interest

The authors declare no competing interests.

Acknowledgements

This research has been conducted using the UK Biobank Resource under Application Number

41655. This work was supported by the Deutsche Forschungsgemeinschaft (DFG, GE 2835/2–

1, GE 2835/9-1, SFB 1451 – Project-ID 431549029) (https://www.dfg.de/), granted to SG. SG

received salary from GE 2835/2–1. MM received salary from GE 2835/9-1. SMB is supported

by the MODS project funded from the programme "Profilbildung 2020" (grant no.

PROFILNRW-2020-107-A), an initiative of the Ministry of Culture and Science of the State

of Northrhine Westphalia (https://www.mkw.nrw/). SMB is supported by the EBRAINS 2.0

Project funded from the European Union's Horizon Europe Programme under the Specific

Grant Agreement No. 101147319. EBRAINS is funded by the Horizon Europe Framework

Programme   (©2023   https://ebrains.eu,   https://research-and-

innovation.ec.europa.eu/funding/funding-opportunities/funding-programmes-and-open-

calls/horizon-europe_en). The funders did not play any role in the study design, data collection

and analysis, decision to publish, or preparation of the manuscript.

Data and code sharing statement

Access to UKB data is explained at https://www.ukbiobank.ac.uk/enable-your-research. The

code used for the machine learning framework (https://doi.org/10.5281/zenodo.7153571) has

been made publicly available at https://github.com/mlnl/cca_pls_toolkit

Bibliography

1. World Health Organization. Noncommunicable diseases: progress monitor 2022. 2022.

2. World Health Organization. Noncommunicable diseases country profiles 2018. 2018.

3. Lloyd-Jones DM, Allen NB, Anderson CAM, Black T, Brewer LC, Foraker RE, et al.
Life's Essential 8: Updating and Enhancing the American Heart Association's Construct
of Cardiovascular Health: A Presidential Advisory from the American Heart
Association. Vol. 146, Circulation. Lippincott Williams and Wilkins; 2022. p. E18–43.

4. Livingston G, Huntley J, Liu KY, Costafreda SG, Selbæk G, Alladi S, et al. Dementia
prevention, intervention, and care: 2024 report of the Lancet standing Commission. Vol.
404, The Lancet. Elsevier B.V.; 2024. p. 572–628.

5. Farina FR, Bridgeman K, Gregory S, Crivelli L, Foote IF, Jutila OEI, et al. Next
generation brain health: transforming global research and public health to promote
prevention of dementia and reduce its risk in young adult populations. Vol. 5, The
Lancet Healthy Longevity. Elsevier Ltd; 2024.

6. Furman D, Campisi J, Verdin E, Carrera-Bastos P, Targ S, Franceschi C, et al. Chronic
inflammation in the etiology of disease across the life span. Nature Medicine.
2019;25(12):1822–32.

7. Beydoun MA, Beydoun HA, Gamaldo AA, Teel A, Zonderman AB, Wang Y.
Epidemiologic studies of modifiable factors associated with cognition and dementia:
systematic review and meta-analysis. BMC public health. 2014;14(1):1–33.

8. Opel N, Thalamuthu A, Milaneschi Y, Grotegerd D, Flint C, Leenings R, et al. Brain
structural abnormalities in obesity: relation to age, genetic risk, and common psychiatric
disorders: Evidence through univariate and multivariate mega-analysis including 6420
participants from the ENIGMA MDD working group. Molecular Psychiatry.
2021;26(9):4839–52.

9. Medic N, Ziauddeen H, Ersche KD, Farooqi IS, Bullmore ET, Nathan PJ, et al. Increased
body mass index is associated with specific regional alterations in brain structure.
International Journal of Obesity. 2016;40(7):1177–82.

10.   Tüngler A, Auwera SV der, Wittfeld K, Frenzel S, Terock J, Röder N, et al. Body mass
index but not genetic risk is longitudinally associated with altered structural brain
parameters. Scientific Reports. 2021;11(1):1–11.

11.   Ronan L, Alexander-bloch AF, Wagstyl K, Farooqi S, Brayne C, Tyler LK, et al.
Obesity associated with increased brain age from midlife. Neurobiology of Aging.
2016;47:63–70.

12.   Kim H, Kim C, Seo SW, Na DL, Kim HJ, Kang M, et al. Association between body
mass index and cortical thickness: Among elderly cognitively normal men and women.
International Psychogeriatrics. 2015;27(1):121–30.

13. Gurholt TP, Kaufmann T, Frei O, Alnæs D, Haukvik UK, Meer D van der, et al.
Population-based body–brain mapping links brain morphology with anthropometrics
and body composition. Translational Psychiatry [Internet]. 2021;11(1). Available from:
http://dx.doi.org/10.1038/s41398-021-01414-7

14. Caunca MR, Gardener H, Simonetto M, Cheung YK, Alperin N, Yoshita M, et al.
Measures of obesity are associated with MRI markers of brain aging: The Northern
Manhattan Study. Neurology. 2019;93(8):e791–803.

15. Kharabian-Masouleh S, Beyer F, Lampe L, Loeffler M, Luck T, Riedel-Heller SG, et al.
Gray matter structural networks are associated with cardiovascular risk factors in
healthy older adults. Journal of Cerebral Blood Flow and Metabolism. 2018;38(2):360–
72.

16. Chen X, Wen W, Anstey KJ, Sachdev PS. Effects of cerebrovascular risk factors on
gray matter volume in adults aged 60-64 years: A voxel-based morphometric study.
Psychiatry Research - Neuroimaging. 2006;147(2–3):105–14.

17. Vakhrusheva J, Marino B, Stroup TS, Kimhy D. Aerobic Exercise in People with
Schizophrenia: Neural and Neurocognitive Benefits. Current Behavioral Neuroscience
Reports. 2016;3(2):165–75.

18. Gu Y, Brickman AM, Stern Y, Habeck CG, Razlighi QR, Luchsinger JA, et al.
Mediterranean diet and brain structure in a multiethnic elderly cohort. Neurology.
2015;85(20):1744–51.

19. Jacków-Nowicka J, Podgórski P, Bladowska J, Szcześniak D, Rymaszewska J, Zatońska
K, et al. The Impact of Common Epidemiological Factors on Gray and White Matter
Volumes in Magnetic Resonance Imaging–Is Prevention of Brain Degeneration
Possible? Frontiers in Neurology. 2021;12(July):1–11.

20. Xu J, Liu N, Polemiti E, Garcia-Mondragon L, Tang J, Liu X, et al. Effects of urban
living environments on mental health in adults. Nature Medicine. 2023;29(6):1456–67.

21. Firth J, Siddiqi N, Koyanagi A, Siskind D, Rosenbaum S, Galletly C, et al. The Lancet
Psychiatry Commission: a blueprint for protecting physical health in people with mental
illness. The Lancet Psychiatry. 2019;6(8):675–712.

22. Mihalik A, Chapman J, Adams RA, Winter NR, Ferreira FS, Shawe-Taylor J, et al.
Canonical Correlation Analysis and Partial Least Squares for Identifying Brain–
Behavior Associations: A Tutorial and a Comparative Study. Biological Psychiatry:
Cognitive Neuroscience and Neuroimaging. 2022;7(11):1055–67.

23. Mihalik A, Ferreira FS, Moutoussis M, Ziegler G, Adams RA, Rosa MJ, et al. Multiple
Holdouts With Stability: Improving the Generalizability of Machine Learning Analyses
of Brain–Behavior Relationships. Biological Psychiatry. 2020;87(4):368–76.

24. Nicolaisen-Sobesky E, Mihalik A, Kharabian-Masouleh S, Ferreira FS, Hoffstaedter F,
Schwender H, et al. A cross-cohort replicable and heritable latent dimension linking
behaviour to multi-featured brain structure. Communications Biology. 2022;5(1):1–36.

25.   Maleki Balajoo S, Plachti A, Nicolaisen-Sobesky E, Dong D, Hoffstaedter F, Meuth S,
et al. Data-driven identification of a left anterior hippocampus' morphological network
associated with self-regulation. Research Square [Internet]. 2024; Available from:
https://www.researchsquare.com/article/rs-4170788/v1

26.   Han F, Gu Y, Brown GL, Zhang X, Liu X. Neuroimaging contrast across the cortical
hierarchy is the feature maximally linked to behavior and demographics. NeuroImage.
2020;215(April):116853.

27.   Mihalik A, Ferreira FS, Rosa MJ, Moutoussis M, Ziegler G, Monteiro JM, et al. Brain-
behaviour modes of covariation in healthy and clinically depressed young people.
Scientific Reports. 2019;9(1):1–11.

28.   Küppers V, Bi H, Nicolaisen-Sobesky E, Hoffstaedter F, Yeo BTT, Drzezga A, et al.
Lower motor performance is linked with poor sleep quality, depressive symptoms, and
grey matter volume alterations. bioRxiv : the preprint server for biology [Internet]. 2024
Jun; Available from: http://www.ncbi.nlm.nih.gov/pubmed/38895316
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC11185664

29.   Tian YE, Biase MAD, Mosley PE, Lupton MK, Xia Y, Fripp J, et al. Evaluation of
Brain-Body Health in Individuals with Common Neuropsychiatric Disorders. JAMA
Psychiatry. 2023;80(6):567–76.

30.   Beyer F, Masouleh SK, Kratzsch J, Schroeter ML, Röhr S, Riedel-Heller SG, et al. A
metabolic obesity profile is associated with decreased gray matter volume in cognitively
healthy older adults. Frontiers in Aging Neuroscience. 2019;10(JUL):1–14.

31.   Schaare HL, Masouleh SK, Beyer F, Kumral D, Uhlig M, Reinelt JD, et al. Association
of peripheral blood pressure with gray matter volume in 19- to 40-year-old adults.
Neurology. 2019;92(8):E758–73.

32.   Beyer F, Masouleh SK, Huntenburg JM, Lampe L, Luck T, Riedel-Heller SG, et al.
Higher body mass index is associated with reduced posterior default mode connectivity
in older adults. Human Brain Mapping. 2017;38(7):3502–15.

33.   Murray S, Tulloch A, Gold MS, Avena NM. Hormonal and neural mechanisms of food
reward, eating behaviour and obesity. Nature Reviews Endocrinology. 2014;10(9):540–
52.

34.   Hansen JY, Shafiei G, Markello RD, Smart K, Cox SML, Nørgaard M, et al. Mapping
neurotransmitter systems to the structural and functional organization of the human
neocortex. Nature Neuroscience. 2022;25(11):1569–81.

35.   Markello RD, Hansen JY, Liu Z qi, Bazinet V, Shafiei G, Suárez LE, et al. neuromaps :
structural and functional interpretation of brain maps. Nature Methods. 2022;

36.   Corlier F, Hafzalla G, Faskowitz J, Kuller LH, Becker JT, Lopez OL, et al. Systemic
inflammation as a predictor of brain aging: Contributions of physical activity, metabolic
risk, and genetic risk. NeuroImage. 2018 May;172:118–29.

37.   Marsland AL, Gianaros PJ, Kuan DCH, Sheu LK, Krajina K, Manuck SB. Brain
morphology links systemic inflammation to cognitive function in midlife adults. Brain,
Behavior, and Immunity. 2015;48:195–204.

38.   Miller AA, Spencer SJ. Obesity and neuroinflammation: A pathway to cognitive
impairment. Brain, Behavior, and Immunity. 2014 Nov;42:10–21.

39.   Patel V, Edison P. Cardiometabolic risk factors and neurodegeneration: a review of the
mechanisms underlying diabetes, obesity and hypertension in Alzheimer's disease. Vol.
95, Journal of Neurology, Neurosurgery and Psychiatry. BMJ Publishing Group; 2024.
p. 581–9.

40.   Patel Y, Woo A, Shi S, Ayoub R, Shin J, Botta A, et al. Obesity and the cerebral cortex:
Underlying neurobiology in mice and humans. Brain, Behavior, and Immunity. 2024
Jul;119:637–47.

41.   Genon S, Eickhoff SB, Kharabian-Masouleh S. Linking interindividual variability in
brain structure to behaviour. Nature Reviews Neuroscience. 2022;23(5):307–18.

42.   Miller KL, Alfaro-Almagro F, Bangerter NK, Thomas DL, Yacoub E, Xu J, et al.
Multimodal population brain imaging in the UK Biobank prospective epidemiological
study. Nature Neuroscience. 2016;19(11):1523–36.

43.   Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J, et al. UK Biobank: An
Open Access Resource for Identifying the Causes of a Wide Range of Complex
Diseases of Middle and Old Age. PLoS Med. 2015 Mar 31;12(3):e1001779.

44.   Alfaro-Almagro F, Jenkinson M, Bangerter NK, Andersson JLR, Sotiropoulos SN,
Jbabdi S, et al. Image processing and Quality Control for the first 10,000 brain imaging
datasets from UK Biobank. NeuroImage. 2018;166(October 2017):400–24.

45.   Destrieux C, Fischl B, Dale A, Halgren E. Automatic parcellation of human cortical gyri
and sulci using standard anatomical nomenclature. NeuroImage. 2010;53(1):1–15.

46.   Sasse L, Nicolaisen-Sobesky E, Dukart J, Eickhoff SB, Götz M, Hamdan S, et al.
Overview of leakage scenarios in supervised machine learning. J Big Data. 2025 May
29;12(1):135.

47.   Hotelling H. Relations between two sets of variates. Vol. 28, Biometrika. 1936. p. 3/4.

48.   Helmer M, Warrington S, Mohammadi-Nejad AR, Ji JL, Howell A, Rosand B, et al. On
stability of Canonical Correlation Analysis and Partial Least Squares with application to
brain-behavior associations. BioRxiv. 2023;

49.   Vinod HD. Canonical ridge and econometrics of joint production. Journal of
Econometrics. 1976;4(2):147–66.

50.   Hardoon DR, Szedmak S, Shawe-Taylor J. Canonical correlation analysis: An overview
with application to learning methods. Neural Computation. 2004;16(12):2639–64.

51.   Kapoor S, Narayanan A. Leakage and the reproducibility crisis in machine-learning-
based science. Patterns. 2023 Sep;4(9).

52. Monteiro JM, Rao A, Shawe-Taylor J, Mourão-Miranda J. A multiple hold-out
framework for Sparse Partial Least Squares. Journal of Neuroscience Methods.
2016;271:182–94.

53. Alexander-Bloch AF, Shou H, Liu S, Satterthwaite TD, Glahn DC, Shinohara RT, et al.
On testing for spatial correspondence between maps of human brain structure and
function. NeuroImage. 2018;178(June):540–51.

54. Beliveau V, Ganz M, Feng L, Ozenne B, Højgaard L, Fisher PM, et al. A High-
Resolution In Vivo Atlas of the Human Brain's Serotonin System. 2017;37(1):120–8.

55. Savli M, Bauer A, Mitterhauser M, Ding YS, Hahn A, Kroll T, et al. Normative
database of the serotonergic system in healthy subjects using multi-tracer PET.
NeuroImage. 2012;63(1):447–59.

56. Dukart J, Holiga Š, Chatham C, Hawkins P, Forsyth A, Mcmillan R, et al. Cerebral
blood flow predicts differential neurotransmitter activity. 2018;1–11.

57. Lukow PB, Martins D, Veronese M, Vernon AC, McGuire P, Turkheimer FE, et al.
Cellular and molecular signatures of in vivo imaging measures of GABAergic
neurotransmission in the human brain. Communications Biology. 2022;5(1):1–11.

58. Bedard M andre, Aghourian M, Legault-denis C, Postuma RB, Soucy J paul, Gagnon J
françois, et al. Brain Cholinergic Alterations in Idiopathic REM Sleep Behaviour
Disorder: A PET Imaging Study with 18F-FEOBV. Sleep Medicine [Internet]. 2018;
Available from: https://doi.org/10.1016/j.sleep.2018.12.020

59. Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E. A multi-
modal parcellation of human cerebral cortex. Nature Publishing Group.
2016;536(7615):171–8.

60. Margulies DS, Ghosh SS, Goulas A, Falkiewicz M, Huntenburg JM, Langs G, et al.
Situating the default-mode network along a principal gradient of macroscale cortical
organization. Proceedings of the National Academy of Sciences of the United States of
America. 2016;113(44):12574–9.

61. Vaishnavi SN, Vlassenko AG, Rundle MM, Snyder AZ, Mintun MA, Raichle ME.
Regional aerobic glycolysis in the human brain. Proceedings of the National Academy
of Sciences of the United States of America. 2010;107(41):17757–62.

62. Wey HY, Gilbert TM, Zürcher NR, She A, Bhanot A, Taillon BD, et al. Insights into
neuroepigenetics through human histone deacetylase PET imaging. Science
Translational Medicine. 2016 Aug;8(351).

63. Hawrylycz MJ, Lein ES, Guillozet-Bongaarts AL, Shen EH, Ng L, Miller JA, et al. An
anatomically comprehensive atlas of the adult human brain transcriptome. Nature. 2012
Sep;489(7416):391–9.

64. Markello RD, Arnatkeviciute A, Poline B, Fulcher BD. Standardizing workflows in
imaging transcriptomics with the abagen toolbox. 2021;1–27.

65.   Malén T, Karjalainen T, Isojärvi J, Vehtari A, Bürkner PC, Putkinen V, et al. Atlas of
type 2 dopamine receptors in the human brain: Age and sex dependent variability in a
large PET cohort. NeuroImage. 2022 Jul;255:119149.

66.   Sasaki T, Ito H, Kimura Y, Arakawa R, Takano H, Seki C, et al. Quantification of
Dopamine Transporter in Human Brain Using PET with[18] F-FE-PE2I. J Nucl Med. 2012
Jul;53(7):1065–73.

67.   Gallezot JD, Nabulsi N, Neumeister A, Planeta-Wilson B, Williams WA, Singhal T, et
al. Kinetic Modeling of the Serotonin 5-HT$_{1B}$ Receptor Radioligand [[11] C]P943 in
Humans. J Cereb Blood Flow Metab. 2010 Jan;30(1):196–210.

68.   Gallezot JD, Planeta B, Nabulsi N, Palumbo D, Li X, Liu J, et al. Determination of
receptor occupancy in the presence of mass dose: [[11] C]GSK189254 PET imaging of
histamine H$_3$ receptor occupancy by PF-03654746. J Cereb Blood Flow Metab. 2017
Mar;37(3):1095–107.

69.   Smart K, Cox SML, Scala SG, Tippler M, Jaworska N, Boivin M, et al. Sex differences
in [11C]ABP688 binding: a positron emission tomography study of mGlu5 receptors.
Eur J Nucl Med Mol Imaging. 2019 May;46(5):1179–83.

70.   Mueller S, Wang D, Fox MD, Yeo BTT, Sepulcre J, Sabuncu MR, et al. Individual
Variability in Functional Connectivity Architecture of the Human Brain. Neuron. 2013
Feb;77(3):586–95.

71.   Reardon PK, Seidlitz J, Vandekar S, Liu S, Patel R, Lalonde FM, et al. Normative brain
size variation and brain shape diversity in humans. 2018;1227(June):4–8.

72.   DuBois JM, Rousset OG, Rowley J, Porras-Betancourt M, Reader AJ, Labbe A, et al.
Characterization of age/sex and the regional distribution of mGluR5 availability in the
healthy human brain measured by high-resolution [11C]ABP688 PET. Eur J Nucl Med
Mol Imaging. 2016 Jan;43(1):152–62.

73.   Galovic M, Al-Diwani A, Vivekananda U, Torrealdea F, Erlandsson K, Fryer TD, et al.
In vivo NMDA receptor function in people with NMDA receptor antibody encephalitis
[Internet]. 2021 [cited 2025 Aug 5]. Available from:
http://medrxiv.org/lookup/doi/10.1101/2021.12.04.21267226

74.   Galovic M, Erlandsson K, Fryer TD, Hong YT, Manavaki R, Sari H, et al. Validation of
a combined image derived input function and venous sampling approach for the
quantification of [18F]GE-179 PET binding in the brain. NeuroImage. 2021
Aug;237:118194.

75.   McGinnity CJ, Hammers A, Riaño Barros DA, Luthra SK, Jones PA, Trigg W, et al.
Initial Evaluation of[18] F-GE-179, a Putative PET Tracer for Activated *N* -Methyl d-
Aspartate Receptors. J Nucl Med. 2014 Mar;55(3):423–30.

76.   Kantonen T, Karjalainen T, Isojärvi J, Nuutila P, Tuisku J, Rinne J, et al. Interindividual
variability and lateralization of µ-opioid receptors in the human brain: Individual
differences in the µ-opioid receptor system. NeuroImage. 2020;217(April).

77. Turtonen O, Saarinen A, Nummenmaa L, Tuominen L, Tikka M, Armio RL, et al. Adult
Attachment System Links With Brain Mu Opioid Receptor Availability In Vivo.
Biological Psychiatry: Cognitive Neuroscience and Neuroimaging. 2021;6(3):360–9.

78. Vijay A, Cavallo D, Goldberg A, De Laat B, Nabulsi N, Huang Y, et al. PET imaging
reveals lower kappa opioid receptor availability in alcoholics but no effect of age.
Neuropsychopharmacol. 2018 Dec;43(13):2539–47.

79. Laurikainen H, Tuominen L, Tikka M, Merisaari H, Armio RL, Sormunen E, et al. Sex
difference in brain CB1 receptor availability in man. NeuroImage. 2019 Jan;184:834–
42.

80. Normandin MD, Zheng MQ, Lin KS, Mason NS, Lin SF, Ropchan J, et al. Imaging the
Cannabinoid CB1 Receptor in Humans with [^{11}C] OMAR: Assessment of Kinetic
Analysis Methods, Test–Retest Reproducibility, and Gender Differences. J Cereb Blood
Flow Metab. 2015 Aug;35(8):1313–22.

81. Xu T, Nenning KH, Schwartz E, Hong SJ, Vogelstein JT, Goulas A, et al. Cross-species
functional alignment reveals evolutionary hierarchy within the connectome.
NeuroImage. 2020 Dec;223:117346.

82. Sydnor VJ, Larsen B, Bassett DS, Alexander-Bloch A, Fair DA, Liston C, et al.
Neurodevelopment of the association cortices: Patterns, mechanisms, and implications
for psychopathology. Neuron. 2021 Sep;109(18):2820–46.

83. Tai XY, Veldsman M, Lyall DM, Littlejohns TJ, Langa KM, Husain M, et al.
Cardiometabolic multimorbidity, genetic risk, and dementia: a prospective cohort study.
The Lancet Healthy Longevity. 2022;3(6):e428–36.

84. Vergoossen LWM, Jansen JFA, Backes WH, Schram MT. Cardiometabolic
determinants of early and advanced brain alterations: Insights from conventional and
novel MRI techniques. Vol. 115, Neuroscience and Biobehavioral Reviews. Elsevier
Ltd; 2020. p. 308–20.

85. Beck D, Lange AMG de, Pedersen ML, Alnæs D, Maximov II, Voldsbekk I, et al.
Cardiometabolic risk factors associated with brain age and accelerate brain ageing.
Human Brain Mapping. 2021 Feb;43(2):700–20.

86. Pillinger T, D'Ambrosio E, McCutcheon R, Howes OD. Is psychosis a multisystem
disorder? A meta-review of central nervous system, immune, cardiometabolic, and
endocrine alterations in first-episode psychosis and perspective on potential models.
Vol. 24, Molecular Psychiatry. Nature Publishing Group; 2019. p. 776–94.

87. Schaare HL, Kumral D, Uhlig M, Lemcke L, Villringer A. Associations between mental
health, blood pressure and the development of hypertension. Nature Communications.
2023;14(1):1–17.

88. Gusev E, Sarapultsev A. Interplay of G-Proteins and Serotonin in the
Neuroimmunoinflammatory Model of Chronic Stress and Depression: A Narrative
Review. Current Pharmaceutical Design. 2024;30(3):180–214.

89.   O'Brien PD, Hinder LM, Callaghan BC, Feldman EL. Neurological consequences of
888   obesity. The Lancet Neurology. 2017;16(6):465–77.

889   90.   Anisman H, Merali Z, Hayley S. Neurotransmitter, peptide and cytokine processes in
890   relation to depressive disorder: Comorbidity between depression and neurodegenerative
891   disorders. Progress in Neurobiology. 2008;85(1):1–74.

892   91.   Iakunchykova O, Leonardsen EH, Wang Y. Genetic evidence for causal effects of
893   immune dysfunction in psychiatric disorders: where are we? Transl Psychiatry. 2024 Jan
894   26;14(1):63.

895   92.   Leonard BE. Impact of inflammation on neurotransmitter changes in major depression:
896   An insight into the action of antidepressants. Progress in Neuro-Psychopharmacology
897   and Biological Psychiatry. 2014;48:261–7.

898   93.   Nguyen TT, Hulme J, Vo TK, Vo GV. The Potential Crosstalk Between the Brain and
899   Visceral Adipose Tissue in Alzheimer's Development. Neurochemical Research.
900   2022;47(6):1503–12.

901   94.   Iakunchykova O, Schirmer H, Roe JM, Sørensen Ø, Wilsgaard T, Hopstock LA, et al.
902   Longitudinal and concurrent C-reactive protein and diet associations with cognitive
903   function in the population-based Tromsø study. Journal of Alzheimer's Disease. 2025
904   Mar;104(2):403–13.

905   95.   Cotter DL, Ahmadi H, Cardenas-Iniguez C, Bottenhorn KL, Gauderman WJ,
906   McConnell R, et al. Exposure to multiple ambient air pollutants changes white matter
907   microstructure during early adolescence with sex-specific differences. Commun Med.
908   2024 Aug 1;4(1):155.

909   96.   Kim H, Noh J, Noh Y, Oh SS, Koh SB, Kim C. Gender Difference in the Effects of
910   Outdoor Air Pollution on Cognitive Function Among Elderly in Korea. Front Public
911   Health. 2019 Dec 10;7:375.

912   97.   Kuźma Ł, Struniawski K, Pogorzelski S, Bachórzewska-Gajewska H, Dobrzycki S.
913   Gender Differences in Association between Air Pollution and Daily Mortality in the
914   Capital of the Green Lungs of Poland–Population-Based Study with 2,953,000 Person-
915   Years of Follow-Up. JCM. 2020 Jul 23;9(8):2351.

916   98.   Mo S, Wang Y, Peng M, Wang Q, Zheng H, Zhan Y, et al. Sex disparity in cognitive
917   aging related to later-life exposure to ambient air pollution. Science of The Total
918   Environment. 2023 Aug;886:163980.

919   99.   Oiamo T, Luginaah I. Extricating Sex and Gender in Air Pollution Research: A
920   Community-Based Study on Cardinal Symptoms of Exposure. IJERPH. 2013 Aug
921   22;10(9):3801–17.

922   100.   van Galen KA, Horst KW ter, Serlie MJ. Serotonin, food intake, and obesity. Obesity
923   Reviews. 2021;22(7):1–13.

924   101.   Voigt JP, Fink H. Serotonin controlling feeding and satiety. Behavioural Brain
925   Research. 2015;277:14–31.

926   102. Cordeiro LMS, Rabelo PCR, Moraes MM, Teixeira-Coelho F, Coimbra CC, Wanner
927   SP, et al. Physical exercise-induced fatigue: The role of serotonergic and dopaminergic
928   systems. Brazilian Journal of Medical and Biological Research. 2017;50(12):1–13.

929   103. Frank GKW. Advances from neuroimaging studies in eating disorders. CNS Spectrums.
930   2015;20(4):391–400.

931   104. Penninx BWJH, Lamers F, Jansen R, Berk M, Khandaker GM, De Picker L, et al.
932   Immuno-metabolic depression: from concept to implementation. The Lancet Regional
933   Health - Europe. 2025 Jan;48:101166.

934   105. Matt SM, Gaskill PJ. Where Is Dopamine and how do Immune Cells See it?: Dopamine-
935   Mediated Immune Cell Function in Health and Disease. Journal of Neuroimmune
936   Pharmacology. 2020;15(1):114–64.

Captions

1. Supplementary methods

1.1 Sex-specific analyses.
1.2 Comparison of latent dimensions with demographics.
1.3 Latent dimensions linking risk factors to subcortical and cerebellar volumes.

2. Supplementary results

2.1 Latent dimensions linking cardiometabolic health to raw cortical thickness and raw grey
matter volume.
2.2 Latent dimensions linking cardiometabolic health to proportional cortical thickness and
proportional grey matter volume.
2.3 Latent dimensions linking cardiometabolic health to corrected cortical thickness and
corrected grey matter volume.
2.4 Latent dimension of cardiometabolic health and demographics.
2.5 Latent dimensions linking physical robustness to raw cortical thickness and raw grey matter
volume.
2.6 Latent dimensions linking physical robustness to proportional cortical thickness and
proportional grey matter volume.
2.7 Latent dimensions linking physical robustness to corrected cortical thickness and corrected
grey matter volume.
2.8 Latent dimension of physical robustness and demographics.
2.9 Latent dimensions linking risk factors to subcortical and cerebellar volumes

3. Supplementary tables

Table S1. Variables of risk factors used in this study.
Table S2. Demographic variables used in this study.
Table S3. Brain maps included in neuromaps that were compared with the latent dimension.
Table S4. Statistics of the latent dimensions for raw cortical thickness.
Table S5. Statistics of the latent dimensions for raw grey matter volume.
Table S6. Statistics of the latent dimensions for proportional cortical thickness.
Table S7. Statistics of the latent dimensions for proportional grey matter volume.
Table S8. Statistics of the latent dimensions for corrected cortical thickness.
Table S9. Statistics of the latent dimensions for corrected grey matter volume.
Table S10. Association between the brain loadings of the latent dimension of cardiometabolic
health and neuromaps.
Table S11. Association between the brain loadings of the latent dimension of physical
robustness and neuromaps.
Table S12. Statistics of the latent dimensions for subcortical and cerebellar volumes.

4. Supplementary figures

Figure S1. Distribution of risk factors. Part 1.