Integrative Network Analysis Reveals Novel Moderators of Aβ-Tau Interaction in

Alzheimer's Disease

Akihiro Kitani[1], Yusuke Matsui[1,2]*

[1]Biomedical and Health Informatics Unit, Department of Integrated Health Science,

Nagoya University Graduate School of Medicine, Nagoya, Japan

[2]Institute for Glyco-core Research (iGCORE), Nagoya University, 461-8673 Nagoya,

Aichi, Japan.

*Corresponding author

Yusuke Matsui

E-mail: matsui@met.nagoya-u.ac.jp

Abstract

Background: Although interactions between amyloid-beta and tau proteins have

been implicated in Alzheimer's disease (AD), the precise mechanisms by which these

interactions contribute to disease progression are not yet fully understood. Moreover,

despite the growing application of deep learning in various biomedical fields, its

application in integrating networks to analyze disease mechanisms in AD research

remains limited. In this study, we employed BIONIC, a deep learning-based network

integration method, to integrate proteomics and protein–protein interaction data, with

an aim to uncover factors that moderate the effects of the Aβ-tau interaction on mild

cognitive impairment (MCI) and early-stage AD.

Methods: Proteomic data from the ROSMAP cohort were integrated with

protein–protein interaction (PPI) data using a Deep Learning-based model. Linear

regression analysis was applied to histopathological and gene expression data, and

mutual information was used to detect moderating factors. Statistical significance was

determined using the Benjamini-Hochberg correction ($p < 0.05$).

Results: Our results suggested that astrocytes and GPNMB+ microglia moderate the

Aβ-tau interaction. Based on linear regression with histopathological and gene

expression data, GFAP and IBA1 levels and *GPNMB* gene expression positively

contributed to the interaction of tau with Aβ in non-dementia cases, replicating the

results of the network analysis.

Conclusions: These findings indicate that GPNMB+ microglia moderate the Aβ-tau

interaction in early AD and therefore are a novel therapeutic target. To facilitate further

research, we have made the integrated network available as a visualization tool for the

scientific community (URL: https://igcore.cloud/GerOmics/AlzPPMap).

Keywords

Alzheimer's disease, Tau, Amyloid β, microglia, astrocyte, GPNMB, proteomics,

network integration, moderator, interaction effects

Background

Alzheimer's disease (AD) is the most common form of dementia,

characterized by extracellular amyloid-beta (Aβ) plaques and intracellular tau protein

tangles, which are believed to be central to its pathology.[1–3] Recent studies have

reported an interaction between Aβ and tau proteins in the pathogenesis of AD,

contributing to disease progression.[4–9] For instance, Lee et al.[7] revealed that

Aβ-tau interactions are associated with the propagation of tau. In practice, lecanemab,

an anti-Aβ antibody, which has been shown to slow cognitive decline in patients with

early-stage AD, reduces total tau protein and P-Tau181 levels in cerebrospinal

fluid.[10] This suggests that Aβ contributes to the effect of tau in AD. Moreover, several

phenotypes are associated with the Aβ-tau interaction, such as astrocyte reactivity,

blood pressure, vascular burden, and microglia[11–15]; however, the detailed

mechanisms are not fully understood. Various cellular perturbations have been

identified through single-nucleus RNA sequencing (snRNA-seq) in AD.[16–22] For

example, a recent snRNA-seq study[17] revealed the presence of Trem2-dependent

disease-associated microglia (DAM) and a unique Serpina3n+C4b+ reactive

oligodendrocyte population in 5XFAD mice. In human AD, distinct glial phenotypes

including *IRF8*-driven reactive microglia and oligodendrocytes with impaired

myelination and metabolic adaptation to neuronal degeneration have been observed.

However, the specific cell types and effects of Aβ-tau interactions are yet to be

elucidated.

Network analyses based on gene or protein co-expression have revealed

gene groups and associated pathways correlated with pathological and clinical factors

in AD.[23–25] Additionally, network integration methods based on multi-omics and

other data types have been developed.[26–29] By integrating both physical and

functional interactions simultaneously, network integration analyses can identify

detailed molecular relationships. However, this approach has largely been applied to

cancer. AD is a complex disease involving various pathologies, and network

integration is a promising approach for determining comprehensive physical and

functional interactions involving Aβ and tau in AD.

In this study, we investigated moderating factors in the Aβ-tau interaction

using a systematic network integration approach. The aim of this study was to

elucidate the cell types and moderating factors in the interaction between Aβ and tau

proteins in early-stage AD. Figure 1 shows the graphical abstract of the study. First, we

integrated the proteomics and protein–protein interaction (PPI) networks by using

BIONIC,[30] a network integration algorithm. In the downstream analysis, glial

cell-related proteins, including those associated with microglia and astrocytes, were

enriched around *APP* and *MAPT* subnetworks, which encode the precursors of Aβ

and tau, respectively. Further analyses of microglial subtypes revealed enrichment in

GPNMB+ microglia. As a validation, linear regression analysis based on

histopathological and gene expression data showed that interactions between Aβ and

IBA1, GFAP, and *GPNMB*, were positively related to tau levels. Our findings highlight

that deep learning-based network integration can serve as a novel framework for

elucidating complex disease mechanisms. The integrated network, made available as

a visualization tool for the scientific community, not only enhances our understanding

of the Aβ-tau interaction but also provides a basis for generating experimental

validation ideas and developing more effective treatments for AD. By leveraging this

tool, researchers can gain insights into the intricate molecular interactions at play in

AD, fostering the discovery of new therapeutic targets.

Methods

Proteomics data from the ROSMAP cohort

For the proteomic analysis, ROSMAP tandem mass tag (TMT) proteomics data were

obtained. TMT proteomics is a technique for the simultaneous quantitative analysis of

protein expression in multiple samples using LC-MS/MS and is commonly employed

in AD research.[24,25] The ROSMAP data set included TMT proteomics data for 400

cases with differences in cognitive status (Alzheimer's dementia, other dementia, MCI,

or no impairment). Data normalization and batch correction were conducted according

to previous described procedures.[31] Four outlier samples were excluded from

subsequent analyses based on a principal component analysis. In this study,

ROSMAP TMT data for MCI and early AD cases were included. Early AD was defined

according to previously reported criteria: meeting the NIA-AA core clinical criteria for

probable AD dementia and having a global CDR score of 0.5 to 1.0 and a CDR

memory box score of 0.5 or greater at screening and baseline.[10] Additionally, the

MMSE effectively discriminates between CDR stages 0.5, 1, 2, and 3 but performs

poorly in the separation of CDR stages zero and 0.5. The previously established

MMSE ranges were 30 for no, 26–29 for questionable, 21–25 for mild, 11–20 for

moderate, and 0–10 for severe dementia.[32] We defined early AD as having an

MMSE score of ≥21.

Proteomics and PPI data integration using a Graph Attention Network-based

method

To integrate the proteomics and PPI data, total values were calculated for each gene

symbol in the proteomics dataset. Pearson correlation coefficients were derived from

the proteomics data, retaining only data with Benjamini-Hochberg (BH)-corrected

p-values of <0.05. Normalization was performed to set maximum values to 1 and

minimum values to 0. For PPI, STRING[33] filtering was used to obtain physical

interactions within *Homo sapiens* with confidence scores exceeding 0.7, and

normalization was performed to set the maximum values to 1 and minimum values to 0.

Proteomic and PPI data were integrated using a recently reported method,

BIONIC,[30] which integrates multiple biological networks while preserving

information using graph attention networks.[34] Default values were used for the

BIONIC hyperparameters. For both proteomics and PPI data, 4,787 proteins common

to both datasets were retained. After integrating the two networks, 512-dimensional

features were output for each protein.

Clustering analysis and dimension reduction using features derived from

BIONIC

Considering that the data derived from neural networks are nonlinear, clustering was

performed following a method similar to that used in snRNA-Seq analyses.[35] After

feature calculation using BIONIC and subsequent scaling to Z-scores, Euclidean

distances were computed to create a distance matrix. Then, setting k = 25, the

k-nearest neighbor method was used to calculate an adjacency matrix and form a

graph structure. This graph structure was then subjected to clustering using the

Louvain method[36] (resolution = 1) to classify each protein into clusters.

Cluster centroid analysis using BIONIC-derived features

After implementing BIONIC to evaluate the networks derived from proteomic and PPI

data, a clustering analysis of proteins was performed. To analyze the relationships

between clusters, the average of the feature values for each cluster was calculated as

its centroid. Pearson correlation coefficients and Euclidean distances between the

clusters were calculated. Furthermore, the network of clusters was visualized using

the reciprocal values of the Euclidean distances as the weights of the edges.

Cell type enrichment analysis based on human brain and microglia snRNA-seq

data

For a cell type enrichment analysis, publicly available human brain[37] and

microglia[21,22] snRNA-seq data were used. Cell annotations for each dataset were

based on publicly available data. The AddModuleScore function was performed using

the Seurat package[38] to calculate the average expression levels in each protein

community. The combined score was calculated by multiplying the expression level by

the proportion of positive cells in each community.

Cell type composition analysis based on human brain and microglia

snRNA-seq data

For a cell type composition analysis, publicly available human microglia[21,22]

snRNA-seq data were used. ANOVA was used to compare the frequencies of different

microglial subtypes across various conditions (Control, Aβ+, and Aβ+Tau+). Next, for

the microglial subtypes "Microglia_GPNMB_LPL" and "Microglia_GPNMB_PLAT," a

Tukey HSD test was used to determine specific differences between groups (for cases

where ANOVA showed $p < 0.05$).

Gene Ontology functional enrichment analysis

A Gene Ontology (GO) analysis was used to explore the biological functions of our

targeted gene set. The GO biological process (GObp) dataset version 2023.1 was

downloaded from the Molecular Signatures Database (MsigDB)

(https://www.gsea-msigdb.org/gsea/msigdb). For the analysis, the enrichment

function in the clusterProfiler package was used, and values of p < 0.05, after BH

correction, were considered significant. The Rrvgo[39] software was used to

summarize the GO terms.

Modulators of the Aβ-tau interaction

To explore modulators of the Aβ-tau interaction based on proteomics data, a slightly

modified version of the MINDy algorithm[40] was employed. Briefly, (1) Euclidean

distances for APP and MAPT were calculated using BIONIC features, and common

elements within the top 5% with respect to proximity were selected for analyses. (2)

For each candidate modulator, the top and bottom 35% of proteins with respect to

expression levels were segregated, and mutual information for APP and Tau was

calculated for each group. Mutual information, which measures the interdependence

between two random variables X and Y, was defined using equation (1) as follows:

$$MI(X,Y) = \sum x \in X \sum y \in Y\, p(x,y) \log\big(p(x)p(y)p(x,y)\big)\, \#(1)$$

Here,

p(x,y) represents the joint probability distribution of *X* and *Y*.

p(x) and *p(y)* are the marginal probability distributions of *X* and *Y*, respectively.

The logarithm base 2 was used.

Subsequently, (3) the difference in mutual information between the top and bottom

35% groups was calculated using equation (2):

$$\Delta MI = |MI(MAPTMtop35, APPMtop35) - MI(MAPTMbottom35, APPMbottom35)| \#(2)$$

Here, M_{top35} and $M_{bottom35}$ indicate the APP and MAPT groups in the top 35% and

bottom 35% of protein expression levels, respectively. This differentiation is crucial for

understanding the difference in mutual information between the two groups, allowing

for a clear analysis of the Aβ-Tau interaction modulators. Bootstrapping with 1,000

iterations was then used to compare mutual information, with values of $p < 0.05$

deemed significant.

Next, factors moderating Aβ-Tau interactions were extracted using data obtained from

BIONIC and a network analysis was conducted. Among candidate modulators

identified in the MINDy analysis, HSPA5, which is closely associated with Aβ[41] and

Tau[42], was included in the analysis. The inner product of the BIONIC-derived

features was calculated to determine protein similarity, retaining the top 5% for

network construction. From this network, the top 5% of factors based on Euclidean

distances to *APP*, *MAPT*, and *HSPA5* were retained to form a subnetwork. Community

detection within this subnetwork was performed using the leading eigenvector function

of the iGraph package.[43]

Neuropathological and RNA-seq datasets

The neuropathological and RNA-Seq datasets used in this study were obtained from

previously published works.[44] Neuropathological data include quantitative

immunostaining results for the microglial marker IBA1, astrocyte marker GFAP, Aβ,

tau, and phosphorylated tau. The datasets included 377 samples collected from the

cortical grey (parietal and temporal) and white matter (parietal) as well as the

hippocampus. Additionally, the frequency distributions of age, sex, CERAD scores,

and Braak scores among the participants were evaluated, differentiating between the

no-dementia and dementia groups. Age was compared between the dementia and

no-dementia groups using a t-test, whereas sex, Braak score, and CERAD score were

compared between the groups using Fisher's exact test.

Linear regression analysis for exploring interaction terms

To explore moderating factors in the Aβ-tau interaction, a linear regression analysis

was used to evaluate interaction effects, as described previously.[13] Measurements

were performed on formalin-fixed, paraffin-embedded tissue sections (5 μm). For Aβ,

the percentage of area covered by Aβ immunoreactivity was used, identifying both

Aβ40 and Aβ42. The analysis focused on the percentage of the area covered by Tau2

immunoreactivity, identifying mature neurofibrillary tangles and dystrophic neurites.

AT8, an antibody that recognizes phospho-tau (Ser202, Thr205), was also utilized.

Interaction effects with Aβ were investigated using immunoreactivities of IBA1 and

GFAP and gene expression levels of *GPNMB*. IBA1 was measured as the percentage

of area covered by IBA1 immunoreactivity, GFAP as the percentage of area covered

by GFAP immunoreactivity (identifying activated astrocytes), and *GPNMB* gene

expression levels were quantified as fragments per kilobase of exon per million reads

mapped (FPKM). The interaction effects were evaluated using the *t*-test, with values of

$p < 0.05$ considered significant.

Trajectory analysis of human microglia snRNA-seq data

A trajectory analysis was conducted using human microglial snRNA-seq data and the

Slingshot package. The default parameters for the slingshot function were utilized,

and the getCurves function was used with the following parameters: approx_points =

300, threshold = 0.01, stretch = 0.8, allow.breaks = FALSE, and shrink = 0.99. From

the dataset reported by Sun et al.,[21] only microglia were extracted from immune cells

for clustering. Clustering was performed using the Seurat package, following the

tutorial provided by the Satija Lab

(https://satijalab.org/seurat/articles/pbmc3k_tutorial). The settings were

FindVariableFeatures with nfeatures = 4000, FindNeighbors, RunUMAP with dims =

1:10, and FindClusters with resolution = 0.5. To visualize gene expression over time,

ggplot2 was used and regression curves were generated using generalized additive

models (GAM).

Heatmaps of CERAD and Braak scores in the ROSMAP cohort data

To investigate whether GPNMB protein expression moderates the Aβ-tau interaction

in patients with MCI and early AD in the ROSMAP cohort, correlations between

CERAD and Braak scores were evaluated. The CERAD score in the ROSMAP cohort

is a semi-quantitative measure of neuritic plaque density, classified into four scores

ranging from 4 to 1, corresponding to no AD, possible AD, probable AD, and definite

AD, respectively. The Braak score is a semi-quantitative measure of the severity of

neurofibrillary tangle pathology, ranging from 0 to 6, corresponding to Braak stages

I–VI. For the top 35% and bottom 35% of patients based on GPNMB protein

expression, the frequency of each CERAD and Braak score was visualized using a

heatmap.

Establishment of AlzPPMap

Using the R Shiny app, we developed a tool named AlzPPMap to analyze and

visualize the integrated network. By inputting one or more gene names, the tool allows

users to select a percentage of neighboring genes within the integrated network. The

selected percentage of data is then used to construct and display a subnetwork.

Results

Study design

The analysis workflow, including an overview of the datasets, methods, and

results, is presented in Supplementary Fig. S1. We extracted proteomics data for

patients with MCI or early-stage AD, constructed a co-expression network, and

integrated it with a PPI network using BIONIC, a deep learning-based method. To

characterize the integrated data, we utilized enrichment analysis and whole

snRNA-seq data (Fig. 2). Using proteomics data, we searched for factors associated

with tau and Aβ (Fig. 3), constructed a subnetwork of tau, Aβ, and moderators, and

performed cell type and functional enrichment analyses (Fig. 4). We validated these

findings using immunohistochemical data from a separate cohort (Fig. 5). Additionally,

for the subnetworks, we obtained microglia-specific snRNA-seq data to perform a

microglial cell subtype enrichment analysis (Fig. 6). Finally, we validated these

279   findings using immunohistochemical and gene expression data (Fig. 7).

281   **Integration of protein co-expression and physical PPI networks**

282   We constructed an integrated framework as a basis for analyzing the effects of

PPIs and their moderators in the AD postmortem brain. To this end, we integrated

proteomic data and a physical PPI network to construct a functional-physical PPI

model. We employed BIONIC,[30] a deep learning method designed to integrate

multiple biological networks while preserving their integrity (Fig. 2a). BIONIC uses a

graph attention network[34] to integrate multiple networks, maintain the distinctive

characteristics of each, and generate new features. It shows good performance[30]

and has been used in analyses of gene interactions in cancer.[45]

290   First, a functional protein interaction network was constructed based on

protein abundance data. We compiled postmortem proteome profiles for patients with

MCI and early-stage AD from the dorsolateral prefrontal cortex, which were available

from the Religious Order Study (ROS) and Rush Memory and Aging Project (MAP)

cohort (https://adknowledgeportal.synapse.org/). Next, we generated a PPI network

using the STRING database[33] (https://string-db.org/) and constructed subnetworks

by prioritizing "high confidence" interactions. Using BIONIC, the integrated framework

was embedded into a lower-dimensional vector space, and the embedded coordinates

were obtained in a 512-dimensional space. Our downstream analysis used 512

integrated network features.

Characterizing integrated protein network modules

Next, we characterized the biological context captured by the integrated

network by evaluating the cell types and biological functions associated with the

protein modules. To investigate these modules, we built k-nearest neighbor graphs of

proteins based on the integrated network to obtain clusters (see Methods and Figs. 2a,

b). The cell types and biological processes corresponding to each module were

inferred using publicly available snRNA-seq data[37] and Gene Ontology terms. The

modules in the integrated network contained cell type signatures (Fig. 2c) and were

enriched in biological processes related to these cell types (Supplementary Table S1

and Fig. 2d). For example, module 1 was characterized by endothelial cells, microglia,

and astrocytes, modules 5, 17, 22, and 27 were associated with oligodendrocytes,

modules 9 and 13 with astrocytes, and module 11 with neurons (Fig. 2c). Consistent

with the cell type results, in module 5, terms related to oligodendrocytes, such as

"Ensheathment of neuron" and "Oligodendrocyte differentiation," were identified,

whereas module 11 was predominantly enriched in synapse-related terms (Fig. 2d).

Identification of *APP* and *MAPT* as adjacent modules in the integrated network

We explored the inter-modular relationships within the integrated network by specifically examining modules that included *APP* and *MAPT*, which serve as markers for Aβ and tau, respectively. The *APP* gene was located in module 11 and *MAPT* was located in module 12. Additionally, *APOE*, which plays a significant role in the accumulation and aggregation of Aβ, was assigned to module 1. We then constructed a network of module–module interactions (Fig. 2e, and Supplementary Fig. S2). Notably, these three modules were in close proximity in the network, with a strong correlation between the *APP* and *MAPT* modules (Pearson correlation coefficient = 0.7667, p < 0.001). Additionally, strong correlations were observed between the *APOE* module and both the *APP* (r = 0.6479, p < 0.001) and *MAPT* modules (r = 0.6289, p < 0.001). Furthermore, within the network, the *MAPT* module was closest to the *APP* module based on Euclidean distances; the *APP* module was second closest to the *MAPT* module. These module-to-module correlations in early-stage AD support the interactions between *APOE*, *APP*, and *MAPT*. [46,47]

Next, we evaluated whether similar findings could be obtained without network integration. Employing the STRING database with settings for high confidence and physical interactions alone, no interactions were detected between the *APP* and *MAPT* modules. Similarly, *APOE* did not interact with the other genes (Supplementary

Fig. S3). When examining protein expression data separately, *APP* and *MAPT* did not

show significant correlations (Pearson's correlation coefficient = -0.0465, p = 0.61).

These results underscore the importance of our integrated network approach in

capturing the biological interactions between Aβ and tau, which are crucial for

understanding early-stage AD development.

Stress response factors moderate Aβ-tau interactions

Next, we aimed to identify factors that could moderate the Aβ-tau interaction

based on the integrated network. For this purpose, we prioritized proteins with strong

moderating effects on APP-MAPT interactions against proteins close to the *APP* and

MAPT modules. We utilized the computational framework of Modulator Inference by

Network Dynamics (MINDy)[40] (Fig. 3a). Briefly, MINDy calculates mutual information

for interaction pairs (such as *APP* and *MAPT*), conditioned by the expression of the

modulator, which reflects the modulating effect. In the present analysis, we extracted

the neighboring proteins of the *APP* and *MAPT* modules in the integrated network (see

Methods, "Modulators of the Aβ-tau interaction"). We then stratified the samples from

the proteomics dataset into groups representing the top and bottom 35% of

expression for each neighboring protein and evaluated differences in the magnitude of

mutual information between *APP* and *MAPT* (Fig. 3a). We analyzed 312 neighboring

proteins and identified the 11 most significant modulators (Fig. 3b). These candidates

included factors involved in the cellular stress response (*HSPD1* and *HSPA5*),[48]

ubiquitin-proteasome system (*USP48*, *KBTBD6*, *UBE2D4*, and *STAMBPL1*),[49–51]

and apoptosis and cellular responses (*MAP3K5*).[51] These results suggest that

proteostasis involving stress response genes is related to the Aβ-tau interaction.

360   To further identify proteins contributing to the Aβ-tau interaction from the

moderator candidates, we investigated proteins known to physically interact with Aβ

and/or tau sourced from the Amyloidome[41] and Tau Interactome.[42] Consequently,

Heat Shock Protein Family A (Hsp70) Member 5 (*HSPA5*) was identified. *HSPA5*, a

marker of ER stress associated with AD,[52,53] has been shown to interact in vitro

with Aβ and tau, mitigating their toxicity[54–58]. Although in vivo studies have

identified it as a candidate therapeutic target[59,60], the detailed mechanisms of

action and its relevance in the Aβ-tau interaction are not fully understood[61].

Glial cells are closely related to the APP-MAPT interaction in the integrated

network

371   We further investigated whether stress responses involving *HSPA5* are

involved in Aβ-tau interactions and identified related cell types and biological

pathways. To address this, we exploited the integrated network extensively and

reconstructed *HSPA5* and its neighboring subnetworks as well as *APP* and *MAPT* (Fig.

4a). Note that the subnetwork included other significant moderators, *STAMBPL1* and

MAP3K5, suggesting that these factors, in addition to *HSPA5*, could be associated

with the Aβ-tau interaction. This subnetwork was characterized by functions in

neuronal death, positive regulation of kinase activity, cellular response to chemical

stress, synapse organization, and regulation of autophagy, confirming its relation to

stress response pathways (Fig. 4b, see also Supplementary Table S2).

We classified the subnetwork proteins into six communities based on topology

and then estimated the cell types and biological functions associated with each

community (Figs. 4c, d, and Supplementary Table S3). We found that glial cells were

the primary cell type related to the stress response subnetwork (Fig. 4c). Additionally,

the combined score for each community, calculated by multiplying the expression level

by the proportion of cells, showed enrichment for glial cells (Supplementary Fig. S4a).

Similar results were obtained using snRNA-seq data from two different cohorts[21,22]

and different regions (Supplementary Figs. S4b, c). However, communities 2 and 6

were excluded from this analysis owing to the small number of genes included. In the

APP-related Community 3, oligodendrocytes were enriched in protein folding,

oxidative stress responses, neuronal death, and autophagy. Community 4, which

contained *MAPT* and *HSPA5*, was characterized by microglia and excitatory neurons

with enrichment in kinase activity. Other communities such as communities 1 and 5

were also enriched in glial cells (Figs. 4c, d). Microglia were detected as a feature of

three of the four significantly enriched submodules (Fig. 4c).

Histopathological data analysis validated the roles of microglia and astrocytes

in the Aβ-tau interaction in early AD

To further investigate the effect of Aβ-tau interaction, we examined publicly

available histopathological (ACT study) data.[44] This dataset contains extensive

multimodal histopathological information, including Aβ and tau pathology. The

distributions of age, sex, CERAD score, and Braak score in groups with and without

dementia are shown in Supplementary Fig. S5. Among these parameters, CERAD

and Braak scores showed significant differences between the groups (Fisher's exact

test, $p = 0.00237$ and $p = 0.000468$, respectively). In the no-dementia group, the

average CERAD score was 1.476 and the average Braak score was 3.421. Braak

stages I and II indicate that neurofibrillary tangles are confined mainly to the entorhinal

region of the brain, whereas stages III and IV involve the limbic regions, such as the

hippocampus.

We analyzed the effect of the interaction between tau and Aβ using a linear

modeling approach. Tau2, an antibody for a mature neurofibrillary tangles and

dystrophic neurites, was used as the dependent variable, whereas Aβ, along with

GFAP or IBA1 (activated astrocyte or microglia markers, respectively), was used as

the independent variable. We investigated the interaction between GFAP or IBA1 and

Aβ, stratifying the analysis by the dementia status and across four brain regions (Fig.

5). For GFAP, a significant positive contribution to the interaction was observed in the

hippocampus of both the dementia and no-dementia groups (HIP, β = 43.59, p = 0.022

and β = 24.852 and p = 0.001, respectively) and in the temporal neocortex of the

no-dementia group (TCx, β = 13.873 and p < 0.001). In the case of IBA1, a positive

contribution trend was observed in the HIP of the no-dementia group, though the

relationship was not significant (β = 4.678 and p = 0.338), whereas a significant

positive contribution was observed in the TCx (β = 4.352 and p < 0.001). Conversely,

in the dementia group, a significant negative contribution was noted in the TCx (β =

-11.124, p = 0.04; Fig. 5b and Supplementary Table S4). Analyses of AT8, an antibody

that recognizes phospho-tau (Ser202 and Thr205) showed the same trend in the

no-dementia group: GFAP (β = 33.912 and p = 0.235 in HIP; β = 62.732 and p < 0.001

in TCx) and IBA1 (β = 14.442 and p = 0.36 in HIP; β = 16.344 and p < 0.001 in TCx)

(Supplementary Table S5). Of note, interactions were observed only in the HIP and

TCx regions, which are vulnerable in patients with early-stage AD.[62]

The average CERAD and Braak scores in the no-dementia group mentioned

earlier suggest that, although not exhibiting dementia, this group exhibited mild

progression of AD pathology (Supplementary Figs. S5c, d). Collectively, these results

support the hypothesis that astrocytes and microglia moderate the Aβ-tau interaction

in early-stage AD.

GPNMB+ microglia function as moderators of the Aβ-tau interaction

We further examined microglial subtypes that contribute to Aβ-tau interactions.

Microglia-overexpressing neuroinflammation-related processes can be assigned to

subtypes that exhibit different cellular states under different AD conditions.[63] We

evaluated whether a certain subtype of microglia moderates the Aβ-tau interaction in

early-stage AD. We accessed recently reported snRNA-seq data[22] from frontal

cortex biopsies of 52 patients with hydrocephalus (NPH). This cohort included

samples with confirmed amyloid and tau pathologies, in which AD was in an

asymptomatic stage. Of the 52 living biopsies, Aβ plaques were identified in 19

subjects, Aβ plaques and phosphorylated tau pathology in eight subjects, and no

pathology in the remaining 25 subjects; for subjects with Aβ pathology, the subsequent

onset of AD was determined in longitudinal studies. In total, 892,828 high-quality

single-nucleus profiles were included in analyses. Here, we focused on the microglial

cell populations.

We performed an enrichment analysis of each microglial subtype for the

response-related subnetwork communities (communities 1–6) based on the

integrated network. GPNMB+ microglia were significantly enriched in community 1

(Fig. 6). The expression level and proportion of cells identified as the GPNMB+

microglial subtype was high for the community 1 signature score (Fig. 6a). In addition,

we compared the compositions of cell subtypes across different statuses in the same

cohort. The frequency of GPNMB+ microglia was significantly higher in Aβ+ and

Aβ+Tau+ samples than in the control group (Supplementary Fig. S6 and

Supplementary Table S6). The combined score for Community 1, calculated by

multiplying the expression level and the proportion of cells with positive expression,

further confirmed that the GPNMB+ microglial subtype was the highest ranked among

the examined subtypes (Fig. 6b). Importantly, the GPNMB+ microglial subtype is

expressed in the early stages of AD and expands as the disease progresses.[22]

To further validate the involvement of GPNMB+ microglia in the Aβ-tau

interaction in early-stage AD, we used another large-scale snRNA-seq data set for

microglia subtypes in AD,[21] including transcriptome data for 194,000 single-nuclear

microglia and multiple AD pathological phenotypes in 443 human subjects. The cell

subtypes were annotated based on a previous study.[22] We found the same trend

towards enrichment for GPNMB+ in Module 1 (Figs. 6c, d). These results suggest

that the stress response module moderating Aβ-tau interactions in early AD

corresponds to GPNMB+ microglia.

Aβ-tau stress response mediators are activated prior to GPMNB+ microglial

activation

We further examined the relationship between stress response-related

mediators of the Aβ-tau interaction and GPNMB+ microglia. Previous studies have

reported that ER stress induces GPNMB expression in neurons but not in

microglia.[64] We hypothesized that stress response mediators are involved in the

transition of microglia from a homeostatic state to an activated GPNMB+ state in the

human brain. To test this hypothesis, trajectory analysis of the two snRNA-seq

datasets used in the previous analyses was performed using Slingshot.[65] Briefly, in

this analysis, cells are embedded in the manifold space based on expression

similarity between cells. This trajectory reflects the cellular state and can be treated

as a "pseudo-time" axis. Multiple branched trajectories were identified. For each

trajectory, we identified the GPNMB+ microglial coordinates on the pseudo-time

trajectories (Supplementary Figs. S7a, c).

Next, we analyzed the points at which the candidate Aβ-tau-moderating

factors related to the stress response predicted using MINDy were expressed in the

pseudo-time analysis. In both datasets, the moderator proteins were highly

expressed prior to GPNMB+ microglia in all pseudo-time trajectories (Supplementary

Figs. S7b, d). Similar results were obtained for *HSPA5*, a moderating factor based on

the amyloidome (i.e., the collection of amyloid peptides and related proteins, peptides,

and other molecules)[41] and tau interactome.[42] These results suggest that stress

factors moderating the Aβ-tau interaction may act through the activation of GPNMB+

microglia.

GPMNB protein expression is correlated with the clinical severity of AD

We evaluated the hypothesis that GPMNB+ microglia moderate Aβ-tau

interactions in early AD. In particular, we examined whether *GPMNB* could serve as a

biomarker for the clinical severity of AD.

For this purpose, we assessed brain tissue protein expression and clinical

parameters from the ROSMAP cohort data using CERAD and Braak scores as

indicators of the clinical severity of AD. We stratified the patients into two groups

based on *GPNMB* protein expression: the top 35% and bottom 35%. The *GPNMB*

high- and low-expression groups corresponded to the GPNMB+ microglia-activated

and non-activated groups, respectively. Within the ROSMAP cohort, the CERAD

score is a semi-quantitative measure of neuritic plaques, where 1 indicates definite

AD, 2 is probable AD, 3 is possible AD, and 4 represents no AD. The top 35% showed

greater disease progression based on both CERAD and Braak scores than that of the

bottom 35% (Figs. 7a, b).

GPNMB shows significant Aβ-tau interaction effects in the preclinical stage

We further examined whether *GPNMB* moderates the effect of the Aβ-tau

interaction in early-stage AD. To this end, we assessed the histopathological and AD

diagnostic information from the ACT study as well as gene expression data. We

tested interaction effects of *GPNMB* gene expression on Aβ and tau by linear

regression analyses (Fig. 7c, Supplementary Tables S7 and S8).

There were significant positive interaction effects of *GPNMB* gene expression

on Aβ-tau in the non-dementia group in the HIP and TCx brain regions (β = 0.089, p =

0.02 and β = 0.026, p < 0.001 in HIP and TCx, respectively). Notably, no significant

interaction effects were observed at the same sites in the dementia group.

Collectively, these findings suggest that GPNMB+ microglia and astrocyte moderate

the Aβ-tau interaction in the early stages of AD (Fig. 7d).

Establishment of an integrated network analysis tool, AlzPPMap

Finally, we developed a tool named AlzPPMap (Alzheimer's disease

Proteomics PPI Integrated Network Map) for the analysis and visualization of

integrated networks. This tool enables users to analyze the integrated functional

network derived from proteomics data alongside physical networks derived from PPIs.

Users can select one or more genes of interest and, as demonstrated in this study,

choose a specific percentage of closely related genes within the network based on

their proximity to the selected gene(s). The selected genes are then used to construct,

visualize, and detect communities within a subnetwork. AlzPPMap facilitates the

identification of molecules and molecular pathways that are both functionally and

physically associated with the target genes. The tool can be accessed at:

https://igcore.cloud/GerOmics/AlzPPMap.

Discussion

The number of individuals with AD is expected to continue increasing,[66]

making the development of effective treatments a pressing issue. As mentioned in the

Introduction, whereas Aβ and tau proteins are known to contribute to AD pathology,

the details of their involvement remain unclear.[1,2] The specific interactions between

them have been reported,[4–7] but not the underlying mechanisms. In this study, we

employed a deep learning-based network integration analysis to investigate the

mechanisms of the Aβ-tau interaction in patients with early-stage AD. The study was

conducted in four steps: (1) network integration, (2) identification of moderators and

subnetwork construction, (3) validation analysis using histopathological data, and (4)

development of a user-friendly web application.

In the first phase, we constructed an integrated functional-physical interaction

network where two different data types—protein co-expression and physical

PPIs—were combined into a single network using BIONIC.[30] Interestingly,

enrichment analysis and snRNA-seq data showed that the integrated network

captured relevant cell types and pathways. We also confirmed that Aβ and tau were

present in neighboring modules in the integrated network. Notably, neither

co-expression information nor physical interaction data alone could capture the

Aβ-tau interaction, emphasizing the need to evaluate functional and physical

interactions simultaneously. Co-expression network analyses are widely used in AD

research,[23–25] but it is difficult to represent physical interactions, such as Aβ-tau

interactions or the amyloidome. Therefore, an approach that integrates networks

based on different types of biological information may be effective in elucidating the

complex molecular pathogenesis of AD.

In the second phase, we used the MINDy[40] algorithm to search for factors

that moderate the Aβ-tau interaction using proteomics expression data. We identified

ER stress markers such as *HSPA5*, along with other stress response-related factors,

including *HSPD1*, *USP48*, *KBTBD6*, *UBE2D4*, *STAMBPL1*, and *MAP3K5*, as

potential moderators. Although ER stress responses have been frequently reported to

be associated with AD,[52–60] their detailed mechanisms remain unclear. In this

study, our findings suggested that stress response pathways were associated with

the Aβ-tau interaction, indicating that elucidating how these pathways influence the

progression of AD could be a focus for future research. In particular, understanding

how ER stress responses are involved in the Aβ-tau interaction and transmitted

through glial cells could contribute to the development of novel therapeutic strategies.

We next identified molecules, cell types, and pathways involved in the Aβ-tau

interaction by constructing subnetworks from the factors identified in the second

phase, extracting genes closely related to the two proteins within the integrated

network. The results suggested that glial cells, particularly GPNMB+ microglia, are

involved in the Aβ-tau interaction.

In the third phase, we validated the findings using an independent dataset

with histopathological data from the ACT study.[44] Through linear regression

analysis, we examined the interaction between GFAP, IBA1, and GPNMB with Aβ in

the hippocampus and temporal neocortex in the non-dementia group. Significant

positive interactions were detected, further suggesting that astrocytes and GPNMB+

microglia are moderators of the Aβ-tau interaction in early-stage AD. GPNMB+

microglia, a recently identified subtype, are elevated in the frontal white matter and

cortex of patients with AD.[67] Additionally, *GPNMB* is expressed in activated

microglia[68] and is associated with AD pathology.[22] Furthermore, the expression of

GPNMB in microglia is correlated with recently reported subtypes, such as

disease-associated microglia (DAM)[63] and microglial neurodegenerative phenotype

(MGnD),[69] which are closely associated with plaques and neurodegeneration.

However, their specific functions related to Aβ and tau have not been fully elucidated.

Our results provide new insights into the functions of GPNMB+ microglia in the

pathogenesis of AD.

In summary, network integration analysis has been extensively reported in

cancer research,[26–30] but its use in AD research remains limited. Through this

study, we were able to identify potential moderators of the Aβ-tau interaction by

integrating proteomics and PPI data using BIONIC.[30] Future analyses using a

framework like the one presented in this study, integrating other omics data, could

lead to new insights into the mechanisms of AD and other disease areas. Additionally,

in the last phase, we developed a tool named AlzPPMap to evaluate the integrated

network constructed in this study. This tool enables the analysis and visualization of

relationships between other genes, potentially providing new insights and ideas for

experimental validation.

This study had several limitations. First, except for the GPNMB+ microglial

data[22], all data were derived from postmortem brains. This introduces potential

confounding factors between postmortem effects and AD pathology. Second, in the

analyses summarized in Fig. 4, protein expression was aggregated by genes, not

accounting for multiple isoforms or post-translational modifications, especially in *APP*

and *MAPT*. For example, Aβ40 and Aβ42 are produced from *APP* through different

processes, and Tau in *MAPT* undergoes various phosphorylation modifications,[70]

impacting the disease differently. Histopathological evaluations were used to validate

the integrated protein network analysis results. Third, the *GPNMB* gene expression

analysis was based on bulk RNA-seq data, which may not specifically reflect

GPNMB+ microglia. A detailed analysis of the role of GPNMB+ microglia in AD

requires further experimental studies. Fourth, although we applied a significance

threshold of $P < 0.05$ with the Benjamini-Hochberg correction, there is ongoing

discussion in the field regarding the potential benefits of using a more stringent

threshold. Future studies with larger sample sizes could help to further support our

findings. Nonetheless, our findings offer valuable new insights into the Aβ-tau

interaction and highlight potential directions for the development of AD therapeutics.

List of abbreviations

Alzheimer's disease (AD), amyloid-beta (Aβ), disease-associated microglia (DAM), mild cognitive impairment (MCI), protein–protein interaction (PPI), Religious Order Study and Rush Memory and Aging Project (ROSMAP), single-nucleus RNA sequencing (snRNA-seq), tandem mass tag (TMT)

Declarations

Acknowledgments

We extend our thanks to all of the donors who generously contributed invaluable data for this study. We are also grateful to the researchers who made their data publicly available. The results published here are, in whole or in part, based on data obtained from the AD Knowledge Portal (https://adknowledgeportal.org). The study data were provided by the Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago. Data collection was supported through funding by NIA grants P30AG10161 (ROS), R01AG15819 (ROSMAP; genomics and RNAseq), R01AG17917 (MAP), R01AG30146, R01AG36042 (5hC methylation, ATACseq), RC2AG036547 (H3K9Ac), R01AG36836 (RNAseq), R01AG48015 (monocyte RNAseq) RF1AG57473 (single nucleus RNAseq), U01AG32984 (genomic and whole exome sequencing), U01AG46152 (ROSMAP AMP-AD, targeted proteomics), U01AG46161(TMT

proteomics), U01AG61356 (whole genome sequencing, targeted proteomics,

ROSMAP AMP-AD), the Illinois Department of Public Health (ROSMAP), and the

Translational Genomics Research Institute (genomic). Additional phenotypic data

were obtained from www.radc.rush.edu. Human AD brain single-nucleus RNA

sequencing (snRNA-seq) data were downloaded from the NCBI for Biotechnology

Information Gene Expression Omnibus (https://www.ncbi.nlm.nih.gov/geo/).

Microglial snRNA-seq data were accessed at https://braincelldata.org/resource and

https://doi.org/10.7303/syn2580853. Additionally, the human neuropathological and

RNA sequencing datasets used for the linear regression analysis are available at

NIAGADS (https://www.niagads.org/). Figures were created using Bio-Render.com.

Funding

This work was supported by the Human Glycome Atlas Project (HGA) and JSPS

KAKENHI Grant Number JP20H04282.

Author contributions

Conceptualization and Methodology: M.Y., K.A., Formal Analysis, Investigation, Data

Curation, Writing – Original Draft, Visualization: K.A., Resource, Writing – Review &

Editing, Supervision, Project Administration and Funding Acquisition: M.Y.

Human Ethics and Consent to Participate declarations

This study utilizes publicly available datasets obtained through appropriate

agreements or public data portals, ensuring compliance with ethical guidelines. The

proteomic data used in this study were accessed from the AD Knowledge Portal

(https://adknowledgeportal.org) under the terms of the data use agreement. These

datasets were collected with the informed consent of participants and ethical approval

from the original study institutions, including ROSMAP cohort studies. Additionally,

neuropathological and single-nucleus RNA sequencing (snRNA-seq) datasets used

in this study were obtained from the NCBI Gene Expression Omnibus (GEO) and

NIAGADS (https://www.niagads.org/), where all samples were collected following

approved institutional protocols, with participants providing informed consent for

research and data sharing. For data requiring restricted access, the necessary

approvals and permissions were obtained from the respective data providers. This

study adhered to all relevant ethical regulations for the analysis of secondary human

data.

Competing interests

The authors declare that there are no competing interests.

Availability of data and materials

All data used in this study are publicly accessible. The proteomic data can be

accessed via the AD Knowledge Portal (https://doi.org/10.7303/syn2580853). Human

AD brain snRNA-seq and microglia snRNA-seq datasets are available at the NCBI

Gene Expression Omnibus (GEO) under accession number GSE174367 and via the

following links: https://braincelldata.org/resource and

https://doi.org/10.7303/syn2580853. The human neuropathological and RNA

sequencing datasets employed for the linear regression analysis can be found at

https://www.niagads.org/datasets/ng00059. Additional information required to

reanalyze the data reported in this study is available from the lead author upon

request.

Code availability

The code used can be found on GitHub

(https://github.com/matsui-lab/AlzPPMap/tree/main).

References

1. d'Errico P, Meyer-Luehmann M. Mechanisms of Pathogenic Tau and Aβ Protein
Spreading in Alzheimer's Disease. Front Aging Neurosci [Internet]. 2020;12:265.

Available from: http://dx.doi.org/10.3389/fnagi.2020.00265

2. Knopman DS, Amieva H, Petersen RC, Chételat G, Holtzman DM, Hyman BT, et al.
Alzheimer disease. Nature Reviews Disease Primers [Internet]. 2021 [cited 2023 Nov
24];7:1–21. Available from: https://www.nature.com/articles/s41572-021-00269-y

3. Congdon EE, Ji C, Tetlow AM, Jiang Y, Sigurdsson EM. Tau-targeting therapies for
Alzheimer disease: current status and future directions. Nat Rev Neurol [Internet].
2023;19:715–36. Available from: http://dx.doi.org/10.1038/s41582-023-00883-2

4. Bloom GS. Amyloid-β and tau: the trigger and bullet in Alzheimer disease
pathogenesis. JAMA Neurol [Internet]. 2014;71:505–8. Available from:
http://dx.doi.org/10.1001/jamaneurol.2013.5847

5. Busche MA, Hyman BT. Synergy between amyloid-β and tau in Alzheimer's
disease. Nat Neurosci [Internet]. 2020;23:1183–93. Available from:
http://dx.doi.org/10.1038/s41593-020-0687-6

6. Zhang H, Wei W, Zhao M, Ma L, Jiang X, Pei H, et al. Interaction between Aβ and
Tau in the Pathogenesis of Alzheimer's Disease. Int J Biol Sci [Internet].
2021;17:2181–92. Available from: http://dx.doi.org/10.7150/ijbs.57078

7. Lee WJ, Brown JA, Kim HR, La Joie R, Cho H, Lyoo CH, et al. Regional Aβ-tau
interactions promote onset and acceleration of Alzheimer's disease tau spreading.
Neuron [Internet]. 2022;110:1932-1943.e5. Available from:
http://dx.doi.org/10.1016/j.neuron.2022.03.034

8. Aksman LM, Oxtoby NP, Scelsi MA, Wijeratne PA, Young AL, Alves IL, et al. A
data-driven study of Alzheimer's disease related amyloid and tau pathology
progression. Brain [Internet]. 2023;146:4935–48. Available from:
http://dx.doi.org/10.1093/brain/awad232

9. Josephs KA, Tosakulwong N, Weigand SD, Graff-Radford J, Schwarz CG, Senjem
ML, et al. Flortaucipir PET uncovers relationships between tau and amyloid-β in
primary age-related tauopathy and Alzheimer's disease. Sci Transl Med [Internet].
2024;16:eado8076. Available from:
https://www.science.org/doi/10.1126/scitranslmed.ado8076

10. van Dyck CH, Swanson CJ, Aisen P, Bateman RJ, Chen C, Gee M, et al.
Lecanemab in Early Alzheimer's Disease. N Engl J Med [Internet]. 2023;388:9–21.
Available from: http://dx.doi.org/10.1056/NEJMoa2212948

11. Bellaver B, Povala G, Ferreira PCL, Ferrari-Souza JP, Leffa DT, Lussier FZ, et al.

Astrocyte reactivity influences amyloid-β effects on tau pathology in preclinical Alzheimer's disease. Nat Med [Internet]. 2023;29:1775–81. Available from: http://dx.doi.org/10.1038/s41591-023-02380-x

12. Kim T, Yi D, Byun MS, Ahn H, Jung JH, Kong N, et al. Synergistic interaction of high blood pressure and cerebral beta-amyloid on tau pathology. Alzheimers Res Ther [Internet]. 2022;14:193. Available from: http://dx.doi.org/10.1186/s13195-022-01149-7

13. Coomans EM, van Westen D, Pichet Binette A, Strandberg O, Spotorno N, Serrano GE, et al. Interactions between vascular burden and amyloid-β pathology on trajectories of tau accumulation. Brain [Internet]. 2023; Available from: http://dx.doi.org/10.1093/brain/awad317

14. Yang H-S, Teng L, Kang D, Menon V, Ge T, Finucane HK, et al. Cell-type-specific Alzheimer's disease polygenic risk scores are associated with distinct disease processes in Alzheimer's disease. Nat Commun [Internet]. 2023 [cited 2023 Dec 7];14:1–13. Available from: https://www.nature.com/articles/s41467-023-43132-2

15. Shahidehpour RK, Nelson PT, Katsumata Y, Bachstetter AD. Exploring the link between dystrophic microglia and the spread of Alzheimer's neuropathology. Brain [Internet]. 2024 [cited 2024 Aug 18]; Available from: https://pubmed.ncbi.nlm.nih.gov/39101580/

16. Mathys H, Peng Z, Boix CA, Victor MB, Leary N, Babu S, et al. Single-cell atlas reveals correlates of high cognitive function, dementia, and resilience to Alzheimer's disease pathology. Cell [Internet]. 2023;186:4365-4385.e27. Available from: http://dx.doi.org/10.1016/j.cell.2023.08.039

17. Zhou Y, Song WM, Andhey PS, Swain A, Levy T, Miller KR, et al. Human and mouse single-nucleus transcriptomics reveal TREM2-dependent and TREM2-independent cellular responses in Alzheimer's disease. Nat Med [Internet]. 2020;26:131–42. Available from: http://dx.doi.org/10.1038/s41591-019-0695-9

18. Olah M, Menon V, Habib N, Taga MF, Ma Y, Yung CJ, et al. Single cell RNA sequencing of human microglia uncovers a subset associated with Alzheimer's disease. Nat Commun [Internet]. 2020;11:6129. Available from: http://dx.doi.org/10.1038/s41467-020-19737-2

19. Grubman A, Chew G, Ouyang JF, Sun G, Choo XY, McLean C, et al. A single-cell atlas of entorhinal cortex from individuals with Alzheimer's disease reveals cell-type-specific gene expression regulation. Nat Neurosci [Internet].

2019;22:2087–97. Available from: http://dx.doi.org/10.1038/s41593-019-0539-4

20. Mathys H, Davila-Velderrain J, Peng Z, Gao F, Mohammadi S, Young JZ, et al. Single-cell transcriptomic analysis of Alzheimer's disease. Nature [Internet]. 2019;570:332–7. Available from: http://dx.doi.org/10.1038/s41586-019-1195-2

21. Sun N, Victor MB, Park YP, Xiong X, Scannail AN, Leary N, et al. Human microglial state dynamics in Alzheimer's disease progression. Cell [Internet]. 2023;186:4386-4403.e29. Available from: http://dx.doi.org/10.1016/j.cell.2023.08.037

22. Gazestani V, Kamath T, Nadaf NM, Dougalis A, Burris SJ, Rooney B, et al. Early Alzheimer's disease pathology in human cortex involves transient cell states. Cell [Internet]. 2023;186:4438-4453.e23. Available from: http://dx.doi.org/10.1016/j.cell.2023.08.005

23. Neff RA, Wang M, Vatansever S, Guo L, Ming C, Wang Q, et al. Molecular subtyping of Alzheimer's disease using RNA sequencing data reveals novel mechanisms and targets. Science Advances [Internet]. 2021;7:eabb5398. Available from: https://www.science.org/doi/abs/10.1126/sciadv.abb5398

24. Johnson ECB, Carter EK, Dammer EB, Duong DM, Gerasimov ES, Liu Y, et al. Large-scale deep multi-layer analysis of Alzheimer's disease brain reveals strong proteomic disease-related changes not observed at the RNA level. Nat Neurosci [Internet]. 2022 [cited 2023 Nov 26];25:213–25. Available from: https://www.nature.com/articles/s41593-021-00999-y

25. Bai B, Wang X, Li Y, Chen P-C, Yu K, Dey KK, et al. Deep Multilayer Brain Proteomics Identifies Molecular Networks in Alzheimer's Disease Progression. Neuron [Internet]. 2020;105:975-991.e7. Available from: http://dx.doi.org/10.1016/j.neuron.2019.12.015

26. Cho H, Berger B, Peng J. Compact Integration of Multi-Network Topology for Functional Analysis of Genes. Cell Syst [Internet]. 2016;3:540-548.e5. Available from: http://dx.doi.org/10.1016/j.cels.2016.10.017

27. Chen G, Liu Z-P. Graph attention network for link prediction of gene regulations from single-cell RNA-sequencing data. Bioinformatics [Internet]. 2022 [cited 2023 Dec 4];38:4522–9. Available from: https://academic.oup.com/bioinformatics/article/38/19/4522/6663989

28. Schulte-Sasse R, Budach S, Hnisz D, Marsico A. Integration of multiomics data with graph convolutional networks to identify new cancer genes and their associated

molecular mechanisms. Nature Machine Intelligence [Internet]. 2021 [cited 2023 Dec 4];3:513–26. Available from: https://www.nature.com/articles/s42256-021-00325-y

29. Peng W, Tang Q, Dai W, Chen T. Improving cancer driver gene identification using multi-task learning on graph convolutional network. Brief Bioinform [Internet]. 2022;23. Available from: http://dx.doi.org/10.1093/bib/bbab432

30. Forster DT, Li SC, Yashiroda Y, Yoshimura M, Li Z, Isuhuaylas LAV, et al. BIONIC: biological network integration using convolutions. Nat Methods [Internet]. 2022;19:1250–61. Available from: http://dx.doi.org/10.1038/s41592-022-01616-x

31. ROSMAP_ProteomicsTMT_BatchCorrectionOC.pdf.

32. Perneczky R, Wagenpfeil S, Komossa K, Grimmer T, Diehl J, Kurz A. Mapping scores onto stages: mini-mental state examination and clinical dementia rating. Am J Geriatr Psychiatry [Internet]. 2006;14:139–44. Available from: http://dx.doi.org/10.1097/01.JGP.0000192478.82189.a8

33. Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, et al. STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. Nucleic Acids Res [Internet]. 2019;47:D607–13. Available from: http://dx.doi.org/10.1093/nar/gky1131

34. Veličković P, Cucurull G, Casanova A, Romero A, Liò P, Bengio Y. Graph Attention Networks [Internet]. arXiv [stat.ML]. 2017. Available from: http://arxiv.org/abs/1710.10903

35. Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, et al. Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. Cell [Internet]. 2015;161:1202–14. Available from: http://dx.doi.org/10.1016/j.cell.2015.05.002

36. Blondel VD, Guillaume J-L, Lambiotte R, Lefebvre E. Fast unfolding of communities in large networks. J Stat Mech [Internet]. 2008 [cited 2024 Jun 8];2008:P10008. Available from: https://iopscience.iop.org/article/10.1088/1742-5468/2008/10/P10008/meta

37. Morabito S, Miyoshi E, Michael N, Shahin S, Martini AC, Head E, et al. Single-nucleus chromatin accessibility and transcriptomic characterization of Alzheimer's disease. Nat Genet [Internet]. 2021;53:1143–55. Available from: http://dx.doi.org/10.1038/s41588-021-00894-z

38. Hao Y, Hao S, Andersen-Nissen E, Mauck WM 3rd, Zheng S, Butler A, et al.
Integrated analysis of multimodal single-cell data. Cell [Internet].
2021;184:3573-3587.e29. Available from: http://dx.doi.org/10.1016/j.cell.2021.04.048

39. Sayols S. rrvgo: a Bioconductor package for interpreting lists of Gene Ontology
terms. MicroPubl Biol [Internet]. 2023;2023. Available from:
http://dx.doi.org/10.17912/micropub.biology.000811

40. Wang K, Saito M, Bisikirska BC, Alvarez MJ, Lim WK, Rajbhandari P, et al.
Genome-wide identification of post-translational modulators of transcription factor
activity in human B cells. Nat Biotechnol [Internet]. 2009;27:829–39. Available from:
http://dx.doi.org/10.1038/nbt.1563

41. Andrew RJ, Fisher K, Heesom KJ, Kellett KAB, Hooper NM. Quantitative
interaction proteomics reveals differences in the interactomes of amyloid precursor
protein isoforms. J Neurochem [Internet]. 2019;149:399–412. Available from:
http://dx.doi.org/10.1111/jnc.14666

42. Drummond E, Pires G, MacMurray C, Askenazi M, Nayak S, Bourdon M, et al.
Phosphorylated tau interactome in the human Alzheimer's disease brain. Brain
[Internet]. 2020;143:2803–17. Available from:
http://dx.doi.org/10.1093/brain/awaa223

43. Csárdi G, Nepusz T. The igraph software package for complex network research.
https://www.semanticscholar.org › ..https://www.semanticscholar.org › . [Internet].
2006 [cited 2024 Jun 8]; Available from:
https://www.semanticscholar.org/paper/1d2744b83519657f5f2610698a8ddd177ced4
f5c

44. Miller JA, Guillozet-Bongaarts A, Gibbons LE, Postupna N, Renz A, Beller AE, et
al. Neuropathological and transcriptomic characteristics of the aged brain. Elife
[Internet]. 2017;6. Available from: http://dx.doi.org/10.7554/eLife.31126

45. Li P, Jiang Z, Liu T, Liu X, Qiao H, Yao X. Improving drug response prediction via
integrating gene relationships with deep learning. Brief Bioinform [Internet]. 2024;25.
Available from: http://dx.doi.org/10.1093/bib/bbae153

46. Young CB, Johns E, Kennedy G, Belloy ME, Insel PS, Greicius MD, et al. APOE
effects on regional tau in preclinical Alzheimer's disease. Mol Neurodegener [Internet].
2023;18:1. Available from: http://dx.doi.org/10.1186/s13024-022-00590-4

47. Verghese PB, Castellano JM, Garai K, Wang Y, Jiang H, Shah A, et al. ApoE

influences amyloid-β (Aβ) clearance despite minimal apoE/Aβ association in

physiological conditions. Proc Natl Acad Sci U S A [Internet]. 2013;110:E1807-16.

Available from: http://dx.doi.org/10.1073/pnas.1220484110

48. Stetler RA, Gan Y, Zhang W, Liou AK, Gao Y, Cao G, et al. Heat shock proteins:

cellular and molecular mechanisms in the central nervous system. Prog Neurobiol

[Internet]. 2010;92:184–211. Available from:

http://dx.doi.org/10.1016/j.pneurobio.2010.05.002

49. Das S, Ramakrishna S, Kim K-S. Critical Roles of Deubiquitinating Enzymes in

the Nervous System and Neurodegenerative Disorders. Mol Cells [Internet].

2020;43:203–14. Available from: http://dx.doi.org/10.14348/molcells.2020.2289

50. Genau HM, Huber J, Baschieri F, Akutsu M, Dötsch V, Farhan H, et al.

CUL3-KBTBD6/KBTBD7 ubiquitin ligase cooperates with GABARAP proteins to

spatially restrict TIAM1-RAC1 signaling. Mol Cell [Internet]. 2015;57:995–1010.

Available from: http://dx.doi.org/10.1016/j.molcel.2014.12.040

51. Ichijo H, Nishida E, Irie K, ten Dijke P, Saitoh M, Moriguchi T, et al. Induction of

apoptosis by ASK1, a mammalian MAPKKK that activates SAPK/JNK and p38

signaling pathways. Science [Internet]. 1997;275:90–4. Available from:

http://dx.doi.org/10.1126/science.275.5296.90

52. Ajoolabady A, Lindholm D, Ren J, Pratico D. ER stress and UPR in Alzheimer's

disease: mechanisms, pathogenesis, treatments. Cell Death Dis [Internet]. 2022

[cited 2023 Dec 7];13:1–15. Available from:

https://www.nature.com/articles/s41419-022-05153-5

53. Gorbatyuk MS, Gorbatyuk OS. The Molecular Chaperone GRP78/BiP as a

Therapeutic Target for Neurodegenerative Disorders: A Mini Review. J Genet Syndr

Gene Ther [Internet]. 2013;4. Available from:

http://dx.doi.org/10.4172/2157-7412.1000128

54. Yang Y, Turner RS, Gaut JR. The chaperone BiP/GRP78 binds to amyloid

precursor protein and decreases Abeta40 and Abeta42 secretion. J Biol Chem

[Internet]. 1998;273:25552–5. Available from:

http://dx.doi.org/10.1074/jbc.273.40.25552

55. Hoshino T, Nakaya T, Araki W, Suzuki K, Suzuki T, Mizushima T. Endoplasmic

reticulum chaperones inhibit the production of amyloid-beta peptides. Biochem J

[Internet]. 2007;402:581–9. Available from: http://dx.doi.org/10.1042/BJ20061318

56. Sakono M, Kidani T. ATP-independent inhibition of amyloid beta fibrillation by the
endoplasmic reticulum resident molecular chaperone GRP78. Biochem Biophys Res
Commun [Internet]. 2017;493:500–3. Available from:
http://dx.doi.org/10.1016/j.bbrc.2017.08.162

57. Fontaine SN, Rauch JN, Nordhues BA, Assimon VA, Stothert AR, Jinwal UK, et al.
Isoform-selective Genetic Inhibition of Constitutive Cytosolic Hsp70 Activity Promotes
Client Tau Degradation Using an Altered Co-chaperone Complement. J Biol Chem
[Internet]. 2015;290:13115–27. Available from:
http://dx.doi.org/10.1074/jbc.M115.637595

58. Moll A, Ramirez LM, Ninov M, Schwarz J, Urlaub H, Zweckstetter M. Hsp
multichaperone complex buffers pathologically modified Tau. Nat Commun [Internet].
2022 [cited 2023 Dec 7];13:1–13. Available from:
https://www.nature.com/articles/s41467-022-31396-z

59. Shao Y, Xu Y, Di H, Shi X, Wang Y, Liu H, et al. The inhibition of ORMDL3
prevents Alzheimer's disease through ferroptosis by PERK/ATF4/HSPA5 pathway.
IET Nanobiotechnol [Internet]. 2023;17:182–96. Available from:
http://dx.doi.org/10.1049/nbt2.12113

60. Gao H, Lei X, Ye S, Ye T, Hua R, Wang G, et al. Genistein attenuates memory
impairment in Alzheimer's disease via ERS-mediated apoptotic pathway in vivo and
in vitro. J Nutr Biochem [Internet]. 2022;109:109118. Available from:
http://dx.doi.org/10.1016/j.jnutbio.2022.109118

61. Castanho I, Murray TK, Hannon E, Jeffries A, Walker E, Laing E, et al.
Transcriptional Signatures of Tau and Amyloid Neuropathology. Cell Rep [Internet].
2020;30:2040-2054.e5. Available from: http://dx.doi.org/10.1016/j.celrep.2020.01.063

62. Braak H, Braak E. Neuropathological stageing of Alzheimer-related changes. Acta
Neuropathol [Internet]. 1991;82:239–59. Available from:
http://dx.doi.org/10.1007/BF00308809

63. Keren-Shaul H, Spinrad A, Weiner A, Matcovitch-Natan O, Dvir-Szternfeld R,
Ulland TK, et al. A Unique Microglia Type Associated with Restricting Development of
Alzheimer's Disease. Cell [Internet]. 2017;169:1276-1290.e17. Available from:
http://dx.doi.org/10.1016/j.cell.2017.05.018

64. Noda Y, Tsuruma K, Takata M, Ishisaka M, Tanaka H, Nakano Y, et al. GPNMB
Induces BiP Expression by Enhancing Splicing of BiP Pre-mRNA during the
Endoplasmic Reticulum Stress Response. Sci Rep [Internet]. 2017;7:12160.

Available from: http://dx.doi.org/10.1038/s41598-017-11828-3

65. Street K, Risso D, Fletcher RB, Das D, Ngai J, Yosef N, et al. Slingshot: cell
lineage and pseudotime inference for single-cell transcriptomics. BMC Genomics
[Internet]. 2018;19:477. Available from: http://dx.doi.org/10.1186/s12864-018-4772-0

66. 2024 Alzheimer's disease facts and figures. Alzheimers Dement [Internet].
2024;20:3708–821. Available from: https://pubmed.ncbi.nlm.nih.gov/38689398/

67. Satoh J-I, Kino Y, Yanaizu M, Ishida T, Saito Y. Microglia express GPNMB in the
brains of Alzheimer's disease and Nasu-Hakola disease. Intractable Rare Dis Res
[Internet]. 2019;8:120–8. Available from: http://dx.doi.org/10.5582/irdr.2019.01049

68. Hüttenrauch M, Ogorek I, Klafki H, Otto M, Stadelmann C, Weggen S, et al.
Glycoprotein NMB: a novel Alzheimer's disease associated marker expressed in a
subset of activated microglia. Acta Neuropathol Commun [Internet]. 2018;6:108.
Available from: http://dx.doi.org/10.1186/s40478-018-0612-3

69. Krasemann S, Madore C, Cialic R, Baufeld C, Calcagno N, El Fatimy R, et al. The
TREM2-APOE Pathway Drives the Transcriptional Phenotype of Dysfunctional
Microglia in Neurodegenerative Diseases. Immunity [Internet]. 2017;47:566-581.e9.
Available from: http://dx.doi.org/10.1016/j.immuni.2017.08.008

70. Wegmann S, Biernat J, Mandelkow E. A current view on Tau protein
phosphorylation in Alzheimer's disease. Curr Opin Neurobiol [Internet].
2021;69:131–8. Available from: http://dx.doi.org/10.1016/j.conb.2021.03.003

Figure legends

Fig. 1 A graphical abstract of the study

This study consists of the following four steps: 1. Integration of proteomics and protein–protein interaction (PPI) data, 2. Downstream analysis: Identification of moderators involving APP and MAPT, and construction of subnetworks, 3. Validation using pathological histological evaluation data, 4. Development of a user-friendly web application. In addition to step 2 conducted in this study, users can also construct and visualize subnetworks of genes of interest (including interactions between multiple genes).

Fig. 2 Characterization of integrated protein network modules

(a) Schematic representation of the workflow for integrating proteomics and protein–protein interaction (PPI) data and evaluation of integrated features.

(b) Stochastic neighbor embedding (tSNE) plot displaying protein modules.

(c) Upper panels: Uniform Manifold Approximation and Projection (UMAP) visualizations of single-nucleus RNA sequencing (snRNA-seq) data for patients with Alzheimer's disease (AD) (GSE174367)[37] clustered according to cell types (left) and conditions (right) Lower panels: expression levels of proteins within the identified modules.

(d) Representative Gene Ontology (GO) terms enriched in each protein module.

(e) Network depicting the closeness among modules. Closeness was defined by

averaging BIONIC-derived features within modules, calculating the Euclidean

distances among modules, and taking reciprocals.

Fig. 3 Inference of APP-MAPT modulators using a mutual information-based

method

(a) Graphical representation of the Modulator Inference by Network Dynamics

(MINDy) analysis.

(b) Scatter plot illustrating the relationships between –log10(p-values) and Scores.

Points representing p-values less than 0.05 are highlighted in red and are labeled

with their corresponding gene symbols.

Fig. 4 Glial cells moderate APP and MAPT interactions

(a) Workflow for network reconstruction from BIONIC analysis results, subnetwork

construction using proteins highly associated with intermediary factors, APP and

MAPT, and community detection within the network, followed by cell type

identification and enrichment analyses for each community.

(b) GO enrichment analysis of the subnetwork, categorized by parent terms using the

rrvgo[39] software.

(c) Representative GO terms enriched in each community.

(d) Dotplot displaying expression levels of proteins within each community.

Fig. 5 Histopathological analysis validated the roles of microglia/astrocytes as

Aβ-tau moderators in early Alzheimer's disease

(a) Schematic of the analytical workflow for linear regression analyses. Samples were

sourced from four brain regions: frontal white matter (FWM), hippocampus (HIP),

parietal neocortex (PCx), and temporal neocortex (TCx). RNA sequencing and

immunohistochemistry data were used.[44] A linear regression was performed,

repeated for GFAP and IBA1, and Aβ as predictors of tau interactions.

(b) Significant results of linear regression analyses for each of the four distinct brain

regions, with subjects categorized into no-dementia and dementia groups.

Fig. 6 snRNA-seq Analysis Identifying Enrichment for GPNMB+ Microglia in

Community 1

(a, c) Dot plot showing the expression levels of genes aggregated by community in

each microglial subtype.[21,22]

(b, d) Min-max normalized combined score for community 1, calculated by multiplying

the expression level by the proportion of expressing cells.

Fig. 7 GPNMB+ microglia as a possible moderator of the Aβ-tau interaction

(a, b) Heatmaps displaying the correlations between CERAD and Braak scores in

samples categorized by the top (a) and bottom (b) 35% of GPNMB expression.

1008   (c) Significant results of linear regression analyses for each of the four distinct brain

1009   regions, with subjects categorized into no-dementia and dementia groups.

1010   (d) Schematic representation summarizing the findings supporting the roles of both

1011   astrocytes and GPNMB+ microglia in the Aβ-tau interaction.

1012

1013

1014

Protein Expression and Protein-Protein Interaction Network Integration

MCI
Early AD

Correlation Network

+

PPI

Network Integration

Integrated Features

Downstream Analysis

Web Application

Exploration of Aβ-Tau Interaction Moderating Factor

Pair-wise distance

Modulator Inference

Sub-Network Reconstruction

Validation Analysis

User Input
- Gene symbols (up to 10)
- Percentage of closely related genes

APP   HSPA5   MAPT

Validation of Mediating Factor and Cell Subtype Analysis

RNA-seq   IHC   Linear Model

AlzPPMap: Alzheimer's disease Proteomics-PPI integrated network Map

This application allows you to analyze selected gene's subnetwork for the integrated network. Select a number and genes to visualize the integrated network. The background methodology is described in...

Network   Table   Graph Download

Select genes (up to 10):
HSPA5 APP MAPT

Select top % for HSPA5

Select top % for APP

Select top % for MAPT

RUN

Community 1
Community 3
Community 4
Community 2
Community 5
Community 6

a BIONIC Data Integration

ROSMAP study

MCI (n=97)
Early AD (n=25)

Correlation Network
+
PPI

BIONIC

Integrated Features

k-nearest neighbor algorithm
↓
Louvain algorithm
↓
Cell type Enrichment analysis
GObp Enrichment analysis

b

Cluster
1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28

t-SNE-2

t-SNE-1

c

Diagnosis

ASC
EX
INH
MG
ODC
OPC
PER.END

AD
Control

UMAP_2 / UMAP_1

Percent Expressed
25, 50, 75, 100

Average Expression
2, 1, 0, -1

Identity: PER.END, OPC, ODC, MG, INH, EX, ASC

Features: module_1 ... module_28

d

Cluster 5

GOBP AXON DEVELOPMENT
GOBP REGULATION OF MONOATOMIC ION TRANSMEMBRANE TRANSPORT
GOBP ENSHEATHMENT OF NEURONS
GOBP PROTEIN LOCALIZATION TO CELL PERIPHERY
GOBP GLIOGENESIS
GOBP GLIAL CELL DIFFERENTIATION
GOBP GLIAL CELL DEVELOPMENT
GOBP ACTION POTENTIAL
GOBP SISTER CHROMATID COHESION
GOBP TRANSMISSION OF NERVE IMPULSE
GOBP OLIGODENDROCYTE DIFFERENTIATION
GOBP MITOTIC SISTER CHROMATID COHESION
GOBP OLIGODENDROCYTE DEVELOPMENT
GOBP ESTABLISHMENT OF SISTER CHROMATID COHESION
GOBP AXON ENSHEATHMENT IN CENTRAL NERVOUS SYSTEM

Count
10, 15

p.adjust
0.00075
0.00050
0.00025

GeneRatio: 0.03 0.05 0.07 0.09

Cluster 11

GOBP SYNAPSE ORGANIZATION
GOBP REGULATION OF TRANS SYNAPTIC SIGNALING
GOBP AXON DEVELOPMENT
GOBP VESICLE MEDIATED TRANSPORT IN SYNAPSE
GOBP CELL JUNCTION ASSEMBLY
GOBP REGULATION OF NEURON PROJECTION DEVELOPMENT
GOBP REGULATION OF SYNAPSE STRUCTURE OR ACTIVITY
GOBP ACTIN FILAMENT ORGANIZATION
GOBP POSITIVE REGULATION OF CELL PROJECTION ORGANIZATION
GOBP SYNAPSE ASSEMBLY
GOBP POSTSYNAPSE ORGANIZATION
GOBP NEURON PROJECTION GUIDANCE
GOBP DENDRITE DEVELOPMENT
GOBP SYNAPTIC TRANSMISSION GLUTAMATERGIC
GOBP REGULATION OF SYNAPSE ASSEMBLY

Count
40, 60, 80, 100

p.adjust
4.087516e-24
3.065637e-24
2.043758e-24
1.021879e-24
2.818464e-65

GeneRatio: 0.10 0.15 0.20

e

Edge weight
1.0, 0.8, 0.6, 0.4, 0.2

1 — APOE Cluster
11 — APP Cluster
12 — MAPT Cluster

a

APP

Modulator

MAPT

Top 35%

Bottom 35%

$$\Delta MI = $$
$$|MI(MAPT_{Mtop35}, APP_{Mtop35}) -$$
$$MI(MAPT_{Mbottom35}, APP_{Mbottom35})|$$

$$MI(X, Y) = \sum x \in X \sum y \in Y p(x, y) \log(p(x)p(y)p(x, y))$$

Modulator low

MAPT   APP

Modulator high

MAPT   APP

b

Score vs -log10(p)

FLRT3

HSPD1

USP48

BRD4   RAB30

UBE2D4

RCAN1   KBTBD6

HSPA5

STAMBPL1

MAP3K5

-log10(p)

Score

● p.value < 0.05

a

BIONIC

Genes

Features by BIONIC

Inner Product

Build a Network Using the Top 5% of Inner Product Scores.

Subnetwork from top 5% closest gene connections; clustered by community detection

Public Amyloidome

Modulator Candidates from Fig.2

Public Tau Interactome

APP

HSPA5

MAPT

Calculate Euclidean distance

Cell type	Mechanism	
Oligodendrocyte	Neuron death, Stress response	APP
Neuron, Microglia	Kinase activity	MAPT, HSPA5
Microglia	Migration, Chemotaxis	
OPC, Astrocyte, Microglia	Neuron death, Kinase activity	GPNMB

b Enrichment Analysis

c

d

Community 1

Community 3

Community 4

Community 5

a

Brain samples
(FWM, HIP, PCx, TCx)

RNA-seq

Immunohistochemistry

Gene Expression Profiles

Samples

IHC scores

$$\text{Tau} = \beta_0 + \beta_1 A\beta + \beta_2 M + \beta_3(A\beta \times M) + \varepsilon$$

linear regression for the identification of intermediary factors

b

Hippocampus
(No Dementia)

Astrocyte

Amyloid beta

Interaction effect
$\beta=24.9$, $p=0.001$

Tau

Temporal cortex
(No Dementia)

Astrocyte

Amyloid beta

Interaction effect
$\beta=13.9$, $p<0.001$

Tau

Microglia

Amyloid beta

Interaction effect
$\beta=4.35$, $p<0.001$

Tau

a Heatmap of braaksc and ceradsc (Top 35% GPNMB Expression)

b Heatmap of braaksc and ceradsc (Bottom 35% GPNMB Expression)

c

Hippocampus (No Dementia)

GPNMB gene expression

Amyloid beta → Tau

Interaction effect β=0.089, p=0.02

Temporal cortex (No Dementia)

GPNMB gene expression

Amyloid beta → Tau

Interaction effect β=0.026, p<0.001

d MCI / early stage Alzheimer's disease

Amyloid beta

GPNMB+ Microglia

Astrocyte

Tau