Volatile Organic Compound Specialists in the Phycosphere

Authors: Vaishnavi G. Padaki[1], Xavier Mayali[2], Peter K. Weber[2], Stephen J. Giovannoni[1],

Kaylene Abraham[1], Kerry Jacobs[1], Lindsay Collart[1], Kimberly H. Halsey[1*]

Author Affiliations:

[1] Department of Microbiology, Oregon State University

[2] Lawrence Livermore National Laboratory, Livermore, CA 94550 USA

[*]corresponding author

Abstract:

Labile dissolved organic carbon (LDOC) in the oceans accounts for ~ ¼ of global photosynthesis
and turns over with a half-life of about one day, fueling one of the largest engines of microbial
heterotrophic production on the planet. Volatile organic compounds (VOCs) are poorly
constrained components of LDOC. We detected 78 VOCs totaling 18.5 µM in cultures of the
model diatom *Phaeodactylum tricornutum*, including hydrocarbons usually found in petroleum.
In five individual cocultures with bacteria adapted to grow with this diatom, 1 to 66 VOCs were
depleted. Two of the most active VOC consumers, *Marinobacter* and *Roseibium,* had more VOC
oxidation genes, and attached to the diatom, suggesting VOC specialism. Diatom photosynthesis
increased by up to 29% in the presence of VOC consumers, indicating that VOC consumption by
heterotrophic bacteria in the phycosphere – a region of rapid organic carbon oxidation that
surrounds phytoplankton cells – could impact global rates of primary production.

Main:

The phycosphere, the layer of water immediately surrounding an algal cell, is the site of complex

interactions between phytoplankton and bacterioplankton, which influence carbon and nutrient

cycling in marine ecosystems [1–3]. Microbial activities in the phycosphere are distinct from

processes occurring in bulk water, a result of bacterial taxa with specialized adaptations

populating the region of concentrated chemicals surrounding phytoplankton [4,5]. Direct chemical

transfer from phytoplankton to bacterioplankton is enhanced by their occupation of the narrow

zone of the phycosphere, which allows access to metabolites before they diffuse into the bulk

seawater [2,6]. The ecology of the phycosphere is similar to its terrestrial analog, the rhizosphere;

phylogenetically related bacterial taxa are found across both systems [7], chemotaxis plays a

central role in accessing the narrow zone of high chemical concentration [8,9], and the chemicals

transferred from the primary and secondary producers are functionally similar [2]. For example,

primary metabolites, such as sugars, amino acids, and vitamins, as well as metabolite precursors,

such as organosulfur compounds (dimethylsulfide (DMS) and dimethylsulfoniopropionate) [10,11],

are used as bacterial growth substrates in the phycosphere and rhizosphere [12,13].

Volatile organic compounds (VOCs) are an important subset of the labile dissolved

organic compounds (LDOC) produced by phytoplankton, but VOCs are often overlooked in

metabolomic studies because most are not captured by common methods of analysis such as

solid phase extraction and liquid chromatography. VOCs are relatively low in molecular weight,

hydrophobic, and can diffuse across cell membranes. Some VOCs, such as isoprene and DMS,

are produced by phytoplankton in response to stress [14–16], while others (e.g., acetaldehyde and

methyl iodide) are intermediate chemicals in metabolic pathways [17,18]. Due to their diffusive

properties, these VOCs may escape biochemical transformation and cross the algal cell

membrane into the phycosphere. Other VOCs, including aromatic benzenoids such as benzene,

toluene, and ethylbenzene/xylene (collectively known as BTEX), are produced by phytoplankton

via the shikimate and non-mevalonate pathways [19,20], but the specific reactions involved in

BTEX formation are not known.

Research on VOC transfer between algae and bacteria is sparse but suggests VOCs

released by algae can be growth substrates for marine bacteria. A wide range of VOCs were

detected during the growth of the diatom, *Thalassiosira pseudonana*, and a subset of those

supported the growth of *Pelagibacter ubique* HTCC1062, a member of ubiquitous SAR11

bacterioplankton [18,21]. These publications suggested that VOCs were "public goods" released by

the diatom constitutively and available to any organism able to metabolize them. HTCC1062

metabolizes isoprene, acetone, acetaldehyde, DMS, and cyclohexanol, which are some of the

most well-studied VOCs for their roles in sea-air exchange and atmospheric chemistry [18]. BTEX

are produced by some algae [19], and are growth substrates for a wide variety of bacteria isolated

from oilfields [22,23], soils [24], groundwater [25], and wastewater [26]. Less is known about BTEX

consumption by marine bacteria, which is likely an important sink that limits BTEX emissions

into the atmosphere [27]. Genetic mechanisms underlying bacterial metabolism of some VOCs

(e.g., methanol, DMS, acetone, acetaldehyde) have been elucidated [18,28,29]. The widespread

expression of some of the genes involved, such as acetone/cyclohexanone monooxygenase,

suggests marine bacteria have an important role in mediating VOC sea-air emissions [18].

However, the full diversity of VOCs used by bacteria in the oceans is unknown.

The bacterial community in the phycosphere of marine microalgae may be uniquely

adapted to VOC metabolism, as is the case for methylotrophs in the phyllosphere [30]. The

phycosphere of a modest-sized (~20 μm) diatom extends about 50 – 2000 μm away from the cell

[2]. In this region, chemicals are modeled to be $\geq$50% higher than in bulk seawater [2], where VOC

concentrations are in the pM to nM range. VOC concentration in the phycosphere is partly

controlled by the rate of diffusion from the algal source. The rate of diffusion decreases with

increased molecular size and decreased hydrophobicity and likely increases with phytoplankton

growth rate and bacterial VOC uptake. Some bacteria occupy the phycosphere through

chemotaxis or physical attachment to algal cells [31,32]. Chemotactic bacteria experience transient

exposures to a dynamic organic chemical cocktail in the phycosphere [12], where the composition

depends on the algal producer [1,33], environment [34,35], and antagonistic and mutualistic

interactions [12,36,37], requiring bacterial consumers to efficiently shift physiologies to match

resource availability. Strategies bacteria use to navigate carbon acquisition as they traverse

phycosphere boundaries are not known but may depend on pool sizes and the diversity of

regulatory behavior systems for the uptake and metabolism of various substrates [38]. Bacteria that

can adhere to algae would, in principle, optimize diffusion-mediated VOC uptake by maximizing

areal coverage of VOC flux directly from the algal source.

Here, we investigated VOC transfer from the model diatom, *Phaeodactylum tricornutum,*

to phycosphere bacteria previously isolated from a large-scale *P. tricornutum* production pond.

The bacteria varied in taxonomy, DOC uptake, and attachment to the diatom [12,39,40]. Results

show significant VOC transfer from *P. tricornutum* to its phycosphere bacteria, fueling primary

and secondary production. Extrapolated to larger scales, our findings support the perspective that

phycosphere bacteria are a sink for up to 29% of primary production and play an important role

in limiting VOC accumulation in the surface ocean.

Results:

Phaeodactylum tricornutum produced a wide range of VOCs in culture

During exponential growth of *P. tricornutum*, 78 m/z signals were detected by proton

transfer reaction mass spectrometry (PTR-MS) at concentrations higher than the media control

(Benjamini-Hochberg, Q-value < 0.1, n = 6). VOCs accumulated to a total of 85.1 pM cell^{-1} and

represented nine chemical functional groups (Fig. 1). Hydrocarbons represented about 60% of

the VOCs produced in the exponential phase (54.2 pM cell^{-1}; Supp. table 1). The benzenoid class

of aromatic hydrocarbons was most abundantly produced, including BTEX compounds (m/z

79.11, m/z 93.07, and m/z 107.08), respectively [19], and $C_{11}H_{16}^{+}$ (m/z 149.13). Note that in PTR-

MS, VOCs are detected at their protonated ion masses. Alcohols were the second most

abundantly produced group of VOCs, contributing up to 11% of the total VOC pool (10.20 pM

cell^{-1}). Methanol (m/z 33.03) and, likely, ethanol (m/z 47.04) made up the vast majority of the

accumulated alcohols (Fig.1, Supp. table 1). Consistent with reports that *P. tricornutum* is a

halocarbon producer [41,42], m/z 124.96, corresponding to ethyl bromide, was detected in the

exponential phase. The well-studied VOCs acetone (m/z 59.05), acetaldehyde (m/z 45.03), and

DMS (m/z 68.07) were also detected in the exponential phase (Fig. 1, Supp. table 1).

After *P. tricornutum* entered the stationary growth phase, 58 m/z signals were detected in

concentrations higher than the media control (Benjamini-Hochberg, Q-value < 0.1, n = 6) and

accumulated to 9.01 pM cell^{-1} (Fig. 1), which was only about 10% of the VOC accumulation

during the exponential phase (Fig. 1, Supp. table 1). VOCs detected in both phases of growth

were consistently lower in concentration in the stationary phase. Slower VOC production in

response to cessation of growth, and loss of VOCs from the vented growth flasks, are the most

likely explanations for the concentration differences between growth phases. Hydrocarbons still

represented the majority of the VOCs (84%) but were only 7.59 pM cell^{-1}. Sixteen of the 58 m/z

signals were unique to the stationary phase, but many of these had unknown identities.

The most abundant VOC in both exponential and stationary phases was $C_{11}H_{16}^+$ (m/z

149.13) in the benzenoid superclass and corresponds to dictyopterene, an algal sesquiterpenoid

pheromone. Chemical isomers of dictyopterene include the aromatic hydrocarbons, neopentyl-

benzene, 4-tert-butyl-toluene, and 1-ethyl-2-propylbenzene, which have been identified in plant

leaves [43]. Forty-two m/z signals, including BTEX, were detected in both exponential and

stationary phases (Fig. 1).

VOCs were depleted in P. tricornutum-bacteria cocultures

Five *P. tricornutum*-bacteria cocultures were grown to identify VOCs produced by the

diatom that decreased in concentration in the presence of a bacterium, which would suggest

those VOCs were consumed by the bacterium. Alternatively, decreased VOC concentrations in a

coculture could indicate VOC production was inhibited in the presence of the bacterium. The

five *P. tricornutum*-bacteria cocultures (herein "PT-bacteria genus name") each showed distinct

VOC depletion patterns during exponential growth when compared to axenic *P. tricornutum*

(Fig. 2a, Supp. Fig. 1; PERMANOVA, p-value = 0.03, n = 6). Of the 78 m/z signals produced by

P. tricornutum in exponential phase, 31 to 59 were significantly depleted in four of the five

cocultures (Benjamini-Hochberg, Q-value cut-off ≤ 0.1, n = 6) (Fig. 2a). VOCs were most

strongly depleted in PT-*Marinobacter* (7.09 pM cell^{-1}), followed by PT-*Stappia* (6.51 pM cell^{-1}),

PT-*Roseibium* (5.52 pM cell^{-1}) and PT-*Rhodophyticola* (4.54 pM cell^{-1}) (Fig. 2a, Fig. 4, Supp.

table 1). About half the number of m/z signals were depleted in the PT-*Roseibium* compared to

PT-*Marinobacter*. *Marinobacter* and *Roseibium* were observed to be attached to *P. tricornutum*

(Fig. 3) [39]*, thus the cell-normalized VOC depletion data in PT-*Marinobacter* and PT-*Roseibium*

are based on estimates of bacteria cell densities using flow cytometry for the free-living and

scanning electron microscopy for the attached bacteria (see Methods). In PT-*Yoonia*, only one

m/z signal, 45.03, corresponding to acetaldehyde, was significantly depleted (0.5 pM cell^{-1} (Fig.

2a)). There were 14 m/z signals that did not show significant depletion in any of the five PT-

bacteria cocultures. A lack of depletion could mean that the bacteria could not take up or

metabolize those VOCs, or those VOCs were maintained at a steady state concentration by equal

rates of production and consumption in the coculture. Of those, 8 m/z signals belonged to the

nitrogen-containing class of VOCs (Supp. table 6). Furthermore, neither methanol nor acetone

were depleted in any of the five cocultures.

Hydrocarbons were the most highly depleted group of VOCs in all PT-bacteria cocultures

except PT-*Yoonia*. Depending on the bacterium present, 11-19 of the depleted m/z signals were

classified as hydrocarbons and made up to 70% of the total VOC pool depleted in the cocultures.

PT-*Marinobacter* depleted the most hydrocarbons (5.95 pM cell^{-1}), followed by PT-*Stappia*

(5.44 pM cell $^{-1}$), PT-*Roseibium* (4.48 pM cell^{-1}) and PT-*Rhodophyticola* (3.94 pM cell $^{-1}$) (Fig.

2a). Monoaromatic hydrocarbons, including BTEX, $C_{11}H_{16}^{+}$, and phenylacetylene (m/z 103.05),

were the most strongly depleted hydrocarbons in the PT-bacteria cocultures (Fig. 2a).

In stationary growth phase, the range of depleted VOCs in the PT-bacteria cocultures was

lower, from 0.64-4.97 pM cell^{-1} (average 2.96 ± 0.66 pM cell^{-1}). The depletion patterns between

PT-bacteria cocultures in stationary phase were not distinct from one another (Fig. 2b,

PERMANOVA p-value = 0.22, n = 6). In total, 24 to 55 m/z signals were depleted in the PT-

bacteria cocultures (Benjamini-Hochberg, Q-value < 0.1, n = 6). The number of m/z signals

depleted in PT-*Yoonia* expanded from 1 in the exponential phase to 24 in the stationary phase,

but the amounts of the individual VOCs depleted were low (Fig. 2b, Supp. Table 2 and 3). In

stationary phase, 9-16 m/z signals depleted in PT-bacteria cocultures were classified as

hydrocarbons, and these VOCs were the majority of the depleted VOC pools (80-92%). VOCs

classified as carboxylic acids were the second-most depleted group, followed by ketones (Fig. 2a

and b). Except for hydrocarbons, the magnitude of VOC depletion in PT-bacteria cocultures did

not mirror VOC abundances in axenic *P. tricornutum* in either growth phase (Fig. 1, Supp. tables

1, 2, and 3).

VOC transfer to bacteria comprised up to 30% of primary production

P. tricornutum growth rates, chlorophyll content, and photosynthetic efficiencies were

unaffected by the presence of any of the bacterial species in the PT-bacteria cocultures (Supp.

Table 4). To assess the impact of bacterial VOC uptake on *P. tricornutum* primary production,

short-term $^{14}CO_2$-uptake rates, approximating gross carbon production (GCP) were measured.

GCP in PT-bacterial co-cultures were higher in PT-*Marinobacter,* PT-*Stappia,* and PT-

Rhodophyticola than in axenic *P. tricornutum* (Fig. 4)*.* A hydrocarbon trap that stripped VOCs

from the axenic *P. tricornutum* culture, simulating a highly efficient bacterial VOC sink, caused

GCP to increase 24.9% compared to axenic *P. tricornutum* with no trap (Fig. 4; $p = 0.025$, n =

6). GCP in PT-*Marinobacter,* PT-*Rhodophyticola,* and PT-*Stappia* were 20.1% to 29.3% higher

than in axenic *P. tricornutum,* similar to the hydrocarbon trap-control, and corresponding to 0.27

to 0.41 nmol C cell^{-1} h^{-1} transferred to the bacteria in the form of VOCs (p-values < 0.02, t-tests,

n=3) (Fig. 4). Consistent with the minimal amount of VOC depleted in PT-*Yoonia* there was no

significant difference in CO_2-uptake rates between PT-*Yoonia* and axenic *P. tricornutum* (p-

values 0.23, t-test, n=3). Interestingly, GCP was not stimulated in PT-*Roseibium* until *Roseibium*

entered the second exponential growth phase in its diauxic growth pattern (Fig. 2; note lag

between days 2 and 4). This result suggests that *Roseibium* shifted carbon uptake from primarily

LDOC early in culture growth to primarily VOCs later in culture growth.

VOC depletion correlates with genome content for hydrocarbon metabolism

Of the five bacteria tested*, Marinobacter* and *Roseibium* genomes harbor the greatest

range of hydrocarbon metabolism genes. For example, genes that initiate hydrocarbon oxidation

were identified in *Marinobacter* and *Roseibium,* including *rhdA,* encoding ring hydroxylating

dehydrogenase, *almA*, encoding flavin-binding monooxygenase involved in long-chain alkane

degradation, and *alkB,* encoding alkane monooxygenase. *rhdA* is in a superclass of nonheme iron

enzymes that convert aromatic hydrocarbons to dihydrodiols in the presence of dioxygen and

NADH [44]. The broad substrate range for *rhdA* includes benzene, and the benzene-alkyl

substituted hydrocarbons toluene, ethylbenzene, and xylene, consistent with BTEX depletion

profiles in *Marinobacter* and *Roseibium*. Benzene dioxygenases are a subclass of *rhdA* that

oxidize benzene and toluene to their corresponding dihydrodiols. We manually curated the *rhdA*

hits for *bnzA*, encoding the α-subunit of the benzene 1,2 dioxygenase hydroxylase based on the

conservation of the active site motif Cys-X_1-His-15-to-17 aa-Cys-X_2-His (where X is any amino

acid) [45]. *bnzA* was identified in *Marinobacter*, *Rhodophyticola*, and *Roseibium* (Fig. 2d).

Marinobacter also encodes *bnzB,* the β-subunit of benzene 1,2 dioxygenase hydroxylase (Fig.

2d). The presence of both *bnzA* and *bnzB* subunits increases hydroxylation activity compared to

the activity of either of the subunits alone [46]. Genes facilitating hydrocarbon oxidation in *Stappia*

were not identified. Acetone/cyclohexanone monooxygenase (*acmA*) was identified in

Marinobacter, Roseibium, and Stappia. The broad substrate range of *acmA* likely includes cyclic

ketones, such as valerophenone (m/z 163.11) and E-jasmone (m/z 165.13), corresponding to m/z

signals that were depleted in the PT-*Marinobacter*, PT-*Roseibium*, and PT-*Stappia*. All five

bacteria encode one or two copies of aldehyde dehydrogenase (*aor*), consistent with

acetaldehyde (m/z 45.03) depletion in all five cocultures (Fig. 2).

Marinobacter and *Rhodophyticola* exhibit distinct benzene metabolism

We selected benzene as a representative hydrocarbon to directly test incorporation into

biomass by *Marinobacter* and *Rhodophyticola*. These bacteria differed in the amounts of

benzene depleted in the PT-bacteria cocultures, in their genetic capacities for hydrocarbon

metabolism and chemotaxis, in their attachment physiologies (Fig. 2, 3, 5) [39], and could be

grown without *P. tricornutum* in media prepared from filtrate collected from *P. tricornutum* in

exponential growth ('PTspent'; see methods). *Marinobacter* and *Rhodophyticola* grew to higher

abundances in PT spent medium with added benzene vs. no added benzene (Fig. 5).

Benzene uptake and subsequent incorporation into biomass by *Marinobacter* and

Rhodophyticola were quantified using nanoSIMS. We incubated each bacterium with and

without ^{13}C benzene (60 μM in *Marinobacter* cultures, 36 μM in *Rhodophyticola* cultures) and

^{15}N leucine (50 nM) in f/2+Si artificial seawater medium (ASW), PT spent medium, and in

coculture with *P. tricornutum*. *Marinobacter* increased its growth in ASW with benzene added

as the sole source of carbon (Fig. 5a). Associated C_{net} values of incorporation, measuring the

fraction of a cell's C originating from benzene, increased over the incubation period, reaching

0.03 to 0.15 (average 0.065) on day 7 (Fig. 5b). *Marinobacter* reached higher cell densities in

PTspent with added benzene compared to PTspent with no benzene or ASW with benzene (Fig.

5a (p-value <0.05, n = 3). Consistent with benzene-stimulated growth in PTspent, *Marinobacter*

C_{net} values in that treatment ranged from 0.02 to 0.17, averaging 0.07 on day 4. *Marinobacter*

cell densities were unaffected by benzene addition to PT-*Marinobacter*. Surprisingly,

Marinobacter did not incorporate ^{13}C benzene in the presence of live *P. tricornutum* cells.

Marinobacter C_{net} values in the coculture hovered around zero across the 10-day incubation,

even in the presence of 60 μM externally-supplied benzene (Fig. 5a,b). Thus, *Marinobacter* did

not incorporate measurable quantities of ^{13}C-benzene in the presence of live *P. tricornutum* cells.

N_{net} values from leucine in all *Marinobacter* treatments, including the coculture, were low but

significant (<0.01, Supp. Fig. 5), confirming bacterial activity.

Despite significantly higher *Rhodophyticola* cell densities in PTspent with 36 μM

benzene added compared to the no benzene-added control (Fig. 5c). ^{13}C-benzene addition to

PTspent resulted in low C_{net} values, and only 2% of the population showed enrichment in ^{13}C

benzene on day 6 (Fig. 5d). *Rhodophyticola* may take up and oxidize benzene for energy rather

than biomass generation. No significant difference was observed in *Rhodophyticola* cell densities

grown in the PT coculture with or without added ^{13}C-benzene. Similar to *Marinobacter*, C$_{net}$

values in *Rhodophyticola* grown in the PT coculture were very low. *Rhodophyticola* showed

significant ^{15}N enrichment from leucine in both PTspent and PT-*Rhodophyticola* treatments

(Supp. Fig. 5).

Discussion

Phycospheres are recognized as energy-rich regions of heterotrophic activity where

microbes adapted to this niche benefit from proximity to high concentrations of the organic

carbon resources produced by phytoplankton cells. VOCs are rapidly diffusing molecules that

easily traverse cell membranes and, due to the introduction of new measurement technologies,

are increasingly recognized as complex and important products of phytoplankton metabolism.

Here, we show that the phycosphere concept applies to VOCs and that they appear to be an

important component of photosynthetic carbon transfer to heterotrophic microbial populations in

the region immediately surrounding phytoplankton cells. We also show that some phycosphere

bacteria have traits that can be interpreted as evidence of VOC specialism, enabling them to

access these chemicals whose concentrations are expected to rapidly decrease with distance from

their source.

VOCs were primarily produced during active *P. tricornutum* growth, when over a quarter

of photosynthetically fixed carbon was emitted from *P. tricornutum* as VOCs. Hydrocarbons

were 64% of the total VOC produced, with BTEX accounting for 13% of the emitted VOC pool,

accumulating to 11.8 µM. BTEX production by phytoplankton has been proposed to be of

sufficient magnitude to influence atmospheric chemistry [19], unless bacterial VOC consumption

prevents BTEX accumulation in the surface ocean [27]. The VOC concentrations we measured

could be underestimated because the cultures were grown in vented flasks. However, in the

absence of bacterial sinks, internal VOC concentrations in axenic *P. tricornutum* cells may reach

equilibria with the surrounding medium.

We measured concentrations of VOCs in diatom-bacteria cocultures relative to diatoms

alone to investigate the potential for VOC uptake by bacteria. In the case of benzene, isotopic

labeling confirmed that the bacteria were consumers, but for other VOCs, we cannot eliminate

the possibility that VOC depletion in coculture was caused by inhibition of VOC production by

P. tricornutum. *P. tricornutum* photosynthesis (GCP) was stimulated in cocultures where

depletion of a wide range of VOCs was also observed. Some marine bacteria are known to

metabolize VOCs, including BTEX [21,47], and we show that the bacteria that caused VOC

depletion also harbor genes for some of the depleted compounds; for other depleted compounds,

genetic loci associated with consumption are unknown and cannot be ruled out.

The data we report here indicate a range of VOC specialism (Table 1). Four of the five

bacterial strains harbor multiple hydrocarbon oxidation genes that encode enzymes with broad

substrate ranges. *Marinobacter* and *Roseibium,* which consumed large amounts of hydrocarbons,

had the most genes for hydrocarbon oxidation. *Marinobacter* and *Roseibium* encode few genes

for complex polysaccharide metabolism, and in previous research, incorporated low amounts of

macromolecular DOC into biomass [39]. In our experiments, *Rhodophyticola* and *Stappia,* which

harbor fewer hydrocarbon oxidation genes, depleted smaller amounts of individual hydrocarbons

but a wide range. Mayali et al. reported that *Stappia, Rhodophyticola,* and *Yoonia* used complex

DOC in the 2023 study, in agreement with their genetic capacities for oligosaccharide, starch,

and cellulose degradation. Thus, for this set of traits, among a set of five heterotrophic isolates

from *P. tricornutum* ponds, two strains appeared to be VOC specialists: they used more VOCs

and less carbohydrate DOM, and as discussed below, they were motile, and attached to *P.*

tricornutum. Two strains used lesser amounts of VOCs, are active metabolizers of carbohydrate

DOM, are not motile, and did not attach; and one strain used carbohydrate DOM, only a single

VOC, and attached.

We looked for evidence of phycosphere occupation to understand whether, in the strains

we studied, it was linked to VOC use. The VOC specialists, *Marinobacter* and *Roseibium,* and

one strain that did not use VOCs, *Yoonia*, attached to *P. tricornutum,* but the other strains did

not. *Marinobacter* attached to the diatom in a flat orientation, while *Roseibium* attached to *P.*

tricornutum using polar adhesion (Supp. Fig. 2). It has been reported that chemotaxis and

motility are key traits of hydrocarbon-degrading bacteria. Volatile aromatic hydrocarbons,

including toluene and benzene derivatives, have been shown in previous work to be effective

bacterial chemoattractants, even in the presence of complex organic matter, such as petroleum,

crude oil, and diesel [48–50]. *Marinobacter* and *Roseibium* encode canonical chemotaxis *che* genes

[39], but chemoattractant compounds have not been identified in these bacteria [8,51] (Table 1).

One of the most surprising results in this study was the absence of [13]C-benzene cellular

incorporation in *Marinobacter* grown in coculture with *P. tricornutum,* despite strong [13]C-

benzene incorporation in *Marinobacter* grown in the absence of *P. tricornutum* (i.e., in PTspent

and ASW) and significant [12]C-benzene depletion in PT-*Marninobacter.* We repeated the

experiment and confirmed the results. These unexpected results suggest that in coculture with *P.*

tricornutum, Marinobacter obtained benzene, and possibly other hydrocarbons, directly from the

diatom rather than from the bulk medium. *Marinobacter* and *Roseibium* were sometimes

observed attached to *P. tricornutum.* Bacterial attachment or entry into the phycosphere [8] could

facilitate efficient VOC transfer directly from the diatom to the bacteria, before the chemicals

diffuse into the bulk medium. In coculture, direct VOC uptake from the diatom to *Marinobacter*

may occur via their physical interaction, enabling passive VOC diffusion across the diatom and

bacterial membranes. However, FadL outer membrane transporters can boost BTEX metabolism

in some Gram-negative hydrocarbon degraders [52], and the *Marinobacter* strain used in this study

encodes a FadL protein with 72% nucleotide identity to the well-characterized FadL in

Escherichia coli. In the presence of the diatom, *Marinobacter* may regulate FadL to optimize use

of a wide range of substrates delivered from *P. tricornutum*. However, FadL substrate specificity

and kinetics are not yet known, and FadL may become saturated, inhibiting BTEX uptake (i.e.,

added ^{13}C-benzene in our experiment) from the bulk medium. Bacteria also exhibit varying

preferences for BTEX substrates [53] and the benzenoid collection of hydrocarbons. Benzene may

be of lower value to *Marinobacter* compared to the full collection of VOC substrates from

growing *P. tricornutum,* or benzene may be oxidized to CO_2 in the coculture. Nevertheless, the

contrasting ^{13}C labeling results in *Marinobacter* suggest attachment dynamics in the *P.*

tricornutum phycosphere may have an important role in VOC uptake.

In the model set of phytoplankton and associated bacteria we investigated, phycosphere

resources were exploited by bacteria employing varying strategies: VOC specialists

(*Marinobacter* and *Roseibium*) that were motile and attached, generalists capable of VOC and

macromolecular uptake (*Rhodophyticola* and *Stappia*), and macromolecular specialists (*Yoonia*).

We propose that VOC specialists use chemotaxis to find and colonize the phycosphere, where

proximity to sources of low molecular weight, rapidly diffusing VOCs offers advantages [8].

Unknown is the duration of attachment or how hydrocarbon uptake is differentially regulated in

the presence and absence of *P. tricornutum*. The free-living generalists, *Rhodophyticola* and

Stappia, used VOCs and carbohydrate DOC and had fewer hydrocarbon oxidation genes.

Collectively, VOC use by phycosphere bacteria was surprisingly broad in the range of

compounds used and accounted for up to 29% of primary production by *P. tricornutum.* These

findings add support to previous reports of VOC uptake by bacteria and stimulation of

photosynthesis in phytoplankton co-cultured with VOC oxidizers. Previous reports show that

balanced production and consumption maintain low VOC concentrations in the oceans and have

associated VOC consumption with specialized metabolism in abundant, streamlined, non-motile

cells [54–57] but in the set of bacteria we studied some of the most active VOC utilizers were motile

and attached to the phytoplankton. Here, we show that the flux can be intercepted by VOC

specialists in the phycosphere, reducing ocean VOC accumulation and air-sea transfer. If the

proposed model for interpreting these results, that heterotroph consumption stimulates

photosynthesis because phytoplankton replenish lost VOC resources, is correct, then it is

conceivable that the mechanisms we described result in a global increase in ocean photosynthesis

and are an important conduit of direct transfer of GCP to bacteria that does not rely on the

influence of protistan grazing and virus predation.

Materials and methods:

Culture growth and <u>P. tricornutum</u> physiology:

Phaeodactylum tricornutum strain CCMP 2561 was maintained axenically and in

separate co-cultures with *Roseibium* sp. 13C1, *Yoonia* sp. 4BL, *Marinobacter* sp. 3-2,

Rhodophyticola sp. 6CLA, and *Stappia* sp. ARW1T. Cultures were grown under 12h:12h light:

dark cycles (60-70 µmole photons m^{-2} s^{-1}), at 19°C in f/2+Si medium (ASW) [58]. DAPI (NucBlue,

Thermo Fischer) and fluorescence microscopy were used to check *P. tricornutum* axenicity.

Bacterial and diatom growth were determined in triplicate. Cell densities were measured using a

GUAVA flow cytometer (Millipore; Billerica, MA, USA) 1-2 h into the light phase. VOCs,

chlorophyll content (Chl*a*), and photosynthetic efficiency (F_v/F_m) were measured 4-6 h into the

light phase. Chl*a* was determined from filtering 2-5 ml of culture (GF/F, Whatman, 25 mm),

extracting in 5 ml 90% acetone, and storing at -20°C for 24 h. Chl*a* was quantified by

spectrophotometer (Shimadzu; Kyoto, Japan) [59]. F_v/F_m was measured by fast repetition-rate

fluorometer following 10 min dark acclimation [60]. Short-term carbon fixation rates

(approximating gross carbon production) were measured as in Moore et al. (2020). Axenic *P.*

tricornutum was grown in a closed system with culture headspace recirculating at 80 ml min^{-1} by

peristaltic pump and either directed back into the culture medium or passed first through a

hydrocarbon trap (Marineland Black Diamond Activated Carbon) and then into the medium. *P.*

tricornutum-bacteria cocultures were grown in the same manner with no hydrocarbon trap. All

flasks were recirculated for 24 h prior to inoculation with the cultures to remove background

VOCs. Samples for $^{14}CO_2$-uptake (10 ml) were collected by syringe during the *P. tricornutum*

exponential phase (2-4 x 10^5 cells ml^{-1}), spiked with 2 µCi of ^{14}C-sodium bicarbonate and

incubated 20 min in light or dark.

VOC measurement by PTR-TOF/MS

VOCs were quantitatively measured in axenic *P. tricornutum* and cocultures by PTR-TOF/MS

(Ionicon, Austria), as in Moore et al. (2020). Autoclaved ASW and HPLC-grade water were

measured alongside the samples as blank controls. Culture, ASW, or HPLC water (100 ml) was

transferred to a dynamic stripping chamber [56]and bubbled with breathing-grade air passed first

through a hydrocarbon trap and then through a glass frit in the bottom of the chamber at 50 ml

min^{-1} for 5 min to strip VOCs from the sample. PTR-TOF/MS data were analyzed using PTR

viewer 3.4.3 (Ionicon Analytik). Calibration and binning of m/z signals were performed as in

Moore et al. (2020). By using correlation plots and GLOVOCS [43], contaminants, fragments,

duplicates, and water clusters were removed; m/z signals were putatively identified using

GLOVOCS and Ionicon Viewer 3.4.1.

Benzene uptake and nanoSIMS analyses:

Benzene dose experiments were performed on *Rhodophyticola*, *Marinobacter*, and *Roseibium*.

Stappia could not be reliably grown in the absence of *P. tricornutum,* and *Yoonia* showed no

depletion of benzene in coculture. A saturated benzene solution was prepared in ASW and added

to *P. tricornutum*-spent media (prepared from 0.2 µm filtered spent media collected from axenic

P. tricornutum grown to 2-4 x 10^5 cells ml^{-1} and diluted in 1:10 with ASW) to achieve benzene

concentrations 0-180 µM with no headspace. Cultures were maintained at 19°C in the dark.

^{13}C-benzene uptake was evaluated using nanoSIMS in six different treatments grown in Teflon-

sealed borosilicate vials (*Marinobacter*) or airtight Nalgene bottles (*Rhodophyticola*). Fifty nM

98% ^{15}N leucine (Cambridge Isotopes, MA, USA) was added to all treatments as an independent

measure of cell activity (see Supp. table 9). Sample preparation for nanoSIMS and a brief

explanation of isotopic measurement are explained elsewhere (Supp. document 4). A CAMECA

NanoSIMS 50 was used to measure the isotopic composition of individual cells, explained in

detail elsewhere [61].

SEM imaging

Samples (5 ml) of *Marinobacter* and *Rhodophyticola* cocultured with *P. tricornutum* were collected in mid-exponential phase, treated with 1.25 ml of 2X 5% glutaraldehyde + 2% paraformaldehyde (PFA) in 0.1 M sodium cocadylate buffer (NaCB), incubated for 24 h, filtered on a 0.2 µm polycarbonate filter, and washed three times with NaCB + Milli-Q water. The filtered cells were subjected to critical point drying and then vacuum-dried and gold-coated. Imaging was done using a detector with a 5.1 mm working distance, 1.8 kV voltage, and 20-30 µm aperture diameter at the Electron Microscope facility, LPSC, Oregon State University. The number of bacteria adhered to *P. tricornutum* in coculture was estimated from 50 diatom cells.

Genetic potential for hydrocarbon metabolism

Whole genome sequences of each bacteria were acquired from NCBI (GB accession numbers: *Marinobacter*: PRJNA500125; *Rhodophyticola*: PRJNA441682; Yoonia: PRJNA441685; *Stappia*: PRJNA441689; *Roseibium*: PRJNA441686). Geneious Prime v2023.2.1 and Hidden Markov Models were used to search genomes for proteins initiating hydrocarbon oxidation (i.e., *alkB*, Pfam(PF00487); *almA*, Pfam(PF00743); *rhdA*, Pfam(PF00848); *acmA*, Pfam(PF00743); *aldH*, Pfam(PF00171) [62]. Each hit for *alkB* and *rhdA* was manually curated using Geneious Prime v2023.2.1. Each hit was aligned using MUSCLE protein alignment in Geneious prime v2023.2.1.

Statistics

Statistical analyses and figure construction were done in R studio v.4.1.1, with ggplot2 and Complex heatmap packages [63,64]. Paired student t-tests were used to identify significant

differences between treatments. Benjamini-Hochberg corrections (Q-value cut-off ≤ 0.1)

eliminated false positive m/z signals. Principle component analysis and Permutational

Multivariate Analysis of Variance (PERMANOVA) were used to evaluate VOC accumulation

and depletion patterns between cultures.

Acknowledgments

We thank Christina Ramon for lab assistance and Nicholas Baetge and Chih Ping Lee for

assistance with data analysis. This research was supported by award ID OCE1948163 to K.H.H.

and S.J.G. from the National Science Foundation, Biological Oceanography Program. Work at

LLNL was Supported by the US Department of Energy's (DOE) Genomic Science Program

through the LLNL μBioSpheres Science Focus Area grant # SCW1039 and performed under the

auspices of the DOE under contract DE-AC52-07NA27344.

Author contributions

K.H.H., X.M., and V.G.P. conceived of the study. V.G.P. conducted all experiments with

assistance from K.A., K.J., and L.C. and wrote the first manuscript draft with K.H.H. NanoSIMS

analyses were conducted by X.M. and P.K.W. All authors contributed to final version of the

manuscript.

References:

1.   Fu, H., Uchimiya, M., Gore, J. & Moran, M. A. Ecological drivers of bacterial community
assembly in synthetic phycospheres. *Proc Natl Acad Sci U S A* **117**, 3656–3662 (2020).

2.   Seymour, J. R., Amin, S. A., Raina, J.-B. & Stocker, R. Zooming in on the phycosphere:
the ecological interface for phytoplankton–bacteria relationships. *Nat Microbiol* **2**, 17065
(2017).

3.   Mitchell, J. G., Okubo, A. & Fuhrman, J. A. Microzones surrounding phytoplankton form the basis for a stratified marine microbial ecosystem. *Nature 1985 316:6023* **316**, 58–59 (1985).

4.   Seymour, J. R., Ahmed, T., Durham, W. M. & Stocker, R. Chemotactic response of marine bacteria to the extracellular products of Synechococcus and Prochlorococcus. *Aquatic Microbial Ecology* **59**, 161–168 (2010).

5.   Smriga, S., Fernandez, V. I., Mitchell, J. G. & Stocker, R. Chemotaxis toward phytoplankton drives organic matter partitioning among marine bacteria. *Proc Natl Acad Sci U S A* **113**, 1576–1581 (2016).

6.   Kogure, K., Simidu, U. & Taga, N. Bacterial attachment to phytoplankton in sea water. *J Exp Mar Biol Ecol* **56**, 197–204 (1981).

7.   Ramanan, R. *et al.* Phycosphere bacterial diversity in green algae reveals an apparent similarity across habitats. *Algal Res* **8**, 140–144 (2015).

8.   Raina, J. B. *et al.* Chemotaxis increases metabolic exchanges between marine picophytoplankton and heterotrophic bacteria. *Nature Microbiology 2023 8:3* **8**, 510–521 (2023).

9.   Ugolini, G. S. *et al.* Microfluidic approaches in microbial ecology. *Lab Chip* **24**, 1394–1418 (2024).

10.   Durham, B. P. *et al.* Cryptic carbon and sulfur cycling between surface ocean plankton. *Proc Natl Acad Sci U S A* **112**, 453–457 (2015).

11.   Nouchi, I., Hosono, T. & Sasaki, K. Seasonal changes in fluxes of methane and volatile sulfur compounds from rice paddies and their concentrations in soil water. *Plant Soil* **195**, 233–245 (1997).

12.   Brisson, V. *et al.* Identification of Effector Metabolites Using Exometabolite Profiling of Diverse Microalgae. *mSystems* **6**, (2021).

13.   Russo, A., Pollastri, S., Ruocco, M., Monti, M. M. & Loreto, F. Volatile organic compounds in the interaction between plants and beneficial microorganisms. *J Plant Interact* **17**, 840–852 (2022).

14.   Dawson, R. A. *et al.* The microbiology of isoprene cycling in aquatic ecosystems. *Aquatic Microbial Ecology* **87**, 79–98 (2021).

15.   Shemi, A., Ben-Dor, S., Rotkopf, R., Dym, O. & Vardi, A. Phylogeny and biogeography of the algal DMS-releasing enzyme in the global ocean. *ISME Communications* **3**, (2023).

16.   Asher, E., Dacey, J. W., Ianson, D., Peña, A. & Tortell, P. D. Concentrations and cycling of DMS, DMSP, and DMSO in coastal and offshore waters of the Subarctic Pacific during summer, 2010-2011. *J Geophys Res Oceans* **122**, 3269–3286 (2017).

17. Pozzer, A. C., Gómez, P. A. & Weiss, J. Volatile organic compounds in aquatic
ecosystems – Detection, origin, significance and applications. *Science of the Total*
Environment **838**, (2022).

18. Moore, E. R., Weaver, A. J., Davis, E. W., Giovannoni, S. J. & Halsey, K. H. Metabolism
of key atmospheric volatile organic compounds by the marine heterotrophic bacterium
Pelagibacter HTCC1062 (SAR11). *Environ Microbiol* **24**, 212–222 (2022).

19. Rocco, M. *et al.* Oceanic phytoplankton are a potentially important source of benzenoids
to the remote marine atmosphere. *Communications Earth & Environment 2021 2:1* **2**, 1–8
(2021).

20. Misztal, P. K. *et al.* Atmospheric benzenoid emissions from plants rival those from fossil
fuels. *Scientific Reports 2015 5:1* **5**, 1–10 (2015).

21. Moore, E. R., Davie-Martin, C. L., Giovannoni, S. J. & Halsey, K. H. Pelagibacter
metabolism of diatom-derived volatile organic compounds imposes an energetic tax on
photosynthetic carbon fixation. *Environ Microbiol* **22**, 1720–1733 (2020).

22. Wu, H. jun *et al.* Metagenomic analysis reveals specific BTEX degrading microorganisms
of a bacterial consortium. *AMB Express* **13**, (2023).

23. Bacosa, H. P., Mabuhay-Omar, J. A., Balisco, R. A. T., Omar, D. M. & Inoue, C.
Biodegradation of binary mixtures of octane with benzene, toluene, ethylbenzene or
xylene (BTEX): insights on the potential of Burkholderia, Pseudomonas and Cupriavidus
isolates. *World J Microbiol Biotechnol* **37**, 1–8 (2021).

24. Hocinat, A., Boudemagh, A., Ali-Khodja, H. & Medjemadj, M. Aerobic degradation of
BTEX compounds by Streptomyces species isolated from activated sludge and
agricultural soils. *Arch Microbiol* **202**, 2481–2492 (2020).

25. Khodaei, K., Nassery, H. R., Asadi, M. M., Mohammadzadeh, H. & Mahmoodlu, M. G.
BTEX biodegradation in contaminated groundwater using a novel strain (Pseudomonas
sp. BTEX-30). *Int Biodeterior Biodegradation* **116**, 234–242 (2017).

26. Jiang, B. *et al.* Biodegradation of Benzene, Toluene, Ethylbenzene, and o-, m-, and p-
Xylenes by the Newly Isolated Bacterium Comamonas sp. JB. *Appl Biochem Biotechnol*
176, 1700–1708 (2015).

27. Halsey, K. H. & Giovannoni, S. J. Biological controls on marine volatile organic
compound emissions: A balancing act at the sea-air interface. *Earth Sci Rev* **240**, 104360
(2023).

28. Rix, G. D. *et al.* Analysis of Essential Isoprene Metabolic Pathway Proteins in Variovorax
sp. Strain WS11. *Appl Environ Microbiol* **89**, (2023).

29. Halsey, K. H. *et al.* Biological cycling of volatile organic carbon by phytoplankton and
bacterioplankton. *Limnol Oceanogr* **62**, 2650–2661 (2017).

30. Yurimoto, H. & Sakai, Y. Interaction between C1-microorganisms and plants: contribution to the global carbon cycle and microbial survival strategies in the phyllosphere. *Biosci Biotechnol Biochem* **87**, 1–6 (2022).

31. Samo, T. J. *et al.* Attachment between heterotrophic bacteria and microalgae influences symbiotic microscale interactions. *Environ Microbiol* **20**, 4385–4400 (2018).

32. Di Costanzo, F., Di Dato, V. & Romano, G. Diatom–Bacteria Interactions in the Marine Environment: Complexity, Heterogeneity, and Potential for Biotechnological Applications. *Microorganisms 2023, Vol. 11, Page 2967* **11**, 2967 (2023).

33. Wear, E. K., Carlson, C. A., Windecker, L. A. & Brzezinski, M. A. Roles of diatom nutrient stress and species identity in determining the short- and long-term bioavailability of diatom exudates to bacterioplankton. *Mar Chem* **177**, 335–348 (2015).

34. Liu, S. *et al.* Opportunities and challenges of using metagenomic data to bring uncultured microbes into cultivation. *Microbiome* **10**, (2022).

35. Wang, Y., Zhao, Y., Bollas, A., Wang, Y. & Au, K. F. Nanopore sequencing technology, bioinformatics and applications. *Nature Biotechnology 2021 39:11* **39**, 1348–1365 (2021).

36. Van Tol, H. M., Amin, S. A. & Virginia Armbrust, E. Ubiquitous marine bacterium inhibits diatom cell division. *ISME J* **11**, 31–42 (2017).

37. Helliwell, K. E., Shibl, A. A. & Amin, S. A. The Diatom Microbiome: New Perspectives for Diatom-Bacteria Symbioses. *The Molecular Life of Diatoms* 679–712 (2022) doi:10.1007/978-3-030-92499-7_23/COVER.

38. Balakrishnan, R., de Silva, R. T., Hwa, T. & Cremer, J. Suboptimal resource allocation in changing environments constrains response and growth in bacteria. *Mol Syst Biol* **17**, 10597 (2021).

39. Mayali, X. *et al.* Single-cell isotope tracing reveals functional guilds of bacteria associated with the diatom Phaeodactylum tricornutum. *Nat Commun* **14**, 5642 (2023).

40. Kim, H. *et al.* Bacterial response to spatial gradients of algal-derived nutrients in a porous microplate. *The ISME Journal 2021 16:4* **16**, 1036–1045 (2021).

41. Lang, X. P., He, Z., Yang, G. P. & Dai, G. Physiological responses and altered halocarbon production in Phaeodactylum tricornutum after exposure to polystyrene microplastics. *Ecotoxicol Environ Saf* **268**, 115702 (2023).

42. Cabrita, M. T., Vale, C. & Rauter, A. P. Halogenated Compounds from Marine Algae. *Marine Drugs 2010, Vol. 8, Pages 2301-2317* **8**, 2301–2317 (2010).

43. Yáñez-Serrano, A. M. *et al.* GLOVOCS-Master compound assignment guide for proton transfer reaction mass spectrometry users. *Atmos Environ* **244**, 117929 (2021).

44.   Jakoncic, J., Jouanneau, Y., Meyer, C. & Stojanoff, V. The catalytic pocket of the ring-
hydroxylating dioxygenase from Sphingomonas CHY-1. (2006)
doi:10.1016/j.bbrc.2006.11.117.

45.   Tan, H. M., Tang, H. Y., Joannou, C. L., Abdel-Wahab, N. H. & Mason, J. R. The
Pseudomonas putida ML2 plasmid-encoded genes for benzene dioxygenase are unusual in
codon usage and low in G + C content. *Gene* **130**, 33–39 (1993).

46.   Bagnéris, C., Cammack, R. & Mason, J. R. Subtle Difference between Benzene and
Toluene Dioxygenases of Pseudomonas putida. *Appl Environ Microbiol* **71**, 1570 (2005).

47.   Mangwani, N., Kumari, S. & Das, S. Involvement of quorum sensing genes in biofilm
development and degradation of polycyclic aromatic hydrocarbons by a marine bacterium
Pseudomonas aeruginosa N6P6. *Appl Microbiol Biotechnol* **99**, 10283–10297 (2015).

48.   Olson, M. S., Ford, R. M., Smith, J. A. & Fernandez, E. J. Quantification of bacterial
chemotaxis in porous media using magnetic resonance imaging. *Environ Sci Technol* **38**,
3864–3870 (2004).

49.   Lacal, J. *et al.* Bacterial chemotaxis towards aromatic hydrocarbons in Pseudomonas.
Environ Microbiol **13**, 1733–1744 (2011).

50.   Matilla, M. A., Gavira, J. A. & Krell, T. Accessing nutrients as the primary benefit arising
from chemotaxis. *Curr Opin Microbiol* **75**, 102358 (2023).

51.   Bell, W. & Mitchell, R. Chemotactic and growth responses of marine bacteria to algal
extracellular products. Biol. Bull. 143, 265–277 (1972).

52.   Hearn, E. M., Patel, D. R. & Van Den Berg, B. Outer-membrane transport of aromatic
hydrocarbons as a first step in biodegradation. *Proc Natl Acad Sci U S A* **105**, 8601–8606
(2008).

53.   Chicca, I. *et al.* Degradation of BTEX mixture by a new Pseudomonas putida strain: role
of the quorum sensing in the modulation of the upper BTEX oxidative pathway.
Environmental Science and Pollution Research **27**, 36203–36214 (2020).

54.   Sun, J. *et al.* One Carbon Metabolism in SAR11 Pelagic Marine Bacteria. *PLoS One* **6**,
e23973 (2011).

55.   Simó, R., Cortés-Greus, P., Rodríguez-Ros, P. & Masdeu-Navarro, M. Substantial loss of
isoprene in the surface ocean due to chemical and biological consumption.
Communications Earth & Environment 2022 3:1 **3**, 1–8 (2022).

56.   Davie-Martin, C. L., Giovannoni, S. J., Behrenfeld, M. J., Penta, W. B. & Halsey, K. H.
Seasonal and Spatial Variability in the Biogenic Production and Consumption of Volatile
Organic Compounds (VOCs) by Marine Plankton in the North Atlantic Ocean. *Front Mar*
Sci **7**, (2020).

57. Halsey, K. H. *et al.* Biological cycling of volatile organic carbon by phytoplankton and bacterioplankton. *Limnol Oceanogr* **62**, 2650–2661 (2017).

58. Guillard, R. R. L. & Ryther, J. H. STUDIES OF MARINE PLANKTONIC DIATOMS: I. CYCLOTELLA NANA HUSTEDT, AND DETONULA CONFERVACEA (CLEVE) GRAN. *Can J Microbiol* **8**, 229–239 (1962).

59. Ritchie, R. J. Consistent Sets of Spectrophotometric Chlorophyll Equations for Acetone, Methanol and Ethanol Solvents. *Photosynth Res* **89**, 27–41 (2006).

60. Kolber, Z. S., Prášil, O. & Falkowski, P. G. Measurements of variable chlorophyll fluorescence using fast repetition rate techniques: defining methodology and experimental protocols. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1367**, 88–106 (1998).

61. Dekas, A. E. *et al.* Characterizing Chemoautotrophy and Heterotrophy in Marine Archaea and Bacteria With Single-Cell Multi-isotope NanoSIP. *Front Microbiol* **10**, 455124 (2019).

62. Apweiler, R. *et al.* The Universal Protein Resource (UniProt) 2009. *Nucleic Acids Res.* **37**, D169–D174 (2008).

63. Gu, Z. Complex heatmap visualization. *iMeta* **1**, (2022).

64. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

Figures and tables:

Table 1: Traits of bacteria used in this study

Bacteria	Carbohydrate metabolism[a]	Hydrocarbon metabolism[b]	Motility	Chemotaxis (*che* genes)	Attachment to diatom[c]
Marinobacter	+	+++	Yes	Yes	Yes
Roseibium	+	+++	Yes	Yes	Yes
Rhodophyticola	+++	++	No	No	No
Stappia	+++	++	No	No	No
Yoonia	+++	+	No (?)	Yes	Yes

[a]Summarized from Mayali et al. 2023
[b]Summarized from Fig. 2.
[c]Mayali et al. 2023 and this study

Figure 1: VOC production in P. tricornutum. a. Cell-normalized concentrations of m/z values (rows) ranging from low (light blue) to high (orange) concentrations produced during exponential and stationary phases. Black are m/z values that were not significantly higher than the media control. Functional group classifications for each m/z+1 value are shown in the rightmost column. b. P. tricornutum growth (n=6 independent cultures). Arrows indicate sampling days for VOCs.

Figure 2: Phycosphere bacteria demonstrate widely ranging VOC depletion patterns. a. m/z+1 depletion (rows) in each P. tricornutum–bacteria coculture (columns are labeled with the bacteria grown with P. tricornutum) in exponential (a) and stationary phase (b) relative to axenic P. tricornutum in each of the growth phases. The magnitude of depletion is shown ranging from low (light blue) to high (yellow). Pearson correlation distances were used to cluster depletion patterns between PT-bacteria cocultures (top axes). Pie charts show the proportion of the total depleted VOC concentration in each functional group.(c) Growth of each bacteria (red lines) in coculture with P. tricornutum (blue lines), (n=6). Arrows indicate sampling day for VOCs. d. Genetic potential for hydrocarbon metabolism in the phycosphere bacteria. Number of gene copies is shown for each bacteria. Abbreviations: alkB (alkane-1-monooxygenase), bnzA (benzene dioxygenase alpha-subunit), bnzB (benzene dioxygenase beta-subunit), almA (flavin-binding monooxygenase), aor (aldehyde oxidoreductase), acmA (acetone monooxygenase).

Figure 3. Scanning electron microscopy images showing (a) *Marinobacter* attached to *P. tricronutum* in a flat orientation and (b) *Roseibium* attached to *P. tricornutum* in polar orientation.

Figure 4: Photosynthetic carbon fixation in P. tricornutum in the presence and absence of VOC sinks (hydrocarbon trap or each bacteria in coculture). Points are short term (10 min) carbon fixation rates for each culture or coculture. Box plots show the mean and upper and lower quartile divisions; whiskers show 95% confidence limits. *p-value < 0.01; t-test. Inset shows the total bacterial cell-normalized VOC depletion in each P. tricornutum-bacteria coculture.

Figure 5: Benzene incorporation into Marinobacter and Rhodophyticola biomass (a,c) Bacterial cell densities in the presence (solid lines) and absence (dashed lines) of 13C benzene in ASW (blue), Ptspent (olive), and coculture with P. tricornutum (orange). Asterisks indicate treatments with significant increases in cell densities in the presence of benzene. (b, d) 13C enrichment in Marinobacter and Rhodophyticola in ASW (Marinobacter only), Ptspent, and in coculture with P. tricornutum. Cnet is the fraction of biomass labelled from 13C benzene. (e) Linear regression of bacterial 13C benzene enrichment in PT-Marinobacter (top) and PT-Rhodophyticola (bottom). Student t-test *p-value < 0.05. (f) Expanded y-axis showing bacterial Cnet values in cocultures and inset SEM images of Marinobacter attached to P. tricornutum and Rhodophyticola unattached.

Bacteria ◦ *Yoonia* 4BL ▪ *Roseibium* 13C1 • Marinobacter 3−2 • *Rhodophyticola* 6CLA • *Stappia* ARW1T

Supp. fig. 1: Principal coordinate analysis of depleted VOCs in *P. tricornutum*–bacteria cocultures in exponential (left) and stationary (right) growth phases.

Bacteria
● *Roseibium* ▲ *Marinobacter* ■ *Rhodophyticola*
Treatment ● PTspent ○ ASW

Supp. fig. 2: Final cell densities measured in benzene dose experiments. Each bacterium was inoculated into Ptspent (filled symbols) and ASW media (open symbols) with initial benzene concentrations 0-180 μM. *Roseibium* (circles), *Marinobacter* (triangles), and *Rhodophyticola* (squares).

Supp. fig. 3: Cnet vs Nnet in *Marinobacter* and *Rhodophyticola* measured in three different treatments.

606