Temporal regularities of vocal exchange in Java sparrows

Sota Kikuchi[1][¶], Noriko Kondo[1][¶], and Hiroki Koda[1][*]

[1]Graduate School of Arts and Sciences, The University of Tokyo, Komaba, Tokyo, 153-8902, Japan

[¶]Co-first authors/equal contributions

[*]Corresponding to HK

HK (ORCID: 0000-0002-0927-3473): hkoda@g.ecc.u-tokyo.ac.jp

Abstract

In human turn-taking, speakers alternate at very short intervals while avoiding overlaps. If speakers do not receive a vocal response from another party, they often repeat their utterance after the expected response time has elapsed to elicit a reply. Intra-individual intervals tend to be longer than inter-individual intervals. Such temporal regularity in vocal exchanges has also been observed in social mammals, such as non-human primates. In contrast, vocal exchanges in birds have been studied primarily in the context of songs, with limited research on call-based vocal exchanges. Studies specifically examining intra-individual call intervals are even scarcer. In this study, we investigated vocal exchanges and their temporal patterns in Java sparrows placed face-to-face. The results revealed that they vocalized at very short intervals following the calls of the other individual. However, no significant differences were observed between the inter- and intra-individual intervals. This contrasts with the temporal characteristics of vocal exchanges observed in social mammals that have been studied to date. We propose that a possible explanation for this difference lies in the variation in social group structures between birds and mammals.

Keywords: Vocal exchange, turn-taking, avian call, temporal regularity, behavioural contagion, group formation

Introduction

Human conversation is characterized by a vocal turn-taking phenomenon, in which speakers quickly alternate roles to convey information to each other (e.g. [1]). Turn-taking in conversations is characterized by two key features: i) the preservation of extremely short intervals between turns, and ii) the avoidance of overlapping speech. Notably, inter-individual intervals–the time between the end of one speaker's utterance and the beginning of the next–exhibit a universal temporal regularity, averaging approximately 200 ms regardless of linguistic or cultural differences [2, 3].

As human vocal turn-taking is characterized by consistent short temporal regularity

regardless of semantic content, the vocal exchange of non-human animals can be examined

within the same framework. Numerous studies have explored vocal interactions across various

animal species in an effort to draw parallels with human conversational turn-taking [4,5].

Primate vocal exchanges have been extensively studied, particularly in *contact calls*. *Contact*

calls are common vocalizations exchanged between individuals in a cohesive social group and

have been analyzed in detail to characterize the intervals in vocal exchange. Similar to human

conversation, vocal exchanges in many primate species exhibit temporal regularity,

characterized by stable intervals between individuals' vocalizations and significant avoidance of

overlap (e.g., [6–9]). In addition, it has been reported that the time interval for repeated

vocalizations by the same individual in the absence of a response from other individuals

(repeated transition) is longer than the time interval for vocalizations by different individuals

taking turns (exchange transition) [8,10–15]. A longer time interval in the exchange transition

can be interpreted as the manifestation of a vocalizing individual expecting a response from

another individual. In other words, after calling, the caller anticipates and waits for a response

from another individual, and if no response is received, it calls again to elicit a reaction. This

temporal regularity represents a key similarity between human conversation and animal vocal

exchange and has been reported in socially living non-primate mammals, such as rodents

[19,20], bats [21], elephants [22,23], meerkats (*Suricata suricattaI*) [24], and cetaceans [25–30].

Accordingly, this can be considered a common rule in the communication of social mammals

and is adaptive in socially structured groups. (See the conceptual framework of temporal

regularity in Fig 1.)

Fig 1. Conceptual representation of the mechanism underlying the emergence of temporal

regularity in vocal exchange/turn-taking of various social mammals.

When social mammals are engaged in a reciprocal vocal exchange (A, top), the individual

expects a "response" from the other individual and therefore waits for a response. After the

expected response time had elapsed, the individual repeated the vocalization (A, bottom). Fig

1B illustrates the inter-call intervals of vocal exchange in mammals. The inter-call interval of

the repeated transition t_r is longer than that of the exchange transition t_e (A, B).

Studies of vocal communication in birds have mainly focused on songs, the elaborate

and complex vocalizations that function in territorial defense and mate attraction [31]. Song

interaction studies include male-male counter-singing [32] and male-female duets [33]. The

exchange of calls, on the other hand, has not been as extensively documented as in primates.

However, calls are more appropriate than songs and duets when compared with human

conversational vocal turn-taking and mammalian vocal exchange, as the latter are primarily

restricted to breeding contexts. Some studies have examined the vocal exchange in bird species

from a comparative perspective with that of mammals. In large-billed crows (*Corvus*

macrorhynchos), temporal regularity in the vocal exchange of their contact, *'ka call'* - a fixed

time exchange interval between different callers - was observed [34]. Similarly, the vocal

exchange of calls in parents and offspring in common starlings (*Sturnus vulgaris*) [35] and the

begging calls of barn owls (*Tyto alba*) [36] also confirmed the avoidance of caller vocal overlap,

indicating temporal regularity in these call exchanges, although these studies did not examine

the inter-call intervals in detail. Recently, calls have received more attention, and call exchange

has begun to be studied in zebra finches (*Taeniopygia guttata*) of the family Estrildidae, a

leading model species in song research, providing new insights into avian vocal exchange.

Experimental studies examining vocal call exchange in zebra finches placed face-to-face

confirmed that call overlap was significantly avoided and vocal timing was actively adjusted

[37–40]. The authors of these studies argued that the vocal timing adjustments observed in

zebra finches, which depend on the timing of other individual calls, are analogous to primate

vocal exchanges. Such adjustments in call timing are considered a reflection of social

relationships, such as pair mating, in zebra finches [40], suggesting that ecological adaptations

in vocal interactions are similar to those observed in mammals. Although these findings

highlight the role of vocal timing adjustments in social interactions, research to date has

primarily focused on the intervals between calls produced by different individuals, while studies

on the intervals between repeated transitions remain limited (see [41]).

When considering the role of vocal turn-taking in animal societies, the socio-ecological

structure of the species in question should be taken into account. Social mammals form stable

groups with fixed membership even outside the breeding season, often including unrelated

individuals. In such groups, vocal turn-taking helps maintain group cohesion by enabling

individuals to signal their location and avoid separation through reciprocal vocal responses [42].

Birds, on the other hand, also form social flocks, but these tend to be more fluid and less stable

than mammalian social structures. Within bird flocks, the most salient social units are often

pair-mates, family members, or kin. For example, species such as Greylag geese (*Anser anser*)

form matrilineal alliances [43], yet they do not necessarily maintain exclusive territories with

fixed group membership, as is often observed in mammal groups. Given these socio-ecological

differences, along with the fact that birds in general are capable of traveling much faster than

mammals due to their ability to fly, temporary spatial separation from flock members likely

poses a lower risk for birds than for group-forming social mammals. Accordingly, vocal turn-

taking with an expectation of a response appears to be less essential for birds than for social

mammals as a mechanism for maintaining proximity and group cohesion. As is evident from

recent zebra finch studies, however, the vocal exchange of calls in birds has been increasingly

examined using a framework similar to that applied to primates. Previous studies have shown

that humans, non-human primates, mammals, and birds share common vocal exchange

features—specifically, short inter-individual intervals and longer intra-individual intervals, both

of which help to avoid call overlap. Due to these shared characteristics, it has been hypothesized

that vocal exchange may serve comparable functions across taxa. Nevertheless, as noted above,

fundamental socioecological differences between social mammals and birds may shape their

communicative strategies in distinct ways. Importantly, most avian vocal exchange research has

focused on zebra finches, a key model species in songbird studies. These studies, which

involved controlled face-to-face experiments, demonstrated significant avoidance of vocal

overlap and active adjustments in call timing [37–40]. The authors of these studies proposed

that such timing adjustments, which depend on the calls of others, are analogous to primate-like

vocal turn-taking. Moreover, call timing was found to reflect social relationships such as pair

bonding [40], suggesting that ecological functions of call timing may be more similar across

taxa than previously assumed. However, these studies have mainly focused on inter-individual

call intervals, and research on intra-individual repeated transitions remains limited (see [41]).

In this study, we investigated the temporal regularity of vocal exchanges in the Java

sparrow, *Padda oryzivora*. Java sparrows are small birds belonging to the family Estrildidae

that are native to Java and Bali, Indonesia. These are socially monogamous species that

sometimes form large flocks [44]. Their calls have not been well studied, except for trill calls,

which are given in the context of affiliative and agonistic interactions [45]. There are also short

monosyllabic calls, presumed to function as contact calls [44], and we focus on the exchange of

these calls. As Java sparrows belong to the Estrildidae family, as do zebra finches, we expected

to observe vocal exchange with characteristics similar to those of zebra finches, that is, short

intervals between different birds. In addition, we investigated the within-individual call

intervals to test whether Java sparrows avoid the temporal window in which a 'response call'

from the other individual is most likely to occur, resulting in longer call intervals of the same

individual. More precisely, we investigated the temporal regularity of their vocal exchanges by

experimentally arranging multiple individuals face-to-face in a soundproof chamber, similar to

the experimental settings of studies on marmosets and zebra finches [8,11,46]. We compared

the time intervals between calls exchanged between different individuals and those repeated by

the same individual to examine the characteristics of their temporal distribution.

Methods

Subjects

Six adult Java sparrows (five males and one female, 2–6 years of age) were used in this

study. All individuals were housed separately in identical cages (38 cm high × 31 cm wide × 25

cm deep) within the same housing area, where they could hear and see each other. The housing

environment was maintained with a 12-hour light:12-hour dark (12L:12D) lighting cycle, a

room temperature ranging from 20°C to 28°C, and humidity levels kept above 40%. Adequate

amounts of food and water were provided in each cage, and the subjects had unrestricted access

to them throughout the experiment.

Apparatus

The experiment was performed in a soundproof chamber (2146 cm H × 1378 cm W ×

996 cm D) placed in a room separate from the housing room. Within the chamber, we designed

an experimental space that allowed individual cages to be placed opposite to each other, facing

one another (Fig. 2A). We placed the two cages at two different locations where the two target

birds could audio-visually recognize each other. The horizontal distance between the two cages

was 58 cm. We attached two small microphones (CVM-V02C, COMICA), one directly above

each of the two cages (approximately 10 cm), to allow close-up recording of vocalizations in

each cage. The microphones were connected via cables to an audio interface (UMC1820 U-

PHORIA, Behringer) using a computer (Ubuntu 20.2, CPU: Intel Core 5, RAM 16GB) located

outside the soundproof chamber. The audio signals were recorded at a 32-bit quantization/32

kHz sampling rate.

Fig 2. Schematic illustrations of (A) experimental settings, (B) labelling procedure of calls,

and (C) the generation process of shuffled session data.

(A) The experimental flow of the three phases, along with cage dimensions and spatial

arrangement. (B) The sound spectrogram (top), waveform (middle), and segmental call regions

labelled by evaluators (bottom), obtained from a 2-channel recording. (C) Workflow to generate

pseudo turn-taking data (right). The originally recorded F2F data were replaced in different

recording sessions, keeping the individual pair relationship consistency.

Procedures

A single session of the experiment consisted of three phases (pre-face-to-face phase,

PRE phase; face-to-face phase, F2F phase; and post-face-to-face phase, POST phase) according

to the following procedure (Fig. 2A).

First, the animals were exposed to a 15-minute PRE phase; in the PRE phase, only one

individual (the subject) was introduced into the soundproof room and placed in an individually

compartmented area. The purpose of the PRE phase was to habituate the participants to the

experimental space and encourage spontaneous vocalizations throughout the experimental

session while reducing their stress. The duration of the F2F phase was 15 minutes. In the F2F

phase, the other individual (facing individual) was placed opposite the subject bird facing it.

Finally, the POST phase was set for 15 min. During the POST phase, facing individuals were

removed from the soundproof room with only the preceding individuals remaining, as was the

case during the PRE phase. Four sessions were performed for every combination of two

individuals selected from six individuals (15 different pair sets) for a total of 60 sessions (2700

min) of recording. The order of the pair sets was randomized as much as possible to make sure

that the same individuals did not participate in consecutive experimental sessions.

Acoustic analysis

Labelling vocal data

Before analysis, all vocalizations recorded during the experimental sessions were

labelled (or annotated). Labelling was performed using three processes: segmentation,

classification, and caller identification. All labelling tasks were shared by two independent

evaluators (SK and NK) using Raven Pro version 1.6 [47] or Praat version 6.3.10 [48]. Before

the labelling tasks, the two evaluators checked with each other to agree on the criteria for the

segmentation and classification of vocalization types (Fig 2B).

Segmentation

Two evaluators independently judged the intervals between vocalizations to define the

intervals in the vocal region. Specifically, the onset and offset times of the vocal regions were

logged from the recorded data based on the visual inspection of the waveforms and

spectrograms while listening to the audio data.

Classifications and caller identifications

The detected vocal regions are classified into three types of vocal types (calls, songs,

and trills). Here, *songs* were classified as "vocal sequences that appear in a stereotyped pattern

for an individual," *calls* as "monosyllabic and short-note vocalizations that have no sequential

rules and are not songs," which is described as "contact calls" in [44], and *trills* as "typical

sounds that appear as short-duration repetitive pulses."

Caller identification was performed by comparing the intensities of multiple channels in

the recorded data. The location of the microphone channel with higher intensity was judged to

be closer to that of the target individual.

Labelling validation

To check the validity of the labelling, a subset of the data was extracted: four sessions

of data were selected from 60 sessions to include approximately 1000 vocalizations, and both

evaluators independently labelled the data in the F2F phase. The "agreement" of the labelling

between the two labelled data (label-SK, label-NK) was examined. The agreement rate between

two annotation labels (label-SK and label-NK) for one segment was defined as the accuracy rate

and was calculated as follows: $Acc = \frac{N_{agree}}{N_{all}}$, where Acc represents the accuracy rate, N_{all} is the

total number of segmented vocal regions, and N_{agree} is the number of agreements. . The

accuracy rate for classification and caller identification was 95.75% (of 969/1012 vocalizations).

We also examined whether the onset and offset times differed between the evaluators. The onset

and offset times were recorded in the two labelled datasets (label-SK and label-NK). To

compare the difference in time, we calculated the time difference ($\Delta t := t_{sk} - t_{nk}$) from the

time in label-SK (t_{sk}) to the time in the nearest neighbor label-NK (t_{nk}), for onset and offset,

respectively. The interval estimation of the obtained time difference data (n = 969) was

performed using a simple one-sample t-test. If there was a decision bias to annotate label-SK

earlier (or later) than label-NK, then there would be a "biased" interval estimation that did not

include 0. Otherwise, the interval estimation includes 0. The t-test results estimated the intervals

to be [-0.006, 0.010] and [-0.025, 0.012] for the onset and offset times, respectively, with no

statistically significant value bias compared to 0. Accordingly, this result indicates that there

was no statistical judgment bias for label-SK and NK.

Definition of the call transition sequences

For the analysis of vocal exchange, we only used recordings of the F2F phases. We

define the type of call transition sequence as follows: An "exchange transition" was defined as a

case in which a vocalization by one individual is followed by a vocalization by a different

individual, and a "repeated transition" as two consecutive vocalizations by the same individual.

In exchange transitions, the offset time of the preceding vocalization may be later than the onset

time of the following vocalization when an overlap occurs. Following these definitions, we

defined a "call interval" as the time interval between the offset of the preceding call and the

onset of the subsequent call, regardless of the vocalizing individual.

Statistical procedures

Estimations of interval structures by fitting Gaussian mixture models

To investigate whether the temporal structure of the exchange transitions and repeated

transitions in F2F phases differed statistically, we applied Gaussian mixture models (GMMs) to

the time interval data of each transition. The GMM attempts to estimate the number of mixtures

and parameters (mean and variance), assuming that the data are generated as a mixture of

multiple Gaussian distributions consisting of different parameters. Previous studies on vocal

exchange have shown that 1) the intervals follow logarithms and 2) the data are generated from

two types of Gaussian distributions: a short time interval caused by the association of two

vocalization transitions and a long time interval randomly caused by two independent

vocalizations. Our aim of the GMM analysis here was to isolate only the distribution of "the

time interval of associated two sequential vocalizations," which was our interest. By visually

inspecting the bimodal distribution of the interval data, we fit the model with mixtures of the

two components of Gaussian distributions to log-transformed data to estimate the parameters of

the mixture models. Prior to log transformation, we excluded interval cases with negative

values; that is, where the onset time of the subsequent call preceded the offset time of the

preceding call, and there were only two such cases. Of these two components, the one

estimating the shorter time interval ("short component," hereafter) was interpreted as the

distribution of "the time interval of associated two sequential vocalizations" for each of

exchange and repeated transitions. The other component estimating longer time intervals ("long

component") was interpreted as the distribution of "the time interval between two independent

vocalizations occurring at random." For the GMM fitting, we used `mclust` (v. 6.0.0) [49] in R

(v. 4.2.3) [50].

After estimating the interval structure by fitting GMM, we performed a permutation test

to statistically compare the interval between the exchange and repeated transitions as follows:

First, we generated values from two normal distributions defined by the mean and variance of

the intervals estimated by the GMM (two normal distributions for the exchange and repeat

transitions), and calculated the difference between the generated values. This calculation is

repeated 10,000 times to obtain a distribution of 10,000 values. We then calculated a 95^{th}

percentile interval, defined by the 2.5^{th} and 97.5^{th} percentiles of these values. By examining

whether this interval contained zero, we determined whether a statistically significant difference

existed between the two distributions. We used `rnorm()` function in R to generate data from

the normal distribution.

We further investigated whether the presence of a conspecific influenced the temporal

structure of intervals in repeated transitions. Specifically, we examined whether the intervals

between pre- and post-transition phases differed, using GMM estimation and a permutation test.

The results showed that the distributions of transition intervals in the PRE and POST phases

were remarkably similar. The estimated means (± variance) of the short-latency component

from the GMM were –0.75 ± 0.01 and –0.074 ± 0.334 for the PRE and POST phases,

respectively. The 95th percentile range for the difference in means, obtained via permutation

testing, ranged from –1.802 to 0.489 and included zero, indicating no significant difference

between the two phases. Based on this result, we combined the PRE and POST phases into a

single category, hereafter referred to as the "solo" phase. We then compared the interval

distributions between the solo and face-to-face (F2F) phases using the same GMM estimation

and permutation testing approach to assess whether they differed.

Pseudo-turn-taking data by shuffling the vocal data

The short component observed in the exchange transition in the F2F phases could be

interpreted as the reaction time to the preceding call in the vocal exchange, but we cannot rule

out the possibility that this might have been caused by chance when two individuals vocalized

independently at a certain interval of time. To investigate whether the distribution of the short

component was generated by chance, we compared the pseudo-turn-taking data (generated with

reference to [8]) with the actual exchange transition data. Pseudo-turn-taking data were created

by shuffling and combining data from different sessions to simulate "virtual" vocal exchanges.

Specifically, they were generated by pairing the onsets and offsets of the subject birds' calls

with those of the facing birds, each extracted from separate sessions in the F2F condition. Thus,

while the pairing remained consistent between the actual and pseudo-turn-taking data, the

vocalizations of the subject birds and facing birds occurred independently in the pseudo-turn-

taking condition. (Fig. 2C).

Statistical evaluation of the number of calls across experimental phases

To investigate whether the vocalizations of subject birds were influenced by those of

the facing birds, we compared the number of vocalizations produced by the subject birds across

the three phases (PRE, F2F, and POST) using generalized linear mixed models (GLMM) and

model selection techniques. We fitted a GLMM (error distribution: Poisson distribution, link

function: log) with the number of vocalizations given in each session by the subject birds that

were placed in the experimental space first (i.e., experienced all three experimental phases) as

the explanatory variable, the three experimental phases as the fixed effect term, and the subject

and trial as random intercepts. By comparing the Akaike information criterion (AIC), we

estimated which model outperformed the null model in which the condition was removed from

the fixed effects. GLMM fitting and AIC calculations were performed using the glmer function

of lme4 (v. 1.1-34) [51] and the dredge function of MuMIn (v. 1.47.5) [52] in R.

Ethical note

318   The experiments were reviewed and approved in advance by the Animal Experiment

319   Ethics Committee of the University of Tokyo Graduate School of Arts and Sciences (#2022-1

320   and 2023-8).

Results

Temporal distributions of call transitions

In the F2F phase, we recorded 5062 call transition sequences, of which 1712 were

exchange transitions, and 3350 were repeat transitions. There were only two overlapping

interactions in the exchange transitions. Fig 3 shows the distribution of the call offset onset

intervals for exchange and repeated transitions. The mean with variance of the estimated short

component of GMM are - 0.728 +/- 0.023 and - 0.669 +/- 0.035 for the exchange and repeated

transitions, respectively (Fig 3A and 3B, Table 1; note that the values were log-transformed).

This suggests that there was no statistical difference between the two transitions. Subsequently,

using these estimated parameters, we performed a permutation test to determine the differences

in values between the conditions. The calculated 95th percentile range of the value differences

was from -0.533 to 0.403, which was not biased away from the 0.

When comparing the distribution of repeated transition of 'solo' and F2F phase by

GMM, the estimated means (± variance) of the short-latency component were -0.75 ± 0.009 and

-0.669 ± 0.035, respectively. The 95th percentile range for the difference in means, obtained via

permutation testing, ranged from -0.329 to 0.505. These values include zero, indicating no

significant difference between the two phases. These results suggest that Java sparrows showed

vocal responses to facing individuals, as reflected in the shorter intervals during exchange

transitions. However, the duration of repeated transitions was not affected by the mere presence

of the other bird.

Fig 3. Histogram and estimated probability density function curve (curve line) fitted by

the two modal GMM for the exchange transition (A) and for repeated transitions (B).

The dashed vertical lines represent the positions of the estimated mean parameter values in the

short components of the GMMs. Note that although the data are log-transformed, the x-axis

shows antilogarithmic values in seconds.

Transition	Data size (N)	Components	Mean		Variance	Mixture ratio
Exchange	1710	short	-0.728	(0.187)	0.023	0.322
		long	0.321	(2.094)	0.502	0.678
Repeated	3350	short	-0.669	(0.214)	0.035	0.167
		long	0.578	(3.784)	0.296	0.833

Table 1 The parameter estimations of the GMMs fitting for the data of exchange transition and

repeated transition. The numbers in parentheses in the Mean, and Variance columns represent

antilogarithmic values, converted from logarithms back to seconds to help understanding.

Comparisons with pseudo-turn-taking data

In the pseudo-turn-taking data, we identified 948 exchange transitions and 4168

repeated transitions. There are three overlapping cases of exchange transition. Fig 4 shows the

distribution of call offset-onset intervals for the exchange transitions of the pseudo-turn-taking

data. The distribution of the pseudo-turn-taking data was uniform and monotonic, with no clear

components such as the short component observed in the exchange transition. This strongly

suggests that the short component identified in the exchange transitions originated from the

"response" call of the paired individuals to the preceding call of the other individuals.

Fig 4. Distributions of inter-call intervals of exchange transitions, repeated transitions and

pseudo turn-taking data. The red line represents the exchange transitions, the blue line

represents the repeated transitions, and the green line represents the pseudo-turn-taking data.

The length of the inter-call interval could be theoretically up to 900 seconds, but the X-axis here

is limited to 10 seconds. This is because more than 80% of the inter-call interval was contained

within 10 seconds (90.04% of exchange transition, 83.40% of repeated transition and 82.38% of

pseudo transition).

Comparisons of the number of calls across experimental phases

We recorded 9608 calls during 60 sessions (900 min for each of the three conditions,

totalling 2700 min). In the three experimental phases, the number of calls given by the birds

increased in the F2F phase for five of the six subjects compared to the PRE and POST phases

(Fig 5). The mean numbers of calls given by the birds first introduced into the experimental

space in the pre-, F2F-, and post-phases were 34.5, 50.8, and 33.8, respectively. The GLMM

analysis revealed that calls were most frequently made in the F2F phase (AIC: 4221.3; AIC:

4486.9; Fig. 5).

Fig 5. Total number of calls given by six subjects in three experimental phases.

(A) The average number of calls in each phase. Each line indicates each bird. The female bird is

indicated by the lowercase ID.

Discussion

In this study, we investigated call intervals given by two Java sparrows sequentially

(exchange transitions) and those given by the same individual repeatedly (repeated transitions).

The results showed that the time intervals of the short component did not differ between

exchanged and repeated transitions in Java sparrows. In other words, the birds in our study

repeatedly called at the same time intervals when calls from other individuals were most likely

to occur during exchange transitions. This contrasts with the temporal regularity of turn-

taking/vocal exchanges observed in social mammals, in which individuals remain silent during

time intervals when the likelihood of vocal responses from other individuals is high,

anticipating a response that leads to longer intervals in repeated transitions than in exchange

transitions. However, as evidenced by the increase in the number of calls during the F2F phase,

this does not imply that Java sparrows were calling independently of other individuals. Rather,

it is likely that Java sparrows responded contagiously to the calls of others, although they did

not appear to anticipate a 'response' call.

In call exchange studies in birds, the stable and short time intervals between calls given

by two individuals have been interpreted as vocal exchanges, similar to human turn-taking

behavior [38]. In the present study, the intra-individual call intervals in the exchange transitions

of Java sparrows were concentrated within a short temporal window. However, this result alone

does not allow for the conclusion that the vocal exchange of Java sparrows shares

characteristics similar to turn-taking in human conversations, as no significant difference in

interval length was observed between repeated and exchange transitions. This indicates that,

unlike the vocal turn-taking observed in socially communicative mammals, including humans,

Java sparrows do not expect or wait for a response from other individuals.

Not predicting or expecting a response from the partner does not mean that Java

sparrows give calls independently of the partner's call and do not communicate with each other.

A comparison of the pseudo-turn-taking data and exchange transitions showed a short

component only for the exchange transition. This indicates that they made calls in response to

the preceding calls of the other individual. In addition, Java sparrows made more vocalizations

in the F2F phase than in the standalone PRE and POST phases, although there were individual

differences. This indicates that calls have an interactive communication function. In other words,

Java sparrows vocally communicated with each other without predicting or anticipating a

response from the facing individual, but calling contagiously in response to others'

vocalizations. This is quite different from human conversation, in which turn-taking occurs at

an appropriate time by predicting the vocalizations of the person to whom he or she is speaking

[1].

Why do Java sparrows make contagious calls? We hypothesized that this would

synchronize the movement of the flock. For birds that live in fluid flocks and can fly and move

quickly, separation from a particular flock member is not as fatal as in mammals. Massive

aggregation is an ecological characteristic of collective flocks of small birds, such as sparrows.

In such large flocks, synchronizing the behavior among flock members is important for

increasing the survival rate [53]. Behavioral synchronization has been widely reported in

animals that live in a large mass of individuals: synchronization of vigilance or scanning

behaviors to avoid predators [54,55], escape motion from predators within a flock [56], flight

angles, and speeds between nearby individuals during flight [57]. Acoustic signals can function

as cues for synchronized movements. Spatial coordination by calls has been experimentally

reported in zebra finches flying in a wind tunnel [58]. In our study, we investigated only the

vocal exchange between two individuals in separate cages. To test our hypothesis that

contagious calling facilitates the coordination of flock movement, vocal exchange and

behavioral synchronization among multiple individuals should be investigated in more natural,

flock-like settings. The fact that they gave calls at similar intervals in both transitions suggests

that many overlaps of calls by different individuals may have occurred, but only two cases were

observed. Even in the pseudo-turn-taking data, the number of overlapping cases was as low as

three. Therefore, it is plausible that the small number of overlapping cases was due to the short

length of their calls (mean length of 24.7 msec). However, in Java sparrows, the call duration is

so brief that even if the birds anticipate a response, there may be no need to avoid overlapping.

This can be further clarified in future experiments using larger flocks. Specifically, in vocal

exchanges involving more than two individuals, the chance of call overlap would be expected to

increase simply due to the greater number of potential responders. If birds only respond

reflexively without anticipating the timing of others' calls, such overlap is likely to occur more

frequently. However, if individuals are capable of anticipating responses and coordinating their

vocal timing, they may be able to avoid overlaps even in larger groups. Indeed, such overlap

avoidance has been reported in zebra finches [37–40], where individuals adjust their timing to

avoid calling simultaneously. Whether similar mechanisms operate in Java sparrows in more

complex settings, such as those involving larger flocks, remains an open question.

In our study, Java sparrows were predominantly male, with a single female, and each

individual was housed separately. Consequently, there were no individuals with affiliative

relationships such as pairs. Given these conditions, it must be acknowledged that the social

environment of the birds in our study is a significant simplification of their natural social lives.

In zebra finches housed in mixed-sex groups, vocal exchange between pairs has been shown to

take priority, with vocal exchanges occurring at shorter intervals between pairs than in non-

paired individuals [41]. Additionally, the call interval in mixed-sex groups is shorter than that in

groups consisting only of females [59]. Given that, similar to zebra finches, pair bonds are

fundamental in Java sparrows, it is reasonable to assume that the temporal regularity of vocal

exchange may vary depending on social relationships. Although our study can be regarded as an

investigation of the basic temporal regularity of vocal exchange in the absence of pair bonds,

future research is essential to examine the influence of sex differences and social relationships

on the temporal regularity of vocal exchange.

In general, when two individuals produce consecutive vocalizations in a fixed

timeframe in non-human animals, this tends to be regarded as a vocal exchange with

characteristics similar to turn-taking in human conversations. However, temporal regularities

may differ between mammals and birds, as well as between species, such as zebra finches and

Java sparrows, owing to variations in their ecological adaptations and social group structures.

However, given the differences in results between zebra finches and Java sparrows, it is

currently impossible to draw definitive conclusions on vocal turn-taking in birds. Furthermore,

to enable meaningful comparisons between mammals and birds, research on call exchanges in

birds should be conducted across a wide range of species. We hope that this study will

encourage further research in the field of vocal exchange, ultimately shedding light on the

similarities and differences between avian and mammalian vocal interactions, their functions,

and their evolutionary significance.

Acknowledgements

We are grateful to two anonymous reviewers for drastically improving our manuscript. We

appreciate the administrative support provided by Riko Goto and Sana Kohmoto.

References

1. Sacks H, Schegloff EA, Jefferson G. A simplest systematics for the organization of turn-
taking for conversation. Language. 1974;50(4): 696–735. doi:
https://doi.org/10.1353/lan.1974.0010

2. Levinson SC, Torreira F. Timing in turn-taking and its implications for processing models of
language. Front Psychol. 2015;6: 731. doi: https://doi.org/10.3389/fpsyg.2015.00731

3. Stivers T, Enfield NJ, Brown P, Englert C, Hayashi M, Heinemann T, et al. (2009).
Universals and cultural variation in turn-taking in conversation. Proc Natl Acad Sci
USA. 2009;106(26): 10587–10592. doi: https://doi.org/10.1073/pnas.0903616106

4. Levinson SC. Turn-taking in human communication – Origins and implications for language
processing. Trends in Cogn Sci. 2016;20(1): 6–14. doi:
https://doi.org/10.1016/j.tics.2015.10.010

5. Pika S, Wilkinson R, Kendrick KH, Vernes SC. Taking turns: Bridging the gap between
human and animal communication. Proc R Soc Lond B Biol Sci. 2018;285: 20180598.
doi: https://doi.org/10.1098/rspb.2018.0598

6. Lemasson A, Gandon E, Hausberger M. Attention to elders' voice in non-human primates.
Biol Lett. 2010;6(3): 325–328. doi: https://doi.org/10.1098/rsbl.2009.0875

7. Levréro F, Touitou S, Frédet J, Nairaud B, Guéry J, Lemasson A. Social bonding drives vocal
exchanges in Bonobos. Sci Rep. 2019;9: 711. doi: https://doi.org/10.1038/s41598-018-
36024-9

8. Takahashi DY, Narayanan DZ, Ghazanfar AA. Coupled oscillator dynamics of vocal turn-
taking in monkeys. Curr Biol. 2013;23(21): 2162–2168. doi:
https://doi.org/10.1016/j.cub.2013.09.005

9. Grijseels DM, Fairbank DA, Miller CT. A model of marmoset monkey vocal turn-taking.
Proc R Soc B. 2024;291(2026):20240150. https://doi.org/10.1098/rspb.2024.0150

10. Koda H. Flexibility and context-sensitivity during the vocal exchange of coo calls in wild
Japanese macaques (Macaca fuscata yakui). Behaviour. 2004;141(10): 1279–1296. doi:
https://doi.org/10.1163/1568539042729685

11. Miller CT, Wang X. Sensory-motor interactions modulate a primate vocal behavior:

Antiphonal calling in common marmosets. J Comp Physiol A. 2006;192(1): 27–38. doi:

https://doi.org/10.1007/s00359-005-0043-z

12. Sugiura H. Temporal and acoustic correlates in vocal exchange of coo calls in Japanese

macaques. Behaviour. 1993;24(3–4): 207–225. doi:

https://doi.org/10.1163/156853993X00588

13. Sugiura H. Adjustment of temporal call usage during vocal exchange of coo calls in

Japanese Macaques. Ethology. 2007;113(6): 528–533. doi:

https://doi.org/10.1111/j.1439-0310.2007.01349.x

14. Takahashi DY, Fenley AR, Ghazanfar AA. Early development of turn-taking with parents

shapes vocal acoustics in infant marmoset monkeys. Philos Trans R Soc Lond B Biol

Sci. 2016;371(1693): 20150370. doi: https://doi.org/10.1098/rstb.2015.0370

15. Yamaguchi C, Izumi A, Nakamura K. Temporal rules in vocal exchanges of phees and trills

in common marmosets (Callithrix jacchus). Am J Primatol. 2009;71(7): 617–622. doi:

https://doi.org/10.1002/ajp.20697

16. Bateson MC. Mother-infant exchanges: The epigenesis of conversational interaction. Ann N

Y Acad Sci. 1975;263(1): 102-113. doi: https://doi.org/10.1111/j.1749-

6632.1975.tb41575.x

17. Bosch LT, Oostdijk N, Boves L. On temporal aspects of turn taking in conversational

dialogues. Speech Commun. 2005;47: 80-86. doi:

https://doi.org/10.1016/j.specom.2005.05.009

18. Hilbrink EE, Gattis M, Levinson SC. Early developmental changes in the timing of turn-

taking: A longitudinal study of mother–infant interaction. Front Psychol. 2015;6: 1492.

doi: https://doi.org/10.3389/fpsyg.2015.01492

19. Yosida S, Kobayasi KI., Ikebuchi M, Ozaki R, Okanoya K. Antiphonal vocalization of a

subterranean rodent, the naked mole-rat (Heterocephalus glaber). Ethology.

2007;113(7): 703–710. doi: https://doi.org/10.1111/j.1439-0310.2007.01371.x

20. Yosida S, Okanoya K. Naked mole-rat is sensitive to social hierarchy encoded in antiphonal

vocalization. Ethology. 2009;115(9): 823–831. doi: https://doi.org/10.1111/j.1439-

0310.2009.01677.x

21. Carter GG, Skowronski MD, Faure PA, Fenton B. Antiphonal calling allows individual

discrimination in white-winged vampire bats. Anim Behav. 2008;76(4): 1343–1355.

doi: https://doi.org/10.1016/j.anbehav.2008.04.023

22. Leong KM, Ortolani A, Burks KD, Mellen JD, Savage A. Quantifying acoustic and

temporal characteristics of vocalizations for a group of captive African elephants

Loxodonta Africana. Bioacoustics. 2003;13(3): 213–231. doi:

https://doi.org/10.1080/09524622.2003.9753499

23. Soltis J, Leong K, Savage A. African elephant vocal communication I: Antiphonal calling

behaviour among affiliated females. Anim Behav. 2005;70(3): 579–587. doi:

https://doi.org/10.1016/j.anbehav.2004.11.015

24. Demartsev V, Strandburg-Peshkin A, Ruffner M, & Manser M. Vocal turn-taking in

Meerkat group calling sessions. Curr Biol. 2018;28(22): 3661-3666.e3. doi:

https://doi.org/10.1016/j.cub.2018.09.065

25. Caldwell MC, Caldwell DK. Vocalization of naive captive dolphins in small groups.

Science. 1968;159 (3819): 1121-1123. doi:

https://doi.org/10.1126/science.159.3819.1121

26. Janik VM. Whistle matching in wild bottlenose dolphins. Science. 2000;289(5483): 1355–

1357. doi: https://doi.org/10.1126/science.289.5483.1355

27. Mishima Y, Morisaka T, Ishikawa M, Karasawa Y, Yoshida Y. Pulsed call sequences as

contact calls in Pacific white-sided dolphins (Lagenorhynchus obliquidens). J Acoust

Soc Am. 2019;146(1): 409–424. doi: https://doi.org/10.1121/1.5116692

28. Morisaka T, Yoshida Y, Akune Y, Mishima H, Nishimoto S. (2013). Exchange of

"signature" calls in captive belugas (Delphinapterus leucas). J Ethol. 2013;31(2): 141–

149. doi: https://doi.org/10.1007/s10164-013-0358-0

29. Nakahara F, Miyazaki N. Vocal exchanges of signature whistles in bottlenose dolphins

(Tursiops truncatus). J Ethol. 2011;29: 309–320. doi: https://doi.org/10.1007/s10164-

010-0259-4

30. Schulz TM, Whitehead H, Gero S, Rendell L. Overlapping and matching of codas in vocal

interactions between sperm whales: Insights into communication function. Anim Behav.

2008;76(6): 1977–1988. doi: https://doi.org/10.1016/j.anbehav.2008.07.032

31. Catchpole CK, Slater PJB. Bird Song: Biological Themes and Variations. 2nd ed.

Cambridge: Cambridge University Press; 2008. doi:

https://doi.org/10.1017/CBO9780511754791

32. Hyman J. Countersinging as a signal of aggression in a territorial songbird. Anim Behav.

2003;65(6); 1179-1185. doi: https://doi.org/10.1006/anbe.2003.2175

33. Fortune ES, Rodríguez C, Li D, Ball GF, Coleman MJ. Neural mechanisms for the

coordination of duet singing in wrens. Science. 2011;334(6056): 666-670. doi:

https://doi.org/10.1126/science.1209867

34. Kondo N, Watanabe S, Izawa E. A temporal rule in vocal exchange among large-billed

crows Corvus macrorhynchos in Japan. Ornithol Sci. 2010;9(1): 83–91. doi:

https://doi.org/10.2326/osj.9.83

35. Chaiken M. The ontogeny of antiphonal calling in European starlings. Dev Psychobiol.

1990;23(3): 233–246. doi: https://doi.org/10.1002/dev.420230304

36. Dreiss AN, Ruppli CA, Oberli F, Antoniazza S, Henry I, Roulin A. Barn owls do not

interrupt their siblings. Anim Behav. 2013;86(1): 119–126. doi:

https://doi.org/10.1016/j.anbehav.2013.04.019

37. Benichov JI, Benezra SE, Vallentin D, Globerson E, Long MA, Tchernichovski O. The

forebrain song system mediates predictive call timing in female and male zebra finches.

Curr Biol. 2016;26(3): 309-318. doi: https://doi.org/10.1016/j.cub.2015.12.037

38. Benichov JI, Vallentin D. Inhibition within a premotor circuit controls the timing of vocal

turn-taking in zebra finches. Nat Commun. 2020;11: 211. doi:

https://doi.org/10.1038/s41467-019-13938-0

39. D'Amelio PB, Trost L, ter Maat A. Vocal exchanges during pair formation and maintenance

in the zebra finch (Taeniopygia guttata). Front Zool. 2017;14(1): 13. doi:

https://doi.org/10.1186/s12983-017-0197-x

40. Ter Maat A, Trost L, Sagunsky H, Seltmann S, Gahr M. Zebra finch mates use their

forebrain song system in unlearned call communication. PLoS ONE. 2014;9(10):

e109334. doi: https://doi.org/10.1371/journal.pone.0109334

41. Stowell, D., Gill, L., & Clayton, D. Detailed temporal structure of communication networks

in groups of songbirds. J R Soc Interface. 2016;13(119): 20160296. doi:

https://doi.org/10.1098/rsif.2016.0296

42. Boinski S, Garber PA. On the move: How and why animals travel in groups. Chicago:

University of Chicago Press; 2000.

43. Frigerio D, Weiss B, Kotrschal K. Spatial proximity among adult siblings in greylag geese

(Anser anser): evidence for female bonding? Acta Ethol. 2001;3(2): 121-125. doi:

https://doi.org/10.1007/s102110000028

44. Goodwin D. Estrildid finches of the world. New York: Cornell University Press; 1982.

45. Furutani A, Mori C, Okanoya K. Trill-calls in Java sparrows: Repetition rate determines the

category of acoustically similar calls in different behavioral contexts. Behav

Proc. 2018;157: 68-72. doi: https://doi.org/10.1016/j.beproc.2018.08.010

46. Perez EC, Fernandez MS, Griffith SC, Vignal C, Soula HA. Impact of visual contact on

vocal interaction dynamics of pair-bonded birds. Anim Behav. 2015;107: 125-137. doi:

https://doi.org/10.1016/j.anbehav.2015.05.019

47. Lisa K. Yang Center for Conservation Bioacoustics at the Cornell Lab of Ornithology.

Raven Pro: Interactive Sound Analysis Software (Version 1.6) [Computer

software].2019. Ithaca, NY: The Cornell Lab of Ornithology.

Available: https://ravensoundsoftware.com

48. Boersma P, Weenink D. Praat: Doing phonetics by computer (version 6.3.10) [Computer

software]. 2023. Available: http://www.praat.org

49. Scrucca L, Fop M, Murphy TB, Raftery AE. mclust 5: Clustering, Classification and

Density Estimation Using Gaussian Finite Mixture Models. R Journal. 2016;8(1): 289.

doi: https://doi.org/10.32614/RJ-2016-021

50. R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria:

R Foundation for Statistical Computing; 2023. Available: https://www.R-project.org/

51. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using **lme4**.

J Stat Softw. 2015;67(1): 1-48. doi: https://doi.org/10.18637/jss.v067.i01

52. Barton K. MuMIn: Multi-model inference. 2023. Available: https://CRAN.R-

project.org/package=MuMIn

53. Lindström Å. Finch flock size and risk of hawk predation at a migratory stopover site. Auk.

1989;106(2): 225-232. doi: https://doi.org/10.1093/auk/106.2.225

54. Evans MHR, Lihou KL, Rands, SA. Black-headed gulls synchronise their activity with their

nearest neighbours. Sci Rep. 2018;8(1): 9978. doi: https://doi.org/10.1038/s41598-018-

28378-x

55. Fernández-Juricic E, Siller S, Kacelnik A. Flock density, social foraging, and scanning: An

experiment with starlings. Behav Ecol. 2004;15(3): 371–379. doi:

https://doi.org/10.1093/beheco/arh017

56. Hemelrijk CK, Van Zuidam L, Hildenbrandt H. What underlies waves of agitation in

starling flocks. Behav Ecol Sociobiol. 2015;69(5): 755–764. doi:

https://doi.org/10.1007/s00265-015-1891-3

57. Bajec IL, Heppner FH. Organized flight in birds. Anim Behav. 2009;78(4): 777–789. doi:

https://doi.org/10.1016/j.anbehav.2009.07.007

58. Arnold F, Staniszewski MS, Pelzl L, Ramenda C, Gahr M, Hoffmann S. Vision and vocal

communication guide three-dimensional spatial coordination of zebra finches during

wind-tunnel flights. Nat Ecol Evol. 2022;6(8): 1221-1230. doi:

https://doi.org/10.1038/s41559-022-01800-4

59. Blaich CF, Steury KR, Petttengill P, Mahoney KT, Guha A. Temporal patterns of contact

call interactions in pair-bonded domestic Zebra Finches (Taeniopygia guttata). Bird

Behav. 1996;11(2):59–69.

A

Exchange transition

Inter-call interval of exchange transition (*te*)

Repeated transition

"wait" for response

Inter-call interval of repeated transition (*tr*)

B

Exchange transition (*te*)   Repeated transition (*tr*)

Inter-call interval

A

PRE phase

Microphone (Channel 1)   Microphone (Channel 2)

10cm

38cm

58cm

31cm

F2F phase

POST phase

B

0.1 sec

Channel1

Channel2

Ch.1

Ch.2

Call sequence

Ch.1   Ch.1   Ch.2

Repeated transition   Exchange transition

C

Session 1   →   Pseudo session 1

Session 2   →   Pseudo session 2

A