Improving the Robustness of Phylogenetic Independent

Contrasts: Addressing Abrupt Evolutionary Shifts with Outlier-

and Distribution-Guided Correlation

Zheng-Lin Chen, Rui Huang, Hong-Ji Guo and Deng-Ke Niu[*]

MOE Key Laboratory for Biodiversity Science and Ecological Engineering, College of Life

Sciences, Beijing Normal University, Beijing, 100875, China

Correspondence to be sent to: Deng-Ke Niu, College of Life Sciences, Beijing Normal

University, Beijing, 100875, China; Emails: dkniu@bnu.edu.cn, dengkeniu@hotmail.com

ABSTRACT

Traditional phylogenetically aware correlation methods perform well under gradual

evolutionary processes. However, abrupt evolutionary shifts—or macroevolutionary jumps,

characteristic of punctuated evolution—can produce extreme phylogenetically independent

contrasts (PIC), leading to inflated false positives or increased false negatives in trait

correlation analyses. We introduce O(D)GC (Outlier- and Distribution-Guided Correlation), a

flexible workflow that identifies outliers in PICs using a distribution-free boxplot criterion

and applies Spearman correlation whenever influential outliers are detected. If no outliers are

detected, Pearson correlation is used—automatically for large datasets ($n \geq 30$), or guided by

normality testing in smaller samples. We systematically compared PIC-O(D)GC with five

widely applied phylogenetic correlation methods—PIC-Pearson, PIC-MM, PGLS

(phylogenetic generalized least squares), MR-PMM (multi-response phylogenetic mixed

model), and Corphylo—on 322,000 simulated datasets spanning five evolutionary scenarios

(two shift settings: single-trait shifts and dual-trait co-directional jumps; and three no-shift

gradual evolution settings), including both fixed-depth and randomly located shifts, tested

across 11 shift or noise gradients, three tree sizes (16, 128, 256 tips), and both balanced and

random topologies. Overall, PIC-O(D)GC achieved error rates comparable to—or noticeably

higher than—those of PIC-MM, while yielding substantially lower error rates than most

alternative methods. Under no-shift conditions, it retained power similar to other methods.

Analyses of three empirical datasets likewise showed that PIC-O(D)GC and PIC-MM

corrected shift-induced distortions that misled conventional methods. Moreover, PIC-

O(D)GC offers a conceptually simple framework and incurs markedly lower computational

cost. By design, its correlation-only output provides less mechanistic detail than regression-

based approaches like PGLS. However, when paired with PIC diagnostics, this outlier-guided

strategy highlights evolutionary jumps, distinguishes coupled from decoupled shifts, and—

via clade partitioning or tip pruning—recovers background correlations, offering biologically

informative insights into how punctuated events interact with gradual trends in trait

evolution.

KEYWORDS

Phylogenetic comparative methods; Phylogenetically independent contrasts (PIC); Outlier-

and distribution-guided correlation (PIC-O(D)GC); Abrupt evolutionary shifts; Robust

regression; Trait correlation; Brownian motion (BM) evolution; Outlier detection.

INTRODUCTION

Phylogenetically closer species tend to exhibit more similar traits due to their shared
ancestry, resulting in statistical dependencies among samples (Felsenstein 1985, Garamszegi
2014, Cornwallis and Griffin 2024). Ignoring these statistical dependencies and using
conventional statistical methods can lead to erroneous conclusions. To address this challenge,
evolutionary biologists have developed a range of phylogenetic comparative methods that
explicitly account for these dependencies (Grafen 1989, Lynch 1991, Garamszegi 2014,
O'Meara 2016, Cornwallis and Griffin 2024, Dewar et al. 2025).

The earliest method was phylogenetically independent contrasts (PIC) (Felsenstein
1985). The core principle behind PIC is that, while species at the tips of a phylogenetic tree
are interdependent due to shared ancestry, the evolutionary changes along each branch can be
treated as statistically independent events. In this context, PIC refers to the contrast values
calculated between species pairs along the phylogenetic tree, inferring evolutionary changes
along pairs of branches and standardizing these changes by branch length (or time) to
generate transformed data points. In traditional applications, these PIC values are analyzed
using ordinary least squares (OLS) regression or, equivalently under linear assumptions,
Pearson correlation, to assess associations between traits. This transformation renders
evolutionary data more suitable for conventional statistical methods by removing the
phylogenetic non-independence. Thus, PIC can refer both to the contrast values calculated
from the phylogeny and to the subsequent OLS regression analysis applied to these values.

Applying OLS regression to PIC values assumes that traits evolve according to a
Brownian motion (BM) model, under which trait variation accumulates gradually and
randomly over time. While the computation of PIC values does not strictly require a BM
model, the statistical justification for treating PIC values as independent and identically
distributed (i.i.d.) variables in regression analyses typically relies on the BM assumption or
its extensions. Under this workflow, OLS regression through the origin on PIC values yields
slope estimates, standard errors, and p-values that are numerically identical to those obtained
from phylogenetic generalized least squares (PGLS), assuming BM holds (Rohlf 2001,
Blomberg et al. 2012). PGLS generalizes this approach by directly modeling the phylogenetic
covariance structure through a variance–covariance matrix of the residuals in the linear
model, thereby accommodating phylogenetic autocorrelation among observations (Martins
and Hansen 1997, Pagel 1997, Pagel 1999). This adjustment helps reduce false positives in

hypothesis testing due to shared evolutionary history. Although some PGLS models, such as

Pagel's λ, Ornstein-Uhlenbeck (OU), and Early-burst (EB) models, can handle heterogeneous

evolutionary rates (Hansen 1997, Freckleton et al. 2002, Harmon et al. 2010, Ho and Ane

2014), they fundamentally assume gradual trait changes (Landis et al. 2013). This assumption

can be problematic when dealing with evolutionary jumps caused by singular events.

During biological evolution, certain branches may experience rare and dramatic trait

changes (Uyeda et al. 2011, Landis et al. 2013). Increasingly, studies and data have shown

that abrupt evolutionary shifts, or evolutionary jumps, have a significant impact on

biodiversity evolution (Landis and Schraiber 2017, Gao and Wu 2022, Sumner et al. 2023). A

typical example of this is Felsenstein's worst-case scenario (Felsenstein 1985, Uyeda et al.

2018), where two traits undergo a significant jump near the root of the phylogenetic tree

while evolving independently on other branches. Even if this dramatic change occurred only

once, it could still lead to erroneous conclusions of significant correlation when using

traditional regression-based PIC analyses or PGLS models (Uyeda et al. 2018).

Recently, more studies have focused on identifying jump events and providing more

reasonable explanations for the processes underlying trait evolution. Uyeda et al. (2018)

underscore the need for models like the singular events model, which can accommodate the

complexities and unpredictabilities inherent in evolutionary biology. Some studies suggest

that jump events cause trait changes to exhibit a "fat-tails" distribution, using Levy processes

to handle this distribution (Landis et al. 2013, Duchen et al. 2017). Additionally, LASSO-

based methods within models like OU have been applied to pinpoint shifts along specific

branches of phylogenetic trees (Khabbazian et al. 2016, Bastide et al. 2018, Zhang et al.

2024). Others propose that abrupt shift events lead to outliers, using robust regression to

reduce model sensitivity to violations (Slater and Pennell 2014, Adams et al. 2024). A variety

of mixed-model approaches have also been explored to address model deviations and trait

interdependence (Ives and Helmus 2011, Westoby et al. 2023, Halliwell et al. 2025).

Evolutionary shifts, especially those occurring in branches close to the root of the

phylogenetic tree, typically do not manifest directly as outliers in the trait data but instead

appear as outliers in the PIC values. As a result, robust statistical methods designed to handle

outliers in the original trait data may not be as effective in addressing evolutionary shifts.

Adams et al. (2024) demonstrated that when four different estimators (L1, S, M, and MM)

were integrated into both PGLS and PIC analysis, methods like PIC-MM and PIC-S

performed significantly better in detecting evolutionary relationships, especially when

accounting for outliers. Thus, computing PIC values is essential for effective analysis of the

relationships between traits with evolutionary shifts.

Dramatic changes in some evolutionary branches can result in several outliers in the PIC

dataset (Uyeda et al. 2018), distorting the statistical results of parametric tests. While

nonparametric approaches are generally more robust to such outliers because they do not rely

on specific distributional assumptions, they are not routinely incorporated into traditional

phylogenetic correlation workflows.

To address this gap, we propose a hybrid workflow, Outlier-Guided Correlation (OGC),

for analyzing pairwise trait relationships under diverse evolutionary conditions. The core idea

of OGC is to integrate the PIC framework with a data-driven mechanism that evaluates the

presence of outliers in trait contrasts and selects an appropriate correlation measure

accordingly. For larger datasets, OGC selects Spearman or Pearson correlation depending on

whether outliers are detected. For smaller datasets, where non-normality can significantly

impact results, the workflow incorporates both normality testing and outlier detection,

referred to as Outlier- and Distribution-Guided Correlation (ODGC). Together, these

approaches form a unified workflow collectively referred to as O(D)GC, designed to address

the challenges posed by varying data conditions.

It should be noted that, whereas regression-based methods such as PGLS are commonly

used for both parameter estimation and hypothesis testing, the O(D)GC workflow is

intentionally restricted to null hypothesis testing, focusing on whether significant pairwise

trait correlations are present. Readers should interpret its results within this scope.

MATERIALS AND METHODS

Phylogenetic Tree Construction and Overview of Simulation Strategy

We categorize the construction of phylogenetic trees into two types. The first is the fixed-

balanced tree scenario, where all non-leaf nodes have left and right subtrees with identical

topology and depth, forming a fully symmetric structure. In this setting, a single fixed-

balanced phylogenetic tree with 128 species is used as the basis for all trait simulations.

Specifically, traits are simulated under combinations of evolutionary scenarios and

experimental conditions reflecting different levels of stochastic variation or evolutionary

shifts, with 1,000 replicates per condition, using different random seeds while keeping the

phylogenetic tree fixed. Consequently, for each trait evolutionary scenario-gradient

combination, 1,000 datasets are generated based on the same phylogenetic structure. To

assess the influence of tree size, additional analyses were conducted using 16- and 256-

species fixed-balanced trees, following the same simulation protocol.

The second category is a constrained random-tree scenario (hereafter referred to as the

random-tree scenario), in which a new phylogenetic tree is generated prior to each trait

simulation. In this setting, the root ancestor is forced to bifurcate into two subtrees of equal

size that share an identical topology. This topology is randomly generated under a coalescent

model for each simulation replicate, such that randomness is introduced at the level of

subtree topology. For each replicate, a phylogenetic tree is first constructed, followed by trait

simulation under a given evolutionary scenario and gradient condition. This pair of

operations—tree construction and trait simulation—is repeated 1,000 times using different

random seeds, ensuring variation in both tree topology and simulated trait values.

Subsequently, we evaluated the performance of each trait-correlation inference method

across all evolutionary scenario–gradient combinations using two complementary

benchmarks for defining reference correlation outcomes. For each simulated dataset, the

inferred correlation result produced by a given method was compared with the corresponding

benchmark outcome, and method performance was summarized across the 1,000 replicates

for each scenario–gradient combination.

Depending on the benchmark definition, performance of methods was quantified using

appropriate aggregate metrics, including overall correctness and error-based measures.

Detailed definitions of the benchmarks and the corresponding evaluation metrics are

provided in the following sections.

Felsenstein's Worst-Case Scenario: Deep-root Directional Shifts

Following previous simulation studies of directional trait shifts (Uyeda et al. 2018), we

simulated abrupt shifts occurring on a single deep branch near the root of the phylogeny,

representing a classical worst-case scenario for PIC-based methods. In this setting, trait

evolution follows a BM process on all other branches, but an abrupt directional change is

introduced on one of the branches closest to the root, a configuration known to generate

clusters of extreme contrasts and to challenge standard phylogenetic comparative approaches.

Trait evolution under BM was simulated with unit evolutionary rate ($\sigma^2 = 1$). Directional

shifts were implemented as multiplicative scaling factors applied to trait values rather than

additive offsets. Shift magnitudes ranged from 4^{-3} to 4^7, where a magnitude of $4^0 = 1$

corresponds to the baseline condition with no effective shift. Positive exponents indicate trait

upsizing, whereas negative exponents indicate downsizing. Each shift magnitude was

179 simulated 1,000 times. Random seeds were controlled to ensure reproducibility and, where

180 appropriate, independence between traits.

181  To implement the deep-root shift, the phylogenetic tree was split at the root into two

182 subtrees—a background subtree and a shift subtree—using the extract.clade() function in the

183 **ape** package (version 5.8) (Paradis and Schliep 2019). The background subtree was assigned

184 a baseline root value of 1, whereas the shift subtree was assigned a root value scaled by the

185 specified shift magnitude ($1 \times$ shift magnitude). Trait evolution was then simulated

186 independently on each subtree under BM using the rTraitCont() function from the **ape**

187 package.

189 *Two-trait shared shift (independent traits).*—To evaluate false positive behavior under shared

190 directional shifts, we simulated two independent traits (X_1 and X_2) on the same phylogeny

191 within a regression framework, $X_2 = \beta X_1 + \varepsilon$, with $\beta = 0$. Under this setting, ε itself constitutes

192 the second trait X_2. Both traits evolved independently under a Brownian motion model with

193 identical variance ($\sigma^2 = 1$) and the same root value (1), and were therefore uncorrelated by

194 construction. An abrupt shift of equal magnitude was introduced on the same deep branch in

195 both traits, while the remainder of the tree followed background BM evolution. Independent

196 random seeds were used for X_1 and X_2 to ensure the absence of any true association. This

197 design allowed us to assess whether phylogenetic methods spuriously detect correlations

198 when none exist.

200 *Single-trait shift in the predictor (correlated traits).*—To evaluate false negative behavior,

201 we simulated two positively correlated traits with $X_2 = \beta X_1 + \varepsilon$, where $\beta = 1$ and ε followed a

202 normal distribution with mean 0 and variance 0.01, following the phylogenetic regression

203 framework of Revell (2010) and Mazel et al. (2016). An abrupt directional shift was

204 introduced only in the predictor trait X_1 on the same deep branch using the same set of shift

205 magnitudes as described above, while X_2 remained unchanged. To preserve the predefined

206 correlation structure, X_1 was first simulated across the entire tree, X_2 was constructed as a

207 function of X_1, and X_1 was then re-simulated only within the shift subtree using the same

208 random seed, with the only change being that the subtree root value was shifted by the

209 specified magnitude (i.e., the shift magnitude was added at the subtree root). The final X_1 was

210 assembled by combining background and shift-subtree values, whereas X_2 remained

211 unchanged. This design allowed us to quantify how often true correlations were masked by

212 directional shifts affecting only one trait.

Simulation with Randomly Located Trait Shifts

To evaluate method performance under more biologically realistic conditions, we extended

the simulation framework by allowing the branch experiencing a directional shift to be

randomly selected across the phylogeny. In contrast to shifts fixed at deep branches, this

design captures scenarios in which shifts arise on shallow internal branches or terminal

lineages.

For each replicate, a single branch was sampled uniformly at random from the phylogeny

and assigned a directional shift. Except for the location of the shift, all simulation settings—

including trait evolutionary models, shift implementation, shift magnitudes, and replication

scheme—were identical to those used in the deep-root simulations described above.

As in the deep-root scenario, we considered both two-trait shared shifts in independent

traits (to assess false positive behavior) and single-trait shifts in the predictor for correlated

traits (to assess false negative behavior), with shifts applied to the randomly selected branch

following the same procedures outlined above.

Trait Shifts Introduced at Different Phylogenetic Depths

To evaluate whether method performance depends on the phylogenetic depth at which a

directional shift occurs, we conducted an additional set of simulations on the 128-species

fixed-balanced tree, in which shifts were introduced systematically at predefined

phylogenetic depths. In contrast to the worst-case and random-position-shift simulations, this

analysis was designed as a controlled diagnostic experiment to isolate the effect of shift depth

while holding all other factors constant. Based on preliminary results from the deep-root

(worst-case) simulations (see Results), which indicated that all methods are largely

insensitive to downsizing shifts, we focused exclusively on a representative large-magnitude

upsizing shift that poses the greatest challenge for phylogenetic correlation inference.

Specifically, we examined a shift magnitude of 4^5, which provides a strong but non-extreme

perturbation suitable for isolating depth-dependent effects. For each phylogenetic depth, a

single shift was introduced on one randomly selected branch at that depth. At each shift

depth, simulations were conducted under both the dual-trait shared-shift and the single-trait

(X_1-only) shift scenarios, with 1,000 independent replicates for each scenario.

Simulated Traits Without Shifts: Fixed Correlation with Varying Noise

Although the primary focus of this study is on evaluating PIC-O(D)GC under shift-driven

evolutionary scenarios, we also considered trait simulations without abrupt shifts to assess

whether the workflow behaves comparably to established methods under gradual

evolutionary dynamics. To provide such a baseline, we followed the core simulation

framework used by Revell (2010) and Mazel et al. (2016), in which paired traits X_1 and X_2

were generated under a linear generative model,

$$X_2 = \beta X_1 + \varepsilon,$$

with $\beta = 1$, such that a structural dependence between the two traits is biologically encoded.

The noise term ε was assigned varying variance to modulate the strength of the realized

association. Under this design, these simulations primarily function as power analyses,

assessing how effectively different methods recover an intended association as stochastic

variation increases, while acknowledging that sufficiently large noise can render some

replicates statistically indistinguishable from uncorrelated data.

To reflect different sources of phylogenetic signal under gradual evolutionary dynamics

(i.e., without abrupt shifts), we considered three classes of simulation scenarios that differ in

whether phylogenetic structure arises from the focal trait X_1, from the noise term ε, or from

both.

In the BM–BM scenario, both X_1 and ε evolve under BM, with X_1 having a variance of 1

and ε having variance d. Accordingly, both the signal and noise components carry

phylogenetic structure, and the realized association between X_1 and X_2 can vary substantially

across replicates as noise variance increases.

In the BM–Norm scenario, X_1 evolves under BM with variance 1, whereas ε is drawn

from an independent and identically distributed normal distribution with mean 0 and variance

d. Because the noise term is phylogenetically unstructured, this scenario satisfies the

assumptions of ordinary least squares (OLS) regression and therefore represents the only

setting in which OLS is theoretically appropriate and included for comparison (Revell 2010).

In the Norm–BM scenario, X_1 follows a normal distribution with mean 0 and variance 1,

while ε evolves under BM with variance d. Here, phylogenetic signal arises solely from the

noise term, allowing us to examine how phylogenetically structured error can obscure an

otherwise simple linear relationship.

Across all three scenarios, the noise variance parameter d took values of 4^{-3}, 4^{-2}, 4^{-1}, 4^0,

4^1, 4^2, 4^3, 4^4, 4^5, 4^6, and 4^7. For each scenario and value of d, we conducted 1,000 replicate

simulations, resulting in a total of 33,000 datasets per phylogenetic tree.

We did not include the fourth scenario described by Revell (2010), in which both X_1 and

ε lack phylogenetic structure. Because this study focuses on evaluating method behavior in

settings where phylogenetic signal is present in either the traits or the residual variation,

explicitly simulating a fully non-phylogenetic scenario was not necessary for our purposes.

284   *Empirical Datasets*

To complement the simulation analyses and evaluate the relative performance of PIC-

O(D)GC against other phylogenetic comparative methods under real biological conditions,

we applied these methods to three empirical datasets that may contain abrupt evolutionary

shifts. Two of them were previously analyzed by Adams et al. (2024) in a study of

phylogenetically robust regression.

The first dataset was obtained from the supplementary material of Makino and Kawata

(2019), and includes propagule size, genome size, and a phylogenetic tree for 44 invertebrate

species. Following the original study, both traits were $\log_{10}$-transformed prior to analysis to

reduce skewness and improve normality assumptions. Propagule size in this context refers to

the physical size (cm) of a reproductive unit that is separated from the parent and capable of

developing into a new individual. In ecological and evolutionary studies, propagule size is

widely used as a key indicator of an organism's reproductive strategy, particularly along the

r/K selection continuum (Romiguier et al. 2014). The phylogenetic tree provided by the

authors was used without modification.

The second dataset was originally compiled by Scales et al. (2009) and includes the

relative proportions of fast glycolytic (FG) and slow oxidative (SO) muscle fibers in the limb

muscles of 22 lizard species. This dataset was later used by Uyeda et al. (2018) to illustrate

the limitations of phylogenetic comparative methods, particularly in cases involving abrupt

shifts or convergent trait evolution. We used the phylogenetic tree provided by Uyeda et al.

(2018).

The third dataset includes body mass and population density for 109 species of

Artiodactyla. Trait data were obtained from the PanTHERIA database (accessed June 4,

2025) (Jones et al. 2009). Species with missing values for either trait were excluded. Both

body mass (grams) and population density (individuals per km^2) were $\log_{10}$-transformed prior

to analysis to reduce skewness and better meet the assumptions of linear models and

comparative methods. The phylogenetic tree used for the Artiodactyla clade was extracted

from the time-scaled maximum clade credibility tree for mammals developed by Upham et

al. (2019), available at VertLife.org (accessed on June 4, 2025).

Statistical Methods Compared

To evaluate the performance of PIC-O(D)GC examining the evolutionary relationship

between X_1 and X_2 under both gradual evolution and scenarios involving abrupt evolutionary

shifts, we compared it with several previously established phylogenetic comparative

methods:

1. PIC-Pearson: Pearson correlation applied to PIC values; because it yields the same

direction and significance inferences as an OLS regression on the same PICs, running a

separate PIC-OLS analysis would be redundant, so we used only PIC-Pearson.

2. PGLS: We fitted five alternative evolutionary covariance models: BM (Felsenstein

1985), Pagel's λ (Pagel 1999), OU fixed, OU random (Hansen 1997), and EB (Harmon et al.

2010). For each simulated dataset, we used the Akaike information criterion (AIC) to identify

the best-fitting covariance structure. PGLS results were then reported under the model with

the lowest AIC.

3. Robust phylogenetic regression with both PIC and PGLS frameworks: For each

framework, we utilized four robust estimators: L1, S, M, and MM. These were implemented

as PIC-L1, PIC-S, PIC-M, and PIC-MM for PIC-based methods, and PGLS-L1, PGLS-S,

PGLS-M, and PGLS-MM for PGLS-based methods (Adams et al. 2024).

4. Corphylo: A method that calculates Pearson correlation coefficients for multiple

continuous traits, allowing users to incorporate phylogenetic signal and measurement errors

into the model (Zheng et al. 2009). This method assumes an OU process for trait evolution,

making it particularly suitable for traits that deviate from the BM model. Corphylo serves a

dual purpose: it enables both the detection of phylogenetic signals in individual traits and the

examination of correlations between multiple traits. A key feature of Corphylo is its ability to

include measurement errors of trait data, potentially improving accuracy under real-world

conditions.

5. MR-PMM: Multi-response phylogenetic mixed models (MR-PMM) extend

phylogenetic regression frameworks by jointly modeling multiple traits as responses,

allowing the decomposition of trait covariation into phylogenetic and residual components

(Westoby et al. 2023, Halliwell et al. 2025). Unlike conventional phylogenetic regression,

MR-PMM does not impose a directional predictor–response structure and is therefore well

suited for assessing trait correlations and co-evolution. Although our simulations involve

only two traits rather than higher-dimensional response vectors, MR-PMM is directly

applicable to this bivariate setting and represents a conceptually distinct framework for

evaluating phylogenetic trait associations.

6. Raw-Pearson: The standard Pearson correlation applied directly to the original trait

values, without any phylogenetic correction. Because Pearson (or its OLS equivalent)

assumes i.i.d. residuals, we report Raw-Pearson results only for the BM–Norm scenarios—

the only conditions in our simulations that satisfy this assumption. Although Pearson/OLS

methods are seldom used in phylogenetic comparative analyses due to the non-independence

created by shared ancestry, they remain a useful theoretical benchmark when their statistical

requirements are met (Revell 2010).

356   *Outlier- and Distribution-Guided Phylogenetic Correlation Analysis*

To address the influence of evolutionary shifts and outliers in phylogenetic comparative

analysis, we introduced a novel workflow: PIC-O(D)GC. In this method, the choice between

Pearson and Spearman correlation analysis is guided by the identification of outliers within

the PIC data, with additional consideration of the PIC value distribution in small-sample

settings.

Potential outliers in PICs were identified using the boxplot rule (Tukey 1977), flagging

values beyond $Q1 - 1.5 \times IQR$ or $Q3 + 1.5 \times IQR$, where $Q1$ and $Q3$ are the first and third

quartiles and $IQR = Q3 - Q1$ is the interquartile range. This nonparametric, distribution-free

rule—corresponding to the whiskers of the boxplot—provides a robust and widely used

definition of extreme observations, flagging about 0.7% of normally distributed data as

outliers (Hoaglin et al. 1986). Because the boxplot rule is intentionally conservative, it may

occasionally flag points spuriously due to random variation or over-flag observations in

skewed distributions. However, such misclassification has little practical impact in our

framework, as flagged datasets are analyzed using Spearman correlation, for which a small

number of additional flagged points primarily leads to a modest loss of power while

preserving monotonic associations. In contrast, failing to control even a single influential

outlier in a Pearson analysis can substantially distort, or even reverse, the estimated

correlation.

Outlier detection is performed at the level of individual observations: if a given sample is

flagged as an outlier in any trait, that observation is treated as an outlier overall, and the

corresponding dataset is analyzed using Spearman correlation; otherwise, Pearson correlation

is applied. This rule ensures robustness when one or both traits—whether at the same or

different phylogenetic positions—experience abrupt shifts, and, although not explored here,

the underlying logic can be readily generalized to multivariate settings, including partial

correlation analyses.

For larger datasets with 128 species (127 PIC values), Pearson correlation, despite its

reliance on normality assumptions, is generally robust to minor deviations from normality

due to the central limit theorem. In such cases, the impact of skewness and kurtosis is

reduced by the large sample size, ensuring reliable correlation estimates. Therefore, for these

datasets, the choice of Pearson or Spearman correlation was determined solely by the

presence or absence of outliers.

For smaller datasets with 16 species (15 PIC values), both normality testing and outlier

detection were conducted. This workflow, referred to as PIC-ODGC, dynamically selects the

appropriate correlation method based on data characteristics. If no outliers were detected and

normality assumptions were satisfied, the Pearson correlation was used. In cases where

outliers were present, or normality was violated, Spearman correlation was applied to

minimize the impact of extreme values on the results.

395   ***Benchmark I: Structural-Coupling–Based Evaluation (β-based)***

In simulation-based evaluations, trait associations are often benchmarked against the

structure explicitly specified in the generative model ($X_2 = \beta X_1 + \varepsilon$). Under this approach,

correlations are treated as present or absent according to whether the coupling parameter (β)

is encoded during data generation.

Accordingly, we implemented a generative-model–based benchmark in which simulation

replicates with a coupling parameter $\beta = 1$ were classified as true positives, whereas

replicates with $\beta = 0$ were classified as true negatives. For each replicate, method

performance was evaluated based on whether a statistically significant association between

the two traits was detected.

This benchmark directly assesses a method's ability to recover the structural relationship

encoded by the coupling parameter β in the simulation design and therefore provides a clear

reference for evaluating consistency with that aspect of the generative model. Performance

was summarized using class-specific error rates, including the false positive rate (FPR),

defined as the proportion of $\beta = 0$ replicates incorrectly classified as significant, and the false

negative rate (FNR), defined as the proportion of $\beta = 1$ replicates in which the encoded

relationship was not detected.

413   ***Benchmark II: Data-Pattern–Based Evaluation***

Beyond assumptions encoded in the generative model about trait coupling, a distinct

inferential question concerns whether an association specified during data generation is

statistically supported by the realized data. In stochastic evolutionary processes, the realized

relationship between traits is jointly determined by the coupling parameter β and evolutionary

noise ε, such that even strong generative coupling may yield weak or undetectable

associations when noise variance is large (see Supplementary Methods S1). In empirical data,

β and ε are not separately observable, and inference is therefore based on their combined

effect on realized trait patterns.

To address this distinction, we implemented an additional benchmarking framework that

evaluates whether association is statistically supported in the realized data. Under this

framework, benchmark labels were assigned at the replicate level according to whether

evidence for trait association was present in the realized data.

Specifically, for each simulation replicate, evidence for association was evaluated using

correlation tests applied to normalized branch-level trait changes. Because ancestral states are

recorded in the simulations, branch-level changes can be computed directly as $\Delta X_1/L$ and

$\Delta X_2/L$, where ΔX denotes trait change along each branch and L the corresponding branch

length. Under a Brownian motion evolutionary model, these normalized changes are

independently distributed across branches, providing a valid basis for conventional

correlation testing (Supplementary Methods S1). Replicates exhibiting significant correlation

($p < 0.05$) were classified as benchmark-positive, whereas replicates lacking significant

evidence were classified as benchmark-negative. Method performance was quantified using

classification accuracy, defined as the proportion of replicates for which inferred results

agreed with these data-driven benchmark labels.

This benchmark therefore targets a different inferential question than the β-based

framework. Whereas the β-based benchmark evaluates method performance relative to the

structural coupling specified during data generation, the data-pattern–based benchmark

assesses whether the combined effects of β, ε, and finite sample size yield statistically

supported association in the realized data (Ives 2022).

Programs and Packages

All simulations and analyses were performed in the R environment (version 4.1.3) (R Core

Team 2020) on a Linux system. The **phytools** package (version 2.3.0) (Revell 2024) and **ape**

package (version 5.8) (Paradis and Schliep 2019) were used for tree generation. Simulations

and PIC calculations were performed using the **ape** package in R. PIC values were calculated

with ape∷pic() under a Brownian motion assumption, using the standard contrast-based

pruning procedure rather than global ancestral-state reconstruction. Normality tests were

conducted using the Shapiro-Wilk test (Royston 1992) via the shapiro.test() function from the

stats package (version 4.3.1). Pearson and Spearman correlation analyses were performed

using stats∷cor.test(). All PGLS regressions were conducted using the phylolm() function

from the phylolm package (version 2.6.2) (Ho and Ane 2014). Corphylo was conducted using

ape∷corphylo() function (Zheng et al. 2009). MR-PMM was conducted using the

MCMCglmm() function from the MCMCglmm package (version 2.36) (Sorensen and

Gianola 2002, Hadfield 2010, Hadfield and Nakagawa 2010). Robust phylogenetic regression

were conducted using the Conduct.Robust_PhylogeneticRegression() function from ROBRT

package (version 0.0.0.9000) (Adams et al. 2024).

460   **RESULTS**

Impact of Abrupt Evolutionary Shifts on Data Patterns and Correlation Analysis

To illustrate PIC-ODGC's ability to handle abrupt evolutionary shifts, we simulated two traits

on a fixed-balanced phylogenetic tree with 16 species (Fig. 1a), considering two shift

configurations: (1) simultaneous shifts applied to both traits, which are uncorrelated, and (2)

shifts applied to X_1 only, where X_1 and X_2 are correlated. For each configuration, we selected

three shift levels—baseline, a weak shift, and a strong shift—as representative cases to

illustrate how shift magnitude affects trait distributions (e.g., group separation) and contrasts

(e.g., emergence of isolated outliers).

As shown in Figures 1d and 1f, when both traits undergo an abrupt shift near the root, the

raw trait values form two distinct clusters in the coordinate space, with the separation

between these clusters increasing as the shift magnitude becomes larger. In the PIC

coordinate system, the shared shift manifests as a single outlier in the PIC data, with the

degree of outlier deviation becoming more pronounced as the shift magnitude increases

(Figs. 1e and 1g).

When phylogenetic autocorrelation is ignored, directly applying Pearson or Spearman

correlation to the raw trait data frequently results in false positives, regardless of the shift

magnitude (Figs. 1b, 1d, and 1f). In contrast, controlling for phylogenetic autocorrelation by

using PIC values eliminates false positives when there is no effective shift or when the shift

is small (Figs. 1c and 1e). However, under large upsizing shifts, Pearson correlation becomes

highly sensitive to the outlier, leading to a false positive result and erroneously inferring a

significant positive correlation between X_1 and X_2, akin to non-phylogenetic methods.

Conversely, the Spearman correlation applied to the PIC data under large shifts avoids false

positives by correctly detecting no significant correlation between the two traits (Fig. 1g).

Figure 1. Effects of abrupt shifts on data pattern and correlation analysis. a) Schematic fixed-balanced

phylogenetic tree with 16 species; the thick branch marks the location of an abrupt evolutionary shift. b–c)

Baseline scenario for the dual-shift simulations shown in panels d–g. In this scenario, both traits X_1 and X_2

evolve independently under Brownian motion (BM) without an effective shift (baseline condition;

multiplicative shift factor = 1). d–e) and f–g) Dual-shift simulations in which both traits undergo an abrupt shift

on the marked branch, with increasing shift magnitudes (4-fold and 16-fold, respectively). h–i) Baseline

scenario for the single-trait shift simulation series shown in panels j–m. In this scenario, X_1 evolves under BM

without an effective shift (multiplicative shift factor = 1), and X_2 is generated from X_1 via a fixed linear

relationship with added noise. j–k) Moderate (4-fold) shift in X_1 only. l–m) Strong (64-fold) shift in X_1 only. In

all phylogenetically independent contrast (PIC) plots (c, e, g, i, k, m), the contrast corresponding to the branch

marked by the thick line in panel a is highlighted as an enlarged dot. In panels e, g, k, and m, this contrast

reflects an evolutionary shift; in panels c and i, it serves as a baseline reference. In the weak dual-shift dataset

(e), the highlighted contrast lies approximately 1.8 interquartile ranges (IQRs) above the third quartile (Q3) for

PIC X_1 and 2.1 IQRs above Q3 for PIC X_2. In the strong dual-shift dataset (g), this deviation increases markedly,

reaching approximately 8.8 and 7.8 IQRs above Q3 for PIC X_1 and PIC X_2, respectively. In the weak single-trait

shift dataset (k), the shifted trait shows only moderate deviation in PIC, remaining below the outlier threshold

(i.e., < 1.5 IQRs above Q3). In contrast, in the extreme single-trait shift case (m), PIC X_1 exhibits a pronounced

deviation of approximately 5.9 IQRs above Q3. Pearson and Spearman correlation tests: *** indicates $p <$

0.001; ** indicates $p < 0.01$; * indicates $p < 0.05$; ns indicates not significant ($p \geq 0.05$). Data used to generate

this figure are available in Supplementary Table S1.

In the second set of simulations, where only X_1 experiences an abrupt shift while X_2

follows unperturbed BM, the true correlation between the two traits remains intact outside the

shifted lineage. As in the dual-shift scenario, the raw data exhibit a visible cluster separation

as the shift magnitude increases (Figs. 1j and 1l). Despite the presence of a shift in only one

trait, the PIC data again display a prominent outlier under large upsizing (Fig. 1m). While

Pearson correlation applied to the PIC values is affected by this outlier and fails to detect the

existing correlation—resulting in a false negative (Fig. 1m)—Spearman correlation remains

robust across shift levels, consistently recovering the true positive association (Figs. 1i, 1k,

and 1m).

Together, these results show that large upsizing shifts can produce outliers in PIC data,

which in turn may lead to false negatives or false positives in Pearson-based correlation

analyses, depending on whether the shift affects one or both traits. While we simulated a

deep-branch shift to illustrate maximal distortion, equally large shifts on intermediate or

terminal branches can also generate extreme contrasts and thereby mislead mislead PIC-

based parametric analyses. This reflects an inherent limitation of traditional PIC approaches,

which weigh all contrasts equally without regard to their evolutionary context.

Refining Robust Regression Methods for Comparison with PIC-O(D)GC

To avoid visual and interpretive complexity, we restricted comparisons to representative

implementations within each methodological category rather than exhaustively presenting all

possible variants. Specifically, we evaluated robust regression methods spanning two

methodological types—PGLS- and PIC-based regression—combined with four robust

estimators (L1, S, M, and MM), and selected representative implementations based on

objective performance criteria derived from simulation results.

Under shift conditions, PIC-MM consistently ranked as the best or among the best-

performing robust estimators (Supplementary Result S1). In no-shift simulations, PIC-MM

continued to rank among the stronger-performing methods, although performance differences

across estimators were generally modest. These conclusions were consistent under two

benchmark criteria used to evaluate robust regression, suggesting the observed ranking is not

sensitive to a particular evaluation standard. In line with recent results reported by Adams et

al. (2024), we therefore retain PIC-MM as the primary robust comparator to PIC-O(D)GC.

PIC-OGC: Performance Across Evolutionary Shifts and No-Shift Regimes

We first compared the performance of PIC-OGC with five alternative methods using

simulations with evolutionary shifts on the 128-species fixed-balanced tree, focusing initially

on scenarios with simultaneous shifts in both traits.

Under Felsenstein's worst-case scenario with deep-root directional shifts (Fig. 2a), PIC-

OGC exhibited the strongest control of type I error, maintaining a false positive rate of zero

across all shift magnitudes examined. PIC-MM was similarly insensitive to increasing shift

magnitude, although its false positive rate was slightly higher than that of PIC-OGC,

remaining close to 0.05 across the entire gradient. Other methods—including PIC-Pearson,

which serves as the most direct comparator to PIC-OGC—also showed low false positive

rates under weak shifts. As shift magnitude increased, false positive rates for these methods

rose sharply, and at shift magnitudes $\geq 4^3$, all approached or reached one, indicating near-

complete loss of type I error control.

When trait shifts were introduced at randomly selected locations on the tree (Fig. 2b),

PIC-OGC no longer ranked as the best-performing method across the full shift gradient.

Under weak shifts, PIC-OGC, PIC-MM, and PIC-Pearson jointly exhibited the lowest false

positive rates. However, as shift magnitude increased, PIC-MM consistently maintained the

lowest false positive rate, whereas the false positive rate of PIC-OGC increased markedly,

reaching a maximum of approximately 0.26. The other methods—including PIC-Pearson—

showed rapidly escalating false positive rates with increasing shift magnitude, again

approaching one under strong shifts.

We next examined scenarios with univariate shifts affecting only X_1. Regardless of

whether shifts occurred under the worst-case (Fig. 2c) or randomly located design (Fig. 2d),

PIC-OGC and PIC-MM showed identical performance, maintaining false negative rates of

zero across all shift magnitudes. The remaining four methods also achieved near-zero false

negative rates under weak shifts, but diverged markedly as shift magnitude increased. Under

strong shifts, Corphylo exhibited the lowest false negative rates among these alternatives,

whereas the other methods showed substantially reduced sensitivity.

Figure 2. Performance of six methods across evolutionary shift and no-shift regimes simulated on a 128-species

fixed-balanced tree. Panels a–b report false positive rates under simultaneous shifts in both X_1 and X_2, and

panels c–d report false negative rates when only X_1 is shifted. Panels a and c correspond to Felsenstein's worst-

case scenario, in which directional shifts are introduced on deep-root branches, whereas panels b and d show

results from simulations with randomly located trait shifts. Panels e–g show performance under gradual

evolutionary scenarios without abrupt shifts, in which traits follow the relationship $X_2 = X_1 + \varepsilon$ with increasing

noise variance. Three evolutionary background conditions are considered: BM–BM (panel e), BM–Norm (panel

f), and Norm–BM (panel g). Shift and noise variance magnitudes are expressed on a 4^n scale, where the

horizontal axis shows only the exponent (n) rather than the full value. Additional details on the simulation

design, gradient configurations, and performance metrics are provided in the Materials and Methods. The data

used to generate this figure are available in Supplementary Table S2.

In addition to simulations with abrupt evolutionary shifts, we also examined method

performance under gradual evolutionary scenarios without discrete shifts. These analyses

were conducted on the 128-species fixed-balanced tree, with trait relationships defined as X_2

$= X_1 + \varepsilon$, where the noise term ε varied across 11 gradients of variance (Figs. 2e–g). Three

evolutionary background scenarios were considered: BM–BM, BM–Norm, and Norm–BM.

Across all three no-shift scenarios, overall performance patterns differed markedly from

those observed under shift conditions. In particular, MR-PMM exhibited a distinct and

inconsistent behavior, with its performance relative to other methods fluctuating noticeably

across noise levels and evolutionary settings, setting it apart from the other methods without

a consistent advantage or disadvantage.

With the exception of PGLS under the BM–Norm scenario—where its performance was

clearly degraded—the remaining five methods showed broadly similar behavior across noise

gradients. Under gradual evolution, differences among these methods were modest, and no

approach consistently emerged as clearly superior or inferior across scenarios. This

convergence in performance contrasts with the pronounced method-specific differences

observed under abrupt shifts, underscoring that evolutionary shifts, rather than background

noise alone, are the primary drivers of method divergence.

In the BM–Norm scenario, where the data-generating process conforms the assumptions

of standard linear regression, correlation analysis on raw trait values (Raw-Pearson,

equivalent to OLS) was included as a theoretical benchmark (Supplementary Fig. S1). Under

increasing noise levels, Raw-Pearson closely tracked the performance of PGLS, with both

methods forming a consistently low-performing group. This pattern is consistent with

theoretical expectations: when residuals are phylogenetically independent, PGLS effectively

reduces to OLS (Revell 2010, Symonds and Blomberg 2014).

604   ***Sensitivity of Methods to the Phylogenetic Location of Evolutionary Shifts***

The above results shown in Figures 2b and 3d summarize method performance under

randomly located trait shifts, effectively averaging behavior across all possible phylogenetic

positions. To further characterize method sensitivity to where a shift occurs on the tree, we

conducted an additional set of diagnostic simulations in which shifts were introduced

systematically at each phylogenetic depth. Based on the separation among methods observed

in Figure 2, we selected a representative shift magnitude of 4^5, and applied shifts

independently at each branch level of the 128-species fixed-balanced tree.

Under simultaneous shifts in both traits (Fig. 3a), both PIC-MM and PIC-OGC showed

reduced robustness to shifts occurring near the tips compared to shifts on deeper branches

closer to the root. This pattern indicates that terminal shifts pose a greater challenge for

robust phylogenetic methods than deep-root shifts. Notably, PIC-OGC exhibited stronger

sensitivity to shift location than PIC-MM, with false positive rates increasing more rapidly

toward the tips.

In contrast, under univariate shifts affecting only X_1 (Fig. 3b), PIC-MM and PIC-OGC

consistently maintained false negative rates of zero across all branch levels, indicating stable

power regardless of shift location. Interestingly, the remaining four methods showed

pronounced sensitivity to shift position under univariate shifts, with false negative rates

varying substantially across branch levels.

Figure 3. Sensitivity of method performance to the phylogenetic location of evolutionary shifts. Performance of

six methods evaluated under shifts introduced at different depths of a 128-species fixed-balanced tree. A

representative shift magnitude of 4^5 was applied systematically at each phylogenetic depth. Panel a shows false

positive rates under simultaneous shifts in both traits, whereas panel b shows false negative rates under

univariate shifts affecting only X_1. Additional details of the simulation design and performance metrics are

provided in the Materials and Methods. The data used to generate this figure are available in Supplementary

Table S3.

Performance of the Methods Under Non-BM Evolutionary Conditions

While the application of OLS regression or Pearson correlation on PIC values requires the

assumption that traits evolve under a BM model—so that PICs are statistically i.i.d.—the

computation of PIC values themselves does not strictly rely on BM. Instead, PIC values are

defined as algebraic contrasts on a phylogeny, and remain meaningful under a wide range of

evolutionary models (Garland et al. 1992). It is the statistical interpretation (e.g., *p*-values) of

PIC-OLS or PIC-Pearson that breaks down when the BM assumption is violated. In contrast,

methods such as PIC-MM and PIC-OGC—though still operating on PIC values—employ

robust estimators or nonparametric correlations that do not require the strict assumptions of

BM. Thus, they are retained in our comparison alongside PGLS, MR-PMM, and Corphylo

when evaluating method performance under simulated non-BM conditions. Importantly, this

also means that their performance can be fairly compared to methods that are explicitly

designed to accommodate a range of evolutionary models.

Although all simulations incorporated Brownian-motion–structured components, the

realized tip-level trait patterns may deviate from ideal BM expectations due to evolutionary

shifts, added stochastic noise, or their interaction. As a result, these data can sometimes be

better approximated by alternative covariance structures (e.g., OU, λ, or EB) when fitted

using PGLS. To capture this distinction, we used AIC-based model comparison as a

diagnostic criterion to classify simulated datasets as either BM-consistent or non-BM-

consistent at the tip level under standard PGLS covariance models.

Overall, performance patterns under non-BM conditions (Supplementary Fig. S2) closely

resembled those observed across the full set of simulations (Fig. 2). With the exception of

specific edge cases in which non-BM datasets are not defined (Supplementary Fig. S2a, c, d)

and direct comparisons are therefore not possible, relative method behavior remained largely

unchanged.

Across scenarios involving evolutionary shifts, including both bivariate shifts and

randomly located univariate shifts, PIC-OGC and PIC-MM continued to show strong and

stable performance under non-BM conditions, maintaining low error rates across a wide

range of shift magnitudes (Supplementary Figs. S2a–d). As observed across the full set of

simulations, under randomly located single-trait shifts at large magnitudes, PIC-OGC

exhibited noticeably higher error rates than PIC-MM. Importantly, this deviation was limited

in scope and did not alter the overall rank pattern of relative method performance.

In contrast, under gradual evolutionary scenarios without abrupt shifts, differences

among methods under non-BM conditions were modest, and PIC-OGC and PIC-MM were

not clearly distinguishable from the other approaches (Supplementary Figs. S2e–g). As

observed across the full set of simulations, no method consistently emerged as either superior

or inferior in the absence of shifts, indicating that background evolutionary heterogeneity

alone is insufficient to drive substantial performance divergence.

PIC-O(D)GC: Robust Performance across Tree Structure and Sample Size

To assess the robustness of the results on PIC-O(D)GC performance across evolutionary

shifts and no-shift regimes, we repeated the analyses shown in Figure 2 on additional

phylogenetic settings, including fixed-balanced trees with 16 and 256 species and randomly

generated trees with 128 species. Across these additional tree topologies and sample sizes,

PIC-O(D)GC and PIC-MM consistently ranked among the top-performing methods under

evolutionary-shift scenarios (Supplementary Figs. S3–S5). Although the magnitude of

performance differences varied with shift gradient, tree structure, and sample size—the slight

advantage of PIC-O(D)GC over PIC-MM observed under the 128-species fixed-balanced tree

(Fig. 2a) was no longer evident under other tree configurations—the overall relative ranking

of methods remained unchanged.

Secondary Benchmark Confirms Robustness of PIC-Based Methods

All results reported above are based on the β-based benchmark, which evaluates method

performance relative to the generative coupling between traits. To provide a complementary

perspective, we additionally evaluated all methods using a second, data-pattern–based

benchmark. This alternative framework exploits the availability of ancestral states in

simulations, enabling the computation of statistically independent branchwise changes

($\Delta X_1/L$ and $\Delta X_2/L$) that can be evaluated using standard statistical methods (e.g., Pearson and

Spearman correlations), rather than phylogenetic comparative methods.

When evaluated under the data-pattern–based benchmark, differences relative to the β-

based benchmark are minimal across most simulation settings. In particular, under

evolutionary-shift scenarios and across all three gradual-evolution scenarios (BM–BM, BM–

Norm, and Norm–BM) at low noise levels, the two benchmarks yield largely mirrored

conclusions: methods show low error rates when evaluated against the β-based benchmark

and correspondingly high accuracy under the data-pattern–based benchmark. As noise

variance increases under the tree gradual-evolution scenarios, however, the two benchmarks

begin to diverge. False-negative rates under the β-based benchmark rise sharply, whereas

accuracy under the data-pattern–based benchmark declines more gradually and remains

substantially higher (Fig. 4).

This divergence reflects differences in benchmark definition rather than changes in

method behavior. Under extreme noise, structural coupling defined by β may persist in the

generative model, while realized data provide little statistical support for association (see

Supplementary Methods S1 for details). Methods that correctly infer absent support therefore

remain accurate under the data-pattern–based benchmark even when $\beta = 1$.

Figure 4 provides a schematic illustration of this benchmark divergence using a

representative method (PGLS), while detailed method-specific results under the data-pattern–

based benchmark are reported in Supplementary Figs. S6–S12. Despite these benchmark-

dependent differences, overall performance patterns under the data-pattern–based benchmark

closely mirror those obtained under the β-based benchmark, with relative method rankings

remaining largely unchanged and PIC-O(D)GC and PIC-MM continuing to rank among the

top-performing methods under evolutionary-shift scenarios.

Figure 4. Illustrative comparison of benchmark outcomes for a representative method (PGLS) under a gradual-evolution simulation scenario (BM–BM), simulated on a fixed-balanced tree with 128 species. The figure contrasts evaluations under the *β*-based and data-pattern–based benchmarks and highlights their differing perspectives under noise-dominated conditions. Additional details of the simulation design and the two benchmarks are provided in the Materials and Methods. Detailed method-specific results under the data-pattern–based benchmark are reported in Supplementary Figs. S6–S12. The data used to generate this figure are available in Supplementary Table S4.

Application of PIC-O(D)GC to Selected Empirical Datasets with Trait Shifts

Simulation results, validated against internal benchmarks from the generative model or recorded ancestral states, showed PIC-O(D)GC and PIC-MM to be most robust under abrupt shifts. To evaluate PIC-O(D)GC's performance on real-world data, we analyzed three empirical trait datasets, derived either in full or in part from published phylogenetic resources or comparative studies (Jones et al. 2009, Scales et al. 2009, Makino and Kawata 2019).

Invertebrate propagule size vs. genome size.—Across 44 invertebrate species, the PIC distribution revealed seven extreme contrasts (Fig. 6a). Notably, $\log_{10}$-transformation of the original trait values did not eliminate these deviations, indicating that standard variance-stabilizing transformations are insufficient to remove lineage-specific extremes after phylogenetic correction. These contrasts likely reflect heterogeneous evolutionary shifts occurring along a small number of branches and illustrate how a few extreme observations can exert disproportionate influence on correlation-based comparative analyses (Fig. 5b). In this dataset, methods that explicitly account for extreme contrasts—such as PIC-OGC and

737 PIC-MM—returned non-significant associations, whereas other methods inferred significant

738 correlations. This discrepancy highlights the sensitivity of comparative correlation results to

739 the treatment of extreme contrasts and underscores the importance of explicitly addressing

740 such deviations in phylogenetically informed analyses.

741
Figure 5. Relationship between propagule size and genome size across 44 invertebrates. a) Phylogenetically

independent contrasts (PICs) between $\log_{10}$-transformed propagule size and genome size. Dashed lines indicate

the outlier boundaries defined by the boxplot rule, $Q1 - 1.5 \times IQR$ and $Q3 + 1.5 \times IQR$ (Tukey 1977). Contrasts

exceeding these thresholds in either trait are marked as filled triangles and treated as outliers, while remaining

contrasts are shown as points. b) Statistical significance of the association between genome size and propagule

size as assessed by six comparative methods. Detailed correlation coefficients and p-values, along with the

original trait values and the PIC values from which the scatterplots were derived, are provided in Supplementary

Table S5.

Lizard fast glycolytic vs. slow oxidative fiber proportion.—Across 22 lizard species, the PIC

distribution revealed a single univariate outlier in FG fiber proportion (Fig. 6a),

corresponding to a contrast between *Phrynosoma* and its sister clade (Fig. 6b). Inspection of

the raw data revealed that this contrast is driven almost entirely by a marked reduction in FG

fiber proportion across the four sampled *Phrynosoma* species, indicating a directional

downsizing shift within this lineage. Across the full dataset, phylogenetic comparative

methods produced mixed results (Fig. 6c). While PIC-ODGC and PIC-MM detected a

statistically significant association between SO and FG fiber proportions, other methods such

as PIC-Pearson, PGLS, and MR-PMM failed to do so.

To test the influence of the downsizing shift on methods' performance, we re-ran all

analyses after removing the four *Phrynosoma* species. Following this exclusion, all methods

except MR-PMM returned p-values below 0.05, consistently indicating a negative association

between SO and FG fiber proportions (Supplementary Table S6). This result suggests that the

localized evolutionary shift within *Phrynosoma* both introduces statistical heterogeneity and

suppresses correlation signals under certain comparative frameworks.

Figure 6. Relationship between slow oxidative (SO) and fast glycolytic (FG) muscle fiber proportions in 22

lizard species. a) Phylogenetically independent contrasts (PICs) between the proportions of SO and FG muscle

fibers. Dashed lines indicate the outlier boundaries defined by the boxplot rule, $Q1 - 1.5 \times IQR$ and $Q3 + 1.5 \times$

IQR (Tukey 1977). A contrast exceeding these thresholds in FG fiber proportion is marked as filled triangles

and treated as outliers, while remaining contrasts are shown as points. b) Phylogenetic tree highlighting the

branch associated with the outlier contrast. The arrow marks the lineage corresponding to a pronounced

reduction in FG fiber proportion. c) The *p*-values from multiple phylogenetic comparative methods testing for

association between SO and FG fiber proportions. Detailed correlation coefficients and *p*-values, along with the

original trait values and PIC values from which the scatterplots were derived, are provided in Supplementary

Table S6.

Body mass and population density in even-toed ungulates.—The PIC distribution across 109

species exhibits seven univariate outliers with varying degrees of deviation (Fig. 7a),

reflecting extreme evolutionary contrasts along specific branches. Despite all methods

indicating a negative association between the two traits, their statistical support differs

markedly (Fig. 7b). In particular, two methods sensitive to extreme values, PIC-Pearson and

Corphylo, yield non-significant results, whereas approaches that explicitly account for

outliers, including PIC-OGC and PIC-MM, detect a significant negative association.

Figure 7. Relationship between body mass and population density in even-toed ungulates (Artiodactyla). a)

Phylogenetically independent contrasts (PICs) between $\log_{10}$-transformed body mass and population density for

109 artiodactyl species. Dashed lines indicate the outlier boundaries defined by the boxplot rule, $Q1 - 1.5 \times$

IQR and $Q3 + 1.5 \times IQR$ (Tukey 1977). Contrasts exceeding these thresholds in either trait are marked as filled

triangles and treated as outliers, while remaining contrasts are shown as points. b) p-values from multiple

comparative methods testing for an association between body mass and population density. Detailed correlation

coefficients and p-values, along with the original trait values and PIC values from which the scatterplots were

derived, are provided in Supplementary Table S7.

Because the direction of evolutionary change could not be confidently inferred from the

phylogenetic positions of the most extreme contrast pairs, we adopted a conservative

sensitivity analysis focusing on the two most pronounced univariate deviations in the PIC

distribution (Fig. 7a). These two contrasts represent the most extreme deviations and are

therefore the most likely to distort inference under methods that assume identically

distributed contrasts; weaker departures were retained because their influence on overall

results is expected to be negligible. Specifically, we removed the four terminal taxa involved

in these two extreme contrasts—one associated with body mass (between *Syncerus caffer* and

Ammodorcas clarkei) and the other associated with population density (between *Kobus leche*

and *Kobus megaceros*). Given the large sample size (109 species), excluding these four taxa

had a negligible impact on overall statistical power. Reanalysis after this exclusion yielded

consistent results across all methods, with negative effect estimates and p-values below 0.01

in every case (Supplementary Table S7). These results indicate that a small number of highly

extreme contrasts can substantially attenuate statistical support for otherwise robust trait

associations.

811   ***Runtime Comparison of Phylogenetic Correlation Methods***

To assess computational efficiency, we compared the runtime of PIC-O(D)GC with the other

five phylogenetic comparative methods across six datasets (Fig. 8). Among all methods, PIC-

O(D)GC and PIC-MM showed the smallest increase in runtime as sample size grew,

highlighting their strong scalability. The overall computational cost increased from PIC-

Pearson to PIC-O(D)GC, PIC-MM, and PGLS, with MR-PMM and Corphylo being the two

slowest methods.

Figure 8. Computational time comparison across phylogenetic correlation methods. Average runtime

(in seconds, log10-transformed) was measured for each method across six datasets, including three

empirical datasets used in the above analyses (22, 44, and 109 species) and three simulated datasets

based on fixed-balanced phylogenetic trees (16, 128, and 256 species). All simulated datasets were

generated under the dual-trait shift scenario with an effect size of 4^{-3}. Each method was repeated 100

times per dataset to obtain mean values. All runtime measurements were performed on a Dell server

running CentOS Linux 7.9, equipped with an AMD EPYC 7702 64-core processor and 1 TB of RAM.

Computational time did not always increase with sample size. For instance, both PGLS and Corphylo

ran slower on the 109-species dataset than on the 128-species one. This was likely due to relatively

long tip labels (e.g., Latin species names) in the 109-species phylogeny, which increased the time

required for tree–trait matching. Due to noticeable fluctuations in system load on the shared

computational environment, absolute runtimes varied considerably across replicates. However, the

relative performance ranking of methods was stable and therefore provides a reliable basis for

comparison. The raw timing data used to generate this figure are provided in Supplementary Table

S8.

PIC-Pearson was the fastest methods, benefiting from linear computational complexity.

PIC-O(D)GC also exhibited low computational cost, relying on median-based outlier

detection (O(n)) and, depending on outlier presence, applying either Pearson or Spearman

correlation. When Spearman correlation is used, an O(n log n) cost is incurred due to ranking,

which remains efficient even for large trees. In small datasets, a normality test (e.g., Shapiro–

Wilk) is additionally applied to guide correlation choice, introducing modest overhead that is

most apparent at small sample sizes. PIC-MM required more computation than PIC-O(D)GC,

consistent with its robust two-stage estimation procedure: an initial S-estimation step (up to

$O(n^3)$), followed by iterative M-estimation steps (each $O(n^2)$). PGLS was even slower than

PIC-MM, driven by repeated matrix inversions. The remaining model-based approaches—

MR-PMM, and Corphylo—incurred the highest computational time overall. For example, on

the largest dataset ($n = 256$), PIC-Pearson, PIC-O(D)GC, and PIC-MM completed in 3×10^{-4},

10^{-3}, and 2×10^{-3} seconds, respectively, while PGLS and MR-PMM required 0.12 and 6.5

seconds, respectively. Corphylo was particularly slow, taking 26 seconds on the same dataset.

These results underscore the scalability of PIC-O(D)GC, which achieves high

computational efficiency without compromising robustness. This dual strength makes it

particularly suitable for large-scale phylogenetic comparative analyses.

853   **DISCUSSION**

854   ***Equivalence of Pearson Correlation and OLS in Context***

Both Pearson correlation and OLS regression can be applied to PIC to assess trait

associations in a phylogenetically aware manner (Felsenstein 1985, Felsenstein 1988, Murray

et al. 2020, Horvath et al. 2022, Bai et al. 2023, Wang et al. 2023). When applied to PIC data

under linear assumptions, these two approaches yield mathematically equivalent results in

terms of statistical significance and correlation direction. In this study, we chose to use

Pearson correlation for its computational simplicity and consistency with other nonparametric

methods, such as Spearman correlation, which is a key component of the PIC-O(D)GC

workflow. The equivalence between Pearson correlation and OLS regression ensures logical

and statistical consistency with previous findings while reinforcing the relevance and

robustness of our results within the context of phylogenetic comparative analyses.

866   ***Comparing Statistical Methods for Detecting Trait Correlations***

Our study evaluated the performance of various statistical methods in detecting trait

correlations under diverse evolutionary scenarios, including with and without abrupt shifts.

PIC-O(D)GC and PIC-MM consistently demonstrated robust performance, particularly

excelling under shift scenarios by minimizing false positives as well as false negatives,

thereby maintaining high overall accuracy—consistent with prior evaluations of robust

regression strategies in the presence of model violations and trait shifts (Adams et al. 2024).

In contrast, traditional methods like PIC-Pearson and PGLS, as well as robust variants such

as PGLS-MM, struggled under high shift gradients, highlighting their limitations in

challenging conditions. In no-shift scenarios, both PIC-MM and PIC-O(D)GC performed

comparably to other reliable methods, such as PIC-Pearson, PGLS, and Corphylo,

underscoring that neither method was uniformly superior across all conditions. These

complementary strengths highlight the importance of aligning method choice with expected

data characteristics and analytical goals.

Notably, the abrupt trait shifts introduced in our simulations can be considered an

extreme form of evolutionary heterogeneity—a concept emphasized by Mazel et al. (2016) as

a key source of model violation in phylogenetic regression. While Mazel et al. focused on

rate heterogeneity, our results show that discrete directional shifts can also distort inference

and compromise methods assuming homogeneous evolutionary processes. This underscores

the importance of methods capable of accommodating both continuous and discontinuous

heterogeneity.

888   *Conceptual Simplicity and Practical Accessibility*

In addition to the substantial runtime advantage demonstrated in Figure 8, PIC-O(D)GC also

offers strengths in conceptual simplicity and user accessibility. By dynamically adapting to

the presence or absence of outliers and data normality, it strikes an effective balance between

robustness and interpretability. Its reliance on straightforward statistical principles, such as

Spearman's rank correlation, ensures that the method is logically intuitive and easy to

understand, even for researchers with limited statistical training (Redmond and Keenan

2002). By avoiding rigid assumptions about data distribution, PIC-O(D)GC provides

transparent and interpretable results, reducing the risk of misinterpretation and enhancing its

accessibility to a broader range of biologists.

899   *From Statistical Artifacts to Evolutionary Signals: Interpreting PIC Outliers*

Beyond its utility in assessing trait correlations, the outlier detection step in PIC-O(D)GC

holds promise for identifying evolutionary shifts or jumps. Recent studies have highlighted

the importance of models that accommodate abrupt changes in trait evolution, such as the

singular events model (Uyeda et al. 2018). These rare but significant transitions may leave

identifiable signatures in PIC data (Uyeda et al. 2018), including extreme contrast values, or

pronounced localized changes along specific branches.

The importance of examining PIC scatterplots cannot be overstated. Even when an

overall correlation appears statistically significant, the result may be disproportionately

driven by a few extreme contrast values. Visual inspection of the PIC plot is therefore

essential, not only as a diagnostic check, but also as a guide to more structured interpretation.

When such outliers are present, we recommend analyzing the relationship between two

traits from three complementary perspectives: 1) Overall correlation can be assessed using

robust methods such as PIC-MM or PIC-O(D)GC, which account for outliers and non-

normality while maintaining statistical stability across heterogeneous evolutionary scenarios;

2) Shift-to-shift relationships should be interpreted separately, considering whether shifts are

aligned or opposed in direction, which may reveal patterns of evolutionary coupling or

divergence. 3) Background correlation refers to the relationship between traits during non-

shift, gradual phases of evolution.

When a shift is located deep in the phylogeny—particularly near the root, as seen in

Figure 1 (panels d, f, j, l)—an effective strategy is to partition the data into two clades at the

shift node. This allows separate estimation of trait relationships within each group, providing

insight into background evolutionary dynamics before and after the divergence introduced by

the shift.

For shifts occurring on terminal or near-terminal branches (Fig. 3), we observed that

PIC-O(D)GC, and even PIC-MM, may exhibit reduced sensitivity, reflecting the

disproportionate influence of a small number of localized, large-effect contrasts. In these

cases, removing the taxa involved in the detected shift can be an effective and often

necessary step to limit their impact and recover the underlying background correlation

structure.

Once the influence of shifts is appropriately filtered—whether through clade partitioning

or tip removal—regression-based methods such as PIC-OLS, or PGLS (Felsenstein 1985,

Martins and Hansen 1997, Pagel 1997, Pagel 1999, Ives et al. 2007, Revell 2010) become

better suited for background-phase analysis by leveraging continuous trait variation and

enabling effect-size estimation and model-based inference. In practice, the empirical

examples of body mass and population density in even-toed ungulates and SO and FG muscle

fiber proportions in lizard species illustrate how such regression-based approaches can

provide clear directionality and statistically supported assessments of trait associations once

shift effects are appropriately filtered. These regression-based approaches are preferred in

background-phase analysis due to their interpretability and modeling efficiency. In contrast,

robust contrast-based methods like PIC-O(D)GC or PIC-MM (Adams et al. 2024) may be

overly conservative in this refined context.

Empirical evidence increasingly supports the idea that phenotypic evolution involves

rare but consequential jumps embedded within longer periods of gradual change or stasis.

Such discontinuities have been observed across a range of systems, including body size

evolution in crocodylomorphs (Godoy et al. 2019), bursts of morphological innovation in

lizards (Brennan et al. 2024), shifts in venom gene evolution linked to ecological opportunity

in snakes (Barua and Mikheyev 2020), and abrupt changes in genomic traits such as GC

content and genome size in bacteria (Gao and Wu 2022, Hu et al. 2022, Mahajan and Agashe

2022). These cases underscore the importance of not only identifying where and when such

evolutionary shifts occur, but also understanding how they alter patterns of trait covariation.

In particular, they highlight the need to distinguish among overall correlations across the tree,

background correlations within stable lineages, and shift-to-shift relationships that may

reflect directional evolutionary coupling or divergence between regimes.

The combined use of contrast-based diagnostics such as PIC-O(D)GC and targeted

analyses of trait relationships offers a tractable workflow for detecting and interpreting

evolutionary pulses. Unlike fully model-based methods, PIC analysis leverages branch-level

resolution to identify candidate shifts without prior assumptions about evolutionary regimes.

This approach complements theoretical and algorithmic models such as OU-with-jumps

(Landis et al. 2013), Lévy process-based jump models (Duchen et al. 2017, Landis and

Schraiber 2017), LASSO-regularized shift detection (Khabbazian et al. 2016, Bastide et al.

2018, Zhang et al. 2024), and singular events models (Uyeda et al. 2011, Uyeda et al. 2018),

all of which recognize discrete shifts as critical components of macroevolutionary dynamics.

963   ***Limitations and Future Directions***

While correlation-based methods such as PIC-O(D)GC are robust to non-normal data

and outliers, they are inherently limited to detecting monotonic associations and do not

provide parameter estimates or accommodate multiple predictors. Thus, they should be

viewed as complementary to parametric models like PGLS, which remain essential for

estimating the strength and functional form of evolutionary relationships.

Further refinement of this workflow may be achieved by incorporating phylogenetic

features such as branch lengths and trait-specific evolutionary rates. These additions could

improve the specificity of shift detection and help distinguish genuine evolutionary

transitions from background variation. Integrating PIC outlier diagnostics into hypothesis-

driven macroevolutionary studies may thus provide a more nuanced view of how trait

relationships evolve across lineages.

FUNDING

This study was supported in part by the National Natural Science Foundation of China (grant

number 31671321), with leftover resources used for occasional research expenses.

ACKNOWLEDGMENTS

We sincerely thank Professor Josef Uyeda (Associate Editor) and the anonymous reviewer

for their constructive and insightful comments. Many sentences in this article were drafted

with the assistance of ChatGPT-4 and its later versions, large language models developed by

OpenAI. The authors thoroughly reviewed and revised each sentence and took full

responsibility for the language and content of the entire article.

DATA AVAILABILITY

Supplementary Methods and supplementary analyses are provided in

Supplementary_Methods_S1.pdf and Supplementary_Results_S1.pdf, respectively.

Supplementary Figures S1–S12 are available in Supplementary_Figures.pdf. All data

supporting the figures presented in the main text and supplementary materials, including

Tables S1–S28, are provided in Supplementary_Tables.xlsx. All source code used for

simulations and analyses is available in Code.zip. We also have implemented a suite of

functions for conducting PIC-O(D)GC in the R package PICODGC (Phylogenetic

Independent Contrasts with Outlier- and Distribution-Guided Correlation), freely available at

https://github.com/zhenglinchen/PICODGC.

REFERENCES

Adams R., Cain Z., Assis R., DeGiorgio M. 2024. Robust phylogenetic regression. Syst.

Biol., 73:140-157.

Bai X.-L., Yang D., Sher J., Zhang Y.-B., Zhang K.-Y., Liu Q., Wen H.-D., Zhang J.-L., Slot

M. 2023. Divergences in stem and leaf traits between lianas and coexisting trees in a

subtropical montane forest. Journal of Plant Ecology, 17:rtad037.

Barua A., Mikheyev A.S. 2020. Toxin expression in snake venom evolves rapidly with

constant shifts in evolutionary rates. Proc. R. Soc. B, 287:20200613.

Bastide P., Ané C., Robin S., Mariadassou M. 2018. Inference of adaptive shifts for

multivariate correlated traits. Syst. Biol., 67:662-680.

Blomberg S.P., Lefevre J.G., Wells J.A., Waterhouse M. 2012. Independent contrasts and

PGLS regression estimators are equivalent. Syst. Biol., 61:382-391.

Brennan I.G., Chapple D.G., Keogh J.S., Donnellan S. 2024. Evolutionary bursts drive

morphological novelty in the world's largest skinks. Curr. Biol., 34:3905-3916.e3905.

Cornwallis C.K., Griffin A.S. 2024. A guided tour of phylogenetic comparative methods for

studying trait evolution. Annu. Rev. Ecol. Evol. Syst., 55:181-204.

Dewar A.E., Belcher L.J., West S.A. 2025. A phylogenetic approach to comparative

genomics. Nat. Rev. Genet., 26:395-405.

Duchen P., Leuenberger C., Szilagyi S.M., Harmon L., Eastman J., Schweizer M., Wegmann

D. 2017. Inference of evolutionary jumps in large phylogenies using Levy processes. Syst.

Biol., 66:950-963.

Felsenstein J. 1985. Phylogenies and the comparative method. Amer. Natur., 125:1-15.

Felsenstein J. 1988. Phylogenies and quantitative characters. Annu. Rev. Ecol. Syst., 19:445-

471.

Freckleton R.P., Harvey P.H., Pagel M. 2002. Phylogenetic analysis and comparative data: A

test and review of evidence. Amer. Natur., 160:712-726.

Gao Y., Wu M. 2022. Microbial genomic trait evolution is dominated by frequent and rare

pulsed evolution. Sci. Adv., 8:eabn1916.

Garamszegi L.Z. 2014. Modern Phylogenetic Comparative Methods and Their Application in

Evolutionary Biology: Concepts and Practice. Berlin, Springer.

Garland T., Jr, Harvey P.H., Ives A.R. 1992. Procedures for the analysis of comparative data

using phylogenetically independent contrasts. Syst. Biol., 41:18-32.

Godoy P.L., Benson R.B.J., Bronzati M., Butler R.J. 2019. The multi-peak adaptive

landscape of crocodylomorph body size evolution. BMC Evol. Biol., 19:167.

Grafen A. 1989. The phylogenetic regression. Philos. Trans. R. Soc. Lond. B Biol. Sci.,

326:119-157.

Hadfield J.D. 2010. MCMC methods for multi-response generalized linear mixed models: the

MCMCglmm R package. J. Stat. Softw., 33:1-22.

Hadfield J.D., Nakagawa S. 2010. General quantitative genetic methods for comparative

biology: phylogenies, taxonomies and multi-trait models for continuous and categorical

characters. J. Evol. Biol., 23:494-508.

Halliwell B., Holland B.R., Yates L.A. 2025. Multi-response phylogenetic mixed models:

concepts and application. Biol. Rev., 100:1294-1316.

Hansen T.F. 1997. Stabilizing selection and the comparative analysis of adaptation.

Evolution, 51:1341-1351.

Harmon L.J., Losos J.B., Jonathan Davies T., Gillespie R.G., Gittleman J.L., Bryan Jennings

W., Kozak K.H., McPeek M.A., Moreno-Roark F., Near T.J., Purvis A., Ricklefs R.E.,

Schluter D., Schulte Ii J.A., Seehausen O., Sidlauskas B.L., Torres-Carvajal O., Weir J.T.,

Mooers A.Ø. 2010. Early bursts of body size and shape evolution are rare in comparative

data. Evolution, 64:2385-2396.

Ho L.S.T., Ane C. 2014. A linear-time algorithm for Gaussian and non-Gaussian trait

evolution models. Syst. Biol., 63:397-408.

Hoaglin D.C., Iglewicz B., Tukey J.W. 1986. Performance of Some Resistant Rules for

Outlier Labeling. J Am Stat Assoc, 81:991-999.

Horvath S., Lu A.T., Haghani A., Zoller J.A., Li C.Z., Lim A.R., Brooke R.T., Raj K.,

Serres-Armero A., Dreger D.L., Hogan A.N., Plassais J., Ostrander E.A. 2022. DNA

methylation clocks for dogs and humans. Proc. Natl. Acad. Sci. USA, 119:e2120887119.

Hu E.-Z., Lan X.-R., Liu Z.-L., Gao J., Niu D.-K. 2022. A positive correlation between GC

content and growth temperature in prokaryotes. BMC Genomics, 23:110.

Ives A.R. 2022. Random errors are neither: On the interpretation of correlated data. Methods

Ecol. Evol., 13:2092-2105.

Ives A.R., Helmus M.R. 2011. Generalized linear mixed models for phylogenetic analyses of

community structure. Ecol. Monographs, 81:511-525.

Ives A.R., Midford P.E., Garland T., Jr. 2007. Within-species variation and measurement

error in phylogenetic comparative methods. Syst. Biol., 56:252-270.

Jones K.E., Bielby J., Cardillo M., Fritz S.A., O'Dell J., Orme C.D.L., Safi K., Sechrest W.,

Boakes E.H., Carbone C., Connolly C., Cutts M.J., Foster J.K., Grenyer R., Habib M., Plaster

C.A., Price S.A., Rigby E.A., Rist J., Teacher A., Bininda-Emonds O.R.P., Gittleman J.L.,

Mace G.M., Purvis A. 2009. PanTHERIA: a species-level database of life history, ecology,

and geography of extant and recently extinct mammals. Ecology, 90:2648-2648.

Khabbazian M., Kriebel R., Rohe K., Ané C. 2016. Fast and accurate detection of

evolutionary shifts in Ornstein–Uhlenbeck models. Methods Ecol. Evol., 7:811-824.

Landis M.J., Schraiber J.G. 2017. Pulsed evolution shaped modern vertebrate body sizes.

Proc. Natl. Acad. Sci. USA, 114:13224-13229.

Landis M.J., Schraiber J.G., Liang M. 2013. Phylogenetic analysis using Levy processes:

finding jumps in the evolution of continuous traits. Syst. Biol., 62:193-204.

Lynch M. 1991. Methods for the analysis of comparative data in evolutionary biology.

Evolution, 45:1065-1080.

Mahajan S., Agashe D. 2022. Evolutionary jumps in bacterial GC content. G3

Genes|Genomes|Genetics, 12:jkac108.

Makino T., Kawata M. 2019. Invasive invertebrates associated with highly duplicated gene

content. Mol. Ecol., 28:1652-1663.

Martins E.P., Hansen T.F. 1997. Phylogenies and the comparative method: A general

approach to incorporating phylogenetic information into the analysis of interspecific data.

Amer. Natur., 149:646-667.

Mazel F., Davies T.J., Georges D., Lavergne S., Thuiller W., Peres-Neto P.R. 2016.

Improving phylogenetic regression under complex evolutionary models. Ecology, 97:286-

293.

Murray G.G.R., Charlesworth J., Miller E.L., Casey M.J., Lloyd C.T., Gottschalk M., Tucker

A.W., Welch J.J., Weinert L.A. 2020. Genome reduction is associated with bacterial

pathogenicity across different scales of temporal and ecological divergence. Mol. Biol. Evol.,

38:1570-1579.

O'Meara B. 2016. Phylogenetic Comparative Method. In: Kliman RM editor. Encyclopedia

of Evolutionary Biology. Oxford, Academic Press, p. 254-256.

Pagel M. 1997. Inferring evolutionary processes from phylogenies. Zool. Scr., 26:331-348.

Pagel M. 1999. Inferring the historical patterns of biological evolution. Nature, 401:877-884.

Paradis E., Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics, 35:526-528.

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Redmond A.C., Keenan A.M. 2002. Understanding statistics. Putting p-values into

perspective. J Am Podiatr Med Assoc, 92:297-305.

Revell L.J. 2010. Phylogenetic signal and linear regression on species data. Methods Ecol.

Evol., 1:319-329.

Revell L.J. 2024. phytools 2.0: an updated R ecosystem for phylogenetic comparative

methods (and other things). PeerJ, 12:e16505.

Rohlf F.J. 2001. Comparative methods for the analysis of continuous variables: geometric

interpretations. Evolution, 55:2143-2160.

Romiguier J., Gayral P., Ballenghien M., Bernard A., Cahais V., Chenuil A., Chiari Y.,

Dernat R., Duret L., Faivre N., Loire E., Lourenco J.M., Nabholz B., Roux C., Tsagkogeorga

G., Weber A.A.T., Weinert L.A., Belkhir K., Bierne N., Glémin S., Galtier N. 2014.

Comparative population genomics in animals uncovers the determinants of genetic diversity.

Nature, 515:261-263.

Royston P. 1992. Approximating the Shapiro-Wilk W-test for non-normality. Statistics and

Computing, 2:117-119.

Scales J.A., King A.A., Butler M.A. 2009. Running for your life or running for your dinner:

what drives fiber-type evolution in lizard locomotor muscles? Amer. Natur., 173:543-553.

Slater G.J., Pennell M.W. 2014. Robust regression and posterior predictive simulation

increase power to detect early bursts of trait evolution. Syst. Biol., 63:293-308.

Sorensen D., Gianola D. 2002. Likelihood, Bayesian, and MCMC Methods in Quantitative

Genetics. New York, Springer.

Sumner S., Favreau E., Geist K., Toth A.L., Rehan S.M. 2023. Molecular patterns and

processes in evolving sociality: lessons from insects. Philos Trans R Soc Lond B Biol Sci,

378:20220076.

Symonds M.R.E., Blomberg S.P. 2014. A primer on phylogenetic generalised least squares.

In: Garamszegi LZ editor. Modern Phylogenetic Comparative Methods and Their

Application in Evolutionary Biology: Concepts and Practice. Berlin, Heidelberg, Springer

Berlin Heidelberg, p. 105-130.

Tukey J.W. 1977. Exploratory Data Analysis. Reading, MA, Addison-Wesley.

Upham N.S., Esselstyn J.A., Jetz W. 2019. Inferring the mammal tree: Species-level sets of

phylogenies for questions in ecology, evolution, and conservation. PLoS Biol., 17:e3000494.

Uyeda J.C., Hansen T.F., Arnold S.J., Pienaar J. 2011. The million-year wait for

macroevolutionary bursts. Proc. Natl. Acad. Sci. USA, 108:15908-15913.

Uyeda J.C., Zenil-Ferguson R., Pennell M.W. 2018. Rethinking phylogenetic comparative

methods. Syst. Biol., 67:1091-1109.

Wang B., Tong Z.-Y., Xiong Y.-Z., Wang X.-F., Scott Armbruster W., Huang S.-Q. 2023.

The evolution of flower–pollinator trait matching, and why do some alpine gingers appear to

be mismatched? Ann. Bot., 132:1073-1088.

Westoby M., Yates L., Holland B., Halliwell B. 2023. Phylogenetically conservative trait

correlation: Quantification and interpretation. J Ecol, 111:2105-2117.

Zhang W., Kenney T., Ho L.S.T. 2024. Evolutionary shift detection with ensemble variable

selection. BMC Ecology and Evolution, 24:11.

Zheng L., Ives A.R., Garland T., Larget B.R., Yu Y., Cao K. 2009. New multivariate tests for

phylogenetic signal and trait correlations applied to ecophysiological phenotypes of nine

Manglietia species. Funct. Ecol., 23:1059-1069.

1143