Enhancing Phylogenetic Independent Contrasts: Addressing Abrupt Evolutionary Shifts with Outlier- and Distribution-Guided Correlation

Zheng-Lin Chen, Rui Huang, Hong-Ji Guo and Deng-Ke Niu[*]

MOE Key Laboratory for Biodiversity Science and Ecological Engineering, College of Life Sciences, Beijing Normal University, Beijing, 100875, China

*Correspondence to be sent to: *Deng-Ke Niu, College of Life Sciences, Beijing Normal University, Beijing, 100875, China*; Emails: dkniu@bnu.edu.cn, dengkeniu@hotmail.com

ABSTRACT

Phylogenetic comparative methods are indispensable for analyzing cross-species data while

accounting for evolutionary relationships. Traditional methods like phylogenetically

independent contrasts (PIC) and phylogenetic generalized least squares (PGLS) are

well-suited for gradual evolution under Brownian motion (BM) or similar assumptions.

However, their effectiveness diminishes under abrupt evolutionary shifts or heterogeneous

evolutionary processes. We introduce a hybrid framework, outlier-guided correlation (OGC)

for large datasets and outlier- and distribution-guided correlation (ODGC) for small datasets,

collectively termed O(D)GC, which integrates Pearson and Spearman correlation analyses to

handle outliers and non-normal data dynamically. Using simulations across diverse

evolutionary scenarios, we compared PIC-O(D)GC with a comprehensive range of methods,

including eight robust regression approaches (PIC-MM, PIC-L1, PIC-S, PIC-M, and their

PGLS counterparts); PGLS optimized using five evolutionary models: BM, lambda,

Ornstein–Uhlenbeck random (OU-random), OU-fixed, and Early-burst; Corphylo (an

OU-based method); PIC-Pearson; and two advanced models, phylogenetic generalized linear

mixed models (PGLMM) and multi-response phylogenetic mixed models (MR-PMM). Our

results demonstrate that PIC-O(D)GC and PIC-MM consistently exhibit superior

performance under conditions with evolutionary shifts, maintaining low false positive rates

and high accuracy. In no-shift scenarios, PIC-O(D)GC and PIC-MM perform comparably to

other reliable methods, underscoring their adaptability and robustness across varied datasets.

Furthermore, PIC-O(D)GC offers computational efficiency and flexibility, making it

particularly suitable for large-scale datasets. These findings underscore the value of adaptive,

nonparametric frameworks like PIC-O(D)GC in complementing traditional phylogenetic

approaches. By bridging computational simplicity with methodological robustness, O(D)GC

provides a versatile tool for trait correlation analysis in both stable and dynamic evolutionary

contexts.

KEYWORDS

Phylogenetic comparative methods; Phylogenetically independent contrasts (PIC);

Outlier-guided correlation (OGC); Abrupt evolutionary shifts; Robust regression; Trait

correlation; phylogenetic autocorrelation; Adaptive frameworks.

INTRODUCTION

Phylogenetically closer species tend to exhibit more similar traits due to their shared ancestry,

resulting in phylogenetic autocorrelation among samples (Felsenstein 1985, Garamszegi 2014,

Cornwallis and Griffin 2024). Ignoring this autocorrelation and using conventional statistical

methods can lead to erroneous conclusions. To address this challenge, evolutionary biologists

have developed a range of phylogenetic comparative methods that explicitly account for these

dependencies (Grafen 1989, Lynch 1991, Garamszegi 2014, O'Meara 2016, Cornwallis and

Griffin 2024).

The earliest method was phylogenetically independent contrasts (PIC) (Felsenstein 1985).

The core principle behind PIC is that, while species at the tips of a phylogenetic tree are

interdependent due to shared ancestry, the evolutionary changes along each branch can be

treated as statistically independent events. In this context, PIC refers to the contrast values

calculated between species pairs along the phylogenetic tree, inferring evolutionary changes

along pairs of branches and standardizing these changes by branch length (or time) to generate

transformed data points. These PIC values, typically subjected to ordinary least squares (OLS)

regression, provide insight into the relationships between traits. This approach makes the

evolutionary data more suitable for conventional statistical methods by removing the

phylogenetic non-independence. Thus, PIC can refer both to the contrast values calculated

from the phylogeny and to the subsequent OLS regression analysis applied to these values.

PIC assumes that traits evolve according to a Brownian motion (BM) model, where trait

variation accumulates gradually and randomly over time. Strictly speaking, OLS regression of

PIC is a special case of the phylogenetic generalized least squares (PGLS) method under the

conditions of perfect BM (Rohlf 2001). PGLS extends this framework by incorporating a

variance-covariance matrix for the residuals in the generalized linear model to account for

phylogenetic autocorrelation among samples (Martins and Hansen 1997, Pagel 1997, Pagel

1999). This adjustment helps reduce false positives. Although some PGLS models, such as

Pagel's λ, Ornstein–Uhlenbeck (OU), and Early-burst (EB) models, can handle heterogeneous

evolutionary rates (Hansen 1997, Freckleton et al. 2002, Harmon et al. 2010, Ho and Ane

2014), they fundamentally assume gradual trait changes (Landis et al. 2013). This assumption

can be problematic when dealing with evolutionary jumps caused by singular events.

During biological evolution, certain branches may experience rare and dramatic trait

changes (Maddison and FitzJohn 2015). Increasingly, studies and data have shown that abrupt

evolutionary shifts, or evolutionary jumps, have a significant impact on biodiversity evolution

(Landis and Schraiber 2017, Gao and Wu 2022, Smith et al. 2023, Sumner et al. 2023). A

typical example of this is Felsenstein's worst-case scenario (Felsenstein 1985, Uyeda et al.

2018), where two traits undergo a significant jump near the root of the phylogenetic tree while

evolving independently on other branches. Even if this dramatic change occurred only once, it

could still lead to erroneous conclusions of significant correlation between the two traits using

ordinary linear regression on PIC values or PGLS models (Uyeda et al. 2018).

Recently, more studies have focused on identifying jump events and providing more

reasonable explanations for the processes underlying trait evolution. Uyeda et al. (2018)

underscore the need for models like the singular events model, which can accommodate the

complexities and unpredictabilities inherent in evolutionary biology. Some studies suggest

that jump events cause trait changes to exhibit a "fat-tails" distribution, using Levy processes

to handle this distribution (Landis et al. 2013, Duchen et al. 2017). Others propose that abrupt

shift events lead to outliers, using robust regression to reduce model sensitivity to violations

(Slater and Pennell 2014, Adams et al. 2024). Additionally, phylogenetic generalized linear

mixed models (PGLMM) and Multi-response phylogenetic mixed models (MR-PMM) offer

flexible frameworks to address the complexities of phylogenetic data, including diverse data

types and random effects (Ives and Helmus 2011, Westoby et al. 2023, Halliwell et al. 2024).

PGLMM models both fixed and random effects to account for phylogenetic autocorrelation,

while MR-PMM extends this capability to multiple traits, providing a comprehensive

approach to co-evolutionary analysis. Theoretically, they are capable of handling deviations

from BM.

Evolutionary shifts, especially those occurring in branches close to the root of the

phylogenetic tree, typically do not manifest directly as outliers in the trait data but instead

appear as outliers in the PIC values. As a result, robust statistical methods designed to handle

outliers in the original trait data may not be as effective in addressing evolutionary shifts.

Adams et al. (2024) demonstrated that when four different estimators (L1, S, M, and MM)

were integrated into both PGLS and PIC analysis, methods like PIC-MM and PIC-S

performed significantly better in detecting evolutionary relationships, especially when

accounting for outliers. Thus, computing PIC values is essential for effective analysis of the

relationships between traits with evolutionary shifts.

Dramatic changes in some evolutionary branches can result in several outliers in the PIC

dataset, distorting the statistical results of parametric tests. Nonparametric tests, which

measure the strength and direction of association between two variables without assuming a

specific distribution, can effectively handle data with outliers, reducing false positives.

Therefore, we propose a hybrid framework, Outlier-Guided Correlation (OGC), to optimize

the analysis of trait relationships under diverse evolutionary conditions. This framework

integrates the existing PIC method with a dynamic selection mechanism that evaluates the

presence of outliers in trait contrasts. For larger datasets, OGC dynamically applies Pearson or

Spearman correlation based on the presence of outliers. For smaller datasets, where

non-normality can significantly impact results, the framework incorporates both normality

testing and outlier detection, referred to as Outlier- and Distribution-Guided Correlation

(ODGC). Together, these approaches form a unified framework collectively referred to as

O(D)GC, designed to address the challenges posed by varying data conditions.

Our analysis demonstrates that PIC-O(D)GC effectively reduces false positives in the

presence of strong abrupt shifts, while maintaining high accuracy. When evolutionary shifts

are weak or absent, PIC-O(D)GC performs comparably to established phylogenetic methods,

including PGLS (optimized across five evolutionary models), PIC-Pearson, robust regression

methods, Corphylo (Zheng et al. 2009), PGLMM, and MR-PMM. Its performance closely

aligns with that of PIC-MM, with the added advantage of computational simplicity. This

consistency across a wide range of evolutionary scenarios highlights O(D)GC as a reliable and

scalable tool for analyzing trait correlations in phylogenetic datasets, complementing other

robust methods such as PIC-MM.

MATERIALS AND METHODS

Simulations of Phylogenetic Trees

To account for the influence of phylogenetic tree topology, we simulated two types of

phylogenetic trees. The first type is a fixed, fully balanced phylogenetic tree, where all

simulations were conducted using the same balanced dichotomous tree. The second type

consists of randomly generated phylogenetic trees based on a pure birth model, where the

ancestral root splits into two descendant subtrees of equal size. Asymmetry in each subtree is

introduced by varying the branching rates within each subtree, generating different tree

topologies for each simulation to explore diverse phylogenetic scenarios across replicates.

We set the sample size to 128 species and conducted simulations using both fixed and

random trees. Additionally, to investigate the performance of each model under small sample

conditions, we conducted a supplementary analysis with a sample size of 16 species, using

only the fixed tree topology.

Simulated Data with Abrupt Shifts

Two traits (X_1 and X_2) were independently simulated on a phylogenetic tree, with $X_2 = 0 \times X_1$

$+ \, e$, where both X_1 and e underwent an abrupt shift of equal magnitude on one of the branches

closest to the root, while on all other branches they evolved independently under the BM

model with a variance of 1 (Fig. 1). The shift magnitude ranged across values from 4^{-3} to 4^{7},

including both up-sizing and down-sizing shifts. Specifically, $4^{0} = 1$ represents no shift (i.e.,

no change in magnitude), while positive exponents correspond to increases in trait value, and

negative exponents correspond to decreases in trait value. Each shift gradient was simulated

1000 times, resulting in a total of 11,000 simulations. Despite the two traits being designed as

independent, significant correlations occasionally emerged, likely due to random noise

occurring in the same direction.

Simulated Data without Trait Jumps

We simulated two traits (X_1 and X_2) on the tree with the relationship $X_2 = X_1 + e$, where e

represents a noise term. By adjusting the variance of the noise term e, we varied the correlation

strength between X_1 and X_2. Referencing Revell (2010), our simulations included the

following scenarios:

1. **BM1 & BM1+BM2**: X_1 evolves under a BM process with a variance of 1, while e

follows another BM process with a variance of d, leading to X_2 representing the

combined BM1 and BM2 processes (BM1+BM2).

2. **BM & BM+Norm**: X_1 evolves under a BM process with a variance of 1, and e follows

a normal distribution (mean = 0, variance = d), resulting in X_2 combining the BM

process and the normal distribution (BM+Norm).

3. **Norm & Norm+BM**: X_1 follows a normal distribution (mean = 0, variance = 1), and e

evolves under a BM process (variance = d), making X_2 represent the combination of

the normal distribution and the BM process (Norm+BM).

The d takes values of 4^{-3}, 4^{-2}, 4^{-1}, 4^0, 4^1, 4^2, 4^3, 4^4, 4^5, 4^6, and 4^7 in all the three scenarios.

For each scenario and value of d, we conducted 1000 replicate simulations, totaling 33,000

simulations for each tree type.

This study did not include the fourth scenario from Revell (2010), Norm1 &

Norm1+Norm2, in which Trait X_1 follows a normal distribution with a mean of 0 and a

variance of 1, while X_2 evolves as X_1 plus a noise term e, where e follows a normal distribution

with a mean of 0 and a variance of d. This scenario results in simulated data with little to no

phylogenetic correlation, which, though useful as a control for the absence of phylogenetic

influence, was not included in this study. While the first three simulation scenarios primarily

model data with phylogenetic autocorrelation, large noise term variances in these scenarios

can also lead to data lacking phylogenetic autocorrelation, making a separate control scenario

unnecessary.

Statistical Methods Compared

To evaluate the performance of PIC-OGC examining the evolutionary relationship between X_1

and X_2 amid an abrupt evolutionary jump, we compared it with several previously established

phylogenetic comparative methods:

1. **PIC-Pearson**: Pearson correlation analysis of PIC values, equivalent to OLS

regression of PIC, which is commonly used in phylogenetic comparative analysis.

2. **PGLS**: Using five evolutionary models, BM (Felsenstein 1985), Pagel's λ (Pagel

1999), OU fixed, OU random (Hansen 1997), and EB (Harmon et al. 2010).

3. **Robust phylogenetic regression with both PIC and PGLS frameworks:** For each

framework, we utilized four robust estimators: L1, S, M, and MM. These were

implemented as PIC-L1, PIC-S, PIC-M, and PIC-MM for PIC-based methods, and

PGLS-L1, PGLS-S, PGLS-M, and PGLS-MM for PGLS-based methods (Adams et al.

2024).

4. **Corphylo**: A method that calculates Pearson correlation coefficients for multiple

continuous traits, allowing users to incorporate phylogenetic signal and measurement

errors into the model (Zheng et al. 2009). This method assumes an OU process for trait

evolution, making it particularly suitable for traits that deviate from the BM model.

Corphylo serves a dual purpose: it enables both the detection of phylogenetic signals in

individual traits and the examination of correlations between multiple traits. A key

feature of Corphylo is its ability to include measurement errors of trait data, potentially

improving accuracy under real-world conditions. While our simulations did not

explicitly account for measurement errors, this functionality highlights its potential

robustness in empirical applications.

5. **PGLMM** (Ives and Helmus 2011).

6. **MR-PMM** (Westoby et al. 2023, Halliwell et al. 2024).

Outlier- and Distribution-Guided Phylogenetic Correlation Analysis

To address the influence of evolutionary shifts and outliers in phylogenetic comparative

analysis, we introduced a novel framework: PIC-OGC. In this method, the choice between

Pearson and Spearman correlation analysis is guided by the identification of outliers within the

PIC data. The process starts by calculating the PICs for each trait using the phylogenetic tree

and determining their absolute deviations from the median PIC of the dataset. A PIC is defined

as an outlier if its absolute deviation from the median exceeds $T_i = n \times MAD$, where T_i

represents the threshold for detecting outliers, n is a sample-size-adjusted factor, and *MAD*

(Median Absolute Deviation) is the median of the absolute deviations from the median PIC of

the dataset (Wilcox 2003, Rindskopf and Shiyko 2010, Leys et al. 2013).

For a normal distribution, *MAD* is approximately equal to 0.67449 times the standard

deviation (σ) (Iglewicz and Hoaglin 1993, Wilcox 2003). The expected coverage of the data

under different multiples of *MAD* (Rindskopf and Shiyko 2010) is shown in Table 1, which

also provides the corresponding expected number of outliers. As the threshold increases, the

data coverage percentage approaches 100%, leading to a sharp decline in the expected number

of outliers. This reduction scales inversely with both the threshold and dataset size, reflecting

how stricter thresholds provide greater specificity at the cost of identifying fewer outliers.

Table 1. Data coverage and expected number of outliers under different thresholds for

detecting outliers.

Threshold ($n \times MAD$)	Coverage	Expected number of outliers	
		16 Species	128 Species
3	95.6%	0.7	6
4	99.26%	0.1	0.9
5	99.94%	0.009	0.08
6	99.994%	0.0009	0.008
7	99.9995%	0.00008	0.0006

This table presents the expected number of outliers and the corresponding data coverage

percentages for various outlier detection thresholds, defined as multiples of the Median

Absolute Deviation (*MAD*), under the assumption of a normal distribution. The expected

number of outliers is calculated for datasets with 15 and 127 phylogenetic independent

contrasts (PIC values), corresponding to phylogenetic trees with 16 and 128 species,

respectively.

For our datasets with 16 species (15 PIC values) and 128 species (127 PIC values), we set

thresholds at $5 \times MAD$ and $6 \times MAD$, respectively, to ensure that only data points with

substantial deviations are identified as outliers, thereby minimizing the risk of misclassifying

data points that do not significantly impact the performance of statistical analysis methods.

Using these thresholds, the expected number of falsely identified outliers for normal data is

less than one per hundred simulations, ensuring high specificity in identifying deviations

caused by evolutionary shifts or other phenomena beyond normal variation.

To further evaluate the robustness of our approach, we repeated the study using a more

stringent threshold to define outliers: $6 \times MAD$ for the datasets with 16 species and $7 \times MAD$ for

the datasets with 128 species. Although minor variations in specific metrics were observed

under these stricter thresholds, the overall ranking of the performance of the compared

methods was consistent (see the Results section for details).

If either of the two traits, X_1 or X_2 in a dataset, is identified as containing an outlier, the

entire dataset is flagged as outlier-containing. To further clarify, we considered an example

arising from random variation. For instance, when a data point exhibits extreme deviations,

such as $11 \times MAD$ for X_1 and $1 \times MAD$ for X_2, Pearson correlation may result in a false negative

if such points are not appropriately flagged as outliers. Specifically, if a true positive

correlation exists between X_1 or X_2 but the outlier is not removed, the extreme value can distort

the Pearson correlation coefficient, reducing its sensitivity and leading to an incorrect

conclusion of no correlation. Conversely, if no true correlation exists between X_1 and X_2,

assigning such points to Spearman correlation—even if they are incorrectly categorized as

outliers—would not inherently lead to false positives. This is because Spearman evaluates

rank-based monotonic relationships and is less influenced by the magnitude of extreme values

unless the rankings of the data points coincidentally align in a way that suggests a spurious

monotonic relationship. This example underscores the importance of defining outliers based

on deviations from the data's central tendency, regardless of whether the deviation arises from

evolutionary shifts or ordinary variation. Moreover, the design of OGC, which independently

evaluates outliers for each trait, theoretically enables it to handle scenarios where only one

trait experiences a shift. This adaptability ensures robust performance even in asymmetric

shift conditions, highlighting its versatility across a broader range of evolutionary scenarios.

Specifically, when no outliers are detected in either trait (X_1 or X_2), Pearson correlation is used

to capture the linear relationship. Conversely, when outliers are identified in one or both traits,

Spearman's rank correlation is employed to mitigate the influence of these extreme values,

ensuring that the method remains robust against deviations from normality.

For larger datasets with 128 species (127 PIC values), Pearson correlation, despite its

reliance on normality assumptions, is generally robust to minor deviations from normality due

to the central limit theorem. In such cases, the impact of skewness and kurtosis is reduced by

the large sample size, ensuring reliable correlation estimates. Therefore, for these datasets, the

choice of Pearson or Spearman correlation was determined solely by the presence or absence

of outliers.

For smaller datasets with 16 species (15 PIC values), both normality testing and outlier

detection were conducted. This approach, referred to as PIC-ODGC, dynamically selects the

appropriate correlation method based on data characteristics. If no outliers were detected and

normality assumptions were satisfied, the Pearson correlation was used. In cases where

outliers were present, or normality was violated, Spearman correlation was applied to

minimize the impact of extreme values on the results.

Benchmark for True Trait Correlations in Each Simulation

We aimed to compare the performance of different methods in detecting trait correlations

while minimizing false-positive relationships between X_1 and X_2. An effective method should

not only minimize false positives but also retain the ability to detect genuine, albeit rare,

correlations that might arise due to random noise. Striking this balance is critical, as a method

overly focused on avoiding false positives may miss true, meaningful correlations.

In the simulations without evolutionary shifts, traits X_1 and X_2 were simulated along a

phylogenetic tree, with their relationships defined as $X_2 = X_1 + e$, where e represents a noise

term. If the simulated formula is directly assumed to imply a significant positive correlation

between X_1 and X_2, the expected correlation may diminish or even disappear when the

variance of the noise term e is large, due to the overwhelming influence of noise. An accurate

method, in this case, would identify the dataset as lacking a significant correlation, which

contradicts the original assumption. Such discrepancies would label the method as producing a

false negative result. Conversely, when the simulation explicitly sets no relationship between

the two traits, random noise might generate spurious correlations. If a method accurately

detects such correlations, it would erroneously be marked as producing a false positive result,

penalizing its evaluation unfairly. These possibilities highlight the importance of carefully

assessing correlations rather than assuming them based solely on the model structure. Similar

concerns have been raised by Ives (2022), who emphasized that random errors in correlated

data are not mere noise but can obscure or even invert the expected relationships between traits,

particularly under conditions of high variance.

To define a robust benchmark for assessing true trait correlations, we adopted a

phylogenetically informed approach. True evolutionary relationships between X_1 and X_2 were

inferred by conducting traditional correlation tests on the normalized changes $\Delta X_1 / L$ and

$\Delta X_2 / L$, where L is the branch length. The changes themselves are independent across branches,

ensuring that conventional correlation tests can provide reliable evaluations of trait

relationships. The normalization by branch length serves to account for differences in

evolutionary time scales, making the comparisons more meaningful and biologically

interpretable.

Unlike our previous approach (Chen et al. 2023, Chen and Niu 2024), which relied on

normality testing to select between Pearson and Spearman correlations, this study adopts an

outlier-driven strategy. Outliers were identified as PIC values whose absolute deviation from

the median exceeded a threshold $T_i = n \times MAD$, where n was adjusted based on the dataset size.

This threshold was consistent with the one used in OGC, ensuring comparability in outlier

detection across benchmarks. When no outliers were detected, Pearson correlation was

applied; otherwise, Spearman rank correlation was used. This refined approach not only

reduces the influence of extreme values but also ensures that true evolutionary relationships

between traits with evolutionary shifts are accurately captured.

For smaller datasets with 16 species (15 PIC values), the benchmark also incorporated

normality testing alongside outlier detection to better account for the data's characteristics.

The same threshold applied in ODGC was used to define outliers, maintaining consistency in

detection criteria across different analytical frameworks. If no outliers were identified and the

normality assumption was satisfied, the Pearson correlation was used to assess the true

evolutionary relationships. Conversely, in cases where outliers were present, or normality was

violated, Spearman correlation was employed to mitigate the influence of extreme values.

This dual consideration of outlier presence and normality ensured that the benchmark

remained robust and adaptable to the unique challenges posed by small datasets.

Evaluating Concerns of Benchmark Bias in PIC-O(D)GC Assessment

Despite the robustness of our benchmark, concerns remain about potential biases when

assessing PIC-O(D)GC because the benchmark shares methodological similarities with the

PIC-O(D)GC framework itself. For example, both rely on outlier detection using a shared

threshold (e.g., $T_i = n \times MAD$) and employ the same decision-making process between

Pearson and Spearman correlation based on the presence of outliers or normality testing. Such

alignment raises the possibility that the benchmark could inadvertently favor PIC-O(D)GC by

design. If PIC-O(D)GC exhibits superior performance relative to other methods is evident, it is

prudent to question whether this advantage arises from genuine methodological robustness or

inherent alignment with the benchmark criteria.

To address this concern, we implemented a second benchmark that is entirely independent

of PIC-O(D)GC's design.   In this evaluation framework, we directly used the simulated

formula to assume the presence or absence of correlations between X_1 and X_2. Specifically,

when the relationship was defined as $X_2 = X_1 + e$, we treated X_1 and X_2 as correlated.

Conversely, when the simulation explicitly set X_1 and X_2 as uncorrelated (e.g., $X_2 = 0 \times X_1 + e$),

we used the absence of a correlation as the gold standard.

Although the second benchmark introduces limitations—namely, spurious correlations

caused by random noise could mislabel a method's correct detection as a false positive—its

influence on all methods being compared remains consistent. This uniform impact ensures a

logically unbiased assessment of relative performance, allowing us to validate PIC-O(D)GC's

performance under a complementary and independent criterion.

By applying both benchmarks, we aimed to mitigate potential bias and provide a more

comprehensive evaluation of the methods under consideration. In the Results section, unless

otherwise specified, the comparisons of different methods are based on the first benchmark's

determination of whether trait correlations exist within the data.

Programs and Packages

All simulations and analyses were performed in the R environment (version 4.0.2) (R Core

Team 2020) on a Linux system. The phytools package (version 2.3.0) (Revell 2024) was used

for tree generation. The geiger *package* (version 2.0.11) (Pennell et al. 2014) was used for

simulations. Normality tests were conducted using the Shapiro-Wilk test (Royston 1992) via

the *shapiro.test()* function from the *stats* package (version 4.3.1). Pearson and Spearman

correlation analyses were performed using the *cor.test()* function from the *stats* package. PICs

were calculated using the *pic()* function from the ape package (version 5.8) (Paradis and

Schliep 2019). All PGLS regressions were conducted using the *phylolm()* function from the

phylolm package (version 2.6.2) (Ho and Ane 2014). Corphylo was conducted using the

corphylo() function from the *ape* package (version 5.8) (Zheng et al. 2009). PGLMM and

MR-PMM were both conducted using the *MCMCglmm* function from the *MCMCglmm*

package (version 2.36) (Hadfield and Nakagawa 2010).

RESULTS

Impact of Abrupt Evolutionary Shifts on Data Patterns and Correlation Analysis

To assess PIC-OGC's ability to handle abrupt evolutionary shifts, we simulated two traits on a

balanced phylogenetic tree with 128 species. Each trait underwent a shift near the root, while

other branches evolved independently under a BM model (variance = 1, Fig. 1A). Shift

magnitudes ranged from 4^{-3} to 4^7, including both increases and decreases, with 4^0 representing

no shift. From these simulations, we selected three datasets as case studies to examine how

shifts affect both the overall trait patterns and the resulting PIC contrasts.

As shown in Figures 1B, 1D, and 1F, an abrupt shift in trait evolution causes the raw trait

values to form two distinct clusters in the coordinate space, with the separation between

these clusters increasing as the shift magnitude becomes larger. In the PIC coordinate system,

the shift manifests as an outlier in the PIC data, with the degree of outlier deviation

becoming more pronounced as the shift magnitude increases (Figs. 1C, 1E, and 1G).

When phylogenetic autocorrelation is ignored, directly applying Pearson or Spearman

correlation to the raw trait data frequently results in false positives, regardless of the shift

magnitude (Figs. 1B, 1D, and 1F).In contrast, controlling for phylogenetic autocorrelation by

using PIC values instead of raw trait data eliminates false positives when there is no shift or

when the shift is small (Figs. 1C and 1E). However, under large shifts, Pearson correlation

becomes highly sensitive to the outlier, leading to a false positive result and erroneously

inferring a significant positive correlation between X_1 and X_2, akin to non-phylogenetic

methods. Conversely, the Spearman correlation applied to the PIC data under large shifts

avoids false positives by correctly detecting no significant correlation between the two traits

(Fig. 1G).

Figure 1. Effects of abrupt shifts on data pattern and correlation analysis. (A) A balanced

phylogenetic tree with 128 species, with the thick red line indicating an abrupt shift. (B, C)

Simulation outcomes without any shifts (1-fold shift). (D, E) Simulation outcomes with a

4-fold shift. (F, G) Simulation outcomes with a 16-fold shift. The phylogenetically

independent contrasts (PIC) values affected by the evolutionary shift are highlighted as

enlarged red dots. The simulations illustrate that significant evolutionary shifts lead to the

emergence of outliers in the PIC values. When $3 \times MAD$ (Median Absolute Deviation) is used

as the threshold for detecting outliers, all three PIC datasets contain identifiable outliers. In the

weak-shift PIC dataset (E), the red dot is recognized as an outlier; however, it is not the point

with the greatest deviation from the median. In the strong-shift PIC dataset (G), the red dot

represents the most deviated point, with X_1 at $14 \times MAD$ and X_2 at $15 \times MAD$. This figure

highlights the performance of traditional correlation methods and the varying effectiveness of

parametric and nonparametric correlation analyses of PIC under different shift magnitudes.

The data used to plot this figure are available in Supplementary Table S1

Refining Method Selection for Comparison with PIC-O(D)GC

Directly comparing all 18 parameter-model combinations from the methods listed in the

Materials and Methods section would lead to an overly complex and unclear presentation,

masking critical differences between approaches. To ensure a clear and focused evaluation of

PIC-O(D)GC against other phylogenetic comparative methods, we refined the comparison by

selecting only the best-performing models within each category.

For PGLS, the Akaike information criterion (AIC) was used to identify the optimal

evolutionary model—BM, λ, OU fixed, OU random, or EB—for each dataset. Only the

best-fitting PGLS model for each dataset was included in the final analysis.

For robust phylogenetic regression, which encompasses two methodological types

(PGLS- and PIC-based regression) and four robust estimators (L1, S, M, and MM), we first

evaluated their performance using simulations on the 128-species fixed tree.

Facing evolutionary shifts, PIC-MM stands out as the best method, consistently

maintaining low false positive rates across all shift gradients, with rates remaining stable

within narrow bounds of 0.046–0.062, demonstrating remarkable robustness and reliability

(Fig. 2A). PIC-S also showed stability across different shift magnitudes, with false positive

rates ranging from 0.098 to 0.103. However, PIC-S consistently exhibited higher false positive

rates compared to PIC-MM, indicating inferior control over false positives despite its stability.

Other methods, whether based on PIC or PGLS, showed acceptable performance under weak

shifts (shift ≤ 4), with lower false positive rates. However, their false positive rates sharply

increased to 1 under stronger shifts (shift > 4), indicating their instability in managing abrupt

evolutionary changes effectively. Interestingly, PIC-L1 and PGLS-L1 exhibited lower false

positive rates than PIC-MM under weak shifts (shift ≤ 4), suggesting that the L1 estimator may

have potential advantages in specific contexts.

Accuracy reflects the combined influence of false positive and false negative rates by

measuring the proportion of correct predictions (true positives and true negatives), offering a

holistic view of the methods' overall performance. Figure 2B confirms that PIC-MM is the

most reliable method, achieving both high accuracy and stability across all evolutionary shift

gradients. PIC-S follows as the second-best performer, while PIC-L1 and PGLS-L1 slightly

excel under low shift conditions. The trends in accuracy largely align with the false positive

rates in Figure 2A, underscoring the strong correlation between these two metrics for robust

methods.

Figures 2C-2E show the accuracy of various methods across three no-shift evolutionary

scenarios: BM1 & BM1+BM2, BM & BM+Norm, and Norm & Norm+BM. PIC-MM is

generally among the best-performing methods, though it is occasionally slightly outperformed

in specific cases, such as under high variance in the BM & BM+Norm scenario, where PIC-L1

and PGLS-L1 show a clear, though modest, advantage. PIC-S, on the other hand, is no longer

the second-best performer; in fact, it ranks among the worst in BM1 & BM1+BM2 and Norm

& Norm+BM, showing considerable performance degradation under these conditions.

Figure 2. Performance of eight robust phylogenetic regression methods. This figure shows

their false positive rates (A) and accuracies (B) across evolutionary shift gradients and their

accuracies across three simulation scenarios without abrupt shifts (C-E). The datasets were

simulated using a fixed-balanced tree of 128 species. The data used to plot this figure are

available in Supplementary Table S2. Please refer to the Materials and Methods section for

additional details on the simulation setup, variance gradients, and an overview description of

the eight methods.

When analyzing randomly generated 128-species trees and 16-species fixed trees,

PIC-MM consistently demonstrates the strongest ability to handle evolutionary shifts. Across

the three simulation scenarios without evolutionary shifts, PIC-MM performs at least as well

as, or not significantly worse than, the other methods (Supplementary Table S2). Even when

the outlier detection threshold was adjusted from $6 \times MAD$ to $7 \times MAD$ for 128-species datasets,

PIC-MM maintained its top performance across both random and fixed trees (Supplementary

Table S3). These results highlight PIC-MM's robustness and versatility, making it a reliable

tool for analyzing phylogenetic data under both stable and dynamic evolutionary conditions.

These findings align with the conclusions of Adams et al. (2024). Consequently, PIC-MM was

selected as the representative robust regression method for comparison with PIC-O(D)GC.

Through this optimization and selection process, we reduced the total number of methods

compared with PIC-O(D)GC to six, facilitating a more focused and interpretable evaluation.

PIC-OGC: Robust Performance Across Evolutionary Shifts

We compared the performance of PIC-OGC with six other methods using simulations on the

128-species fixed tree. As shown in Figure 3A, there are significant differences in false

positive rates among the methods. Under weak shifts (shift ≤ 4), PIC-OGC consistently

exhibits the same false positive rates as PIC-Pearson, which is equivalent to traditional

PIC-OLS regression. These rates are the lowest among the seven methods, and this similarity

can be attributed to the fact that weak or downsizing evolutionary shifts do not generate

outliers in the PIC datasets. However, when the shift becomes stronger (> 4), the false positive

rate of PIC-Pearson sharply increases to 1, while PIC-OGC maintains a stable false positive

rate of 0.039. This result demonstrates that PIC-OGC is significantly more robust against

evolutionary shifts than traditional PIC regression methods. PIC-MM also shows robustness

against evolutionary shifts but has slightly higher false positive rates than PIC-OGC. Corphylo

shows similar performance to PIC-OGC and PIC-Pearson under weak shifts (shift ≤ 4), but its

false positive rates experience a sharp increase to 1 under stronger shifts, similar to

PIC-Pearson. PGLMM and PGLS also show low false positive rates under weak shifts (shift ≤

4) but experience sharp increases to 1 under stronger shifts (shift > 4). MR-PMM performs the

worst, with the highest or among the highest false positive rates across 10 out of 11 shift

gradients.

Figure 3B displays the accuracy of the same seven methods under the same 11

evolutionary shift gradients. Accuracy here serves as a comprehensive metric, reflecting both

false positive and false negative rates. Notably, PIC-OGC demonstrates the highest accuracy

or shares the top position across all shift gradients. PIC-MM also performs exceptionally well,

with accuracy very similar to PIC-OGC. The other methods show acceptable performance

under weak shifts but exhibit unacceptably low accuracy under stronger shifts. Despite PGLS

results being derived from selecting the best-fitting model from five candidate models for each

simulation, its performance still falls significantly short of PIC-OGC and PIC-MM.

Figure 3. False positive rates (A) and accuracies (B) of seven methods on simulations across

evolutionary shift gradients simulated on a fixed balanced tree of 128 species. The data used to

plot this figure, including false positive rates and accuracy metrics, are available in

Supplementary Table 4. Please refer to the Materials and Methods section for additional

details on the simulation setup, variance gradients, and an overview description of the seven

methods.

Fixed trees represent a fully balanced phylogenetic structure, where all simulations are

conducted using the same balanced dichotomous tree. This consistent topology ensures

uniformity in the branching structure, resulting in homogenized data. In contrast, random trees

are generated based on a pure birth model, introducing asymmetry by varying the branching

rates within each subtree. This variability results in diverse tree topologies across simulations,

reflecting a wider range of phylogenetic scenarios and introducing additional heterogeneity in

the data. In our analysis with 128-species random trees (Supplementary Table S4), most

methods showed performance very similar to their results under fixed tree simulations, with

PIC-OGC and PIC-MM consistently demonstrating the best performance overall. However,

the slight yet consistent edge that PIC-OGC held over PIC-MM across all shift gradients under

fixed tree simulations was no longer observed with random trees. Instead, each method

exhibited slight advantages under specific gradients, emphasizing their complementary

strengths in handling the complexities introduced by varying phylogenetic structures.

Additionally, MR-PMM exhibited a significant improvement under low shift gradients (shift

≤ 4). While it had the worst performance with fixed trees, it became the method with the

lowest false positive rate and highest accuracy in weak conditions, outperforming all other

methods.

All the above results were obtained using a $6 \times MAD$ threshold to define outliers. Using a

$7 \times MAD$ threshold yielded highly similar results, with PIC-OGC and PIC-MM remaining the

top-performing methods (Supplementary Table S5).

PIC-OGC: Strong Performance in Simulations Without Abrupt Shifts

The analysis here focuses on simulations conducted on a 128-species fixed tree, where two

traits exhibit a relationship of $X_2 = X_1 + e$, with e representing random noise varying across 11

gradients of variance displayed on the x-axis of Figure 4. These simulations aim to evaluate

the methods' performance in detecting relationships between evolutionary traits under

conditions without abrupt shifts.

In the BM1 & BM1+BM2 scenario (Fig. 4A), the three PIC-based methods (PIC-OGC,

PIC-MM, and PIC-Pearson), along with Corphylo, PGLS, and PGLMM, maintain comparably

high accuracy across all variance gradients, demonstrating their robustness in handling traits

with strong phylogenetic signals. MR-PMM, however, exhibits a distinct pattern, showing a

significant decline in accuracy at intermediate variance gradients while maintaining

comparable performance at both low and high variance levels.

In the BM & BM+Norm scenario (Fig. 4B), the three PIC-based methods (PIC-OGC,

PIC-MM, and PIC-Pearson) and Corphylo clearly outperform MR-PMM, PGLS, and

PGLMM.

In the Norm & Norm+BM scenario (Fig. 4C), all methods achieve very high accuracy

(close to 1) at low variance gradients ($\leq 4^2$), except for Corphylo, which underperforms at the

lowest variance gradient (4^{-3}). As variance increases to moderately high gradients, the

accuracies of the six methods—PIC-OGC, PIC-MM, PIC-Pearson, Corphylo, PGLS, and

PGLMM—decline significantly and consistently. In contrast, MR-PMM shows a unique

pattern distinct from the other methods, with neither a clear advantage nor disadvantage in

overall performance but demonstrating context-specific strengths.

Across all three simulation scenarios, the three PIC-based methods (PIC-OGC, PIC-MM,

and PIC-Pearson) consistently exhibit near-optimal performance, highlighting their

robustness in handling phylogenetic traits without abrupt shifts.

Figure 4. Accuracy of seven methods on simulations without abrupt shifts. Simulations were

conducted on a fixed balanced tree of 128 species, with relationships between traits defined as

$X_2 = X_1 + e$, where e represents random noise, across three scenarios: (A) BM1 & BM1+BM2,

(B) BM & BM+Norm, and (C) Norm & Norm+BM. The data used to plot this figure are

available in Supplementary Table S4. Please refer to the Materials and Methods section for

additional details on the simulation setup, variance gradients, and an overview description of

the seven methods.

Results from simulations on the 128-species random trees show overall trends consistent

with those observed on the fixed tree, reaffirming the strong performance of the three

PIC-based methods across diverse phylogenetic topologies (Supplementary Table S4). While

their overall performance remains excellent, PIC-OGC shows a slight drop at two specific

variance gradients (4^4 and 4^5) of the last simulation scenario (Norm & Norm+BM), where it no

longer ranks among the top three methods. These results indicate a minor, context-dependent

limitation of PIC-OGC in random tree simulations.

The above results on methods' accuracies detecting relationships between evolutionary

traits under conditions without abrupt shifts were obtained using a $6 \times MAD$ threshold to define

outliers. Using a $7 \times MAD$ threshold yielded highly similar results, with the three PIC-based

methods remaining the top-performing methods (Supplementary Table S5).

Performance of the Methods in Detecting Relationships Beyond BM Assumptions

Given that the computation of PIC values relies strictly on the assumption of a BM model, it is

logically problematic for non-BM data. As a result, PIC-Pearson, which uses Pearson

correlation analysis on PIC values, is excluded from further comparisons in such cases

because its theoretical foundation no longer holds for non-BM data. However, PIC-MM and

PIC-OGC, despite being rooted in PIC calculations, are retained for further analysis due to

their strong performance in earlier results (Figs. 3–4), raising the question of whether their

robustness extends to non-BM conditions. PGLS is included because it is a more flexible

framework that can adapt to different evolutionary models by incorporating the appropriate

covariance structure, making it suitable for non-BM data. Corphylo is retained as it assumes

an OU process for trait evolution, making it suitable for data that deviate from BM

assumptions. PGLMM and MR-PMM are also retained because they do not strictly rely on

BM assumptions, and their flexibility in modeling phylogenetic relationships and handling

complex data may allow them to outperform PIC-based methods under non-BM conditions.

Garland et al. (1992) clarified that the application of PIC is not strictly limited to gradual,

clock-like evolution as modeled by BM. Instead, PIC remains applicable to non-BM scenarios,

provided that branch lengths and evolutionary models are appropriately transformed to

account for deviations from BM assumptions. Building on this foundational flexibility, our

study evaluates the performance of PIC-OGC and PIC-MM under untransformed conditions,

testing whether these methods can deliver reliable results even without adjustments to branch

lengths or models.

To classify simulated data as either BM or non-BM condition, we calculated the AIC

values from PGLS regression models fitted under five evolutionary models: BM, lambda, OU

random, OU fixed, and EB. The data were classified as BM condition if the BM model yielded

the lowest AIC; otherwise, they were considered non-BM condition.

Generally, the two PIC-based methods, PIC-OGC and PIC-MM, maintained their

superiority under non-BM conditions involving evolutionary shifts (Figs. 5A–5B). PIC-OGC

performed exceptionally well, achieving almost the lowest false positive rates and highest

accuracy across all shift gradients. PIC-MM also demonstrated strong overall performance,

though it exhibited a high false positive rate and low accuracy at a specific shift gradient (4).

Despite being first optimized for performance by selecting the best-fitting evolutionary model,

PGLS exhibited the weakest performance among retained methods under non-BM conditions,

showing high false positive rates and low accuracy, particularly at higher shift gradients.

In scenarios without evolutionary shifts (Fig. 5C), none of the six retained methods

showed a consistent advantage or disadvantage. Performance varied across different variance

levels, with no single method consistently emerging as the clear best or worst. Similar trends

were observed when using random trees (Supplementary Table S6) or adjusting the threshold

for defining outliers (Supplementary Table S7), reinforcing the robustness of these

conclusions across varying tree structures and outlier detection criteria.

Figure 5. Performance of six methods across datasets that deviate from BM assumptions. (A).

The false positive rates of six methods under datasets with evolutionary shifts. (B). The

accuracy of the methods under the same conditions. (C) combines results from datasets

without evolutionary shifts (three no-shift simulation scenarios) under non-BM conditions.

All the datasets were simulated using a fixed-balanced tree of 128 species. They were

classified as BM or non-BM conditions based on AIC values from PGLS regression models

fitted under five evolutionary models (BM, lambda, OU random, OU fixed, and EB). Data

were classified as BM condition if the BM model had the lowest AIC; otherwise, they were

considered non-BM condition. The data used to plot this figure are available in Supplementary

Table S6. Similar trends were observed when analyzing randomly generated 128-species trees

(Supplementary Table S6). Please refer to the Materials and Methods section for additional

details on the simulation setup, variance gradients, and an overview description of the six

methods.

Comparison of the accuracies under conditions with and without shifts (Fig. 3B vs Fig.

4A–C; Fig. 5B vs Fig. 5C; see also Supplementary Tables S2–S7) reveals that methods other

than PIC-OGC and PIC-MM effectively handle low shift and low variance conditions but

generally perform poorly under medium-to-high shift or variance conditions. Notably, their

ability to cope with medium-to-high variance is significantly better than their performance

under medium-to-high shifts.

PIC-ODGC: Robust Performance with Small Samples

For small samples consisting of 16 species (15 PIC values), the choice between Pearson and

Spearman correlation is influenced not only by the presence or absence of outliers but also by

whether the data follow a normal distribution. To address this, we enhanced PIC-OGC to

PIC-ODGC, incorporating both outlier detection and normality testing.

Using a fixed tree with 16 species, we replicated the analyses conducted with the

128-species fixed tree. Two thresholds for defining outliers, $5 \times MAD$ and $6 \times MAD$, were

applied. The conclusions drawn from the 128-species dataset regarding PIC-OGC and

PIC-MM were consistently confirmed in the 16-species dataset, with PIC-ODGC and

PIC-MM emerging as the top-performing methods (see Supplementary Tables S2–S7). This

consistency highlights the robustness of these methods, even under small sample conditions.

Addressing Concerns about PIC-Based Performance

PIC-O(D)GC and PIC-MM, being rooted in PIC calculations, might initially raise concerns

about whether their observed strong performance stems from a bias due to the gold

standard's reliance on calculating $\Delta X_1 / L$ and $\Delta X_2 / L$, a process that closely mirrors the PIC

estimation procedure. This concern is particularly pronounced for PIC-O(D)GC, as it also

employs the same outlier detection thresholds and criteria for selecting Pearson or Spearman

correlation.

To address potential biases in the original benchmark, we validated PIC-O(D)GC and

PIC-MM using a second, entirely independent benchmark. Unlike the first benchmark, this

alternative framework does not rely on any assumptions or calculations related to PIC or any

specific method. Instead, it directly determines the presence or absence of correlations based

on the simulated formula: X_1 and X_2 are treated as correlated when defined as $X_2 = X_1 + e$, and

uncorrelated when defined as $X_2 = 0 \times X_1 + e$. Although this benchmark may classify spurious

correlations caused by random noise as false positives, its design is method-agnostic,

ensuring no bias toward or against any specific approach.

Using the second benchmark, we compared the methods in their false positive rates

under shift conditions and false negative rates under no-shift conditions.

First, we observed no substantial differences in the performance of various methods

between 128-species random and fixed trees.

Among the eight phylogenetically robust methods, PIC-MM consistently demonstrated

the lowest false positive rates, reaffirming its superior position within this group

(Supplementary Table S8). Under no-shift conditions with 128 species, PIC-MM performed

similarly to three comparable methods (PIC-M, PGLS-M, and PGLS-MM), all significantly

outperforming the other four methods. However, with 16 species, none of the eight methods

exhibited clear superiority in terms of false negative rates.

For the false positive rates under shift conditions, PIC-OGC and PIC-MM performed

almost identically, both achieving top performance with consistently low false positive rates

(Supplementary Table S9). In the three no-shift evolutionary scenarios, neither PIC-OGC nor

PIC-MM consistently achieved the lowest false negative rates, but both demonstrated

generally acceptable performance. Specifically, MR-PMM exhibited significantly higher

false negative rates in the BM1 & BM1+BM2 scenario and markedly lower false negative

rates in the Norm & Norm+BM scenario, while the other methods, including PIC-OGC and

679 PIC-MM, showed no substantial differences. In the BM & BM+Norm scenario, PGLS and

680 PGLMM performed slightly better than the other methods.

681  For the 16-species fixed tree, PIC-ODGC's false positive rates were similar to those of

682 PIC-MM, outperforming all other methods (Supplementary Table S9). Under the three

683 no-shift evolutionary scenarios, none of the seven methods showed distinct advantages or

684 disadvantages.

685  Under non-BM conditions, the performance of various methods was largely consistent

686 across the 128-species fixed tree, 128-species random trees, and the 16-species fixed tree. In

687 scenarios with evolutionary shifts, PIC-O(D)GC and PIC-MM maintained stable and low

688 false positive rates, reaffirming their robustness in handling such conditions. However, under

689 no-shift conditions, the false negative rates of PIC-ODGC and PIC-MM were comparable to

690 those of other methods, showing no clear advantage or disadvantage.

691  The use of two entirely different benchmarks for defining the presence or absence of

692 correlations between X_1 and X_2 did not lead to differences in the ranking of method

693 performance. This consistency highlights that the advantages demonstrated by PIC-O(D)GC

694 and PIC-MM across various scenarios, including their stable and low false positive rates

695 under evolutionary shifts, are not merely a result of their apparent similarity to the first

696 benchmark. Instead, these strengths are driven by their inherent robustness and adaptability

697 in handling phylogenetic data under varying evolutionary conditions. Moreover, the

698 alignment of method performance trends under both benchmarks reinforces the reliability of

699 the first benchmark. This consistency suggests that the first benchmark is not biased in favor

700 of any specific method but rather serves as a fair and effective framework for evaluating

701 phylogenetic comparative methods.

703 **DISCUSSION**

Equivalence of Pearson Correlation and OLS in Context

This study employed Pearson correlation to evaluate PIC relationships, which is

mathematically equivalent to the OLS regression commonly used in previous studies.

Pearson correlation provides the same statistical significance and directionality as OLS

regression, ensuring that our results are directly comparable with prior work. We chose to

use Pearson correlation for its computational simplicity and consistency with other

nonparametric methods, such as Spearman correlation, which is a key component of the

PIC-O(D)GC framework. The equivalence between Pearson correlation and OLS regression

ensures logical and statistical consistency with previous findings while reinforcing the

relevance and robustness of our results within the context of phylogenetic comparative

analyses.

Limitations of Traditional Data Transformations in Phylogenetic Correlation Analysis

A potential concern readers may raise is the complexity of the PIC-O(D)GC framework.

Given its focus on handling non-normal data distributions and outliers, why not simply apply

data transformations (e.g., logarithmic or square root transformations) to standardize the data

before using the traditional PIC-OLS regression? Such transformations are well-known for

their ability to address skewness and improve normality, potentially achieving comparable

results with much less complexity.

While data transformations are a commonly used strategy, their application to PIC data

in phylogenetic analyses faces several critical limitations:

1. Negative Values in PIC Data. PIC calculations often result in negative values, which

cannot be directly transformed using logarithmic or square root functions. Adding a

constant to shift the data into a positive range is a potential workaround, but this

approach introduces arbitrary biases, especially when the data span a wide range of

values.

2. Ineffectiveness of Yeo-Johnson Transformation. The Yeo-Johnson transformation is

designed to handle both positive and negative values, making it seemingly suitable for

phylogenetic data. However, its effectiveness in achieving normality is severely

limited for PIC data. In our prior studies using random tree simulations with 128

species, none of the datasets $\Delta X_1/L$ or $\Delta X_2/L$, showed normal distributions. Even after

applying the Yeo-Johnson transformation, fewer than 1% of the datasets achieved

normality (Chen and Niu 2024). Moreover, most phylogenetic trees in real-world

research are non-balanced, closely resembling the random trees used in our

simulations. Such trees inherently generate $\Delta X_1/L$ and $\Delta X_2/L$ values, as well as PIC

data, that deviate significantly from normality. This highlights not only the inherent

challenges of achieving normality in phylogenetic datasets but also the practical

limitations of relying on transformations like Yeo-Johnson to address these challenges

effectively.

3. Outlier Sensitivity. Traditional data transformations focus on normalizing

distributions but do not explicitly address outliers, which can significantly impact the

performance of correlation analyses. PIC-O(D)GC, by dynamically selecting Pearson

or Spearman correlation based on the presence of outliers, provides a tailored

approach that is inherently robust to extreme values.

749 Comparing Statistical Methods for Detecting Trait Correlations

Our study evaluated the performance of various statistical methods in detecting trait

correlations under diverse evolutionary scenarios, including with and without abrupt shifts.

PIC-O(D)GC and PIC-MM consistently demonstrated robust performance, particularly

excelling under shift scenarios by minimizing false positives and maintaining high accuracy.

In contrast, traditional methods like PIC-Pearson and PGLS, as well as robust variants such

as PGLS-MM, struggled under high shift gradients, highlighting their limitations in

challenging conditions.

In no-shift scenarios, PIC-MM performed comparably to other reliable methods, such as

PIC-L1, PGLS-L1, and Corphylo, highlighting the role of estimator choice within

regression-based methods. Meanwhile, PIC-O(D)GC maintained strong performance

through its hybrid correlation framework, independent of estimator-specific considerations.

The resilience of PIC-O(D)GC in adapting to outliers and non-normal data further supports

its utility across stable and dynamic datasets.

Advantages of PIC-OGC: Computational Efficiency and Simplicity

Among robust phylogenetic regression estimators, MM estimators were recommended by

Adams et al. (2024) for their strong performance across various scenarios. Consistent with

their findings, we observed that PIC-MM effectively controls false positive rates in datasets

with shifts. Similarly, PIC-O(D)GC exhibits comparable robustness, aligning closely with

PIC-MM in both false positive rate and accuracy.

However, the computational efficiency of PIC-O(D)GC offers a distinct advantage over

existing approaches, particularly the PIC-MM method. Both methods share an initial

computational complexity of $O(n)$ for calculating PIC values, where n represents the number

of taxa. The key difference lies in the subsequent steps:

1. PIC-MM employs iterative adjustments of model parameters, requiring k iterations to

converge. Each iteration involves recalculating residuals and adjusting weights,

resulting in a total complexity of $O(k \cdot n)$.

2. In contrast, PIC-OGC identifies outliers by evaluating the deviation of each PIC

value from the median. If no outliers are detected, a parametric correlation method

(Pearson) is applied; otherwise, a nonparametric method (Spearman) is used. The

outlier detection step is computationally lightweight $O(n)$ and does not impact the

overall complexity, which remains $O(n\log n)$, driven by the sorting step in

Spearman's rank correlation.

PIC-OGC's lower computational complexity provides a clear advantage in scalability,

making it particularly well-suited for large phylogenetic datasets. This efficiency, combined

with its robustness to abrupt evolutionary changes, positions PIC-O(D)GC as a scalable and

practical alternative to MM-based methods for phylogenetic comparative analyses.

For small datasets, the OGC framework is extended to PIC-ODGC by incorporating

normality testing to address the increased sensitivity of parametric methods to non-normal

distributions. The Shapiro-Wilk test, commonly used for normality assessment, adds an

additional computational complexity of $O(n^2)$ due to pairwise comparisons. As a result,

PIC-ODGC has a higher computational complexity compared to PIC-MM. However, this

complexity is negligible in practice for small sample sizes, such as datasets with 16 species,

where the additional computation does not impose a significant burden. While PIC-ODGC

maintains the computational simplicity and robustness of PIC-OGC, the added step of

normality testing introduces an additional layer of decision-making that may not always be

necessary. Nonetheless, its adaptive framework ensures accuracy across diverse scenarios,

making it a valuable extension of PIC-OGC for scenarios involving small datasets or

heightened sensitivity to data distribution characteristics.

Another notable advantage of PIC-O(D)GC is its simplicity and accessibility. By

dynamically adapting to the presence or absence of outliers and data normality,

PIC-O(D)GC effectively balances robustness and interpretability. Its reliance on

straightforward statistical concepts, such as Spearman's rank correlation, makes it easier to

understand and interpret, even for researchers with limited statistical training. By avoiding

assumptions about data distribution, PIC-O(D)GC offers intuitive and transparent results,

reducing the risk of misinterpretation and enhancing accessibility for a broader range of

biologists.

Advantages of Correlation-Based Methods over Regression-Based Approaches

Correlation-based methods such as PIC-O(D)GC, Corphylo, and MR-PMM have notable

advantages over regression-based approaches like PIC-OLS, PGLS, and PGLMM. The latter

methods, similar to OLS regression, inherently assume a directional causal relationship,

where the independent variable is considered error-free, or its errors are negligible (McArdle

1988, Osborne and Waters 2002, Casson and Farmer 2014, Jarantow et al. 2023). This

assumption implies that the values of the independent variable are precise, while the

dependent variable may include a degree of error. However, when the independent variable

is subject to measurement error, this can lead to biased regression coefficients—a

phenomenon known as "measurement error bias."

In contrast, correlation-based methods do not require assumptions about the

directionality of causal relationships and are unaffected by measurement errors in either

variable. This characteristic makes correlation-based methods particularly robust in scenarios

where the precision of independent variables cannot be guaranteed. While correlation-based

approaches cannot fully address all statistical challenges, their focus on detecting

associations rather than causation makes them more reliable under conditions where

measurement errors are present in both traits.

Limitations of Correlation-Based and Nonparametric Methods in Phylogenetic

Analysis

While correlation-based methods like PIC-O(D)GC offer notable advantages, including

robustness to outliers and flexibility in handling non-normal data, they share certain

limitations inherent to this class of methods. These methods, by design, cannot capture

non-monotonic relationships, which limits their ability to detect more complex evolutionary

patterns. This constraint applies to all correlation-based approaches, as they inherently focus

on assessing monotonic associations between variables.

Moreover, correlation-based methods, including PIC-O(D)GC, are not equipped to

handle interactions or analyze multiple predictors simultaneously, a key capability of

regression-based approaches. For instance, regression methods such as PGLS can model

complex relationships between traits and account for confounding factors, providing a more

comprehensive framework for studying evolutionary mechanisms. By relying solely on

pairwise correlations, methods like PIC-O(D)GC are inherently limited in their ability to

unravel multi-trait co-evolutionary dynamics.

Another critical limitation of nonparametric methods, including PIC-O(D)GC, lies in

their reliance on testing null hypotheses. While rejecting null models is informative for

identifying significant associations, it offers limited insight into the underlying evolutionary

processes. Parameter-based methods, such as regression models, provide richer information

through the estimation of meaningful parameters. For example, regression coefficients (β) in

PGLS models directly quantify the influence of one trait on another, enabling researchers to

test mechanistic hypotheses about co-evolution and trait evolution rates.

Despite these limitations, correlation-based methods remain valuable tools for

addressing specific challenges in phylogenetic data analysis. They should not be viewed as

replacements for parametric models but rather as complementary approaches that provide

851 robust and computationally efficient solutions, particularly in scenarios with non-normal

852 data distributions, outliers, or small sample sizes.

853 **FUNDING**

854 This study was supported in part by the National Natural Science Foundation of China (grant

855 number 31671321), with leftover resources used for occasional research expenses.

856 **ACKNOWLEDGMENTS**

857 Many sentences in this article were written with the assistance of ChatGPT-4 (OpenAI 2024).

858 The authors thoroughly reviewed and revised each sentence and took full responsibility for the

859 language and content of the entire article.

860 **DATA AVAILABILITY**

861 Supplementary Tables S1–S8 are provided in the file Supplemental_Tables.xlsx. The raw data

862 for the simulated phylogenetic analyses, including trait correlation data for 16-species fixed

863 trees, 128-species fixed trees, and 128-species random trees across shift and non-shift

864 scenarios, are available in RawData.zip. The scripts used to generate the results presented in

865 the main text— covering phylogenetic tree construction and trait data simulation

866 (Simulation_01_FixedBalancedTree.R and Simulation_02_RandomTree.R), outlier detection

867 (Analysis_05_MAD.R), regression model fitting (Analysis_01_PIC.R,

868 Analysis_02_PGLS_RPR.R, Analysis_03_Corphylo.R, Analysis_04_PGLMM_MRPMM.R),

869 are provided in Code.zip.

871 **REFERENCES**

872 Adams R., Cain Z., Assis R., DeGiorgio M. 2024. Robust phylogenetic regression. Syst.

873 Biol., 73:140-157.

Casson R.J., Farmer L.D. 2014. Understanding and checking the assumptions of linear

regression: a primer for medical researchers. Clin. Exp. Ophthalmol., 42:590-596.

Chen Z.-L., Guo H.-J., Niu D.-K. 2023. Dependent variable selection in phylogenetic

generalized least squares regression analysis under Pagel's lambda model.

bioRxiv:2023.005.2021.541623.

Chen Z.-L., Niu D.-K. 2024. Optimizing variable selection in phylogenetic eigenvector

regression for trait correlation analysis. bioRxiv:2024.2004.2014.589420.

Cornwallis C.K., Griffin A.S. 2024. A guided tour of phylogenetic comparative methods for

studying trait evolution. Annu. Rev. Ecol. Evol. Syst., 55:181-204.

Duchen P., Leuenberger C., Szilagyi S.M., Harmon L., Eastman J., Schweizer M., Wegmann

D. 2017. Inference of evolutionary jumps in large phylogenies using Levy processes. Syst.

Biol., 66:950-963.

Felsenstein J. 1985. Phylogenies and the comparative method. Am. Nat., 125:1-15.

Freckleton R.P., Harvey P.H., Pagel M. 2002. Phylogenetic analysis and comparative data:

A test and review of evidence. Amer. Natur., 160:712-726.

Gao Y., Wu M. 2022. Microbial genomic trait evolution is dominated by frequent and rare

pulsed evolution. Sci. Adv., 8:eabn1916.

Garamszegi L.Z. 2014. Modern Phylogenetic Comparative Methods and Their Application

in Evolutionary Biology: Concepts and Practice. Berlin, Springer.

Garland T., Jr, Harvey P.H., Ives A.R. 1992. Procedures for the analysis of comparative data

using phylogenetically independent contrasts. Syst. Biol., 41:18-32.

Grafen A. 1989. The phylogenetic regression. Philos. Trans. R. Soc. Lond. B Biol. Sci.,

326:119-157.

Hadfield J.D., Nakagawa S. 2010. General quantitative genetic methods for comparative

biology: phylogenies, taxonomies and multi-trait models for continuous and categorical

characters. J. Evol. Biol., 23:494-508.

Halliwell B., Holland B.R., Yates L.A. 2024. Multi-response phylogenetic mixed models:

concepts and application. bioRxiv:2022.2012.2013.520338.

Hansen T.F. 1997. Stabilizing selection and the comparative analysis of adaptation.

Evolution, 51:1341-1351.

Harmon L.J., Losos J.B., Jonathan Davies T., Gillespie R.G., Gittleman J.L., Bryan Jennings

W., Kozak K.H., McPeek M.A., Moreno-Roark F., Near T.J., Purvis A., Ricklefs R.E.,

Schluter D., Schulte Ii J.A., Seehausen O., Sidlauskas B.L., Torres-Carvajal O., Weir J.T.,

Mooers A.Ø. 2010. Early bursts of body size and shape evolution are rare in comparative

data. Evolution, 64:2385-2396.

Ho L.S.T., Ane C. 2014. A linear-time algorithm for Gaussian and non-Gaussian trait

evolution models. Syst. Biol., 63:397-408.

Iglewicz B., Hoaglin D. 1993. Volume 16: How to Detect and Handle Outliers. ASQ Quality

Press.

Ives A.R. 2022. Random errors are neither: On the interpretation of correlated data. Methods

Ecol. Evol., 13:2092-2105.

Ives A.R., Helmus M.R. 2011. Generalized linear mixed models for phylogenetic analyses of

community structure. Ecol. Monographs, 81:511-525.

Jarantow S.W., Pisors E.D., Chiu M.L. 2023. Introduction to the use of linear and nonlinear

regression analysis in quantitative biological assays. Curr. Protoc., 3:e801.

Landis M.J., Schraiber J.G. 2017. Pulsed evolution shaped modern vertebrate body sizes.

Proc Natl Acad Sci U S A, 114:13224-13229.

Landis M.J., Schraiber J.G., Liang M. 2013. Phylogenetic analysis using Levy processes:

finding jumps in the evolution of continuous traits. Syst Biol, 62:193-204.

Leys C., Ley C., Klein O., Bernard P., Licata L. 2013. Detecting outliers: Do not use

standard deviation around the mean, use absolute deviation around the median. J. Exp. Soc.

Psychol., 49:764-766.

Lynch M. 1991. Methods for the analysis of comparative data in evolutionary biology.

Evolution, 45:1065-1080.

Maddison W.P., FitzJohn R.G. 2015. The unsolved challenge to phylogenetic correlation

tests for categorical characters. Syst. Biol., 64:127-136.

Martins E.P., Hansen T.F. 1997. Phylogenies and the comparative method: A general

approach to incorporating phylogenetic information into the analysis of interspecific data.

Amer. Natur., 149:646-667.

McArdle B.H. 1988. The structural relationship: regression in biology. Can. J. Zool.,

66:2329-2339.

O'Meara B. 2016. Phylogenetic Comparative Method. In: Kliman RM editor. Encyclopedia

of Evolutionary Biology. Oxford, Academic Press, p. 254-256.

OpenAI. 2024. ChatGPT-4. https://www.openai.com/.

Osborne J.W., Waters E. 2002. Four assumptions of multiple regression that researchers

should always test. Pract. Assess. Res. Eval., 8:2.

Pagel M. 1997. Inferring evolutionary processes from phylogenies. Zool. Scr., 26:331-348.

Pagel M. 1999. Inferring the historical patterns of biological evolution. Nature, 401:877-884.

Paradis E., Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics, 35:526-528.

Pennell M.W., Eastman J.M., Slater G.J., Brown J.W., Uyeda J.C., FitzJohn R.G., Alfaro

M.E., Harmon L.J. 2014. geiger v2.0: an expanded suite of methods for fitting

macroevolutionary models to phylogenetic trees. Bioinformatics, 30:2216-2218.

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Revell L.J. 2010. Phylogenetic signal and linear regression on species data. Methods Ecol.

Evol., 1:319-329.

Revell L.J. 2024. phytools 2.0: an updated R ecosystem for phylogenetic comparative

methods (and other things). PeerJ, 12:e16505.

Rindskopf D., Shiyko M. 2010. Measures of Dispersion, Skewness and Kurtosis. In:

Peterson P, Baker E, McGaw B editors. International Encyclopedia of Education (Third

Edition). Oxford, Elsevier, p. 267-273.

Rohlf F.J. 2001. Comparative methods for the analysis of continuous variables: geometric

interpretations. Evolution, 55:2143-2160.

Royston P. 1992. Approximating the Shapiro-Wilk W-test for non-normality. Statistics and

Computing, 2:117-119.

Slater G.J., Pennell M.W. 2014. Robust regression and posterior predictive simulation

increase power to detect early bursts of trait evolution. Syst Biol, 63:293-308.

Smith E.G., Surm J.M., Macrander J., Simhi A., Amir G., Sachkova M.Y., Lewandowska M.,

Reitzel A.M., Moran Y. 2023. Micro and macroevolution of sea anemone venom phenotype.

Nat. Commun., 14:249.

Sumner S., Favreau E., Geist K., Toth A.L., Rehan S.M. 2023. Molecular patterns and

processes in evolving sociality: lessons from insects. Philos Trans R Soc Lond B Biol Sci,

378:20220076.

Uyeda J.C., Zenil-Ferguson R., Pennell M.W. 2018. Rethinking phylogenetic comparative

methods. Syst. Biol., 67:1091-1109.

Westoby M., Yates L., Holland B., Halliwell B. 2023. Phylogenetically conservative trait

correlation: Quantification and interpretation. J Ecol, 111:2105-2117.

Wilcox R.R. 2003. 3 - Summarizing data. In: Wilcox RR editor. Applying Contemporary

Statistical Techniques. Burlington, Academic Press, p. 55-91.

Zheng L., Ives A.R., Garland T., Larget B.R., Yu Y., Cao K. 2009. New multivariate tests

for phylogenetic signal and trait correlations applied to ecophysiological phenotypes of nine

Manglietia species. Funct. Ecol., 23:1059-1069.