TITLE

Genetic Diversity In Gene Regulatory Interactions Underlies The Response To Soil

Drying In The Model Grass *Brachypodium Distachyon*

SHORT TITLE

Gxe In Gene Regulation

AUTHORS

Jie Yun[1], Angela C. Burnett[2,a], Alistair Rogers[2,b], David L. Des Marais[1,c]

[1]Civil & Environmental Engineering, Massachusetts Institute of Technology, Cambridge, MA, 02139, USA

[2]Environmental & Climate Sciences Department, Brookhaven National Laboratory, Upton, NY 11973, USA

[a]Current Affiliation: Advanced Research + Invention Agency, Alan Turing Institute, British Library, 96 Euston Road, London, NW1 2DB, UK angie.burnett@aria.org.uk

[b]Current Affiliation: Climate & Ecosystem Sciences Division, Lawrence Berkeley National, Laboratory, Berkeley, CA 94720, USA alistair@lbl.gov

[c]Corresponding author: dldesmar@mit.edu

ABSTRACT

Gene expression is a quantitative trait under the control of genetic and environmental factors and their interaction, so-called GxE. Understanding the mechanisms driving GxE is fundamental for ensuring stable crop performance across environments, and for predicting the response of natural populations to climate change. Gene expression is regulated through complex molecular networks, however environmental and genotypic effects on genome-wide regulatory networks are rarely considered. In this study, we model genome-scale gene expression variation between two natural accessions of the model grass *Brachypodium distachyon* and their response to soil drying. We identified genotypic, environmental, and GxE responses in physiological, metabolic, and gene expression traits. We then identified gene regulation conservation and variation among conditions and genotypes, simplified as co-expression clusters found unique in or conserved across library types. Putative gene regulatory interactions are inferred as network edges with a graphical modelling approach, resulting in hypotheses about gene-gene interactions specific to -- or with higher affinity in -- one genotype, one environmental treatment, or in one genotype under treatment. We further find that some gene-gene interactions are conserved across conditions such differential expression is apparently transmitted to the target gene. These variably detected edges cluster together in co-expression modules, suggestive of different constraints or selection strength acting on specific pathways. We further applied our graphical modeling approach to identify putative, environmentally dependent regulatory mechanisms of leaf glucose content as an exemplar metabolite. Our study highlights an approach to identify variable features of gene regulatory networks and thereby identify key components for later genomic intervention to elucidate function or modulate environmental response. Our results also suggest possible targets of evolutionary change in gene regulatory networks associated with environmental plasticity.

INTRODUCTION

Gene expression represents a unique class of quantitative, complex traits. Like all quantitative traits, the transcript abundance of an individual gene can take a wide range of values, can show variation in those values among samples from a population, and this variation may be controlled by a number of both intrinsic and extrinsic factors [1–4]. And, because gene expression is a component of many quantitative traits such as morphology, life history, behavior, and physiology [5], using tools from quantitative genetics to study gene expression can provide insight into the molecular control of these higher-order traits [6].

Classically, variation in a quantitative trait can be modeled as a function of genetic differences among sampled individuals, of the effect of the environment on those individuals (i.e., phenotypic plasticity), and of the interaction between genotype and environment, GxE [7]. Recently, several studies have used expression QTL (eQTL) mapping to identify genetic loci that are associated with differences in the abundance of transcripts among natural accessions of plants [8–11]; these genetic variants describe heritable, constitutive differences in expression among genotypes. Abundant studies have likewise demonstrated the strong effects that differences in environment can have on the expression of single genes or suites of genes (e.g.,[12–15]).

Perhaps owing to the important role afforded phenotypic plasticity in plant ecology and evolution [16], hypotheses about its molecular basis pre-date modern genomic science [17] and remain central to studies of plant molecular biology, evolution, and biotechnology[18–24]. Recent studies assessing the prospects for biotechnology to enhance plant response to environmental stress often begin with transcriptomic, proteomic, or metabolomic assays of molecular responses to simulated stressors [25–29]. These -omics studies are predicated on the hypothesis that regulators of transcriptional activity offer attractive targets to manipulate suites of downstream molecular processes and, ultimately, organism-scale responses to stress (e.g., as described in [25,30]). However, identifying genes which can be manipulated to improve performance in the field while not also incurring a fitness penalty (so-called yield drag) remains challenging [31]. The molecular variants driving natural genetic variation in environmental response – GxE – represent a means to study how small genetic changes drive environmentally

responsive physiological and developmental traits [7]. Because such genetic variants are

naturally occurring and have likely been exposed to generations of natural selection in

wild populations, understanding their function *in vivo* may provide insight into how to

design synthetic systems to improve stress tolerance without incurring yield drag.

Genetic variants affecting GxE also represent the population-level variation on which

natural selection might act to optimize fitness in response to climate change [32] and are

thus of interest for predicting the future of natural plant communities.

GxE in plant gene expression has been reported in previous studies, most often

at the level of individual genes or at global scale treating each gene as an independent

variable [33,34]. Such an approach can be confounded by the nature of gene regulation, in

which suites of genes are often under common regulatory control [35]; as a result, treating

each gene as independent is difficult to justify statistically. Genomic features such as

promoter and enhancer elements, post-transcriptional modification, and chromatin

accessibility [36–40], add further complexity to co-variance among transcript abundance *in*

vivo. Collectively, the interaction between genes at the transcriptional level can be

conceptualized as a gene regulation network (GRN), within which there are

dependencies of expression among genes and the possible flow of information through

a cell [41,42]. The topology of GRNs implies some regulators directly regulate or are

"upstream" of many genes and may therefore exhibit pleiotropic effects when perturbed.

GRNs, and the sub-networks, modules, or communities that comprise them, may be

seen as an integrated molecular phenotype that allows for identifying transcriptional

associations with physiological, metabolic, or developmental functions, thereby

improving biological interpretability [43].

The dependency of gene co-expression and the resulting topology of GRNs on

environmental variation is poorly understood [44]. By extension, we have a limited

understanding of how environmentally responding networks evolve and might be

successfully manipulated using biotechnology to improve crop resilience. For example,

genes exhibiting GxE in their expression might be clustered into modules, as it is

hypothesized that environmental pressure results in the evolution of modules such that

organismal sub-systems can function independently [45]. A study on *Caenorhabditis*

elegans found that genes exhibiting GxE in their expression have longer promoters than

genome averages, with a higher density of regulatory motifs [46]. An extension of this

work also proposed that GxE genes might be associated with a greater number of *trans-*

acting regulators than genes showing simple genotypic differential expression [47].

Previously, we showed that the number of genes connecting to and the influence of

genes showing GxE in expression are lower under drought stress but higher under cold

stress compared to a random set of genes [48], again suggesting that the topology of

GRNs could have been shaped by – or possibly shaped the efficacy of -- past selection

in response to environmental conditions. However, that study, like most studies of

environmentally dependent gene expression, assumed that co-expression relationships

are invariant across genotypes or environmental treatments, which limits our ability to

understand the evolution of complex gene expression phenotypes [42,49].

Here, we use two inbred natural accessions of the model grass *Brachypodium*

distachyon [50] to study the physiological and transcriptomic patterns of GxE in response

to soil water deficit. *B. distachyon* is closely related to economically significant grasses

such as wheat, barley, and oat, and it has the advantages of small stature and small

genome size [50,51]. Moreover, as a natural species with no history of artificial selection, it

may exhibit higher phenotypic plasticity as compared to domesticated crops selected in

stable environments with high resource input [7]. Indeed, we previously documented GxE

in leaf water and metabolite content in *B. distachyon* response to soil drying[52]. *B.*

distachyon populations are found throughout the Mediterranean where they most likely

exhibit a winter annual strategy – germinating in the fall or spring and setting seed

before the typically hot, dry summer months[52,53]. We use highly replicated samples of

factorial combinations of genotype and environment, harnessing variation from

stochastic variance of gene expression and micro-environmental heterogeneity among

replicates to reconstruct regulatory interactions [54,55].

RESULTS

Physiological responses to soil drying

We first performed a drying experiment in which we measured a series of traits daily on

two inbred natural accessions as soil water content was experimentally decreased from

85% of field capacity to 50% of field capacity. These measurements suggest that both

accessions began reducing whole plant water usage starting on the fourth day of the

dry-down (Figure 1A, t-test adjusted by Tukey, p=0.048) and thereafter experienced a

similar magnitude of soil drying stress with most assayed traits exhibiting clear

differences between control and drying treatments by Day 6. Carbon assimilation

(Figure 1C; p= 0.0379) and stomatal conductance (Figure 1D; p= 0.0265) were

significantly lower in droughted plants starting on day 5. Total plant fresh weight started

to decrease in the droughted plants on day 6 (Figure 1E; p-value<0.0001) and exhibited

significantly lower dry weights (Figure 1F; p-value=0.0018) on day 6. Dry root:shoot

ratio is slightly higher in drought on day 7 (Figure 1G; p-value= 0.0143). Leaf hydraulic

potential is marginally more negative in droughted plants (Figure 1B; p-value=0.09) on

day 6 and is significantly more negative under drought on Day 7 (p-value<0.0001).

Relative chlorophyll content exhibits lower but non-significant values under drought

(Figure 1J; F-test, p-value=0.08). Linear mixed models show that both accessions

respond similarly along this dry-down, with the exception of leaf glucose and fructose

contents, which showed significant increase starting on Day 5 in Bd3-1 (z-test,

p<0.0001) and Day 6 in Bd21 (p<0.0001; Figure 1H and Supplementary Figure 1).

Figure 1. Comparison of physiological and metabolic traits between control and dry-
down in two accessions from Day 0 to Day 7. (Generalized) Linear models were
determined based on AIC score. Random-effects components of the models were
based on likelihood ratio tests. Differences of means shown here with asterisks, were
determined using contrasts between the two treatments at each time point in each
genotype by Tukey's test. N = 9 for each point.

Figure 2. Leaf blade traits of Bd21 and Bd3-1 on Day 6. The p-values of effect terms were determined using (generalized) linear (mixed) models, by removing each term and performing likelihood ratio test (anova and Chisq test, with maximum likelihood mode if models have random factors). DW = Dry Weight of tissue. Glc Equiv = glucose equivalents. N = 48-58 for each combination of accession and treatment.

We next performed a more highly replicated experiment (n=48-58 for each accession in each treatment) focused on contrasts between the two accessions at soil water contents of 85% and 55%. These plants constitute the sampling for all following analyses. Consistent with the trend observed before, tissue water content decreases under drought stress in both genotypes, as does the leaf photosynthesis rate (Figure 2).

Leaf metabolite content was affected by soil drying. Leaf starch content decreased on Day 6 (Figure 2G; Chisq test and p<1.6e-10). Sucrose shows no treatment effects in the leaf (Figure 2E; p=0.48). The degradation of starch and the catabolism of sucrose might contribute to the accumulation of glucose and fructose, both of which are strongly elevated in our Day 6 samples (Figure 2C, D; both p<2.2e-16). However, leaf glucose increases to a larger degree in Bd3-1 than in Bd21, shown as a significant GxE response (Figure 2C, p=0.0015). Another important group of

carbohydrates in cool-season Pooid grasses like *Brachypodium* are fructans, which may

serve as a short-term storage carbohydrate when sucrose exceeds demand [56]. Here,

we assayed high-degree polymerized fructan, which showed detectable decreases in

leaves on Day 6 (Figure 2H; $p < 1.6e-7$), similar to what we observed for starch. Low-

degree polymerized fructan also decrease slightly (Figure 2F; $p = 0.055$), similar to

sucrose. In general, most carbohydrates show a genotypic effect with higher values in

Bd3-1. Amino acids per unit leaf mass was higher in droughted plants on Day 6 (Figure

2J; $p < 2.2e-16$). Total protein content increased in Bd3-1 and marginally increased in

Bd21 (Figure 2I; $p = 0.00015$), showing a GxE response ($p = 0.034$).

Factors affecting the transcript abundance of individual genes

Figure 3. Gene expression diversity. **A)** PCA shows clustering of RNASequencing
library by accession and environmental treatment. **B)** Factors affecting the transcript
abundance of individual genes. Each point represents a gene model in the *B.
distachyon* Bd21 genome. Lfc = Log-fold change of transcript abundance in response to
soil drying. Colors indicate the significance of treatment effect on Bd21 and Bd3-1
samples; genes with a significant GxE effect are highlighted in red and two genes
discussed in the text are circled. Boxplots show two genes with large GxE effect: **C)**
Bradi3g00920 and **D)** Bradi1g65130, gene expression reaction norm based on
log2(norm_count+1). (Red Bd21d, blue Bd21c, orange Bd3-1d, purple Bd3-1c)

The transcripts of 29,740 gene models were identified in our RNASequencing data, representing 91.4% of those in the Bd21 v3.0 reference. Principle component analysis shows clear grouping of samples by library type (Figure 3A), in which the first PC separates the two genotypes and explains 15.5% of total sample variance and the second PC separates the two treatments and explains 6.4% of total variance.

26,388 genes passed our filtering criteria and are included in the following analysis of by-gene differential expression. Complete DGE gene list is available in Supplementary Data Set 1. At an FDR=0.05, 4725 genes are constitutively higher, and 4087 genes are constitutively lower in Bd21 as compared to Bd3-1 (i.e., show a significant genotypic effect). 3111 genes are constitutively higher, and 2832 genes are lower in drought than control (environmental effect); among these, 3066 genes exhibit both significant genotype and environment effects (hereafter "G+E genes"). In Bd21, 4024 and 3983 genes are up- or down-regulated, respectively, under drought. In Bd3-1, 3997 and 4010 genes are up- or down-regulated, respectively, under drought (Figure 3B). 538 genes exhibit significant GxE interaction at FDR = 0.1 (Figure 3B; hereafter "GxE genes"). Among these GxE genes, 107 are up-regulated and 66 are down-regulated in both accessions but with differing magnitudes of response in the two accessions; most such genes exhibit more dramatic changes in Bd3-1 (Figure 3B). 135 genes are uniquely up-regulated and 99 uniquely down-regulated in response to drying in Bd3-1 (so-called "conditional neutrality;"[7]); similarly, 39 genes are uniquely up - regulated and 60 uniquely down-regulated in Bd21. Only 21 genes are expressed in different directions in response to the drought treatment in the two accessions.

Of the two genes showing most striking GxE effect, Bradi3g00920 showed a 3.52 log-fold increase under drought stress in Bd21 (Wald test corrected by BH, p_{adj} = 5.55e-68) but increases only 1.05 log-fold increase in Bd3-1 (p_{adj}=3.08e-5). This leads to an additional 2.48 lfc in addition to genotypic and environmental effect, named GxE lfc (p_{adj}=1.20e-12, Figure 3C). The gene is annotated to be Leucine rich repeat family protein. Similarly, Bradi1g65130 showed a significant 4.32 log-fold decrease in Bd3-1 in response to drying (p_{adj}=2.90e-8) but only 0.84 log-fold decrease in Bd21 (p_{adj}=0.13,

Figure 3D), with a 3.48 GxE lfc. The gene is predicted to be a cyclase/dehydrase family

protein.

There are 40 annotated transcription factors among the 538 GxE genes,

including a predicted MYB-related transcription factor with the largest GxE Lfc of -1.5

among them. GO terms enriched among GxE genes include processes related to

chlorophyll, starch and organic substances, amino acids, translation, and protein folding

processes, and response to stimulus (Supplementary Data Set 2).

Variation in gene-gene interaction within co-expression modules

The high replication used in our RNA sequencing experiment (N = 48-58, subsampled

to 48 samples for each library type) affords us the power to estimate co-expression

relationships independently in each of the four library types (Bd21 control watering,

Bd21 drought, Bd3-1 control, Bd3-1 drought) using WGCNA. Genes in a module might

represent a group of genes commonly activated by a set of regulatory factors and

sharing molecular functions [43]. Depending on the parameter settings implemented in

WGCNA we identified 14-24 modules in Bd21 control, 14-18 modules in Bd21 drought,

14-16 modules in Bd3-1 control, and 15-20 modules in Bd3-1 drought (Figure 4).

We first identify modules that overlap among the four library types using a high

dimensional Fisher's exact test [57], testing the hypothesis that a set of modules identified

within or between library types share significantly more genes than expected. These

cliques of overlapping module are interrogated for four general patterns. In the first

pattern, we reason that a clique comprised of one module from each of the four library

types may represent gene co-expression relationships that are conserved across

genotype and treatment (Figure 4, teal polygons). We find 40 such modules comprising

20 overlapping cliques, among which 36 modules show significant GO enrichments.

Clique 1 is comprised of four modules that have common GO terms of circadian rhythm.

Clique 8 is comprised of a chain of seven partially overlapping cliques enriched for

similar core cellular processes including protein translation, photosynthesis, electron

transport chain, photosystem, and glycogen metabolic process (Table 1). Some

modules in conserved cliques are highly correlated with physiological traits. In

particular, Bd3-1d21 in Clique 8-6 is correlated with daily water usage (p-value=4.4e-5)

while Bd3-1d4 in Clique 8-5 is correlated with both daily water usage (p-value=1.5e-4),

and tissue water content (p-value=1.9e-4). Both modules are enriched with GO terms

associated with photosynthesis, and Bd3-1d12 in Clique 8-8 is correlated with leaf

glucose content (p-value=6.4e-5) and partially overlaps with a Clique 8-7 enriched for

glycogen metabolic process.

The second pattern is cliques comprised of two modules from the same genotype

but with different treatments (Figure 4, yellow polygons). We find three such cliques,

which may represent modules that are only coherent in one genotype. For example,

Clique 8-9 solely comprises modules in Bd21 and is enriched with functions such as

response to wounding and phenylpropanoid biosynthetic processes (including *B.*

distachyon orthologs of C4H, PAL, 4CL and F3H, F3'H and an R2R3 MYB transcription

factor which regulates phenylpropanoid biosynthesis in other species, see

Supplementary Data Set 3-4). Clique 8-12 is comprised solely of modules in Bd3-1 and

is enriched in chlorophyll and pigment metabolic process and terpenoid biosynthetic

processes; it contains five NAC regulators/domain related genes, as well as a

chlorophyll A-B binding protein and a putative senescence-inducible chloroplast stay-

green protein.

The third type of clique is treatment specific, composed of two overlapping

modules from different genotypes but the same environmental condition (Figure 4,

green polygons). We identified one such case, Clique 3. Both modules in this clique are

correlated with leaf glucose content (p-value=9.4e-8 and 1.6e-6 for Bd21d and Bd3-1d,

respectively). Although the two modules have no GO terms in common, Bd21d13 is

enriched with a GO term of response to water and contains two RAB18 orthologs

(Responding to ABA), an ortholog of rice ASR5 (abscisic acid-, stress- and ripening-

induced gene), and one IAR4. Many genes in Bd21d13 -- including the two RAB18

genes -- have very low expression in the control condition.

Finally, a clique may be comprised of modules either from a single library type or

from three library types and excluding the fourth library type; we interpret this pattern to

represent modules exhibiting GxE (Figure 4, pink or orange polygons). There are three

cliques which are unique to a single module in a single library type. Among these,

Clique 2 is comprised solely by a module identified in Bd21d and, although not enriched

313 for any GO terms, the module is highly correlated with glucose content (p-value=1.1e-

314 5). The annotations of several genes in this clique relate to abscisic acid (ABA)

315 signaling and regulation. Moreover, the proximal promoters of genes in this module are

316 enriched with a predicted bZIP motif (p-value = 2.6e-7). Clique 5 comprises only one

317 module from Bd21c and is enriched with GO term of phosphatidylethanolamine

318 biosynthesis process. Clique 7 is identified in Bd21c, enriched with GO terms of

319 glutathione metabolic process and response to chemical. We also identified three

320 cliques comprised of modules from three library types with the fourth condition

321 unrepresented. Among these, Clique 8-11 lacks a Bd3-1d module while the three

322 included modules have GO terms in common related to cell wall biogenesis. Many of

323 the genes in these modules are not included in any reconstructed module in Bd3-1d.

Figure 4. Conservation of gene co-expression modules among genotypes and treatments. Nodes represent modules identified in each of the four library types, with the size of the node corresponding to the number of genes in each module (red, Bd21d; blue, Bd21c; orange, Bd3-1d; purple, Bd3-1c). Clusters of modules enclosed by a polygon -- cliques -- show significant gene overlap using the Fisher's Exact Test(Data are available in Supplementary Data Set 5); polygon color denotes the pattern of module conservation among libraries (teal, four-module stable overlap; pink, three-module overlap; green, two-module of same environment overlap; yellow, two-module of same genotype overlap; and orange, module uniquely identified in a single library type) with the transparency of the color proportional to the number of genes in common. Underlined module names are shown as inferred network in Figure 5.

Table 1. Common GO terms at lowest level among modules in overlapping groups
(based on all module versions FDR=0.1) and correlated traits (based on moderate size
module version, in specific library type, only $p<0.001$ are reported, not corrected by
multiple test). DGE enrichment in modules are shown (based on constraint module size
version, only $p<0.001$ are reported, not corrected by multiple test, and the percentage is
shown if > 75%). Cliques with no common GO terms or correlated traits identified are
not shown. Complete lists are available in Supplementary Data Set 6-8.

Name (# of included modules)	Common GO	Correlated Trait ($p<1e-03$)	DGE Enrichment ($p<1e-03$; percentage when >75%)
Clique 1 (4)	circadian rhythm		Bd3-1c6: E
Clique 2 (1)		glucose (1.1e-05)	E (97%),GxE
Clique 3 (2)		Bd21d13: Glucose (9.4e-08) Bd3-1d8: Glucose (1.6e-06)	Bd21d13: E (100%); Bd3-1d8: E (92%),GxE
Clique 4 (3)		Bd3-1d19: leaf water content (8.5e-04)	Bd21c13: G (76%); Bd3-1c2: E,G,GxE; Bd3-1d19: G (97%)
Clique 5 (1)	phosphatidylethanolamine biosynthetic process		
Clique 7 (1)	glutathione metabolic process; alternative respiration; response to chemical		GxE
Clique 8-1 (4)	ribosomal small subunit assembly; translation; photosynthetic electron transport in photosystem II; photorespiration; ATP synthesis coupled proton transport; reductive pentose-phosphate cycle; electron transport coupled proton transport, respiratory electron transport chain		
Clique 8-2 (4)	electron transport chain, cellular nitrogen compound biosynthetic process, cellular respiration		
Clique 8-3 (4)	translation		
Clique 8-4 (4)	cytoplasmic translation; regulation of actin polymerization or depolymerization; ribosomal large subunit biogenesis, protein maturation		
Clique 8-5 (4)	Photorespiration; reductive pentose-phosphate cycle	Bd3-1d4: Leaf water content (1.5e-04) daily water usage (1.9e-04)	Bd21c2: E,G; Bd21d5: E (82%);
Clique 8-6 (4)	response to light stimulus; photosynthesis, light harvesting in photosystem I; photosystem II stabilization; pigment biosynthetic process	Bd3-1d21: daily water usage (4.4e-05)	Bd21c2: E,G; Bd21d5: E (82%); Bd3-1c3: E; Bd3-1d21: E,G,GxE
Clique 8-7 (4)	glycogen metabolic process		Bd21c8: E,G; Bd21d11: E,G,GxE; Bd3-1c5: E,G,GxE; Bd3-1d21: E,G,GxE
Clique 8-8 (4)		Bd3-1d12: Glucose (6.4e-05)	Bd21c8: E,G; Bd21d11: E,G,GxE; Bd3-1c5: E,G,GxE; Bd3-1d12: E,GxE
Clique 8-9 (2)	response to wounding; phenylpropanoid biosynthetic process; regulation of defense response; monocarboxylic acid metabolic process		Bd21c5: G,GxE; Bd21d2: E,G,GxE
Clique 8-11 (3)	plant-type primary cell wall biogenesis; plant-type secondary cell wall biogenesis; cellulose biosynthetic process		Bd21c9: E,G (100%); Bd21d6: E,G (95%); Bd3-1c10: E,G (82%)
Clique 8-12 (2)	terpenoid biosynthetic process; chlorophyll metabolic process; pigment metabolic process	Bd3-1d21: daily water usage (4.4e-05)	Bd3-1c14: E,G (76%) Bd3-1d21: E,G,GxE

Inferring regulatory interactions within modules

The preceding analyses demonstrate that the gene composition of some co-expression

modules vary as a function of genotype, treatment, and GxE, while other modules are

conserved across genotypes and treatments. We next assess a possible cause of

module-level variation: variation in regulatory interactions among genes in modules. We

use graphical models to infer the conditional independence of genes to estimate causal

relationships among them, here working with the simplifying assumption that the

regulator(s) of a focal gene is in the same module as that gene. We focus on three

exemplar modules. Since the regulation is based on estimates of transcript abundance,

gene "interaction" might represent either direct transcription factor-target regulation or

indirect regulation such as arising from through system-level phenomena such as

metabolism or cell cycle similar as described [58].

A

B

C

Figure 5. Inferred directed interactions within modules. **A)** Module 12 in Bd21d, **B)**
Module 14 in Bd3-1d and **C)** Module 10 in Bd21d inferred using the UTI-GSP method.
Nodes are genes and edges are inferred causal interactions. Edges with two arrows
indicate either the direction of interaction is uncertain or that the interaction is inferred to
be in both directions. Second row of each node shows annotation of orthologs in
Arabidopsis or rice. Glucose is included as a node in in module Bd3-1d12 as an
example of a module with high correlation between leaf glucose content and the module
eigengene (Table 1). Genes exhibiting a significant pattern of GxE in our DGE analyses
are highlighted with red outline.

373
Bd21d12 is the sole module in Clique 2, and many of its genes relate to ABA

biosynthesis and signaling. Two of these are predicted to act upstream many other

genes in the module (Figure 5A red fill): Bradi1g13760, an ortholog of AtNCED9, a 9-

cis-epoxycarotenoid dioxygenase, and Bradi3g52660, orthologous to AtCYP707A1, with

inferred ABA 8'-hydroxylase activity. The module also contains genes whose orthologs

are ABA-induced or ABA-dependent regulators (Figure 5A orange fill), including

Bradi4g39520 (orthologous to a transcription repressor, AtAITR3), Bradi3g50220

(orthologous to ABA-responsive AtATHB-7), Bradi2g15940 (orthologous to AtGBF2, an

ABA regulated bZIP transcription factor), and Bradi2g41950 and Bradi2g18510

(orthologous to AtABI1 and AtABI2, respectively). Other genes are associated with

drought response in Arabidopsis (Figure 5A blue fill), including stomatal movement

(Bradi2g20660, Bradi3g39800), wax biosynthesis (Bradi2g23740), and a late embryo

abundant (LEA) protein (Bradi1g51770). The gene with strongest GxE effect in our

dataset, Bradi3g00920 (Figure 3C), has the highest eigenvalue in this module.

Module 12 in Bd3-1d is correlated with leaf glucose content and is a member of

Clique 8-8 which share three modules with Clique 8-7 enriched for glycogen metabolic

processes (Figure 5B). Bradi1g55600 is inferred to be upstream of many other genes in

this module (Figure 5B grey fill); its Arabidopsis ortholog is an early light-induced protein

(ELIP2) and regulates chlorophyll synthesis [59]. Paralogs of Bradi1g55600 are also

present in this module. Downstream in the inferred network, there are genes whose

Arabidopsis orthologs encode enzymes in sugar and related metabolic pathways

(Figure 5B green fill), including Bradi3g33930 (pyruvate kinase), Bradi1g22050 (6-

phosphogluconolactonase in OPP pathway), Bradi5g14580 (endoglucanase in cell wall

degradation), Bradi2g56700 (galactosyltransferase in proteoglycan synthesis), and

Bradi1g18650 (4-alpha-glucanotransferase in starch degradation). Above, we reported

that leaf glucose content is correlated with this module; when we include glucose as a

node in the module, we find that it is predicted to directly interact with Bradi1g22050, the

6-phosphogluconolactonase. Bd3-1d12 also contains Bradi4g15010, whose Arabidopsis

ortholog is predicted to have starch synthesizing and degrading activity (LESV).

Additional genes in this module have predicted functions in secondary metabolism

(Figure 5B purple fill), including orthologs of an Arabidopsis MYB family transcription

factor involved in flavonol biosynthesis (Bradi1g64687), flavonoid 3'-hydroxylase

(Bradi3g04750), flavanone 2-hydroxylase (Bradi2g48450) and a 4-coumarate: CoA

ligase in the phenylpropanoid pathway (Bradi3g52350). Notably, several GxE genes

with unknown functions are connected in this module.

Finally, we report on module Bd21d 10 in Clique 1, that is conserved in all four

library types and is enriched in genes relate to circadian processes (Figure 5C, yellow

fill). There are five predicted MYB TFs in this module, including Bradi3g51757

(orthologous to AtRVE6 and OsRVE4), Bradi2g62067 (orthologous to OsLUX),

Bradi5g19380 (orthologous to OsMYB511, AtRVE7), and Bradi3g16515 (orthologous to

OsLHY/CCA1), Bradi1g29680 (orthologous to OsLHY in rice and AtRVE2/ ERP1 in

Arabidopsis). Additional genes in this module are orthologous to CONSTANS (CO) and

related proteins in Arabidopsis. Studies suggest genes in circadian-controlled pathways

show conserved oscillation over 24 hour intervals [60], suggesting that stochastic variance

among individuals may have provided the signal for reconstructing these conserved

interactions in our data.

Genes predicted to interact show similar or different patterns of differential gene

expression

Genes in modules are enriched with certain types of DGEs as shown in Table 1.

However, within a module, gene expression variation between two genes connected by

an edge in our graphs (Figure 5) can be dependent on: genotype (G change, e.g.,

regulator and target gene show no DGE and G DGE respectively, see Supplementary

File 1 for testing method and illustrative examples), environment (E change, e.g.,

regulator and target gene show no DGE and E DGE respectively), genotype by

environment (GxE change, e.g., regulator and target gene show no DGE and GxE DGE

respectively) or no effect (e.g., regulator gene and target gene both show no DGE). We

summarize the different types in the inferred gene regulation network in modules in

Supplementary Table 1 and Supplementary Figure 2 (based on modules in Figure 5).

For example, in module Bd21d12 that only exists in Bd21 under drought condition, of 41

total edges, there are two edges showing no effect of expression changes, five edges

showing G changes, 11 edges showing E changes, 12 edges showing G+E changes,

11 edges showing GxE changes. While in Bd21 specific module Bd21c5, of 62 total

edges, there are 23 edges showing no DGE changes, 25 edges showing G changes,

nine edges showing E changes, eight edges showing G+E changes, and no edges

showing GxE changes.

Genetic and Environmental Effects on Interactions among Genes

An edge between nodes in our inferred gene regulation network represents a

hypothesis about the regulatory interaction between two genes. The high biological

replication of our RNASequencing across four library types enables us to fit linear

models to explore the effects of genotype, environment, and their interaction on the

expression relationship between pairs of genes. There are four patterns of interest in

our dataset.

First, the regulatory interaction between two genes may be present under one or

some library types (combinations of genotype and treatment) but absent in others; this

pattern would be diagnosed by identifying significant slope change such that the

correlation between these two genes is no longer significant. One possible explanation

for this pattern is environmentally dependent regulatory interaction (Figure 6B). In ABA

module Bd21d12, described above, the regression of Bradi4g40850 (unknown) on

upstream genes shows a significant E effect of slope from gene Bradi3g50220 (ATHB-7

which is regulated in an ABA-dependent manner). Similarly, in drought specific module

Bd21d13, the regression of Bradi1g37410 (which encodes a dehydrin family protein) on

upstream genes suggests that Bradi1g44480 (dehydrogenase E1 component domain

containing protein) regulates Bradi1g37410 only in drought (Supplementary Figure 3A).

Across modules, we see clustering of such context-dependent edges as reflected in

module Bd21d13 in clique 3 – reconstructed only under drought conditions – containing

32 edges of which 14 are environmentally dependent (44%) (Supplementary Table 1).

A regulatory interaction between two genes also may be observed in only one of

the two genotypes (Figure 6D). For example, in Bd21 specific module Bd21c5, the

regression of Bradi1g15840 (putative allene oxide cyclase in jasmonic acid synthesis)

on upstream genes suggest Bradi3g44990 (putative cell-wall invertase) regulates

Bradi1g15840 only in Bd21 (Figure 6D). Consistent with the observation that in Clique

8-9 specific to Bd21, we find that Bd21c5 has 12 edges showing a significant effect of

genotype among its among 65 edges (18%) (Supplementary Table 1).

We might also observe differences among the slopes of the library types such

that a regression is only significant in a single library type, indicative of GxE for a given

pairwise regulatory interaction (Figure 6A). In module Bd21d12 discussed above, the

regression of Bradi3g00920 (Leucine Rich Repeat family protein) on upstream genes

suggests that Bradi2g15940 (GBF2) interacts with Bradi3g00920 with a significant GxE

effect of the slope (Figure 6A). Module Bd21d12 is found only in Bd21 under drought

stress and has many genes associated with ABA signaling.

In the second general pattern, the interaction between two genes is observed to

change in magnitude between library types; this might be caused if the binding

interaction between a transcription factor and a regulatory motif is stronger in one

genotype than a second genotype (Figure 6E). Such an interaction would appear as a

change in the slope of the regression of target gene on its putative regulator but without

a complete loss of the correlation. As one example, Bradi4g20520 (unknown function) is

predicted to regulate Bradi3g19730 (jacalin-like lectin domain containing protein) in

Bd21c6 in all library types but with a greater slope in the Bd21 libraries than in Bd3-1

libraries.

Third, the baseline expression of genes might be different across library types

(Figure 6G), possibly due to the presence of cofactors or the general resource and

energy availability of the cell; this type of regulation may be detected as a difference in

intercept among library types in our multiple regression analysis. In module Bd21c10,

Bradi3g45230 (RNA-metabolising metallo-beta-lactamase) regression on genes

including Bradi2g10880 (LNK3) shows the regulation is similar in all library types, but

the baseline in Bd3-1 is constitutively higher in both treatments. This pattern is observed

as a higher intercept in Bd3-1 libraries which leads to genetically dependent expression

of the downstream gene. Such intercept differences as a function of environment,

genotype, or GxE are widely found among the pairwise interactions identified in our

data.

The final pattern is no difference in the pairwise regulation between genes

among library types but an apparent transmission of a regulatory response between the

genes. This pattern is manifested as neither a significant slope change nor an intercept

effect in the regression, and a common differential expression between putative

regulator and its target gene. One such case is Bradi5g18277 (serine-type

endopeptidase) and Bradi1g53680 (zinc transporter 3 precursor) in module Bd31c16

(Figure 6F). In module Bd31d12, GxE genes are found more likely to share a regulatory

interaction with other genes exhibiting GxE than non GxE genes in the module (t-test, p-

value = 0.013; see Supplementary Figure 3B for an example).

Finally, we use our approach to study the transcriptional control of leaf glucose

content, which shows a significant GxE effect in our assays (Figure 1H) and is highly

correlated with four module eigengenes (Bd21d12, Bd21d13, Bd3-1d8, and Bd3-1d12;

Table 1). Nine genes are inferred to regulate glucose when it is included as a node in

these modules. Among these, we find that Bradi4g39520 only exhibits an interaction

with glucose under drought stress (Figure 6C); the glucose content is same between

two genotypes and not dependent on the expression level of Bradi4g39520 under

control conditions. Bradi4g39520 is an ortholog of AITR3 in Arabidopsis, ABA-induced

transcription repressor that acts as feedback regulator in ABA signaling.

Figure 6. Predicted and actual examples of regulations between pairs of genes selected. **A-B, D-G)** show regulation slope changes, transmission, and regulation intercept changes. These are illustrated with linear regression between regulator and target, the expression level of each gene, and allele specific expression (ASE) types. ASE types include constitutive regulation (-) and environment specific (xE) detected in full model (see method), or regulation detected in control (c) and drought (d) condition. P-value <0.1 and not corrected by multiple tests. **C)** regression of log-transformed leaf glucose content on upstream identified gene Bradi4g39520 as a putative regulator of leaf glucose content. Transcript abundances are log2 (norm_count+1).

Identifying allele-specific expression in target genes

We next used allele specific expression assays to infer whether changes in regulatory

interactions attributed to genetic or GxE effects in transcript abundance arise from

factors acting in *cis* or in *trans*. In the above examples of the genotype-dependent slope

between the regulation from Bradi3g44990 to Bradi1g15840 (Figure 6D), Bradi1g15840

is found to be *cis*-regulated (Supplementary Figure 4B). In the example of GxE

regulation of Bradi3g00920 by Bradi2g15940 (Figure 6A), we identified Bradi3g00920 is

regulated by a *cis*- acting factor under drought stress (Supplementary Figure 4A).

Moreover, in the example of an apparent transmission from Bradi5g18277 to

Bradi1g53680 (Figure 6F), we observed that transcript variation of Bradi1g53680 is

controlled by a *trans*- acting variant (Supplementary Figure 4C) as expected. As for the

example of Bradi3g45230 being regulated by Bradi2g10880 with an effect of genotype

on the intercept of the regression (Figure 6G), Bradi3g45230 is found to be *cis*-

regulated (Supplementary Figure 4D). It might be an example of variation in an

enhancer or similar cis-regulatory element leading to base level expression difference.

DISCUSSION

Gene expression has long been a focus of both evolutionary and functional genetic

studies. Here, we aimed to integrate these two perspectives by studying the

environmental and genetic control of regulatory interactions between genes using highly

replicated RNASequencing libraries of *Brachypodium distachyon* accessions exposed

to moderate soil drying and control conditions.

Here, we extend past work showing that the expression of individual genes can

vary as a function of genotype and environmental treatment to demonstrate variation in

interactions between genes. One example highlighted throughout the analyses in this

study is Bradi3g00920, which shows a pattern of GxE in its gene expression. By

developing a directed graphical network for the module in which the gene is located, we

infer that Bradi3g00920 is relatively downstream of several other genes putatively

regulating it. Multiple regression of its gene expression suggests that Bradi3g00920 is

regulated by Bradi2g19540 only under drought in Bd21. We infer that this pattern

suggests a possible cause of the observed GxE in gene expression for Bradi3g00920.

Further, allele-specific expression reveals that this gene is associated with a *cis*-
regulated expression pattern.

We next consider how a differentially expressed gene may transmit its
expression pattern through its interactions with other genes. Here, we refer to the
correlated pattern of differential expression as "transmission." As shown in the example
in Figure 6F, an inferred target gene (Bradi1g53680) shows the same dependency on a
putative regulator (Bradi5g18277) across all library types. Indeed, the target gene is
found to be *trans-* regulated, suggesting that the genetic variant(s) causing the
differential expression is in the direct regulator or upstream of it. It suggests the
interpretation of *trans-* acting loci in QTL studies might include what we here call gene
expression transmission and not a protein coding change in the trans-acting regulator.
The transmission of the expression is also a direct example showing certain genetic
variants can affect more than one genes, and that the expression of individual genes
should not be considered independent variables.

Variation in gene-gene interactions observed in modules could lead to a module
not being reconstructed in certain combinations of genotype and environment. For
example, 44% of the inferred edges in module Bd21d13 exhibit environmentally
dependent regulation and, perhaps as a result, this module is found only under drought
stress. In contrast, in modules that have fewer pairwise gene-gene regulatory variations,
or when gene expression states are transmitted faithfully from upstream nodes in the
network, these modules may be more likely observed under all sampled conditions.

Genetic variants affecting expression, whether constitutive differences between
varieties (in our study, those with a significant G effect) or GxE, may be acted on by
selection if they ultimately have an effect on phenotype. Past work addressed the
possibility of spatial separation across a network between a genetic variant and its
effect[35,61], or between the perception of an environmental stimulus and the downstream
transcriptional response[62]. The network context of a genetic variant is important for
evolutionary studies because the number and functions of genes which interact with a
variable gene influences the pleiotropy of that genetic variant[63]. Recent work considered
the topology and/or pleiotropy of gene expression in the context of the evolution of
plasticity[48,64]; however, most studies do not account for the prospect of pleiotropic

effects of these genetic variants and, as a result, variation in the network topology itself.

We specifically consider such regulatory interaction variation here. The large

RNASequencing dataset reported here also allowed us to estimate gene co-expression

independently in each combination of genotype and environmental treatment, thus

relaxing the assumption of an invariant network topology across conditions used

previously[48,64,65]. We note that other approaches to account for network structure, such

as mapping covariance or principle components as traits in QTL analysis [3,11,66] may

detect similar regulatory structure.

Changes in gene-gene interactions are found to be clustered in some modules,

suggesting a spatial pattern of regulatory variation in networks and supporting the

hypothesis that modularity allows for reduced pleiotropic effects of mutations[63]. A

related hypothesis is that certain regulatory pathways are under weaker purifying

selection, thus leading to accumulation genetic of variation in a subset of modules.

Consistent with this latter hypothesis, we observed that variable interactions are more

common in modules not associated with essential biological pathways (e.g., translation

and ATP synthesis processes). The small overlap between variable modules and the

chain structure in Figure 4 is representative of these patterns. Previously, we observed

a similar pattern in which individual genes exhibiting expression GxE were localized at

the periphery of a gene co-expression network under drought[48]. The pattern we observe

here is similar, but now at the level of gene-gene regulatory interactions.

While we have highlighted a number of cases where the interactions among

genes vary as a function of genotype, environment, and GxE, we do observe several

modules that, while conserved among library types, are enriched for genes that respond

to stress. Clique 8-6, for example (Figure 4), is comprised of one module from each of

the four library types but is significantly enriched for genes that individually show a

significant response to soil drying stress in our study. These gene lists are also enriched

for GO terms associated with the light reactions of photosynthesis. Thus, while the gene

interaction network of the pathway forms a system whose interactions are constant

across various conditions and genotypes, this system is nonetheless environmentally

responsive. The mechanism of this response -- whether via active regulation or more

systems-level metabolic differences -- cannot be determined with this type of data but

we highlight this difference from the environmentally dependent gene-gene interactions

that we observe in, e.g., Clique 3 (Figure 4).

Our study also revealed a functional connection between cellular and whole-plant

traits to specific gene regulatory features. First, modules associated with traits or

enriched with GO terms represent hypotheses about the transcriptional control of key

drought responses. However, even in modules representing housekeeping pathways,

fewer than 50% of genes have annotations; our network-based approach therefore

represents to discover gene function, particularly for genes in the overlapping

intersection between modules. Second, certain traits, like leaf glucose content, might be

controlled by several pathways, as we found to be associated with four modules (Table

1). We also show how these associations may change in different environments as

some modules only exist under drought stress.

Collectively, our work highlights how natural selection may modify gene

regulation networks to have new functions or environmental responses and may also

shed light on how we can modify traits through breeding. First, a gene's expression can

be modified at various steps of its regulation, e.g., at its direct regulator or further

upstream. Conversely, modification of a gene's expression to be responsive to an

environmental factor will likely require accounting for gene-gene interactions; our study

shows one way to identify such targets. Second, regulatory variation or environmentally

responsive features seem to accumulate in specific modules. Variants whose effects

are restricted to such modules may have fewer pleiotropic effects and thereby be more

likely to be retained in populations. Third, it might be possible to change plasticity to

certain stressors without having pleiotropic effects in other environments; modules with

many GxE edges but few G edges might be fruitful targets to such avoid trade-offs. On

the other hand, targeting modifications to modules with E edges but few GxE edges

may prevent rank-changing interactions when breeding for a range of environments[19,20].

While our results suggest how natural or artificial selection may shape, or be shaped by,

physiological and regulatory responses to the environment, how these factors directly

contribute to drought tolerance and lifetime fitness cannot be determined in the chamber

growth experiments like those performed here. Future studies need to build the

connection from the current transient gene expression and regulation characterization to
fitness in the natural environment.
648
METHODS
Experimental growth conditions and drought treatments
The experiment protocol was adapted from a study by Des Marais et al.[67] Two
accessions of *Brachypodium distachyon* (Bd21, Bd3-1) seeds were placed at 4 °C for
seven days to synchronize seedling germination. After that, the seeds were planted in
Deepot D16H pots (Stuewe&Sons) filled with Profile porous ceramic rooting media
(Profile Products, Buffalo Grove, IL). The dry weight of each pot (DW) and its field
capacity for water (FC) were recorded. FC was determined by saturating a pot with
water and allowing it to drain under gravity overnight and calculating the weight
difference from DW; this was used as the basis for the subsequent controlled soil dry
down. Plants were grown in growth chambers with 50% humidity, 500 μmol photons m^{-2}
s^{-1} light intensity, and 25°C day/20°C night (12h/12h) and bottom watered with tap water
(pH=5-6) every other day throughout the experiment, supplemented with DYNA-GRO
GROW Liquid Plant Food 7-9-5 (Dyna-Gro, Richmond, CA, USA) over the first 32 days.
On day 33 after sowing, control plants were watered to 85% FC every night while dry-
down plants were allowed to decrease their soil water content to 50% FC over 7 days
(5% decrease each day). The plants were divided into two groups with Group 1 used to
characterize plant responses along the dry-down process, and Group 2 used for gene
expression and metabolite content.
On days 0 (85%FC), 2 (75%FC), 4 (65% FC), 5 (60% FC), 6 (55% FC), and 7
(50% FC) nine individuals of each genotype from each treatment (control and drying)
were used for gas exchange measurements at 3–5 hours after lights-on, non-
photochemical quenching (NPQ) at 5-6 hours after lights-on, and destructive sampling
for leaf hydraulic potential, leaf metabolites, and total plant biomass at 6-8 hours after
lights-on. Our preliminary analyses determined that gas exchange and leaf water
potentials of droughted plants began to deviate from water controls on days 5 and 6 of
the dry-down; accordingly, we selected day 6 for more detailed analysis. 48-58
replicates of leaf tissue in each genotype and treatment were measured for gas

exchange and photosynthesis at 1.5-5 hours after lights-on on day 6 and harvested for

RNA extraction and metabolites assay between 6-7.5 hours after lights-on (equivalent to

mid-day). Sampling of individuals was randomized to avoid confounding time-of-day

harvest with genotype or treatment.

Physiological and metabolic measurements

Net CO_2 assimilation rate (A), and stomatal conductance (g_{sw}) of the youngest fully

expanded leaf of a selected tiller were determined using a Li-6800 (LI-COR

Biosciences, Lincoln, NE). The LiCor chamber was set to a flow speed of 200 µmol s^{-1},

humidity at 50%, CO_2 concentration at 400 µmol mol^{-1}, fan speed of 10,000 rpm, and

photosynthetic photon flux density (PPFD) at 1000 µmol (m^{-2} s^{-1}). Leaves were left for 5

min to stabilize in the chamber before each reading. Photochemical quenching (qP),

non-photo-chemical quenching (qN), and Nonphotochemical chlorophyll fluorescence

quenching (NPQ), PSII maximum efficiency of light adapted leaves (fv'/fm') were

measured on leaves at noon; Parameters follow Li-6800 protocol of Determining PSII

efficiency. Leaf chlorophyll content was estimated by taking the average of two leaves

measured with MultispeQ V2.0 prior to lights-on in the growth chamber.

The leaf water potential was measured by the Scholander Pressure Chamber

Model 670 (PMS Instrument Company, Albany, OR) on the youngest fully expanded

leaf of sampled plants.

Above-ground tissue was excised at soil level with a razor blade. Roots were

cleaned with tap water to gently remove growth media and dried with tissue paper.

Fresh tissue biomass was recorded before samples were dried for 48 h at 80°C and

then weighed for dry biomass.

Statistical analyses

For the traits collected during dry-down (Supplementary Data Set 9), most traits were

fitted into linear mixed models using lm and lmer [68] in R[69]. Glucose, fructose were fitted

with generalized linear mixed models with Gamma distribution in glm and glmer [68], with

family=Gamma(link= "inverse"). Accession (two levels), Treatment (two levels), Harvest

day (six levels), and their interactions, were modeled as fixed factors and freeze dry

session (four levels) and assay batches (five levels) for metabolites were modeled as
random factors. Model selection was first performed on the fixed-effect parts of each
model, based on maximum likelihood Akaike information criterion (AIC) using the
dredge function in the MuMin package [70]. Random-effects factors of each model are
based on maximum likelihood of models using the anova function. Models are shown in
Supplementary Table 2. Throughout the dry-down process, the pairwise significance
test between treatments for each accession on each harvesting day (Figure 1 and
Supplementary Figure 1) was estimated with a t-test (or Z-test for glm, glmer) adjusted
by Tukey using emmeans function [71] in R based on the model selected. Outliers were
removed before model fitting. The results are visualized using ggplot2 and ggpubr[72,73].

Similarly, leaf trait data collected on Day 6 (Supplementary Data Set 10) were fit
with (generalized) linear mixed models using lm, lmer, glm, or glmer, as appropriate,
with fixed factors of Accession, Treatment, and their interaction, and random factors of
growth batch, freeze dry sessions, and assay batch for metabolites (Figure 2). The p-
values shown are based on likelihood ratio test (Chisq test) of dropping each term using
anova function.

725 Transcriptome sampling and processing
Leaf tissue flash frozen on liquid nitrogen was ground using an MP FastPrep-24™
Classic bead beating grinder with CoolPrep™ 24 x 2mL Adapter (MP Biomedicals)
loaded with dry ice. Approximately 100mg of tissue was used for RNA extraction using
Spectrum Plant Total RNA Kit (Sigma-Aldrich, USA) according to the manufacturer
instructions, followed by DNase treatment (On-Column DNase I Digestion Set, Sigma-
Aldrich, USA). The quality and quantity of the RNA samples were determined both by
absorbance measurements (Nanodrop 1000, NanoDrop Technologies) and with a Tape
Station (4200 TapeStation System, Agilent). 25ng of RNA were processed for High-
Throughput 3' Digital Gene Expression [74,75] by the MIT BioMicro Center, pooled and
then sequenced across three flowcells on an Illumina Novaseq (NovaSeq6000
Sequencing System, Illumina) generating, on average, 3.9±1.3 million raw reads per
sample.

Raw sequence reads were trimmed by Trimmomatic [76] and Cutadapt [77] to
remove adapters and poly(A) tails. All reads were mapped to the *B. distachyon* Bd21
primary transcriptome sequence
(Bdistachyon_556_v3.2.transcript_primaryTranscriptOnly, downloaded from
Phytozome) and the annotated sequence of the whole Bd21 plastome, which is
available at Genbank (LT558597 accession). Read alignments from Bowtie2 [78] were
corrected by RNA-Seq by Expectation-Maximization (RSEM) [79] to account for non-
uniquely aligned reads. Bowtie2 alignment is done with seedlength -L 25 and mismatch
allowance -n 2 which allows mismatches of Bd3-1 read alignments. Alignment with --
estimate-rspd specifies the reads position distributions that are highly 5' or 3'- biased [79].
Reads with the same unique molecular identifier and aligned to the same genes were
removed to reduce PCR biases during library generation [74,75] prior to adjustment in
RSEM. RNASeq reads from Bd21 and Bd3-1 mapped to the Bd21 references at
77.0±1.7% and 78.1±1.8 %, respectively, suggesting no genotypic effect on mapping.
Reads which did not map to the genome during this first round were mapped onto
ribosomal RNA, after which the total mapping rates increased to ~96%.
754
Differential gene expression (DGE)
RNASeq reads were normalized using Relative Log Expression normalization
implemented by DESeq2 [80]; these normalized data with $\log_2$ (Count+1) transformation
are used for downstream analyses. The grouping is visualized with PCA using
factoextra package in R.[81] In DGE step, genes with >0 counts in more than 10 samples
are kept. DESeq2 fit the transcript count of each gene into the generalized linear model
Transcript Count ~ Genotype + Treatment (+ Genotype * Treatment) or Transcript
Count ~ Treatment, stratified by genotype or Transcript Count ~ genotype, stratified by
treatment. Each gene is fitted into a negative binomial distribution with mean μ and
variance α. Results in Figure 3 were visualized with ggplot in R. [72,73]
765
Identifying patterns of variation in clusters of gene-gene interactions
For this and following analyses, we removed transcripts exhibiting low variance (<0.1)
among replicates in each library type; in total, 16,207 genes are included. Co-

expression modules in each library type (Bd21 Control, Bd21 Drought, Bd3-1 Control,

Bd3-1 Drought) were generated using Weighted Gene Co-expression Network Analysis

(WGCNA)[82]. For WGCNA, the starting correlation matrix is unsigned, as both negative

and positive correlations are of interest in this study. Weight power=5 is added to the

correlation matrix to achieve a scale-free adjacency matrix. The topology similarity

matrices in the Bd21c, Bd3-1d, and Bd3-1c data sets are scaled to have a similar

distribution to Bd21d, which we chose arbitrarily as the reference library type.

Dendrograms were generated from topology similarity matrices, and modules were

generated by cutting at three parameter settings: (1) moderate size modules, used for

most analysis below, high dissimilarity (height= NULL(0.99) with a dynamic hybrid

algorithm (minClusterSize =30, deepsplit=2, pamStage = FALSE, mergeheight=0.25);

(2) conservative (smaller) modules, used for causal inference analyses discussed

below, cutHeight = 0.984, minClusterSize = 15, deepSplit = 3, pamStage = FALSE,

mergeheight=0.25; and (3) liberal (larger) modules, cutHeight = NULL(0.99),

minClusterSize = 30, deepSplit = 1, pamStage = FALSE, mergeheight=0.25. The three

different settings are used to better understand the effect of threshold choice within

each library type. In brief, our WGCN analyses recovered 15, 24, and 14 modules in

Bd21c in the three settings, respectively; 14, 18, and 14 modules in Bd21d in the three

settings, respectively; 14, 16, and 14 modules in Bd3-1c in three settings, respectively;

and 15, 20, and 15 modules in Bd3-1d in the three settings, respectively. Gene

membership is summarized in Supplementary Data Set 9.

We next addressed metrics of module preservation across our four library types,

Bd21c, Bd21d, Bd3-1c, and Bd3-1d. First, we assessed the degree to which the lists of

genes in a given module are conserved among our two genotypes and two

environmental conditions using a test for membership overlap. We required that module

overlap must be significant in all three parameter settings (moderate, conservative, and

liberal, described above, as shown in Figure 4, three modules with unstable overlapping

are not shown). To examine the overlap in gene module membership among library

types we used the SuperExact test [57] and a p-value cut-off of with $< 10^{-12}$ with the

alternative hypothesis that the size intersect between two groups or among more

groups is larger than expected by chance (Supplementary Data Set 5). Hypergraphs

representing module overlap networks -- here, termed "cliques" for subnetworks -- were

visualized using the R package igraph [83,84].

Second, we tested for enrichment of transcripts exhibiting significant DGE in our

co-expression modules using Fisher's Exact Tests [57]. Significant enrichments (p-value

<0.001 uncorrected by test number) are reported in Supplementary Data Set 6 and

summarized in Table 1. Here the enrichment is reported for modules estimated from our

conservative parameter settings to be consistent with following network inference

analysis.

Third, biological roles of each module were inferred using GO term enrichment,

by inspecting gene annotations, and by testing for module correlation with the sampled

physiological and metabolic traits. Module GO enrichment is conducted on using the

PANTHER classification[85] and implemented in rbioapi[86], with Fisher's exact statistical

overrepresentation tests of GO biological process. Only the significantly enriched GO

terms at FDR = 0.1 and the lowest-rank GO categories are reported. Here the

enrichment is reported if positive in modules of any of the parameter settings

(Supplementary Data Set 7). For common GO term comparison among modules in an

overlapping clique, all GO categories are considered (summarized in Table 1). Common

genes and annotations in overlapping cliques are list in Supplementary Data Set 3-4.

Gene annotations are from MapMan (https://mapman.gabipd.org/mapmanstore) based

on the Phytozome database. Correlations between module eigengenes (i.e., their first

principal component) and traits was performed in each library type. The correlation p-

values are reported in Supplementary Data Set 11 and summarized in Table 1 based on

moderate criteria modules, uncorrected for the number of tests.

Finally, we tested the hypothesis that genes within modules share common DNA

motifs corresponding to known transcription factor binding sites. All possible 6-mer

sequences were tested for enrichment in the -500 to +200 upstream window regions[87]

relative to each gene's predicted transcriptional start site in the Bd21 reference genome,

in the module using AME in MEME suite[88]. The AME parameters were: --verbose 1 --

scoring avg --method fisher --hit-lo-fraction 0.25 --evalue-report-threshold 10.0, with the

control set based on all the genes. The Arabidopsis DAPSeq database [89] was used as a

reference to interpret enriched motif sequence identified. Here the enrichment is

reported for moderate criteria modules.

Causal network inference in modules

Methods such as WGCNA, based on correlations in transcript abundance, cannot

provide information about the direction of regulatory interactions. Our high replication

provides the power to model genes as random variables and infer conditional

independence of gene expression to represent possible causality. Here, we use the

Unknown-Targeted Interventional Greedy Sparest Permutation (UT-IGSP) algorithm [90]

to infer regulatory causality, representing a group of genes as a directed acyclic graph

(DAG). We use all genes in a given WGCNA module – estimated from one library type -

- as priors to reduce latent variables in the model, relying on the simplifying assumption

that there are no edges among genes between modules. We include both samples that

the module is inferred from (here, representing "observational" data) as well as samples

from all other library types (here, representing "interventional data"), representing the

assumption that most of the gene regulation structure is the same between library

types. In these analyses, we use the conservative module criteria (most restrictive

inclusion in the WGCNA analyses) and limit the analysis to modules with number of

genes fewer than 70% of samples in the specific library type. When testing for modules

that are conserved within a treatment or within a genotype, modules with as many as 70

genes can be used because more samples are available for inference. We use stability

selection [91] as implemented in Belyaeva et al [92] to minimize the effect of our selected

hyperparameters. Edges with a selection probability higher than upper 0.95 (or 0.97 for

bigger conserved modules to keep sparse edges, and 0.95 for modules with glucose as

a node) quantile are kept. DAGs were exported from R using the rgraphviz and

visualized using Graphviz Visual Editor [93].

Identifying patterns of variation in gene-gene interactions

We extended our analysis of gene expression responses using a regression approach

that explicitly models the magnitude and direction of putative interaction between genes

based on their transcript abundances. For a putative regulator and target pair identified

in our DAGs, we regress the target gene transcript abundance on the regulator

transcript abundance, and test whether the slope and intercept coefficients of

regression changes as a function of genotype or environmental treatment. When there

are multiple inferred regulators for a target gene, a multivariate linear regression is

applied.

For the hypothesis that gene x_i $(i = 1, \ldots, n)$ regulates gene y, to infer the

intercept or slope differences between genotypes, treatment, or their interactions, we

modeled their (log$_2$ -transformed) gene expression x_i and y as:

869
$$y = \gamma_0 + \gamma_g g + \gamma_e e + \gamma_{ge} g * e + \sum_{i=1}^{n} (\beta_{0,i} + \beta_{g,i} g + \beta_{e,i} e + \beta_{ge,i} g * e) * x_i$$

Here g, e are used to differentiate the dataset used to train the parameters (i.e., g, e are

"0,0", "0,1", "1,0", "1,1" for Bd21 control, Bd21 drought, Bd3-1 control and Bd3-1 drought

respectively). Thus, consistent with the DGE test using DESeq2, above, a significant p-

value of γ_g, β_g suggest a genotypic effect on regression intercept and slopes,

γ_e, β_e suggest an environmental effect on intercept and slopes, and γ_{ge}, β_{ge} suggest a

GxE effect of intercept and slopes. Only genes with regression R^2 greater than >0.5 are

considered, as estimated with the Car package in R. [94] P-value of <0.05 is reported to

be significant without multiple test correction. Note that if the transcript abundances are

standardized before fitting our regression models, the slope differentiation test should

closely match the results of CILP tests introduce by Lea et al[66] which we include as

Supplementary Figure 5.

Allele specific expression

We used allele-specific expression to assess whether GxE in transcript abundances

were caused by cis- or trans-acting variants. Bd21 served as a pollen donor on

emasculated Bd3-1 flowers, resulting in ten F1 hybrid seeds. To confirm successful

outcrossing in this largely selfing species, we extracted DNA from leaves (DNeasy Plant

Pro Kit; Qiagen, MD, US) and tested for a known length polymorphism on a 3% agarose

gel (data not shown). We performed a dry-down experiment on these F1s and extracted

leaf RNA, as above. 100ng of RNA for each of 10 samples along with 12 samples (three

replicated of each genotype and condition) from the initial experiment were aliquoted for

sequencing. For DGE, and to identify SNPs between alleles, we generated libraries

using NEB Ultra II Directional RNA with Poly(A) Selection (New England Biolabs,

Ipswich, MA). Libraries were sequenced on an Illumina Novaseq generating 34.1 ± 13.7

million raw reads per sample.

To generate a reference diploid transcriptome, RNAseq reads of each Bd3-1

sample were aligned to the Bd21 reference genome (Bdistachyon_556_v3.2) based on

a RNAseq short variant discovery pipeline [95], with STAR alignment [96] of two pass and

alignSJoverhangMin=30 and alignSJDBoverhangMin=10, variants calling GATK [97] and

the flag of dont-use-soft-clipped-bases. The variants called in each sample were then

filtered to find SNPs and in/dels with homozygous alternative genotype mapping. To

reduce false discovery only variants identified in more than half of the replicates of each

genotype were kept. The synthesized Bd21 or Bd3-1 vcf together with Bd21 genome

and annotation gff was then used to generate *de novo* Bd21 or Bd3-1 genome (only

exons are corrected) using g2gtools [98], and a diploid genome with emase [99]. The reads

from all the samples were mapped to this diploid transcriptome using bowtie[100] and a

count matrix was estimated from alignments using emase (an alternate pipeline

employing bowtie2 followed by RSEM provides similar results; not shown) to account for

the multi-mapping particularly accounting for allelic expression. Isoforms are not

differentiated in this study.

Genes with different counts between two alleles are considered to have

sequence variants but sparse differences in a gene might not accurately differentiate

two alleles. Accordingly, we included 3963 genes that have more than >1 count of the

allele in one accession and <20% total count of the allele in the opposing accession in

all samples, modified from Gould et al [101]. The RNASeq count matrix was normalized

using RLE in DESeq2, as above. All genes (>0 in more than 10 samples, including

allele-indifferentiable genes) are included in the normalization step with allele-

distinguishable genes shown as allelic expression and other genes as total expression

to ensure the normalization factors are calculated based on most genes for the

comparable of genes with allelic variants. Possible read-mapping bias was assessed as

in Lovell et al. [102] shown in Supplementary Figure 6. PCA of samples are shown in

Supplementary Figure 7.

We employed a pipeline based on DESeq2 as introduced by Lovell et al [102].

Briefly, the transcript abundance of a target gene is fit into full model

$log_2 q_{ij} = \beta_0 + \beta_T T_i + \beta_A A_i + \beta_G G_i + \beta_{AG} A_i * G_i + \beta_{TG} T_i * G_i + \beta_{AT} T_i * A_i + \beta_{AGT} A_i * T_i * G_i$

where q_{ij} is gene expression of sample i for gene j, T_i is treatment (control: 0, drought:

1), A_i is allele type (Bd21: 0, Bd3-1: 1), G_i is generation (F1: 0, Parents: 1). Accordingly,

β_A tests for cis-acting regulation; β_{AG} tests for trans-acting; β_{AT} tests for cis-by-

environment-acting; and β_{AGT} tests for trans-by-environment-acting. Here, the approach

was also modified to perform tests in each environmental condition (as a "half model").

To be specific, each test of cis/trans is done separately in each condition because of our

limited power and low expression of the gene in samples in some conditions; GxE

cannot be directly tested in this analysis. P-values <0.1 is reported to be significant

without multiple test correction. Results are summarized in Supplementary Data Set 12.

Accession numbers

RNASequencing data for DGE and network analysis and ASE from this article can be

found in the GenBank data libraries under GSE267880 and GSE267878. All source

code used in this study is available online

(https://github.com/jyjiey/GxE_in_gene_regulation_network).

SUPPLEMENTARY DATA

Supplementary Data Set 1. Gene DGE list

Supplementary Data Set 2. GxE DGE gene GO enrichment

Supplementary Data Set 3. Module common genes

Supplementary Data Set 4. Module common gene annotation sorted

Supplementary Data Set 5. Module overlap p-value and intersect gene number

Supplementary Data Set 6. Module G,E,GxE DGE gene enrichment p-value

Supplementary Data Set 7. Module GO enrichment

Supplementary Data Set 8. Module traits correlation p-value

Supplementary Data Set 9. Traits data along dry-down

Supplementary Data Set 10. Traits data on day6 matching RNA samples

Supplementary Data Set 11. Gene module membership

Supplementary Data Set 12. ASE type

Supplementary Figure 1. Other physiological traits during dry-down in addition to

Figure 1

Supplementary Figure 2. Inferred directed interactions within selected modules with

gene DGE and edge DGE changes labeled

Supplementary Figure 3. Examples of regulations between pairs of genes selected in

addition to Figure 6

Supplementary Figure 4. ASE of the genes in Figure 6

Supplementary Figure 5. Heatmap comparing CILP and correlation variation from

regression method

Supplementary Figure 6. Mean reads of Bd21 allelic count vs. mean reads of Bd3-1

allelic count across all samples

Supplementary Figure 7. PCA of allelic gene expression

Supplementary Table 1. Percentage of E, G, G+E and GxE DGE change and slope

change among edges identified in small modules with inferred directed interactions

Supplementary Table 2. Models selected for physiological traits during dry-down

reported in Figure 1 and Supplementary Figure 1

Supplementary File 1. Direct test of DGE change between two genes

ACKNOWLEDGEMENTS

This work was supported by a grant from the MIT Jameel Water and Food Systems Lab

(JWAFS) and NSF CAREER Award (IOS 2239070) to DLD. JY was supported, in part,

by a graduate research fellowship from MIT JWAFS. ACW & AR were supported

through the United States Department of Energy contract no. DE-SC0012704 to

Brookhaven National Laboratory. We acknowledge the MIT SuperCloud and Lincoln

Laboratory Supercomputing Center for providing (HPC, database, consultation)

resources that have contributed to the research results reported within this paper. We

also acknowledge the computational support of Massachusetts Green High-

Performance Computer the Commonwealth Computational Cloud for Data Driven

Biology (C3DDB).

AUTHOR CONTRIBUTIONS

DLD and JY designed the research; JY, ACB, AR and DLD performed research; JY and DLD design the analytic and computational tools, analyzed data. DLD and JY wrote the paper with contributions from AR and ACB.

REFERENCE

1. Hill, M. S., Vande Zande, P. & Wittkopp, P. J. Molecular and evolutionary processes generating variation in gene expression. *Nature Reviews Genetics* 1–13 (2020) doi:10.1038/s41576-020-00304-w.

2. Wittkopp, P. J. & Kalay, G. Cis-regulatory elements: molecular mechanisms and evolutionary processes underlying divergence. *Nat Rev Genet* **13**, 59–69 (2012).

3. Veltsos, P. & Kelly, J. K. The quantitative genetics of gene expression in Mimulus guttatus. *PLOS Genetics* **20**, e1011072 (2024).

4. Huang, W., Carbone, M. A., Lyman, R. F., Anholt, R. R. H. & Mackay, T. F. C. Genotype by environment interaction for gene expression in Drosophila melanogaster. *Nat Commun* **11**, 5451 (2020).

5. National Academies of Sciences, E. *Next Steps for Functional Genomics: Proceedings of a Workshop*. (2020). doi:10.17226/25780.

6. Rockman, M. V. Reverse engineering the genotype–phenotype map with natural genetic variation. *Nature* **456**, 738–744 (2008).

7. Des Marais, D. L., Hernandez, K. M. & Juenger, T. E. Genotype-by-Environment Interaction and Plasticity: Exploring Genomic Responses of Plants to the Abiotic Environment. *Annual Review of Ecology, Evolution, and Systematics* **44**, 5–29 (2013).

8. Lowry, D. B. *et al.* Expression Quantitative Trait Locus Mapping across Water Availability Environments Reveals Contrasting Associations with Genomic Features in Arabidopsis. *The Plant Cell* **25**, 3266–3279 (2013).

9. Lovell, J. T. *et al.* The genomic landscape of molecular responses to natural drought stress in Panicum hallii. *Nat Commun* **9**, 5213 (2018).

10. Josephs, E. B. *et al.* The Evolutionary Forces Shaping Cis- and Trans-Regulation of Gene Expression within a Population of Outcrossing Plants. *Molecular Biology and Evolution* **37**, 2386–2393 (2020).

11. Josephs, E. B., Lee, Y. W., Stinchcombe, J. R. & Wright, S. I. Association mapping reveals the role of purifying selection in the maintenance of genomic variation in gene expression. *Proceedings of the National Academy of Sciences* **112**, 15390–15395 (2015).

12. Varoquaux, N. *et al.* Transcriptomic analysis of field-droughted sorghum from seedling to maturity reveals biotic and metabolic responses. *Proceedings of the National Academy of Sciences* **116**, 27124–27132 (2019).

13. Winkelmüller, T. M. *et al.* Gene expression evolution in pattern-triggered immunity within Arabidopsis thaliana and across Brassicaceae species. *The Plant Cell* (2021) doi:10.1093/plcell/koab073.

14. Wu, T.-Y. *et al.* Evolutionarily conserved hierarchical gene regulatory networks for plant salt stress response. *Nat. Plants* **7**, 787–799 (2021).

15. Reynoso, M. A. *et al.* Evolutionary flexibility in flooding response circuitry in angiosperms. *Science* **365**, 1291–1295 (2019).

16. Bradshaw, A. D. Evolutionary Significance of Phenotypic Plasticity in Plants. in *Advances in Genetics* (eds. Caspari, E. W. & Thoday, J. M.) vol. 13 115–155 (Academic Press, 1965).

17. Smith, H. Signal perception, differential expression within multigene families and the molecular basis of phenotypic plasticity. *Plant, Cell & Environment* **13**, 585–594 (1990).

18. Fox, R. J., Donelson, J. M., Schunter, C., Ravasi, T. & Gaitán-Espitia, J. D. Beyond buying time: the role of plasticity in phenotypic adaptation to rapid environmental change. *Philosophical Transactions of the Royal Society B: Biological Sciences* **374**, 20180174 (2019).

19. Xiong, W. *et al.* Increased ranking change in wheat breeding under climate change. *Nat. Plants* **7**, 1207–1212 (2021).

20. Choquette, N. E. *et al.* Environment-specific selection alters flowering-time plasticity
and results in pervasive pleiotropic responses in maize. *New Phytologist* **238**, 737–
749 (2023).

21. Levis, N. A. & Pfennig, D. W. Evaluating 'Plasticity-First' Evolution in Nature: Key
Criteria and Empirical Approaches. *Trends Ecol Evol* **31**, 563–574 (2016).

22. Rispail, N. *et al.* Multi-Environmental Trials Reveal Genetic Plasticity of Oat
Agronomic Traits Associated With Climate Variable Changes. *Front. Plant Sci.* **9**,
(2018).

23. Des Marais, D. L. & Juenger, T. E. Pleiotropy, plasticity, and the evolution of plant
abiotic stress tolerance. *Annals of the New York Academy of Sciences* **1206**, 56–79
(2010).

24. Remold, S. K. & Lenski, R. E. Pervasive joint influence of epistasis and plasticity on
mutational effects in Escherichia coli. *Nat Genet* **36**, 423–426 (2004).

25. Wei, S. *et al.* A transcriptional regulator that boosts grain yields and shortens the
growth duration of rice. *Science* **377**, eabi8455 (2022).

26. Da Ros, L. *et al.* Multi-omics atlas of combinatorial abiotic stress responses in
wheat. *The Plant Journal* **116**, 1118–1135 (2023).

27. Shu, J. *et al.* Transcriptomic, proteomic, metabolomic, and functional genomic
approaches of Brassica napus L. during salt stress. *PLOS ONE* **17**, e0262587
(2022).

28. Xu, K. *et al.* Proteomic analysis reveals the molecular mechanism underlying the
cold acclimation and freezing tolerance of wheat (Triticum aestivum L.). *Plant Sci*
318, 111242 (2022).

29. Amiour, N. *et al.* The use of metabolomics integrated with transcriptomic and
proteomic studies for identifying key steps involved in the control of nitrogen
metabolism in crops such as maize. *Journal of Experimental Botany* **63**, 5017–5033
(2012).

30. Zhang, H., Zhu, J., Gong, Z. & Zhu, J.-K. Abiotic stress responses in plants. *Nat Rev*
Genet **23**, 104–119 (2022).

31. Passioura, J. B. Translational research in agriculture. Can we do it better? *Crop*
Pasture Sci. **71**, 517–528 (2020).

32. Nicotra, A. B. *et al.* Plant phenotypic plasticity in a changing climate. *Trends Plant*
Sci **15**, 684–692 (2010).

33. Des Marais, D. L. *et al.* Physiological Genomics of Response to Soil Drying in
Diverse Arabidopsis Accessions[W][OA]. *Plant Cell* **24**, 893–914 (2012).

34. Shin, J. H. *et al.* Transcriptomic changes due to water deficit define a general
soybean response and accession-specific pathways for drought avoidance. *BMC*
Plant Biol **15**, 26 (2015).

35. Liu, X., Li, Y. I. & Pritchard, J. K. Trans Effects on Gene Expression Can Drive
Omnigenic Inheritance. *Cell* **177**, 1022-1034.e6 (2019).

36. Control of Transcription Factor Activity. in *Plant Genes, Genomes and Genetics*
149–160 (John Wiley & Sons, Ltd, 2015). doi:10.1002/9781118539385.ch10.

37. Transcription Factors Interpret cis-Regulatory Information. in *Plant Genes, Genomes*
and Genetics 133–148 (John Wiley & Sons, Ltd, 2015).
doi:10.1002/9781118539385.ch9.

38. Making mRNAs – Control of Transcription by RNA Polymerase II. in *Plant Genes,*
Genomes and Genetics 121–131 (John Wiley & Sons, Ltd, 2015).
doi:10.1002/9781118539385.ch8.

39. Marand, A. P., Eveland, A. L., Kaufmann, K. & Springer, N. M. cis-Regulatory
Elements in Plant Development, Adaptation, and Evolution. *Annu Rev Plant Biol* **74**,
111–137 (2023).

40. Jones, D. M. & Vandepoele, K. Identification and evolution of gene regulatory
networks: insights from comparative studies in plants. *Current Opinion in Plant*
Biology **54**, 42–48 (2020).

41. Alon, U. Network motifs: theory and experimental approaches. *Nat Rev Genet* **8**,
450–461 (2007).

42. Lachowiec, J., Queitsch, C. & Kliebenstein, D. J. Molecular mechanisms governing
differential robustness of development and environmental responses in plants. *Ann*
Bot **117**, 795–809 (2016).

43. Thompson, D., Regev, A. & Roy, S. Comparative analysis of gene regulatory
networks: from network reconstruction to evolution. *Annu Rev Cell Dev Biol* **31**,
399–428 (2015).

44. Sun, Y. & Dinneny, J. R. Q&A: How do gene regulatory networks control
environmental responses in plants? *BMC Biology* **16**, 38 (2018).

45. Wagner, G. P., Pavlicev, M. & Cheverud, J. M. The road to modularity. *Nature*
Reviews Genetics **8**, 921–931 (2007).

46. Grishkevich, V. *et al.* A genomic bias for genotype–environment interactions in C.
elegans. *Molecular Systems Biology* **8**, 587 (2012).

47. Grishkevich, V. & Yanai, I. The genomic determinants of genotype × environment
interactions in gene expression. *Trends Genet* **29**, 479–487 (2013).

48. Des Marais, D. L., Guerrero, R. F., Lasky, J. R. & Scarpino, S. V. Topological
features of a gene co-expression network predict patterns of natural diversity in
environmental response. *Proceedings of the Royal Society B: Biological Sciences*
284, 20170914 (2017).

49. Luscombe, N. M. *et al.* Genomic analysis of regulatory network dynamics reveals
large topological changes. *Nature* **431**, 308–312 (2004).

50. Scholthof, K.-B. G., Irigoyen, S., Catalan, P. & Mandadi, K. K. Brachypodium: A
Monocot Grass Model Genus for Plant Biology. *The Plant Cell* **30**, 1673–1694
(2018).

51. Draper, J. *et al.* Brachypodium distachyon. A New Model System for Functional
Genomics in Grasses. *Plant Physiol* **127**, 1539–1555 (2001).

52. Des Marais, D. L., Lasky, J. R., Verslues, P. E., Chang, T. Z. & Juenger, T. E.
Interactive effects of water limitation and elevated temperature on the physiology,
development and fitness of diverse accessions of Brachypodium distachyon. *New*
Phytologist **214**, 132–144 (2017).

53. López-Alvarez, D. *et al.* Environmental niche variation and evolutionary
diversification of the Brachypodium distachyon grass complex species in their native
circum-Mediterranean range. *American Journal of Botany* **102**, 1073–1088 (2015).

54. Elowitz, M. B., Levine, A. J., Siggia, E. D. & Swain, P. S. Stochastic Gene
Expression in a Single Cell. *Science* **297**, 1183–1186 (2002).

55. Cortijo, S., Bhattarai, M., Locke, J. C. W. & Ahnert, S. E. Co-expression Networks
From Gene Expression Variability Between Genetically Identical Seedlings Can
Reveal Novel Regulatory Relationships. *Front. Plant Sci.* **11**, (2020).

56. Livingston, D. P., Hincha, D. K. & Heyer, A. G. Fructan and its relationship to abiotic
stress tolerance in plants. *Cell. Mol. Life Sci.* **66**, 2007–2023 (2009).

57. Wang, M., Zhao, Y. & Zhang, B. Efficient Test and Visualization of Multi-Set
Intersections. *Sci Rep* **5**, 16923 (2015).

58. Xiong, Y. *et al.* Glucose–TOR signalling reprograms the transcriptome and activates
meristems. *Nature* **496**, 181–186 (2013).

59. Tzvetkova-Chevolleau, T. *et al.* The light stress-induced protein ELIP2 is a regulator
of chlorophyll synthesis in Arabidopsis thaliana. *The Plant Journal* **50**, 795–809
(2007).

60. Global Profiling of Rice and Poplar Transcriptomes Highlights Key Conserved
Circadian-Controlled Pathways and cis-Regulatory Modules. *PLOS ONE* **6**, e16907
(2011).

61. Marchand, G. *et al.* Bridging physiological and evolutionary time-scales in a gene
regulatory network. *New Phytol* **203**, 685–696 (2014).

62. Chevin, L.-M., Leung, C., Le Rouzic, A. & Uller, T. Using phenotypic plasticity to
understand the structure and evolution of the genotype-phenotype map. *Genetica*
150, 209–221 (2022).

63. Wagner, G. P. & Zhang, J. The pleiotropic structure of the genotype–phenotype
map: the evolvability of complex organisms. *Nature Reviews Genetics* **12**, 204–213
(2011).

64. Laitinen, R. A. E. & Nikoloski, Z. Genetic basis of plasticity in plants. *J Exp Bot* **70**,
739–745 (2019).

65. Hämälä, T., Gorton, A. J., Moeller, D. A. & Tiffin, P. Pleiotropy facilitates local
adaptation to distant optima in common ragweed (Ambrosia artemisiifolia). *PLOS*
Genetics **16**, e1008707 (2020).

66. Lea, A. *et al.* Genetic and environmental perturbations lead to regulatory
decoherence. *eLife* **8**, e40538 (2019).

67. Des Marais, D. L., Razzaque, S., Hernandez, K. M., Garvin, D. F. & Juenger, T. E.
Quantitative trait loci associated with natural diversity in water-use efficiency and
response to soil drying in Brachypodium distachyon. *Plant Science* **251**, 2–11
(2016).

68. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models

Using lme4. *Journal of Statistical Software* **67**, 1–48 (2015).

69. R Core Team (2022). R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

70. Bartoń, K. MuMIn: Multi-Model Inference. (2023).

71. Lenth, R. V. *et al.* emmeans: Estimated Marginal Means, aka Least-Squares Means.

(2024).

72. ggplot2 Based Publication Ready Plots. https://rpkgs.datanovia.com/ggpubr/.

73. ggplot2: Elegant Graphics for Data Analysis | SpringerLink.

https://link.springer.com/book/10.1007/978-3-319-24277-4.

74. Xiong, Y. *et al.* A Comparison of mRNA Sequencing with Random Primed and 3′-

Directed Libraries. *Scientific Reports* **7**, 14626 (2017).

75. Soumillon, M., Cacchiarelli, D., Semrau, S., Oudenaarden, A. van & Mikkelsen, T. S.

Characterization of directed differentiation by high-throughput single-cell RNA-Seq.

bioRxiv 003236 (2014) doi:10.1101/003236.

76. Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for

Illumina Sequence Data. Bioinformatics, btu170.

77. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet.journal* **17**, 10–12 (2011).

78. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat*

Methods **9**, 357–359 (2012).

79. Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data

with or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

80. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

81. Kassambara, A. & Mundt, F. factoextra: Extract and Visualize the Results of

Multivariate Data Analyses. (2020).

82. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation

network analysis. *BMC Bioinformatics* **9**, 559 (2008).

83. Csárdi, G. & Nepusz, T. The igraph software package for complex network research.

in (2006).

84. Csárdi, G. *et al.* igraph: Network Analysis and Visualization. (2023).

85. Thomas, P. D. *et al.* PANTHER: A Library of Protein Families and Subfamilies

Indexed by Function. *Genome Res.* **13**, 2129–2141 (2003).

86. Rezwani, M., Pourfathollah, A. A. & Noorbakhsh, F. rbioapi: user-friendly R interface

to biologic web services' API. *Bioinformatics* **38**, 2952–2953 (2022).

87. Ksouri, N. *et al.* Tuning promoter boundaries improves regulatory motif discovery in

nonmodel plants: the peach example. *Plant Physiology* **185**, 1242–1258 (2021).

88. McLeay, R. C. & Bailey, T. L. Motif Enrichment Analysis: a unified framework and an

evaluation on ChIP data. *BMC Bioinformatics* **11**, 165 (2010).

89. O'Malley, R. C. *et al.* Cistrome and Epicistrome Features Shape the Regulatory

DNA Landscape. *Cell* **165**, 1280–1292 (2016).

90. Squires, C., Wang, Y. & Uhler, C. Permutation-Based Causal Structure Learning

with Unknown Intervention Targets. Preprint at

https://doi.org/10.48550/arXiv.1910.09007 (2020).

91. Meinshausen, N. & Bühlmann, P. Stability Selection. *Journal of the Royal Statistical*

Society Series B: Statistical Methodology **72**, 417–473 (2010).

92. Belyaeva, A. *et al.* Causal network models of SARS-CoV-2 expression and aging to

identify candidates for drug repurposing. *Nat Commun* **12**, 1024 (2021).

93. Rgraphviz. *Bioconductor* http://bioconductor.org/packages/Rgraphviz/.

94. Fox, J. *et al.* car: Companion to Applied Regression. (2023).

95. RNAseq short variant discovery (SNPs + Indels). *GATK*

https://gatk.broadinstitute.org/hc/en-us/articles/360035531192-RNAseq-short-

variant-discovery-SNPs-Indels- (2023).

96. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15

(2013).

97. Auwera, G. A. V. der & O'Connor, B. D. *Genomics in the Cloud: Using Docker,*

GATK, and WDL in Terra. (O'Reilly Media, Inc., 2020).

98. g2gtools documentation. https://churchill-lab.github.io/g2gtools/.

99. Raghupathy, N. *et al.* Hierarchical analysis of RNA-seq reads improves the accuracy

of allele-specific expression. *Bioinformatics* **34**, 2177–2184 (2018).

100. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-
efficient alignment of short DNA sequences to the human genome. *Genome Biology*
10, R25 (2009).
101. Gould, B. A., Chen, Y. & Lowry, D. B. Gene regulatory divergence between
locally adapted ecotypes in their native habitats. *Molecular Ecology* **27**, 4174–4188
(2018).
102. Lovell, J. T. *et al.* Drought responsive gene expression regulatory divergence
between upland and lowland ecotypes of a perennial C4 grass. *Genome Res* **26**,
510–518 (2016).

1236
1237
1238

SUPPLEMENTARY INFORMATION

1240

1241

Supplementary Figure 1. Other physiological traits during dry-down in addition to Figure 1. Comparison of physiological and metabolic traits between control and dry-down in two accessions from Day 0 to Day 7. (Generalized) Linear models were determined based on AIC score. Random-effects components of the models were based on

1246   likelihood ratio tests. Based on the selected model, differences of means shown here

1247   with asterisks, were determined using contrasts between the two treatments at each

1248   time point in each genotype by Tukey's test. N = 9 for each point.

1249

A

B

C

1257   Supplementary Figure 2. Inferred directed interactions within selected modules with

1258   gene DGE and edge DGE changes, corresponding to **A)** Module 12 in Bd21d, **B)**

1259   Module 14 in Bd3-1d and **C)** Module 10 in Bd21d shown in Figure 5, with node colors

1260   are based on gene DGE and edge colors are based on edge DGE changes (blue: E,

1261   green: G, brown: G+E, orange: GxE, black/white: none).

1262

1263

1264   Supplementary Figure 3. Examples of regulations between pairs of genes selected in

1265   addition to Fig.6 **A)** shows regulation E slope changes, and **B)** shows transmission of

1266   GxE DGE. They are illustrated with linear regression between regulator and target, their

1267   expression levels based on log2(norm_count+1). Bd21c, Bd21d, Bd3-1c, Bd3-1d are

1268   colored with blue, red, purple, orange, respectively.

1269

1270

1271 Supplementary Figure 4. ASE of the genes in Figure 6 (Bradi3g00920, Bradi1g15840,

1272 Bradi1g53680, Bradi3g45230). F1 is expression in hybrid and F0 is expression in

1273 parents. Values are normalized count. Bd21c, Bd21d, Bd3-1c, Bd3-1d are colored with

1274 blue, red, purple, orange, respectively.

1275

Supplementary Figure 5. Heatmap comparing CILP and correlation variation from
regression method based on gene pairs with correlation >0.75 in any of the library
types. The number represent the # of the cases found in the intersect. The three
number in the parenthesis are p-value of fisher's exact test, the proportion of the genes
in the intersect in the correlation category, and the proportion of the genes in the
intersect in the CILP category. The color is based on p-value.

1283

1284 Supplementary Figure 6. mean reads of Bd21 allelic count vs. mean reads of Bd3-1

1285 allelic count across all samples of allele specific genes suggest no alignment bias to

1286 Bd21 or Bd3-1

1287

1288

1289 Supplementary Figure 7. PCA of allelic gene counts colored based on samples

1290 genotype and treatment (after normalization)

1291

1292 Supplementary Table 1. Percentage of E, G, G+E and GxE DGE change, or slope

1293 change identified in edges in small module (<40 genes) inferred gene regulation

1294 network (e.g., Figure 5).

module no.	total edges	edge DGE change				edge rewiring %			
		G	E	G+E	GxE	G	E	G+E	GxE
bd21c_13	67	25	4	10	6	10%	12%	3%	9%
bd21c_14	52	19	8	5	0	8%	17%	0%	6%
bd21c_17	37	16	4	6	0	5%	5%	3%	5%
bd21c_21	30	8	4	12	0	10%	20%	3%	7%
bd21c_22(part of 9)	22	5	3	3	0	5%	23%	9%	5%
bd21c_23(part of 10)	25	5	3	10	0	4%	8%	4%	0%
bd21d_12	41	5	11	12	11	34%	15%	5%	12%
bd21d_13	32	2	11	7	5	44%	3%	6%	6%
bd21d_16 (part of 5)	23	6	9	6	1	17%	4%	4%	17%
bd31d_12	68	15	16	11	6	10%	9%	0%	1%
bd31d_14	51	17	3	8	0	10%	4%	4%	4%
bd31d_18(part of 21)	32	4	10	10	2	13%	19%	9%	9%
bd31d_19	108	44	15	18	4	7%	24%	5%	6%
bd31c_13	79	8	4	0	0	8%	5%	3%	13%
bd21c_5	65	25	9	8	0	5%	18%	2%	9%
bd31c_10	55	11	19	10	0	7%	9%	2%	11%
bd31c_16	27	13	1	3	0	15%	19%	0%	15%

1295

1296 Supplementary Table 2. Models used for physiological traits during dry-down reported in

1297 both Figure 1 and Supplementary Figure 1. (Generalized) Linear models were

1298 determined based on AIC score. Random-effects components of the models were

1299 based on likelihood ratio tests.

No.	Trait	Model selected
1	Water usage	Accession +Treatment+harvest.day+ Accession*harvest.day+ Treatment*harvest.day
2	Hydraulic potential	Accession +Treatment+harvest.day+ Treatment*harvest.day
3	Net photosynthesis rate	Accession +Treatment+harvest.day+ Treatment*harvest.day
4	gsw	Treatment+harvest.day+ Treatment*harvest.day

5	Fresh weight	Accession +Treatment+harvest.day+ Treatment*harvest.day
6	Dry weight	Accession +Treatment+harvest.day+ Treatment*harvest.day
7	Dry root shoot ratio	Accession +Treatment+harvest.day+ Treatment*harvest.day
8	Glucose	Accession + Treatment +harvest.day+ Accession* Treatment +Treatment*harvest.day
9	fv'/fm'	Accession
10	Relative chlorophyll content	Accession +harvest.day
11	Fructose	Accession + Treatment +harvest.day+ Accession* harvest.day + Treatment*harvest.day+(1| DW.Plate)
12	Sucrose	Accession
13	Low DP fructan	-
14	Starch	Accession + Treatment +harvest.day+ Accession* harvest.day + Treatment*harvest.day+(1| DW.Plate..)
15	High DP fructan	harvest.day
16	Protein	Accession + Treatment +harvest.day+Treatment*harvest.day
17	Amino acid	Treatment +harvest.day+Treatment*harvest.day

1300

Supplementary File 1 Direct test of DGE change between two genes

Direct test of DGE change between two genes

We fit pairs of genes in regressions to formally test whether DGE changes occur

between them. To be specific, a vector of log2 -transformed transcript abundance for

two correlated genes were fit with the following model:

$$x = gene * g * e$$

where x is a combined vector of gene expression of two genes, and variable

$gene$ (1,2) differentiates which the gene of gene expression value x. Thus, the

regression coefficients of $gene * g$ which differ significantly from 0 suggest that effects

of genotype on transcript abundances differ between the two genes, and similarly

coefficients of $gene * e$ which differ significantly from 0 suggest that effects of treatment

environment on transcript abundances differ between the two genes, and significant

coefficients for $gene * g * e$ suggests a GxE effect on transcript abundance. P-values

<0.001 is reported to be significant without multiple test corrections. Results are shown

in Supplementary Figure 4. This method avoids the indirect comparison by DGEs of two

genes, but generally consistent with it (see Supplementary Figure 4).

1317

1318   Examples illustrating gene-gene interaction. **A-C)** DGE G changes; **E-G)** DGE GxE

1319   changes. In each type of variation, the three plots show examples of different responses

1320   (**A,D**), direction (**B,E**), and sensitivity(**C,F**), respectively. Red and blue dots show two

1321   gene expression levels respectively, in four library types.

1322