1 Title

2 Context-dependent coordination of movement in *Tribolium castaneum* larvae

3 Running title

4 Beetle larvae tune movements to context

5 Authors

6 Bella Xu Ying[1,2,3], *Maarten F. Zwart[1,2,3], **Stefan R. Pulver[1,2,3]

7 Affiliations

[1] Institute of Behavioural and Neural Sciences, University of St Andrews, St Andrews,
United Kingdom
[2] Centre of Biophotonics, University of St Andrews, St Andrews, United Kingdom
[3] School of Psychology and Neuroscience, University of St Andrews, St Andrews,
United Kingdom

*Correspondence: mfz@st-andrews.ac.uk
**Correspondence: sp96@st-andrews.ac.uk

15 Key words

16 *Tribolium castaneum*, locomotion, kinematics, motor control, neuroethology, central
17 nervous system

18 Summary statement

19 Red flour beetle larvae walk with a legged wave gait and use their tails as anchors to
20 climb inclines and tunnel into flour.

Abstract

Insect pests, like the red flour beetle *Tribolium castaneum*, destroy up to 20% of stored grain products worldwide, making them a significant threat to food security. Their success hinges upon adapting their movements to unpredictable, heterogeneous environments like flour. *Tribolium* is well developed as a genetic model system; however, little is known about their natural locomotion and how their nervous systems coordinate adaptive movement. Here, we employed videographic whole-animal and leg tracking to assess how *Tribolium* larvae locomote over different substrates and analyze their gait kinematics across speeds. Unlike many hexapods, larvae employed a bilaterally symmetric, posterior-to-anterior wave gait during fast locomotion. At slower speeds, coordination within thoracic segments was disrupted, although intersegmental coordination remained intact. Moreover, larvae used terminal abdominal structures (pygopods) to support challenging movements, such as climbing overhangs. Pygopod placement coincided with leg swing initiation, suggesting a stabilizing role as adaptive anchoring devices. Surgically lesioning the connective between thoracic and abdominal ganglia impaired pygopod engagement and led to escalating impairments in flat-terrain locomotion, climbing and tunnelling. These results suggest that effective movement in *Tribolium* larvae requires thoracic-abdominal coordination, and that larval gait and limb recruitment is context-dependent. Our work provides the first kinematic analysis of *Tribolium* larval locomotion and gives insights into its neural control, creating a foundation for future motor control research in a genetically tractable beetle that jeopardizes global food security.

Introduction

Stored product insects are devastating agricultural pests. These animals have been estimated to cause 9% of postharvest loss in developed countries and 20% in developing countries (reviewed in (Phillips and Throne, 2010)). The red flour beetle, *Tribolium castaneum (Tribolium),* is representative of many stored product pest insects presenting a major challenge to global food security. Larvae and adults consume wheat products like flour by efficiently moving over and tunnelling through these unstable, shifting substrates (Good, 1936). Therefore, understanding how *Tribolium* adapt their movements to heterogeneous and challenging terrain could provide

insights into conserved principles of motor control whilst aiding the development of
new pest control approaches.

Animals use various strategies for adapting movement to the environments in which
they live, continuously monitoring and adjusting movements on a cycle-by-cycle basis
to cope with unexpected obstacles and/or drastic variations in the physical properties
of 3-dimensional terrain. Studying the neural control of movement adaptability has
been a long-standing focus of motor systems research across multiple species.
Terrestrial animals with internal, external, or hydrostatic skeletons have all evolved
different strategies for movement control in different contexts (reviewed in insects
(Mantziaris et al., 2020; van Griethuijsen and Trimmer, 2014), mammals (Leiras et al.,
2022) and birds (Daley, 2018)). Some animals deploy limbs (Hooper et al., 2009;
Picker et al., 2012; Triphan et al., 2010), some utilize soft outer cuticular features
(Digumarti et al., 2019; Booth et al., 2024) and others adapt body parts not typically
involved in locomotion to solve motor problems (Shield et al., 2021). Similarly,
Tribolium exhibit a diverse behavioral repertoire in response to different sensorimotor
cues, and adult anatomy has been well-described. Previous work has shown that
Tribolium can move through a variety of stored grain, but prefer finely grained, whole
wheat flour over coarser bran (Đukić et al., 2020). Moreover, adults can navigate in
response to attractive and aversive odors (Đukić et al., 2021; Gerken et al., 2018),
shapes (Semeao et al., 2011), light levels (Duehl et al., 2011), colors (Prakashini et
al., 2023), and different habitat cues (Romero et al., 2010). Both larvae and adults
have articulated, jointed limbs (Good, 1936), with larvae being composed of both soft
ventral cuticle and semi-hardened dorsal cuticle. While anatomical studies have begun
to characterize the larval neuromuscular system (Schultheis et al., 2019), knowledge
concerning the specific kinematics of locomotion and how the *Tribolium* nervous
system recruits thoracic limbs and abdominal segments to meet different locomotor
demands is extremely limited.

Past studies have prioritized adult anatomy and behavior, although larval instars are
equally capable of maneuvering and colonizing stored grain. Here we characterize
locomotor parameters in late-instar *Tribolium* larvae to gain insights into how these
animals adapt movements to their environment. We analyze how larvae negotiate
different substrates and combine whole-animal and leg tracking to characterize and

quantify leg kinematics underlying a range of locomotor speeds. We find larvae have

an ethologically relevant preference for fibrous substrates and walk primarily with a

posterior-to-anterior-propagating bilaterally symmetric gait, which shifts towards

uncoordinated exploratory movements at slower speeds. Severing the connectives

between thoracic and abdominal segments impairs walking, climbing, and tunneling

but does not disrupt fundamental leg kinematics. In addition, we characterize a

pygopod planting behavior, abolished in animals with severed connectives, which is

used for stability and propulsion under different motor contexts (challenged locomotion

vs tunnelling). These results provide first insights into the biomechanical

underpinnings of *Tribolium* movement and the neural basis of ethologically relevant

motor programs in an insect species with direct impacts on global food security.

Materials and methods

Animals

Wild-type *Tribolium castaneum* (San Bernardino strain) were reared in 500ml

borosilicate bottles with perforated lids (VWR SCOT291182809) at 29°C in organic

whole wheat flour and 5% dried active yeast per weight. 4-5[th] instar larvae

approximating 4-6mm in length were used in all experiments.

Statistics

Main text results and box plots are given as mean ±SE, and box whiskers are ±SD

unless otherwise stated. Details of statistical analysis are included under each

experiment's subsection below and in figure legends.

Substrate testing

Basic locomotor ability was tested on twelve substrates – white printer paper,

cardboard, parafilm, lens paper, Sylgard™, glass, velvet, sandpaper, plastic, packing

foam, white flour, whole wheat flour, 1%, 5% and 10% agarose. Ten larvae were placed

in the center of the substrate and recorded for 1 minute. Movies were recorded with a

mobile camera (SM-A415F/DSN, Samsung, Seoul, South Korea) at 30fps and

analyzed with Fiji (Fiji is Just ImageJ) (Schindelin et al., 2012). Larval head position

was manually tracked for the first 15 seconds (since larvae on preferred substrates

often walked out of frame within this time window) with the Manual Tracking plugin (https://imagej.net/ij/plugins/track/track.html) for Fiji.

Free locomotion assay

30 larvae were placed in groups of 3-4 in a 18x18cm arena lined with white printer paper and recorded for 2 minutes in darkness with a custom-built imaging setup using an MQ013MG-E2 XIMEA CMOS (Ximea GmbH, Münster, Germany) camera with a 35mm C-mount lens (Thorlabs Inc, NJ, USA). Animals were illuminated with 850nm infrared light using a Thorlabs M850L3 LED (Thorlabs Inc, NJ, USA). Larvae were tracked with FIMTrack (Risse et al., 2014). Statistical analysis on biases in headsweep direction was performed as an intercept-only general linear model (GLM) in R.

Gait characterization

Ten larvae were placed on white printer paper under an Olympus SZX10 (Olympus Microscopes, Tokyo, Japan) dissection microscope at 0.8x magnification using a 0.5x objective. Animals were recorded with an MQ042RG-CM XIMEA CMOS (Ximea GmbH, Münster, Germany) camera at 60fps. 3-5 straight walk cycles were extracted and manually tracked with Fiji's Manual Tracking plugin. Gait dynamics and locomotor parameters were analyzed with Python scripts for individual legs; cycle period was calculated as 1 over average cycle frequency. A Savitzky-Golay low-pass filter was applied to manually tracked velocities to discretize swing durations. Stance duration was calculated as the period between consecutive swings. Duty cycle was calculated as average swing/stance duration per segment over average cycle period per animal. Intrasegmental phase difference represents the phase difference between left and right legs within each thoracic segment. Intersegmental phase difference represents phase difference of left legs in each segment relative to left T3.

Coherence and phase analysis was performed with MATLAB application *Groundswell* (https://github.com/JaneliaSciComp/Groundswell). A fast Fourier transform was performed on the anchor leg's velocity trace to determine its spectral composition. The anchor's fundamental frequency was determined as the frequency with greatest power density (i.e., during swing phases) in the power spectrum. If the highest power frequency was ambiguous (i.e., appeared as blocks of plateaus on the spectrum), the middle value within the plateau was chosen, and the higher frequency block was used

144 in cases with two adjacent blocks. Coherence and phase relationships of the other five

145 legs to the anchor was compared at the determined fundamental frequency.

146 *Groundswell* uses coherence magnitude as a measure of the linear relationship

147 between two waveforms at the determined fundamental frequency, and calculates a

148 coherence magnitude threshold with the formula:

149
$$C\_mag\_thresh = \sqrt{1 - a^{\frac{1}{(\text{dof}/2 - 1)}}} \quad (1)$$

150 Where degrees of freedom (dof) = 2*R*K, in which R is the number of observations of

151 the waveforms together and K is the number of tapers. This threshold determines the

152 minimum coherence magnitude required for significant coherence between two

153 signals with α = 0.05.

Preliminary incline assay

Ten larvae were individually placed on custom-built platforms of varying incline (180°

– flat substrate, 135° – slight upwards incline, 90° – vertical wall, 45° – overhang).

Locomotion was filmed with an MQ022RG-CM XIMEA CMOS (Ximea GmbH, Münster,

Germany) camera for 1 minute at 30fps with a Thorlabs OSL1 Fiber Illuminator

(Thorlabs Inc, NJ, USA) with gooseneck lights at lowest intensity. Raw videos were

optimized for FIMTrack in Fiji with custom-written macros. Statistical analysis on mean

instantaneous velocity was performed as one-way ANOVA. Path curviness was

calculated with the formula:

163
$$(\text{Total distance travelled})/(\text{Total displacement}) * \text{Max displacement} \quad (2)$$

Connective severing assay

In preliminary control assays, six larvae were anesthetized on ice and sequestered

between 0.1mm tungsten pins ventral side up on a dissection dish. A cut was made

with fine microdissection scissors on the cuticle line separating A1 and A2 segments.

The blades were inserted at a ~45° angle to the cuticle and pulled upwards to sever

the connective. Sham larvae were given lateral lesions on A1. Control larvae were not

lesioned but anesthetized on ice for equal time. Surgeries were verified via fillet

microdissection post-experimentally. Larvae without clear, visible connective severing

were discarded from results. Behavior was filmed with the setup outlined in *Free*

locomotion assay 2 hours after surgery*;* raw videos were processed in Fiji and tracked
with FIMTrack. Statistical analysis on mean instantaneous velocity was performed as
a one-way ANOVA.

For following incline and tunnelling experiments, surgeries were repeated on seven
larvae; sham animals' lateral lesions alternated between left and right sides to
minimize biases in turning. Different animals were used for incline and tunnelling
assays to avoid excessive fatigue on lesioned animals.

Top-down connective severing incline assay

Same methodology as *Preliminary incline assay,* excluding 90° due to short lifespans
of severed animals. 2 larvae (1 control and 1 severed) could not be tracked with
FIMTrack,   thus   mean   velocity   was   obtained   with   wrMTrck
(https://www.phage.dk/plugins/wrmtrck.html) and their tracks are therefore absent in
Fig. 5A. Statistical analysis on mean instantaneous velocity was performed as one-
way ANOVAs within each inclination with Bonferroni corrected post-hocs.

Up-close connective severing incline assay

Control, sham and severed larvae were individually placed onto a walled 70mm x 3mm
paper platform on a custom-built rig; incline was adjusted with a micromanipulator and
verified with a protractor. Locomotion was filmed until the animal walked out of view
with a MQ042RG-CM XIMEA CMOS (Ximea GmbH, Münster, Germany) camera
connected with a 1"-32 C-mount (Thorlabs Inc, NJ, USA) to a Canon FR1048KWN
lens (Canon Inc, Tokyo, Japan). If the animal remained immobile, filming was stopped
at the experimenter's discretion. 180° videos were filmed at 100fps and illuminated by
an overhead Thorlabs M850L3 850nm light (Thorlabs Inc, NJ, USA). 45° overhang
trials were illuminated from the bottom with a Stemmer Imaging CCS LDL100 850nm
light (Stemmer Imaging Ltd, Tongham, UK) and limited to 50fps by the field of view.
135° and 90° were not tested due to the short lifespans of severed animals. For
analysis, velocity was tracked with FIMTrack; pygopod planting events and start and
stop frames of leg swings were manually tracked with Fiji's Manual Tracking plugin.
The constricted arena walls promoted exploration, therefore to effectively analyze
locomotor parameters and gait kinematics, walk cycles following the T3 to T1
sequence were extracted and stance durations above median ± 1SD were discarded

to filter out pause periods. Mean percentage of sequences extracted out of total tracked time (including pauses) per animal were: 55.4% (control 180°), 47.2% (sham 180°), 34.2% (severed 180°), 28.3% (control 45°), 24.9% (sham 45°) and 26.0% (severed 45°). Cycle period, stance duration and duty cycle retained their definitions as in _Gait characterization,_ but swing duration was calculated as the time between swing start and swing stop frames.

A pygopod plant event was defined as a visible extension of the pygopods, forming almost a blunt triangular point, into the substrate, regardless of whether the abdomen is in contact with the substrate. The Freedman-Diaconis rule was used in Fig. 5R, using the formula:

$$(2 * IQR)/(3\sqrt{n}) \quad (3)$$

to calculate x-axis bin length. Planting rate was used for Fig. 5T to normalize for variable video lengths, calculated as number of events over total tracked time. Pearson's correlations were applied for regression analyses.

Tunneling assay

Post-surgery animals were allowed to wake and begin walking on paper before testing. A circular dish was filled with ~10mm of flour; control, sham and severed larvae were individually placed and filmed with an MQ0113RG-E2 XIMEA CMOS (Ximea GmbH, Münster, Germany) at 15fps under a Nikon SMZ745T dissection scope at 0.63x magnification. Filming stopped at 2 minutes regardless of how deep larvae tunneled. Body lengths at time 0s and 120s were measured in Fiji. Statistics was performed as a Kruskal-Wallis test and Dunn post-hoc analysis with Bonferroni correction (given non-normal data distribution).

Results

Tribolium larvae locomote most efficiently over fibrous substrates

Tribolium beetles take approximately 35 days at 29ºC to undergo complete metamorphosis, comprising four distinct developmental stages – egg, larval, pupal, and adult (Devi and Devi, 2015) (Fig. 1A). The larval stage is characterized by three thoracic segments, each supported by a pair of legs, and nine abdominal segments.

Larvae possess two small, rounded ventral pygopods in the most posterior abdominal
segment, and two spiked urogomphi dorsally (Fig. 1B).

We first established larval substrate preference based on ease of locomotion. We
found larvae crawl best on textured, fibrous, wood-based substrates such as paper,
cardboard, and lens paper (Fig. 1C) (Movie 1). In contrast, they had substantial
difficulty gripping smooth surfaces like plastic and glass, thus preventing forwards
momentum. Similarly, their legs slipped on or stuck to deformable substrates like
Sylgard™ and agarose, also impeding coordinated progress (Fig. 1D,E). Notably, on
these substrates, we observed pygopod planting for substrate engagement in an
anchoring motion, which, when released, propelled the larvae forwards (Movie 1). This
abdominal engagement was rarely observed on fibrous substrates. Surprisingly,
crawling was heavily impeded over whole wheat flour and white flour (Fig. 1C-E), as
larvae attempted to tunnel upon contact with these substrates.

Larvae exhibit diverse motor behaviors during free locomotion

Next, movement parameters during free locomotion were analyzed to quantify the
spectrum of larval motor capabilities. Paper was used as the default substrate in
subsequent experiments given larval preference, accessibility, and ease of imaging.
Tribolium larvae exhibited heterogeneous walking patterns, and favored edge
exploration, i.e., they rarely strayed from arena edges once they found them (Fig. 2A).
Additionally, we observed persistent burrowing attempts at the paper edges during
preliminary trials in which the paper substrate was not fully adhered to the arena
platform.

Locomotion on paper had a mean instantaneous velocity of 1.8 ± 0.1 mm s^{-1} (range =
0.9 to 3.0 mm s^{-1}; larval sizes ranged 4-6 mm), and animals displayed a spectrum of
intra-animal velocity variability during only 2 minutes of exploration (Fig. 2B). Faster
animals were not constantly fast, but instead varied speed throughout locomotion,
whereas slower animals were limited to lower speed ranges (Fig. 2C). Larvae spent
equal time walking straight (49.6%) and turning (50.4%) (Fig. 2D). Animals decelerated
to turn but rarely stopped completely, thus staying immobile only 0.5% of the time (Fig.
2D). Finally, while some larvae showed a bias in turning direction, many engaged
equally in left and right headsweeps during exploration (p = .869; Fig. 2E).

We observed a larval motor program we termed "backtrack and redirect" (Movie 2),
defined as a headsweep/turn in one direction, followed by visible backwards
locomotion, and a contralateral headsweep/turn and forwards locomotion (Fig. 2F),
which on average lasted 3-6 seconds. This behavior followed a distinct sequence of
instantaneous velocity and body angle changes. First, velocity dropped as the animal
turned (body angle deviating from 180º), followed by an increase in velocity at 180º
body angle due to backtracking, and ended with another deceleration and deviation
from 180º in the opposite direction from the animal's original trajectory (Fig. 2G). While
63.3% of larvae displayed this motor program at least once during 2 minutes of free
locomotion (Fig. 2H), some displayed it a maximum of 8 times, and a third never
engaged in this behavior.

Gait coordination and leg kinematics during locomotion is speed-dependent

To characterize natural gait, leg kinematics and the coordination between thoracic
segments, we tracked legs during straight walking bouts on paper. We found *Tribolium*
larvae walked with a bilaterally symmetric posterior-to-anterior wave gait (Fig. 3A).
Then, we determined the frequency of locomotion for each animal by performing a
Fourier transform on leg velocities and selecting the frequency with highest power of
an anchor leg (left T1). We defined swing phases by a leg's lifting and planting on the
substrate and stance phases as the duration in which legs are in contact with the
ground. An average cycle period was then calculated for each leg, then averaged per
animal. After a swing, legs usually stanced next to the legs of the segment in front,
and these often began their swing phase before the previous contacted the ground
(Fig. 3A). This pattern was observed most prominently within the T2 to T1 transition
as their swing phases overlapped more visibly (Fig. 3B-D).

Fast (Fig. 3B,D,F) and slow (Fig. 3C,E,G) walk cycles differed markedly. In faster
animals, typical walk cycles averaged 0.2 ± 0.007s and were initiated by a T3 swing
phase, followed by T2 with a phase delay of approximately 120º, then T1 with another
120º delay. These delays were largely consistent between legs and across animals,
independent of speed (Fig. 3H). In fast animals, many legs were significantly coherent
with each other (Fig. 3J). In slower animals, average cycle period was 0.7 ± 0.1s, and
some legs often swung alone during walk cycles (Fig. 3E). Furthermore, not all six
legs were required to swing during each cycle. Absent leg swings consequently

impaired intrasegmental coordination; phase differences between contralateral leg
pairs increased (binned mean for 5 slowest larvae = 41.0 ± 7.5º) compared to faster
animals (binned mean for 5 fastest larvae = 15.1 ± 3.4º) (Fig. 3I). As such, fewer legs
were significantly coherent to each other in slower animals (Fig. 3J), but the T3 to T1
sequence was largely preserved (Fig. 3H). Finally, analysis of all swing and stance
phases against cycle period showed stance duration increased with cycle period,
whilst swing duration was constant (Fig. 3K). Therefore, swing and stance duty cycles,
defined as the proportion of time normalized to the cycle period, showed opposing
exponential relationships with cycle period (Fig. 3L).

Thoracic-abdominal coordination is crucial for stable, effective locomotion

To begin to understand the neural control of movement in *Tribolium* larvae, we severed
the connective between thoracic and abdominal ganglia (Fig. 4A,B) to assess the
effects of interrupting ascending and descending connections between brain and
suboesophageal ganglion (SOG) and ventral nerve cord (VNC) on unrestrained
locomotion. Severed animals travelled significantly less far (mean = 5.83 ± 0.58cm)
on paper than controls (mean = 13.3 ± 2.0cm, p = 0.020) and sham treated animals
(mean = 14.4 ± 2.0cm, p = .008; Fig. 4C,D). Moreover, they displayed a narrower
range of body bending angles (range = 115.8 to 224.1º) than controls (range = 122.7
to 245.4º) and shams (range = 111.0 to 273.2º) (Fig. 4E). To see how these
impairments affected performance during a locomotor challenge, we first performed
preliminary experiments gauging control larvae performance on different inclinations:
180º (flat surface), 135º (uphill), 90º (vertical wall), and 45º (overhang) (Fig. S1). We
found control animals walked progressively less far with greater deviations from 180º
but were still able to cling onto and climb 45º overhangs. Interestingly, larval paths
became straighter, and fewer animals approached arena edges with greater inclination
(Fig. S1A-D,F).

Next, we used inclines as a motor challenge for severed animals to investigate how
disconnecting the abdominal nervous system would affect clinging and climbing during
challenged locomotion. Control, sham, and severed animals were tested on 180º, 135º
and 45º. Severed animals exhibited prominent clinging and climbing impairments –
42.9% at 135º and 71.4% at 45º of larvae failed to cling on and fell (represented as
0mm s^{-1} points for velocity in Fig. 5B). Severed animals that could cling and climb

seemed to have straightening paths and had slower speeds with increasing incline, as was observed in control and sham treated animals (Fig. 5A-H, Fig. S1G). As expected, severed speeds were significantly lower for baseline 180° (control-severed, p < .001; sham-severed, p = .002) and 135° (control-severed, p < .001; sham-severed, p = .007). Although some severed animals had curved postural phenotypes, this was unrelated to their speed of locomotion (curved individuals ranked 3rd, 4th and 5th out of 7 total severed larvae in speed). Nonsignificant differences at 45° may reflect an overall speed limitation at such extreme inclinations.

To assess leg kinematics and coordination differences across treatments and inclinations, we observed locomotor performance on 180° and 45° inclines on an up-close incline behavioral rig as opposed to the top-down view (Movie 3). At 45°, 14.3% of severed animals fell, and 42.9% clung on but remained immobile. Consequently, these animals were excluded from locomotor and kinematic analysis. Notably, no control or sham animals fell at any inclination in either experiment. To effectively compare locomotion across conditions, walk cycles following the T3 to T1 gait sequence were analyzed. Severed animals were slower overall but followed locomotor parameter trends in control and sham for 180°. However, they were indistinguishable from controls and shams at 45° as overall locomotion slowed. Swing/stance durations and duty cycles followed the same trends with speed observed previously (Fig. 3K-L) independent of inclination; stance duration was modulated, whilst swing duration remained constant across speeds (Fig. 5I-J,M-N). Stride lengths increased linearly with speed at both 180° (r^2 = .754, p < .001; Fig. 5K) and 45° (r^2 = .387, p < .001; Fig. 5O), and intersegmental differences were proportional across treatments and inclinations (Fig. 5L,P).

At 45°, only control and sham larvae engaged their pygopods to assist climbing (Movie 3), which suggested climbing required descending inputs to the abdomen. Moreover, larvae consistently attempted to maintain abdominal contact with the substrate, suggesting sustained contraction of body wall muscles whilst climbing. Pygopod planting events were defined as visible pygopod extensions, and a change from a parallel alignment with the substrate to a perpendicular pointed orientation into the substrate (Fig. 5Q). Analysis of their relation to leg kinematics in all walk cycles showed that most pygopod planting events coincided temporally with the initiation of

leg swing phases (Fig. 5R), and pygopod planting-triggered analysis showed the
sequence from T3 (blue) to T1 (magenta) as seen by the peaks in the distributions
(Fig. 5R). Moreover, pygopod engagement was preceded by an increase in velocity
and followed by a decrease (Fig. 5S), consistent with the association between
pygopod planting events and leg swings. Although most control and sham larvae
engaged their pygopods on the overhang (mean rate = 0.76Hz, range = 0 to 1.20Hz),
this did not significantly correlate with their maximum velocity (r^2 = .123, p = .092) (Fig.
5T).

Abdominal inputs are essential for tunnelling

As evidenced, abdominal inputs (ascending or descending) facilitated free locomotion
on flat surfaces and during locomotor challenges. Although clinging and climbing
behavior may assist *Tribolium* in navigating and infesting food storage systems, we
wanted to probe the effects of severing thoracic-abdominal connectives on tunnelling
into their food source. Larvae (control, sham and severed) were individually dropped
onto a dish filled with approximately 10mm deep whole wheat flour and visible body
length was measured 0s and 120s afterwards (Fig. 6A-F). Control and sham animals
immediately tunneled into flour on contact, as observed previously during substrate
preference tests, with 100% of control and 75% of sham larvae being fully submerged
after 2 minutes (Fig. 6G). Strikingly, 0% of severed animals tunneled effectively into
flour, and on average, 90.7% of their original body lengths was exposed after 2 minutes
(Fig. 6H). Severed animals still visibly displaced flour with their legs, albeit without
making any progress (Fig. 6F). Therefore, the abdominal segments are essential for
larval submergence into flour.

Discussion

To study and understand the *Tribolium* larval abdomen's roles in locomotion and its
relationship to the legs, we first characterized the larva's general locomotor
performance across a range of conditions. We found larvae relied on their legs for
unchallenging locomotion on preferred substrates, which mostly consisted of fibrous
materials. This supports the hypothesis that their original habitat consisted of tree bark
(Dawson, 1977; Good, 1936), and explains why wet, slippery surfaces present a
challenge. While adults are capable of locomotion across flour (Pai, 2010), tunnelling

is the primary motor program for navigating their food source during pre-pupal instars.
This behavior reflects ethologically relevant photophobic preferences (Duehl et al.,
2011) and their propensity to tunnel deeper when approaching pupation (King and
Dawson, 1972). The locomotor patterns underlying their exploratory behavior consist
of straight runs and continuous turning, with little to no periods of immobility. This
differs from *Drosophila* larvae, which combine repeated straight runs with pauses
followed by headsweeps, redirecting them to a new trajectory (Lahiri et al., 2011;
Suster and Bate, 2002). *Tribolium* larvae also show no evidence of handedness in
their exploratory movements, again in contrast with *Drosophila* adults (Buchanan et
al., 2015) and larvae (Wosniack et al., 2022) but also other insects such as ants (Hunt
et al., 2014) and cockroaches (Camhi and Johnson, 1999). This may reflect the
requirements for *Tribolium* to navigate a 3-dimensional environment of constantly
shifting substrate. We also found a larval preference for edge exploration, which may
represent photophobicity, centrophobicity or a thigmotactic drive to maximize body
contact with their environment to mimic how they are naturally enveloped in substrate.
This behavior is also found in *Drosophila* adults, but not for centrophobic or
thigmotactic reasons (Soibam et al., 2012).

The most common gait in insects is the alternating tripod gait, in which the first and
third leg on one side of the body move together with the contralateral second leg
(Wilson, 1966). *Tribolium* larvae, however, use a travelling wave gait, involving
posterior-to-anterior sequences of swings propagating from T3 to T1 across all walking
speeds. Their contralateral leg pairs move synchronously, and phase delays between
segments are largely maintained across speeds. This bilaterally symmetric travelling
wave gait is found in the larvae of other beetles, including leaf beetles (Gustafson and
Chaboo, 2009; Zurek et al., 2015a). Since environmental conditions do not differ
between larval and adult stages in these animals, the wave gait could therefore reflect
developmental constraints on the circuit architecture driving locomotion in the
Cucujiformia infraorder of beetles. However, the travelling wave gait pattern is found
more widely. For example, in krill and crayfish, it generates the forces required for
locomotion in an aquatic medium (Mulloney and Smarandache-Wellmann, 2012;
Zhang et al., 2014), and in some millipedes it is used to generate the required forces
for burrowing in soil (Manton, 1961). A similar bilaterally symmetrical gait has been
observed in tardigrades, which is employed in a substrate-dependent manner.

Tardigrades more commonly use a tetrapod gait, but transition to a wave gait on softer
and therefore unpredictable surfaces (Nirody et al., 2021). Similarly, the tobacco
hornworm caterpillar, *Manduca sexta,* usually adopts a "progressive" posterior-to-
anterior wave gait during horizontal crawling on stiff substrates, like *Tribolium* larvae,
but transition to "non-progressive" gaits on soft substates (Metallo et al., 2020).
Another example is the dung beetle *Pachysoma endroedyi* 'galloping' gait, in which
the anterior two leg pairs move in a travelling wave, with the hind leg pair dragging
behind them and contributing little to forwards motion (Smolka et al., 2013). In these
desert dwelling species, in-phase contralateral coordination may be energetically
favorable for traversing constantly shifting terrains like sand, a constraint they might
share with aquatic species. These considerations are equally relevant for *Tribolium,*
as they consume and inhabit a constantly deformable substrate. This gait may become
particularly beneficial during *Tribolium* pre-pupal stages, when larvae tunnel deeper
for warmth and lower population densities to prepare for pupation (King and Dawson,
1972).

Gait adjustments tune movement to the requirements of different terrains and speeds
of locomotion (Bässler and Büschges, 1998; Bender et al., 2011; Graham, 1972;
Ramdya et al., 2017; Schilling et al., 2013; Smolka et al., 2013). Many animals display
speed-dependent gait transitions. For example, adult *Drosophila* gaits have been
shown to transition from a pentapodal wave (one leg is in swing phase at a time) to a
tetrapodal (two legs, but not bilateral pairs) and finally tripodal wave (three legs) by
modulating stance phases as the animal progresses from slow to fast speeds
(DeAngelis et al., 2019). We therefore also characterized the kinematic changes
accompanying different speeds of locomotion in *Tribolium* larvae. They similarly have
increasing numbers of limbs in swing phase at a time during increasingly fast
locomotion, but do not appear to adopt a gait other than their bilaterally symmetrical
metachronal wave. In addition, *Tribolium* larvae increase walking speeds by reducing
stance duration. This is a widely observed locomotor strategy used across phyla, from
humans (Grillner et al., 1979), to cats (Frigon et al., 2014), to *Drosophila* (DeAngelis
et al., 2019) and tardigrades (Nirody et al., 2021). Moreover, larval stride length
increases linearly with speed, also adhering to a highly conserved locomotor strategy
across invertebrates and vertebrates (Granatosky and McElroy, 2022; Hooper, 2012;

Strang and Steudel, 1990). A similar in-depth analysis of adult *Tribolium* gait
kinematics in the future will allow further comparison.

The *Tribolium* VNC is unfused, with a separate ganglion in each body segment,
chained together by longitudinal connectives, and each thoracic ganglion innervating
a leg pair (Farnworth et al., 2022a; Hunnekuhl et al., 2020). This anatomical
arrangement is found across arthropods (Richter et al., 2010) and tardigrades (Mayer
et al., 2013), and suggests gait variation across and within species might be derived
from one single modifiable control strategy (Nirody, 2021). This raises the question
whether the neural control of its behavior shares similarities with that of the more well-
studied and soft-bodied *Drosophila* larva, which is legless and moves by means of
abdominal peristaltic waves (Lahiri et al., 2011), and has a segmentally fused central
nervous system (Truman and Bate, 1988).

Qualitatively observed pygopod planting during unrestricted motor programs
(backtrack and redirect, see Fig. 2F) first suggested a prominent role for the abdomen
in assisting natural movement. We present several lines of evidence that pygopods
act to support locomotion in challenging conditions. First, while fast animals crawling
on level fibrous surfaces do not necessitate pygopod engagement, they do while
climbing 45º overhangs. Furthermore, pygopod planting events are largely temporally
locked to leg swing phases, during which the animal needs to maintain static stability,
and these events are preceded by a temporary increase in crawling speed. This shows
a correlation between situations that compromise the animal's stability and likelihood
of pygopod engagement. Furthermore, the weak, non-significant correlation between
pygopod planting rate and maximum velocity further implicates its role in stability over
a mechanism to boost/propel overall speed. Second, severing the A1-A2 connective,
thus removing descending and ascending inputs between the brain, thorax and
abdominal segments, resulted in lower speeds during walking and climbing. Some
severed animals dragged their abdomen during walking (Movie 3), which may
contribute to decreasing velocity as the lack of abdominal sensory inputs may impair
its ability to support itself during locomotion. However, impairments on inclines varied,
ranging from some severed animals falling off mild (135º) inclines to a select few
matching control and sham performance on 45º overhangs. The latter may reflect the
overhang limiting the feasible speeds larvae can produce in this motor challenge. Like

the variability in inter-animal velocities in normal larvae, the severity of severing is
likely equally dependent on the individual. Nevertheless, severed animals never
engaged their pygopods, and most tended to fall off 45º overhangs. These animals'
inability to engage their posterior segments therefore particularly impaired their
locomotor capabilities under challenging conditions. Furthermore, this phenotype is
unlikely due to a failure to coordinate leg movements, since severed animals fall into
the same speed-dependent distributions of kinematic patterns including swing and
stance duration, duty cycle, stride length and intrasegmental phase difference as
control and sham larvae. This suggests that abdominal regions are not required for
fundamental leg coordination but provide a crucial support role in larvae.

Other species that have abdominal anchoring points with potential stabilizing roles
include *Labidomera clivicollis* larvae, which use pygopods during body extension
whilst climbing tree branches (Gustafson and Chaboo, 2009); and blackfly larvae,
which are adapted to anchor into riverbeds to avoid getting swept away by strong
currents (Jamnback and Frempong-Boadu, 1966). The *Tribolium* larval anatomical
adaptations we describe that confer stability and allow the animal to stay upright may
similarly help evade predation, for example by the parasitoid wasp *Holepyris sylvanidis*
(Amante et al., 2018), either by enhancing locomotion efficiency in shifting terrain or
avoiding falling off tree branches in their original habitat. The use of posterior terminal
organs for substrate engagement during locomotion is also found in other soft-bodied
animals including *Drosophila* larvae (Booth et al., 2024), *Gastrophysa viridula*, which
deploy their pygopods phase-locked to leg movements during the locomotor cycle
(Zurek et al., 2015b); and *Manduca sexta*, whose locomotor cycle starts with the
terminal prolegs lifting off the substrate (van Griethuijsen and Trimmer, 2014). Our
results elucidate a functional role for the pygopods, which were previously only
phenotypically mentioned in the *iBeetle* RNA interference (RNAi) screens (Hakeemi et
al., 2022; Schmitt-Engel et al., 2015).

The contribution of abdominal segments is likely to extend beyond offering physical
support during locomotion. During motor challenges like overhangs, larvae may send
ascending information from abdominal sensory neurons (e.g., regarding reduced body
contact with substrate) to sensorimotor integration centers for sensory feedback.
These may then integrate and send back descending information regarding internal

representations (e.g., current geotactic location, motor state) to make a higher order
decision to plant the pygopods to increase stability. Interruption of communications to
and from the abdomen and thus lack of stability in severed animals may explain those
that fall off or fail to climb. However, the severed animals that could cling and climb
followed the same leg kinematics as controls and shams, therefore lack of abdominal
input appears to not disrupt thoracic movements.

Furthermore, our data suggest the posterior segments have prominent functions in
tunnelling alongside stability. Tunnelling is a key motor program for navigation,
feeding, development, and reproduction in their natural habitat, and the inability to
tunnel in severed larvae suggests a crucial role for abdominal contractions, pygopod
recruitment, and/or ascending sensory feedback in coordinating effective tunnelling.
The abdomen may be recruited to generate pushing forces to submerge larvae in flour,
similar to millipedes (Manton, 1961). Equally, it may aid exploratory movements such
as the backtrack and redirect behavior (Fig. 2F) to search for new food sources.
Investigating the structure of both ascending and descending inputs passing through
the larval *Tribolium* A1-A2 connective is therefore a promising avenue to explore the
neural basis of its behavior and could be a possible pinch point to disrupt the animal's
ability to infest grain stores.

Finally, *Tribolium* is an attractive emerging model system in behavioral neuroscience,
combining a robust experimental toolkit with an extensive body of work describing its
development and physiology. As the first coleopteran species to have a fully
sequenced genome (~16,500 genes), 47% of which are orthologous to insects and
vertebrates, *Tribolium* are already highly favored in the field of developmental genetics
(Richards et al., 2008). Their genetic toolkit includes the GAL4-UAS system (Schinko
et al., 2010), CRISPR/Cas9 genome editing (Adrianos et al., 2018; Gilles et al., 2015),
balancer chromosomes, and stable injectable RNAi gene silencing (Tomoyasu et al.,
2008). Forward genetic screens uncovered previously unknown genetic regulators of
leg and body wall muscle development (Schultheis et al., 2019), and recent studies
have described the structure and function of the *Tribolium* cryptonephridial system,
which is essential for the animal's exceptional tolerance to desiccation (Naseem et al.,
2023). The structures and locations of brain regions in larval and adult *Tribolium*
showed broad similarities to other insect model organisms (Farnworth et al., 2022b;

Garcia-Perez et al., 2021; He et al., 2019). In addition, comparison of *Tribolium* and *Drosophila* larval neuromuscular systems revealed resemblances in somatic muscle composition (Schultheis et al., 2019). Its similarities to both ancestral and highly derived insects therefore make it an attractive comparative model.

Overall, this study has emphasized how the kinematic analysis of movement and underlying neural control provides insights into natural behaviors that govern *Tribolium*'s success as a pest. We identified the importance of abdominal segments and pygopods to both locomotor challenges and tunnelling, suggesting that studying the structure and function of its connectives has important potential implications for behavioral neuroscience and food security. How far our findings extend to earlier instars and adults remains a promising question for future study.

Acknowledgements

We thank Barry Denholm for providing *Tribolium* stocks, Adam Taylor for *Groundswell*, Balwant Singh for building the up-close incline rig, Chris Sutherland for statistical advice and Pulver Lab and Zwart Lab members for constant support.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: B.X.Y., M.F.Z. and S.R.P.; Methodology: B.X.Y.; Investigation: B.X.Y.; Formal Analysis: B.X.Y. and M.F.Z.; Visualization: B.X.Y.; Writing: B.X.Y., M.F.Z. and S.R.P; Funding Acquisition: B.X.Y.; Supervision: M.F.Z. and S.R.P.

Funding

This work was supported by the Institute of Behavioural and Neural Sciences' Undergraduate Research Assistant Scheme to B.X.Y.

Data and resource availability

Raw experimental movies are available at the University of St Andrews Research Portal: doi:10.17630/a97be752-0c97-42a2-ada4-745c5acff342. Analyzed files are available at Zenodo: doi:10.5281/zenodo.11520087. Analysis code for gait and incline results are available upon request.

References

Adrianos, S., Lorenzen, M. and Oppert, B. (2018). Metabolic pathway interruption: CRISPR/Cas9-mediated knockout of tryptophan 2,3-dioxygenase in *Tribolium castaneum*. *J Insect Physiol* **107**, 104–109.

Amante, M., Russo, A., Schöller, M. and Steidle, J. L. M. (2018). Olfactory host location and host preference of Holepyris sylvanidis (Brèthes) (Hymenoptera: Bethylidae), a parasitoid of *Tribolium confusum* Jacquelin du Val and *T. castaneum* (Herbst) (Coleoptera: Tenebrionidae). *Journal of Stored Products Research* **78**, 105–109.

Bässler, U. and Büschges, A. (1998). Pattern generation for stick insect walking movements—multisensory control of a locomotor program. *Brain Research Reviews* **27**, 65–88.

Bender, J. A., Simpson, E. M., Tietz, B. R., Daltorio, K. A., Quinn, R. D. and Ritzmann, R. E. (2011). Kinematic and behavioral evidence for a distinction between trotting and ambling gaits in the cockroach Blaberus discoidalis. *Journal of Experimental Biology* **214**, 2057–2064.

Booth, J. H., Meek, A. T., Kronenberg, N. M., Pulver, S. R. and Gather, M. C. (2024). Optical mapping of ground reaction force dynamics in freely behaving *Drosophila melanogaster* larvae. *eLife* **12**,.

Buchanan, S. M., Kain, J. S. and De Bivort, B. L. (2015). Neuronal control of locomotor handedness in *Drosophila*. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 6700–6705.

Camhi, J. M. and Johnson, E. N. (1999). High-frequency steering maneuvers mediated by tactile cues: antennal wall-following in the cockroach. *Journal of Experimental Biology* **202**, 631–643.

Daley, M. A. (2018). Understanding the agility of running birds: Sensorimotor and
mechanical factors in avian bipedal locomotion. *Integr Comp Biol* **58**, 884–893.

Dawson, P. S. (1977). Life history strategy and evolutionary history of *Tribolium* flour
beetles. *Evolution* **31**, 226–229.

DeAngelis, B. D., Zavatone-Veth, J. A. and Clark, D. A. (2019). The manifold
structure of limb coordination in walking *Drosophila*. *eLife* **8**, e46409.

Devi, M. B. and Devi, N. V. (2015). Biology of rust-red flour beetle, *Tribolium*
castaneum (Herbst) (Tenebrionidae:Coleoptera). *Ind. Jour. of Entomol.* **77**, 81.

Digumarti, K. M., Trimmer, B., Conn, A. T. and Rossiter, J. (2019). Quantifying
dynamic shapes in soft morphologies. *Soft Robot* **6**, 733–744.

Duehl, A. J., Cohnstaedt, L. W., Arbogast, R. T. and Teal, P. E. A. (2011). Evaluating
Light Attraction to Increase Trap Efficiency for *Tribolium castaneum*
(Coleoptera: Tenebrionidae). *Journal of Economic Entomology* **104**, 1430–
1435.

Đukić, N., Andrić, G., Ninkovic, V., Pražić Golić, M., Kljajić, P. and Radonjić, A.
(2020). Behavioural responses of *Tribolium castaneum* (Herbst) to different
types of uninfested and infested feed. *Bulletin of Entomological Research* **110**,
550–557.

Đukić, N., Andrić, G., Glinwood, R., Ninkovic, V., Andjelković, B. and Radonjić,
A. (2021). The effect of 1-pentadecene on *Tribolium castaneum* behaviour:
Repellent or attractant? *Pest Management Science* **77**, 4034–4039.

Farnworth, M. S., Bucher, G. and Hartenstein, V. (2022a). An atlas of the developing
Tribolium castaneum brain reveals conservation in anatomy and divergence in
timing to *Drosophila melanogaster*. *Journal of Comparative Neurology* **530**,
2335–2371.

Farnworth, M. S., Bucher, G. and Hartenstein, V. (2022b). An atlas of the developing
Tribolium castaneum brain reveals conservation in anatomy and divergence in
timing to Drosophila melanogaster. *J Comp Neurol* 10.1002/cne.25335.

Frigon, A., D'Angelo, G., Thibaudier, Y., Hurteau, M.-F., Telonio, A., Kuczynski, V.
and Dambreville, C. (2014). Speed-dependent modulation of phase variations
on a step-by-step basis and its impact on the consistency of interlimb
coordination during quadrupedal locomotion in intact adult cats. *Journal of*
Neurophysiology **111**, 1885–1902.

Garcia-Perez, N. C., Bucher, G. and Buescher, M. (2021). Shaking hands is a
homeodomain transcription factor that controls axon outgrowth of central
complex neurons in the insect model *Tribolium. Development* **148**, dev199368.

Gerken, A. R., Scully, E. D. and Campbell, J. F. (2018). Red Flour Beetle
(Coleoptera: Tenebrionidae) Response to Volatile Cues Varies With Strain and
Behavioral Assay. *Environmental Entomology* **47**, 1252–1265.

Gilles, A. F., Schinko, J. B. and Averof, M. (2015). Efficient CRISPR-mediated gene
targeting and transgene replacement in the beetle *Tribolium castaneum*.
Development **142**, 2832–2839.

Good, N. E. (1936). The flour beetles of the Genus *Tribolium*.

Graham, D. (1972). A behavioural analysis of the temporal organisation of walking
movements in the 1st instar and adult stick insect *(Carausius morosus)*. *J.*
Comp. Physiol. **81**, 23–52.

Granatosky, M. C. and McElroy, E. J. (2022). Stride frequency or length? A
phylogenetic approach to understand how animals regulate locomotor speed.
J Exp Biol **225**, jeb243231.

Grillner, S., Halbertsma, J., Nilsson, J. and Thorstensson, A. (1979). The
adaptation to speed in human locomotion. *Brain Res* **165**, 177–182.

Gustafson, G. T. and Chaboo, C. S. (2009). Ambulatory use of ventro-lateral
abdominal ampullae by larvae of *Labidomera clivicollis* (Kirby) (Coleoptera:
Chrysomelidae: Chrysomelinae). *cole* **63**, 357–363.

Hakeemi, M. S., Ansari, S., Teuscher, M., Weißkopf, M., Großmann, D., Kessel, T.,
Dönitz, J., Siemanowski, J., Wan, X., Schultheis, D., et al. (2022). Screens

in fly and beetle reveal vastly divergent gene sets required for developmental
processes. *BMC Biology* **20**, 38.

He, B., Buescher, M., Farnworth, M. S., Strobl, F., Stelzer, E. H., Koniszewski, N.
D., Muehlen, D. and Bucher, G. (2019). An ancestral apical brain region
contributes to the central complex under the control of foxQ2 in the beetle
Tribolium. *eLife* **8**, e49065.

Hooper, S. L. (2012). Body size and the neural control of movement. *Current Biology*
22, R318–R322.

Hooper, S. L., Guschlbauer, C., Blümel, M., Rosenbaum, P., Gruhn, M., Akay, T.
and Büschges, A. (2009). Neural control of unloaded leg posture and of leg
swing in stick insect, cockroach, and mouse differs from that in larger animals.
J Neurosci **29**, 4109–4119.

Hunnekuhl, V. S., Siemanowski, J., Farnworth, M. S., He, B. and Bucher, G.
(2020). Immunohistochemistry and Fluorescent Whole Mount RNA In Situ
Hybridization in Larval and Adult Brains of *Tribolium*. *Methods Mol Biol* **2047**,
233–251.

Hunt, E. R., O'Shea-Wheller, T., Albery, G. F., Bridger, T. H., Gumn, M. and Franks,
N. R. (2014). Ants show a leftward turning bias when exploring unknown nest
sites. *Biol. Lett.* **10**, 20140945.

Jamnback, H. and Frempong-Boadu, J. (1966). Testing blackfly larvicides in the
laboratory and in streams. *Bull World Health Organ* **34**, 405–421.

King, C. E. and Dawson, P. S. (1972). Population biology and the *Tribolium* model.
Evol. Biol. **5**, 133–227.

Lahiri, S., Shen, K., Klein, M., Tang, A., Kane, E., Gershow, M., Garrity, P. and
Samuel, A. D. T. (2011). Two alternating motor programs drive navigation in
Drosophila larva. *PLOS ONE* **6**, e23180.

Leiras, R., Cregg, J. M. and Kiehn, O. (2022). Brainstem circuits for locomotion.
Annu Rev Neurosci **45**, 63–85.

Manton, S. M. (1961). The evolution of Arthropodan locomotory mechanisms. Part 71.
Functional requirements and body design in Colobognatha (diplopoda),
together with a comparative account of diplopod burrowing techniques, trunk
musculature and segmentation. *Journal of the Linnean Society of London,*
Zoology **44**, 383–462.

Mantziaris, C., Bockemühl, T. and Büschges, A. (2020). Central pattern generating
networks in insect locomotion. *Dev Neurobiol* **80**, 16–30.

Mayer, G., Martin, C., Rüdiger, J., Kauschke, S., Stevenson, P. A., Poprawa, I.,
Hohberg, K., Schill, R. O., Pflüger, H.-J. and Schlegel, M. (2013). Selective
neuronal staining in tardigrades and onychophorans provides insights into the
evolution of segmental ganglia in panarthropods. *BMC Evol Biol* **13**, 230.

Metallo, C., Mukherjee, R. and Trimmer, B. A. (2020). Stepping pattern changes in
the caterpillar Manduca sexta: the effects of orientation and substrate. *Journal*
of Experimental Biology **223**, jeb220319.

Mulloney, B. and Smarandache-Wellmann, C. (2012). Neurobiology of the
crustacean swimmeret system. *Prog Neurobiol* **96**, 242–267.

Naseem, M. T., Beaven, R., Koyama, T., Naz, S., Su, S., Leader, D. P., A. Klaerke,
D., Calloe, K., Denholm, B. and Halberg, K. V. (2023). NHA1 is a
cation/proton antiporter essential for the water-conserving functions of the
rectal complex in *Tribolium castaneum. Proceedings of the National Academy*
of Sciences **120**, e2217084120.

Nirody, J. A. (2021). Universal features in panarthropod inter-limb coordination during
forward walking. *Integrative and Comparative Biology* **61**, 710–722.

Nirody, J. A., Duran, L. A., Johnston, D. and Cohen, D. J. (2021). Tardigrades
exhibit robust interlimb coordination across walking speeds and terrains.
Proceedings of the National Academy of Sciences **118**, e2107289118.

Pai, A. (2010). *Tribolium*. In *Encyclopedia of Animal Behavior*, pp. 446–452. Elsevier.

Phillips, T. W. and Throne, J. E. (2010). Biorational approaches to managing stored-
product insects. *Annual Review of Entomology* **55**, 375–397.

Picker, M., Colville, J. F. and Burrows, M. (2012). A cockroach that jumps. *Biol Lett*
8, 390–392.

Prakashini, V. Y., Srinivasan, G., Shanthi, M. and Prabakaran, K. (2023).
Preference of Red Flour Beetle *Tribolium castaneum* (Herbst) Towards Colour
Cues. *Indian Journal of Entomology* 445–447.

Ramdya, P., Thandiackal, R., Cherney, R., Asselborn, T., Benton, R., Ijspeert, A.
J. and Floreano, D. (2017). Climbing favours the tripod gait over alternative
faster insect gaits. *Nat Commun* **8**, 14494.

Richards, S., Gibbs, R. A., Weinstock, G. M., Brown, S. J., Denell, R., Beeman, R.
W., Gibbs, R., Beeman, R. W., Brown, S. J., Bucher, G., et al. (2008). The
genome of the model beetle and pest *Tribolium castaneum*. *Nature* **452**, 949–
955.

Richter, S., Loesel, R., Purschke, G., Schmidt-Rhaesa, A., Scholtz, G., Stach, T.,
Vogt, L., Wanninger, A., Brenneis, G., Döring, C., et al. (2010). Invertebrate
neurophylogeny: suggested terms and definitions for a neuroanatomical
glossary. *Front Zool* **7**, 29.

Risse, B., Otto, N., Berh, D., Jiang, X. and Klämbt, C. (2014). FIM imaging and
FIMtrack: Two new tools allowing high-throughput and cost effective locomotion
analysis. *J Vis Exp* 52207.

Romero, S. A., Campbell, J. F., Nechols, J. R. and With, K. A. (2010). Movement
Behavior of Red Flour Beetle: Response to Habitat Cues and Patch
Boundaries. *Environmental Entomology* **39**, 919–929.

Schilling, M., Hoinville, T., Schmitz, J. and Cruse, H. (2013). Walknet, a bio-inspired
controller for hexapod walking. *Biol Cybern* **107**, 397–419.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch,
T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an
open-source platform for biological-image analysis. *Nat Methods* **9**, 676–682.

Schinko, J. B., Weber, M., Viktorinova, I., Kiupakis, A., Averof, M., Klingler, M.,
Wimmer, E. A. and Bucher, G. (2010). Functionality of the GAL4/UAS system
in *Tribolium* requires the use of endogenous core promoters. *BMC*
Developmental Biology **10**, 53.

Schmitt-Engel, C., Schultheis, D., Schwirz, J., Ströhlein, N., Troelenberg, N.,
Majumdar, U., Dao, V. A., Grossmann, D., Richter, T., Tech, M., et al. (2015).
The iBeetle large-scale RNAi screen reveals gene functions for insect
development and physiology. *Nat Commun* **6**, 7822.

Schultheis, D., Weißkopf, M., Schaub, C., Ansari, S., Dao, V. A., Grossmann, D.,
Majumdar, U., Hakeemi, M. S., Troelenberg, N., Richter, T., et al. (2019). A
large scale systemic RNAi screen in the red flour beetle *Tribolium castaneum*
identifies novel genes involved in insect muscle development. *G3*
Genes|Genomes|Genetics **9**, 1009–1026.

Semeao, A. A., Campbell, J. F., Whitworth, R. J. and Sloderbeck, P. E. (2011).
Response of *Tribolium castaneum* and *Tribolium confusum* adults to vertical
black shapes and its potential to improve trap capture. *Journal of Stored*
Products Research **47**, 88–94.

Shield, S., Jericevich, R., Patel, A. and Jusufi, A. (2021). Tails, flails, and sails: How
appendages improve terrestrial maneuverability by improving stability.
Integrative and Comparative Biology **61**, 506–520.

Smolka, J., Byrne, M. J., Scholtz, C. H. and Dacke, M. (2013). A new galloping gait
in an insect. *Current Biology* **23**, R913–R915.

Soibam, B., Mann, M., Liu, L., Tran, J., Lobaina, M., Kang, Y. Y., Gunaratne, G. H.,
Pletcher, S. and Roman, G. (2012). Open-field arena boundary is a primary
object of exploration for Drosophila. *Brain Behav* **2**, 97–108.

Strang, K. T. and Steudel, K. (1990). Explaining the scaling of transport costs: the
role of stride frequency and stride length. *Journal of Zoology* **221**, 343–358.

Suster, M. L. and Bate, M. (2002). Embryonic assembly of a central pattern generator
without sensory input. *Nature* **416**, 174–178.

Tomoyasu, Y., Miller, S. C., Tomita, S., Schoppmeier, M., Grossmann, D. and
Bucher, G. (2008). Exploring systemic RNA interference in insects: a genome-
wide survey for RNAi genes in *Tribolium* . *Genome Biology* **9**, R10.

Triphan, T., Poeck, B., Neuser, K. and Strauss, R. (2010). Visual targeting of motor
actions in climbing *Drosophila*. *Curr Biol* **20**, 663–668.

Truman, J. W. and Bate, M. (1988). Spatial and temporal patterns of neurogenesis in
the central nervous system of *Drosophila melanogaster*. *Dev Biol* **125**, 145–
157.

van Griethuijsen, L. I. and Trimmer, B. A. (2014). Locomotion in caterpillars. *Biol*
Rev Camb Philos Soc **89**, 656–670.

Wilson, D. M. (1966). Insect walking. *Annu Rev Entomol* **11**, 103–122.

Wosniack, M. E., Festa, D., Hu, N., Gjorgjieva, J. and Berni, J. (2022). Adaptation
of *Drosophila* larva foraging in response to changes in food resources. *Elife* **11**,
e75826.

Zhang, C., Guy, R. D., Mulloney, B., Zhang, Q. and Lewis, T. J. (2014). Neural
mechanism of optimal limb coordination in crustacean swimming. *Proc. Natl.*
Acad. Sci. U.S.A. **111**, 13840–13845.

Zurek, D. B., Gorb, S. N. and Voigt, D. (2015a). Locomotion and attachment of leaf
beetle larvae Gastrophysa viridula (Coleoptera, Chrysomelidae). *Interface*
Focus **5**, 20140055.

Zurek, A. J., Witczak, S., Dulinski, M., Wachniew, P., Rozanski, K., Kania, J.,
Postawa, A., Karczewski, J. and Moscicki, W. J. (2015b). Quantification of
anthropogenic impact on groundwater-dependent terrestrial ecosystem using

geochemical and isotope tools combined with 3-D flow and transport modelling.
Hydrology and Earth System Sciences **19**, 1015–1033.

Xu Ying, B., Zwart, M.F. and Pulver, S.R. (2024) Context-dependent coordination of
movement in *Tribolium castaneum* larvae. Dataset. *University of St Andrews*
Research Portal. https://doi.org/10.17630/a97be752-0c97-42a2-ada4-
745c5acff342.

Xu Ying, B., Pulver, S. R., & Zwart, M. F. (2024). Data for Xu Ying et al., 2024.
Zenodo. https://doi.org/10.5281/zenodo.11520088.

Figure legends

Figure 1: *Tribolium castaneum* larval locomotion is most efficient on fibrous
substrates. (A) Diagrammatic representation of *Tribolium castaneum* life cycle. (B)
Morphology of 4th-5th instar larva under a 10x objective. Full body image at 2x
magnification and insets at 5x magnification showing three thoracic leg pairs, nine
abdominal segments, pygopods (white arrow) and urogomphi (red arrow). Scale bars
= 1mm. (C) Larval paths over select substrates during first 15 seconds of free
locomotion. Scale bar = 10mm. (D) Mean instantaneous velocity across fifteen
substrates, ordered highest to lowest. (E) As (D), for total distance travelled. White
squares = mean, boxes = mean ±SE, whiskers = ±SD overlayed on raw data. N = 10.

Figure 2: Larvae display inter- and intra-animal variability in free locomotion
parameters. (A) Larval locomotor paths across white paper for 2 minutes. Colored
paths represent fast (blue), intermediate (yellow) and slow (red) larvae. Example
larvae were selected based on highest, lowest and intermediate mean instantaneous
velocity out of 30 larvae. Scale bar = 5cm. (B) Violin plot showing intra-animal
variability in instantaneous velocity. White squares = mean. (C) Instantaneous velocity
traces across 2 minutes of colored animals in (A) highlighting inter- and intra-animal
variability. (D) Proportion of time spent in distinct locomotor behaviors: straight = 180°
body angle and >0mm s^{-1} velocity, bent = non-180° body angle and >0mm s^{-1} velocity,
still = 0mm s^{-1} velocity regardless of body angle. Whiskers = mean ±SE, n = 30. (E)
Bottom: total left and right headsweep events by individual larvae. Left headsweep =
≥210° body angle, right headsweep = ≤150° body angle as defined by FIMTrack. Top:

histogram of headsweep directional difference (left minus right headsweep count);
intercept-only GLM, p = .869 for hypothesis that left headsweep probability out of total
is significantly different from 0. (F) 3 frame sequence during a backtrack and redirect
behavior. Frames are color-coded for total time (3.73s); temporal larval trajectory is
overlayed beneath current frame (white). Note the center frame showing more
posterior body position relative to preceding frames, indicative of "backtracking". Scale
bar = 5mm. (G) Instantaneous velocity and bodybending angle against time during a
single backtrack and redirect behavior sequence. Gray area highlighting backwards
locomotion. (H) Total number of backtrack and redirect events during two minutes of
free locomotion. N = 30.

Figure 3: Gait coordination and leg kinematic coherence are largely speed
dependent. (A) Left: example top-down view of *Tribolium* larva with labelled legs
across three thoracic segments: T3 (blue), T2 (green) and T1 (magenta). Scale bar =
1mm. Right: select frames of tracked leg positions showing progression from T3 to T1
in one representative walk cycle. Note the bilateral symmetry as intrasegmental leg
pairs swing. Scale bar = 1mm. (B,C) Example velocity traces of legs for (B) fast and
(C) slow animals across three walk cycles. Note the greater intersegmental
coordination in (B) than (C). (D,E) Swing (black) and stance (white) phases in the
same (D) fast and (E) slow animals in (B,C). (F,G) Polar plots of coherence phase and
magnitude of waveforms in (B,C) using T1L as anchor/0° for (F) fast and (G) slow
animals. Negative phase values temporally precede positive ones. Dark grey dotted
inner circle indicates α = 0.05 for coherence magnitude. (H) Scatter of phase difference
between left legs of each segment and T3L across fundamental frequencies (proxy for
walking speed). (I) Scatter of phase difference between legs of each segment. (J)
Percentage of significantly coherent legs (out of 5) to anchor (T1L) per animal. (K)
Mean swing and stance durations and (L) duty cycles for each segment (color-coded)
across 3-5 walk cycles as a function of mean cycle period. N = 10.

Figure 4: Severing thoracic-abdominal connectives impairs locomotion and
restricts bending. (A,B) Surgical manipulations for (A) sham: lateral lesion between
A1-A2 and (B) severed: ventral severing of the A1-A2 connectives. (C) Walk paths on
paper across treatments for 1 minute. Scale bar = 2cm. (D) Mean instantaneous
velocity across treatments. Boxes = mean ±SE, whiskers = ±SD; * p < .05, One-way

ANOVA, n = 6. (E) Frequency plot of all body angles throughout 1 minute of locomotion

at 15fps, binned every 10º. Note the exponential y axis (log10).

Figure 5: Abdominal inputs are not required for thoracic coordination and

pygopods provide stability in locomotion. (A) Walk paths of control, sham, and

severed larvae on top-down incline rig. Scale bar = 2cm. (B) Mean instantaneous

velocity across treatments and inclinations. Boxes = mean ±SE, whiskers = ±SD; ***

p < .001, ** p < .01, One-way ANOVA, n = 7. (C-H) Representative swing (black) and

stance (white) phases of walk cycles following T3 to T1 sequence on up-close incline

rig. Note the prolonged stance phases in severed at baseline 180º. (I-P) Scatter plots

comparing locomotor parameters across surgical manipulations for 180º and 45º: (I,M)

swing (filled shapes) and stance (empty shapes) duration, (J,N) swing and stance duty

cycle, (K,O) stride length and (L,P) time difference to T3 (intersegmental difference).

All values are averaged per segment (squares, circles, triangles); velocity is averaged

per animal. (K,O) Stride length correlates significantly with velocity at 180º (r^2 = .754,

p < .001), and 45º (r^2 = .387, p < .001), pooled across all animals. (Q) Frames showing

pygopod relaxation parallel to substrate (black arrow) and planting into substrate

(white arrow). Scale bar = 1mm. (R) Histogram of normalized pygopod planting event

count relative to initiation of closest leg swing across all walk cycles, binned according

to the Freedman–Diaconis rule (see Methods). Note the order of distribution peaks

highlights predominant T3 to T1 gait even in unfiltered walk cycles. Total number of

pygopod plant events analyzed = 117 pooled across 13 larvae. (S) Normalized mean

instantaneous velocity (black trace) relative to pygopod planting events (dotted line);

individual velocities across 13 larvae (control and sham combined) in grey. (T)

Pygopod plant rate correlates insignificantly and weakly with maximum velocity (r^2 =

.123, p = .092), pooled across all animals. N = 7 per treatment.

Figure 6: Abdominal inputs are crucial for larval tunnelling into natural food

source. (A-F) Top-down frames at 0s and 120s after placement on flour for (A,B)

control, (C,D) sham and (E,F) severed animals. Severed animals struggle to tunnel;

black arrow in (F) indicates displacement of flour from thoracic leg movements despite

no body submergence. Scale bar = 2mm. (G) Change in raw body length between 0s

and 120s across treatments. (H) Percentage of original body length exposed after

120s. Whiskers = mean ±SE; *** $p < .001$, ** $p < .01$, Kruskal-Wallis test. N = 8 per

treatment.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure S1: Increasing incline becomes progressive locomotor challenge for *Tribolium* larvae.
(A-D) Left: diagrams of larval orientations relative to angled platforms (white rectangles). Right: walk paths of larvae on respective platforms. Note, 45º is an overhang. Asterisks indicate starting position. (E) Mean instantaneous velocity across inclinations. Boxes = mean ±SE, whiskers = SD; *** $p<.001$, * $p<.05$, One-way ANOVA, n = 10. (F) Path curviness across inclinations calculated via (total distance / displacement) * max distance from start for preliminary control trials in (A-E), n = 10. (G) Path curviness for experimental trials in Fig. 5A,B across inclinations and surgical manipulations, n = 7.