Context-dependent coordination of movement in *Tribolium castaneum* larvae

Authors

Bella Xu Ying[1,2,3] zxy1@st-andrews.ac.uk

Maarten Frans Zwart[1,2,3]* mfz@st-andrews.ac.uk

Stefan Robert Pulver[1,2,3]* sp96@st-andrews.ac.uk

*Corresponding authors

[1]Institute of Behavioural and Neural Sciences

[2]Centre of Biophotonics

[3]School of Psychology and Neuroscience, University of St Andrews, St Andrews, KY16 9JP, United Kingdom

Abstract

Stored product insect pests, like *Tribolium castaneum* beetles, cause 20% of postharvest loss. However, how their nervous systems coordinate adaptive movements for successful infestation is unknown. Here, we assess how *Tribolium* larvae locomote over different substrates and analyze their gait kinematics across speeds. Unlike many hexapods, larvae employ a bilaterally symmetric, posterior-to-anterior wave gait during fast locomotion. During slow locomotion, thoracic intrasegmental coordination is disrupted, whilst intersegmental coordination is preserved. Additionally, terminal abdominal structures (pygopods) support challenging locomotion, such as climbing overhangs. The onset of pygopod engagement coincides with leg swing initiation, suggesting a stabilizing role. Surgically severing the connective between thoracic and abdominal ganglia impaired pygopod engagement and impeded flat-surface locomotion, climbing, and tunnelling without interrupting leg kinematics. These results suggest that thoracic-abdominal coordination underlies effective movement, and gait/limb recruitment is context-dependent. Our work provides the first kinematic analysis of *Tribolium* larval locomotion and insights into its neural control.

Introduction

Stored product insects are devastating agricultural pests. These animals have been estimated to cause 9% of postharvest loss in developed countries and 20% in developing countries (reviewed in[1]). The red flour beetle, *Tribolium castaneum (Tribolium),* is representative of many stored product pest insects presenting a major challenge to global food security. Larvae and adults consume wheat products like flour by efficiently moving over and tunnelling through these unstable, shifting substrates[2]. Therefore, understanding how *Tribolium* adapt their movements to heterogeneous and challenging terrain could provide insights into conserved principles of motor control whilst aiding the development of new pest control approaches.

Animals use various strategies for adapting movement to the environments in which they live, continuously monitoring and adjusting movements on a cycle-by-cycle basis to cope with unexpected obstacles and/or drastic variations in the physical properties of 3-dimensional terrain. Studying the neural control of movement adaptability has

been a long-standing focus of motor systems research across multiple species.
Terrestrial animals with internal, external, or hydrostatic skeletons have all evolved
different strategies for movement control in different contexts (reviewed in insects[3,4],
mammals[5] and birds[6]). Some animals deploy limbs[7–9], some utilize soft outer cuticular
features[10,11] and others adapt body parts not typically involved in locomotion to solve
motor problems[12]. Similarly, *Tribolium* exhibit a diverse behavioral repertoire in
response to different sensorimotor cues, and adult anatomy has been well-described.
Previous work has shown that *Tribolium* can move through a variety of stored grain,
but prefer finely grained, whole wheat flour over coarser bran[13]. Moreover, adults can
navigate in response to attractive and aversive odors[14,15], shapes[16], light levels[17],
colors[18], and different habitat cues[19]. Both larvae and adults have articulated, jointed
limbs[2], with larvae being composed of both soft ventral cuticle and semi-hardened
dorsal cuticle. While anatomical studies have begun to characterize the larval
neuromuscular system[20], knowledge concerning the specific kinematics of locomotion
and how the *Tribolium* nervous system recruits thoracic limbs and abdominal
segments to meet different locomotor demands is extremely limited.

Past studies have prioritized adult anatomy and behavior, although larval instars are
equally capable of maneuvering and colonizing stored grain. Here we characterize
locomotor parameters in late-instar *Tribolium* larvae to gain insights into how these
animals adapt movements to their environment. We analyze how larvae negotiate
different substrates and combine whole-animal and leg tracking to characterize and
quantify leg kinematics underlying a range of locomotor speeds. We find larvae have
an ethologically relevant preference for fibrous substrates and walk primarily with a
posterior-to-anterior-propagating bilaterally symmetric gait, which shifts towards
uncoordinated exploratory movements at slower speeds. Severing the connectives
between thoracic and abdominal segments impairs walking, climbing, and tunneling
but does not disrupt fundamental leg kinematics. In addition, we characterize a
pygopod planting behavior, abolished in animals with severed connectives, which is
used for stability and propulsion under different motor contexts (challenged locomotion
vs tunnelling). These results provide first insights into the biomechanical
underpinnings of *Tribolium* movement and the neural basis of ethologically relevant
motor programs in an insect species with direct impacts on global food security.

Results

Tribolium castaneum life cycle, morphology, and substrate preference

Tribolium beetles take approximately 35 days at 29ºC to undergo complete metamorphosis, comprising four distinct developmental stages – egg, larval, pupal, and adult[21] (Fig. 1a). The larval stage is characterized by three thoracic segments, each supported by a pair of legs, and nine abdominal segments. Larvae possess two small, rounded ventral pygopods in the most posterior abdominal segment, and two spiked urogomphi dorsally (Fig. 1b).

We first established larval substrate preference based on ease of locomotion. We found larvae crawl best on textured, fibrous, wood-based substrates such as paper, cardboard, and lens paper (Fig. 1c). In contrast, they had substantial difficulty gripping smooth surfaces like plastic and glass, thus preventing forwards momentum. Similarly, their legs slipped on or stuck to deformable substrates like Sylgard™ and agarose, also impeding coordinated progress (Fig. 1d-e). Notably, on these substrates, we observed pygopod planting for substrate engagement in an anchoring motion, which, when released, propelled the larvae forwards (Supplementary Movie 1). This abdominal engagement was rarely observed on fibrous substrates. Surprisingly, crawling was heavily impeded over whole wheat flour and white flour (Fig. 1c-e), as larvae attempted to tunnel upon contact with these substrates.

Figure 1: *Tribolium castaneum* life stages, larval morphology, and locomotor capability over different substrates. (a) Diagrammatic representation of *Tribolium castaneum* life cycle. **(b)** Morphology of 4th-5th instar larva under a 10x objective. Full body image at 2x magnification and insets at 5x magnification showing three thoracic leg pairs, nine abdominal segments, pygopods (white arrow) and urogomphi (red arrow). Scale bars = 1mm. **(c)** Larval paths over select substrates during first 15 seconds of free locomotion. Scale bar = 10mm. **(d)** Box plot of mean instantaneous velocity across fifteen substrates, ordered highest to lowest. **(e)** As **(d)**, for total distance travelled. White squares = mean, boxes = mean ±SE, whiskers = ±SD overlayed on raw data; n = 10.

Quantifying larval locomotor parameters

Next, movement parameters during free locomotion were analyzed to quantify the spectrum of larval motor capabilities. Paper was used as the default substrate in subsequent experiments given larval preference, accessibility, and ease of imaging. *Tribolium* larvae exhibited heterogeneous walking patterns, and favored edge exploration, i.e., they rarely strayed from arena edges once they found them (Fig. 2a). Additionally, we observed persistent burrowing attempts at the paper edges during preliminary trials in which the paper substrate was not fully adhered to the arena platform.

Locomotion on paper had a mean instantaneous velocity of 1.8 ± 0.1 mm/s (range = 0.9 to 3.0 mm/s; larval sizes ranged 4-6 mm), and animals displayed a spectrum of intra-animal velocity variability during only 2 minutes of exploration (Fig. 2b). Faster animals were not constantly fast, but instead varied speed throughout locomotion, whereas slower animals were limited to lower speed ranges (Fig. 2c). Larvae spent equal time walking straight (49.6%) and turning (50.4%) (Fig. 2d). Animals decelerated to turn but rarely stopped completely, thus staying immobile only 0.5% of the time (Fig. 2d). Finally, most larvae did not show a bias in turning direction, engaging equally in left and right headsweeps during exploration (Fig. 2e).

We observed a larval motor program we termed "backtrack and redirect" (Supplementary Movie 2), defined as a headsweep/turn in one direction, followed by visible backwards locomotion, and a contralateral headsweep/turn and forwards locomotion (Fig. 2f), which on average lasted 3-6 seconds. This behavior followed a distinct sequence of instantaneous velocity and body angle changes. First, velocity dropped as the animal turned (body angle deviating from 180º), followed by an increase in velocity at 180º body angle due to backtracking, and ended with another deceleration and deviation from 180º in the opposite direction from the animal's original trajectory (Fig. 2g). While 63.3% of larvae displayed this motor program at least once during 2 minutes of free locomotion (Fig. 2h), some displayed it a maximum of 8 times, and a third never engaged in this behavior.

Figure 2: Inter- and intra-animal variability in free locomotion parameters. (a) Select larvae's paths of free locomotion across white paper for 2 minutes. Colored paths represent fast (blue), intermediate (yellow) and slow (red) larvae based on mean instantaneous velocity. Scale bar = 5cm. **(b)** Violin plot showing intra-animal variability in instantaneous velocity. White squares = mean. **(c)** Instantaneous velocity traces across 2 minutes of colored animals in **(a)** highlighting inter- and intra-animal variability. **(d)** Proportion of time spent in distinct locomotor behaviors: straight = 180° body angle and >0mm/s velocity, bent = non-180° body angle and >0mm/s velocity, still = 0mm/s velocity regardless of body angle. Whiskers = mean ±SE, n = 30. **(e)** Total headsweep events by individual larvae. Left headsweep = ≥210° body angle, right headsweep = ≤150° body angle. **(f)** 3 frame sequence during a backtrack and redirect behavior. Frames are color-coded for total time (3.73s); temporal larval trajectory is overlayed beneath current frame (white). Note the center frame showing more posterior body position relative to preceding frames, indicative of "backtracking". Scale bar = 5mm. **(g)** Instantaneous

velocity and bodybending angle against time during backtrack and redirect behavior sequence. Gray area highlighting backwards locomotion. **(h)** Total number of backtrack and redirect events during two minutes of free locomotion, n = 30.

Gait coordination and leg kinematics across walking speeds

To characterize natural gait, leg kinematics and the coordination between thoracic segments, we tracked legs during straight walking bouts on paper. We found *Tribolium* larvae walked with a bilaterally symmetric posterior-to-anterior wave gait (Fig. 3a). Then, we determined the frequency of locomotion for each animal by performing a Fourier transform on leg velocities and selecting the frequency with highest power of an anchor leg (left T1). We defined swing phases by a leg's lifting and planting on the substrate and stance phases as the duration in which legs are in contact with the ground. An average cycle period was then calculated for each leg. After a swing, legs usually stanced next to the legs of the segment in front, and these often began their swing phase before the previous contacted the ground (Fig. 3a). This pattern was observed most prominently within the T2 to T1 transition as their swing phases overlapped more visibly (Fig. 3b-d).

Fast (Fig. 3b, d, f) and slow (Fig. 3c, e, g) walk cycles differed markedly. In faster animals, typical walk cycles averaged 0.2 ± 0.007s and were initiated by a T3 swing phase, followed by T2 with a phase delay of approximately 120°, then T1 with another 120° delay. These delays were largely consistent between legs and across animals, independent of speed (Fig. 3h). In fast animals, many legs were significantly coherent with each other (Fig. 3j). In slower animals, average cycle period was 0.7 ± 0.1s, and some legs often swung alone during walk cycles (Fig. 3e). Furthermore, not all six legs were required to swing during each cycle. Absent leg swings consequently impaired intrasegmental coordination; phase differences between contralateral leg pairs increased (binned mean for 5 slowest larvae = 41.0 ± 7.5°) compared to faster animals (binned mean for 5 fastest larvae = 15.1 ± 3.4°) (Fig. 3i). As such, fewer legs were significantly coherent to each other in slower animals (Fig. 3j), but the T3 to T1 sequence was largely preserved (Fig. 3h). Finally, analysis of all swing and stance phases against cycle period showed stance duration increased with cycle period, whilst swing duration was constant (Fig. 3k). Therefore, swing and stance duty cycles,

defined as the proportion of time normalized to the cycle period, showed opposing exponential relationships with cycle period (Fig. 3l).

Figure 3: Gait coordination and leg kinematic coherence are largely speed dependent.
(a) Left: example top-down view of *Tribolium* larva with labelled legs across three thoracic segments: T3 (blue), T2 (green) and T1 (magenta). Scale bar = 1mm. Right: select frames of tracked leg positions showing progression from T3 to T1 in one representative walk cycle. Note the bilateral symmetry as leg pairs swing. Scale bar = 1mm. **(b-c)** Example velocity traces of legs for **(b)** fast and **(c)** slow animals across three walk cycles. Note the greater intersegmental coordination in **(b)** than **(c)**. **(d-e)** Swing (black) and stance (white) phases in the same **(d)** fast and **(e)** slow animals in **(b-c)**. **(f-g)** Polar plots of coherence phase and magnitude of waveforms in **(b-c)** using T1L as anchor/0° for **(f)** fast and **(g)** slow animals. Negative phase values temporally precede positive ones. Dark grey dotted inner circle indicates α = 0.05 for coherence magnitude. **(h)** Scatter of phase difference between left legs of each segment and T3 across fundamental frequencies (proxy for walking speed). **(i)** Scatter of phase difference between legs of each segment. **(j)** Percentage of significantly coherent legs (out of 5) to anchor (T1L). **(k)** Mean swing and stance durations and **(l)** duty cycles for individual legs as a function of cycle period averaged across 3-5 tracked cycles per leg. N = 10.

Effects of severing thoracic-abdominal connectives on locomotion and climbing

To begin to understand the neural control of movement in *Tribolium* larvae, we severed the connective between thoracic and abdominal ganglia (Fig. 4a-b) to assess the effects of interrupting ascending and descending connections between brain and suboesophageal ganglion (SOG) and ventral nerve cord (VNC) on unrestrained locomotion. Severed animals travelled significantly less far (mean = 5.83 ± 0.58cm) on paper than controls (mean = 13.3 ± 2.0cm, $p = 0.020$) and sham treated animals (mean = 14.4 ± 2.0cm, $p = 0.008$; Fig. 4c-d). Moreover, they displayed a narrower range of body bending angles (range = 115.8 to 224.1º) than controls (range = 122.7 to 245.4º) and shams (range = 111.0 to 273.2º) (Fig. 4e). To see how these impairments affected performance during a locomotor challenge, we first performed preliminary experiments gauging control larvae performance on different inclinations (Supplementary Figure 1): 180º (flat surface), 135º (uphill), 90º (vertical wall), and 45º (overhang). We found control animals walked progressively less far with greater deviations from 180º but were still able to cling onto and climb 45º overhangs. Interestingly, larval paths became straighter, and fewer animals approached arena edges with greater inclination.

Figure 4: Severing thoracic-abdominal connectives impairs locomotion and restricts bending. (a-b) Surgical manipulations for **(a)** sham: lateral lesion between A1-A2 and **(b)** severed: ventral severing of the A1-A2 connectives. **(c)** Walk paths on paper across treatments for 1 minute. Scale bar = 2cm. **(d)** Box plots of mean instantaneous velocity. Boxes = mean ±SE, whiskers = ±SD; * p<.05, One-way ANOVA, n = 6. **(e)** Frequency plot of all body angles throughout 1 minute of locomotion, binned every 10º. Note the exponential y axis (log10).

Supplementary Figure 1: Increasing incline becomes progressive locomotor challenge for *Tribolium* larvae. (a-d) To the left, diagrams of larval orientations relative to angled platforms and to the right, walk paths of larvae at varying inclines. Note 45º is an overhang. Asterisks indicate starting position. **(e)** Box plot of mean instantaneous velocity across inclinations. Boxes = mean ±SE, whiskers = SD; *** p<.001, * p<.05, One-way ANOVA, n = 10.

Next, we used inclines as a motor challenge for severed animals to investigate how disconnecting the abdominal nervous system would affect clinging and climbing during challenged locomotion. Control, sham, and severed animals were tested on 180º, 135º and 45º. Severed animals exhibited prominent clinging and climbing impairments – 42.9% at 135º and 71.4% at 45º of larvae failed to cling on and fell (represented as 0mm/s points for velocity in Fig. 5b). Severed animals that could cling and climb had straightening paths and slower speeds with increasing incline, as was observed in control and sham treated animals (Fig. 5a-h). As expected, severed speeds were significantly lower for baseline 180º (control-severed, p<.001; sham-severed, p=.002) and 135º (control-severed, p<.001; sham-severed, p=.007). Nonsignificant differences at 45º may reflect an overall speed limitation at such extreme inclinations.

To assess leg kinematics and coordination differences across treatments and inclinations, we observed locomotor performance on 180º and 45º inclines on an up-close incline behavioral rig as opposed to the top-down view (Supplementary Movie 3, Supplementary Movie 4). At 45º, 14.3% of severed animals fell, and 42.9% clung on but remained immobile. Consequently, these animals were excluded from locomotor and kinematic analysis. Notably, no control or sham animals fell at any inclination in either experiment. To effectively compare locomotion across conditions, walk cycles following the T3 to T1 gait sequence were analyzed. Severed animals were slower overall but followed locomotor parameter trends in control and sham for 180º. However, they were indistinguishable from controls and shams at 45º as overall locomotion slowed. Swing/stance durations and duty cycles followed the same trends with speed observed previously (Fig. 3k-l) independent of inclination; stance duration was modulated, whilst swing duration remained constant across speeds (Fig. 5i-j, m-n). Stride lengths increased linearly with speed at both 180º (r^2 = .754, p<.001; Fig. 5k) and 45º (r^2 = .387, p<.001; Fig. 5o), and intersegmental differences were proportional across treatments and inclinations (Fig. 5l, p).

At 45º, only control and sham larvae engaged their pygopods to assist climbing (Supplementary Movie 4), which suggested climbing required descending inputs to the abdomen. Moreover, larvae consistently attempted to maintain abdominal contact with the substrate, suggesting sustained contraction of body wall muscles whilst climbing. Pygopod planting events were defined as visible pygopod extensions, and a change

from a parallel alignment with the substrate to a perpendicular pointed orientation into it (Fig. 5q). Analysis of their relation to leg kinematics in all walk cycles showed that most pygopod planting events coincided temporally with the initiation of leg swing phases (Fig. 5r), and pygopod planting-triggered analysis showed the sequence from T3 (blue) to T1 (magenta) as seen by the peaks in the distributions (Fig. 5r). Moreover, pygopod engagement was preceded by an increase in velocity and followed by a decrease (Fig. 5s), consistent with the association between pygopod planting events and leg swings. Although most control and sham larvae engaged their pygopods on the overhang (mean rate = 0.76Hz, range = 0 to 1.20Hz), this did not significantly correlate with their maximum velocity (r^2 = .123, p = .092) (Fig. 5t).

Figure 5: Abdominal inputs are not required for thoracic coordination and pygopods

provide stability in locomotion. (a) Walk paths of control, sham, and severed larvae on top-down incline rig. Scale bar = 2cm. **(b)** Box plots of mean instantaneous velocity of tracked larvae. Boxes = mean ±SE, whiskers = ±SD; *** p<.001, ** p<.01, One-way ANOVA, n = 7. **(c-h)** Representative swing (black) and stance (white) phase plots of walk cycles following T3 to T1 sequence on up-close incline rig. Note the prolonged stance phases in severed at baseline 180º. **(i-p)** Scatter plots comparing locomotor parameters: **(i, m)** swing/stance duration, **(j, n)** swing/stance duty cycle, **(k, o)** stride length and **(l, p)** time difference to T3 (intersegmental difference). All values are averaged per leg; velocity is mean across walk cycles. **(k, o)** Stride length correlates significantly with velocity at 180º (r^2 = .754, p < .001), and 45º (r^2 = .387, p < .001). **(q)** Frames showing pygopod relaxation parallel to substrate (black arrow) and planting into substrate (white arrow). Scale bar = 1mm. **(r)** Histogram of normalized pygopod planting event count relative to initiation of closest leg swing across all walk cycles, binned according to the Freedman–Diaconis rule (see Methods). Note the order of distribution peaks highlights predominant T3 to T1 gait even in unfiltered walk cycles. Total number of leg movements = 117 per leg. **(s)** Normalized mean instantaneous velocity (black trace) relative to pygopod planting events (dotted line); individual velocities across 13 larvae (control and sham combined) in grey. **(t)** Pygopod plant rate correlates insignificantly and weakly with maximum velocity (r^2 = .123, p = .092). N = 7 per treatment.

Abdominal contributions in tunnelling

As evidenced, abdominal inputs (ascending or descending) facilitated free locomotion on flat surfaces and during locomotor challenges. Although clinging and climbing behavior may assist *Tribolium* in navigating and infesting food storage systems, we wanted to probe the effects of severing thoracic-abdominal connectives on tunnelling into their food source. Larvae (control, sham and severed) were individually dropped onto a dish filled with approximately 6mm deep whole wheat flour and visible body length was measured 0s and 120s afterwards (Fig. 6a-f). Control and sham animals immediately tunneled into flour on contact, as observed previously during substrate preference tests, with 100% of control and 75% of sham larvae being fully submerged after 2 minutes (Fig. 6g). Strikingly, 0% of severed animals tunneled effectively into flour, and on average, 90.7% of their original body lengths was exposed after 2 minutes

(Fig. 6h). Severed animals still visibly displaced flour with their legs, albeit without

making any progress (Fig. 6f). Therefore, the abdominal segments are essential for

larval submergence into flour.

Figure 6: Abdominal inputs are crucial for larval tunnelling into natural food source. (a-

f) Top-down frames at 0s and 120s after placement on flour for **(a-b)** control, **(c-d)** sham and

(e-f) severed animals. Severed animals struggle to tunnel; black arrow indicates displacement

of flour from thoracic leg movements despite no body submergence. Scale bar = 2mm. **(g)**

Change in body length between 0s and 120s across treatments. **(h)** Percentage of original

body length exposed after 120s. Whiskers = mean ±SE; *** p<.001, ** p<.01, Kruskal-Wallis

test, n = 8.

Discussion

To study and understand the *Tribolium* larval abdomen's roles in locomotion and its

relationship to the legs, we first characterized the larva's general locomotor

performance across a range of conditions. We found larvae relied on their legs for

unchallenging locomotion on preferred substrates, which mostly consisted of fibrous

materials. This supports the hypothesis that their original habitat consisted of tree

bark[2,22], and explains why wet, slippery surfaces present a challenge. While adults are

capable of locomotion across flour[23], tunnelling is the primary motor program for

navigating their food source during pre-pupal instars. This behavior reflects

ethologically relevant photophobic preferences[17] and their propensity to tunnel deeper

when approaching pupation[24]. The locomotor patterns underlying their exploratory
behavior consist of straight runs and continuous turning, with little to no periods of
immobility. This differs from *Drosophila* larvae, which combine repeated straight runs
with pauses followed by headsweeps, redirecting them to a new trajectory[25,26].
Tribolium larvae also show no evidence of handedness in their exploratory
movements, again in contrast with *Drosophila* adults[27] and larvae[28] but also other
insects such as ants[29] and cockroaches[30]. This may reflect the requirements for
Tribolium to navigate a 3-dimensional environment of constantly shifting substrate. We
also found a larval preference for edge exploration, which may represent
photophobicity, centrophobicity or a thigmotactic drive to maximize body contact with
their environment to mimic how they are naturally enveloped in substrate. This
behavior is also found in *Drosophila* adults, but not for centrophobic or thigmotactic
reasons[31].

The most common gait in insects is the alternating tripod gait, in which the first and
third leg on one side of the body move together with the contralateral second leg[32].
Tribolium larvae, however, use a travelling wave gait, involving posterior-to-anterior
sequences of swings propagating from T3 to T1 across all walking speeds. Their
contralateral leg pairs move synchronously, and phase delays between segments are
largely maintained across speeds. This bilaterally symmetric travelling wave gait is
found in the larvae of other beetles, including leaf beetles[33,34]. Since environmental
conditions do not differ between larval and adult stages in these animals, the wave
gait could therefore reflect developmental constraints on the circuit architecture driving
locomotion in the Cucujiformia infraorder of beetles. However, the travelling wave gait
pattern is found more widely. For example, in krill and crayfish, it generates the forces
required for locomotion in an aquatic medium[35,36], and in some millipedes it is used to
generate the required forces for burrowing in soil[37]. A similar bilaterally symmetrical
gait has been observed in the dung beetle *Pachysoma endroedyi*. In their 'galloping'
gait, the anterior two leg pairs move in a travelling wave, with the hind leg pair dragging
behind them and contributing little to forwards motion[38]. In these desert dwelling
species, in-phase contralateral coordination may be energetically favorable for
traversing constantly shifting terrains like sand, a constraint they might share with
aquatic species. These considerations are equally relevant for *Tribolium,* as they
consume and inhabit a constantly deformable substrate. This gait may become

particularly beneficial during *Tribolium* pre-pupal stages, when larvae tunnel deeper
for warmth and lower population densities to prepare for pupation[24].

Many animals display speed-dependent gait transitions. For example, adult
Drosophila gaits have been shown to transition from a pentapodal wave (one leg is in
swing phase at a time) to a tetrapodal (two legs, but not bilateral pairs) and finally
tripodal wave (three legs) by modulating stance phases as the animal progresses from
slow to fast speeds[39]. Gait adjustments tune movement to the requirements of different
terrains and speeds of locomotion[38,40–44]. We therefore also characterized the
kinematic changes accompanying different speeds of locomotion in *Tribolium* larvae.
They similarly have increasing numbers of limbs in swing phase at a time during
increasingly fast locomotion, but do not appear to adopt a gait other than their
bilaterally symmetrical metachronal wave. Moreover, larval stride length increases
linearly with speed, following a highly conserved locomotor strategy across
invertebrates and vertebrates[45–47]. A similar in-depth analysis of adult *Tribolium* gait
kinematics in the future will allow further comparison.

The *Tribolium* VNC is unfused, with a separate ganglion in each body segment,
chained together by longitudinal connectives, and each thoracic ganglion innervating
a leg pair[48,49]. This anatomical arrangement is found across arthropods[50] and
tardigrades[51], and suggests gait variation across and within species might be derived
from one single modifiable control strategy[52]. This raises the question whether the
neural control of its behavior shares similarities with that of the more well-studied and
soft-bodied *Drosophila* larva, which is legless and moves by means of abdominal
peristaltic waves[26], and has a segmentally fused central nervous system[53].

Qualitatively observed pygopod planting during unrestricted motor programs
(backtrack and redirect, see Fig. 2f) first suggested a prominent role for the abdomen
in assisting natural movement. We present several lines of evidence that pygopods
act to support locomotion in challenging conditions. First, while fast animals crawling
on level fibrous surfaces do not necessitate pygopod engagement, they do while
climbing 45º overhangs. Furthermore, pygopod planting events are largely temporally
locked to leg swing phases, during which the animal needs to maintain static stability,
and these events are preceded by a temporary increase in crawling speed. This shows
a correlation between situations that compromise the animal's stability and likelihood

of pygopod engagement. Furthermore, the weak, non-significant correlation between
pygopod planting rate and maximum velocity further implicates its role in stability over
a mechanism to boost/propel overall speed. Second, severing the A1-A2 connective,
thus removing descending and ascending inputs between the brain, thorax and
abdominal segments, resulted in lower speeds during walking and climbing. Some
severed animals dragged their abdomen during walking (Supplementary Movie 3),
which may contribute to decreasing velocity as the lack of abdominal sensory inputs
may impair its ability to support itself during locomotion. However, impairments on
inclines varied, ranging from some severed animals falling off mild (135º) inclines to a
select few matching control and sham performance on 45º overhangs. The latter may
reflect the overhang limiting the feasible speeds larvae can produce in this motor
challenge. Like the variability in inter-animal velocities in normal larvae, the severity of
severing is likely equally dependent on the individual. Nevertheless, severed animals
never engaged their pygopods, and most tended to fall off 45º overhangs. These
animals' inability to engage their posterior segments therefore particularly impaired
their locomotor capabilities under challenging conditions. Furthermore, this phenotype
is unlikely due to a failure to coordinate leg movements, since severed animals fall into
the same speed-dependent distributions of kinematic patterns including swing and
stance duration, duty cycle, stride length and intrasegmental phase difference as
control and sham larvae. This suggests that abdominal regions are not required for
fundamental leg coordination but provide a crucial support role in larvae.

Other species that have abdominal anchoring points with potential stabilizing roles
include *Labidomera clivicollis* larvae, which use pygopods during body extension
whilst climbing tree branches[33]; and blackfly larvae, which are adapted to anchor into
riverbeds to avoid getting swept away by strong currents[54]. The *Tribolium* larval
anatomical adaptations we describe that confer stability and allow the animal to stay
upright may similarly help evade predation, for example by the parasitoid wasp
Holepyris sylvanidis[55], either by enhancing locomotion efficiency in shifting terrain or
avoiding falling off tree branches in their original habitat. The use of posterior terminal
organs for substrate engagement during locomotion is also found in other soft-bodied
animals including *Drosophila* larvae[11], *Gastrophysa viridula*, which deploy their
pygopods phase-locked to leg movements during the locomotor cycle[56]; and *Manduca*
sexta, whose locomotor cycle starts with the terminal prolegs lifting off the substrate[4].

Our results elucidate a functional role for the pygopods, which were previously only
phenotypically mentioned in the *iBeetle* RNA interference (RNAi) screens[57,58].

The contribution of abdominal segments is likely to extend beyond offering physical
support during locomotion. During motor challenges like overhangs, larvae may send
ascending information from abdominal sensory neurons (e.g., regarding reduced body
contact with substrate) to sensorimotor integration centers for sensory feedback.
These may then integrate and send back descending information regarding internal
representations (e.g., current geotactic location, motor state) to make a higher order
decision to plant the pygopods to increase stability. Interruption of communications to
and from the abdomen and thus lack of stability in severed animals may explain those
that fall off or fail to climb. However, the severed animals that could cling and climb
followed the same leg kinematics as controls and shams, therefore lack of abdominal
input appears to not disrupt thoracic movements.

Furthermore, our data suggest the posterior segments have prominent functions in
tunnelling alongside stability. Tunnelling is a key motor program for navigation,
feeding, development, and reproduction in their natural habitat, and the inability to
tunnel in severed larvae suggests a crucial role for abdominal contractions, pygopod
recruitment, and/or ascending sensory feedback in coordinating effective tunnelling.
The abdomen may be recruited to generate pushing forces to submerge larvae in flour,
similar to millipedes[37]. Equally, it may aid exploratory movements such as the
backtrack and redirect behavior (Fig. 2f) to search for new food sources. Investigating
the structure of both ascending and descending inputs passing through the larval
Tribolium A1-A2 connective is therefore a promising avenue to explore the neural
basis of its behavior and could be a possible pinch point to disrupt the animal's ability
to infest grain stores.

Finally, *Tribolium* is an attractive emerging model system in behavioral neuroscience,
combining a robust experimental toolkit with an extensive body of work describing its
development and physiology. As the first coleopteran species to have a fully
sequenced genome (~16,500 genes), 47% of which are orthologous to insects and
vertebrates, *Tribolium* are already highly favored in the field of developmental
genetics[59]. Their genetic toolkit includes the GAL4-UAS system[60], CRISPR/Cas9
genome editing[61,62], balancer chromosomes, and stable injectable RNAi gene

silencing[63]. Forward genetic screens uncovered previously unknown genetic regulators of leg and body wall muscle development[20], and recent studies have described the structure and function of the *Tribolium* cryptonephridial system, which is essential for the animal's exceptional tolerance to desiccation[64]. The structures and locations of brain regions in larval and adult *Tribolium* showed broad similarities to other insect model organisms[65–67]. In addition, comparison of *Tribolium* and *Drosophila* larval neuromuscular systems revealed resemblances in somatic muscle composition[20]. Its similarities to both ancestral and highly derived insects therefore make it an attractive comparative model.

Overall, this study has emphasized how the kinematic analysis of movement and underlying neural control provides insights into natural behaviors that govern *Tribolium*'s success as a pest. We identified the importance of abdominal segments and pygopods to both locomotor challenges and tunnelling, suggesting that studying the structure and function of its connectives has important potential implications for behavioral neuroscience and food security. How far our findings extend to earlier instars and adults remains a promising question for future study.

Methods

Animals

Wild-type *Tribolium castaneum* (San Bernardino strain) were reared in 500ml borosilicate bottles with perforated lids (VWR SCOT291182809) at 29°C in organic wholewheat flour and 5% dried active yeast per weight. 4-5th instar larvae approximating 4-6mm in length were used in all experiments.

Statistics

Main text results and box plots are given as mean ±SE, and box whiskers are ±SD unless otherwise stated. Details of statistical analysis are included under each experiment's subsection below.

Substrate testing

Basic locomotor ability was tested on twelve substrates – white printer paper, cardboard, parafilm, lens paper, Sylgard™, glass, velvet, sandpaper, plastic, packing

foam, white flour, whole wheat flour, 1%, 5% and 10% agarose. Ten larvae were placed in the center of the substrate and recorded for 1 minute. Movies were recorded with a mobile camera (SM-A415F/DSN, Samsung, Seoul, South Korea) at 30fps and analyzed with Fiji (Fiji is Just ImageJ)[68]. Larval head position was manually tracked for the first 15 seconds (since larvae on preferred substrates often walked out of frame within this time window) with the Manual Tracking plugin (https://imagej.net/ij/plugins/track/track.html) for Fiji.

Free locomotion assay

30 larvae were placed in groups of 3-4 in a 18x18cm arena lined with white printer paper and recorded for 2 minutes in darkness with a custom-built imaging setup using an MQ013MG-E2 XIMEA CMOS (Ximea GmbH, Münster, Germany) camera with a 35mm C-mount lens (Thorlabs Inc, NJ, USA). Animals were illuminated with 850nm infrared light using a Thorlabs M850L3 LED (Thorlabs Inc, NJ, USA). Larvae were tracked with FIMTrack[69].

Gait characterization

Ten larvae were placed on white printer paper under an Olympus SZX10 (Olympus Microscopes, Tokyo, Japan) dissection microscope at 0.8x magnification using a 0.5x objective. Animals were recorded with an MQ042RG-CM XIMEA CMOS (Ximea GmbH, Münster, Germany) camera at 60fps. 3-5 straight walk cycles were extracted and manually tracked with Fiji's Manual Tracking plugin. Gait dynamics and locomotor parameters were analyzed with Python scripts for individual legs; cycle period was calculated as 1 over average cycle frequency. A Savitzky-Golay low-pass filter was applied to manually tracked velocities to discretize swing durations. Stance duration was calculated as the period between consecutive swings. Duty cycle was calculated as average swing/stance duration over average cycle period. Intrasegmental phase difference represents the phase difference between left and right legs within each thoracic segment. Intersegmental phase difference represents phase difference of left legs in each segment relative to left T3.

Coherence and phase analysis was performed with MATLAB application *Groundswell* (https://github.com/JaneliaSciComp/Groundswell). A fast Fourier transform was performed on the anchor leg's velocity trace to determine its spectral composition. The

anchor's fundamental frequency was determined as the frequency with greatest power

density (i.e., during swing phases) in the power spectrum. If the highest power

frequency was ambiguous (i.e., appeared as blocks of plateaus on the spectrum), the

middle value within the plateau was chosen, and the higher frequency block was used

in cases with two adjacent blocks. Coherence and phase relationships of the other five

legs to the anchor was compared at the determined fundamental frequency.

Groundswell uses coherence magnitude as a measure of the linear relationship

between two waveforms at the determined fundamental frequency, and calculates a

coherence magnitude threshold with the formula:

516
$$C\_mag\_thresh = \sqrt{1 - a^{\frac{1}{(\text{dof}/2-1)}})}$$

Where degrees of freedom (dof) = 2*R*K, in which R is the number of observations of

the waveforms together and K is the number of tapers. This threshold determines the

minimum coherence magnitude required for significant coherence between two

signals with $\alpha = 0.05$.

Preliminary incline assay

Ten larvae were individually placed on custom-built platforms of varying incline (180°

– flat substrate, 135° – slight upwards incline, 90° – vertical wall, 45° – overhang).

Locomotion was filmed with an MQ022RG-CM XIMEA CMOS (Ximea GmbH, Münster,

Germany) camera for 1 minute at 30fps with a Thorlabs OSL1 Fiber Illuminator

(Thorlabs Inc, NJ, USA) with gooseneck lights at lowest intensity. Raw videos were

optimized for FIMTrack in Fiji with custom-written macros. Statistical analysis on mean

instantaneous velocity was performed as one-way ANOVA.

Connective severing assay

In preliminary control assays, six larvae were anesthetized on ice and sequestered

between 0.1mm tungsten pins ventral side up on a dissection dish. A cut was made

with fine microdissection scissors on the cuticle line separating A1 and A2 segments.

The blades were inserted at a ~45° angle to the cuticle and pulled upwards to sever

the connective. Sham larvae were given lateral lesions on A1. Control larvae were not

lesioned but anesthetized on ice for equal time. Surgeries were verified via fillet

microdissection post-experimentally. Larvae without clear, visible connective severing were discarded from results. Behavior was filmed with the setup outlined in *Free locomotion assay* 2 hours after surgery*;* raw videos were processed in Fiji and tracked with FIMTrack. Statistical analysis on mean instantaneous velocity was performed as a between-subjects one-way ANOVA.

For following incline and tunnelling experiments, surgeries were repeated on seven larvae; sham animals' lateral lesions alternated between left and right sides to minimize biases in turning. Different animals were used for incline and tunnelling assays to avoid excessive fatigue on lesioned animals.

Top-down connective severing incline assay

Same methodology as *Preliminary incline assay,* excluding 90° due to short lifespans of severed animals. 2 larvae (1 control and 1 severed) could not be tracked with FIMTrack, thus mean velocity was obtained with wrMTrck (https://www.phage.dk/plugins/wrmtrck.html) and their tracks are therefore absent in Fig. 5a. Statistical analysis on mean instantaneous velocity was performed as a repeated-measures ANOVA with Bonferroni corrected post-hocs.

Up-close connective severing incline assay

Control, sham and severed larvae were individually placed onto a walled 70mm x 3mm paper platform on a custom-built rig; incline was adjusted with a micromanipulator and verified with a protractor. Locomotion was filmed until the animal walked out of view with a MQ042RG-CM XIMEA CMOS (Ximea GmbH, Münster, Germany) camera connected with a 1"-32 C-mount (Thorlabs Inc, NJ, USA) to a Canon FR1048KWN lens (Canon Inc, Tokyo, Japan). If the animal remained immobile, filming was stopped at the experimenter's discretion. 180° videos were filmed at 100fps and illuminated by an overhead Thorlabs M850L3 850nm light (Thorlabs Inc, NJ, USA). 45° overhang trials were illuminated from the bottom with a Stemmer Imaging CCS LDL100 850nm light (Stemmer Imaging Ltd, Tongham, UK) and limited to 50fps by the field of view. 135° and 90° were not tested due to the short lifespans of severed animals. For analysis, velocity was tracked with FIMTrack; pygopod planting events and start and stop frames of leg swings were manually tracked with Fiji's Manual Tracking plugin. The constricted arena walls promoted exploration, therefore to effectively analyze

locomotor parameters and gait kinematics, walk cycles following the T3 to T1 sequence were extracted and stance durations above median ± 1SD were discarded to filter out pause periods. Mean percentage of sequences extracted out of total tracked time (including pauses) per animal were: 55.4% (control 180°), 47.2% (sham 180°), 34.2% (severed 180°), 28.3% (control 45°), 24.9% (sham 45°) and 26.0% (severed 45°). Cycle period, stance duration and duty cycle retained their definitions in _Gait characterization,_ but swing duration was calculated as the time between swing start and swing stop frames.

A pygopod plant event was defined as a visible extension of the pygopods, forming almost a blunt triangular point, into the substrate, regardless of whether the abdomen is in contact with the substrate. The Freedman-Diaconis rule was used in Fig. 5r, using the formula: (2*IQR) / 3√n to calculate x-axis bin length. Planting rate was used for Fig. 5t to normalize for variable video lengths, calculated as number of events over total tracked time. Pearson's correlations were applied for regression analyses.

Tunneling assay

Post-surgery animals were allowed to wake and begin walking on paper before testing. A circular dish was filled with ~10mm of flour; control, sham and severed larvae were individually placed and filmed with an MQ0113RG-E2 XIMEA CMOS (Ximea GmbH, Münster, Germany) at 15fps under a Nikon SMZ745T dissection scope at 0.63x magnification. Filming stopped at 2 minutes regardless of how deep larvae tunneled. Body lengths at time 0s and 120s were measured in Fiji. Statistics was performed as a Kruskal-Wallis test and Dunn post-hoc analysis with Bonferroni correction (given non-normal data distribution).

Data availability

Source data is available at: 10.5281/zenodo.11520088 (analyzed files) and 10.17630/a97be752-0c97-42a2-ada4-745c5acff342 (raw movies).

Code availability

Analysis code is available at: 10.5281/zenodo.11520206.

References

References

1.   Phillips, T. W. & Throne, J. E. Biorational Approaches to Managing Stored-

Product Insects. *Annual Review of Entomology* **55**, 375–397 (2010).

2.   Good, N. E. The Flour Beetles of the Genus *Tribolium*. (1936)

doi:10.22004/AG.ECON.164672.

3.   Mantziaris, C., Bockemühl, T. & Büschges, A. Central pattern generating

networks in insect locomotion. *Dev Neurobiol* **80**, 16–30 (2020).

4.   van Griethuijsen, L. I. & Trimmer, B. A. Locomotion in caterpillars. *Biol Rev Camb*

Philos Soc **89**, 656–670 (2014).

5.   Leiras, R., Cregg, J. M. & Kiehn, O. Brainstem Circuits for Locomotion. *Annu Rev*

Neurosci **45**, 63–85 (2022).

6.   Daley, M. A. Understanding the Agility of Running Birds: Sensorimotor and

Mechanical Factors in Avian Bipedal Locomotion. *Integr Comp Biol* **58**, 884–893

(2018).

7.   Hooper, S. L. *et al.* Neural control of unloaded leg posture and of leg swing in

stick insect, cockroach, and mouse differs from that in larger animals. *J Neurosci*

29, 4109–4119 (2009).

8.   Triphan, T., Poeck, B., Neuser, K. & Strauss, R. Visual targeting of motor actions

in climbing *Drosophila*. *Curr Biol* **20**, 663–668 (2010).

9.   Picker, M., Colville, J. F. & Burrows, M. A cockroach that jumps. *Biol Lett* **8**, 390–

392 (2012).

10.   Digumarti, K. M., Trimmer, B., Conn, A. T. & Rossiter, J. Quantifying Dynamic

Shapes in Soft Morphologies. *Soft Robot* **6**, 733–744 (2019).

11. Booth, J. H., Meek, A. T., Kronenberg, N. M., Pulver, S. R. & Gather, M. C.

Optical mapping of ground reaction force dynamics in freely behaving *Drosophila*

melanogaster larvae. *eLife* **12**, (2024).

12. Shield, S., Jericevich, R., Patel, A. & Jusufi, A. Tails, Flails, and Sails: How

Appendages Improve Terrestrial Maneuverability by Improving Stability.

Integrative and Comparative Biology **61**, 506–520 (2021).

13. Đukić, N. *et al.* Behavioural responses of Tribolium castaneum (Herbst) to

different types of uninfested and infested feed. *Bulletin of Entomological*

Research **110**, 550–557 (2020).

14. Gerken, A. R., Scully, E. D. & Campbell, J. F. Red Flour Beetle (Coleoptera:

Tenebrionidae) Response to Volatile Cues Varies With Strain and Behavioral

Assay. *Environmental Entomology* **47**, 1252–1265 (2018).

15. Đukić, N. *et al.* The effect of 1-pentadecene on Tribolium castaneum behaviour:

Repellent or attractant? *Pest Management Science* **77**, 4034–4039 (2021).

16. Semeao, A. A., Campbell, J. F., Whitworth, R. J. & Sloderbeck, P. E. Response of

Tribolium castaneum and Tribolium confusum adults to vertical black shapes and

its potential to improve trap capture. *Journal of Stored Products Research* **47**,

88–94 (2011).

17. Duehl, A. J., Cohnstaedt, L. W., Arbogast, R. T. & Teal, P. E. A. Evaluating Light

Attraction to Increase Trap Efficiency for Tribolium castaneum (Coleoptera:

Tenebrionidae). *Journal of Economic Entomology* **104**, 1430–1435 (2011).

18. Prakashini, V. Y., Srinivasan, G., Shanthi, M. & Prabakaran, K. Preference of Red

Flour Beetle *Tribolium castaneum* (Herbst) Towards Colour Cues. *Indian Journal*

of Entomology 445–447 (2023) doi:10.55446/IJE.2021.151.

19. Romero, S. A., Campbell, J. F., Nechols, J. R. & With, K. A. Movement Behavior

of Red Flour Beetle: Response to Habitat Cues and Patch Boundaries.

Environmental Entomology **39**, 919–929 (2010).

20. Schultheis, D. *et al.* A Large Scale Systemic RNAi Screen in the Red Flour Beetle

Tribolium castaneum Identifies Novel Genes Involved in Insect Muscle

Development. *G3 Genes|Genomes|Genetics* **9**, 1009–1026 (2019).

21. Devi, M. B. & Devi, N. V. Biology of rust-red flour beetle, *Tribolium castaneum*

(Herbst) (Tenebrionidae:Coleoptera). *Ind. Jour. of Entomol.* **77**, 81 (2015).

22. Dawson, P. S. Life History Strategy and Evolutionary History of *Tribolium* Flour

Beetles. *Evolution* **31**, 226–229 (1977).

23. Pai, A. *Tribolium*. in *Encyclopedia of Animal Behavior* 446–452 (Elsevier, 2010).

doi:10.1016/B978-0-08-045337-8.00030-9.

24. King, C. E. & Dawson, P. S. Population biology and the Tribolium model. *Evol.*

Biol. **5**, 133–227 (1972).

25. Suster, M. L. & Bate, M. Embryonic assembly of a central pattern generator

without sensory input. *Nature* **416**, 174–178 (2002).

26. Lahiri, S. *et al.* Two Alternating Motor Programs Drive Navigation in *Drosophila*

Larva. *PLOS ONE* **6**, e23180 (2011).

27. Buchanan, S. M., Kain, J. S. & De Bivort, B. L. Neuronal control of locomotor

handedness in *Drosophila*. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 6700–6705 (2015).

28. Wosniack, M. E., Festa, D., Hu, N., Gjorgjieva, J. & Berni, J. Adaptation of

Drosophila larva foraging in response to changes in food resources. *Elife* **11**,

e75826 (2022).

29. Hunt, E. R. *et al.* Ants show a leftward turning bias when exploring unknown nest

sites. *Biol. Lett.* **10**, 20140945 (2014).

30. Camhi, J. M. & Johnson, E. N. High-frequency steering maneuvers mediated by

tactile cues: antennal wall-following in the cockroach. *Journal of Experimental*

Biology **202**, 631–643 (1999).

31. Soibam, B. *et al.* Open-field arena boundary is a primary object of exploration for

Drosophila. *Brain Behav* **2**, 97–108 (2012).

32. Wilson, D. M. Insect walking. *Annu Rev Entomol* **11**, 103–122 (1966).

33. Gustafson, G. T. & Chaboo, C. S. Ambulatory Use of Ventro-Lateral Abdominal

Ampullae by Larvae of *Labidomera clivicollis* (Kirby) (Coleoptera: Chrysomelidae:

Chrysomelinae). *cole* **63**, 357–363 (2009).

34. Zurek, D. B., Gorb, S. N. & Voigt, D. Locomotion and attachment of leaf beetle

larvae *Gastrophysa viridula* (Coleoptera, Chrysomelidae). *Interface Focus* **5**,

20140055 (2015).

35. Mulloney, B. & Smarandache-Wellmann, C. Neurobiology of the crustacean

swimmeret system. *Prog Neurobiol* **96**, 242–267 (2012).

36. Zhang, C., Guy, R. D., Mulloney, B., Zhang, Q. & Lewis, T. J. Neural mechanism

of optimal limb coordination in crustacean swimming. *Proc. Natl. Acad. Sci.*

U.S.A. **111**, 13840–13845 (2014).

37. Manton, S. M. The Evolution of Arthropodan Locomotory Mechanisms. Part 71.

Functional Requirements and Body Design in Colobognatha (diplopoda),

Together with a Comparative Account of Diplopod Burrowing Techniques, Trunk

Musculature and Segmentation. *Journal of the Linnean Society of London,*

Zoology **44**, 383–462 (1961).

38. Smolka, J., Byrne, M. J., Scholtz, C. H. & Dacke, M. A new galloping gait in an

insect. *Current Biology* **23**, R913–R915 (2013).

39. DeAngelis, B. D., Zavatone-Veth, J. A. & Clark, D. A. The manifold structure of

limb coordination in walking *Drosophila*. *eLife* **8**, e46409 (2019).

40. Graham, D. A behavioural analysis of the temporal organisation of walking

movements in the 1st instar and adult stick insect *(Carausius morosus)*. *J. Comp.*

Physiol. **81**, 23–52 (1972).

41. Bässler, U. & Büschges, A. Pattern generation for stick insect walking

movements—multisensory control of a locomotor program. *Brain Research*

Reviews **27**, 65–88 (1998).

42. Bender, J. A. *et al.* Kinematic and behavioral evidence for a distinction between

trotting and ambling gaits in the cockroach Blaberus discoidalis. *Journal of*

Experimental Biology **214**, 2057–2064 (2011).

43. Schilling, M., Hoinville, T., Schmitz, J. & Cruse, H. Walknet, a bio-inspired

controller for hexapod walking. *Biol Cybern* **107**, 397–419 (2013).

44. Ramdya, P. *et al.* Climbing favours the tripod gait over alternative faster insect

gaits. *Nat Commun* **8**, 14494 (2017).

45. Strang, K. T. & Steudel, K. Explaining the scaling of transport costs: the role of

stride frequency and stride length. *Journal of Zoology* **221**, 343–358 (1990).

46. Hooper, S. L. Body size and the neural control of movement. *Current Biology* **22**,

R318–R322 (2012).

47. Granatosky, M. C. & McElroy, E. J. Stride frequency or length? A phylogenetic

approach to understand how animals regulate locomotor speed. *J Exp Biol* **225**,

jeb243231 (2022).

48. Hunnekuhl, V. S., Siemanowski, J., Farnworth, M. S., He, B. & Bucher, G.

Immunohistochemistry and Fluorescent Whole Mount RNA In Situ Hybridization

in Larval and Adult Brains of *Tribolium*. *Methods Mol Biol* **2047**, 233–251 (2020).

49. Farnworth, M. S., Bucher, G. & Hartenstein, V. An atlas of the developing

Tribolium castaneum brain reveals conservation in anatomy and divergence in

timing to *Drosophila melanogaster*. *Journal of Comparative Neurology* **530**,

2335–2371 (2022).

50. Richter, S. *et al.* Invertebrate neurophylogeny: suggested terms and definitions

for a neuroanatomical glossary. *Front Zool* **7**, 29 (2010).

51. Mayer, G. *et al.* Selective neuronal staining in tardigrades and onychophorans

provides insights into the evolution of segmental ganglia in panarthropods. *BMC*

Evol Biol **13**, 230 (2013).

52. Nirody, J. A. Universal Features in Panarthropod Inter-Limb Coordination during

Forward Walking. *Integrative and Comparative Biology* **61**, 710–722 (2021).

53. Truman, J. W. & Bate, M. Spatial and temporal patterns of neurogenesis in the

central nervous system of Drosophila melanogaster. *Dev Biol* **125**, 145–157

(1988).

54. Jamnback, H. & Frempong-Boadu, J. Testing blackfly larvicides in the laboratory

and in streams. *Bull World Health Organ* **34**, 405–421 (1966).

55. Amante, M., Russo, A., Schöller, M. & Steidle, J. L. M. Olfactory host location and

host preference of Holepyris sylvanidis (Brèthes) (Hymenoptera: Bethylidae), a

parasitoid of Tribolium confusum Jacquelin du Val and T. castaneum (Herbst)

(Coleoptera: Tenebrionidae). *Journal of Stored Products Research* **78**, 105–109

(2018).

56. Zurek, A. J. *et al.* Quantification of anthropogenic impact on groundwater-

dependent terrestrial ecosystem using geochemical and isotope tools combined

with 3-D flow and transport modelling. *Hydrology and Earth System Sciences* **19**,

1015–1033 (2015).

57. Schmitt-Engel, C. *et al.* The iBeetle large-scale RNAi screen reveals gene

functions for insect development and physiology. *Nat Commun* **6**, 7822 (2015).

58. Hakeemi, M. S. *et al.* Screens in fly and beetle reveal vastly divergent gene sets

required for developmental processes. *BMC Biology* **20**, 38 (2022).

59. Richards, S. *et al.* The genome of the model beetle and pest *Tribolium*

castaneum. *Nature* **452**, 949–955 (2008).

60. Schinko, J. B. *et al.* Functionality of the GAL4/UAS system in *Tribolium* requires

the use of endogenous core promoters. *BMC Developmental Biology* **10**, 53

(2010).

61. Gilles, A. F., Schinko, J. B. & Averof, M. Efficient CRISPR-mediated gene

targeting and transgene replacement in the beetle Tribolium castaneum.

Development **142**, 2832–2839 (2015).

62. Adrianos, S., Lorenzen, M. & Oppert, B. Metabolic pathway interruption:

CRISPR/Cas9-mediated knockout of tryptophan 2,3-dioxygenase in Tribolium

castaneum. *J Insect Physiol* **107**, 104–109 (2018).

63. Tomoyasu, Y. *et al.* Exploring systemic RNA interference in insects: a genome-

wide survey for RNAi genes in Tribolium. *Genome Biology* **9**, R10 (2008).

64. Naseem, M. T. *et al.* NHA1 is a cation/proton antiporter essential for the water-

conserving functions of the rectal complex in Tribolium castaneum. *Proceedings*

of the National Academy of Sciences **120**, e2217084120 (2023).

65. He, B. *et al.* An ancestral apical brain region contributes to the central complex

under the control of foxQ2 in the beetle Tribolium. *eLife* **8**, e49065 (2019).

66. Garcia-Perez, N. C., Bucher, G. & Buescher, M. Shaking hands is a

homeodomain transcription factor that controls axon outgrowth of central

complex neurons in the insect model Tribolium. *Development* **148**, dev199368

(2021).

67. Farnworth, M. S., Bucher, G. & Hartenstein, V. An atlas of the developing

Tribolium castaneum brain reveals conservation in anatomy and divergence in

timing to Drosophila melanogaster. *J Comp Neurol* 10.1002/cne.25335 (2022)

doi:10.1002/cne.25335.

68. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis.

Nat Methods **9**, 676–682 (2012).

69. Risse, B., Otto, N., Berh, D., Jiang, X. & Klämbt, C. FIM Imaging and FIMtrack:

Two New Tools Allowing High-throughput and Cost Effective Locomotion

Analysis. *J Vis Exp* 52207 (2014) doi:10.3791/52207.

70. Xu Ying, B., Zwart, M.F. and Pulver, S.R. (2024) 'Context-dependent coordination

of movement in *Tribolium castaneum* larvae'. Dataset. University of St Andrews

Research Portal. https://doi.org/10.17630/a97be752-0c97-42a2-ada4-

745c5acff342

End Notes

Acknowledgements

This project was supported by the Institute of Behavioural and Neural Sciences
Undergraduate Research Assistant Scheme at the University of St Andrews. We thank
Barry Denholm for providing *Tribolium* stocks and advice early in the project, Adam
Taylor from Janelia Research Campus for *Groundswell*, Balwant Singh for help
building the up-close incline rig, and members of the Pulver and Zwart Labs for
constant support.

Author contributions

B.X.Y., M.F.Z. and S.R.P. conceived experiments; B.X.Y. reared animals, performed
all experiments and analyzed data; M.F.Z. and B.X.Y. analyzed gait data; M.F.Z. and

S.R.P. performed surgeries; B.X.Y. prepared figures and drafted manuscript; S.R.P.
and M.F.Z. revised manuscript; S.R.P and M.F.Z. supervised the project.

793 Conflicts of interest

794 The authors declare no conflicts of interest.