# The GTPase-activating protein CG42795 is a potent neuronal regulator of ageing in *Drosophila melanogaster*

Gergő Falcsik[1,2*], Fanni Kersztes[1,2*], Tünde Mónika Balog[1], Tibor Kovács[1#]

[1]Department of Genetics, Eötvös Loránd University (ELTE), Budapest, Hungary;

[2]Doctoral School of Biology, Institute of Biology, ELTE Eötvös Loránd University, Pázmány Péter sétány 1/C, Budapest, H-1117, Hungary;

\* These authors contributed equally to this work
\# Correspondence: tibor.kovacs@ttk.elte.hu ; Tel.: 00-36-1-372-25-00/8694

Keywords: autophagy, ageing, Rab2, GTPase-activating proteins

## Abstract

Macroautophagy (hereafter autophagy) is the process by which cellular components to be degraded are isolated from the cytoplasm by a double membrane (the mature vesicle, called the autophagosome). Its contents are degraded by acid hydrolases after fusion with the lysosome. This lysosomal degradation process is essential for maintaining cellular homeostasis. Dysfunctional autophagy can lead to the accumulation of cytotoxic protein forms that contribute to the onset of age-related diseases. However, autophagy activity declines as we age/throughout life. Stimulating its function during lifespan is particularly important in post-mitotic cells such as neurons, where cells cannot divide to replace the dead cells. Our research group aims to find new autophagy activation sites to stimulate the efficiency of acid degradation in neurons during ageing. One possibility is the stimulation of membrane fusion events, which are necessary for autophagic degradation, through the activation of small GTPase enzymes. Our previous results have shown that neuron-specific overexpression of the activated form of the Rab2 small GTPase has autophagy and lifespan-enhancing effects. In the present experiment, we used an RNA interference screen to investigate whether silencing 12 GTPase-activating proteins (GAPs) belonging to the TBC1 domain family can enhance Rab2 activation in the *Drosophila* nervous system. Several of the GAPs studied increased the number of Rab2-positive structures, 5 of which were selected for further screening. Our results suggest that neuronal silencing of *CG42795* has autophagy-enhancing effects, the ability to improve the locomotor ability of animals, and prolong lifespan. Our research has shown that the study of GAPs could be a promising new avenue of research for ageing researchers. Further analysis of CG42795 could lead to the development of potential autophagy activators.

# Introduction

Cells utilise degradation processes called autophagy or cellular self-digestion that removes damaged organelles and protein aggregates, also creates new building blocks. Autophagy levels must be tightly regulated to maintain a normal cellular homeostasis [1], [2]. Autophagy includes the processes of microautophagy, chaperone-mediated autophagy (CMA) and macroautophagy, all of which have essential but distinct functions. Microautophagy directly engulfs cytoplasmic components by the invagination of the lysosomal membrane [3]. CMA can only degrade soluble proteins that are selectively and directly translocated to the lysosome. All substrate proteins contain a KFERQ-like motif that is recognized by a chaperone complex (containing hsc70) in the cytoplasm. It interacts with the lysosomal associated membrane protein type 2A (LAMP-2A) to assist the translocation step [4]. Macroautophagy (here after autophagy) is a type of catabolic degradation in which cytoplasmic compartments are sequestered within a double-membrane vesicle, the autophagosome, which fuses with lysosomes to deliver the degradation enzymes and create the environment in which the enzymes can function and eliminate the particles [5].

The location of phagophore formation is still not well understood, although the endoplasmic reticulum (ER) appears to be key, although mitochondrial, Golgi and plasma membrane origins have also been described [6], [7]. Phagophore formation is initiated by the activation of the Atg1 (autophagy-related 1; ULK in mammals) kinase complex, which phosphorylates and activates other proteins required for autophagosome biogenesis [8]. The TOR (target of rapamycin; TORC1 in mammals) pathway, which is the master regulator of intracellular metabolism, is responsible for regulating the Atg1 kinase complex. If the pathway detects elevated levels of certain nutrients, amino acids or insulin, it blocks autophagy by inhibiting the Atg1 (autophagy-related 1) kinase complex, since the cell has enough building blocks and energy for protein synthesis. However, if the cell is starving and the TOR signaling pathway is switched off, the Atg1 kinase complex can be active and new autophagosomes can form, allowing the cell to obtain building blocks from the breakdown of its own materials, because protein synthesis must be continuous in a cell, even in the case of nutrient deficiency. This regulatory mechanism shows that even in basic homeostatic processes, autophagy is essential for cell survival, and does not merely play a recycling role [9].

The early stages of autophagosome biogenesis also involve the Vps34/PI3K complex, which converts phosphatidylinositol (PI) to phosphatidylinositol 3-phosphate (PI3P) on the phagophore membrane. PI3P is an autophagosome membrane-specific PI, so this pattern is recognised by other proteins required for autophagy and recruited to the forming membrane [10], [11]. One example is Atg2, which forms a complex with Atg18. The Atg2-Atg18 complex channels lipids into the phagophore membrane, increasing its surface area. Small vesicles containing Atg9 transmembrane proteins are also incorporated into the membrane, which also plays a role in its growth. In addition to PI3P, another important identity-determining element is added to the membrane of the phagophore and later the autophagosome, namely the Atg8 (LC3B in mammals, Atg8a in *Drosophila*), a ubiquitin-like protein. The ubiquitin ligase-like conjugation system places a phosphatidylethanolamine lipid anchor to the free Atg8 proteins, and then these lipidated Atg8 proteins are then anchored to the phagophore membrane. Amino

acid cleavage of the Atg4 protein at the C-terminus of Atg8 is required for lipid anchor placement. Atg8 can contact the adapter protein p62 (*Drosophila* ortholog Ref(2)P). Many substances marked for selective degradation are processed by Atg8-p62 (and other adaptors that also linked to Atg8) proteins [12]. Mutant, fluorescent forms of these two proteins are often used by researchers as autophagy markers.

The formation, transport and fusion of vesicles during the degradation pathway is a finely regulated, intricate process that requires several proteins with different functions. These include small GTPase, coat, motor, docking and folding proteins and many more [8].

Optimal levels of autophagy are required for cell survival, and impairment of autophagy is associated with age-related diseases such as neurodegeneration and cancer [13], [14]. Autophagic activity has been shown to decline over the lifespan, contributing to the accumulation of toxic protein aggregates in cells, causing them to lose functionality and viability [15]–[18]. This is particularly problematic in post-mitotic cells such as neurons, where damaged and lost neurons cannot be replaced [19]. It was also found that the neuronal autophagic process is usually inhibited in the downstream steps, leading to the accumulation of undigested particles in autolysosomes [20].

As autophagy gradually declines with age, leading to neuronal death and neurodegenerative diseases, we should identify novel targets to enhance autophagy. One possible way to improve autophagic efficiency is to enhance lysosomal fusion events, a step that is typically impaired with age. The HOPS and SNARE complexes regulate the fusion steps with the help of small GTPase proteins [21]. Previously, we have studied three small GTPase proteins in detail: Rab2, Rab7 and Arl8, all of which have important and evolutionarily conserved roles in degradation. Of these Rab2 can induce not only autophagosome-lysosome but also Golgi vesicle-lysosome fusion steps. The function of small GTPase proteins includes membrane fusion, tracking, recruitment of proteins and membrane docking [22]. Small GTPases act like molecular switches, switching between active GTP-bound and inactive GDP-bound forms [23].There are two types of enzymes that regulate small GTPases: GAP enzymes (GTPase-activating proteins) induce GTP hydrolysis and inactivate the protein, while GEFs (guanine nucleotide exchange factors) promote GTP binding and activation [24].

Our previous results have shown that Rab2 activation has a positive effect on lifespan, movement and autophagic activity. Constitutively active Rab2 was also found to be beneficial in the *Drosophila* model of Parkinson's disease, where it could extend lifespan, improve degradation and reduce the amount of toxic alpha-synuclein particles in the brain [25].

We are currently looking for GAP enzymes that regulate Rab2 activity. By silencing specific GAPs, we can inhibit one of the inactivating enzymes in the process, thereby activating the small GTPase protein. This will have a similar effect to constitutively active Rab2 and increase autophagy, lifespan and locomotion. We used an RNA interference screen to investigate 12 GAPs belonging to the TBC1 domain family to find Rab2 regulators.

# Materials and Methodes

<u>Strains used for experiments and their maintenance</u>

We used *Drosophila melanogaster* for our experiments. The strains have been obtained from the Bloomington Drosophila Stock Centre (BDSC) and have also been generated by crossing animals from the ELTE Department of Genetics Strain Collection (EDSC) (**Table 1**). The strains used were maintained on a standard corn-sugar-agar medium. All our experiments were carried out on animals kept at 29°C. The 12 GAP genes silenced in the experiment are referred to by their CG number or as GAP-RNAi. For control animals we used an isogenic $w^{1118}$ (*isowhite*) strain simulating the wild-type *Drosophila*, and for certain experiments an additional RNAi-control strain (GFP-RNAi) were used (**Table 1**). This may be necessary because the double-stranded RNA expressed in RNAi strains can itself induce non-specific effects by activating the RISC complex. The RNAi control is designed to simulate the non-specific effects of RNAi by producing extra targetless double-stranded RNA compared to the *isowhite* (wild type with white eyes) control [26]. However, this GFP-RNAi control strain cannot be used in experiments where the expression of the GFP protein is required (for example, in the case of several of our fluorescent microscopy experiments).

| Name | Genotype | Origin |
|---|---|---|
| control / isogenic white | $w^{1118}$ | BDSC: 5905 |
| GFP-RNAi (RNAi-control) | y[1] sc[*] v[1]; P{y[+t7.7] v[+t1.8] =VALIUM22-EGFP.shRNA.4}attP2 | BDSC: 41551 |
| CG1695 (GAP-RNAi) | y[1] v[1]; P{y[+t7.7] v[+t1.8] =TRiP.HMJ24138}attP40 | BDSC: 62898 |
| CG4041 II (GAP-RNAi) | y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8] =TRiP.GL00239}attP2/TM3, Sb[1] | BDSC: 35332 |
| CG4041 III (GAP-RNAi) | y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8] =TRiP.GL00239}attP2/TM3, Sb[1] | BDSC: 35332 |
| CG5344 (GAP-RNAi) | y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8] =TRiP.HMC06579}attP40 | BDSC: 77447 |
| CG6182 (GAP-RNAi) | y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8] =TRiP.HMC06532}attP40 | BDSC: 77400 |
| CG7112 (GAP-RNAi) | y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8] =TRiP.HMS01132}attP2/TM3, Sb[1] | BDSC: 34976 |
| CG7324 (GAP-RNAi) | y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8] =TRiP.HMS00722}attP2 | BDSC: 32929 |
| CG8085 (GAP-RNAi) | y[1] v[1]; P{y[+t7.7] v[+t1.8] =TRiP.JF03085}attP2 | BDSC: 28670 |
| CG11727 (GAP-RNAi) | y[1] v[1]; P{y[+t7.7] v[+t1.8] =TRiP.HMS01818}attP40 | BDSC: 38350 |
| CG12241 (GAP-RNAi) | y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8] =TRiP.HMS00612}attP2/TM3, Sb[1] | BDSC: 33729 |
| CG31935 (GAP-RNAi) | y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8] =TRiP.HMC05513}attP40 | BDSC: 64495 |
| CG42612 (GAP-RNAi) | y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8] =TRiP.HMC06174}attP40 | BDSC: 66313 |
| CG42795 (GAP-RNAi) | y[1] v[1]; P{y[+t7.7] v[+t1.8] =TRiP.HMJ23781}attP40/CyO | BDSC: 62385 |
| DdcGal4 | w[1118]; P{Ddc-GAL4.L}4.36 | BDSC: 7009 |

| DdcGal4-Rab2-YFP | w[1118]; DdcGal4; Rab2-YFP/ST6 | EDSC |
|---|---|---|
| cgGal4 (II.) | w[1118]; P{Cg-GAL4.A}2 | EDSC |
| cgGal4-Rab2-YFP | w[1118]; cgGal4; Rab2-YFP/ST6 | created by us |
| Rab2-YFP | y[1] w[*]; P{w[+mC]=UAST-YFP.Rab2}l(3)neo38[02]/TM3, Sb[1] | BDSC |
| ST6 | w*; Sp; Sb / ST6 | EDSC |
| 3x mCh-Atg8a, DdcGal4 | w[1118]; DdcGal4; 3x mCh-Atg8a | EDSC |
| Act<CD2<Gal4,GFP (V40) | hsFlp; pAct<CD2<Gal4 UAS-nlsGFP | gift from Gábor Juhász |

*Table 1: Name, genotype, and place of origin of the strains used in our experiments.*

For clonal systems, we used the strain Act<CD2<Gal4,GFP (V40) *(Table 1)*. The essence of this strain is that between the coding sequence of the Gal4 protein and the actin promoter that drives it, there is a flip out cassette that prevents transcription of the gene. There are FRT sequences at both ends of the flip out cassette where the FLP protein can loop out this sequence, allowing the Gal4 protein to be transcribed again. The FLP protein can be activated by heat shock (hsFLP) in random cells. Where the FLP has been activated by heat shock and the Gal4 protein can be transcribed, these cells express GFP as well as the GAP-RNAi sequence, but in the other cells the transgenes are not expressed in the absence of Gal4, these cells can be considered as "wild type" control cells [27].

Fluorescence microscopy

We studied the effects of GAP-RNA interference in two types of tissues, adult *Drosophila* brains and larval fat bodies. Adult female flies were decapitated after anaesthetized with ether, the heads were dissected and fixed in formaldehyde (4%) diluted in PBS (phosphate-buffered saline) for 20 min at room temperature, then washed 3 x 5 times in PBS. We then dissected the brains, also in PBS, and covered the samples with Hoechst solution containing PBS:glycerol (ratio 1:4) before microscopy. To examine fat bodies, the fat body lobes of larvae at the L3 feeding stage (L3F) were dissected in PBS without fixation. Each sample was made from approximately 5-6 larvae. After dissection the samples were immediately covered with Hoechst solution containing PBS:glycerol (ratio 1:4). For starvation experiments, L3 feeding stage (L3F) larvae were transferred to a 20% sucrose solution (protein starvation) for 2 hours prior to dissection. Then they were dissected in PBS without fixation and covered with Hoechst solution containing PBS:glycerol (ratio 1:4).

Microscopic images were taken using a Zeiss Axioimager Z1 microscope equipped with ApoTome2. The brains were photographed at 200x magnification, while the fat bodies were photographed at 400x magnification. The images were taken across the entire thickness of the organ, at a distance of 1 μm per slice. During data analysis, we merged the slices of the organ so that all the fluorescently labelled structures could be counted. We used ZEN 3.5 (Blue edition v3.5.093) and ImageJ software to analyze the images.

In the experiments with the clonal systems, the larvae were transferred to 29°C for 18 hours prior to dissection, and except for this period, the larvae were kept at 25°C. For LysoTracker Red (L7528) staining of acidic compartments in fat bodies, we diluted LysoTracker Red 1:1000

in PBS. After staining for 2 minutes, we washed the samples twice in PBS for 2-2 minutes before covering. The samples were covered with Hoechst solution in PBS:glycerol (ratio 1:4).

## Immunohistochemistry

We used immunohistochemistry to study the quantitative changes in ubiquitinated structures that are present in the brain of the fruit flies. For the experiment, we used the brains of 7-day-old female *Drosophila*. This was done as described by Kovács et al [20]. The samples were prepared as described for fluorescent microscopy and fixed in formaldehyde (4%) for 40 minutes at room temperature. After fixation, we did the washing with PBT (3x 10 minutes, PBS + 0.1% Tween20). Then, we dissected the brains and blocked them for 1 hour in 1ml PBT + 50µl FBS (fetal bovine serum) solution. The samples were then incubated with the primary antibody: anti-ubiquitin (mouse, 1:500, Merck, ST1200) at 5°C for 2 days. Excess primary antibody was washed out with PBT (3x 20 minutes), followed by another blocking for 20 minutes, after which we incubated the brains with the secondary antibody: Alexa Fluor 488 anti-mouse (1:500, Life Technologies, A11001) for 2 hours at room temperature. We also washed out the excess secondary antibody in several steps as described for the primary antibody, and then, during post-dissection, we removed the remaining tissue from other organs and tracheas from the surface of the brains before microscopy. The nuclei were labeled with Hoechst containing PBS:glycerol (1:4) nuclei stain.

## Western blot

To measure quantitative changes in certain proteins, we used western blotting. For the measurement, we isolated protein from the heads of 7-day-old female animals. The animals were anesthetized with ether and then decapitated. We collected 15-15 heads from each strain in Eppendorf tubes. Proteins were extracted from the heads in a 1:1 mixture of 96 µl Fly Lysis Buffer and 2xLaemmli buffer (Bio-Rad, 1610737) using a homogenizing rod and then stored at -20°C. We performed the Western blot according to the method described by Billes et al [28]. We applied 15µl of each sample. The samples were run on a 4-20% Mini-PROTEAN® TGX™ gradient gel and then blotted onto a nitrocellulose membrane (Kisker Biotech, 40520100). After that, we blocked the samples in 3% milk powder solution (BioRad 170–6404 Blotting-Grade Blocker) for 1 hour, and then applied the primary antibodies: anti-Ref(2)P/p62 (rabbit, 1:2000 [29]), α-Tub84B (internal control) (mouse, 1:1000, Sigma, T6199), anti-Atg8a (rabbit, 1:2000 [30], [31]). We incubated the sample in the antibody solutions at 5°C overnight. The next day, the excess (unbound) primary antibody was washed off the nitrocellulose with TBST, 3 x 10 min washes. Then we incubated the nitrocellulose in the secondary antibodies: anti-rabbit IgG alkaline phosphatase (1:1000, Sigma, A3687), anti-mouse IgG alkaline phosphatase (1:1000, Sigma, A5153) for one hour. Detection was done with NBT-BCIP solution (1:50, Sigma, 72091). We used the ImageJ software to evaluate the results. The levels of p62 and Atg8a were corrected by the change measured for α-Tub84B. Western blotting is a semi-quantitative technique. Therefore, we only recorded the direction of changes, not their extent.

## Lifespan analysis

The goal of this experiment is to compare the average lifespan of each gene silenced strain to the wild type animals. The measurement was performed with 5 parallel tubes, 15 female and 15 male animals of each genotype were placed in vials. Every two days, we transferred the *Drosophila* to fresh feeding vials. Each day until all the fruit flies were dead, we recorded the number and sex of animals that died in each tube.

## Climbing test

In this experiment, we can compare the fitness and state of the nervous system of the studied animals as they age. In a small diameter, high glass tube, *Drosophila* start to climb upwards after being knocked down (negative geotaxis). The speed of perception and retention of the stimulus is proportional to the state of the fly's nervous system. The short-term climbing test measures the ability and speed of the animals to perceive the stimuli [32]. We measured the climbing ability of fruit flies at 7, 14, 21 and 28 days of age. Measurements were made with two parallel tubes. We anesthetized each genotype with $CO_2$, then separated the sexes and placed 10-10 animals in each tube. The animals were then rested at 29°C for 1 hour. After resting, the climbing test was repeated three times. There was a 30-minute break between each test. Each trial was videotaped and used to evaluate how each genotype was performing. Short-term climbing takes place to the first boundary line (6 cm) and measures how many of 10 animals reach the boundary line within 20 seconds.

## Statistics and data representation

We used RStudio (2021.09.0 Build 351) to perform the statistical analysis of the experimental data. The first step was to test the distribution of the data. In the case of a normal distribution, we used the t-test, in the case of a non-normal distribution, we used the Mann-Whitney U-test. For the results of the clonal systems, we performed the statistics with a paired t-test in the case of a normal distribution (Two Sample t-test if the two data set has similar variance, Welch Two Sample t-test if the two data set didn't have similar variance), and a paired Mann-Whitney U-test in the case of a non-normal distribution. The confidence level of significance is indicated by the stars on the graphs: **p<0,05: \*, p<0,01: \*\*, p<0,001: \*\*\***. Black stars indicate significance compared to wild type (*isowhite*) control; brown stars indicate significance compared to RNAi control (*GFP-RNAi*).

We created box plots using the RStudio program to represent the evaluated data. In the box plot, the box represents the values belonging to the interquartile range (50% of the data), and the black line in the boxes represents the median. The lower horizontal line shows the minimum value, and the upper horizontal line shows the maximum value. Outliers in the data set are indicated by empty circles. The black dots on the graph represent a value for a single sample, e.g. the evaluation of a microscopic image, except for the climbing essay, where one dot represents a measurement, e.g. the climbing ability of 10 animals in percentages. In the case of box plots, the data have been normalized. We generated the Kaplan-Meier curves of lifespan assay measurements and analyzed data using SPSS22.0 software. The data type of the lifespan assay cannot be displayed using boxplot, so we used a bar chart instead. The standard deviation is shown in the tracks at the top of the bars.

# Results

The amount of active Rab2 proteins can be enhanced by specific GAPs depletion

This part of our research focuses on the Rab2 protein, investigating the effect of silencing 12 GAP proteins potentially involved in the regulation of Rab2 and thus in the network regulating autophagic-degradation pathways *(Figure 1)*.

*Figure 1: GAP screen. Enhancing autophagic degradation efficiency through silencing regulatory GAPs. Rab2-specific GAP activates the hydrolysis of GTP, thereby converting membrane-bound, active Rab2 proteins into an inactive, soluble form (top figure). By silencing the right GAP, Rab2 remains active and the efficiency of autophagic degradation pathways can be increased (bottom figure). *In several organs, fusion of the autophagosome and vesicles containing Golgi-derived hydrolase enzymes with the lysosome are both linked to Rab2. Stimulation of both fusion events can increase degradation efficiency.*

According to our research plan, we started to screen which GAP silencing can increase the amount of active Rab2 proteins in *Drosophila* larval fat bodies and nervous system. We started with the fat body screen because the *Drosophila* larval fat body is a favored tissue for studying autophagy, since their cells are large and show changes in autophagic structures well. For this screen, we crossed the RNAi strains with *cgGal4; Rab2-YFP* virgin, female flies. The cgGal4 is specifically expressed in the larval fat body. We performed the assay in two conditions, using well-fed and starved animals. As a result of starvation, autophagy is activated in the animals, while under well-fed conditions, few autophagic vesicles can be detected [33]. Of the 12 genes 4 GAP-RNAi (*CG6182, CG8085, CG31935* and *CG42612*) were effective in well-fed larvae *(Figure 2: A-B)*, and in starving larvae 6 (*CG1695, CG8085, CG11727, CG12241, CG31935, CG42795*) *(Figure 2: B)* of them were able to significantly increase the amount of active Rab2 proteins. In the well-fed larvae, the GAP silenced strains showed some interesting patterns. While there were hardly any Rab2-positive vesicles in the control strain, many small structures appeared after silencing of *CG8085 (Figure 2: D')*. In *CG31935*, much fewer, but huge vesicles were visible, mainly at the periphery of the cells *(Figure 2: D")*. Also in fat-bodies, we generated clonal cells by somatic recombination using the Act<CD2<Gal4,GFP (V40) system

*(Table 1)*. Because of the mechanism described in the materials and methods section, the Gal4 protein sequence driving the GFP and RNA interference genes is expressed only in clonal cells in these animals. Surrounding cells in which the Gal4 protein is not transcribed are considered wild-type control cells. The amount of acidic degradative vesicles was assessed using LysoTracker-Red staining. LysoTracker-Red marks such acidic organelles as lysosomes, autolysosomes and late endosomes. As with the Rab2 screen in fat bodies, we also tested the larvae here in both well-fed and starving conditions. In well-fed animals, silencing of 4 GAP genes (*CG1695*, *CG7324*, *CG11727*, *CG42612*) increased the amount of acidic vesicles in the clone cells *(Figure 2: C)*. For starving larvae 2 GAP silencing (*CG11727* and *CG31935*) increased, 4 silencing (*CG1695*, *CG7324*, *CG11727*, *CG42612*) decreased the number of acidic vesicles in clone cells compared to control cells *(Figure 2: D)*. In some cases, there was a large increase in the number of acid-degrading vesicles *(Figure 2: D)*, which may indicate that the degradative processes were upregulated. This can be beneficial for breaking down harmful structures in cells.

*Figure 2: GAP silencing can increase the number of Rab2 positive structures in Drosophila brain and larval fat body. A-A"*: *Images of changes in the amount of Rab2-positive structures in larval fat cells due to certain GAP silencing. All three fluorescent images captured from well-fed larvae. **B**: Diagrams showing the evaluated data measured in from larval fat bodies after GAP silencing. In well-fed larvae: left, and in starving larvae: right plot. **C**: Clonal cells expressing GFP and RNA, silencing the gene of interest in the fat body of well-fed larvae, on the left. And the evaluated data on the right. **D:** Clonal cells showing a significant increase in acidic vesicles (red spots). The evaluated data is on the right. **Levels of significance: * = p<0.05; ** = p<0.01; *** = p<0.001***

Autophagic degradation may be enhanced by silencing regulating GAPs in the brain of Drosophila

In the neuron screen, the individual RNA interference lines were crossed with *DdcGal4; Rab2-YFP* animals. This tissue-specific promoter (Ddc) only allows Gal4 expression in the dopaminergic and serotonergic neurons of the brain [34]. Our focus was on the mushroom body (MB) region of the brain, which contains a large number of dopaminergic and serotonergic neurons and is suitable for modeling neurodegenerative diseases of old age [35-37]. We examined the cell body and axonal regions of neurons separately because neurons are highly polarized cells and there can be large differences between the two cell regions *(Figure 3: A-B')*. Of the 12 GAP proteins, 7 GAP silencing (*CG1695, CG5344, CG7324, CG11727, CG12241, CG31935* and *CG42795*) were able to increase the amount of active Rab2 protein at the cells axonal area, and 6 of them (*CG1695, CG5344, CG11727, CG12241, CG31935* and *CG42795*) at the pericarions *(Figure 3: C)*. These results of the two RNAi screen suggest that these genes may be regulators of Rab2 protein activity, either directly or indirectly, in

*Drosophila* neurons and larval fat bodies. Thereby their downregulation could be a cause of the increase of the Rab2 protein active form. Disabling these GAP genes might lead to enhanced autophagic-degradation of harmful structures in cells.

***Figure 3: GAP silencing can increase the number of Rab2 positive structures in Drosophila brain and larval fat body. A, A':*** *Fluorescent microscopic images of fruit fly dopaminergic and serotonergic neurons in the mushroom body. The images show changes in the number of Rab2-YFP positive vesicles (well separated green spots) in the axonal area of the neurons.* ***B, B':*** *Fluorescence microscopy images of changes in the abundance of Rab2-positive structures in the cell body of neurons.* ***C:*** *Boxplots depicting changes in the amount of active Rab2 protein in axonal and cell body regions following each gene silencing of GAPs.* ***Levels of significance: * = p<0.05; ** = p<0.01; *** = p<0.001***

Autophagy activity in the nervous system was assessed by measurement of the mCherry-Atg8a reporter protein (fluorescence microscopy) and the amount of ubiquitinated aggregates by immunohistochemistry. mCherry-Atg8a marks the autophagic vesicles from the forming phagophore to the autolysosome [38]. Atg8a (lipidated form) showed a significant decrease in

several cases (*CG1695*, *CG7324*, *CG11727*, *CG31935* and *CG42795*), which may indicate more efficient autophagic-vesicle fusion and degradation in silenced strains. This decrease in lipidated Atg8a is also coincides with our previous observations with Rab2 constitutive activation experiments [25]. But there was an exception, the silencing of *CG12241*, where a significant increase could be seen in contrast to other GAP-RNAi-s *(Figure 4: A)*. This could mean an upregulation of phagophore formation or a less efficient autophagic-vesicular degradation. Larger protein aggregates and defective organelles that can no longer be degraded by the proteasome are dealt with the macrophagic pathway, so measuring the amount of these structures can also be a good indicator [39]. The amount of ubiquitinated aggregates was significantly reduced in 10 GAP silenced strains (*CG1695*, *CG4041(II)*, *CG5344*, *CG7112*, *CG7324*, *CG8085*, *CG11727*, *CG31935*, *CG42612* and *CG42795*) in *Drosophila* brain *(Figure 4: B)*. These results could be explained by the increased activity of autophagy, which eliminates the ubiquitinated structures marked for degradation. With Western blot we determined the amount of Atg8a (soluble and lipidated form) and Ref(2)P (/p62) proteins isolated from the heads of *Drosophila* from 5 chosen genotype (*CG1695*, *CG7324*, *CG8085*, *CG31935* and *CG42795*), and the level of α-Tub84B was also measured as a control *(Figure 4: C)*. There are two forms of the Atg8a protein, the soluble (Atg8a-I) and the lipidated, membrane-bound (Atg8a-II) which is the active form [40]. The ortholog of this protein is the human LC3B [33]. On the gel picture, the level of Atg8a-II (lipidated form) is consistent with the microscopic results, the amount of protein decreases in the RNAi strains. Ref(2)P/p62 protein is degraded during autophagic-lysosomal degradation, therefore it is a good indicator of the efficacy of autophagy [38]. The p62 level is also decreased in all RNAi strains (taking into account the calculation corrected for the amount of α-Tub84B), which according to the evaluation is significant in the case of *CG1695* and *CG42795*, suggesting that autophagy is activated in these cases.

***Figure 4: Silencing of certain GAP genes in the Drosophila brain may lead to increased autophagic activity. A:*** *Presence of mCherry-Atg8a (lipidated form) containing autophagic structures in Drosophila brain imaged by fluorescence microscopy. The well separated red spots are the individual autophagic vesicles labeled with the fusion protein marker. And the boxplot representing the evaluated data.* ***B:*** *Presence of ubiquitinated aggregates in fruit fly brains labeled by immunohistochemistry after GAP silencing. The boxplot presenting the evaluated data of each silenced strains. Levels of significance:* \* *= p<0.05;* \*\* *= p<0.01;* \*\*\* *= p<0.001. Black* \*: *significance compared to wild type animals. Brown* \*: *significance compared to RNAi control animals.* ***C:*** *Western blot image of neural protein isolated from 5 chosen GAP silenced strains. Atg8a I = soluble form of Atg8a; Atg8a II = lipidated, bound form of Atg8a.*

Neuron specific silencing of GAPs can extend lifespan in fruit flies

For physiological studies, we selected 5 promising GAP silenced strains (*CG1695*, *CG7324*, *CG8085*, *CG31935* and *CG42795*) that performed well in previous experiments and started with these first. The constitutively active form of the Rab2 protein was able to increase the lifespan of *Drosophila*. If we succeed in silencing GAPs that inactivate Rab2, we can expect a similar result. We also performed the silencing specifically for dopaminergic and serotonergic neurons. Thus, the effects of other tissues or neuron types do not interfere with the experiment. In both sexes *CG8085-* and *CG42795*-RNAi significantly increased the fruit flies average lifespan *(Figure 3: A, A')*. Silencing of the other three GAP genes only shows changes in one sex. *CG7324*-RNAi increases the average lifespan of males, but there is no significant change in the lifespan of females. *CG1695*-RNAi shows a significant decrease in males and an

increasing trend in females. *CG31935*-RNAi significantly increases lifespan in females but causes no changes in males. These results show that some GAP silencing can increase the average lifespan of *Drosophila*, but for other genes there are differences between sexes. For the 5 GAP silenced strains mentioned above, we also performed a climbing test. This experiment will show the state of the neurons in each of the studied strains studied at different ages. This is tested by observing how quickly the animals respond to the stimulus and how they can retain it [32]. Some GAP silencing improved, while in one case the climbing ability of the animals was significantly impaired in old age. In females, *CG7324*- and *CG1695*-RNAi significantly increased climbing ability, however in males these only could make an improvement compared to the RNA interference control. An additional gene-silencing (*CG42795*-RNAi) could enhance the climbing ability. Silencing *CG31935* significantly impairs the climbing ability of animals in both sexes. We can say that silencing of certain GAP genes may be able to increase climbing ability in old animals, but we found large differences between the sexes and some silencing had the opposite negative effect.

*Figure 3: Silencing of CG8085 and CG42795 lead to an increased average lifespan of these Drosophila strains compared to wild type animals. A: Kaplan-Meier lifespan curves of individual gene-silenced strains. Curves showing significant increases in both sexes are highlighted by the smaller graphs in the lower left corner. A': Average lifespan plots for each*

*mutant strain, females at top, males at bottom. **B:** Changes in short-term climbing ability of female, GAP silenced animals. **B':** Changes in short-term climbing ability of male, GAP silenced Drosophila. Levels of significance: \* = p<0.05; \*\* = p<0.01; \*\*\* = p<0.001. Black \*: significance compared to wild type animals. Brown \*: significance compared to RNAi control animals.*

## Discussions

In the present study, we investigated the role of 12 GTPase-activating proteins (GAPs) belonging to the TBC1 domain family in Rab2 and autophagy regulation. Silencing of several GAPs was found to affect the amount of membrane-bound form of Rab2 (Figure 2 and Figure 3). The role of GAPs was investigated in larval fat bodies and adult animals' nervous system. Silencing of *CG1695*, *CG11727*, and *CG31935* genes in both organs resulted in Rab2 activation. However, in several cases, differences were found between GAP silencing in larval siliques and in adult brains. *CG5344*-RNAi and *CG7324*-RNAi only increased the number of Rab2-positive structures in the nervous system. In contrast, decreased expression of *CG6182*, *CG8085*, *CG42612* altered the amount of YFP-Rab2 reporter only in fat bodies.

The scilencing of GAPs, it was found that in the nervous system, there is no significant difference between Rab2 positive vesicles measured in the cell body and axonal sections (Figure 3). Alone, *CG7324*-RNAi caused a considerable increase in Rab2 only in the axonal part, but the enhancing effect also showed a tendency to increase in the nuclear part of the neurons. However, we observed significant Rab2 regulatory differences in GAP silencing in fat bodies between fed and starved conditions (Figure 2). *CG6185* and *CG42612* increased the amount of Rab2-positive structures in well-fed samples only. In contrast, silencing of *CG4041*, *CG11727*, *CG12241*, and *CG42795* genes increased Rab2 activity only in starved samples (lysosome amount was significantly increased by *CG42612*, *CG11727*, and *CG31935* silencing in both conditions tested). Similar condition-dependent differences were also observed for other autophagy regulators. The *Drosophila* orthologue of myotubularin-type phosphatase 14 (MTMR14) inhibits basal autophagy exclusively in fat bodies, whereas Mtmr6 (common orthologue of human MTMR6,-7,-8) is a specific inhibitor of stress-induced autophagy [41]. Comparing the two examined organs, the roles of two GAPs, *CG11727* and *CG31935*, seem to be common. While, for example, *CG42795* is only able to play its role in the nervous system. According to the Fly Cell Atlas, *CG42795* is expressed at lower levels in larval tissues, and in adult animals it is most intensively expressed in the nervous system (inter-, motor-, sensory-neurons and glia cells). This may explain why its silencing had a phenotypic result mainly in the nervous system. The expression of *CG11727* is high in the adult nervous system and in L3 larvae [42], [43]. In *Drosophila*, Rab2 is known to regulate the fusion of various types of vesicles with lysosomes and their transport in different organs. In fat bodies, Rab2 regulates the fusion of Golgi derived vesicles with lysosomes but does not affect autophagosome - lysosome fusion [44]. In contrast, in larval neurons, eye primordia and muscle, Rab2 promotes autophagosome-lysosome fusion [45], [46]. We investigated the effects of 5 selected GAPs silencing on the animals' lifespan and locomotion ability. *CG8085*-RNAi and *CG42795*-RNAi extended the lifespan in both sexes, whereas *CG7324*-RNAi had a positive effect only in males

(Figure 4). The silencing of *CG42795* and *CG7324* positively affected the animals' locomotion ability, compared to *CG8085*-RNAi, which impaired locomotion ability. Taken together, posttranscriptional inhibition of *CG42795* was the most beneficial for activating neuronal autophagy and vitality in animals. In the future, it could be useful to investigate the role of the human orthologue of the gene TBC1D30 (TBC domain family member 30) in activating Rab2 in mammals. Experimental studies have not previously investigated the regulatory role of TBC1D30 in autophagy (Rab2), but it plays a critical role in the processing of mature insulin [47]. It could also be a negative regulator of Rab8 and Rab3 smallGTPases [48], [49]. Rab8 is responsible for transport between the trans-Golgi network and the plasma membrane, while Rab3 regulates synaptic vesicle storage and neurotransmitter release in synapses [50], [51]. In the future, it could also be interesting to investigate the role of *CG42795* in the regulation of *Drosophila* Rab3 and Rab8 orthologs. Further study of *CG42795* and TBC1D30 may also help to understand the regulation of autophagy and Rab-specific small GTPases. Additional experiments may reveal a new potent lifespan and autophagy regulatory point.

The *Drosophila* TBC1 domain family includes 25 proteins, of which we have investigated the roles of 12. It might be worthwhile in the future to investigate the regulatory role of the additional 13 GAPs on Rab2 and autophagy. In our previous research, we have also investigated the effects of the neuronal activation of Rab7 and Arl8. Arl8 was found to have a positive effect on neuronal activation similar to Rab2 activation. Autophagy was activated in animals, locomotor function improved, and lifespan extended. In contrast, overexpression of the constitutively active form of Rab7 in dopaminergic and serotonergic neurons had the opposite effect to the activation of Rab2 and Arl8 [25]. In the future, it might be worth investigating the regulatory points of Arl8 and Rab7 similarly to our current study and finding regulatory junctions that can co-activate Arl8a and Rab2 without affecting Rab7 activity.

Our results have highlighted that several GAPs can regulate a small GTPase. Similarly, the same GAP can inhibit many different small GTPases. Therefore, in the future, it is worthwhile to investigate the effects of the functions of the investigated GAPs on other Rab enzymes and to explore the compensatory mechanisms among GAPs with the same regulatory function.

## Author Contributions

G.F. and F.K. performed experiments on *Drosophila* samples, analysed data and wrote the manuscript. T.M.B. performed experiments on *Drosophila* samples and analysed data. T.K. designed experiments, analysed data, and wrote the manuscript.

## Fundings

This work was supported by the OTKA grant (Hungarian Scientific Research Fund) PD 143786 to TK. TK supported by the University Excellence Fund of Eötvös Loránd University, Budapest, Hungary (ELTE). GF and FK were supported by the DKOP-23 Doctoral Excellence Program of the Ministry for Culture and Innovation from the source of the National Research, Development and Innovation Fund.

## Acknowledgments

*Drosophila* strains and reagents were kindly provided by Gábor Juhász. The authors also thank Regina Preisinger, Beatrix Supauer, Erzsébet Gatyás for the excellent technical assistance.

## References

[1]     N. Mizushima, B. Levine, A. M. Cuervo, and D. J. Klionsky, "Autophagy fights disease through cellular self-digestion.," *Nature*, vol. 451, no. 7182, pp. 1069–1075, Feb. 2008, doi: 10.1038/nature06639.

[2]     B. Levine and G. Kroemer, "SnapShot: Macroautophagy.," *Cell*, vol. 132, no. 1, pp. 162.e1-162.e3, Jan. 2008, doi: 10.1016/j.cell.2007.12.026.

[3]     L. Wang, D. J. Klionsky, and H.-M. Shen, "The emerging mechanisms and functions of microautophagy.," *Nat. Rev. Mol. Cell Biol.*, vol. 24, no. 3, pp. 186–203, Mar. 2023, doi: 10.1038/s41580-022-00529-z.

[4]     A. C. Massey, C. Zhang, and A. M. Cuervo, "Chaperone-mediated autophagy in aging and disease.," *Curr. Top. Dev. Biol.*, vol. 73, pp. 205–235, 2006, doi: 10.1016/S0070-2153(05)73007-6.

[5]     D. J. Klionsky and P. Codogno, "The mechanism and physiological function of macroautophagy.," *J. Innate Immun.*, vol. 5, no. 5, pp. 427–33, 2013, doi: 10.1159/000351979.

[6]     M. Graef, J. R. Friedman, C. Graham, M. Babu, and J. Nunnari, "ER exit sites are physical and functional core autophagosome biogenesis components.," *Mol. Biol. Cell*, vol. 24, no. 18, pp. 2918–31, Sep. 2013, doi: 10.1091/mbc.E13-07-0381.

[7]     M. Hamasaki *et al.*, "Autophagosomes form at ER-mitochondria contact sites.," *Nature*, vol. 495, no. 7441, pp. 389–93, Mar. 2013, doi: 10.1038/nature11910.

[8]     H. Nakatogawa, "Mechanisms governing autophagosome biogenesis.," *Nat. Rev. Mol. Cell Biol.*, vol. 21, no. 8, pp. 439–458, 2020, doi: 10.1038/s41580-020-0241-0.

[9]     M. Laplante and D. M. Sabatini, "mTOR signaling at a glance," *J. Cell Sci.*, vol. 122, no. 20, pp. 3589–3594, Oct. 2009, doi: 10.1242/jcs.051011.

[10]    E. L. Axe *et al.*, "Autophagosome formation from membrane compartments enriched in phosphatidylinositol 3-phosphate and dynamically connected to the endoplasmic reticulum.," *J. Cell Biol.*, vol. 182, no. 4, pp. 685–701, Aug. 2008, doi: 10.1083/jcb.200803137.

[11]    J. Cheng *et al.*, "Yeast and mammalian autophagosomes exhibit distinct phosphatidylinositol 3-phosphate asymmetries.," *Nat. Commun.*, vol. 5, p. 3207, 2014, doi: 10.1038/ncomms4207.

[12]    H. Nakatogawa, K. Suzuki, Y. Kamada, and Y. Ohsumi, "Dynamics and diversity in autophagy mechanisms: lessons from yeast.," *Nat. Rev. Mol. Cell Biol.*, vol. 10, no. 7, pp. 458–67, Jul. 2009, doi: 10.1038/nrm2708.

[13]    D. J. Klionsky *et al.*, "Autophagy in major human diseases.," *EMBO J.*, vol. 40, no. 19, p. e108863, Oct. 2021, doi: 10.15252/embj.2021108863.

[14]    Y. Aman *et al.*, "Autophagy in healthy aging and disease.," *Nat. aging*, vol. 1, no. 8,

pp. 634–650, Aug. 2021, doi: 10.1038/s43587-021-00098-4.

[15] A. Donati *et al.*, "Age-related changes in the autophagic proteolysis of rat isolated liver cells: effects of antiaging dietary restrictions.," *J. Gerontol. A. Biol. Sci. Med. Sci.*, vol. 56, no. 9, pp. B375-83, Sep. 2001, doi: 10.1093/gerona/56.9.b375.

[16] M. L. Tóth *et al.*, "Longevity pathways converge on autophagy genes to regulate life span in Caenorhabditis elegans.," *Autophagy*, vol. 4, no. 3, pp. 330–8, Apr. 2008, doi: 10.4161/auto.5618.

[17] T. Vellai, M. L. Tóth, and A. L. Kovács, "Janus-faced autophagy: a dual role of cellular self-eating in neurodegeneration?," *Autophagy*, vol. 3, no. 5, pp. 461–463, 2007, doi: 10.4161/auto.4282.

[18] A. Simonsen, R. C. Cumming, A. Brech, P. Isakson, D. R. Schubert, and K. D. Finley, "Promoting basal levels of autophagy in the nervous system enhances longevity and oxidant resistance in adult Drosophila.," *Autophagy*, vol. 4, no. 2, pp. 176–84, Feb. 2008, doi: 10.4161/auto.5269.

[19] G. Konstantinidis and N. Tavernarakis, "Molecular Basis of Neuronal Autophagy in Ageing: Insights from Caenorhabditis elegans.," *Cells*, vol. 10, no. 3, Mar. 2021, doi: 10.3390/cells10030694.

[20] T. Kovács *et al.*, "A conserved MTMR lipid phosphatase increasingly suppresses autophagy in brain neurons during aging.," *Sci. Rep.*, vol. 12, no. 1, p. 21817, Dec. 2022, doi: 10.1038/s41598-022-24843-w.

[21] D. M. Izmaylov and L. K. Obukhova, "Geroprotector effectiveness of melatonin: investigation of lifespan of Drosophila melanogaster.," *Mech. Ageing Dev.*, vol. 106, no. 3, pp. 233–240, Jan. 1999, doi: 10.1016/s0047-6374(98)00105-5.

[22] H. Stenmark, "Rab GTPases as coordinators of vesicle traffic.," *Nat. Rev. Mol. Cell Biol.*, vol. 10, no. 8, pp. 513–25, Aug. 2009, doi: 10.1038/nrm2728.

[23] S. Toma-Fukai and T. Shimizu, "Structural Insights into the Regulation Mechanism of Small GTPases by GEFs.," *Molecules*, vol. 24, no. 18, Sep. 2019, doi: 10.3390/molecules24183308.

[24] J. L. Bos, H. Rehmann, and A. Wittinghofer, "GEFs and GAPs: critical elements in the control of small G proteins.," *Cell*, vol. 129, no. 5, pp. 865–877, Jun. 2007, doi: 10.1016/j.cell.2007.05.018.

[25] J. Szinyákovics, F. Keresztes, E. A. Kiss, G. Falcsik, T. Vellai, and T. Kovács, "Potent New Targets for Autophagy Enhancement to Delay Neuronal Ageing.," *Cells*, vol. 12, no. 13, Jun. 2023, doi: 10.3390/cells12131753.

[26] N. Alic, M. P. Hoddinott, A. Foley, C. Slack, M. D. W. Piper, and L. Partridge, "Detrimental effects of RNAi: a cautionary note on its use in Drosophila ageing studies.," *PLoS One*, vol. 7, no. 9, p. e45367, 2012, doi: 10.1371/journal.pone.0045367.

[27] J. E. A. Lee, N. J. Cranna, A. S. Chahal, and L. M. Quinn, "Genetic systems to investigate regulation of oncogenes and tumour suppressor genes in Drosophila.," *Cells*, vol. 1, no. 4, pp. 1182–1196, Dec. 2012, doi: 10.3390/cells1041182.

[28] V. Billes *et al.*, "AUTEN-67 (Autophagy Enhancer-67) Hampers the Progression of

Neurodegenerative Symptoms in a Drosophila model of Huntington's Disease.," *J. Huntingtons. Dis.*, vol. 5, no. 2, pp. 133–47, 2016, doi: 10.3233/JHD-150180.

[29] K. Pircs *et al.*, "Advantages and Limitations of Different p62-Based Assays for Estimating Autophagic Activity in Drosophila," *PLoS One*, vol. 7, no. 8, 2012, doi: 10.1371/journal.pone.0044214.

[30] S. Takáts *et al.*, "Autophagosomal Syntaxin17-dependent lysosomal degradation maintains neuronal function in Drosophila," *J. Cell Biol.*, vol. 201, no. 4, pp. 531–539, 2013, doi: 10.1083/jcb.201211160.

[31] P. Nagy *et al.*, "Atg17/FIP200 localizes to perilysosomal Ref(2)P aggregates and promotes autophagy by activation of Atg1 in Drosophila," *Autophagy*, vol. 10, no. 3, pp. 453–467, 2014, doi: 10.4161/auto.27442.

[32] J. W. Gargano, I. Martin, P. Bhandari, and M. S. Grotewiel, "Rapid iterative negative geotaxis (RING): a new method for assessing age-related locomotor decline in Drosophila.," *Exp. Gerontol.*, vol. 40, no. 5, pp. 386–95, May 2005, doi: 10.1016/j.exger.2005.02.005.

[33] V. B. Varga, F. Keresztes, T. Sigmond, T. Vellai, and T. Kovács, "The evolutionary and functional divergence of the Atg8 autophagy protein superfamily.," *Biol. Futur.*, vol. 73, no. 4, pp. 375–384, Dec. 2022, doi: 10.1007/s42977-022-00123-6.

[34] T. Riemensperger *et al.*, "A single dopamine pathway underlies progressive locomotor deficits in a Drosophila model of Parkinson disease.," *Cell Rep.*, vol. 5, no. 4, pp. 952–60, Nov. 2013, doi: 10.1016/j.celrep.2013.10.032.

[35] H. Bolus, K. Crocker, G. Boekhoff-Falk, and S. Chtarbanova, "Modeling Neurodegenerative Disorders in Drosophila melanogaster.," *Int. J. Mol. Sci.*, vol. 21, no. 9, Apr. 2020, doi: 10.3390/ijms21093055.

[36] A. Mershin, E. Pavlopoulos, O. Fitch, B. C. Braden, D. V Nanopoulos, and E. M. C. Skoulakis, "Learning and memory deficits upon TAU accumulation in Drosophila mushroom body neurons.," *Learn. Mem.*, vol. 11, no. 3, pp. 277–87, 2004, doi: 10.1101/lm.70804.

[37] Y. Aso *et al.*, "The neuronal architecture of the mushroom body provides a logic for associative learning.," *Elife*, vol. 3, p. e04577, Dec. 2014, doi: 10.7554/eLife.04577.

[38] P. Lőrincz, C. Mauvezin, and G. Juhász, "Exploring Autophagy in Drosophila.," *Cells*, vol. 6, no. 3, Jul. 2017, doi: 10.3390/cells6030022.

[39] H. Liu *et al.*, "Aberrant mitochondrial morphology and function associated with impaired mitophagy and DNM1L-MAPK/ERK signaling are found in aged mutant Parkinsonian LRRK2(R1441G) mice.," *Autophagy*, vol. 17, no. 10, pp. 3196–3220, Oct. 2021, doi: 10.1080/15548627.2020.1850008.

[40] N. Mizushima and T. Yoshimori, "How to interpret LC3 immunoblotting.," *Autophagy*, vol. 3, no. 6, pp. 542–545, 2007, doi: 10.4161/auto.4600.

[41] A. Manzéger *et al.*, "Condition-dependent functional shift of two Drosophila Mtmr lipid phosphatases in autophagy control.," *Autophagy*, vol. 17, no. 12, pp. 4010–4028, 2021, doi: 10.1080/15548627.2021.1899681.

[42] H. Li *et al.*, "Fly Cell Atlas: A single-nucleus transcriptomic atlas of the adult fruit

fly.," *Science*, vol. 375, no. 6584, p. eabk2432, Mar. 2022, doi: 10.1126/science.abk2432.

[43] J. B. Brown *et al.*, "Diversity and dynamics of the Drosophila transcriptome.," *Nature*, vol. 512, no. 7515, pp. 393–399, Aug. 2014, doi: 10.1038/nature12962.

[44] P. Lörincz *et al.*, "Rab2 promotes autophagic and endocytic lysosomal degradation," *J. Cell Biol.*, vol. 216, no. 7, pp. 1937–1947, 2017, doi: 10.1083/jcb.201611027.

[45] V. K. Lund, K. L. Madsen, and O. Kjaerulff, "Drosophila Rab2 controls endosome-lysosome fusion and LAMP delivery to late endosomes.," *Autophagy*, vol. 14, no. 9, pp. 1520–1542, 2018, doi: 10.1080/15548627.2018.1458170.

[46] N. Fujita *et al.*, "Genetic screen in drosophila muscle identifies autophagy-mediated T-tubule remodeling and a Rab2 role in autophagy," *Elife*, vol. 6, pp. 1–30, 2017, doi: 10.7554/eLife.23367.

[47] V. A. PARSONS, S. VADLAMUDI, A. H. MOXLEY, and K. L. MOHLKE, "25-OR: Role for TBC1D30 in Secretion of Mature Insulin," *Diabetes*, vol. 70, no. Supplement_1, pp. 25-OR, Jun. 2021, doi: 10.2337/db21-25-OR.

[48] J. Peränen, "Rab8 GTPase as a regulator of cell shape.," *Cytoskeleton (Hoboken).*, vol. 68, no. 10, pp. 527–539, Oct. 2011, doi: 10.1002/cm.20529.

[49] J. Kolic *et al.*, "Differential Effects of Voclosporin and Tacrolimus on Insulin Secretion From Human Islets.," *Endocrinology*, vol. 161, no. 11, Nov. 2020, doi: 10.1210/endocr/bqaa162.

[50] L. A. Huber, S. Pimplikar, R. G. Parton, H. Virta, M. Zerial, and K. Simons, "Rab8, a small GTPase involved in vesicular traffic between the TGN and the basolateral plasma membrane.," *J. Cell Biol.*, vol. 123, no. 1, pp. 35–45, Oct. 1993, doi: 10.1083/jcb.123.1.35.

[51] G. Fischer von Mollard *et al.*, "rab3 is a small GTP-binding protein exclusively localized to synaptic vesicles.," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 87, no. 5, pp. 1988–1992, Mar. 1990, doi: 10.1073/pnas.87.5.1988.