1   **A transcription factor quintet orchestrating bundle sheath expression in rice**

3   Lei Hua*, Na Wang, Susan Stanley, Ruth M. Donald, Satish Kumar Eeda, Kumari Billakurthi, Ana

4   Rita Borba and Julian M. Hibberd*

6   Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge CB2 3EA,

7   United Kingdom

*For correspondence

Email for contact: jmh65@cam.ac.uk, lh556@cam.ac.uk

Keywords: *Oryza sativa*; C_4 rice; Bundle Sheath; Sulfite Reductase; Promoter; cis-elements

Abstract

C_4 photosynthesis has evolved in over sixty plant lineages and improves photosynthetic efficiency

by ~50%. One unifying character of C_4 plants is photosynthetic activation of a compartment such

as the bundle sheath, but gene regulatory networks controlling this cell type are poorly understood.

In Arabidopsis a bipartite MYC-MYB transcription factor module restricts gene expression to these

cells but in grasses the regulatory logic allowing bundle sheath gene expression has not been

defined. Using the global staple and C_3 crop rice we identified the *SULFITE REDUCTASE*

promoter as sufficient for strong bundle sheath expression. This promoter encodes an intricate *cis*-

regulatory logic with multiple activators and repressors acting combinatorially. Within this

landscape we identified a distal enhancer activated by a quintet of transcription factors from the

WRKY, G2-like, MYB-related, IDD and bZIP families. This module is necessary and sufficient to

pattern gene expression to the rice bundle sheath. Oligomerisation of the enhancer and fusion to

core promoters containing Y-patches allowed activity to be increased 220-fold. This enhancer

generates bundle sheath-specific expression in *Arabidopsis* indicating deep conservation in

function between monocotyledons and dicotyledons. In summary, we identify an ancient, short,

and tuneable enhancer patterning expression to the bundle sheath that we anticipate will be useful

for engineering this cell type in various crop species.

Introduction

In plants and animals significant progress has been made in understanding transcription factor networks responsible for the specification of particular cell types. In animals, for example, homeobox transcription factors define the body plan of an embryo (Lewis 1978; Krumlauf 1994), and cardiac cell fate is specified by a collective of five transcription factors comprising Pnr and Doc that act as anchors for dTCF, pMad and Tin (Junion et al. 2012). In plants the INDETERMINATE DOMAIN (IDD) transcription factors work together with SCARECROW and SHORTROOT to specify endodermal formation in the root (Moreno-Risueno et al. 2015; Drapek et al. 2017), PHLOEM EARLY (PEAR) and VASCULAR-RELATED NAC DOMAIN (VND) transcription factors permit production of phloem and xylem vessel respectively (Kubo et al. 2005; Miyashima et al. 2019),and basic helix–loop–helix (bHLH) transcription factors determine differentiation of guard cells (MacAlister et al. 2006; Ohashi-Ito and Bergmann 2006; Pillitteri et al. 2006; Kanaoka et al. 2008). Moreover, transcription factor networks that integrate processes as diverse as responses to external factors such as pathogens and abiotic stresses(Nakashima et al. 2009; Tsuda and Somssich 2015), or internal events associated with the circadian clock (McClung 2006; Nagel and Kay 2012) and hormone signalling (Depuydt and Hardtke 2011; Verma et al. 2016) have also been identified. Transcription factor activity is decoded by short *cis*-acting DNA sequences known as enhancers. The binding of multiple transcription factors to enhancers thus controls transcription and the spatiotemporal patterning of gene expression. For example, the Block C enhancer interacts with the core promoter to activate expression of FLOWERING LOCUS T (FT) in long days (Adrian et al. 2010; Liu et al. 2014), and a distant upstream enhancer controls expression of the *TEOSINTE BRANCHED1* locus in maize responsible for morphological differences compared with the wild ancestor teosinte (Stam et al. 2002; Clark et al. 2006). In contrast to the above examples, transcription factors and cognate *cis*-elements responsible for the operation of cell types in grasses once specified have not been defined (Weber et al. 2016; Schmitz et al. 2022).

Given the increased specialisation of organs evident since the colonisation of land this lack of understanding of gene regulatory networks controlling cell specific gene expression is striking. For example, in the liverwort *Marchantia polymorpha* the photosynthetic thallus contains seven cell types (Wang et al. 2023), while leaves of *Oryza sativa* (rice) and *Arabidopsis thaliana* possess at least fifteen and seventeen populations of cells as defined by single-cell sequencing respectively (Wang et al. 2021). In leaves of these angiosperms, particular cell types are specialised for photosynthesis and so whilst photosynthesis gene expression is induced by light in all major cell types of the rice leaf the response is greater in spongy and palisade mesophyll cells compared with guard, mestome and bundle sheath cells (Swift et al. 2023). In the case of the bundle sheath, these cells carry out photosynthesis, but are specialised to allow water transport from veins to mesophyll, sulphur assimilation and nitrate reduction (Leegood 2008; Aubry et al. 2014b; Hua et al. 2021). And, strikingly in multiple lineages, the bundle sheath has been dramatically repurposed

during evolution to become fully photosynthetic and allow the complex C_4 pathway to operate

(Sage 2004).

Compared with the ancestral C_3 state, plants that use C_4 photosynthesis operate higher light,

water and nitrogen use efficiencies (Makino et al. 2003; Sage 2004; Mitchell and Sheehy 2006). It

is estimated that introducing the C_4 pathway into C_3 rice would allow a 50% increase in yield

(Mitchell and Sheehy 2006; Hibberd et al. 2008), but it requires multiple photosynthesis genes to

be expressed in the bundle sheath, including enzymes that decarboxylate C_4 acids to release CO_2

around RuBisCO, organic acid transporters, components of the Calvin-Benson-Bassham cycle,

RuBisCO activase, and enzymes of starch biosynthesis (Kajala et al. 2011; Aubry et al. 2014a;

Ermakova et al. 2021). In summary, although the bundle sheath is found in all angiosperms and

associated with multiple processes fundamental to leaf function, the molecular mechanisms

responsible for directing expression to this cell type, including in global staple crops, remain

undefined. We therefore studied the bundle sheath to better understand the complexity of gene

regulatory networks that operate to maintain function of a cell type once it has been specified. Rice

was chosen as it a global crop, and identifying how it patterns gene expression to the bundle

sheath could facilitate engineering of this cell type.

We hypothesized that analysis of endogenous patterns of gene expression in the rice bundle

sheath would allow us to identify a strong and early-acting promoter for this cell type. Once such a

promoter was identified we also hypothesised that it could be used to initiate an understanding of

the *cis*-regulatory logic that allows gene expression to be patterned to this cell type in grasses. We

tested twenty-five promoters from rice genes that transcriptome sequencing indicated were highly

expressed in these cells. Of these, four specified preferential expression in the bundle sheath, and

one derived from the *SULFITE REDUCTASE* (*SiR*) gene generated strong bundle sheath

expression from plastochron 3 leaves onwards. Truncation analysis showed that bundle sheath

expression pattern from the *SiR* promoter is mediated by a short distal enhancer and a pyrimidine

patch in the core promoter. This bundle sheath module is cryptic until other enhancers acting to

both constitutively activate and repress expression in mesophyll cells are removed. The enhancer

is composed of a quintet of *cis*-elements recognised by their cognate transcription factors from the

WRKY121, GLK2, MYBS1, IDD and bZIP families. These transcription factors act synergistically

and are sufficient to drive expression of the strong bundle sheath *SiR* promoter.

Results

The *SiR* promoter directs expression to the rice bundle sheath

To identify sequences allowing robust expression in rice bundle sheath cells we used data derived from laser capture microdissection of bundle sheath strands and mesophyll cells from mature leaves. Promoter sequence from seven of the most strongly expressed genes in bundle sheath strands (**Supplemental Figure 1A**) were cloned, fused to the β-glucoronidase (GUS) reporter and transformed into rice. Although five of these promoters (*MYELOBLASTOSIS*, *MYB*; *HOMOLOG OF E. COLI BOLA*, *bolA*; *GLUTAMINE SYNTHETASE 1*, *GS1*; *STRESS RESPONSEIVE PROTEIN*, *SRP*; *ACYL COA BINDING PROTEIN*, *ARP*) led to GUS accumulation, it was restricted to veins **(Supplemental Figure 1B, 1C)**. And, for the *SULFATE TRANSPORTER 3;1* and *3;3* (*SULT3;1* and *SULT3;3*) promoters, no staining was observed (**Supplemental Figure 1B, 1C**). The approach of cloning promoters from bundle sheath strands therefore appeared to be more efficient at identifying sequences capable of driving expression in veins. We therefore optimised a procedure allowing bundle sheath cells to be separated from veins (Hua and Hibberd, 2019) and produced high quality transcriptomes from mesophyll, bundle sheath and vascular bundles (Hua et al. 2021). From these data eighteen genes whose transcripts were more abundant in bundle sheath cells compared with both veins and mesophyll cells were identified (**Supplemental Figure 2A**). When the promoter from each gene was fused to GUS and transformed into rice, those from *ATP-SULFURYLASE 1B*, *ATPS1b*; *SULFITE REDUCTASE*, *SiR*; *HIGH ARSENIC CONTENT1.1*, *HAC1.1*; and *FERREDOXIN*, *Fd* were sufficient to generate expression in the bundle sheath (**Supplemental Figure 2B)**. However, *ATPS1b* and *Fd* also displayed weak activity in the mesophyll, and the *HAC1.1* promoter also led to GUS accumulation in epidermal and vascular cells. Thus, only the *SiR* promoter drove strong expression in the bundle sheath with no GUS detected in other cells (**Supplemental Figure 2B, 2C**). An additional six promoters (*SOLUBLE INORGANIC PYROPHOSPHATASE*, *PPase*; *PLASMA MEMBRANE INTRINSIC PROTEIN1;1*, Os*PIP1;1*; *PLASMA MEMBRANE INTRINSIC PROTEIN1;3*, Os*PIP1;3*; *ACTIN-DEPOLYMERIZING FACTOR*, *ADF*; *PEPTIDE TRANSPORTER PTR2*, *PTR2*; *NITRATE REDUCTASE2*, *NIA2*) generated expression in vascular bundles, and eight promoters produced no staining (**Supplemental Figure 2B, 2C**). In summary, most candidate promoters failed to generate expression that was specific to bundle sheath cells, but the region upstream of the rice *SiR* gene was able to do so. We therefore selected the *SiR* promoter for further characterization.

The *SiR* promoter drives strong and early expression in bundle sheath cells

Sequence upstream of the *SiR* gene comprising nucleotides -2571 to +42 relative to the predicted translational start site was sufficient to generate expression in the rice bundle sheath. To allow faster analysis of sequences responsible for this output we domesticated the sequence by removing four *Bsa*I and *Bpi*I sites such that it was compatible with the modular Golden Gate cloning system. When this modified sequence was placed upstream of the GUS reporter it also

generated bundle sheath preferential accumulation **(Figure 1A)**. Fusion to a nuclear-targeted
mTurquoise2 fluorescent protein confirmed that the *SiR* sequence was sufficient to direct
expression to bundle sheath cells, and also revealed expression in the longer nuclei of veinal cells
(Figure 1B). Expression from the domesticated and non-domesticated sequences was not
different **(Figure 1C)**. Compared with 0.58 nmol 4-MU/min/mg protein previously reported for the
Zoysia japonica PHOSPHOENOLCARBOXYKINASE (PCK) promoter (Emmerling 2018) activity
from the *SiR* promoter was at least 36% higher. Designer Transcription Activator-Like Effector
(dTALEs) and cognate Synthetic TALE-Activated Promoters (STAPs) amplify expression and allow
multiple transgenes to be driven from a single promoter (Brückner et al., 2015; Danila et al., 2022).
We therefore tested whether bundle sheath expression mediated by the *SiR* promoter is
maintained and strengthened by the dTALE-STAP system. Stable transformants showed bundle
sheath specific expression **(Supplemental Figure 3A, 3B)**, and GUS activity was ~18-fold higher
than that from the endogenous *SiR* promoter (**Supplemental Figure 3C**). We conclude that the
SiR promoter is compatible with the dTALE-STAP system and its activity can be strengthened. We
also investigated when promoter activity was first detected during leaf development and discovered
that GUS as well as fluorescence from mTurquoise2 were visible in 5-20mm long fourth leaves at
plastochron 3 (**Supplemental Figure 4**). This was not the case for the *ZjPCK* promoter even when
a dTALE was used to amplify expression (**Supplemental Figure 4**). We conclude that the *SiR*
promoter initiates expression in the bundle sheath before the *ZjPCK* promoter, and that it is also
able to sustain higher levels of expression in this cell type.

A distal enhancer and Y-patch necessary for expression in the bundle sheath
The *SiR* promoter contains a highly complex *cis* landscape **(Figure 1D)** comprising at least 638
predicted motifs from 56 transcription factor families (**Supplemental table 1**). We therefore
designed a 5' truncation series to investigate regions necessary for expression in the bundle
sheath (**Figure 1E**). Deleting nucleotides -2571 to -2180 and -1490 to -980 led to a statistically
significant reduction and then increase in MUG activity respectively but neither truncation
abolished preferential accumulation of GUS in the bundle sheath **(Figure 1E-F).** However, when
nucleotides -980 to -394 upstream of the predicted translational start site were removed GUS was
no longer detectable in bundle sheath cells (**Figure 1E-F**). Consistent with this, MUG assays
showed a statistically significantly reduction in activity when these nucleotides were absent **(Figure
1G)**. Thus, nucleotides spanning -980 to -394 of the *SiR* promoter are necessary for bundle sheath
specific expression.
To test whether this region is sufficient for bundle sheath specific expression we linked it to the
minimal CaMV35S core promoter. Although weak GUS signal was detected in a few veinal cells,
this was not the case for the bundle sheath **(Figure 1E-G)**. We conclude that sequence in two
regions of the promoter (from -394 to +42 and from -980 to -394) interact to specify expression to
the bundle sheath. To better understand this interaction we next generated unbiased 5' and 3'

deletions. This second deletion series further reinforced the notion that the *SiR* promoter contains
a complex *cis*-regulatory landscape. For example, when nucleotides -980 to -829 were removed
very weak GUS staining was observed and the MUG assay confirmed that activity was significantly
reduced to 11% that of the full-length promoter **(Figure 2, Supplemental Figure 5).** We conclude
that nucleotides -980 to -829 from the *SiR* promoter are necessary for tuning expression in the leaf.
When nucleotides -829 to -700 were removed GUS appeared in mesophyll cells **(Supplemental**
Figure 5). Truncating nucleotides -613 to -529 abolished GUS accumulation (**Supplemental**
Figure 5). The 3' deletion that removed nucleotides -251 to +42 also stopped accumulation of
GUS in both bundle sheath and mesophyll cells (**Figure 2A-C, Supplemental Figure 5)**. Notably,
when the distal region required for bundle sheath expression (-980 to -829) was combined with
nucleotides -251 to +42 these two regions were sufficient for patterning to this cell type (**Figure 2**).
Having identified a region in the *SiR* promoter that was necessary and sufficient for patterning to
the bundle sheath, we next used phylogenetic shadowing and yeast one hybrid analysis to better
understand the *cis*-elements and *trans*-factors responsible. Analysis of *cis*-elements in the *SiR*
promoter that are highly conserved in grasses identified a short region located from nucleotides -
588 to -539 that contained an *ETHYLENE INSENSITIVE3-LIKE 3* (*EIL3*) transcription factor
binding site (**Supplemental Figure 6A&B**). Whilst deletion of this motif had no detectable effect of
patterning to the bundle sheath (**Supplemental Figure 6C**) the level of expression was reduced
(**Supplemental Figure 6D**). We infer that the *EIL3* motif positively regulates activity of the *SiR*
promoter but is not responsible for cell specificity. These data are consistent with the promoter
truncation analysis that showed nucleotides -613 to -529 containing this motif were not required for
bundle sheath specific expression, but instead function as a constitutive activator (**Supplemental**
Figure 5). When yeast one hybrid was used to search for transcription factors capable of binding
the *SiR* promoter, sixteen were identified (**Supplemental Figure 7A, 7B**). For each, cognate
binding sites were present. This included TCP21 and OsOBF1 that can bind to TCP motifs and
Ocs/bZIP elements respectively. Consistent with the outcome of deleting the *EIL3* motif, three EIL
transcription factors interacted with nucleotides -899 to -500 (**Supplemental Figure 7B, 7C**).
Examination of transcript abundance in mature leaves showed that most of these transcription
factors were expressed in both bundle sheath and mesophyll cells (**Supplemental Figure 7D**)
implying that combinatorial interactions with cell specific factors are likely required for bundle
sheath specific expression from the *SiR* promoter.
207
The enhancer contains four subregions that simultaneously activate in bundle sheath and
repress in mesophyll cells
The truncation analysis above identified two short regions comprising nucleotides -980 to -829
and -251 to +42 that were necessary and sufficient for expression in the rice bundle sheath
(**Figure 2, Supplemental Figure 8**). Sequence spanning nucleotides -251 to +42 includes both the
annotated 5' untranslated region but also likely contains core promoter elements (**Supplemental**

Figure 9A). Re-analysis of publicly available data identified two major transcription start sites at
positions -91 (TSS1) and -41 (TSS2) (**Supplemental Figure 9A**). Although no canonical TATA-box
was evident in this region, a TATA-box variant was detected at position -130 (5'-ATTAAA-3') (Civáň
and Švec 2009) that could be responsible for transcription from TSS1. Moreover, upstream of TSS2
is a putative pyrimidine patch (Y-patch) that represents an alternate but common TC-rich core
promoter motif in plant genomes (Civáň and Švec 2009) (**Supplemental Figure 9A**). Scanning
sequence from -251 to +42 for core promoter elements also identified MTE (Motif Ten Element),
BREu (TFIIB Recognition Element upstream) and DCE-S-I (Downstream Core Element S-I) motifs
associated with eukaryotic core promoters (**Supplemental Figure 9B**). We therefore assume the
region upstream of TSS1 and TSS2 contains the core promoter elements. When consecutive
deletions to this sequence were made, statistically significant reductions in MUG activity were
evident but there was no impact on accumulation of GUS in the bundle sheath. Interestingly, when
the Y-patch was retained but the TATA-box like motif removed, low levels of GUS specific to the
bundle sheath were apparent (**Supplemental Figure 9C, 9D, 9E, 9F**), and deletion of the Y-patch
completely abolished GUS staining (**Supplemental Figure 9C, 9D, 9E, 9F**). Consistent with the Y-
patch being important for bundle sheath expression, when core promoters from other genes
(*PIP1;1, NRT1.1A*) containing a Y-patch were linked to the distal enhancer from *SiR* bundle sheath
expression was detected **(Figure 3A,3B)**, but this was not the case for genes with only a TATA-
box **(Figure 3A,3B)**. GUS activity was higher from the *PIP1;1* core promoter that contains more Y-
patches. Overall, we conclude that the TATA-box like motif is not required for expression in the
bundle sheath, but the Y-patch is necessary for this patterning and in combination with a distal
enhancer comprising nucleotides -980 to -829 it is sufficient for expression in this cell type.

We assessed the distal enhancer for transcription factor binding sites. The FIMO algorithm
identified motifs associated with WRKY, G2-like, MYB-related, MADS, DOF, IDD, ARR, SNAC
(Stress-responsive NAC) families. PlantPAN (Chow et al. 2018), which includes historically
validated *cis*-elements, found an additional Dc3 Promoter Binding Factor (DPBF) binding site for
group A bZIP transcription factors **(Figure 3D)**. Seven consecutive deletions spanning this
enhancer region and hereafter termed subregions a-g were generated **(Figure 3D)**. Although
veinal expression persisted when subregions a, b and d were absent, deletion of subregions a, b, d
and f resulted in loss of GUS from bundle sheath cells (**Figure 3E-3G**). MUG analysis showed that
deletion of all four regions significantly reduced promoter activity **(Figure 3G)**. In contrast, deletions
of nucleotides -938 to -923 (subregion c), -904 to 873 (subregion e), and -853 to -829 (subregion g)
had no impact on the patterning **(Supplemental Figure 10)**. The subregions necessary for
expression in the bundle sheath contained unique binding sites for WRKY, G2-like, MYB-related,
IDD, NAC and bZIP (DPBF) transcription factors. To examine the significance of these regions in
the context of full-length *SiR* promoter, consecutive deletions from subregion a to f were generated
(Supplemental Figure 11A). Deletion of subregion a, d or f, led to GUS accumulating primarily in
mesophyll cells whereas removal of subregion b, c or e, caused GUS staining in both mesophyll

cells and bundle sheath cells **(Supplemental Figure 11B)**. No significant changes in GUS activity

were observed in these deletion lines **(Supplemental Figure 11C)**. We conclude that that the

distal enhancer generates expression in the bundle sheath due to four distinct sub-regions, and

that nucleotides between -980 to -853 also function as repressors of mesophyll expression by

interacting with nucleotides -829 to -251.

WRKY, G2-like, MYB-related, IDD and bZIP transcription factors activate the distal enhancer

To gain deeper insight how the distal enhancer operates we employed multiple approaches

including transactivation assays, co-expression analysis and site directed mutagenesis. The distal

enhancer contained WRKY, G2-like, MYBR, IDD, SNAC and bZIP (DPBF) motifs **(Figure 4A,**

Supplemental Figure 12). We therefore cloned rice transcription factors from each family and

used them as effectors in transient assays **(Supplemental Figure 13)**. WRKY121, GLK2, MYBS1,

IDD2/3/4/6/10, and bZIP3/4/9/10/11 transcription factors led to the strongest activation of

expression from the bundle sheath enhancer **(Figure 4B**, **Supplemental Figure 14A-14D)**,

whereas the stress-responsive NAC transcription factors targeting a SNAC motif that overlaps a

bZIP (DPBF) motif, activated less strongly than bZIP factors **(Supplemental Figure 14E)**. We

therefore conclude that the SNAC motif is not important for activity of the bundle sheath enhancer.

Effector assays using pairwise combinations of transcription factors showed synergistic activation

from the distal enhancer when GLK2 and IDD3,4,6,10 were co-expressed **(Figure 4C)**.

Co-expression analysis using a cell-specific leaf developmental gradient dataset revealed that

GLK2, MYBS1 and IDD4,6,10 transcription factors that bind the G2-like, MYB-related and IDD

motifs respectively were more abundant in mesophyll cells **(Figure 4D)**. However, the bZIP9, IDD2

and WRKY121 transcription factors strongly correlated with *SiR* transcript abundance and were

preferentially expressed in bundle sheath cells **(Figure 4D)**. To test whether bZIP9, IDD2 and

WRKY121 are sufficient to pattern *SiR* expression to specific cells, we mis-expressed the single

transcription factor bZIP9, both bZIP9 and IDD2, and all three (bZIP9 and IDD2 and WRKY121) in

the mesophyll **(Supplemental Figure 15A, 15C, 15E)**. Mis-expression of bZIP alone induced GUS

expression from the bundle sheath enhancer in some mesophyll cells **(Figure 4E, Supplemental**

Figure 15B), and mis-expression of both bZIP9 and IDD2 induced greater expression in mesophyll

cells **(Figure 4E, Supplemental Figure 15D)**. Strikingly, the expression of bZIP9 and IDD2 and

WRKY121 in mesophyll cells fully activated expression in this cell type **(Figure 4E, Supplemental**

Figure 15F). We conclude that one or two transcription factors are weakly sufficient, but all three

together effectively interact with the distal enhancer in bundle sheath cells to drive *SiR* expression.

We next mutated WRKY, G2-like, MYB-related, IDD, and bZIP motifs. With the exception of the

WRKY site that had no statistically robust effect, mutations in each of these motifs diminished or

abolished enhancer activity in the bundle sheath **(Figure 5A-5C)**.

In order to test whether the WRKY, G2-like, MYB-related IDD, and bZIP (DPBF) sites are

sufficient to pattern expression to rice bundle sheath cells we concatenated them and fused them

to the core promoter of *SiR* (**Figure 5E**). GUS staining was evident in the bundle sheath (**Figure**

5F). Fusion to the *PIP1;1* core promoter maintained bundle sheath expression and resulted in an

~5 fold increase in activity (**Figure 5E&F**). Oligomerisation of the enhancer by repeating it three or

five times increased bundle sheath specific expression 23 or 58-fold respectively when fused to

SiR core promoter (**Figure 5E-5G**), and this effect was amplified 90 and 224-fold when fused with

the *PIP1;1* core (**Figure 5E-5G**). Synthetic promoters, created by oligomerising this enhancer and

combining it with core promoters that contain Y-patches enabled fine-tuning of bundle sheath-

specific expression in rice. When an oligomerised version of the enhancer was linked to the *SiR*

core promoter and placed in *A. thaliana*, it generated strong expression in bundle sheath cells

(**Figure 5H, Supplemental Figure 16**). Collectively our data indicate that transcription factors

belonging to the WRKY, G2-like, MYB-related, IDD and bZIP (DPBF) families act cooperatively to

decode distinct *cis*-elements in a distal enhancer of the *SiR* promoter, and that this transcription

factor collective represent an ancient and highly conserved mechanism allowing bundle sheath

specific gene expression in in both monocotyledons and dicotyledons.

Discussion

Expression of multiple genes in the rice bundle sheath is not associated with close upstream enhancers

Gene expression is determined by interactions between elements in the core promoter allowing basal levels of transcription (Juven-Gershon and Kadonaga 2009; Haberle and Stark 2018) with more distal *cis*-regulatory modules (Spitz and Furlong 2012; Shlyueva et al. 2014; Ray-Jones and Spivakov 2021). Such *cis*-regulatory modules include enhancers and silencers that act as hubs receiving input from multiple transcription factors and so allow gene expression to respond spatially and temporally to both internal and external stimuli (Li et al. 2007; Buecker and Wysocka 2012). After testing 25 promoters, we discovered that the majority were not capable of driving expression in the rice bundle sheath, and this included ten that generated no detectable activity of GUS in leaves. In all cases we had cloned sequence between -3191 and -960 nucleotides upstream of the predicted translational start site and so these data demonstrate that the core promoter and any enhancers in these regions are not sufficient to direct expression to rice bundle sheath cells. Combined with the paucity of previously reported promoters active in this cell type (Nomura et al. 2005a; Lee et al. 2021) these data argue either for long range upstream enhancers (Studer et al. 2011; Liu et al. 2015; Li et al. 2019; Yan et al. 2019; Zhao et al. 2022) or other regulatory mechanisms being important to specify expression in the bundle sheath. Possibilities include transcription factor binding sites in introns that impact on transcription start site and strongly enhance gene expression (Rose et al. 2008; Gallegos and Rose 2019; Rose 2019), or in exons where because such sequences specify amino acid sequence as well as binding of *trans*-factors, they have been termed duons (Stergachis et al. 2013). Functional analysis showed that duons can pattern expression to the bundle sheath of the C_4 plant *Gynandropsis gynandra* (Reyna-Llorens et al. 2018), and it is notable that a genome-wide analysis of transcription factor binding sites in grasses revealed genes preferentially expressed in bundle sheath cells tended to contain transcription factor binding sites in their coding sequence (Burgess et al. 2019). It therefore appears possible that gene expression in the bundle sheath is commonly encoded by non-canonical architecture perhaps based on duons rather than more traditional enhancer elements upstream of the core promoter.

Despite the above, we discovered four promoters capable of driving expression in the rice bundle sheath, and each was associated with a gene important in sulphur metabolism. For example, *ATPS*, *SiR* and *Fd* all participate in the first two steps of sulphate reductive assimilation, while *HAC1;1* encodes an arsenate reductase important in the detoxification of arsenate using glutathione that is a product of sulphur assimilation. Collectively, these data further support the notion that the rice bundle sheath cell is specialised in sulphur assimilation (Hua et al. 2021).

Two distinct genetic networks governing expression in bundle sheath cells

The only other promoter for which both *cis*-elements and *trans*-factors that are necessary and sufficient to pattern bundle sheath expression have been reported is from the dicotyledonous

model *A. thaliana*. Here, a bipartite MYC-MYB module upstream of the *MYB76* gene is responsible

for this output (Dickinson et al. 2020). MYB76 forms part of a network governing glucosinolate

biosynthesis in *A. thaliana*, and so it is notable that the gene regulatory network we report in rice is

also associated with sulphur metabolism. However, rather than the bipartite transcription factor

network that regulates bundle sheath expression in *A. thaliana*, in rice we report a quintet of

transcription factors controlling *SiR* **(Figure 6C&6D)**. The enhancer controlling bundle sheath *SiR*

expression in rice comprises four distinct regions recognised by transcription factors belonging to

the WRKY, G2-like, MYB-related, IDD and bZIP families **(Figure 6D)**. As loss of the G2-like, MYB-

related, IDD and bZIP motifs all reduced expression in the bundle sheath, this implies they act co-

operatively - a notion further supported by the fact that GLK2 and IDD3,4,6,10 synergistically

activated promoter output in a transient assay. It is of course possible that other motifs in the

enhancer such as MADS, DOFs and ARRs act as modulators to tune the level of bundle sheath

expression. In fact, single nuclei sequencing of rice and sorghum during photomorphogenesis

identified DOFs as important for the evolution of C_4 gene bundle sheath expression (Swift,

Luginbuhl et al. 2023). For the *PIP1;1* and *NRT1.1A* genes, whose transcripts preferentially

accumulate in the bundle sheath, the core promoters were not able to generate bundle sheath

expression, but they contain a Y-patch and the WRKY, G2-like, MYB-related, IDD and bZIP

enhancer is present in intronic sequence **(Supplemental Figure 17).** It is therefore possible that

this regulatory system controls their expression. Moreover, for two promoters from rice (*Fd* and

HAC1;1) and two from other species (*ZjPCK* and *FtGLDP*) that are sufficient to drive expression to

the bundle sheath contain Y-patches and the cognate *cis*-elements for WRKY, G2-like, MYB-

related, IDD and bZIP transcription factors (**Supplementary Figure 17**).

The distal enhancer in the *SiR* promoter operates in conjunction with the core promoter that

contains two transcription start sites, one with an upstream TATA-box and the other a TC-rich

element known as a pyrimidine (Y) patch **(Supplemental Figure 9A)**. The TATA-box is found in

metazoans and plants and allows recognition by the pre-initiation complex (Smale and Kadonaga

2003), but in plants computational analysis showed that many promoters lack a TATA-box and

instead contain a Y-patch (Yamamoto et al. 2007a, 2007b; Bernard et al. 2010). These genes tend

to be relatively steadily expressed and associated with protein metabolism (Bernard et al., 2010),

and presence of a Y-patch can increase core promoter strength (Jores et al. 2021). For *SiR,* whilst

the TATA-box is not required, the Y-patch is needed for expression in the bundle sheath. Notably,

core promoters with a higher number or longer Y-patches tended to drive stronger expression, and

showed that in plants cell specific gene expression can be tuned by selecting different core

promoters.

The regulatory network comprising the Y-patch and distal enhancer enabling bundle sheath

expression of *SiR* is embedded within a complex *cis*-regulatory landscape with distinct regions

encoding activating and repressing activities (**Figure 6A&6B**). For example, the distal enhancer

(nucleotides -980 to -829) activating expression in the bundle sheath overlaps with sequence

(nucleotides -900 to -700) that suppresses expression in the mesophyll. Notably, the distal
enhancer is both essential for mesophyll repression and also sufficient to drive bundle sheath
specific expression (**Figure 6A**). In addition to controlling cell specificity, this complexity likely also
facilitates the tuning of expression to environmental conditions. For instance, the *EIL* motif (position
-572 to -552) is recognised by *ETHYLENE-INSENSITIVE LIKE* transcription factors that respond to
sulphur deficiency (Maruyama-Nakashita et al. 2006; Dietzen et al. 2020). As transcripts encoding
EIL accumulate in both bundle sheath and mesophyll cells in response to sulphate deficiency it
seems likely that transcription factors repressing expression in the mesophyll respond in a dynamic
manner. In addition to *EIL*, the yeast one hybrid analysis identified seven other families of
transcription factor families that can bind the *SiR* promoter. Many play documented roles during
abiotic or biotic stress, with for example OBF1, ERF3, NAP and FLP acting during low-temperature
or drought responses (Shimizu et al. 2005; Zhang et al. 2013, 2022; Chen et al. 2014; Qu et al.
2022), while TCP21, EREBP1, ERF3, ERF72, ERF83 are involved in both abiotic and biotic stress
(Lin et al. 2007; Jisha et al. 2015; Zhang et al. 2016, 2018; Tezuka et al. 2019; Jung et al. 2021).
Consistent with previous *in silico* analysis (Kurt et al. 2022) the presence of multiple AP2/ERF and
EIL transcription factors binding sites suggests that *SiR* is likely subject to control from ethylene
signalling (Binder 2020) and also of transcription factors that respond to ABA and jasmonic acid
biosynthesis and signalling (Yaish et al. 2010; Chen et al. 2014; Jisha et al. 2015; Zhang et al.
2016, 2018). Together this implies that multiple phytohormone signalling pathways converge on
the *SiR* promoter. These data are similar to those reported for the *SHORTROOT* promoter in *A.*
thaliana roots where a complex network of activating and repressing *trans*-factors also tunes
expression (Sparks et al. 2016). It is also notable that the architecture we report for the bundle
sheath enhancer of *SiR* appears of similar complexity to the collective of five transcription factors
used to specify cardiac mesoderm in *Drosophila melanogaster* and vertebrates (Junion et al.,
2012). For the five transcription factors that bind the cardiac mesoderm enhancer, the order and
positioning of motifs (motif grammar) is flexible. However, this is not always the case, with for
example output from the human *interferon-beta* (*INF-β*) enhancer demanding a conserved
grammar (Thanos and Maniatis 1995; Panne 2008). Further work will be needed to determine if the
bundle sheath enhancer reported here for rice is more similar to one of these models, or indeed, as
reported for the Drosophila *eve stripe 2* enhancer, operates as a billboard in different tissues to
determine patterning of expression (Kulkarni and Arnosti 2003).
411

Using the *SiR* promoter to engineer the rice bundle sheath
In addition to bundle sheath cells being important for sulphur assimilation (Leegood 2008; Aubry
et al. 2014b; Hua et al. 2021) they have also been implicated in nitrate assimilation, the control of
leaf hydraulic conductance and solute transport (Hua et al. 2021) and the systemic response to
high light (Xiong et al. 2021). Moreover, in one of the most striking examples of a cell type being
repurposed for a new function, bundle sheath cells have repeatedly been rewired to allow the

evolution of C_4 photosynthesis (Sage 2004). To engineer these diverse processes, specific and

tuneable promoters for this cell are required. However, identification of sequence capable of driving

specific expression to bundle sheath strands has previously been limited to *A. thaliana* and C_4

species. For example, the *SCARECROW* (Cui et al. 2014), *SCL23* (Cui et al. 2014), *SULT2;2*

(Kirschner et al. 2018) and *MYB76* promoters (Dickinson et al. 2020) are derived from *A. thaliana*,

whilst the *Glycine Decarboxylase P-protein (GLDP)* promoter is from the C_4 dicotyledon *Flaveria*

trinervia (Engelmann et al. 2008; Wiludda et al. 2012)*.* In rice, only the C_4 *Zoysia japonica PCK*

and the C_4 *Flaveria trinervia GLDP* promoters are known to pattern expression to the bundle

sheath (Nomura et al. 2005c; Lee et al. 2021). Both are capable of conditioning expression in this

cell type, but are weak, turn on late during leaf development and the molecular basis underpinning

their ability to restrict expression to the bundle sheath has not been defined. It has therefore not

been possible to rationally design or tune expression to this important cell type in rice. The

architecture of the *SiR* promoter we report here now provides an opportunity to engineer the

bundle sheath.

In summary, from analysis of the ~2600 nucleotide *SiR* promoter we identify an enhancer

comprising 81 nucleotides that with the Y-patch is sufficient to drive expression to bundle sheath

cells. Moreover, we show that output can be tuned via two approaches. First, oligomerising the

distal enhancer can drastically increase expression. Second, combining it with different core

promoters achieved the same output, and correlated with length of the Y-patch present. Our

identification of a minimal promoter that drives expression in bundle sheath cells of rice now

provides a tool to allow this important cell type to be manipulated. Cell specific manipulation of

gene expression has many perceived advantages. For example, when constitutive promoters have

been used to drive gene expression gene silencing and reduction of plant fitness due to metabolic

penalties (Glick 1995; Que et al. 1997). In contrast, tissue specific promoters allow targeted gene

expression either spatially or at particular developmental stages and so allow increased precision

in trait engineering (Kummari et al. 2020). The *SiR* promoter and the bundle sheath *cis*-regulatory

module that we identify thus provide insights into mechanisms governing cell specific expression in

rice, and may also contribute to our ability to engineer and improve cereal crops.

Materials and methods

Plant material and growth conditions

Kitaake (*O. sativa* ssp. *japonica*) was transformed using *Agrobacterium tumefaciens* with the

following modifications as described previously (Hiei et al., 2008). Mature seeds were sterilized

with 2.5% (v/v) sodium hypochlorite for 15 mins, and calli were induced on NB medium with 2 mg/L

2,4-D at 30 ^{o}C in darkness for 3-4 weeks. Actively growing calli were then co-incubated with *A.*

tumefaciens strain LBA4404 in darkness at 25^{o}C for 3 days, they were selected on NB medium

supplied with 35 mg/L hygromycin B for 4 weeks, and those that proliferated placed on NB medium

with 10 mg/L hygromycin B for 4 weeks at 28 ^{o}C under continuous light. Plants resistant to

hygromycin were planted in 1:1 mixture of topsoil and sand and placed in a greenhouse at the

Botanic Gardens, University of Cambridge under natural light conditions but supplemented with a

minimum light intensity of 390 µmol m^{-2} s^{-1}, a humidity of 60%, temperatures of 28^{o}C and 23^{o}C

during the day and night respectively, and a photoperiod of 12 h light, 12 h dark. Subsequent

generations were grown in a growth cabinet in 12 h light/12 h dark, at 28 ^{o}C, a relative humidity of

65%, and a photon flux density of 400 µmol m^{-2} s^{-1}.

Cloning and construct preparation, and motif analysis

The 2613-bp promoter DNA fragment of *SULFITE REDUCTASE* (*SiR,* MSU7 ID:

LOC_Os05g42350, RAP-DB ID: Os05g0503300) was originally amplified from Kitaake genomic

DNA, with forward primer (5'-3') "CACCATGCTTGACCATGTGGACTC" and reverse primer (5'-3')

"ACGGAACCCGTGGAACTC". Gel-purified PCR product was cloned into a Gateway pENTR^{TM}

vector to generate *pENTR-SiRpro* using pENTR™/D-TOPO™ Cloning Kit (Invitrogen), then the

promoter was recombined into the pGWB3 expression vector and fused with *GUS* gene using LR

reaction. The resultant vector was transformed into *A. tumefaciens* strain LBA4404 and

transformed into Kitaake. To engineer the *SiRpro* such that it is compatible with the Golden gate

system, four *Bsa*I or *Bpi*I restriction enzyme recognition sites at -214, -298, -1468, and -2309 were

mutated from T to A and cloned into the *pAGM9121* vector using Golden Gate level 0 cloning

reactions and then into a level 0 PU module EC14328, which was used for driving *kzGUS*

(intronless GUS) and *H2B-mTurquoise2* reporter genes via Golden gate reaction and using *Tnos*

as a terminator. A five prime deletion series was generated using EC14328 as the template and

prepared as level 0 PU modules, and a three prime deletion series prepared as level 0 P modules.

The minimal *CaMV35S* promoter was used as the U module, and they were linked with *kzGUS* and

terminated with *Tnos.*

To test *SiRpro* in the dTALE/STAP system, the 42-bp coding region were excluded and the

2571-bp resultant fragment placed into a level 0 PU module EC14330 and was used to drive

dTALE1. Two reporters were used. For the GUS reporter *kzGUS* was linked with *STAP62* and

terminated with *Tnos.* In the fluorescent reporter construct, a chloroplast targeting peptide fused to

mTurquoise 2 was linked with *STAP 4* and terminated with *Tact2.* In both constructs, *pOsAct1*

driving *HYG* (Hygromycin resistant gene) was terminated with *Tnos* and used as the selection

marker during rice transformation.

The Find Individual Motif Occurrences (FIMO) tool (Grant et al. 2011) from the Multiple Em for

Motif Elucidation (MEME) suite v.5.4.0 (Bailey et al. 2015) was used to search for individual motifs

within the promoter sequences using default parameters with "--thresh" of "1e-3". Position weight

matrix of 656 non-redundant plant motifs and 13 RNA polymerase II (POLII) core promoter motifs

were obtained from JASPAR (https://jaspar.elixir.no/downloads/) (Fornes et al. 2020). To cluster the

transcription factor binding motifs, the RSAT matrix-clustering tool (Castro-Mondragon et al. 2017)

was run on all 656 non-redundant plant motifs using the default parameters, which yield 51 motif

clusters, these clusters were further divided based on transcription factor families (Supplemental

table 2).

Analysis of GUS and fluorescent reporters

In all cases, to account for position effects associated with transformation via *A. tumefaciens*,

multiple T_0 lines were assessed for each construct. GUS staining was performed as described

previously (Jefferson et al., 1987) with the following minor modifications. Leaf tissue was fixed in

90% (v/v) acetone overnight at 4 ^{o}C after washing with 100 mM phosphate buffer (pH 7.0). Leaf

samples were transferred into 1 mg/ml 5-bromo-4-chloro-3-indolyl glucuronide (X-Gluc) GUS

staining solution, subjected to 2 mins vacuum infiltration 5 times, and then incubated at 37 ^{o}C for

between 1 and 168 hours. Chlorophyll was cleared further using 90% (v/v) ethanol overnight at

room temperature. Cross sections were prepared manually using a razor blade and images were

taken using an Olympus BX41 light microscopy. Quantification of GUS activity was performed

using a fluorometric MUG assay (Jefferson et al., 1987). ~200 mg mature leaves from transgenic

plants were frozen in liquid nitrogen and ground into fine powder with a Tissuelyser (Qiagen).

Soluble protein was extracted in 1 mL of 50 mM phosphate buffer (pH 7.0) supplemented with 0.1%

[v/v] Triton X-100 and cOmplete™ Protease Inhibitor Cocktail (half tablet per 50 mL). Protein

concentration then determined using a Qubit protein assay kit (Invitrogen). The MUG fluorescent

assay was performed in duplicates with 20 µl protein extract in MUG assay buffer (50mM

phosphate buffer (pH 7.0), 10 mM EDTA-Na_2, 0.1% [v/v] Triton X-100, 0.1% [w/v] N-

lauroylsarcosine sodium, 10 mM DTT, 2 mM 4-methylumbelliferyl-β-D-glucuronide (MUG)) in a 200

µl total volume. The reaction was conducted at 37 $^\circ$C in GREINER 96 F-BOTTOM microtiter plate

using a CLARIOstar plate reader. 4-Methylumbelliferone (4-MU) fluorescence was recorded every

2 minutes for 20 cycles with excitation at 360 nm and emission detected at 450 nm. 4-MU

concentration was determined based on a standard curve of ten 4-MU standards placed in the

same plate. GUS enzymatic rates were calculated by averaging the slope of MU production from

each of the duplicate reactions.

In order to visualize mTurquoise2 mature leaves were dissected into 2-cm sections, leaf

epidermal cells were removed by scraping the leaf surface with a razor blade and then mounted

with deionized water. 5-mm and middle sections of 2-cm young tissue of the fourth leaves were dissected and mounted with deionized water directly. Imaging was then performed using a Leica TCS SP8 confocal laser-scanning microscope using a 20x air objective. mTurquoise2 fluorescence was excited at 442 nm with emission at 471–481 nm, chlorophyll autofluorescence was excited at 488 nm with emission at 672–692 nm.

Yeast one hybrid, protoplast isolation and transactivation assay

The yeast one hybridisation assay was performed by Hybrigenics (https://www.hybrigenics-services.com/). Fragments were synthesized and used as bait. Rice leaf and root cDNA libraries were used as prey. The number of clones screened and concentration of 3-AT were as follows: fragment 1, 70.2 million clones screened with 0 mM 3-AT; fragment 2, 61.5 million clones screened with 0 mM 3-AT; fragment 3, 68.4 million clones screened with 20 mM 3-AT; fragment 4, 57.4 million clones screened with 100 mM 3-AT; fragment 5, 94.2 million clones screened with 200 mM 3-AT.

Rice leaf protoplasts and PEG-mediated transformation were performed as described previously (Page et al. 2019). Golden gate level 1 modules for transformation were isolated using ZymoPURE™ II Plasmid Midiprep Kit, *ZmUBIpro::GUS-Tnos* was used as transformation control. Transcription factor coding sequences were amplified using rice leaf cDNA, with *Bsa*I and *Bpi*I sites mutated, and cloned into Golden gate SC level 0 modules. They were assembled into a level 1 module with a *ZmUBIpro* promoter and *Tnos* terminator module. Nucleotides -980 to -829 with the endogenous core promoter (nucleotide -250 to +42) were fused with the *LUC* reporter to generate output of transcription activity. In each transformation, 2 µg of transformation control plasmids, 5 µg of reporter plasmids, and 5 µg of effector plasmids per transcription factor were combined and mixed with 170 µl protoplasts. After incubation on the benchtop for overnight protein was extracted using passive lysis buffer, GUS activity was determined with 20 µl of protein sample and MUG fluorescent assay as described above, LUC activity was measured with 20 µl of protein sample and 100 µl of LUC assay reagent (Promega) using Clariostar plate reader. Transcriptional activity from the promoter was calculated as LUC luminescence/rate of MUG accumulation.

Acknowledgements

This work was funded by the Bill and Melinda Gates Foundation C$_4$ Rice grant awarded to the University of Oxford (2015-2019 - OPP1129902, and 2019-2024 INV-002970). For the purpose of open access, the authors have applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

Author contributions

L.H. and J.M.H. conceived the work. J.M.H. guided execution of experiments and oversaw the

project. L.H., N.W., S.S., R.D., K.B., and A.R.B. did the experiments and analysed the data. L.H.

and J.M.H. wrote the manuscript with input from all authors.

Declaration of interests

The authors declare no competing interests.

Figure legends

Figure 1. Nucleotides -980 to -394 of the *SiR* promoter are necessary for bundle sheath

expression. (A) Domesticated *SiR* promoter generates strong GUS staining in bundle sheath. (B)

mTurquoise2 signal driven by the domesticated *SiR* promoter in nuclei (indicated by yellow arrows)

of bundle sheath cells (marked by yellow dashed lines) and vein cells in mature leaves, red

indicates chlorophyll autofluorescence. **(C)** The fluorometric 4-methylumbelliferyl-β-D-glucuronide

(MUG) assay shows no statistically significant difference between the endogenous and

domesticated *SiR* promoter activity. **(D)** Landscape of transcription factor binding sites in the *SiR*

promoter using the Find Individual Motif Occurrences (FIMO) program. The likelihood of match

between 656 plant nonredundant known transcription factor motifs in the *SiR* promoter is shown by

transcription factor families (Supplemental table 1). **(E)** Schematics showing 5' truncations. **(F)**

Representative images of leaf cross sections from transgenic lines after GUS staining. Zoomed-in

images of lateral veins shown in right panels, the staining duration is displayed in the bottom-left

corner, bundle sheath cells highlighted with dashed red line, scale bars = 50 μm. **(G)** Promoter

activity determined by the fluorometric 4-methylumbelliferyl-β-D-glucuronide (MUG) assay. Data

were subjected to a pairwise Wilcoxon test with Benjamini-Hochberg correction. Lines with

differences in activity that were statistically significant (adjusted *P*<0.05) labelled with different

letters. Median catalytic rate of GUS indicated with red line, n indicates total number of transgenic

lines assessed.

Figure 2. A distal enhancer and the core promoter that are necessary and sufficient for

bundle sheath expression (**A**) Schematics showing deletions of nucleotides -980 to -849 and -

119 to +42. (**B**) Representative image of leaf cross sections of transgenic lines after GUS staining.

Zoomed-in images of lateral veins shown in right panels, the staining duration is displayed in the

bottom-left corner, bundle sheath cells highlighted with dashed red line, scale bars = 50 μm. (**C**)

Promoter activity determined by the fluorometric 4-methylumbelliferyl-β-D-glucuronide (MUG)

assay, data subjected to pairwise Wilcoxon test. Lines with differences in activity that were

statistically significant (adjusted *P*<0.05) labelled with different letters. Median catalytic rate of GUS

indicated with red line, n indicates total number of transgenic lines assessed.

Figure 3. The Y-patch and four distinct regions in the distal enhancer are required for
bundle sheath specific expression. (A-C) Nucleotides -980 and -829 from the *SiR* promoter
pattern expression to the bundle sheath when linked with the *PIP1;1* and *NRT1.1A* core promoters
containing Y-patches. **(A)** Prediction of Y-patch and TATA-box sequences in core promoters of
PIP1;1, *NRT1.1A*, *PIP1;3* and *ATPSb*. **(B)** Representative cross sections of transgenic rice leaves
after GUS staining, zoomed-in image of lateral veins shown in the right panel, bundle sheath cells
highlighted with white dashed lines, the staining duration is displayed in the bottom-left corner,
scale bars = 50 µm. **(C)** Promoter activity determined by the fluorometric 4-methylumbelliferyl-β-D-
glucuronide (MUG) assay. **(D)** Schematics showing transcription factor binding sites between
nucleotides -980 and -829. **(E)** Schematics showing consecutive deletions between nucleotides -
980 and -829 fused to the GUS reporter. **(F)** Representative images of cross sections from
transgenic lines after GUS staining, zoomed-in images of lateral veins shown in right panels, the
staining duration is displayed in the bottom-left corner, bundle sheath cells highlighted with red
dashed lines, scale bars = 50 µm. **(G)** Promoter activity determined by the fluorometric 4-
methylumbelliferyl-β-D-glucuronide (MUG) assay. In **C&G**, data were subjected to pairwise
Wilcoxon test with Benjamini-Hochberg correction. Lines with differences in activity that were
statistically significant (adjusted *P*<0.05) labelled with different letters. Median catalytic rate of GUS
indicated with red line, n indicates total number of transgenic lines assessed.

Figure 4. WRKY, G2-like, MYB-related, IDD and bZIP transcription factors interact and
activate with the distal enhancer. (A) Schematics showing transcription factor binding sites
between nucleotides -980 and -829 which are likely required for bundle sheath specific expression.
(B) Effector assays showing that each transcription factor activates expression from the distal
enhancer. **(C)** Effector assays showing synergistic activation from the distal enhancer when GLK2
and IDD3,4,6,10 were co-expressed. Data subjected to pairwise Wilcoxon test with Benjamini-
Hochberg correction in B&C. Lines with differences in activity that were statistically significant
(adjusted *P*<0.05) labelled with different letters. **(D)** Transcript abundance of transcription factors in
bundle sheath strands (BSS) and mesophyll (M) cells during maturation. Leaf developmental stage
S2 to S7 represent base of the 4[th] leaf at the 6th, 8th, 9th, 10th, 13th and 17th day after sowing. (**E**)
Representative images of transgenic lines misexpressing WRKY121, IDD2 and bZIP9 in mesophyll
cells, staining duration is displayed in the bottom-left corner, zoom-in of mesophyll shown in right
panel, red arrows indicate GUS expressing mesophyll cells.

Figure 5. Oligomerisation of bundle sheath enhancer increases bundle sheath expression.
Schematics showing site-directed mutagenesis of WRKY, G2-like, MYBR, IDD and bZIP motifs,
mutated nucleotides highlighted in red (A), and constructs to test impact of oligomerization of
enhancer (E). **(B&F)** Representative images of cross sections from transgenic lines after GUS
staining, zoomed-in images of lateral veins shown in right panel, the staining duration is displayed

in the bottom-left corner, bundle sheath cells highlighted with red dashed lines, scale bars = 50 µm.

(C&G) Promoter activity determined by the fluorometric 4-methylumbelliferyl-β-D-glucuronide

(MUG) assay. Data subjected to pairwise Wilcoxon test with Benjamini-Hochberg correction. Lines

with differences in activity that were statistically significant (adjusted P<0.05) labelled with different

letters. Median catalytic rate of GUS indicated with red line, n indicates total number of transgenic

lines assessed, n indicates total number of transgenic lines assessed. (H) Paradermal view of

Arabidopsis leaf expressing GUS under the control of 3x BS enhancer combined with OsSiR core

promoter, the staining duration is displayed in the bottom-left corner. M indicate mesophyll, BS for

bundle sheath, V for vein. Zoomed in images shown on right.

Figure 6. Model of mechanism underpinning bundle sheath expression from *SiR* promoter.

(A) Schematic with location of Bundle Sheath (BS) enhancer, constitutive activator and mesophyll

repressor. (B) Bundle sheath expression is a result of the enhancer, constitutive activators and

mesophyll repressor acting in concert. Schematic indicating how the enhancer operates within a

broader *cis*-regulatory landscape. (C&D) Model depicting transcription factors and cognate *cis*-

elements responsible for bundle sheath expression.

Supplementary Figure legend titles

Supplemental Figure 1. Analysis of seven rice promoters identified after analysis of transcripts

that accumulate preferentially in bundle sheath strands.

Supplemental Figure 2. Identification of bundle sheath specific promoters after analysis of

transcripts that accumulate preferentially in Kitaake bundle sheath compared with mesophyll cells.

Supplemental Figure 3. The domesticated *SiR* promoter combined with the dTALE/STAP system

drives strong expression in the rice bundle sheath.

Supplemental Figure 4. The rice *SiR* promoter drives expression in bundle sheath cells earlier

than the *Zoysia japonica PCK* promoter.

Supplemental Figure 5. Impact of 5' and 3' deletions between on patterning of GUS from the *SiR*

promoter.

Supplemental Figure 6. The evolutionarily conserved *ETHYLENE INSENSITIVE3-LIKE* (*EIL*)

binding site regulates the expression level but not cell specificity of the *SiR* promoter.

Supplemental Figure 7. Identification of transcription factors interacting with the *SiR* promoter.

Supplemental Figure 8. Nucleotides -980 to -829 in combination with nucleotide -251 to +42

produce bundle sheath specific expression.

Supplemental Figure 9. Nucleotides -251 to -1 likely serve as a core promoter.

Supplemental Figure 10. Subregions in the distal enhancer not required for bundle sheath

specific expression between nucleotides -980 to -829.

Supplemental Figure 11. Regions between nucleotides -980 and -829 that in combination with

nucleotides -828 to -252 repress mesophyll expression.

Supplemental Figure 12. Nucleotide sequence in the distal enhancer.

Supplemental Figure 13. Transcription factors used in transactivation assay.

Supplemental Figure 14. Effector assay showing the effect of WRKY (A), G2-like (B), IDD (C),

MYB-related (D), SNAC and bZIP (E) transcription factors on the distal enhancer.

Supplemental Figure 15. Impact of mis-expression of bZIP9, IDD2 and WRKY121 in mesophyll

cells on GUS expression pattern driven by bundle sheath enhancer.

Supplemental Figure 16. The bundle sheath enhancer produces bundle sheath specific

expression in Arabidopsis leaves.

Supplemental Figure 17. WRKY, G2-like, MYB-related, IDD and bZIP transcription factor binding

sites identified by FIMO program in *Fd*, *HAC1;1*, *PIP1.1*, *NRT1.1A*, *ZjPCK* and *FtGLDP* promoters.

References

Adrian J, Farrona S, Reimer JJ, Albani MC, Coupland G, and Turck F. Cis-regulatory elements and chromatin
state coordinately control temporal and spatial expression of FLOWERING LOCUS T in arabidopsis.
Plant Cell. 2010:**22**(5):1425–1440. https://doi.org/10.1105/tpc.110.074682

Aubry S, Kelly S, Kümpers BMCC, Smith-Unna RD, and Hibberd JM. Deep Evolutionary Comparison of Gene
Expression Identifies Parallel Recruitment of Trans-Factors in Two Independent Origins of C_4
Photosynthesis. PLoS Genet. 2014a:**10**(6):e1004365. https://doi.org/10.1371/journal.pgen.1004365

Aubry S, Smith-Unna RD, Boursnell CM, Kopriva S, and Hibberd JM. Transcript residency on ribosomes
reveals a key role for the *Arabidopsis thaliana* bundle sheath in sulfur and glucosinolate metabolism.
Plant Journal. 2014b:**78**(4):659–673. https://doi.org/10.1111/tpj.12502

Bailey TL, Johnson J, Grant CE, and Noble WS. The MEME Suite. Nucleic Acids Res. 2015:**43**:39–49.
https://doi.org/10.1093/nar/gkv416

Bernard V, Brunaud V, and Lecharny A. TC-motifs at the TATA-box expected position in plant genes: A
novel class of motifs involved in the transcription regulation. BMC Genomics. 2010:**11**(1):1–15.
https://doi.org/10.1186/1471-2164-11-166/TABLES/4

Binder BM. Ethylene signaling in plants. J Biol Chem. 2020:**295**(22):7710.
https://doi.org/10.1074/JBC.REV120.010854

Buecker C and Wysocka J. Enhancers as information integration hubs in development: lessons from
genomics. Trends Genet. 2012:**28**(6):276. https://doi.org/10.1016/J.TIG.2012.02.008

Burgess SJ, Reyna-Llorens I, Stevenson SR, Singh P, Jaeger K, and Hibberd JM. Genome-Wide Transcription
Factor Binding in Leaves from C3 and C4 Grasses. Plant Cell. 2019:**31**(10):2297–2314.
https://doi.org/10.1105/TPC.19.00078

Castro-Mondragon JA, Jaeger S, Thieffry D, Thomas-Chollier M, and Van Helden J. RSAT matrix-clustering:
dynamic exploration and redundancy reduction of transcription factor binding motif collections.
Nucleic Acids Res. 2017:**45**(13):e119–e119. https://doi.org/10.1093/NAR/GKX314

Chen X, Wang Y, Lv B, Li J, Luo L, Lu S, Zhang X, Ma H, and Ming F. The NAC family transcription factor
OsNAP confers abiotic stress response through the ABA pathway. Plant Cell Physiol. 2014:**55**(3):604–
619. https://doi.org/10.1093/PCP/PCT204

Chow C-N, Lee T-Y, Hung Y-C, Li G-Z, Tseng K-C, Liu Y-H, Kuo P-L, Zheng H-Q, and Chang W-C. PlantPAN3.0:
a new and updated resource for reconstructing transcriptional regulatory networks from ChIP-seq
experiments in plants. Nucleic Acids Res. 2018. https://doi.org/10.1093/nar/gky1081

Civáň P and Švec M. Genome-wide analysis of rice (Oryza sativa L. subsp. japonica) TATA box and Y Patch
promoter elements. Genome. 2009:**52**(3):294–297. https://doi.org/10.1139/G09-
001/SUPPL_FILE/G09-001.DOC

Clark RM, Wagler TN, Quijada P, and Doebley J. A distant upstream enhancer at the maize domestication
gene tb1 has pleiotropic effects on plant and inflorescent architecture. Nat Genet. 2006:**38**(5):594–
597. https://doi.org/10.1038/ng1784

Cui H, Kong D, Liu X, and Hao Y. SCARECROW, SCR-LIKE 23 and SHORT-ROOT control bundle sheath cell fate
and function in Arabidopsis thaliana. The Plant Journal. 2014:**78**(2):319–327.
https://doi.org/10.1111/TPJ.12470

Danila F, Schreiber T, Ermakova M, Hua L, Vlad D, Lo SF, Chen YS, Lambret-Frotte J, Hermanns AS, Athmer
B, et al. A single promoter-TALE system for tissue-specific and tuneable expression of multiple genes
in rice. Plant Biotechnol J. 2022. https://doi.org/10.1111/PBI.13864

Depuydt S and Hardtke CS. Hormone signalling crosstalk in plant growth regulation. Curr Biol. 2011:21(9).
https://doi.org/10.1016/J.CUB.2011.03.013

Dickinson PJ, Kneřová J, Szecówka M, Stevenson SSR, Burgess SJ, Mulvey H, Bågman A-MM, Gaudinier A,
Brady SM, and Hibberd JM. A bipartite transcription factor module controlling expression in the
bundle sheath of Arabidopsis thaliana. Nat Plants. 2020:6(12):1468–1479.
https://doi.org/10.1038/s41477-020-00805-w

Dietzen C, Koprivova A, Whitcomb SJ, Langen G, Jobe TO, Hoefgen R, and Kopriva S. The Transcription
Factor EIL1 Participates in the Regulation of Sulfur-Deficiency Response. Plant Physiol.
2020:184(4):2120–2136. https://doi.org/10.1104/pp.20.01192

Drapek C, Sparks EE, and Benfey PN. Uncovering Gene Regulatory Networks Controlling Plant Cell
Differentiation. Trends in Genetics. 2017:33(8):529–539. https://doi.org/10.1016/J.TIG.2017.05.002

Emmerling J. Studies into the Regulation of C4 Photosynthesis – Towards Factors Controlling Bundle Sheath
Expression and Kranz Anatomy Development. Doctoral thesis, Heinrich Heine University, Düsseldorf.
2018:(September).

Engelmann S, Wiludda C, Burscheidt J, Gowik U, Schlue U, Koczor M, Streubel M, Cossu R, Bauwe H, and
Westhoff P. The Gene for the P-Subunit of Glycine Decarboxylase from the C4 Species Flaveria
trinervia: Analysis of Transcriptional Control in Transgenic Flaveria bidentis (C4) and Arabidopsis (C3).
Plant Physiol. 2008:146(4):1773–1785. https://doi.org/10.1104/PP.107.114462

Ermakova M, Arrivault S, Giuliani R, Danila F, Alonso-Cantabrana H, Vlad D, Ishihara H, Feil R, Guenther
M, Borghi GL, et al. Installation of C$_4$ photosynthetic pathway enzymes in rice using a single construct.
Plant Biotechnol J. 2021:19(3):575–588. https://doi.org/10.1111/pbi.13487

Fornes O, Castro-Mondragon JA, Khan A, Van Der Lee R, Zhang X, Richmond PA, Modi BP, Correard S,
Gheorghe M, Baranašić D, et al. JASPAR 2020: Update of the open-Access database of transcription
factor binding profiles. Nucleic Acids Res. 2020:48(D1):D87–D92.
https://doi.org/10.1093/nar/gkz1001

Gallegos JE and Rose AB. An intron-derived motif strongly increases gene expression from transcribed
sequences through a splicing independent mechanism in Arabidopsis thaliana. Scientific Reports 2019
9:1. 2019:9(1):1–9. https://doi.org/10.1038/s41598-019-50389-5

Glick BR. Metabolic load and heterologous gene expression. Biotechnol Adv. 1995:13(2):247–261.
https://doi.org/10.1016/0734-9750(95)00004-A

Grant CE, Bailey TL, and Noble WS. FIMO: Scanning for occurrences of a given motif. Bioinformatics.
2011:27(7):1017–1018. https://doi.org/10.1093/bioinformatics/btr064

Haberle V and Stark A. Eukaryotic core promoters and the functional basis of transcription initiation. Nat
Rev Mol Cell Biol. 2018:19(10):621. https://doi.org/10.1038/S41580-018-0028-8

Hibberd JM and Covshoff S. The Regulation of Gene Expression Required for C$_4$ Photosynthesis. Annu Rev
Plant Biol. 2010:61(1):181–207. https://doi.org/10.1146/annurev-arplant-042809-112238

Hibberd JM, Sheehy JE, and Langdale JA. Using C4 photosynthesis to increase the yield of rice-rationale
and feasibility. Curr Opin Plant Biol. 2008:11(2):228–231. https://doi.org/10.1016/j.pbi.2007.11.002

Hua L, Stevenson SR, Reyna-Llorens I, Xiong H, Kopriva S, and Hibberd JM. The bundle sheath of rice is
conditioned to play an active role in water transport as well as sulfur assimilation and jasmonic acid
synthesis. The Plant Journal. 2021:**107**(1):268–286. https://doi.org/10.1111/TPJ.15292

Jisha V, Dampanaboina L, Vadassery J, Mithöfer A, Kappara S, and Ramanan R. Overexpression of an
AP2/ERF Type Transcription Factor OsEREBP1 Confers Biotic and Abiotic Stress Tolerance in Rice. PLoS
One. 2015:**10**(6). https://doi.org/10.1371/JOURNAL.PONE.0127831

Jores T, Tonnies J, Wrightsman T, Buckler ES, Cuperus JT, Fields S, and Queitsch C. Synthetic promoter
designs enabled by a comprehensive analysis of plant core promoters. Nature Plants 2021 7:6.
2021:**7**(6):842–855. https://doi.org/10.1038/s41477-021-00932-y

Jung SE, Bang SW, Kim SH, Seo JS, Yoon H Bin, Kim YS, and Kim JK. Overexpression of OsERF83, a Vascular
Tissue-Specific Transcription Factor Gene, Confers Drought Tolerance in Rice. Int J Mol Sci.
2021:**22**(14). https://doi.org/10.3390/IJMS22147656

Junion G, Spivakov M, Girardot C, Braun M, Gustafson EH, Birney E, and Furlong EEM. A transcription
factor collective defines cardiac cell fate and reflects lineage history. Cell. 2012:**148**(3):473–486.
https://doi.org/10.1016/J.CELL.2012.01.030

Juven-Gershon T and Kadonaga JT. Regulation of gene expression via the core promoter and the basal
transcriptional machinery. 2009. https://doi.org/10.1016/j.ydbio.2009.08.009

Kajala K, Covshoff S, Karki S, Woodfield H, Tolley BJ, Dionora MJA, Mogul RT, Mabilangan AE, Danila FR,
Hibberd JM, et al. Strategies for engineering a two-celled C4 photosynthetic pathway into rice. J Exp
Bot. 2011:**62**(9):3001–3010. https://doi.org/10.1093/JXB/ERR022

Kanaoka MM, Pillitteri LJ, Fujii H, Yoshida Y, Bogenschutz NL, Takabayashi J, Zhu JK, and Torii KU.
SCREAM/ICE1 and SCREAM2 Specify Three Cell-State Transitional Steps Leading to Arabidopsis
Stomatal Differentiation. Plant Cell. 2008:**20**(7):1775–1785. https://doi.org/10.1105/TPC.108.060848

Kirschner S, Woodfield H, Prusko K, Koczor M, Gowik U, Hibberd JM, and Westhoff P. Expression of
SULTR2;2, encoding a low-affinity sulphur transporter, in the Arabidopsis bundle sheath and vein cells
is mediated by a positive regulator. J Exp Bot. 2018:**69**(20):4897–4906.
https://doi.org/10.1093/jxb/ery263

Krumlauf R. Hox genes in vertebrate development. Cell. 1994:**78**(2):191–201.
https://doi.org/10.1016/0092-8674(94)90290-9

Kubo M, Udagawa M, Nishikubo N, Horiguchi G, Yamaguchi M, Ito J, Mimura T, Fukuda H, and Demura T.
Transcription switches for protoxylem and metaxylem vessel formation. Genes Dev.
2005:**19**(16):1855–1860. https://doi.org/10.1101/GAD.1331305

Kulkarni MM and Arnosti DN. Information display by transcriptional enhancers. Development.
2003:**130**(26):6569–6575. https://doi.org/10.1242/DEV.00890

Kummari D, Palakolanu SR, Kishor PBK, Bhatnagar-Mathur P, Singam P, Vadez V, and Sharma KK. An
update and perspectives on the use of promoters in plant genetic engineering. J Biosci. 2020:**45**(1):1–
24. https://doi.org/10.1007/S12038-020-00087-6/TABLES/7

Kurt F, Filiz E, and Aydın A. Sulfite Reductase (SiR) Gene in Rice (Oryza sativa): Bioinformatics and
Expression Analyses Under Salt and Drought Stresses. J Plant Growth Regul. 2022:**41**(6):2246–2260.
https://doi.org/10.1007/S00344-021-10438-8/FIGURES/9

Lee DY, Hua L, Khoshravesh R, Giuliani R, Kumar I, Cousins A, Sage TL, Hibberd JM, and Brutnell TP.
Engineering chloroplast development in rice through cell-specific control of endogenous genetic
circuits. Plant Biotechnol J. 2021:**19**(11):2291–2303. https://doi.org/10.1111/PBI.13660

Leegood RC. Roles of the bundle sheath cells in leaves of C_3 plants. J Exp Bot. 2008:**59**(7):1663–1673.
https://doi.org/10.1093/jxb/erm335

Lewis EB. A gene complex controlling segmentation in Drosophila. Nature 1978 276:5688.
1978:**276**(5688):565–570. https://doi.org/10.1038/276565a0

Li E, Liu H, Huang L, Zhang X, Dong X, Song W, Zhao H, and Lai J. Long-range interactions between proximal
and distal regulatory regions in maize. Nat Commun. 2019:**10**(1). https://doi.org/10.1038/S41467-
019-10603-4

Li L, Zhu Q, He X, Sinha S, and Halfon MS. Large-scale analysis of transcriptional cis-regulatory modules
reveals both common features and distinct subclasses. Genome Biol. 2007:**8**(6).
https://doi.org/10.1186/GB-2007-8-6-R101

Lin R, Zhao W, Meng X, and Peng YL. Molecular cloning and characterization of a rice gene encoding
AP2/EREBP-type transcription factor and its expression in response to infection with blast fungus and
abiotic stresses. Physiol Mol Plant Pathol. 2007:**70**(1–3):60–68.
https://doi.org/10.1016/J.PMPP.2007.06.002

Liu L, Adrian J, Pankin A, Hu J, Dong X, Von Korff M, and Turck F. Induced and natural variation of
promoter length modulates the photoperiodic response of FLOWERING LOCUS T. Nat Commun.
2014:**5**. https://doi.org/10.1038/ncomms5558

Liu L, Du Y, Shen X, Li M, Sun W, Huang J, Liu Z, Tao Y, Zheng Y, Yan J, et al. KRN4 Controls Quantitative
Variation in Maize Kernel Row Number. PLoS Genet. 2015:**11**(11):e1005670.
https://doi.org/10.1371/JOURNAL.PGEN.1005670

MacAlister CA, Ohashi-Ito K, and Bergmann DC. Transcription factor control of asymmetric cell divisions
that establish the stomatal lineage. Nature 2006 445:7127. 2006:**445**(7127):537–540.
https://doi.org/10.1038/nature05491

Makino A, Sakuma H, Sudo E, and Mae T. Differences between Maize and Rice in N-use Efficiency for
Photosynthesis and Protein Allocation.

Maruyama-Nakashita A, Nakamura Y, Tohge T, Saito K, and Takahashi H. *Arabidopsis* SLIM1 is a central
transcriptional regulator of plant sulfur response and metabolism. Plant Cell. 2006:**18**(11):3235–3251.
https://doi.org/10.1105/tpc.106.046458

McClung CR. Plant circadian rhythms. Plant Cell. 2006:**18**(4):792–803.
https://doi.org/10.1105/TPC.106.040980

Mitchell PL and Sheehy JE. Supercharging rice photosynthesis to increase yield. New Phytol.
2006:**171**(4):688–693. https://doi.org/10.1111/J.1469-8137.2006.01855.X

Miyashima S, Roszak P, Sevilem I, Toyokura K, Blob B, Heo Jok, Mellor N, Help-Rinta-Rahko H, Otero S,
Smet W, et al. Mobile PEAR transcription factors integrate positional cues to prime cambial growth.
Nature. 2019:**565**(7740):490–494. https://doi.org/10.1038/S41586-018-0839-Y

Moreno-Risueno MA, Sozzani R, Yardimci GG, Petricka JJ, Vernoux T, Blilou I, Alonso J, Winter CM, Ohler
U, Scheres B, et al. Transcriptional control of tissue formation throughout root development. Science
(1979). 2015:**350**(6259):426–430.
https://doi.org/10.1126/SCIENCE.AAD1171/SUPPL_FILE/TABLE_S6.XLSX

Nagel DH and Kay SA. Complexity in the wiring and regulation of plant circadian networks. Curr Biol.
2012:**22**(16). https://doi.org/10.1016/J.CUB.2012.07.025

Nakashima K, Ito Y, and Yamaguchi-Shinozaki K. Transcriptional Regulatory Networks in Response to
Abiotic Stresses in Arabidopsis and Grasses. Plant Physiol. 2009:**149**(1):88.
https://doi.org/10.1104/PP.108.129791

**Nomura M, Higuchi T, Ishida Y, Ohta S, Komari T, Imaizumi N, Miyao-Tokutomi M, Matsuoka M, and
Tajima S**. Differential expression pattern of C4 bundle sheath expression genes in rice, a C3 plant.
Plant Cell Physiol. 2005a:**46**(5):754–761. https://doi.org/10.1093/PCP/PCI078

**Nomura M, Higuchi T, Ishida Y, Ohta S, Komari T, Imaizumi N, Miyao-Tokutomi M, Matsuoka M, and
Tajima S**. Differential expression pattern of C4 bundle sheath expression genes in rice, a C3 plant.
Plant Cell Physiol. 2005b:**46**(5):754–761. https://doi.org/10.1093/PCP/PCI078

Ohashi-Ito K and Bergmann DC. Arabidopsis FAMA Controls the Final Proliferation/Differentiation Switch
during Stomatal Development. Plant Cell. 2006:**18**(10):2493–2505.
https://doi.org/10.1105/TPC.106.046136

Page MT, Parry MAJ, and Carmo-Silva E. A high-throughput transient expression system for rice. Plant Cell
Environ. 2019:**42**(7):2057–2064. https://doi.org/10.1111/PCE.13542

Panne D. The enhanceosome. Curr Opin Struct Biol. 2008:**18**(2):236–242.
https://doi.org/10.1016/J.SBI.2007.12.002

Pillitteri LJ, Sloan DB, Bogenschutz NL, and Torii KU. Termination of asymmetric cell division and
differentiation of stomata. Nature 2006 445:7127. 2006:**445**(7127):501–505.
https://doi.org/10.1038/nature05467

Qu X, Zou J, Wang J, Yang K, Wang X, and Le J. A Rice R2R3-Type MYB Transcription Factor OsFLP Positively
Regulates Drought Stress Response via OsNAC. Int J Mol Sci. 2022:**23**(11).
https://doi.org/10.3390/IJMS23115873

Que Q, Wang HY, English JJ, and Jorgensen RA. The Frequency and Degree of Cosuppression by Sense
Chalcone Synthase Transgenes Are Dependent on Transgene Promoter Strength and Are Reduced by
Premature Nonsense Codons in the Transgene Coding Sequence. Plant Cell. 1997:**9**(8):1357.
https://doi.org/10.1105/TPC.9.8.1357

Ray-Jones H and Spivakov M. Transcriptional enhancers and their communication with gene promoters.
Cellular and Molecular Life Sciences. 2021:**78**(19–20):6453. https://doi.org/10.1007/S00018-021-
03903-W

Reyna-Llorens I, Burgess SJ, Reeves G, Singh P, Stevenson SR, Williams BP, Stanley S, and Hibberd JM.
Ancient duons may underpin spatial patterning of gene expression in C4 leaves. Proc Natl Acad Sci U S
A. 2018:**115**(8):1931–1936.
https://doi.org/10.1073/PNAS.1720576115/SUPPL_FILE/PNAS.1720576115.SD07.XLSX

Rose AB. Introns as Gene Regulators: A Brick on the Accelerator. Front Genet. 2019:**9**(FEB).
https://doi.org/10.3389/FGENE.2018.00672

Rose AB, Elfersi T, Parra G, and Korf I. Promoter-proximal introns in Arabidopsis thaliana are enriched in
dispersed signals that elevate gene expression. Plant Cell. 2008:**20**(3):543–551.
https://doi.org/10.1105/TPC.107.057190

Sage RF. The evolution of C_4 photosynthesis. New Phytologist. 2004:**161**(2):341–370.
https://doi.org/10.1111/j.1469-8137.2004.00974.x

Schmitz RJ, Grotewold E, and Stam M. Cis-regulatory sequences in plants: Their importance, discovery, and
future challenges. Plant Cell. 2022:**34**(2):718–741. https://doi.org/10.1093/PLCELL/KOAB281

Shimizu H, Sato K, Berberich T, Miyazaki A, Ozaki R, Imai R, and Kusano T. LIP19, a basic region leucine
zipper protein, is a Fos-like molecular switch in the cold signaling of rice plants. Plant Cell Physiol.
2005:**46**(10):1623–1634. https://doi.org/10.1093/PCP/PCI178

Shlyueva D, Stampfel G, and Stark A. Transcriptional enhancers: from properties to genome-wide
predictions. Nat Rev Genet. 2014:**15**(4):272–286. https://doi.org/10.1038/NRG3682

Smale ST and Kadonaga JT. The RNA polymerase II core promoter. Annu Rev Biochem. 2003:**72**(Volume 72,
2003):449–479. https://doi.org/10.1146/ANNUREV.BIOCHEM.72.121801.161520/CITE/REFWORKS

**Sparks EE, Drapek C, Gaudinier A, Li S, Ansariola M, Shen N, Hennacy JH, Zhang J, Turco G, Petricka JJ, et
al.** Establishment of Expression in the SHORTROOT-SCARECROW Transcriptional Cascade through
Opposing Activities of Both Activators and Repressors. Dev Cell. 2016:**39**(5):585–596.
https://doi.org/10.1016/j.devcel.2016.09.031

Spitz F and Furlong EEM. Transcription factors: from enhancer binding to developmental control. Nat Rev
Genet. 2012:**13**(9):613–626. https://doi.org/10.1038/NRG3207

Stam M, Belele C, Dorweiler JE, and Chandler VL. Differential chromatin structure within a tandem array
100 kb upstream of the maize b1 locus is associated with paramutation. Genes Dev.
2002:**16**(15):1906–1918. https://doi.org/10.1101/GAD.1006702

**Stergachis AB, Haugen E, Shafer A, Fu W, Vernot B, Reynolds A, Raubitschek A, Ziegler S, LeProust EM,
Akey JM, et al.** Exonic transcription factor binding directs codon choice and impacts protein evolution.
Science. 2013:**342**(6164):1367. https://doi.org/10.1126/SCIENCE.1243490

Studer A, Zhao Q, Ross-Ibarra J, and Doebley J. Identification of a functional transposon insertion in the
maize domestication gene tb1. Nat Genet. 2011:**43**(11):1160. https://doi.org/10.1038/NG.942

Swift J, Luginbuehl LH, Schreier TB, Donald RM, Lee TA, Nery JR, Ecker JR, and Hibberd JM. Single nuclei
sequencing reveals C4 photosynthesis is based on rewiring of ancestral cell identity networks. bioRxiv.
2023:2023.10.26.562893. https://doi.org/10.1101/2023.10.26.562893

Tezuka D, Kawamata A, Kato H, Saburi W, Mori H, and Imai R. The rice ethylene response factor OsERF83
positively regulates disease resistance to Magnaporthe oryzae. Plant Physiol Biochem. 2019:**135**:263–
271. https://doi.org/10.1016/J.PLAPHY.2018.12.017

Thanos D and Maniatis T. Virus induction of human IFNβ gene expression requires the assembly of an
enhanceosome. Cell. 1995:**83**(7):1091–1100. https://doi.org/10.1016/0092-8674(95)90136-1

Tsuda K and Somssich IE. Transcriptional networks in plant immunity. New Phytol. 2015:**206**(3):932–947.
https://doi.org/10.1111/NPH.13286

Verma V, Ravindran P, and Kumar PP. Plant hormone-mediated regulation of stress responses. BMC Plant
Biol. 2016:**16**(1):1–10. https://doi.org/10.1186/S12870-016-0771-Y/FIGURES/1

Wang L, Wan MC, Liao RY, Xu J, Xu ZG, Xue HC, Mai YX, and Wang JW. The maturation and aging
trajectory of Marchantia polymorpha at single-cell resolution. Dev Cell. 2023:**58**(15):1429-1444.e6.
https://doi.org/10.1016/J.DEVCEL.2023.05.014

Wang Y, Huan Q, Li K, and Qian W. Single-cell transcriptome atlas of the leaf and root of rice seedlings.
Journal of Genetics and Genomics. 2021:**48**(10):881–898. https://doi.org/10.1016/J.JGG.2021.06.001

Weber B, Zicola J, Oka R, and Stam M. Plant Enhancers: A Call for Discovery. Trends Plant Sci.
2016:**21**(11):974–987. https://doi.org/10.1016/J.TPLANTS.2016.07.013

Wiludda C, Schulze S, Gowik U, Engelmann S, Koczor M, Streubel M, Bauwe H, and Westhoff P. Regulation
of the Photorespiratory GLDPA Gene in C4 Flaveria: An Intricate Interplay of Transcriptional and
Posttranscriptional Processes. Plant Cell. 2012:**24**(1):137. https://doi.org/10.1105/TPC.111.093872

Xiong H, Hua L, Reyna-Llorens I, Shi Y, Chen KM, Smirnoff N, Kromdijk J, and Hibberd JM. Photosynthesis-
independent production of reactive oxygen species in the rice bundle sheath during high light is
mediated by NADPH oxidase. Proc Natl Acad Sci U S A. 2021:**118**(25):e2022702118.
https://doi.org/10.1073/PNAS.2022702118/SUPPL_FILE/PNAS.2022702118.SD04.XLSX

Yaish MW, El-Kereamy A, Zhu T, Beatty PH, Good AG, Bi YM, and Rothstein SJ. The APETALA-2-like
transcription factor OsAP2-39 controls key interactions between abscisic acid and gibberellin in rice.
PLoS Genet. 2010:**6**(9). https://doi.org/10.1371/JOURNAL.PGEN.1001098

Yamamoto Y, Ichida H, Abe T, Suzuki Y, Sugano S, and Obokata J. Differentiation of core promoter
architecture between plants and mammals revealed by LDSS analysis. Nucleic Acids Res.
2007a:**35**(18):6219. https://doi.org/10.1093/NAR/GKM685

Yamamoto YY, Ichida H, Matsui M, Obokata J, Sakurai T, Satou M, Seki M, Shinozaki K, and Abe T.
Identification of plant promoter constituents by analysis of local distribution of short sequences. BMC
Genomics. 2007b:**8**(1):1–23. https://doi.org/10.1186/1471-2164-8-67/FIGURES/9

Yan W, Chen D, Schumacher J, Durantini D, Engelhorn J, Chen M, Carles CC, and Kaufmann K. Dynamic
control of enhancer activity drives stage-specific gene expression during flower morphogenesis. Nat
Commun. 2019:**10**(1). https://doi.org/10.1038/S41467-019-09513-2

Zhang C, Ding Z, Wu K, Yang L, Li Y, Yang Z, Shi S, Liu X, Zhao S, Yang Z, et al. Suppression of Jasmonic Acid-
Mediated Defense by Viral-Inducible MicroRNA319 Facilitates Virus Infection in Rice. Mol Plant.
2016:**9**(9):1302–1314. https://doi.org/10.1016/J.MOLP.2016.06.014

Zhang C, Zhang J, Liu H, Qu X, Wang J, He Q, Zou J, Yang K, and Le J. Transcriptomic analysis reveals the
role of FOUR LIPS in response to salt stress in rice. Plant Mol Biol. 2022:**110**(1–2):37–52.
https://doi.org/10.1007/S11103-022-01282-9/FIGURES/7

Zhang H, Zhang J, Quan R, Pan X, Wan L, and Huang R. EAR motif mutation of rice OsERF3 alters the
regulation of ethylene biosynthesis and drought tolerance. Planta. 2013:**237**(6):1443–1451.
https://doi.org/10.1007/S00425-013-1852-X

Zhang X, Bao Y, Shan D, Wang Z, Song X, Wang Z, Wang J, He L, Wu L, Zhang Z, et al. Magnaporthe oryzae
Induces the Expression of a MicroRNA to Suppress the Immune Response in Rice. Plant Physiol.
2018:**177**(1):352. https://doi.org/10.1104/PP.17.01665

Zhao H, Yang M, Bishop J, Teng Y, Cao Y, Beall BD, Li S, Liu T, Fang Q, Fang C, et al. Identification and
functional validation of super-enhancers in Arabidopsis thaliana. Proc Natl Acad Sci U S A.
2022:**119**(48):e2215328119.
https://doi.org/10.1073/PNAS.2215328119/SUPPL_FILE/PNAS.2215328119.SD02.XLSX

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6