A transcription factor quartet orchestrating bundle sheath expression in rice

Lei Hua, Na Wang, Susan Stanley, Ruth Donald, Kumari Billakurthi, Ana Rita Borba and Julian M. Hibberd

Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge CB2 3EA, United Kingdom

Email for contact: jmh65@cam.ac.uk

Keywords: *Oryza sativa*; C_4 rice; Bundle Sheath; Sulfite Reductase; Promoter; cis-elements

Abstract

The C_4 pathway has evolved in over sixty lineages of plants and improves photosynthetic efficiency

by ~50%. One unifying character of C_4 plants is photosynthetic activation of a compartment such as

the bundle sheath, but gene regulatory networks controlling this cell type are poorly understood. For

example, in *Arabidopsis thaliana* a bipartite MYC-MYB transcription factor module restricts gene

expression to this cell type but in grasses the regulatory logic allowing bundle sheath gene

expression has not been defined. Using the global staple and C_3 crop rice we identified the *SULFITE*

REDUCTASE promoter as sufficient for strong expression in this cell type. This promoter encodes

an intricate *cis*-regulatory logic with multiple activators and repressors acting in a combinatorial

manner. However, within this landscape we identified a distal enhancer comprising a module

activated by quartet of transcription factors from the G2-like, MYB-related, IDD and bZIP families.

This module is necessary and sufficient to pattern gene expression to the rice bundle sheath, and

oligomerisation allowed activity to be increased 23-fold. In summary, we identify a short tunable

enhancer patterning expression to the rice bundle sheath that we anticipate will be useful for

engineering this cell type. The transcription factor module responsible for this patterning is distinct

from that previously reported in *A. thaliana*.

Introduction

Multicellularity allows increased specialisation between cell-types, and since the colonisation of
land the complexity of photosynthetic structures found in plants appears to have increased. For
example, in the liverwort *Marchantia polymorpha* the photosynthetic thallus contains seven cell types
of which mature somatic cells and initials at the base of chambers contain the most chloroplasts
(Wang et al. 2023). Leaves of the angiosperm *Arabidopsis thaliana* possess at least nine
anatomically distinct cell types and single-cell sequencing has identified an additional eight cell
populations (Wang et al. 2021). Similarly, leaves of rice display a complex structure featuring at least
ten cell types, with five more cell clusters predicted from single-cell sequencing (Wang et al. 2021).
Photosynthesis gene expression in species such as rice is induced by light in all major cell types of
the leaf and whilst this response is greatest in spongy and palisade mesophyll cells it is also evident
in guard, mestome and bundle sheath cells (Swift et al. 2023). The bundle sheath constitutes a ring
of photosynthetic cells around the vasculature (Esau, 1965) whose patterns of gene expression
appear to condition these cells to allow water transport from veins to mesophyll cells, as well as
sulphur assimilation and nitrate reduction (Leegood 2008; Aubry et al. 2014b; Hua et al. 2021). In
some species, the bundle sheath is also specialised to accumulate metabolites such as
glucosinolates that deter attack (Rosa and Rodrigues 1999; Rask et al. 2000). And, strikingly in
multiple other lineages this specialised cell type has been repurposed during evolution to become
fully photosynthetic and allow the complex C_4 pathway to operate.

Although the bundle sheath is found in all angiosperms and associated with multiple processes
fundamental to leaf function, the molecular mechanisms responsible for directing expression to this
cell type are poorly understood (Hibberd and Covshoff 2010; Kajala et al. 2011a). For example, few
promoters capable of directing expression to this cell type have been reported (Schuler et al. 2016;
Cui 2021). Moreover, only one transcription factor network comprising a MYC-MYB module that is
necessary and sufficient to drive gene expression to the bundle sheath has been defined(Dickinson
et al. 2020a). To our knowledge, no transcription factors or *cis*-elements responsible for decoding
their activity have been shown to be capable of driving gene expression to the bundle sheath of
grasses. In fact, in cereal crops such as rice only two promoters that pattern gene expression to the
bundle sheath have been reported. The first is the *PHOSPHOENOLPYRUVATE CARBOXYKINASE*
promoter derived from *Zoysia japonica* (*ZjPCK)* (Nomura et al. 2005a; Wang et al. 2017a; Danila et
al. 2022a), and the second a *GLYCINE DECARBOXYLASE P-SUBUNIT* from *Flaveria trinervia*
(*FtGLDP)* (Lee et al. 2021). Both promoters direct expression in mature cells of developed leaves
but are weak, and the regulatory logic allowing them to generate bundle sheath expression is not
well understood.

Identifying how rice patterns gene expression to the bundle sheath would provide insight into the
underlying gene regulatory networks but also facilitate engineering of this cell type. One such
example is associated with attempts to install C_4 photosynthesis into rice (Mitchell and Sheehy 2006).
Compared with the ancestral C_3 state, plants that use C_4 photosynthesis operate higher light, water

and nitrogen use efficiencies (Makino et al. 2003; Sage 2004a; Mitchell and Sheehy 2006). It is

estimated that introducing C_4 photosynthetic metabolism into C_3 rice would allow a 50% increase in

yield (Mitchell and Sheehy 2006; Hibberd et al. 2008). C_4 photosynthesis requires multiple

photosynthesis genes to be expressed in the bundle sheath, including enzymes that decarboxylate

C_4 acids to release CO_2 around RuBisCO, organic acid transporters, components of the Calvin-

Benson-Bassham cycle, RuBisCO activase, and enzymes of starch biosynthesis (Kajala et al. 2011b;

Aubry et al. 2014a; Ermakova et al. 2021). Whilst expression of genes can be driven from

misexpression of transcription factors such as designer Transcription Activator-Like Effectors

(dTALEs) (Danila et al. 2022b) this still demands a promoter able to direct robust expression. In

addition to driving strong expression of structural photosynthesis genes in the rice bundle sheath,

introducing the C_4 pathway into C_3 leaves demands an expansion of the chloroplast and

mitochondrial compartments compared with the ancestral C_3 state (Sage 2004a; Wang et al. 2017b;

Lee et al. 2021). Because biogenesis of these organelles takes place early in leaf development

(Wang et al. 2017a; Lee et al. 2021) a promoter capable of driving expression at these early stages

is also desirable. Thus, the exogenous *ZjPCK* and *FtGLDP* promoters currently limit our ability to

engineer processes in the rice bundle sheath.

We hypothesized that analysis of endogenous patterns of gene expression in the rice bundle

sheath would allow us to identify a strong and early-acting promoter for this cell type. Once such a

promoter was identified we also hypothesised that it could be used to initiate an understanding of

the *cis*-regulatory logic that allows gene expression to be patterned to this cell type in grasses. We

tested twenty-five promoters from rice genes that transcriptome sequencing indicated were highly

expressed in these cells. Of these, four promoters specified preferential expression in the bundle

sheath, and one derived from the *SULFITE REDUCTASE* (*SiR*) gene generated strong bundle

sheath expression from plastochron 3 leaves. Truncation analysis showed that bundle sheath

expression pattern from the *SiR* promoter is mediated by a short distal enhancer. This bundle sheath

module is cryptic until other enhancers acting to both constitutively activate and repress expression

in mesophyll cells are removed. The enhancer is composed of a quartet of *cis*-elements recognised

by their cognate transcription factors from the GLK2, MYBS1, IDD and bZIP families. These four

transcription factors act synergistically and are sufficient to drive expression of the strong bundle

sheath *SiR* promoter.

Results

The *SiR* promoter directs expression to the rice bundle sheath

To identify sequences allowing robust expression in rice bundle sheath cells we used data derived

from laser capture microdissection of bundle sheath strands and mesophyll cells from mature leaves.

Promoter sequence from seven of the most strongly expressed genes in bundle sheath strands

(**Supplemental Figure 1A**) were cloned, fused to the β-glucoronidase (GUS) reporter and

transformed into rice. Although five of these promoters (*MYELOBLASTOSIS*, *MYB*; *HOMOLOG OF*

E. COLI BOLA, *bolA*; *GLUTAMINE SYNTHETASE 1*, *GS1*; *STRESS RESPONSEIVE PROTEIN*,

SRP; *ACYL COA BINDING PROTEIN*, *ARP*) led to GUS accumulation, it was restricted to veins

(**Supplemental Figure 1B, 1C**). And, for the *SULFATE TRANSPORTER 3;1* and *3;3* (*SULT3;1* and

SULT3;3) promoters, no staining was observed (**Supplemental Figure 1B, 1C**). The approach of

cloning promoters from bundle sheath strands therefore appeared to be more efficient at identifying

sequences capable of driving expression in veins. We therefore optimised a procedure allowing

bundle sheath cells to be separated from veins (Hua and Hibberd, 2019) to produce high quality

transcriptomes from mesophyll, bundle sheath and vascular bundles of rice (Hua et al. 2021). From

these data eighteen genes whose transcripts were more abundant in bundle sheath cells compared

with both veins and mesophyll cells were identified (**Supplemental Figure 2A**). When each of these

promoters was fused to GUS and transformed into rice, those from *ATP-SULFURYLASE 1B*,

ATPS1b; *SULFITE REDUCTASE*, *SiR*; *HIGH ARSENIC CONTENT 1.1*, *HAC1.1*; and

FERREDOXIN, *Fd* were sufficient to generate expression in the bundle sheath (**Supplemental**

Figure 2B). However, *ATPS1b* and *Fd* also displayed weak activity in the mesophyll, and the

HAC1.1 promoter led to GUS accumulation in epidermal and vascular cells as well as the bundle

sheath. Thus, only the *SiR* promoter drove strong expression in the bundle sheath with no GUS

detected in other cells (**Supplemental Figure 2B, 2C**). An additional six promoters (*SOLUBLE*

INORGANIC PYROPHOSPHATASE, *PPase*; *PLASMA MEMBRANE INTRINSIC PROTEIN 1;1*,

Os*PIP1;1*; *PLASMA MEMBRANE INTRINSIC PROTEIN 1;3*, Os*PIP1;3*; *ACTIN-DEPOLYMERIZING*

FACTOR, *ADF*; *PEPTIDE TRANSPORTER PTR2*, *PTR2*; *NITRATE REDUCTASE*, *NIA2*)

generated expression in vascular bundles, and eight promoters produced no staining

(**Supplemental Figure 2B, 2C**). In summary, most candidate promoters failed to generate

expression that was specific to bundle sheath cells, but the region upstream of the rice *SiR* gene

was able to do so. We therefore selected the *SiR* promoter for further characterization.

The *SiR* promoter drives strong and early expression in bundle sheath cells

Sequence upstream of the *SiR* gene comprising nucleotides -2571 to +42 relative to the predicted

translational start site was sufficient to generate expression in the rice bundle sheath. To allow faster

analysis of sequences responsible for this output we domesticated the sequence by removing four

*Bsa*I and *Bpi*I sites such that it was compatible with the modular Golden Gate cloning system. When

this modified *SiR_{pro}* sequence was placed upstream of the GUS reporter it also generated bundle

sheath preferential accumulation **(Figure 1A)**. Fusion to a nuclear-targeted mTurquoise2 fluorescent

protein confirmed that the *SiR* sequence was sufficient to direct expression to bundle sheath cells,

and also revealed expression in the longer nuclei of veinal cells **(Figure 1B)**. There were no

detectable differences in levels of expression between domesticated and non-domesticated

sequences **(Figure 1C)**. Compared with 0.58 nmol 4-MU/min/mg protein previously reported for the

ZjPCK$_{pro}$ (Emmerling 2018) activity from the *SiR* promoter was at least 36% higher. Designer

Transcription Activator-Like Effector (dTALEs) and cognate Synthetic TALE- Activated Promoters

(STAPs) allow multiple transgenes to be driven by a single promoter (Brückner et al., 2015; Danila

et al., 2022). To test whether bundle sheath expression mediated by the *SiR* promoter can be

maintained and strengthened by the dTALE-STAP system, it was placed upstream of the dTALE

sequence. STAP62 and STAP4 were fused to GUS and a chloroplast targeted mTurquoise2

respectively. Stable transformants harbouring these reporters showed bundle sheath specific

expression **(Supplemental Figure 3A, 3B)**. GUS activity mediated by dTALE and STAP62 was ~18-

fold higher than GUS driven by the endogenous *SiR* promoter (**Supplemental Figure 3C**). We

conclude that *SiR* is compatible with the dTALE-STAP system and its activity can be strengthened.

We also investigated when activity from the *SiR* promoter was first detected during leaf development

and discovered that GUS as well as fluorescence from mTurquoise2 were visible in 5-20mm long

fourth leaves at Plastochron 3 (**Supplemental Figure 4**). This was not the case for *ZjPCK$_{pro}$* even

when a dTALE was used to amplify expression (**Supplemental Figure 4**). We conclude that the *SiR*

promoter initiates expression in the bundle sheath before the *ZjPCK* promoter, and that it is also able

to sustain higher levels of expression in this cell type.

A distal enhancer and Y-patch necessary for expression in the bundle sheath

Analysis of the *SiR* promoter sequence revealed a highly complex *cis* landscape **(Figure 1D)**

comprising at least 638 motifs from 56 transcription factor families (**Supplemental table 1**). We

therefore designed a 5' truncation series to investigate regions necessary for expression in the

bundle sheath (**Figure 1E**). Deleting nucleotides -2571 to -2180 and -1490 to -980 led to a

statistically significant reduction and then increase in MUG activity respectively but neither truncation

abolished preferential accumulation of GUS in the bundle sheath **(Figure 1E-F).** However, when

nucleotides -980 to -394 upstream of the predicted translational start site were removed GUS was

no longer detectable in bundle sheath cells (**Figure 1E-F**). Consistent with this, MUG assays showed

a statistically significantly reduction in activity when these nucleotides were absent **(Figure 1G)**.

Thus, nucleotides spanning -980 to -394 of the *SiR* promoter are necessary for bundle sheath

specific expression.

To test whether this region is sufficient for bundle sheath specific expression we linked it to the

minimal CaMV35S core promoter. Although weak GUS signal was detected in a few veinal cells it

failed to do so in the bundle sheath **(Figure 1E-G)**. We conclude that sequence in two regions of the

promoter (from -394 to +42 and from -980 to -394) interact to specify expression to the bundle sheath.

To better understand this interaction we next generated unbiased 5' and 3' deletions. This second

deletion series further reinforced the notion that the *SiR* promoter contains a complex *cis*-regulatory

landscape. For example, when nucleotides -980 to -829 were removed very weak GUS staining was

observed and the MUG assay confirmed that activity was significantly reduced to 11% of the full-

length promoter (**Figure 2, Supplemental Figure 5**). We conclude that nucleotides -980 to -829

from the *SiR* promoter are necessary for tuning expression in the leaf. However, when nucleotides -

829 to -700 were removed GUS appeared in mesophyll cells (**Supplemental Figure 5**). Truncating

nucleotides -613 to -529 abolished GUS accumulation (**Supplemental Figure 5**). The 3' deletion

that removed nucleotides -251 to +42 also stopped accumulation of GUS in both bundle sheath and

mesophyll cells (**Figure 2A-C, Supplemental Figure 5**). Notably, when the distal region required for

bundle sheath expression (-980 to -829) was combined with nucleotides -251 to +42 these two

regions were sufficient for patterning to this cell type (**Figure 2**).

Having identified a region in the *SiR* promoter that was necessary and sufficient for patterning to

the bundle sheath, we next used phylogenetic shadowing and yeast one hybrid analysis to better

understand the *cis*-elements and *trans*-factors responsible. Analysis of *cis*-elements in the *SiR*

promoter that are highly conserved in grasses identified a short region located from nucleotides -588

to -539 that contained an *ETHYLENE INSENSITIVE3-LIKE 3* (*EIL3*) transcription factor binding site

(**Supplemental Figure 6A&B**). Whilst deletion of this motif had no detectable effect of patterning to

the bundle sheath (**Supplemental Figure 6C**) the level of expression was reduced (**Supplemental**

Figure 6D). We infer that the *EIL3* motif positively regulates activity of the *SiR* promoter but is not

responsible for cell specificity. These data are consistent with the promoter truncation analysis that

showed nucleotides -613 to -529 containing this motif were not required for bundle sheath specific

expression, but instead function as a constitutive activator (**Supplemental Figure 5**). When yeast

one hybrid was used to search for transcription factors capable of binding the *SiR* promoter, sixteen

were identified (**Supplemental Figure 7A, 7B**). For each, cognate binding sites were present. This

included TCP21 and OsOBF1 that can bind to TCP motifs and bZIP_A or Ocs elements respectively.

Consistent with the outcome of deleting the *EIL3* motif, three EIL transcription factors interacted with

nucleotides -899 to -500 (**Supplemental Figure 7B, 7C**). Examination of transcript abundance in

mature leaves showed that most of these transcription factors were expressed in both bundle sheath

and mesophyll cells (**Supplemental Figure 7D**) implying that combinatorial interaction with cell

specific factors may be required for bundle sheath specific expression from the *SiR* promoter.

The distal enhancer contains a tetrapartite *cis*-regulatory for bundle sheath expression

The truncation analysis above identified two short regions comprising nucleotides -980 to -829

and -251 to +42 that were necessary and sufficient for expression in the rice bundle sheath (**Figure**

2). Sequence spanning nucleotides -251 to +42 includes both the annotated 5' untranslated region

but also likely contains core promoter elements (**Supplemental Figure 8A**). Re-analysis of publicly

available data identified two major transcription start sites at positions -91 (TSS1) and -41 (TSS2)

(**Supplemental Figure 8A**). Although no canonical TATA-box was evident in this region, at position

-130 a TATA-box variant was detected (5'-ATTAAA-3') (Civáň and Švec 2009) that could be

responsible for transcription from TSS1. Moreover, upstream of TSS2 is a putative pyrimidine patch

(Y-patch) that represents an alternate but common TC-rich core promoter motif (Civáň and Švec

2009) (**Supplemental Figure 8A**). Scanning sequence from -251 to +42 for core promoter elements

also identified MTE (Motif Ten Element), BREu (TFIIB Recognition Element upstream) and DCE-S-

I (Downstream Core Element S-I) motifs associated with eukaryotic core promoters (**Supplemental**

Figure 8B). We therefore assume the region upstream of TSS1 and TSS2 contained the core

promoter elements. When consecutive deletions to this sequence were made, statistically significant

reductions in MUG activity were evident but there was no impact on accumulation of GUS in the

bundle sheath. Interestingly, when the Y-patch was retained but the TATA-box like motif removed,

low levels of GUS specific to the bundle sheath were apparent (**Supplemental Figure 8C, 8D, 8E,**

8F), and deletion of the Y-patch completely abolished GUS staining (**Supplemental Figure 8C, 8D,**

8E, 8F). Consistent with the Y-patch being important for bundle sheath expression, when core

promoters from other genes (*PIP1;1, NRT1.1A*) containing a Y-patch were linked to the distal

enhancer from *SiR* bundle sheath expression was detected **(Figure 3A,3B)**, but this was not the

case for genes with only a TATA-box **(Figure 3A,3B)**. GUS activity was higher from the *PIP1;1* core

promoter that contains more Y-patch motifs. Overall, we conclude that the TATA-box like motif is not

required for expression in the bundle sheath, but in contrast the Y-patch is necessary for this

patterning and in combination with a distal enhancer comprising nucleotides -980 to -829 is sufficient

for expression in this cell type.

We next assessed the distal enhancer for transcription factor binding sites. The FIMO algorithm

identified motifs associated with WRKY, G2-like, MYB-related, MADS, DOF, IDD, ARR, SNAC

(Stress-responsive NAC) families. PlantPAN (Chow et al. 2018), which includes historically validated

cis-elements, found an additional *Dc3 Promoter Binding Factor* (*DPBF*) binding site for group A bZIP

transcription factors **(Figure 3D)**. Seven consecutive deletions spanning this enhancer region and

hereafter termed subregions a-g were then generated **(Figure 3D)**. Deletion of subregions a, b, d

and f resulted in loss of GUS from bundle sheath cells, although veinal expression persisted when

subregions a, b and d were absent (**Figure 3E-3G**). MUG analysis showed that deletion of all four

regions significantly reduced promoter activity **(Figure 3G)**. In contrast, deletions of nucleotides -938

to -923 (subregion c), -904 to 873 (subregion e), and -853 to -829 (subregion g) had no impact on

either the pattern or level of expression. Subregions a, b, c and f that were necessary for expression

in the bundle sheath contained unique binding sites for WRKY, G2-like, MYB-related, IDD, NAC and

bZIP (DPBF) transcription factors. To examine the significance of these regions in the context of full-

length *SiR* promoter, consecutive deletions from subregion a to f were generated **(Supplemental**

Figure 10A). Deletion of subregion a, d or f, led to GUS accumulating primarily in mesophyll cells

whereas removal of subregion b, c or e, caused GUS staining in both mesophyll cells and bundle

sheath cells **(Supplemental Figure 10B)**. No significant changes in GUS activity were observed in

these deletion lines **(Supplemental Figure 10C)**. We conclude that that the distal enhancer
generates expression in the bundle sheath due to four distinct sub-regions and that other nucleotides
between -980 to -853 function as repressors of mesophyll expression.

G2-like, MYB-related, IDD and bZIP transcription factors activate the distal enhancer

To gain deeper insight how the distal enhancer operates we employed multiple approaches
including transactivation assay, co-expression analysis and site directed mutagenesis. The distal
enhancer contained WRKY, G2-like, MYBR, IDD, SNAC and bZIP (DPBF) motifs **(Figure 4A)**. We
therefore cloned rice transcription factors from each family and used them as effectors in transient
assays. WRKY121, GLK2, MYBS1, IDD2/3/4/6/10, and bZIP9/10/11 transcription factors activated
expression from the bundle sheath enhancer, whereas the stress-responsive NAC transcription
factors did not **(Figure 4B, Supplemental Figure 11)**. We therefore conclude that the SNAC motif
is not important for activity of the bundle sheath enhancer. Effector assays using pairwise
combinations of transcription factors showed synergistic activation from the distal enhancer when
GLK2 and IDD3,4,6,10 were co-expressed **(Figure 4C)**.
Co-expression analysis using a cell-specific leaf developmental gradient dataset revealed that
GLK2, MYBS1 and IDD4,6,10 transcription factors that bind the G2-like, MYB-related and IDD motifs
respectively were more abundant in mesophyll cells **(Figure 4D)**. However, the WRKY121, IDD2,
and bZIP9/10/11 transcription factors correlated with *SiR* transcript abundance and were
preferentially expressed in bundle sheath cells **(Figure 4D)**. To test whether WRKY121, IDD2, and
bZIP9/10/11 are sufficient to pattern *SiR* expression to specific cells, we mis-expressed them in the
mesophyll in the presence of the *SiR* distal enhancer and Y-patch **(Supplemental Figure 12)**.
Presence of these three factors in mesophyll cells activated expression in this cell type **(Figure 4E)**
suggesting at least one directly interacts with the distal enhancer in bundle sheath cells to drive *SiR*
expression. We next mutated WRKY, MYB-related, IDD, and bZIP motifs. Although mutating the
WRKY did not affect bundle sheath expression pattern, mutations in the MYB-related, IDD, and bZIP
(DPBF) motifs abolished it **(Supplemental Figure 13)**, and so we conclude that the WRKY motif is
not necessary for bundle sheath expression.
In order to test whether the regions containing the G2-like, MYB-related IDD, and bZIP (DPBF)
sites are sufficient to pattern expression to rice bundle sheath cells we concatenated them and fused
them to the core promoter of *SiR* (**Figure 5A**). GUS staining was evident in the bundle sheath
(**Figure 5B**). Oligomerisation of this minimal module increased expression by 23-fold (**Figure 5C**).
Collectively our data indicate that transcription factors belonging to the G2-like, MYB-related, IDD
and bZIP (DPBF) families act cooperatively to decode distinct *cis*-elements in a distal enhancer of
the *SiR* promoter. This regulatory module is both necessary and sufficient for bundle sheath-specific
expression in rice.

Discussion

Expression of multiple genes in the rice bundle sheath is not associated with close upstream

enhancers

Gene expression is determined by interactions between elements in the core promoter allowing

basal levels of transcription (Juven-Gershon and Kadonaga 2009; Haberle and Stark 2018) with

more distal *cis*-regulatory modules (Spitz and Furlong 2012; Shlyueva et al. 2014; Ray-Jones and

Spivakov 2021). Such *cis*-regulatory modules include enhancers and silencers that act as hubs

receiving input from multiple transcription factors and so allow gene expression to respond spatially

and temporally to both internal and external stimuli (Li et al. 2007; Buecker and Wysocka 2012).

After 25 promoters were tested we discovered that the majority were not capable of driving

expression in the rice bundle sheath, and this included ten that generated no detectable activity of

GUS in leaves. In all cases we had cloned sequence between -3191 and -960 nucleotides upstream

of the predicted translational start site and so these data demonstrate that the core promoter and

any enhancers in these regions are not sufficient to direct expression to rice bundle sheath cells.

Combined with the paucity of previously reported promoters active in this cell type (Nomura et al.

2005b; Lee et al. 2021) these data argue either for long range upstream enhancers (Studer et al.

2011; Liu et al. 2015; Li et al. 2019; Yan et al. 2019; Zhao et al. 2022) or other regulatory mechanisms

being important to specify expression in the bundle sheath. Possibilities include transcription factor

binding sites in introns that impact on transcription start site and strongly enhance gene expression

(Rose et al. 2008; Gallegos and Rose 2019; Rose 2019) or in exons where because such sequences

specify amino acid sequence as well as binding of *trans*-factors, they have been termed duons

(Stergachis et al. 2013) Functional analysis showed that duons can pattern expression to the bundle

sheath of the C_4 plant *Gynandropsis gynandra* (Reyna-Llorens et al. 2018), and it is notable that a

genome-wide analysis of transcription factor binding sites in grasses revealed genes preferentially

expressed in bundle sheath cells tended to contain transcription factor binding sites in their coding

sequence (Burgess et al. 2019). It therefore appears possible that gene expression in the bundle

sheath is commonly encoded by non-canonical architecture perhaps based on duons rather than

more traditional enhancer elements upstream of the core promoter.

Despite the above, we discovered four promoters capable of driving expression in the rice bundle

sheath, and each was associated with a gene important in sulphur metabolism. For example, *ATPS*,

SiR and *Fd* all participate in the first two steps of sulphate reductive assimilation, while *HAC1;1*

encodes an arsenate reductase important in the detoxification of arsenate using glutathione that is

a product of sulphur assimilation. Collectively, these data further support the notion that the rice

bundle sheath cell is specialised in sulphur assimilation (Hua et al. 2021).

Two distinct genetic networks governing expression in bundle sheath cells

The only other promoter for which both *cis*-elements and *trans*-factors that are necessary and

sufficient to pattern bundle sheath expression have been reported is from *A. thaliana*. Here, a

bipartite MYC-MYB module upstream of the *MYB76* gene is responsible for this output (Dickinson et
al. 2020b). MYB76 forms part of a network governing glucosinolate biosynthesis in *A. thaliana*, and
so it is notable that the gene regulatory network we report in rice is also associated with sulphur
metabolism. However, rather than the bipartite transcription factor network that regulates bundle
sheath expression in *A. thaliana,* in rice we report a quartet of transcription factors controlling *SiR*
(Figure 6A&6B). The enhancer controlling bundle sheath *SiR* expression in rice comprises four
distinct regions recognised by transcription factors belonging to the G2-like, MYB-related, IDD and
bZIP families **(Figure 6B)**. As loss of the MYB-related, IDD and bZIP motifs abolished expression,
this implies they act co-operatively a notion further supported by the fact that GLK2 and IDD3,4,6,10
synergistically activated promoter output in a transient assay **(Figure 4C)**. It is possible that other
motifs in the enhancer such as MADS, DOFs and ARR act as modulators to tune the level of bundle
sheath expression. In fact, single nuclei sequencing of rice and sorghum during photomorphogenesis
identified DOFs as important for bundle sheath expression (Swift, Luginbuhl et al. 2023).
Interestingly, the three other promoters (*ATPSb, Fd* and *HAC1;1*) we identified as being sufficient to
drive reporter expression to the rice bundle sheath, and also known bundle sheath promoters (*ZjPCK*
and *FtGLDP*) contain the cognate *cis*-elements for G2-like, MYB-related, IDD and bZIP transcription
factors (**Supplementary Figure 14**).

The distal enhancer in the *SiR* promoter operates in conjunction with the core promoter that
contains two transcription start sites, one with an upstream TATA-box and the other a TC-rich
element known as a pyrimidine (Y) patch **(Supplemental Figure 8A)**. The TATA-box is found in
metazoans and plants and allows recognition by the pre-initiation complex (Smale and Kadonaga
2003), but in plants computational analysis showed that many promoters lack a TATA-box and
instead contain a Y-patch (Yamamoto et al. 2007a, 2007b; Bernard et al. 2010). These genes tend
to be relatively steadily expressed and associated with protein metabolism (Bernard et al., 2010),
and presence of a Y-patch can increase core promoter strength (Jores et al. 2021). For *SiR,* whilst
the TATA-box is not required, the Y-patch is needed for expression in the bundle sheath.

This regulatory network comprising the Y-patch and distal enhancer enabling bundle sheath
expression of *SiR* is embedded within a complex *cis*-regulatory landscape upstream of *SiR* with
distinct regions encoding activating and repressing activities (**Figure 6**). For example, the distal
enhancer (nucleotides -980 to -829) activating expression in the bundle sheath is proximate to
sequence (nucleotides -829 to -700) that suppresses expression in the mesophyll. Notably, the distal
enhancer is both essential for mesophyll repression but also sufficient to drive bundle sheath specific
expression. In addition to controlling cell specificity, this complexity likely also facilitates the tuning
of expression to environmental conditions. For instance, the *EIL* motif (position -572 to -552) is
recognised by *ETHYLENE-INSENSITIVE LIKE* (*EIL*) transcription factors that respond to sulphur
deficiency (Maruyama-Nakashita et al. 2006; Dietzen et al. 2020). As transcripts encoding *EIL*
accumulate in both bundle sheath and mesophyll cells in response to sulphate deficiency it seems
likely that transcription factors repressing expression in these mesophyll cells respond in a dynamic

manner. In addition to *EIL*, the yeast one hybrid analysis identified seven other families of transcription factor families that can bind the *SiR* promoter. Many play documented roles during abiotic or biotic stress, with for example OBF1, ERF3, NAP and FLP acting during low-temperature or drought responses (Shimizu et al. 2005; Zhang et al. 2013, 2022; Chen et al. 2014; Qu et al. 2022), while TCP21, EREBP1, ERF3, ERF72, ERF83 are involved in both abiotic and biotic stress (Lin et al. 2007; Jisha et al. 2015; Zhang et al. 2016, 2018; Tezuka et al. 2019; Jung et al. 2021). Consistent with previous *in silico* analysis (Kurt et al. 2022) the presence of multiple AP2/ERF and EIL transcription factors binding sites suggests that *SiR* is likely subject to control from ethylene signalling (Binder 2020) and also of transcription factors that respond to ABA and jasmonic acid biosynthesis and signalling (Yaish et al. 2010; Chen et al. 2014; Jisha et al. 2015; Zhang et al. 2016, 2018). Together this implies that multiple phytohormone signalling pathways converge on the *SiR* promoter. These data are similar to those reported for the *SHORTROOT* promoter in *A. thaliana* roots where a complex network of activating and repressing *trans*-factors also tunes expression (Sparks et al. 2016). Such complexity in the regulatory landscape may enable robustness in maintaining cell specific gene expression, such that individual mutations are less likely to completely abolish the required pattern of expression. It could also allow a fine tuning of gene expression both temporally as well as spatially in response to environmental perturbations. As sulphite reductases are preferentially expressed in the bundle sheath of most species analysed to date (Hua et al. 2021) it will be interesting in the future to determine if this regulatory landscape is shared across these species.

Using the *SiR* promoter to engineer the rice bundle sheath

In addition to bundle sheath cells being important for sulphur assimilation (Leegood 2008; Aubry et al. 2014b; Hua et al. 2021) they have also been implicated in nitrate assimilation, and the control of leaf hydraulic conductance and solute transport (Hua et al. 2021) and the systemic response to high light (Xiong et al. 2021). Moreover, in one of the most striking examples of a cell type being repurposed for a new function, bundle sheath cells have repeatedly been rewired to allow the evolution of C_4 photosynthesis (Sage 2004b). To engineer these diverse processes, specific and tuneable promoters for this cell are required. However, identification of sequence capable of driving specific expression to bundle sheath strands has previously been limited to *A. thaliana* and C_4 species. For example, the *SCARECROW* (Cui et al. 2014), *SCL23* (Cui et al. 2014), *SULT2;2* (Kirschner et al. 2018) and *MYB76* promoters (Dickinson et al. 2020a) derive from *A. thaliana*, whilst the *Glycine Decarboxylase P-protein (GLDP)* promoter is from the C_4 dicotyledon *Flaveria trinervia* (Engelmann et al. 2008; Wiludda et al. 2012). In rice, only the C_4 *Zoysia japonica PCK* and the C_4 *Flaveria trinervia GLDP* promoters are known to pattern expression to the bundle sheath (Nomura et al. 2005c; Lee et al. 2021; Danila et al. 2022b). Both are capable of conditioning expression in this cell type, but are weak, turn on at late leaf developmental stages and the molecular basis

underpinning their ability to restrict expression to the bundle sheath has not been defined. It has

therefore not been possible to rationally design or tune expression to this important cell type in rice.

From analysis of the ~2600 nucleotide *SiR* promoter we identify an enhancer comprising 81

nucleotides that is sufficient to drive expression to bundle sheath cells. Moreover, we show that

output can be tuned via two approaches. First, oligomerising the distal enhancer can drastically

increase expression. Second, combining it with different core promoters achieved the same output,

and this appears to correlate with length of the Y-patch in each promoter. Our identification of a

minimal promoter that drives expression in bundle sheath cells of rice now provides a tool to allow

this important cell type to be manipulated. Cell specific manipulation of gene expression has many

perceived advantages. For example, when constitutive promoters have been used to drive gene

expression gene silencing and reduction of plant fitness due to metabolic penalties can ensue (Glick

1995; Que et al. 1997). In contrast, tissue specific promoters allow targeted gene expression either

spatially or at particular developmental stages and so allow increased precision in trait engineering

and (Kummari et al. 2020). The *SiR* promoter and the bundle sheath *cis*-regulatory module that we

identify provides insights into mechanisms governing cell specific expression in rice, and may also

contribute to our ability to engineer and improve cereal crops.

Materials and methods

Plant material and growth conditions

Kitaake (*O. sativa* ssp. *japonica*) was transformed using *Agrobacterium tumefaciens* with the

following modifications as described previously (Hiei et al., 2008). Mature seeds were sterilized with

2.5% (v/v) sodium hypochlorite for 15 mins, and calli were induced on NB medium with 2 mg/L 2,4-

D at 30 $^{\circ}$C in darkness for 3-4 weeks. Actively growing calli were then co-incubated with *A.*

tumefaciens strain LBA4404 in darkness at 25°C for 3 days, they were selected on NB medium

supplied with 35 mg/L hygromycin B for 4 weeks, and those that proliferated placed on NB medium

with 10 mg/L hygromycin B for 4 weeks at 28 $^{\circ}$C under continuous light. Plants resistant to

hygromycin were planted in 1:1 mixture of top-soil and sand and placed in a greenhouse at the

Botanic Gardens, University of Cambridge under natural light conditions but supplemented with a

minimum light intensity of 390 µmol m^{-2} s^{-1}, a humidity of 60%, temperatures of 28°C and 23°C

during the day and night respectively, and a photoperiod of 12 h light, 12 h dark. Subsequent

generations were grown in a growth cabinet in 12 h light/12 h dark, at 28 $^{\circ}$C, a relative humidity of

65%, and a photon flux density of 400 µmol m^{-2} s^{-1}.

Cloning and construct preparation, and motif analysis

The 2613-bp promoter DNA fragment of *SULFITE REDUCTASE* (*SiR,* MSU7 ID:

LOC_Os05g42350, RAP-DB ID: Os05g0503300) was originally amplified from Kitaake genomic

DNA, with forward primer (5'-3') "CACCATGCTTGACCATGTGGACTC" and reverse primer (5'-3')

"ACGGAACCCGTGGAACTC". Gel-purified PCR product was cloned into a Gateway pENTR^{TM}

vector to generate *pENTR-SiRpro* using pENTR™/D-TOPO™ Cloning Kit (Invitrogen), then the

promoter was recombined into the pGWB3 expression vector and fused with *GUS* gene using LR

reaction. The resultant vector was transformed into *A. tumefaciens* strain LBA4404 and transformed

into Kitaake. To engineer the *SiRpro* such that it is compatible with the Golden gate system, four

*Bsa*I or *Bpi*I restriction enzyme recognition sites at -214, -298, -1468, and -2309 were mutated from

T to A and cloned into the *pAGM9121* vector using Golden Gate level 0 cloning reactions and then

into a level 0 PU module EC14328, which was used for driving *kzGUS* (intronless GUS) and *H2B-*

mTurquoise2 reporter genes via Golden gate reaction and using *Tnos* as a terminator. A five prime

deletion series was generated using EC14328 as the template and prepared as level 0 PU modules,

and a three prime deletion series prepared as level 0 P modules. The minimal *CaMV35S* promoter

was used as the U module, and they were linked with *kzGUS* and terminated with *Tnos*.

To test *SiRpro* in the dTALE/STAP system, the 42-bp coding region were excluded and the 2571-

bp resultant fragment placed into a level 0 PU module EC14330 and was used to drive *dTALE1*.

Two reporters were used. For the GUS reporter *kzGUS* was linked with *STAP62* and terminated with

Tnos. In the fluorescent reporter construct, a chloroplast targeting peptide fused to *mTurquoise 2*

was linked with *STAP 4* and terminated with *Tact2*. In both constructs, *pOsAct1* driving *HYG*

(Hygromycin resistant gene) was terminated with *Tnos* and used as the selection marker during rice

transformation.

The Find Individual Motif Occurrences (FIMO) tool (Grant et al. 2011) from the Multiple Em for Motif

Elucidation (MEME) suite v.5.4.0 (Bailey et al. 2015) was used to search for individual motifs within

the promoter sequences using default parameters with "--thresh" of "1e-3". Position weight matrix of

656 non-redundant plant motifs and 13 RNA polymerase II (POLII) core promoter motifs were

obtained from JASPAR (https://jaspar.elixir.no/downloads/) (Fornes et al. 2020). To cluster the

transcription factor binding motifs, the RSAT matrix-clustering tool (Castro-Mondragon et al. 2017) was

run on all 656 non-redundant plant motifs using the default parameters, which yield 51 motif clusters,

these clusters were further divided based on transcription factor families (Supplemental table 2).

Analysis of GUS and fluorescent reporters

GUS staining was performed as described previously (Jefferson et al., 1987) with the following

minor modifications. Leaf tissue was fixed in 90% (v/v) acetone overnight at 4 $^{\circ}$C after washing with

100 mM phosphate buffer (pH 7.0). Leaf samples were transferred into 1 mg/ml 5-bromo-4-chloro-

3-indolyl glucuronide (X-Gluc) GUS staining solution, subjected to 2 mins vacuum infiltration 5 times,

and then incubated at 37 $^{\circ}$C for between 1 and 168 hours. Chlorophyll was cleared further using 90%

(v/v) ethanol overnight at room temperature. Cross sections were prepared manually using a razor

blade and images were taken using an Olympus BX41 light microscopy. Quantification of GUS

activity was performed using a fluorometric MUG assay (Jefferson et al., 1987). ~200 mg mature

leaves from transgenic plants were frozen in liquid nitrogen and ground into fine powder with a

Tissuelyser (Qiagen). Soluble protein was extracted in 1 mL of 50 mM phosphate buffer (pH 7.0)

supplemented with 0.1% [v/v] Triton X-100 and cOmplete™ Protease Inhibitor Cocktail (half tablet

per 50 mL). Protein concentration then determined using a Qubit protein assay kit (Invitrogen). The

MUG fluorescent assay was performed in duplicates with 20 µl protein extract in MUG assay buffer

(50mM phosphate buffer (pH 7.0), 10 mM EDTA-Na$_2$, 0.1% [v/v] Triton X-100, 0.1% [w/v] N-

lauroylsarcosine sodium, 10 mM DTT, 2 mM 4-methylumbelliferyl-β-D-glucuronide (MUG)) in a 200

µl total volume. The reaction was conducted at 37 $^{\circ}$C in GREINER 96 F-BOTTOM microtiter plate

using a CLARIOstar plate reader. 4-Methylumbelliferone (4-MU) fluorescence was recorded every 2

minutes for 20 cycles with excitation at 360 nm and emission detected at 450 nm. 4-MU

concentration was determined based on a standard curve of ten 4-MU standards placed in the same

plate. GUS enzymatic rates were calculated by averaging the slope of MU production from each of

the duplicate reactions.

In order to visualize mTurquoise2 mature leaves were dissected into 2-cm sections, leaf

epidermal cells were removed by scraping the leaf surface with a razor blade and then mounted with

deionized water. 5-mm and middle sections of 2-cm young tissue of the fourth leaves were dissected

and mounted with deionized water directly. Imaging was then performed using a Leica TCS SP8

confocal laser-scanning microscope using a 20x air objective. mTurquoise2 fluorescence was

excited at 442 nm with emission at 471–481 nm, chlorophyll autofluorescence was excited at 488
nm with emission at 672–692 nm.

Yeast one hybrid, protoplast isolation and transactivation assay

The yeast one hybridisation assay was performed by Hybrigenics (https://www.hybrigenics-
services.com/). Fragments were synthesized and used as bait. Rice leaf and root cDNA libraries
were used as prey. The number of clones screened and concentration of 3-AT were as follows:
fragment 1, 70.2 million clones screened with 0 mM 3-AT; fragment 2, 61.5 million clones screened
with 0 mM 3-AT; fragment 3, 68.4 million clones screened with 20 mM 3-AT; fragment 4, 57.4 million
clones screened with 100 mM 3-AT; fragment 5, 94.2 million clones screened with 200 mM 3-AT.

Rice leaf protoplasts and PEG-mediated transformation were performed as described previously
(Page et al. 2019). Golden gate level 1 modules for transformation were isolated using ZymoPURE™
II Plasmid Midiprep Kit, *ZmUBIpro::GUS-Tnos* was used as transformation control. Transcription
factor coding sequences were amplified using rice leaf cDNA, with *Bsa*I and *Bpi*I sites mutated, and
cloned into Golden gate SC level 0 modules. They were assembled into a level 1 module with a
ZmUBIpro promoter and *Tnos* terminator module. Nucleotides -980 to -829 with the endogenous
core promoter (nucleotide -250 to +42) were fused with the *LUC* reporter to generate output of
transcription activity. In each transformation, 2 μg of transformation control plasmids, 5 μg of reporter
plasmids, and 5 μg of effector plasmids per transcription factor were combined and mixed with 170
μl protoplasts. After incubation on the benchtop for overnight protein was extracted using passive
lysis buffer, GUS activity was determined with 20 μl of protein sample and MUG fluorescent assay
as described above, LUC activity was measured with 20 μl of protein sample and 100 μl of LUC
assay reagent (Promega) using Clariostar plate reader. Transcriptional activity from the promoter
was calculated as LUC luminescence/rate of MUG accumulation.

Acknowledgements

This work was funded by the Bill and Melinda Gates Foundation C_4 Rice grant awarded to the
University of Oxford (2015-2019 (OPP1129902) and 2019-2024 (INV-002970). For the purpose of
open access, the authors have applied a Creative Commons Attribution (CC BY) licence to any
Author Accepted Manuscript version arising from this submission.

Author contributions

L.H. and J.M.H. conceived the work. J.M.H. guided execution of experiments and oversaw the
project. L.H., N.W., S.S., R.D., K.B., and A.R.B. did the experiments and analysed the data. L.H. and
J.M.H. wrote the manuscript with input from all authors.

Declaration of interests

The authors declare no competing interests.

Figure legends

Figure 1. Nucleotides -980 to -394 of the *SiR* promoter are necessary for bundle sheath expression. (A) Domesticated *SiR* promoter generates strong GUS staining in bundle sheath. **(B)** mTurquoise2 signal driven by the domesticated *SiR* promoter in nuclei (indicated by yellow arrows) of bundle sheath cells (marked by yellow dashed lines) and vein cells in mature leaves, red indicates chlorophyll autofluorescence. **(C)** The fluorometric 4-methylumbelliferyl-β-D-glucuronide (MUG) assay shows no statistically significant difference between the endogenous and domesticated *SiR* promoter activity. **(D)** Landscape of transcription factor binding sites in the *SiR* promoter using the Find Individual Motif Occurrences (FIMO) program. The likelihood of match between 656 plant nonredundant known transcription factor motifs in the *SiR* promoter is shown by transcription factor families (Supplemental table 1). **(E)** Schematics of 5' truncations. **(F)** Representative images of leaf cross sections from transgenic lines after GUS staining. Zoomed-in images of lateral veins shown in right panels, bundle sheath cells were highlighted with white dash line, scale bars = 50 μm. **(G)** Promoter activity determined by the fluorometric 4- methylumbelliferyl-β-D-glucuronide (MUG) assay. Data subjected to a Kruskal-Wallis test, median indicates median value of catalytic rate of GUS, n indicates total number of transgenic lines assessed.

Figure 2. A distal enhancer and the core promoter are necessary and sufficient for bundle sheath gene expression (A) Schematics showing deletions of nucleotides -980 to -849 and -119 to +42. **(B)** Representative image of leaf cross sections of transgenic lines after GUS staining. Zoomed-in images of lateral veins shown in right panels, bundle sheath cells were highlighted with white dash line, scale bars = 50 μm. **(C)** Promoter activity determined by the fluorometric 4-methylumbelliferyl-β-D-glucuronide (MUG) assay, data subjected to a Kruskal-Wallis test, median indicates median value of catalytic rate of GUS protein, n indicates total number of transgenic lines assessed.

Figure 3. The Y-patch and four short regions in the distal enhancer are required for bundle sheath specific expression. (A-C) Nucleotides -980 and -829 from the *SiR* promoter pattern expression to the bundle sheath when linked with the *PIP1;1* and *NRT1.1A* core promoters containing Y-patches. **(A)** Prediction of Y-patch and TATA-box sequences in core promoters of *PIP1;1*, *NRT1.1A*, *PIP1;3* and *ATPSb*. **(B)** Representative cross sections of transgenic rice leaves after GUS staining, zoomed-in image of lateral veins shown in the right panel, bundle sheath cells highlighted with white dashed lines, scale bars = 50 μm. **(C)** Promoter activity determined by the fluorometric 4- methylumbelliferyl-β-D-glucuronide (MUG) assay. Data were subjected to a Kruskal-Wallis test, median indicates median value of catalytic rate of GUS protein, n indicates total number of transgenic lines assessed. **(D)** Schematics showing transcription factor binding sites between nucleotides -980 and -829. **(E)** Schematics showing consecutive deletions between nucleotides -980 and -829 fused to the GUS reporter. **(F)** Representative images of cross sections from transgenic

lines after GUS staining, zoomed-in images of lateral veins shown in right panels, bundle sheath

cells highlighted with white dashed lines, scale bars = 50 μm. **(G)** Promoter activity determined by

the fluorometric 4-methylumbelliferyl-β-D-glucuronide (MUG) assay. Data were subjected to a

Kruskal-Wallis test, median indicates median value of catalytic rate of GUS protein, n indicates total

number of transgenic lines assessed.

Figure 4. G2-like, MYB-related, IDD and bZIP transcription factors interact and activate the

distal enhancer. (A) Schematics showing transcription factor binding sites between nucleotides -

980 and -829 which are likely required for bundle sheath specific expression. **(B)** Effector assays

showing that each transcription factor activates expression from the distal enhancer. **(C)** Effector

assays showing synergistic activation from the distal enhancer when GLK2 and IDD3,4,6,10 were

co-expressed. **(D)** Transcript abundance of transcription factors in bundle sheath strands (BSS) and

mesophyll (M) cells during maturation. Leaf developmental stage S2 to S7 represent base of the 4[th]

leaf at the 6th, 8th, 9th, 10th, 13th and 17th day after sowing. **(E)** Representative image of transgenic

lines expressing WRKY121, IDD2 and bZIP9 in mesophyll cells, zoomed-in image of lateral veins

shown in right panel, bundle sheath cells highlighted with white dashed lines.

Figure 5. Oligomerisation of the distal enhancer increases bundle sheath expression. (A)

Schematic showing the distal enhancer comprises three short regions and oligomerisation this

enhancer, fusing it to the *SiR* core promoter sequence. **(B)** Representative images of cross sections

from transgenic lines after GUS staining, zoomed-in image of lateral veins shown in right panel,

bundle sheath cells highlighted with white dashed lines, scale bars = 50 μm. **(C)** Promoter activity

determined by the fluorometric 4-methylumbelliferyl-β-D-glucuronide (MUG) assay. Data subjected

to a Kruskal-Wallis test, median indicates median value of catalytic rate of GUS protein, n indicates

total number of transgenic lines assessed.

Figure 5. Model depicting mechanism underpinning bundle sheath expression from the *SiR*

promoter in rice. (**A&B**) Schematics illustrating the bundle sheath cis regulatory module (BS-CRM)

present in the distal enhancer, along with a Y-patch, and *trans*-acting factors that can bind and act

as activators. (**C**) Schematic indicating how the enhancer operates within a broader *cis*-regulatory

landscape including a constitutive activator and mesophyll repressor.

Supplementary Figure legend titles

Supplemental Figure 1. Analysis of seven rice promoters identified after analysis of transcripts that
accumulate preferentially in bundle sheath strands**.**

Supplemental Figure 2. Identification of bundle sheath specific promoters after analysis of
transcripts that accumulate preferentially in Kitaake bundle sheath compared with mesophyll cells.

Supplemental Figure 3. The domesticated *SiR* promoter combined with the dTALE/STAP system
drives strong expression in the rice bundle sheath.

Supplemental Figure 4. The rice *SiR* promoter drives expression in bundle sheath cells earlier than
the *Zoysia japonica PCK* promoter.

Supplemental Figure 5. Impact of 5' and 3' deletions between on patterning of GUS from the *SiR*
promoter.

Supplemental Figure 6. The evolutionarily conserved *ETHYLENE INSENSITIVE3-LIKE* (*EIL*)
binding site regulates the expression level but not cell specificity of the *SiR* promoter*.*

Supplemental Figure 7. Identification of transcription factors interacting with the *SiR* promoter*.*

Supplemental Figure 8. Nucleotides -251 to -1 likely serve as a core promoter.

Supplemental Figure 9. Subregions in the distal enhancer not required for bundle sheath specific
expression between nucleotides -980 to -829.

Supplemental Figure 10. Regions between nucleotides -980 and -829 that in combination with
nucleotides -828 to -252 repress mesophyll expression.

Supplemental Figure 11. Summary of transactivation assay using rice protoplasts.

Supplemental Figure 12. Schematic showing mis-expression of WRKY121, IDD2 and bZIP9 in
mesophyll cells using the maize *PEPC* promoter driving dTALE1 in combination with STAP56, 45
and 62.

Supplemental Figure 13. Site-directed mutagenesis of transcription factor binding sites in
subregions a, b, d and f.

Supplemental Figure 14. G2-like, MYB-related, IDD and bZIP transcription factor binding sites
identified by FIMO program in *HAC1;1*, *ATPSb*, *Fd*, *ZjPCK* and *FtGLDP* promoters. Regions <300
bp containing all four motif families highlighted in red.

References

Aubry S, Kelly S, Kümpers BMCC, Smith-Unna RD, and Hibberd JM. Deep Evolutionary Comparison of Gene Expression Identifies Parallel Recruitment of Trans-Factors in Two Independent Origins of C$_4$ Photosynthesis. PLoS Genet. 2014a:**10**(6):e1004365. https://doi.org/10.1371/journal.pgen.1004365

Aubry S, Smith-Unna RD, Boursnell CM, Kopriva S, and Hibberd JM. Transcript residency on ribosomes reveals a key role for the *Arabidopsis thaliana* bundle sheath in sulfur and glucosinolate metabolism. Plant Journal. 2014b:**78**(4):659–673. https://doi.org/10.1111/tpj.12502

Bailey TL, Johnson J, Grant CE, and Noble WS. The MEME Suite. Nucleic Acids Res. 2015:**43**:39–49. https://doi.org/10.1093/nar/gkv416

Bernard V, Brunaud V, and Lecharny A. TC-motifs at the TATA-box expected position in plant genes: A novel class of motifs involved in the transcription regulation. BMC Genomics. 2010:**11**(1):1–15. https://doi.org/10.1186/1471-2164-11-166/TABLES/4

Binder BM. Ethylene signaling in plants. J Biol Chem. 2020:**295**(22):7710. https://doi.org/10.1074/JBC.REV120.010854

Buecker C and Wysocka J. Enhancers as information integration hubs in development: lessons from genomics. Trends Genet. 2012:**28**(6):276. https://doi.org/10.1016/J.TIG.2012.02.008

Burgess SJ, Reyna-Llorens I, Stevenson SR, Singh P, Jaeger K, and Hibberd JM. Genome-Wide Transcription Factor Binding in Leaves from C3 and C4 Grasses. Plant Cell. 2019:**31**(10):2297–2314. https://doi.org/10.1105/TPC.19.00078

Castro-Mondragon JA, Jaeger S, Thieffry D, Thomas-Chollier M, and Van Helden J. RSAT matrix-clustering: dynamic exploration and redundancy reduction of transcription factor binding motif collections. Nucleic Acids Res. 2017:**45**(13):e119–e119. https://doi.org/10.1093/NAR/GKX314

Chen X, Wang Y, Lv B, Li J, Luo L, Lu S, Zhang X, Ma H, and Ming F. The NAC family transcription factor OsNAP confers abiotic stress response through the ABA pathway. Plant Cell Physiol. 2014:**55**(3):604–619. https://doi.org/10.1093/PCP/PCT204

Chow C-N, Lee T-Y, Hung Y-C, Li G-Z, Tseng K-C, Liu Y-H, Kuo P-L, Zheng H-Q, and Chang W-C. PlantPAN3.0: a new and updated resource for reconstructing transcriptional regulatory networks from ChIP-seq experiments in plants. Nucleic Acids Res. 2018. https://doi.org/10.1093/nar/gky1081

Civáň P and Švec M. Genome-wide analysis of rice (Oryza sativa L. subsp. japonica) TATA box and Y Patch promoter elements. Genome. 2009:**52**(3):294–297. https://doi.org/10.1139/G09-001/SUPPL_FILE/G09-001.DOC

Cui H. Challenges and Approaches to Crop Improvement Through C3-to-C4 Engineering. Front Plant Sci. 2021:**12**:715391. https://doi.org/10.3389/FPLS.2021.715391/BIBTEX

Cui H, Kong D, Liu X, and Hao Y. SCARECROW, SCR-LIKE 23 and SHORT-ROOT control bundle sheath cell fate and function in Arabidopsis thaliana. The Plant Journal. 2014:**78**(2):319–327. https://doi.org/10.1111/TPJ.12470

Danila F, Schreiber T, Ermakova M, Hua L, Vlad D, Lo SF, Chen YS, Lambret-Frotte J, Hermanns AS, Athmer B, et al. A single promoter-TALE system for tissue-specific and tuneable expression of multiple genes in rice. Plant Biotechnol J. 2022a:**20**(9):1786–1806. https://doi.org/10.1111/PBI.13864

Danila F, Schreiber T, Ermakova M, Hua L, Vlad D, Lo SF, Chen YS, Lambret-Frotte J, Hermanns AS, Athmer B, et al. A single promoter-TALE system for tissue-specific and tuneable expression of multiple genes in rice. Plant Biotechnol J. 2022b. https://doi.org/10.1111/PBI.13864

Dickinson PJ, Kneřová J, Szecówka M, Stevenson SR, Burgess SJ, Mulvey H, Bågman AM, Gaudinier A,
Brady SM, and Hibberd JM. A bipartite transcription factor module controlling expression in the
bundle sheath of Arabidopsis thaliana. Nat Plants. 2020a:**6**(12):1468–1479.
https://doi.org/10.1038/s41477-020-00805-w

Dickinson PJ, Kneřová J, Szecówka M, Stevenson SSR, Burgess SJ, Mulvey H, Bågman A-MM, Gaudinier A,
Brady SM, and Hibberd JM. A bipartite transcription factor module controlling expression in the
bundle sheath of Arabidopsis thaliana. Nat Plants. 2020b:**6**(12):1468–1479.
https://doi.org/10.1038/s41477-020-00805-w

Dietzen C, Koprivova A, Whitcomb SJ, Langen G, Jobe TO, Hoefgen R, and Kopriva S. The Transcription
Factor EIL1 Participates in the Regulation of Sulfur-Deficiency Response. Plant Physiol.
2020:**184**(4):2120–2136. https://doi.org/10.1104/pp.20.01192

Emmerling J. Studies into the Regulation of C4 Photosynthesis – Towards Factors Controlling Bundle Sheath
Expression and Kranz Anatomy Development. Doctoral thesis, Heinrich Heine University, Düsseldorf.
2018:(September).

Engelmann S, Wiludda C, Burscheidt J, Gowik U, Schlue U, Koczor M, Streubel M, Cossu R, Bauwe H, and
Westhoff P. The Gene for the P-Subunit of Glycine Decarboxylase from the C4 Species Flaveria
trinervia: Analysis of Transcriptional Control in Transgenic Flaveria bidentis (C4) and Arabidopsis (C3).
Plant Physiol. 2008:**146**(4):1773–1785. https://doi.org/10.1104/PP.107.114462

Ermakova M, Arrivault S, Giuliani R, Danila F, Alonso-Cantabrana H, Vlad D, Ishihara H, Feil R, Guenther
M, Borghi GL, et al. Installation of C$_4$ photosynthetic pathway enzymes in rice using a single construct.
Plant Biotechnol J. 2021:**19**(3):575–588. https://doi.org/10.1111/pbi.13487

Fornes O, Castro-Mondragon JA, Khan A, Van Der Lee R, Zhang X, Richmond PA, Modi BP, Correard S,
Gheorghe M, Baranašić D, et al. JASPAR 2020: Update of the open-Access database of transcription
factor binding profiles. Nucleic Acids Res. 2020:**48**(D1):D87–D92.
https://doi.org/10.1093/nar/gkz1001

Gallegos JE and Rose AB. An intron-derived motif strongly increases gene expression from transcribed
sequences through a splicing independent mechanism in Arabidopsis thaliana. Scientific Reports 2019
9:1. 2019:**9**(1):1–9. https://doi.org/10.1038/s41598-019-50389-5

Glick BR. Metabolic load and heterologous gene expression. Biotechnol Adv. 1995:**13**(2):247–261.
https://doi.org/10.1016/0734-9750(95)00004-A

Grant CE, Bailey TL, and Noble WS. FIMO: Scanning for occurrences of a given motif. Bioinformatics.
2011:**27**(7):1017–1018. https://doi.org/10.1093/bioinformatics/btr064

Haberle V and Stark A. Eukaryotic core promoters and the functional basis of transcription initiation. Nat
Rev Mol Cell Biol. 2018:**19**(10):621. https://doi.org/10.1038/S41580-018-0028-8

Hibberd JM and Covshoff S. The Regulation of Gene Expression Required for C$_4$ Photosynthesis. Annu Rev
Plant Biol. 2010:**61**(1):181–207. https://doi.org/10.1146/annurev-arplant-042809-112238

Hibberd JM, Sheehy JE, and Langdale JA. Using C4 photosynthesis to increase the yield of rice-rationale
and feasibility. Curr Opin Plant Biol. 2008:**11**(2):228–231. https://doi.org/10.1016/j.pbi.2007.11.002

Hua L, Stevenson SR, Reyna-Llorens I, Xiong H, Kopriva S, and Hibberd JM. The bundle sheath of rice is
conditioned to play an active role in water transport as well as sulfur assimilation and jasmonic acid
synthesis. The Plant Journal. 2021:**107**(1):268–286. https://doi.org/10.1111/TPJ.15292

Jisha V, Dampanaboina L, Vadassery J, Mithöfer A, Kappara S, and Ramanan R. Overexpression of an
AP2/ERF Type Transcription Factor OsEREBP1 Confers Biotic and Abiotic Stress Tolerance in Rice. PLoS
One. 2015:**10**(6). https://doi.org/10.1371/JOURNAL.PONE.0127831

Jores T, Tonnies J, Wrightsman T, Buckler ES, Cuperus JT, Fields S, and Queitsch C. Synthetic promoter
designs enabled by a comprehensive analysis of plant core promoters. Nature Plants 2021 7:6.
2021:**7**(6):842–855. https://doi.org/10.1038/s41477-021-00932-y

Jung SE, Bang SW, Kim SH, Seo JS, Yoon H Bin, Kim YS, and Kim JK. Overexpression of OsERF83, a Vascular
Tissue-Specific Transcription Factor Gene, Confers Drought Tolerance in Rice. Int J Mol Sci.
2021:**22**(14). https://doi.org/10.3390/IJMS22147656

Juven-Gershon T and Kadonaga JT. Regulation of gene expression via the core promoter and the basal
transcriptional machinery. 2009. https://doi.org/10.1016/j.ydbio.2009.08.009

Kajala K, Covshoff S, Karki S, Woodfield H, Tolley BJ, Dionora MJA, Mogul RT, Mabilangan AE, Danila FR,
Hibberd JM, et al. Strategies for engineering a two-celled C4 photosynthetic pathway into rice. J Exp
Bot. 2011a:**62**(9):3001–3010. https://doi.org/10.1093/jxb/err022

Kajala K, Covshoff S, Karki S, Woodfield H, Tolley BJ, Dionora MJA, Mogul RT, Mabilangan AE, Danila FR,
Hibberd JM, et al. Strategies for engineering a two-celled C4photosynthetic pathway into rice. J Exp
Bot. 2011b:**62**(9):3001–3010. https://doi.org/10.1093/jxb/err022

Kirschner S, Woodfield H, Prusko K, Koczor M, Gowik U, Hibberd JM, and Westhoff P. Expression of
SULTR2;2, encoding a low-affinity sulphur transporter, in the Arabidopsis bundle sheath and vein cells
is mediated by a positive regulator. J Exp Bot. 2018:**69**(20):4897–4906.
https://doi.org/10.1093/jxb/ery263

Kummari D, Palakolanu SR, Kishor PBK, Bhatnagar-Mathur P, Singam P, Vadez V, and Sharma KK. An
update and perspectives on the use of promoters in plant genetic engineering. J Biosci. 2020:**45**(1):1–
24. https://doi.org/10.1007/S12038-020-00087-6/TABLES/7

Kurt F, Filiz E, and Aydın A. Sulfite Reductase (SiR) Gene in Rice (Oryza sativa): Bioinformatics and
Expression Analyses Under Salt and Drought Stresses. J Plant Growth Regul. 2022:**41**(6):2246–2260.
https://doi.org/10.1007/S00344-021-10438-8/FIGURES/9

Lee DY, Hua L, Khoshravesh R, Giuliani R, Kumar I, Cousins A, Sage TL, Hibberd JM, and Brutnell TP.
Engineering chloroplast development in rice through cell-specific control of endogenous genetic
circuits. Plant Biotechnol J. 2021:**19**(11):2291–2303. https://doi.org/10.1111/PBI.13660

Leegood RC. Roles of the bundle sheath cells in leaves of C_3 plants. J Exp Bot. 2008:**59**(7):1663–1673.
https://doi.org/10.1093/jxb/erm335

Li E, Liu H, Huang L, Zhang X, Dong X, Song W, Zhao H, and Lai J. Long-range interactions between proximal
and distal regulatory regions in maize. Nat Commun. 2019:**10**(1). https://doi.org/10.1038/S41467-
019-10603-4

Li L, Zhu Q, He X, Sinha S, and Halfon MS. Large-scale analysis of transcriptional cis-regulatory modules
reveals both common features and distinct subclasses. Genome Biol. 2007:**8**(6).
https://doi.org/10.1186/GB-2007-8-6-R101

Lin R, Zhao W, Meng X, and Peng YL. Molecular cloning and characterization of a rice gene encoding
AP2/EREBP-type transcription factor and its expression in response to infection with blast fungus and
abiotic stresses. Physiol Mol Plant Pathol. 2007:**70**(1–3):60–68.
https://doi.org/10.1016/J.PMPP.2007.06.002

Liu L, Du Y, Shen X, Li M, Sun W, Huang J, Liu Z, Tao Y, Zheng Y, Yan J, et al. KRN4 Controls Quantitative
Variation in Maize Kernel Row Number. PLoS Genet. 2015:**11**(11):e1005670.
https://doi.org/10.1371/JOURNAL.PGEN.1005670

Makino A, Sakuma H, Sudo E, and Mae T. Differences between Maize and Rice in N-use Efficiency for
Photosynthesis and Protein Allocation.

Maruyama-Nakashita A, Nakamura Y, Tohge T, Saito K, and Takahashi H. *Arabidopsis* SLIM1 is a central
transcriptional regulator of plant sulfur response and metabolism. Plant Cell. 2006:**18**(11):3235–3251.
https://doi.org/10.1105/tpc.106.046458

Mitchell PL and Sheehy JE. Supercharging rice photosynthesis to increase yield. New Phytol.
2006:**171**(4):688–693. https://doi.org/10.1111/J.1469-8137.2006.01855.X

Nomura M, Higuchi T, Ishida Y, Ohta S, Komari T, Imaizumi N, Miyao-Tokutomi M, Matsuoka M, and
Tajima S. Differential Expression Pattern of C_4 Bundle Sheath Expression Genes in Rice, a C_3 Plant.
Plant Cell Physiol. 2005a:**46**(5):754–761. https://doi.org/10.1093/pcp/pci078

Nomura M, Higuchi T, Ishida Y, Ohta S, Komari T, Imaizumi N, Miyao-Tokutomi M, Matsuoka M, and
Tajima S. Differential expression pattern of C4 bundle sheath expression genes in rice, a C3 plant.
Plant Cell Physiol. 2005b:**46**(5):754–761. https://doi.org/10.1093/PCP/PCI078

Nomura M, Higuchi T, Ishida Y, Ohta S, Komari T, Imaizumi N, Miyao-Tokutomi M, Matsuoka M, and
Tajima S. Differential expression pattern of C4 bundle sheath expression genes in rice, a C3 plant.
Plant Cell Physiol. 2005c:**46**(5):754–761. https://doi.org/10.1093/PCP/PCI078

Page MT, Parry MAJ, and Carmo-Silva E. A high-throughput transient expression system for rice. Plant Cell
Environ. 2019:**42**(7):2057–2064. https://doi.org/10.1111/PCE.13542

Qu X, Zou J, Wang J, Yang K, Wang X, and Le J. A Rice R2R3-Type MYB Transcription Factor OsFLP Positively
Regulates Drought Stress Response via OsNAC. Int J Mol Sci. 2022:**23**(11).
https://doi.org/10.3390/IJMS23115873

Que Q, Wang HY, English JJ, and Jorgensen RA. The Frequency and Degree of Cosuppression by Sense
Chalcone Synthase Transgenes Are Dependent on Transgene Promoter Strength and Are Reduced by
Premature Nonsense Codons in the Transgene Coding Sequence. Plant Cell. 1997:**9**(8):1357.
https://doi.org/10.1105/TPC.9.8.1357

Rask L, Andréasson E, Ekbom B, Eriksson S, Pontoppidan B, and Meijer J. Myrosinase: Gene family
evolution and herbivore defense in Brassicaceae. Plant Mol Biol. 2000:**42**(1):93–114.
https://doi.org/10.1023/A:1006380021658/METRICS

Ray-Jones H and Spivakov M. Transcriptional enhancers and their communication with gene promoters.
Cellular and Molecular Life Sciences. 2021:**78**(19–20):6453. https://doi.org/10.1007/S00018-021-
03903-W

Reyna-Llorens I, Burgess SJ, Reeves G, Singh P, Stevenson SR, Williams BP, Stanley S, and Hibberd JM.
Ancient duons may underpin spatial patterning of gene expression in C4 leaves. Proc Natl Acad Sci U S
A. 2018:**115**(8):1931–1936.
https://doi.org/10.1073/PNAS.1720576115/SUPPL_FILE/PNAS.1720576115.SD07.XLSX

Rosa EAS and Rodrigues PMF. Towards a more sustainable agriculture system: The effect of glucosinolates
on the control of soil-borne diseases. Journal of Horticultural Science and Biotechnology.
1999:**74**(6):667–674.
https://doi.org/10.1080/14620316.1999.11511170/ASSET//CMS/ASSET/B3CB692A-CEFD-4DF3-8FBE-
D467F93CF12C/14620316.1999.11511170.FP.PNG

Rose AB. Introns as Gene Regulators: A Brick on the Accelerator. Front Genet. 2019:**9**(FEB).
https://doi.org/10.3389/FGENE.2018.00672

Rose AB, Elfersi T, Parra G, and Korf I. Promoter-proximal introns in Arabidopsis thaliana are enriched in
dispersed signals that elevate gene expression. Plant Cell. 2008:**20**(3):543–551.
https://doi.org/10.1105/TPC.107.057190

Sage RF. The evolution of C4 photosynthesis. New Phytologist. 2004a:**161**(2):341–370.

Sage RF. The evolution of C_4 photosynthesis. New Phytologist. 2004b:**161**(2):341–370.
https://doi.org/10.1111/j.1469-8137.2004.00974.x

Schuler ML, Mantegazza O, and Weber APM. Engineering C4 photosynthesis into C3 chassis in the
synthetic biology age. The Plant Journal. 2016:**87**(1):51–65. https://doi.org/10.1111/TPJ.13155

Shimizu H, Sato K, Berberich T, Miyazaki A, Ozaki R, Imai R, and Kusano T. LIP19, a basic region leucine
zipper protein, is a Fos-like molecular switch in the cold signaling of rice plants. Plant Cell Physiol.
2005:**46**(10):1623–1634. https://doi.org/10.1093/PCP/PCI178

Shlyueva D, Stampfel G, and Stark A. Transcriptional enhancers: from properties to genome-wide
predictions. Nat Rev Genet. 2014:**15**(4):272–286. https://doi.org/10.1038/NRG3682

Smale ST and Kadonaga JT. The RNA polymerase II core promoter. Annu Rev Biochem. 2003:**72**(Volume 72,
2003):449–479. https://doi.org/10.1146/ANNUREV.BIOCHEM.72.121801.161520/CITE/REFWORKS

Sparks EE, Drapek C, Gaudinier A, Li S, Ansariola M, Shen N, Hennacy JH, Zhang J, Turco G, Petricka JJ, et
al. Establishment of Expression in the SHORTROOT-SCARECROW Transcriptional Cascade through
Opposing Activities of Both Activators and Repressors. Dev Cell. 2016:**39**(5):585–596.
https://doi.org/10.1016/j.devcel.2016.09.031

Spitz F and Furlong EEM. Transcription factors: from enhancer binding to developmental control. Nat Rev
Genet. 2012:**13**(9):613–626. https://doi.org/10.1038/NRG3207

Stergachis AB, Haugen E, Shafer A, Fu W, Vernot B, Reynolds A, Raubitschek A, Ziegler S, LeProust EM,
Akey JM, et al. Exonic transcription factor binding directs codon choice and impacts protein evolution.
Science. 2013:**342**(6164):1367. https://doi.org/10.1126/SCIENCE.1243490

Studer A, Zhao Q, Ross-Ibarra J, and Doebley J. Identification of a functional transposon insertion in the
maize domestication gene tb1. Nat Genet. 2011:**43**(11):1160. https://doi.org/10.1038/NG.942

Swift J, Luginbuehl LH, Schreier TB, Donald RM, Lee TA, Nery JR, Ecker JR, and Hibberd JM. Single nuclei
sequencing reveals C4 photosynthesis is based on rewiring of ancestral cell identity networks. bioRxiv.
2023:2023.10.26.562893. https://doi.org/10.1101/2023.10.26.562893

Tezuka D, Kawamata A, Kato H, Saburi W, Mori H, and Imai R. The rice ethylene response factor OsERF83
positively regulates disease resistance to Magnaporthe oryzae. Plant Physiol Biochem. 2019:**135**:263–
271. https://doi.org/10.1016/J.PLAPHY.2018.12.017

Wang L, Wan MC, Liao RY, Xu J, Xu ZG, Xue HC, Mai YX, and Wang JW. The maturation and aging
trajectory of Marchantia polymorpha at single-cell resolution. Dev Cell. 2023:**58**(15):1429-1444.e6.
https://doi.org/10.1016/J.DEVCEL.2023.05.014

Wang P, Khoshravesh R, Karki S, Tapia R, Balahadia CP, Bandyopadhyay A, Quick WP, Furbank R, Sage TL,
and Langdale JA. Re-creation of a Key Step in the Evolutionary Switch from C_3 to C_4 Leaf Anatomy.
Current Biology. 2017a:**27**(21):3278-3287.e6. https://doi.org/10.1016/j.cub.2017.09.040

Wang P, Khoshravesh R, Karki S, Tapia R, Balahadia CPP, Bandyopadhyay A, Quick WPP, Furbank R, Sage
TLTL, and Langdale JAJA. Re-creation of a Key Step in the Evolutionary Switch from C_3 to C_4 Leaf
Anatomy. Current Biology. 2017b:**27**(21):3278-3287.e6. https://doi.org/10.1016/j.cub.2017.09.040

Wang Y, Huan Q, Li K, and Qian W. Single-cell transcriptome atlas of the leaf and root of rice seedlings.
Journal of Genetics and Genomics. 2021:**48**(10):881–898. https://doi.org/10.1016/J.JGG.2021.06.001

Wiludda C, Schulze S, Gowik U, Engelmann S, Koczor M, Streubel M, Bauwe H, and Westhoff P. Regulation
of the Photorespiratory GLDPA Gene in C4 Flaveria: An Intricate Interplay of Transcriptional and
Posttranscriptional Processes. Plant Cell. 2012:**24**(1):137. https://doi.org/10.1105/TPC.111.093872

Xiong H, Hua L, Reyna-Llorens I, Shi Y, Chen KM, Smirnoff N, Kromdijk J, and Hibberd JM. Photosynthesis-
independent production of reactive oxygen species in the rice bundle sheath during high light is
mediated by NADPH oxidase. Proc Natl Acad Sci U S A. 2021:**118**(25):e2022702118.
https://doi.org/10.1073/PNAS.2022702118/SUPPL_FILE/PNAS.2022702118.SD04.XLSX

Yaish MW, El-Kereamy A, Zhu T, Beatty PH, Good AG, Bi YM, and Rothstein SJ. The APETALA-2-like
transcription factor OsAP2-39 controls key interactions between abscisic acid and gibberellin in rice.
PLoS Genet. 2010:**6**(9). https://doi.org/10.1371/JOURNAL.PGEN.1001098

Yamamoto Y, Ichida H, Abe T, Suzuki Y, Sugano S, and Obokata J. Differentiation of core promoter
architecture between plants and mammals revealed by LDSS analysis. Nucleic Acids Res.
2007a:**35**(18):6219. https://doi.org/10.1093/NAR/GKM685

Yamamoto YY, Ichida H, Matsui M, Obokata J, Sakurai T, Satou M, Seki M, Shinozaki K, and Abe T.
Identification of plant promoter constituents by analysis of local distribution of short sequences. BMC
Genomics. 2007b:**8**(1):1–23. https://doi.org/10.1186/1471-2164-8-67/FIGURES/9

Yan W, Chen D, Schumacher J, Durantini D, Engelhorn J, Chen M, Carles CC, and Kaufmann K. Dynamic
control of enhancer activity drives stage-specific gene expression during flower morphogenesis. Nat
Commun. 2019:**10**(1). https://doi.org/10.1038/S41467-019-09513-2

Zhang C, Ding Z, Wu K, Yang L, Li Y, Yang Z, Shi S, Liu X, Zhao S, Yang Z, et al. Suppression of Jasmonic Acid-
Mediated Defense by Viral-Inducible MicroRNA319 Facilitates Virus Infection in Rice. Mol Plant.
2016:**9**(9):1302–1314. https://doi.org/10.1016/J.MOLP.2016.06.014

Zhang C, Zhang J, Liu H, Qu X, Wang J, He Q, Zou J, Yang K, and Le J. Transcriptomic analysis reveals the
role of FOUR LIPS in response to salt stress in rice. Plant Mol Biol. 2022:**110**(1–2):37–52.
https://doi.org/10.1007/S11103-022-01282-9/FIGURES/7

Zhang H, Zhang J, Quan R, Pan X, Wan L, and Huang R. EAR motif mutation of rice OsERF3 alters the
regulation of ethylene biosynthesis and drought tolerance. Planta. 2013:**237**(6):1443–1451.
https://doi.org/10.1007/S00425-013-1852-X

Zhang X, Bao Y, Shan D, Wang Z, Song X, Wang Z, Wang J, He L, Wu L, Zhang Z, et al. Magnaporthe oryzae
Induces the Expression of a MicroRNA to Suppress the Immune Response in Rice. Plant Physiol.
2018:**177**(1):352. https://doi.org/10.1104/PP.17.01665

Zhao H, Yang M, Bishop J, Teng Y, Cao Y, Beall BD, Li S, Liu T, Fang Q, Fang C, et al. Identification and
functional validation of super-enhancers in Arabidopsis thaliana. Proc Natl Acad Sci U S A.
2022:**119**(48):e2215328119.
https://doi.org/10.1073/PNAS.2215328119/SUPPL_FILE/PNAS.2215328119.SD02.XLSX

Figure 1

Figure 2

A

-980 +42

-829 +42

-980 -251 min35S

-980 -829 -251 +42

B

60h

166h

166 h

43h

C

nmol MU min^{-1}(mg protein)$^{-1}$

	Median	n
a	0.142	24
c	0.016	25
b	0.038	24
b	0.079	29

Figure 4

Figure 5

A

BS enhancer

-980 -938 -923 -904 -873 -853

G2-like MYBR IDD bZIP

3 x BS enhancer

B

48h

4h

C

	Median	n
a	0.034	24
b	0.837	22

nmol MU min^{-1}(mg protein)$^{-1}$

0.0001 0.001 0.01 0.1 1 10 100

Figure 6

A

-2571

BS enhancer

-980 -829

Y-patch

-62 -42 +42

B

GLK2 MYBS1 IDD3/4/6/10 bZIP3/4/9/10

Pol II

BS enhancer

Y Patch

C

BS enhancer + Activators + M silencer = SiR expression