1

Main Manuscript for

Synchronicity transitions determine coupling fluctuations in a model of coupled oscillators with plasticity

Satoshi Kuroki [a*] and Kenji Mizuseki [a]

[a] Department of Physiology, Osaka Metropolitan University Graduate School of Medicine, Asahimachi 1-4-3, Abeno-ku, Osaka, 545-8585, Japan, +81-6-6645-3717

* Satoshi Kuroki, Department of Physiology, Osaka Metropolitan University Graduate School of Medicine, Asahimachi 1-4-3, Abeno-ku, Osaka, 545-8585, Japan, +81-6-6645-3717, skuroki@omu.ac.jp

Email: skuroki@omu.ac.jp

Author Contributions: S.K.: Conceptualization, Investigation, Visualization, Writing - original draft, Writing – review and editing, Funding acquisition; K.M.: Supervision, Writing – review and editing, Funding acquisition

Competing Interest Statement: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Classification: Systems Neuroscience

Keywords: Network reorganization, Plasticity, EI balance, Synchronization

This PDF file includes:

Main Text
Figures 1 to 5

1
Abstract

Sleep and rest are crucial to knowledge abstraction and creativity. During these periods,
synapses undergo plastic modification, while neuronal activities are collectively synchronized,
accompanied by large shifts in the excitation-inhibition (EI) balance. These phenomena differ
from the learning processes that occur during task engagement. However, the detailed
mechanism by which synchronized neuronal activity reshapes neural circuits through plasticity is
not yet fully understood. The Kuramoto model is used to study the collective synchronization of
oscillators, including neurons. We previously proposed the EI-Kuramoto model, in which an EI
balance was implemented in the Kuramoto model. The model alters its synchronicity based on
the EI balance of the interaction strength. In this study, we further developed the EI-Kuramoto
model by implementing Hebbian and homeostatic plasticity, leading to a plastic EI-Kuramoto (pEI-
Kuramoto) model. Models with robust inhibition consistently showed desynchronization and
stable coupling. Contrastingly, weaker inhibition produced bistable synchronization, in which
couplings with intermediate strengths fluctuated, whereas the strongest couplings remained
stable. Our findings suggest that the dynamic interplay between network activity and plasticity
explains why stronger synapses in the brain exhibit greater stability than their weaker
counterparts. Moreover, this mechanism facilitated network reorganization during sleep and rest.

Significance Statement

The reorganization of neuronal circuits during sleep and rest is essential for knowledge
abstraction and creativity. These states involve synaptic plasticity and collective neuronal
synchronization, accompanied by significant shifts in the excitation-inhibition (EI) balance.
However, the precise mechanism through which synchronized activity reshapes neural circuits via
plasticity remains unclear. We developed a novel computational model, the pEI-Kuramoto model,
which incorporates both coupling plasticity and EI balance. Our model demonstrates that robust
inhibition leads to desynchronization and stable coupling. In contrast, weaker inhibition induces
bistable synchronization, in which the intermediate-strength couplings fluctuate, whereas the
strongest couplings remain stable. These findings provide a fundamental mechanism for network
reorganization during sleep and at rest.

30
Main Text

32
Introduction
Empirical evidence suggests that brain activity during sleep and rest promotes memory
consolidation, abstraction of knowledge, and the creation of new ideas (1, 2) through the
reorganization of neural circuits (3, 4). This reorganization should differ from learning while
performing a task. Indeed, synaptic plasticity during sleep and rest improves task performance (5,
6).
Neuronal synchronization occurs in two distinct states. Collective neuronal firing is
desynchronized during active wakefulness, including task engagement, movement, and
attentional states, as well as during rapid eye movement (REM) sleep, which are referred to as
brain-activated states. Contrastingly, neuronal firing synchronizes during non-REM sleep and
quiet wakefulness, which are referred to as brain-deactivated states (7, 8). The transition between
these states is associated with shifts in the excitation-inhibition (EI) balance. Firing rates of
inhibitory neurons and synaptic conductance of inhibitory synapses are significantly elevated
during brain-activated states, while those of excitatory synapses show comparatively smaller
changes (9–12). These observations suggest that the extent to which collective neuronal
synchronization is accompanied by EI balance and synaptic plasticity during brain-deactivated
states can influence network reorganization. However, the detailed mechanisms by which

neuronal synchronization in the brain-deactivated state reshapes the neural network structure
remain unclear.
The collective synchronization of oscillators, including neurons, has been studied using a model
of coupled oscillators. Among these, the Kuramoto model is considered to be the simplest
framework (13, 14), and numerous variants have been proposed. One variant, the EI-Kuramoto
model, captures the EI balance by dividing oscillators into excitatory and inhibitory groups and
assigning four interaction types between them (15, 16). We have previously shown that tuning
these four strengths toggles the network between synchronized, desynchronized, and bistable
states, potentially reflecting a neuronal synchronization mechanism (16). That study, however,
kept all coupling weight—which represent pair-specific interaction strengths—fixed and identical,
and therefore did not address how the EI balance and its associated dynamical states influence
plastic changes in the coupling weights.
In this study, we focused on Hebbian and homeostatic plasticities. Hebbian plasticity, based on
the principle "fire together, wire together," strengthens synapses between neurons that fire in
close temporal proximity (17). However, homeostatic plasticity regulates neuronal excitability and
synaptic strength to maintain network stability, thereby preventing excessive excitation or
quiescence (18). The interplay between these plasticity mechanisms enables the emergence of
diverse neuronal functions and firing patterns (19–22). Several Kuramoto model studies have
implemented plasticity, enabling coupling weights to adapt based on the oscillator activity to
examine their impact on system synchronization (23–33).
In this study, we refined the EI-Kuramoto model by integrating Hebbian and homeostatic plasticity
into the coupling weights between excitatory oscillator unit pairs (referred to as plastic EI, pEI-
Kuramoto model). We observed that the network maintained robustness and minimal fluctuations
when inhibition was strong, corresponding to the desynchronized state. Conversely, with weaker
inhibition, corresponding to the bistable state, couplings with moderate weights tended to
fluctuate, whereas couplings with strong weights remained stable. These findings on the network
dynamics of the pEI-Kuramoto model may provide important insights into the mechanisms of
neural networks in a brain-deactivated state.
78
Results
Establishment of the pEI-Kuramoto model The pEI-Kuramoto model comprised two oscillator
groups: excitatory units, referred to as excUnits, and inhibitory units, referred to as inhUnits. These
specific terms clarified that units represented excitatory and inhibitory factors in this model, not
actual neurons or synapses. Four interaction strengths were assigned between the groups, with
only excitatory-excitatory couplings J_{ee} being distributed and plastic. The other constant interaction
strengths were represented as K_{ei}, K_{ie}, and K_{ii} (Fig 1A; see Materials and Methods). The
interactions, which are modeled as zero phase lag sine functions, are purely attractive within
excUnits and purely repulsive within inhUnits. Consequently, the terms "excitatory" and "inhibitory"
are used synonymously with "attractive" and "repulsive," respectively.
First, we investigated how the distribution of pair-specific excitatory-excitatory coupling weights J_{ee}
without plasticity, affected the dynamics of the pEI-Kuramoto model. In our previous study, we
showed that the Kuramoto model with EI balance (EI-Kuramoto model) exhibited three distinct
dynamic states—synchronized, bistable, and desynchronized—depending on the balance between
the constant values of the interaction strengths (16). In this study, J_{ee} followed a Gaussian
distribution with a mean of μ_{Jee} and a standard deviation of σ_{Jee}, where σ_{Jee} was set to 0, 1/10, and
1/2 of μ_{Jee} (Fig. 1B). The collective synchronization was quantified using the order parameter R,
which represents the magnitude of the average vector between the oscillator units. Regardless of
the standard deviation, all systems with a distributed J_{ee} also exhibited synchronized (high R),
bistable (alternating high and low R), and desynchronized (low R) dynamics as observed with
constant ($\sigma_{Jee} = 0$) values (Fig. 1C).
Next, we incorporated plasticity into J_{ee} by applying Hebbian potentiation and homeostatic
regularization plasticity rules (see Materials and Methods). Among the various methods available
for implementing Hebbian plasticity in the Kuramoto model, we initially selected the simplest
method: the cosine (or sine) function (SI Appendix, Fig. S1A; see Materials and Methods) (23–25,

28, 32, 33). In this process, when a pair of excUnits exhibits closer phases, their J_{ee} coupling weight
is potentiated. Simultaneously, the J_{ee} distribution was rescaled to maintain the initial mean and
standard deviation, thereby preventing overpotentiation (homeostatic regularization plasticity; see
Materials and Methods). The system dynamics were evaluated using the parameter set ($\mu_{Jee} = 1$,
$\sigma_{Jee} = 0.1$, $K_{ei} = 3$, $K_{ie} = 3$, $K_{ii} = 3$) at which the system exhibited a desynchronized state in the
absence of J_{ee} plasticity. Under the cosine rule, unit pairs with smaller phase differences acquire a
larger J_{ee} weight. This rule causes the J_{ee} distribution, which was initially Gaussian, to become
bimodal (Fig. S1B). Plasticity also influences the distribution of oscillator units, making the
distributions of both phase and phase difference become bimodal (SI Appendix, Fig. S1C),
consistent with previous studies that incorporated the Hebbian rule into the Kuramoto model (23,
28). This bimodal distribution of the coupling weight deviates from the long-tailed, skewed
distributions commonly observed in biological synaptic weights (12, 34, 35).
To address this discrepancy, we introduced an exponential-based Hebbian plasticity rule to adjust
the decay rate (Sparseness γ) of the gain for the phase difference (Fig. 1D). We applied this rule
to a range of parameter sets and evaluated the resulting J_{ee} distribution using the skewness,
kurtosis, and dip values, which quantifies the distribution bimodality (36). Although these
distributions initially followed a Gaussian pattern, they transitioned into various bimodal or skewed
forms (Fig. 1E and F). We also quantified synchronization of the excUNits using the mean and
standard deviation of the order parameter R_{exc} through time, denoted as $E[R_{exc}]_t$ and $S[R_{exc}]_t$,
respectively. Plasticity in J_{ee} affects synchronization (Fig. 1G), resulting in uniform, bimodal, or
unimodal phase and phase-difference distributions depending on the parameters (Fig. S1C).
Moreover, the boundary of the statistical features of the J_{ee} distribution (Fig 1E) appears to coincide
with the synchronization boundary (Fig 1G). For further analysis, we chose a parameter set that
resulted in a long-tailed skewed J_{ee} distribution (Fig. 1F, top, magenta circle), which more closely
reflects the synaptic weight distributions reported in the brain (12, 34, 35).
129
Coupling fluctuation increases in bistable synchronization state We then investigated how the
dynamic state affects coupling weight, focusing on the role of inhibition, which varies significantly
across different brain states (9–12), resulting in changes in EI balance (7, 37). To investigate this,
we altered the EI balance of the pEI-Kuramoto model by manipulating the interaction strength of
the inhUnits. We integrated K_{ie} and K_{ii} into a single parameter, K_i, to simplify the simulation following
the approach used in our previous study (16). We then examined how the J_{ee} fluctuations were
associated with the K_i values, quantified as the standard deviation of J_{ee} over time ($S[J_{ee}]_t$; see
Materials and Methods). By analyzing the time course of R_{exc}, we determined the dynamic state of
the system, as previously described (16).
When $K_i < 0.4$, both the average $S[J_{ee}]_t$ across the unit pairs ($E\{S[J_{ee}]_t\}_n$) and the standard deviation
of $S[J_{ee}]_t$ between the pairs ($S\{S[J_{ee}]_t\}_n$) were almost zero (Fig. 2A and B), indicating no plastic
changes in J_{ee}. In this K_i range, the system was in a synchronized state characterized by high
$E[R_{exc}]_t$ and low $S[R_{exc}]_t$ (Fig. 2C and D). The coupling weights between individual pairs exhibited
no fluctuation over time (Fig. 2E, K_i: 0.1; SI Appendix, Movie S1), and the clustered couplings
remained unchanged (Fig. 2F, K_i: 0.1).
As K_i increased, $E\{S[J_{ee}]_t\}_n$ also increased, peaking at $K_i = 0.7$, then decreased, and eventually
stabilized around $K_i = 1.5$ (Fig. 2A). Meanwhile, the $S\{S[J_{ee}]_t\}_n$ continued to increase (Fig. 2B). For
K_i values above 1.5, the system entered a desynchronized state characterized by low $E[R_{exc}]_t$ and
$S[R_{exc}]_t$ values (Fig. 2C and D). Higher or lower couplings remained stable over time, whereas those
of intermediate strength fluctuated (Fig. 2E, K_i: 1.5; SI Appendix, Movie S2). Most clustered
couplings at a given time point disappeared over time, leaving only a small subset that remained
clustered (Fig. 2F, K_i: 1.5).
In the intermediate range ($0.4 < K_i < 1.5$), the system exhibits a bistable state with high $S[R_{exc}]_t$ (Fig.
2D). Notably, the $S[J_{ee}]_t$ was significantly higher in this bistable state (Fig. 2A). Similar to the
desynchronized state ($K_i > 1.5$), the coupling weights of the higher or lower unit pairs were stable,
whereas the intermediate-weighted couplings fluctuated more. However, the number of these
intermediate fluctuating couplings increased (Fig. 2E and F, K_i: 0.8 and 1.0; SI Appendix, Movies

S3 and S4). Thus, transitions in the dynamic state are accompanied by changes in the coupling
weight fluctuations. In particular, the bistable state tended to exhibit larger fluctuations.
Note that Hebbian plasticity affects the dynamic state of the pEI-Kuramoto model (SI Appendix,
Fig. S2A), slightly shifting the bifurcation points between the synchronized and bistable states and
the bistable and desynchronized states toward larger K_i values. Owing to this effect, a small shift
from the eigenvalue-based classification of dynamic states by Kuroki & Muzuseki (2025) was
expected. However, the overall trend of the state transition from synchronized to bistable and then
to desynchronized remained consistent across different Hebbian learning rates. This behavior is
consistent with previous reports on Kuramoto models incorporating cosine-shaped Hebbian
plasticity (28, 33). Contrastingly, σ_{Jee} did not affect the dynamic states (SI Appendix, Fig. S2B),
which is consistent with the results shown in Fig. 1C.

Strong-weighted couplings are stable while intermediate couplings fluctuated We further
investigated coupling weight fluctuations under conditions of slightly weakened inhibition from the
desynchronized state, assuming intermediate brain states between alert brain-activated and -
deactivated states. We analyzed the individual J_{ee} coupling weights at selected points of K_i (K_i =
0.8, 1.0, and 1.5, red crosses in Fig. 2A–D, except K_i = 0.1). Histograms of $E[J_{ee}]_t$ (mean of J_{ee} over
time for each pair of units) and $S[J_{ee}]_t$ also tended to shift toward higher values as K_i decreased
(Fig. 3A and B). A scatter plot of $E[J_{ee}]_t$ and $S[J_{ee}]_t$ formed a characteristic "bow-shaped" distribution
(Fig. 3C). In this two-dimensional distribution, data points near the minimum and maximum $E[J_{ee}]_t$
values had lower $S[J_{ee}]_t$ values, whereas intermediate $E[J_{ee}]_t$ showed a higher $S[J_{ee}]_t$. This pattern
was consistent across all distributions for different K_i values. A similar phenomenon, the
stabilization of strong couplings (e.g., synapses with large mushroom-shaped spines), has been
observed in brain synapses (38–40).
Furthermore, the $S[J_{ee}]_t$ values tended to increase at intermediate $E[J_{ee}]_t$ values with decreasing K_i
(Fig. 3C), suggesting that K_i can regulate the fluctuations in individual J_{ee} plastic couplings. At high
K_i values in desynchronized states, the couplings stabilize, whereas at low K_i values in bistable
states, the couplings of intermediate weights become unstable. The high J_{ee} coupling remained
stable with small fluctuations, even when K_i was low. These dynamics appear to be a mechanism
that preserves the coupling of particularly strong pairs and resets pairs of moderate weight,
facilitating the reorganization of the network structure (Fig. 2E and 2F).

Factors determining coupling weight properties To determine which parameters determine the
individual J_{ee} coupling properties shown in Fig. 3C, we explored the parameters involved in J_{ee}
fluctuations. We analyzed the relationship between the individual J_{ee} coupling properties ($E[J_{ee}]_t$ vs.
$S[J_{ee}]_t$ distribution shown in Fig. 3C) and the candidate parameters (Fig. 4). The couplings were
roughly categorized based on whether $E[J_{ee}]_t$ exceeded 0.018 (high $E[J_{ee}]_t$ group) or not (low $E[J_{ee}]_t$
group) for statistical comparison because the trend reversed around this point.
For the differences in natural frequency (ω_{exc}) of excUnit pairs, smaller natural frequency
differences were associated with smaller $S[J_{ee}]_t$ (Fig. 4A and D). Pairs with larger natural frequency
differences tended to be in the low-$E[J_{ee}]_t$ group, and $S[J_{ee}]_t$ appeared to increase with increasing
natural frequency differences. However, most pairs in the high $E[J_{ee}]_t$ group had relatively small
natural frequency differences. The initial phase differences between the pairs were also significant
(Fig. 4B and E). In the low $E[J_{ee}]_t$ group, the initial phase differences were uniform, whereas in the
high $E[J_{ee}]_t$ group, several pairs exhibited small initial phase differences. Additionally, we examined
the effect of the initial coupling weight of J_{ee}, which showed no noticeable bias (Fig. 4C) and
followed a normal distribution (Fig. 4F). Considering these results, pairs with small differences in
natural frequency and initial phase positions maintained more stable and larger J_{ee} weights (Fig.
4G and H). Contrastingly, the initial J_{ee} weights were not significant.

Theoretical analysis of quasi-steady coupling weight distribution We investigated the
theoretical mechanism underlying the numerical simulation results for coupling weights in a quasi-
steady state, where values exhibit minimal fluctuation (SI Appendix, Text S1). Although a true static
steady weight distribution does not exist in the desynchronized or bistable states due to the non-

constant phases of each oscillator unit, we can consider a quasi-steady weight distribution within
the desynchronized condition, where the phase and weight distributions evolve on a sufficiently
slow timescale. Under this assumption, the coupling weights do not change as a result of the plastic
modulation of Eqs. M11 and M14, which lead to the following relationship (SI Appendix, Eq. S3).

215
$$J_{ee,kj} = \hat{J}_{ee,kj} + c \exp\left(-\gamma|\Delta\theta_{kj}|\right) \tag{1}.$$

216
$$c = \frac{\alpha(1-\beta)}{\beta}$$

The first term, $\hat{J}_{ee,kj}$, represents the value of $J_{ee,kj}$ within the normalized distribution of J_{ee} by mean
$\hat{\mu}_{Jee}$ and variance $\hat{\sigma}_{Jee}$. This second term is the Hebbian potentiation component, scaled by a ratio
between Hebbian and homeostatic plasticity learning rates c. This equation indicates that the
weight $J_{ee,kj}$ depends on the distributions of both the overall J_{ee} distribution and the phase
differences between unit k and j $\Delta\theta_{kj}$. As this update rule is iterated, the influence of the second
$\Delta\theta$-dependent Hebbian term is sequentially incorporated into the J_{ee} distribution. Furthermore, the
distribution of $\Delta\theta$ is dependent on the distribution of θ (SI Appendix, Text S1). Therefore, when
the weight dynamics are sufficiently slow, the J_{ee} distribution is determined by the phase
distribution. Indeed, the characteristics of the J_{ee} distribution appear to be correlated with the
synchronization state of the system, which reflects the shape of the phase distribution (Fig 1E,
1F, SI Appendix, Fig S1). Specifically, a uniform phase distribution (desynchronized state)
corresponds to a left-skewed, long-tailed J_{ee} distribution. A bimodal phase distribution
corresponds to a bimodal J_{ee} distribution. A unimodal phase distribution (synchronized state)
corresponds to a right-skewed J_{ee} distribution.
Next, we considered the change in the phase difference for a pair of excUnits $\Delta\theta_{kj}$. Assuming $J_{ee,kj}$
and $\Delta\theta_{kj}$ are constant, the following relationship can be derived from Eqs. M9 and S3 (SI
Appendix, Eq. S18):

234
$$-\Delta\omega + F_{inh} - F_{exc}^{(-kj)} = -2\left\{\hat{J}_{ee,kj} + c \exp\left(-\gamma|\Delta\theta_{kj}|\right)\right\}\sin\left(\Delta\theta_{kj}\right) \tag{2}.$$

$\Delta\omega$ is natural frequency difference between unit k and j, F_{inh} is interaction strength from the
inhUnits, and $F_{exc}^{(-kj)}$ is interaction strength from the excUnits except unit k and j. The right-hand
side of this equation represents the interaction strength between the unit k and j, while the left-
hand side represents the net forces of the other factors. A stable fixed point for $\Delta\theta_{kj}$ exists where
the functions for the left and right-hand sides intersect (SI Appendix, Fig. S3). On the right-hand
side, the attractive force is driven by both the homeostatic set-point, $\hat{J}_{ee,kj}$, and the Hebbian
potentiation term. Near $\Delta\theta\approx0$, the Hebbian effect is strong, leading to a large attractive force. As
the phase difference $|\Delta\theta|$ increases, this Hebbian potentiation weakens and the coupling weight
relative position within the J_{ee} distribution falls to a lower, causing reduction of the overall
attractive force. The left-hand side represents the net external drive on the pair. This drive is
composed of the natural frequency difference $\Delta\omega$ and the R_{exc} and R_{inh}-dependent term $F_{inh} -$
$F_{exc}^{(-kj)}$, whose direction can be either attractive or repulsive depending on the phase configuration
of the system. (SI Appendix, Text S1). While the influence of F_{inh} and $F_{exc}^{(-kj)}$ is limited in a
desynchronized state where R_{exc} and R_{inh} are near zero, their impact becomes significant in the
bistable state, where these order parameters fluctuate periodically. Sufficient fluctuation in these
forces can overcome the attractive force represented by the right-hand side, causing the stable
fixed point to either disappear or be significantly displaced. This leads to large fluctuations in the
phase difference between the unit pair.
253
Cyclic modulation of inhibition Finally, we varied K_i to mimic the sleep-wake cycle (7). J_{ee}
fluctuated (Fig. 5A, SI Appendix, Movie S5) according to the modulation of K_i (Fig. 5B, top panel).
R_{exc} alternated between the desynchronized and bistable states (Fig. 5B, middle), and $E\{S[J_{ee}]_t\}_n$
changed accordingly (Fig. 5B, bottom). When K_i was high, the system was in a desynchronized
state, and $E\{S[J_{ee}]_t\}_n$ was small. Conversely, when K_i was low, the system exhibits a bistable state
and $E\{S[J_{ee}]_t\}_n$ increases. These dynamics can be considered as modeling how the circuit structure

is stabilized during the brain-activated state and destabilized during the brain-deactivated state,
thereby promoting circuit reorganization.
262
Discussion
In this study, we extended the previously developed EI-Kuramoto model by incorporating Hebbian
and homeostatic plasticity to establish the pEI-Kuramoto model and analyze its dynamics. We
propose that this model represents a continuous transition between the stabilization and
reorganization of network structures, modulated by the desynchronized-bistable state cycle
through inhibition (see Figs 2–5). This study introduces a possible mechanism for preserving
strong couplings and reorganizing network structures during brain-deactivated states, as viewed
through the dynamics of collective oscillators.
271
Potential mechanism for network reorganization Preserving robust couplings while
randomizing weaker ones simplifies the network by rendering them sparse. The choice of
plasticity parameters, which yield a long-tailed coupling weight distribution similar to that
observed in the brain (Fig. 1E and F) and mirror brain dynamics during the brain-deactivated
state (9–12), appears to facilitate network sparsification. The selective preservation of strong
synaptic connections is also frequently observed in the brain (38–40) and is understood through
molecular mechanisms, including the dynamics of NMDA receptors (5, 41) and actin filaments
(42). Our findings introduce a potential mechanism for sustaining strong coupling when viewed
from another aspect of the dynamic system of collective oscillators. In addition, at the system
level, only a few populations of neuronal activity are stable, while others drift even in familiar
environments (43–45). This process aligns with the pruning method in artificial neural networks,
which facilitates knowledge generalization (46, 47). This pruning-like mechanism is particularly
relevant when synaptic downregulation is dominant during brain-deactivated states (5, 41), as
fluctuations in couplings can lead to the weakening and initialization of the couplings.
Furthermore, fluctuations in network connectivity can promote both orderly and random network
reorganization. In our model, bistable states under low inhibition conditions promoted fluctuating
coupling weights between excUnits. As the inhibition decreases, the process transitions from
structured coupling weight changes governed by the Hebbian rule to nearly random changes
driven by the large phase fluctuations of the units (see Supplementary Videos 2–5). It has also
been proposed that the brain enhances neuronal circuit reorganization during brain-deactivated
state (3, 4). Combined with unknown mechanisms to evaluate and stabilize reorganized circuits
(48) such as dopaminergic signaling during non-REM sleep (49), we may approach how new
creative ideas or inexperienced problem-solving methods are generated.
295
Dynamics of pEI-Kuramoto model From the stability analysis of the phase difference between
excUnit pair (Eq. 2), the characteristic bow-shaped relationship between the mean ($E[J_{ee}]_t$) and
standard deviation ($S[J_{ee}]_t$) of coupling weights (Fig. 3C) can be explained. Pairs of strong
couplings maintain a small and stable phase difference ($\Delta\theta \approx 0$). This ensures that the Hebbian
potentiation term, $c \exp(-\gamma|\Delta\theta_{kj}|)$, and $\hat{J}_{ee,kj}$, associated with relative position in the J_{ee}
distribution, remains consistently high. Consequently, these synapses sustain a high $E[J_{ee}]_t$ with
very small fluctuations, resulting in a low $S[J_{ee}]_t$. These pairs form the top-left end of the bow in
Fig 3C. On the other side in the plot, the phases of unit pairs with very weak couplings evolve
almost independently. The Hebbian potentiation effect is negligible, and their synaptic weights
remain at a low baseline value determined by homeostatic pressures. As a result, $S[J_{ee}]_t$ at a
particular time window depends primarily on the natural frequency difference $\Delta\omega$. The largest
$S[J_{ee}]_t$ is found in pairs with intermediate coupling strength. For these couplings, the
synchronizing force is in close competition with the desynchronizing drives from their natural
frequency difference $\Delta\omega$. The system is critically poised: small fluctuations in $\Delta\theta$ can tip the
balance between Hebbian potentiation and homeostatic depression, causing the direction of
weight change to reverse frequently. This instability, where the fixed point for $\Delta\theta$ may appear and
disappear intermittently, drives large fluctuations in the coupling weight J_{ee}. This mechanism
produces a high variance at intermediate mean weights, forming the arch of the bow. From a

dynamics perspective, this mechanism may explain a characteristic feature of biological
synapses: larger synapses are more stable than smaller ones (38–40).
The fluctuation of J_{ee} when K_i is low can be attributed to the system dynamics becoming bistable,
where the system alternates between synchronization and desynchronization. It causes
fluctuations in the phase distribution, which is the primary factor driving the changes in J_{ee}.
Furthermore, in this state, the periodic fluctuation of the order parameters R_{exc}, R_{inh} constant
changes in the interaction term, $F_{inh} - F_{exc}^{(-kj)}$ (Eq. 2). This dynamic instability also strongly
perturbs the phase differences Δθ of moderately coupled pairs more.
The Eq. 2 suggests that unit pairs with small initial frequency Δω and phase differences Δθ are
more robust and more likely to return to a stable, synchronized state even after large
perturbations. This provides an explanation for the results shown in Figure 4. This finding
resonates with observations in the brain, where neurons that exhibit synchronized firing prior to
learning (analogous to a small Δθ) are more likely to be grouped into the same assembly during
learning (50–52). However, it remains unclear whether these neurons also share similar intrinsic
firing frequencies, which would correspond to a small Δω in our model. Although the effective
interaction function in our simulation setting (Fig. 1F, magenta circle), designed to approximate
biological synaptic distributions (12, 34, 35), deviates slightly from a pure sine wave by providing
a stronger restoring force near the stable point (Fig. S3), the enhancement of interaction strength
near Δθ=0 by the Hebbian effect is limited. Thus, it appears that this property stems primarily
from the underlying sinusoidal nature of the coupling, rather than from the modifications
introduced by plasticity.
The quasi-steady J_{ee} distribution is determined by the phase difference distribution (Fig 1E, 1F,
and SI Appendix, Fig S1), as it is a transformation of the Δθ distribution via the Hebbian function
(Eq. 1). In this study, the exponential rule $c \exp\left(-\gamma \left| \Delta\theta_{kj} \right|\right)$ transforms the uniform phase
difference distribution into a hyperbolic-like weight distribution, with a probability density
proportional to $1/J_{ee}$ (Appendix, Text S1). This framework suggests that, by modifying the
Hebbian function, it is possible to generate the J_{ee} distribution into an arbitrary form, such as a
log-normal distribution that observed in the brain (12, 34, 35).

Limitations While the Kuramoto model provides foundational insights, it is considerably simpler
than actual neuronal behavior and has certain limitations. First, our model simplifies neuronal
interactions to mere attractive and repulsive forces, overlooking the pulse-like nature, interaction
delay, and intricate coupling relationships of real neurons (53). Second, the model
underrepresents the complexity of neuroplasticity, such as the influence of neurotransmitters and
internal conditions (54), plasticity in inhibitory neurons (55, 56), and short-term plasticity including
adaptation (57). Lastly, the model does not distinguish between brain states, such as full
wakefulness, non-REM sleep, and REM sleep, which are believed to play unique roles in learning
and neural reorganization (3, 4). Addressing these limitations is essential for a more accurate and
comprehensive understanding of the neural dynamics and plasticity.

Future directions Future research can consider multiple directions to expand the system further.
In this study, we observed a behavior in which strong couplings were retained and intermediate
couplings were randomly perturbed at low inhibition. Creativity may require a mechanism for
evaluating and stabilizing randomly reorganized circuits (48, 58). This mechanism is not yet fully
understood; however, dopamine activity during rest (49) may be a key factor. Another important
question is how changes in overall synchrony during sleep affect the interactions between
functionally distinct networks, such as the hippocampus and cerebral cortex. Exploring the
influence of synchrony transitions on interregional plasticity could shed light on how memory
traces originally stored in the hippocampus gradually become integrated or transferred to cortical
areas (59). Moreover, our model did not specifically consider the dimensionality of information,
but understanding how the brain represents and selects multidimensional information is crucial for
cognition (60). We must consider the biases, interactions, and plasticity rules required to
represent and select multidimensional information from the external world. Clarifying these

mechanisms will contribute to a deeper understanding of how neural circuits are reorganized to
support abstraction and creativity.
369
370
371 **Materials and Methods**
372
373 **pEI-Kuramoto model** The original Kuramoto model (13, 14, 61) is expressed as:

$$374 \qquad \frac{d\theta_j(t)}{dt} = \omega_j + \frac{K}{N}\sum_k^N \sin\left(\theta_k(t) - \theta_j(t)\right) \qquad \text{(M1)},$$

375 where $\theta_j(t)$ is the phase of unit j at time t, ω_j is the natural frequency of unit $\dot{j}$ following a
376 probability distribution $g(\omega)$, N is the number of units, and K is the interaction strength among the
377 oscillation units. In the original Kuramoto model, K is constantly and uniformly applied to all unit
378 pairs. The order parameter R and the mean phase ψ are then calculated from the mean vector of
379 the oscillators as:

$$380 \qquad Re^{i\psi} = \frac{1}{N}\sum_j^N e^{i\theta_j} \qquad \text{(M2)},$$

381 where i denotes an imaginary unit. Given Euler's formula and Eq (2),

$$382 \qquad \frac{1}{N}\sum_k^N \sin(\theta_k - \theta_j) = \frac{1}{N}\sum_k^N \text{Im}\left[e^{i\theta_k}e^{-i\theta_j}\right] = \text{Im}\left[Re^{i\psi}e^{-i\theta_j}\right] = R\sin(\psi - \theta_j) \qquad \text{(M3)}.$$

383 Considering, Eq (3), Eq (1) becomes

$$384 \qquad \frac{d\theta_j}{dt} = \omega_j + KR\sin(\psi - \theta_j) \qquad \text{(M4)}.$$

385
In our previous EI-Kuramoto model (16), the oscillation units were categorized into two groups:
excitatory and inhibitory units, referred to as excUnits and inhUnits, respectively. Employing these
terms clarified that the units in the model represented excitatory and inhibitory factors,
distinguishing them from actual neurons or synapses. The interaction strength of the sine wave
functions with zero phase lag is characterized by purely attractive interactions in excUnits and
purely repulsive interactions in inhUnits. Hence, the terms "excitatory" and "inhibitory" are used
interchangeably with "attractive" and "repulsive," respectively. The interaction strength K is
divided into four quantifiable parameters: K_{ee} (attractive strength between excUnits), K_{ii} (repulsive
strength between inhUnits), K_{ei} (attractive strength from excUnits to inhUnits), and K_{ie} (repulsive
strength from inhUnits to excUnits). K_{ee}, K_{ii}, K_{ei}, and K_{ie} are distinct positive real numbers that are
uniformly applicable across all the unit pairs. Consequently, Eq (M4) becomes:

$$397 \qquad \frac{d\theta_{j,exc}}{dt} = \omega_{j,exc} + K_{ee}R_{exc}\sin(\psi_{exc} - \theta_{j,exc}) - K_{ie}R_{inh}\sin(\psi_{inh} - \theta_{j,exc}) \qquad \text{(M5)}$$

398
399 for excUnits, and

$$400 \qquad \frac{d\theta_{j,inh}}{dt} = \omega_{j,inh} + K_{ei}R_{exc}\sin(\psi_{exc} - \theta_{j,inh}) - K_{ii}R_{inh}\sin(\psi_{inh} - \theta_{j,inh}) \qquad \text{(M6)}$$

401
402 for inhibUnits, where exc and inh represent the parameters for excUnits and inhUnits,
403 respectively. The order parameter and mean phase of each excUnit and inhUnit are

$$404 \qquad R_{exc}e^{i\psi_{exc}} = \frac{1}{N_{exc}}\sum_j^{N_{exc}} e^{i\theta_{j,exc}} \qquad \text{(M7)}$$

405 and

$$406 \qquad R_{inh}e^{i\psi_{inh}} = \frac{1}{N_{inh}}\sum_j^{N_{inh}} e^{i\theta_{j,inh}} \qquad \text{(M8)}.$$

In the present study, we developed a pEI-Kuramoto model (Fig. 1A). A pivotal modification of our
previous EI-Kuramoto model is that the constant and uniform interaction strength value K_{ee} is
replaced with the heterogeneous and plastic coupling weight (interaction strength of an individual
unit pair) matrix J_{ee}. The initial distribution of J_{ee} follows the Gaussian distribution
$g(J_{ee}) \sim \mathcal{N}(\mu_{Jee}/N_{exc}, (\sigma_{Jee}/N_{exc})^2)$, where μ_{Jee} and σ_{Jee} are the mean and standard deviation of
the J_{ee} distribution, respectively. These were scaled with the inverse of N_{exc} to adjust for the
system size. The other interaction strengths, K_{ii}, K_{ei}, and K_{ie}, were uniform and consistently
applicable across all units. Eq (M5) becomes

416
$$\frac{d\theta_{j,exc}}{dt} = \omega_{j,exc} + \sum_{k}^{N_{exc}} J_{ee,kj} \sin(\theta_{k,exc} - \theta_{j,exc}) - K_{ie}R_{inh} \sin(\psi_{inh} - \theta_{j,exc}) \tag{M9}.$$

417 $J_{ee,kj}$ is the coupling weight between the excUnits k (pre) and j (post).
418
Given the rotational symmetry, when the mean natural frequencies of excUnits and inhUnits are
equal, $\bar{\omega} = \bar{\omega}_{exc} = \bar{\omega}_{inh}$, the mean natural frequencies can be set to 0 rad/s. This is equivalent to
rotating a coordinate by the mean natural frequency $\bar{\omega}$ (61). For all simulations in this study,
$N_{exc} = 80$, $N_{inh} = 20$, total simulation duration $T = 50$ s, time step $dt = 0.01$ s, $g(\omega)$ of both
excUnits and inhUnits followed Gaussian distribution $g(\omega) \sim \mathcal{N}(0, 0.01)$ for both excUnits and
inhUnits. The range of time to evaluate the mean of the order parameters R_{exc} and R_{inh} was from
T/2 to T to exclude initial fluctuations. In Fig. 2–5, K_{ii} and K_{ie} are integrated with the same
parameter K_i to simplify the simulation.

427
$$K_i = K_{ie} = K_{ii} \tag{M10}.$$

428
429 At each time step, the J_{ee} values were first modulated by Hebbian potentiation and then rescaled
430 using homeostatic regularization rules. The Hebbian potentiation plasticity is implemented as an
431 exponential function:

432
$$\frac{dJ_{ee,kj}}{dt} = \alpha \exp(-\gamma |\Delta\theta_{kj}|) \tag{11}.$$

433
$$J_{Hebb} \leftarrow J_{ee,kj} + \alpha \exp(-\gamma |\Delta\theta_{kj}|) \tag{M11}.$$

434 α is a learning rate for the Hebbian plasticity. Note that α should scale proportionally with an
435 inverse of N_{exc} as $\alpha \propto 1/N_{exc}$ to maintain consistency across different system sizes. γ determines
436 a sharpness exponential decay from the peak. $\Delta\theta_{kj}$ is the phase difference between excUnit k
437 and j

438
$$\Delta\theta_{kj} = \theta_{k,exc} - \theta_{j,exc} \left(-\pi \leq \Delta\theta_{kj} < \pi\right) \tag{M12}.$$

439 The implementation of Hebbian plasticity with a cosine function (SI Appendix, Fig. S1) is defined

440
$$\frac{dJ_{ee,kj}}{dt} = \frac{\alpha}{2}\{\cos(\Delta\theta_{kj}) + 1\} \tag{13}.$$

441
$$J_{Hebb} \leftarrow J_{ee,kj} + \frac{\alpha}{2}\{\cos(\Delta\theta_{kj}) + 1\} \tag{M13}.$$

442 In any Hebbian learning rule in this paper, $dJ_{ee,kj}/dt = dJ_{ee,jk}/dt$ holds due to the symmetry with
443 respect to the sign of $\Delta\theta_{kj}$.
444
445 After the Hebbian modulations, J_{ee} was rescaled with the homeostatic regularization plasticity rule
446 to prevent overpotentiation. Plasticity was implemented as follows:

447
$$\hat{J}_{ee,kj} = \frac{\hat{\sigma}_{Jee}}{S[J_{ee}]_n}(J_{ee,kj} - E[J_{ee}]_n) + \hat{\mu}_{Jee}$$

448
$$J_{ee,kj} \leftarrow \beta\hat{J}_{ee,kj} + (1-\beta)J_{ee,kj} \tag{14}.$$

449
$$\hat{J}_{ee,kj} = \frac{\hat{\sigma}_{Jee}}{S[J_{Hebb}]_n}(J_{Hebb} - E[J_{Hebb}]_n) + \hat{\mu}_{Jee}$$

450
$$J_{ee,kj} \leftarrow \beta\hat{J}_{Hebb} + (1-\beta)J_{Hebb} \tag{M14}.$$

$J_{ee,kj}$ is scaled adjusting to initial μ_{Jee} and σ_{Jee} settings, $\hat{\mu}_{Jee} = \mu_{Jee}/N_{exc}$, $\hat{\sigma}_{Jee} = \sigma_{Jee}/N_{exc}$. $E[J_{ee}]_n$
and $S[J_{ee}]_n$ are the calculated mean and standard deviation of the J_{ee} distribution among the units

at every time point immediately after Hebbian modification, respectively. β is a learning rate for
the regularization plasticity. This process rescales the J_{ee} distribution toward distribution with
mean $\hat{\mu}_{Jee}$ and standard deviation $\hat{\sigma}_{Jee}$.

Cyclic inhibitory modulation In Fig. 5, K_i is modulated as a sine function
$$K_i = A_s \sin(\omega_s t) + C_s \tag{M15},$$
where A_s (= 0.32), ω_s (= 0.004π), and C_s (= 1.4) are the amplitude, frequency, and baseline
constant for the sine modulation, respectively. The other parameters for interaction strengths are
held constant to simplify the simulation; μ_{Jee} = 1, σ_{Jee} = 0.1, and K_{ei} = 3.

Statistics $E[\cdot]_t$ (also $E\{\cdot\}_t$) and $S[\cdot]_t$ are the mean and the standard deviation over time,
respectively, where t indicates the number of time points. $E[\cdot]_n$ and $S[\cdot]_n$ are the mean and the
standard deviation among unit pairs, respectively, where n indicates the number of unit pairs.
Where · represents a certain variable. For Fig. 5B lower panel, the time window ($\pm$15 sec from
the time point) was set.

Descriptive statistics (mean, standard deviation, kurtosis, and skewness) and statistical tests
were performed using Python Numpy and Scipy libraries. The Dip test (36) was performed using
the Python Diptest Library (https://github.com/RUrlus/diptest). The dip value is the maximum
distance between the actual data distribution and the fitting result of the monomodal Gaussian
distribution. If the data distribution is close to a monomodal Gaussian distribution, the dip value is
low. If is bimodal, then its value is high. Hierarchical clustering was performed using the linkage
function of the Python Scipy library.

Code Availability Statement The scripts for this study can be found at Google Colaboratory:
https://colab.research.google.com/drive/1v8JF97G9YdwUOh_Wuh9i_TvuGa42fLAu?usp=sharing

Acknowledgments

We would like to thank Susumu Setogawa for their insightful contributions to the discussion. This
study was supported by JSPS KAKENHI [grant 24K18243 to S.K.; grant 23H02788 to K.M.] and
the Takeda Science Foundation [to K.M.]. The funding sources had no involvement in the study
design, data collection, interpretation, or the decision to submit this paper for publication.

References

1.   G. Wallas, *The art of thought* (Jonathan Cape, 1926).
2.   J. G. Jenkins, K. M. Dallenbach, Obliviscence during sleep and waking. *Am. J. Psychol.*
35, 605 (1924).
3.   N. Landmann, et al., The reorganisation of memory during sleep. *Sleep Med. Rev*. **18,**
531–541 (2014).
4.   P. A. Lewis, G. Knoblich, G. Poe, How memory replay in sleep boosts creative problem-
solving. *Trends Cogn. Sci.* **22**, 491–503 (2018).
5.   H. Norimoto, et al., Hippocampal ripples down-regulate synapses. *Science* **359**, 1524–
1527 (2018).
6.   A. Goto, et al., Stepwise synaptic plasticity events drive the early phase of memory
consolidation. *Science* **374,** 857–863 (2021).
7.   K. D. Harris, A. Thiele, Cortical state and attention. *Nat. Rev. Neurosci*. **12**, 509–523
(2011).
8.   M. Steriade, R. R. Llinás, The functional states of the thalamus and the associated
neuronal interplay. *Physiol. Rev*. **68**, 649–742 (1988).
9.   B. Haider, M. Häusser, M. Carandini, Inhibition dominates sensory responses in the
awake cortex. *Nature* **493,** 97–100 (2013).

10. H. Miyawaki, K. Diba, Regulation of hippocampal firing by network oscillations during
sleep. *Curr. Biol*. **26**, 893–902 (2016).
11. H. Alfonsa, et al., Intracellular chloride regulation mediates local sleep pressure in the
cortex. *Nat. Neurosci*. **26**, 64–78 (2022).
12. K. Mizuseki, G. Buzsáki, Preconfigured, skewed distribution of firing rates in the
hippocampus and entorhinal cortex. *Cell Rep*. **4**, 1010–1021 (2013).
13. Y. Kuramoto, Chemical oscillations, waves, and turbulence (Springer, 1984).
14. Y. Kuramoto, Self-entrainment of a population of coupled non-linear oscillators in
International Symposium on Mathematical Problems in Theoretical Physics (Springer, 1975), pp.
420–422.
15. E. Montbrió, D. Pazó, Kuramoto model for excitation-inhibition-based oscillations. *Phys.*
Rev. Lett. **120,** 244101 (2018).
16. S. Kuroki, K. Mizuseki, Excitation-inhibition balance controls synchronization in a simple
model of coupled phase oscillators. *Neural Comput*. **37**, 1353–1372 (2025).
17. D. O. Hebb, The organization of behavior; a neuropsychological theory (Wiley, 1949).
18. G. G. Turrigiano, The self-tuning neuron: synaptic scaling of excitatory synapses. ***Cell***
135, 422–435 (2008).
19. T. Toyoizumi, M. Kaneko, M. P. Stryker, K. D. Miller, Modeling the dynamic interaction of
Hebbian and homeostatic plasticity. *Neuron* **84**, 497–510 (2014).
20. S. Kim, D. Jung, S. Royer, Place cell maps slowly develop via competitive learning and
conjunctive coding in the dentate gyrus. *Nat. Commun*. **11**, 4550 (2020).
21. S. Benas, X. Fernandez, E. Kropff, Modeled grid cells aligned by a flexible attractor. *eLife*
(2024).
22. S. N. Weber, H. Sprekeler, Learning place cells, grid cells and invariances with excitatory
and inhibitory plasticity. *eLife* **7**, e34560 (2018).
23. T. Aoki, T. Aoyagi, Co-evolution of phases and connection strengths in a network of
phase oscillators*. Phys. Rev. Lett.* **102**, 034101 (2009).
24. J. C. Bronski, et al., The stability of fixed points for a Kuramoto model with Hebbian
interactions. *Chaos* **27**, 053110 (2017).
25. J. Fialkowski, et al., Heterogeneous nucleation in finite-size adaptive dynamical networks.
Phys. Rev. Lett. **130**, 067402 (2023).
26. S. Y. Ha, J. Lee, Z. Li, J. Park, Emergent dynamics of Kuramoto oscillators with adaptive
couplings: Conservation law and fast learning*. SIAM J. Appl. Dyn. Syst*. **17**, 1560–1588 (2018).
27. S. Y. Ha, S. E. Noh, J. Park, Synchronization of Kuramoto oscillators with adaptive
couplings. *SIAM J. Appl. Dyn. Syst.* **15**, 162–194 (2016).
28. R. K. Niyogi, L. Q. English, Learning-rate-dependent clustering and self-development in a
network of coupled phase oscillators. *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* **80**, 066213
(2009).
29. J. Park, D. Poyato, J. Soler, Filippov trajectories and clustering in the Kuramoto model
with singular couplings. *J. Eur. Math. Soc.* **23**, 3193–3278 (2021).
30. C. B. Picallo, H. Riecke, Adaptive oscillator networks with conserved overall coupling:
Sequential firing and near-synchronized states. *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* **83**,
036206 (2011).
31. D. Poyato, Filippov flows and mean-field limits in the kinetic singular Kuramoto model.
arXiv [math.AP] (2019).
32. Q. Ren, J. Zhao, Adaptive coupling and enhanced synchronization in coupled phase
oscillators. *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* **76**, 016207 (2007).
33. P. Seliger, S. C. Young, L. S. Tsimring, Plasticity and learning in a network of coupled
phase oscillators. *Phys. Rev. E Stat. Nonlin. Soft Matter Phys*. **65**, 041906 (2002).
34. G. Buzsáki, K. Mizuseki, The log-dynamic brain: how skewed distributions affect network
operations. *Nat. Rev. Neurosci.* **15**, 264–278 (2014).
35. M. Iwase, K. Diba, E. Pastalkova, K. Mizuseki, Dynamics of spike transmission and
suppression between principal cells and interneurons in the hippocampus and entorhinal cortex.
Hippocampus **34**, 393–421 (2024).

36. J. A. Hartigan, P. M. Hartigan, The dip test of unimodality. *Ann. Stat.* **13**, 70–84 (1985).
37. A. C. Nobre, S. Kastner, Eds., The Oxford Handbook of Attention (Oxford University
Press, 2014).
38. A. J. G. D. Holtmaat, et al., Transient and persistent dendritic spines in the neocortex in
vivo. *Neuron* **45**, 279–291 (2005).
39. M. Matsuzaki, N. Honkura, G. C. R. Ellis-Davies, H. Kasai, Structural basis of long-term
potentiation in single dendritic spines. *Nature* **429**, 761–766 (2004).
40. J. Grutzendler, N. Kasthuri, W.-B. Gan, Long-term dendritic spine stability in the adult
cortex. *Nature* **420**, 812–816 (2002).
41. G. Tononi, C. Cirelli, Sleep and synaptic down-selection. *Eur. J. Neurosci.* **51**, 413–421
(2020).
42. H. Kasai, et al., Learning rules and persistence of dendritic spines. *Eur. J. Neurosci*. **32**,
241–249 (2010).
43. L. Bollmann, P. Baracskay, F. Stella, J. Csicsvari, Sleep stages antagonistically modulate
reactivation drift. *Neuron* **113,** 1446-1459.e6 (2025).
44. S. P. Vaidya, G. Li, R. A. Chitwood, Y. Li, J. C. Magee, Formation of an expanding
memory representation in the hippocampus. *Nature Neurosci.*, in press.
45. Y. Ziv, et al., Long-term dynamics of CA1 hippocampal place codes. *Nat. Neurosci.* **16**,
264–266 (2013).
46. D. T. Campbell, Blind variation and selective retention in creative thought as in other
knowledge processes. *Psychol. Rev.* **67**, 380–400 (1960).
47. S. N. Gomperts, F. Kloosterman, M. A. Wilson, VTA neurons coordinate with the
hippocampal reactivation of spatial experience. *eLife* **4**, e05360 (2015).
48. E. M. Izhikevich, Dynamical systems in neuroscience: The geometry of excitability and
bursting (MIT Press, 2010).
49. Z. Brzosko, S. B. Mierau, O. Paulsen, Neuromodulation of spike-timing-dependent
plasticity: Past, present, and future. *Neuron* **103**, 563–581 (2019).
50. Y. K. Wu, C. Miehl, J. Gjorgjieva, Regulation of circuit organization and function through
inhibitory synaptic plasticity. *Trends Neurosci*. **45,** 884–898 (2022).
51. C. Q. Chiu, A. Barberis, M. J. Higley, Preserving the balance: diverse forms of long-term
GABAergic synaptic plasticity. *Nat. Rev. Neurosci*. **20**, 272–281 (2019).
52. D. M. Blitz, K. A. Foster, W. G. Regehr, Short-term synaptic plasticity: a comparison of
two synapses. *Nat. Rev. Neurosci*. **5,** 630–640 (2004).
53. R. E. Beaty, M. Benedek, P. J. Silvia, D. L. Schacter, Creative cognition and brain
network dynamics. *Trends Cogn. Sci.* **20**, 87–95 (2016).
54. J. G. Klinzing, N. Niethard, J. Born, Mechanisms of systems memory consolidation during
sleep. *Nat. Neurosci*. **22**, 1598–1610 (2019).
55. O. Barak, Recurrent neural networks as versatile tools of neuroscience research. *Curr.
Opin. Neurobiol.* **46**, 1–6 (2017).
56. J. A. Acebrón, L. L. Bonilla, C. J. Pérez Vicente, F. Ritort, R. Spigler, The Kuramoto
model: A simple paradigm for synchronization phenomena. *Rev. Mod. Phys.* **77**, 137–185 (2005).
602
603
1. J. G. Jenkins, K. M. Dallenbach, Obliviscence during sleep and waking. *Am. J. Psychol.* **35**,
605 (1924).
2. G. Wallas, *The art of thought* (Jonathan Cape, 1926).
3. N. Landmann, *et al.*, The reorganisation of memory during sleep. *Sleep Med. Rev.* **18**, 531–
541 (2014).
4. P. A. Lewis, G. Knoblich, G. Poe, How memory replay in sleep boosts creative problem-
solving. *Trends Cogn. Sci.* **22**, 491–503 (2018).
5. H. Norimoto, *et al.*, Hippocampal ripples down-regulate synapses. *Science* **359**, 1524–1527
(2018).
6. A. Goto, *et al.*, Stepwise synaptic plasticity events drive the early phase of memory
consolidation. *Science* **374**, 857–863 (2021).

7.   K. D. Harris, A. Thiele, Cortical state and attention. *Nat. Rev. Neurosci.* **12**, 509–523 (2011).
8.   M. Steriade, R. R. Llinás, The functional states of the thalamus and the associated neuronal
interplay. *Physiol. Rev.* **68**, 649–742 (1988).
9.   B. Haider, M. Häusser, M. Carandini, Inhibition dominates sensory responses in the awake
cortex. *Nature* **493**, 97–100 (2013).
10. H. Miyawaki, K. Diba, Regulation of hippocampal firing by network oscillations during sleep.
Curr. Biol. **26**, 893–902 (2016).
11. H. Alfonsa, *et al.*, Intracellular chloride regulation mediates local sleep pressure in the cortex.
Nat. Neurosci. **26**, 64–78 (2022).
12. K. Mizuseki, G. Buzsáki, Preconfigured, skewed distribution of firing rates in the
hippocampus and entorhinal cortex. *Cell Rep.* **4**, 1010–1021 (2013).
13. Y. Kuramoto, *Chemical oscillations, waves, and turbulence* (Springer, 1984).
14. Y. Kuramoto, Self-entrainment of a population of coupled non-linear oscillators in
International Symposium on Mathematical Problems in Theoretical Physics, (Springer, 1975),
pp. 420–422.
15. E. Montbrió, D. Pazó, Kuramoto model for excitation-inhibition-based oscillations. *Phys. Rev.
Lett.* **120**, 244101 (2018).
16. S. Kuroki, K. Mizuseki, Excitation-inhibition balance controls synchronization in a simple
model of coupled phase oscillators. *Neural Comput.* **37**, 1353–1372 (2025).
17. D. O. Hebb, *The organization of behavior; a neuropsychological theory* (Wiley, 1949).
18. G. G. Turrigiano, The self-tuning neuron: synaptic scaling of excitatory synapses. *Cell* **135**,
422–435 (2008).
19. T. Toyoizumi, M. Kaneko, M. P. Stryker, K. D. Miller, Modeling the dynamic interaction of
Hebbian and homeostatic plasticity. *Neuron* **84**, 497–510 (2014).
20. S. Kim, D. Jung, S. Royer, Place cell maps slowly develop via competitive learning and
conjunctive coding in the dentate gyrus. *Nat. Commun.* **11**, 4550 (2020).
21. S. Benas, X. Fernandez, E. Kropff, Modeled grid cells aligned by a flexible attractor. *eLife*
(2024).
22. S. N. Weber, H. Sprekeler, Learning place cells, grid cells and invariances with excitatory
and inhibitory plasticity. *Elife* **7**, e34560 (2018).
23. T. Aoki, T. Aoyagi, Co-evolution of phases and connection strengths in a network of phase
oscillators. *Phys. Rev. Lett.* **102**, 034101 (2009).
24. J. C. Bronski, *et al.*, The stability of fixed points for a Kuramoto model with Hebbian
interactions. *Chaos* **27**, 053110 (2017).
25. J. Fialkowski, *et al.*, Heterogeneous nucleation in finite-size adaptive dynamical networks.
Phys. Rev. Lett. **130**, 067402 (2023).
26. S. Y. Ha, J. Lee, Z. Li, J. Park, Emergent dynamics of Kuramoto oscillators with adaptive
couplings: Conservation law and fast learning. *SIAM J. Appl. Dyn. Syst.* **17**, 1560–1588
(2018).
27. S. Y. Ha, S. E. Noh, J. Park, Synchronization of Kuramoto oscillators with adaptive
couplings. *SIAM J. Appl. Dyn. Syst.* **15**, 162–194 (2016).
28. R. K. Niyogi, L. Q. English, Learning-rate-dependent clustering and self-development in a
network of coupled phase oscillators. *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* **80**,
066213 (2009).
29. J. Park, D. Poyato, J. Soler, Filippov trajectories and clustering in the Kuramoto model with
singular couplings. *J. Eur. Math. Soc.* **23**, 3193–3278 (2021).
30. C. B. Picallo, H. Riecke, Adaptive oscillator networks with conserved overall coupling:
Sequential firing and near-synchronized states. *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.*
83, 036206 (2011).
31. D. Poyato, Filippov flows and mean-field limits in the kinetic singular Kuramoto model. *arXiv
[math.AP]* (2019).
32. Q. Ren, J. Zhao, Adaptive coupling and enhanced synchronization in coupled phase
oscillators. *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* **76**, 016207 (2007).

33.   P. Seliger, S. C. Young, L. S. Tsimring, Plasticity and learning in a network of coupled phase
oscillators. *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* **65**, 041906 (2002).

34.   G. Buzsáki, K. Mizuseki, The log-dynamic brain: how skewed distributions affect network
operations. *Nat. Rev. Neurosci.* **15**, 264–278 (2014).

35.   M. Iwase, K. Diba, E. Pastalkova, K. Mizuseki, Dynamics of spike transmission and
suppression between principal cells and interneurons in the hippocampus and entorhinal
cortex. *Hippocampus* **34**, 393–421 (2024).

36.   J. A. Hartigan, P. M. Hartigan, The dip test of unimodality. *Ann. Stat.* **13**, 70–84 (1985).

37.   A. C. Nobre, S. Kastner, Eds., *The Oxford Handbook of Attention* (Oxford University Press,
2014).

38.   A. J. G. D. Holtmaat, *et al.*, Transient and persistent dendritic spines in the neocortex in vivo.
Neuron **45**, 279–291 (2005).

39.   M. Matsuzaki, N. Honkura, G. C. R. Ellis-Davies, H. Kasai, Structural basis of long-term
potentiation in single dendritic spines. *Nature* **429**, 761–766 (2004).

40.   J. Grutzendler, N. Kasthuri, W.-B. Gan, Long-term dendritic spine stability in the adult cortex.
Nature **420**, 812–816 (2002).

41.   G. Tononi, C. Cirelli, Sleep and synaptic down-selection. *Eur. J. Neurosci.* **51**, 413–421
(2020).

42.   H. Kasai, *et al.*, Learning rules and persistence of dendritic spines. *Eur. J. Neurosci.* **32**, 241–
249 (2010).

43.   L. Bollmann, P. Baracskay, F. Stella, J. Csicsvari, Sleep stages antagonistically modulate
reactivation drift. *Neuron* **113**, 1446-1459.e6 (2025).

44.   S. P. Vaidya, G. Li, R. A. Chitwood, Y. Li, J. C. Magee, Formation of an expanding memory
representation in the hippocampus. *Nature Neuroscience* 1–9 (2025).

45.   Y. Ziv, *et al.*, Long-term dynamics of CA1 hippocampal place codes. *Nat. Neurosci.* **16**, 264–
266 (2013).

46.   B. Bartoldson, A. Morcos, A. Barbu, G. Erlebacher, The generalization-stability tradeoff in
neural network pruning in *Advances in Neural Information Processing Systems*, H.
Larochelle, M. Ranzato, R. Hadsell, M. F. Balcan, H. Lin, Eds. (Curran Associates, Inc.,
2020), pp. 20852–20864.

47.   T. Hoefler, D. Alistarh, T. Ben-Nun, N. Dryden, A. Peste, Sparsity in deep learning: pruning
and growth for efficient inference and training in neural networks. *J. Mach. Learn. Res.* **22**,
10882–11005 (2021).

48.   D. T. Campbell, Blind variation and selective retention in creative thought as in other
knowledge processes. *Psychol. Rev.* **67**, 380–400 (1960).

49.   S. N. Gomperts, F. Kloosterman, M. A. Wilson, VTA neurons coordinate with the
hippocampal reactivation of spatial experience. *Elife* **4**, e05360 (2015).

50.   G. Dragoi, S. Tonegawa, Preplay of future place cell sequences by hippocampal cellular
assemblies. *Nature* **469**, 397–401 (2011).

51.   J.-E. K. Miller, I. Ayzenshtat, L. Carrillo-Reid, R. Yuste, Visual stimuli recruit intrinsically
generated cortical ensembles. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E4053-61 (2014).

52.   A. Luczak, P. Barthó, K. D. Harris, Spontaneous events outline the realm of possible sensory
responses in neocortical populations. *Neuron* **62**, 413–425 (2009).

53.   E. M. Izhikevich, *Dynamical systems in neuroscience: The geometry of excitability and
bursting* (MIT Press, 2010).

54.   Z. Brzosko, S. B. Mierau, O. Paulsen, Neuromodulation of spike-timing-dependent plasticity:
Past, present, and future. *Neuron* **103**, 563–581 (2019).

55.   Y. K. Wu, C. Miehl, J. Gjorgjieva, Regulation of circuit organization and function through
inhibitory synaptic plasticity. *Trends Neurosci.* **45**, 884–898 (2022).

56.   C. Q. Chiu, A. Barberis, M. J. Higley, Preserving the balance: diverse forms of long-term
GABAergic synaptic plasticity. *Nat. Rev. Neurosci.* **20**, 272–281 (2019).

57.   D. M. Blitz, K. A. Foster, W. G. Regehr, Short-term synaptic plasticity: a comparison of two
synapses. *Nat. Rev. Neurosci.* **5**, 630–640 (2004).

58. R. E. Beaty, M. Benedek, P. J. Silvia, D. L. Schacter, Creative cognition and brain network
722   dynamics. *Trends Cogn. Sci.* **20**, 87–95 (2016).
723   59. J. G. Klinzing, N. Niethard, J. Born, Mechanisms of systems memory consolidation during
724   sleep. *Nat. Neurosci.* **22**, 1598–1610 (2019).
725   60. O. Barak, Recurrent neural networks as versatile tools of neuroscience research. *Curr. Opin.*
726   *Neurobiol.* **46**, 1–6 (2017).
727   61. J. A. Acebrón, L. L. Bonilla, C. J. Pérez Vicente, F. Ritort, R. Spigler, The Kuramoto model: A
728   simple paradigm for synchronization phenomena. *Rev. Mod. Phys.* **77**, 137–185 (2005).
729

Figures and Tables

Figure 1. EI-Kuramoto model with distributed J$_{ee}$ and plasticity

(A) Schematic representation of the pEI-Kuramoto model

Left: Oscillator units are excitatory (warm-colored) or inhibitory (cool-colored), each with its own

natural frequency (shown by color gradients). Right: J$_{ee}$ functions attractively between excUnits

(red arrows), with each unit pair having its own coupling weight and undergoing plastic

modification (the right matrix). K_{ei} is the attractive interaction strength from excUnit to inhUnit
(magenta arrow). K_{ii} is the repulsive strength between inhUnits (blue arrows). K_{ie} is the repulsive
strength from inhUnit to excUnit (cyan arrow). K_{ei}, K_{ii}, and K_{ie} are constant and applied uniformly
across all units. (B) Distribution of J_{ee} coupling weights with different σ_{Jee}
$\mu_{Jee} = 1$, $N_{exc} = 80$. (C) R_{exc} time traces of synchronized, bistable, and desynchronized states with
each σ_{Jee} (D) Exponential rules of Hebbian potentiation plasticity
The J_{ee} weight of each pair of units is potentiated based on the phase difference between the
units. (E) Validation of the J_{ee} distribution with various plasticity parameters
Skewness (top panels), kurtosis (middle), and dip values (bottom). The parameters used for
evaluation were $\mu_{Jee} = 1$, $\sigma_{Jee} = 0.1$, $K_{ei} = 3$, $K_{ie} = 3$, $K_{ii} = 3$, corresponding to a desynchronized
state. (F) J_{ee} distribution of exponential Hebbian plasticity rules with various parameter settings
The magenta, green, and yellow circles correspond to the parameter settings in (E). (G)
Validation of the synchronization states with various plasticity parameters. $E[R_{exc}]_t$ and $S[R_{exc}]_t$
denote the time-averaged mean and standard deviation, respectively, of the order parameter for
excUnits, R_{exc}.
752

Figure 2. Changes in J_{ee} fluctuation with K_i modification

 (A–D) (A) $E\{S[J_{ee}]_t\}_n$, (B) $S\{S[J_{ee}]_t\}_n$, (C) $E[R_{exc}]_t$, and (D) $S[R_{exc}]_t$ as a function of K_i

 Red crosses indicate each value in the K_i conditions shown in (E) and (F). (E) Time course of J_{ee}
 weights in different K_i conditions. The order of unit pairs is sorted by $E[J_{ee}]_t$ for each K_i value. (F)
 The upper panels show R_{exc} traces and time points of the left J_{ee} matrices. The red and blue
 vertical lines indicate the time points (red and blue are local maximum and local minimum,
 respectively, for K_i = 0.8 and 1.0; constant 5 s interval for K_i = 0.1 and 1.5). The lower panels
 show the J_{ee} weight matrix at selected time points (local minimum or maximum for K_i = 0.8 and
 1.0, constant 5 s intervals for K_i = 0.1 and 1.5). The order of units is rearranged by the result of
 hierarchical clustering for the leftmost panel (around t: 24-25). E and F share the color bar.

Figure 3. Individual J_{ee} characterization of the pEI-Kuramoto model

(A) $E[J_{ee}]_t$ histogram (B) $S[J_{ee}]_t$ histogram (C) $E[J_{ee}]_t$ vs. $S[J_{ee}]_t$ scatter plot for different K_i values

773
Figure 4. Factors for J$_{ee}$ fluctuation
(A–C) E[J$_{ee}$]$_t$ vs. S[J$_{ee}$]$_t$ scatter plot with color coding of (A) the natural frequency ω$_{exc}$ differences,
(B) the initial phase differences, and (C) the initial J$_{ee}$ weights over different K$_i$
The red dotted line in the upper left panel indicates E[J$_{ee}$] = 0.018. (D–F) Violin plots provide
distribution densities based on whether the threshold E[J$_{ee}$]$_t$ ≥ 0.18 or not for (D) natural
frequency ω$_{exc}$ differences, (E) initial phase differences, and (F) initial J$_{ee}$ weights across different
K$_i$
The red and blue horizontal lines indicate the mean and median values, respectively. The red
vertical lines indicate the standard errors. The numbers in the upper right corner of each panel
are the p-values from the Mann–Whitney U test. (G and H) Scatter plots of natural frequency
differences and initial phase differences with color coding of (G) E[J$_{ee}$]$_t$ and (H) S[J$_{ee}$]$_t$ (K$_i$ = 1.5)
The Mann–Whitney U test was used for statistical analysis in panels D-F. The numbers of unit
pairs were as follows: for K$_i$=0.8, <0.18: n=5968, ≥0.18: n=352; for K$_i$=1.0, <0.18: n=5920, ≥0.18:
n=400; for K$_i$=1.5, <0.18: n=5852, ≥0.18: n=468. The statistics were as follows: (D, natural
frequency difference), K$_i$=0.8, z=24.7, p<0.001; K$_i$=1.0, z=26.8, p<0.001; K$_i$=1.5, z=27.8, p<0.001.
(E, initial phase difference) K$_i$=0.8, z=9.20, p<0.001; K$_i$=1.0, z=16.4, p<0.001; K$_i$=1.5, z=16.7,
p<0.001. (F, initial J$_{ee}$ weight) K$_i$=0.8, z=-0.046, p=0.96; K$_i$=1.0, z=0.046, p=0.96; K$_i$=1.5, z=0.17,
p=0.87.
792

794
795 **Figure 5. J_{ee} fluctuation changes with K_i cyclic modulation**
796 (A) Time trace of each J_{ee} weight sorted by $E[J_{ee}]_t$. (B) Time traces of (top) K_i modulation, (middle)
797 R_{exc}, and (low) $E\{S[J_{ee}]_t\}_n$
798
799

Supporting Information for
Synchronicity transitions determine coupling fluctuations in a
model of coupled oscillators with plasticity

Satoshi Kuroki and Kenji Mizuseki

Satoshi Kuroki
Email: skuroki@omu.ac.jp

This PDF file includes:

> Supporting text S1
> Figures S1 to S3
> Legends for Movies S1 to S5

Other supporting materials for this manuscript include the following:

> Movies S1 to S5

Supporting Information Text

Text S1. Derivation of the quasi-steady coupling weight distribution

The combined changes from Hebbian and homeostatic plasticity (Eq M11 and M14) are summarized as follows:

$$J_{ee,kj}(t + \Delta t) = J_{ee,kj}(t) + \alpha \exp(-\gamma|\Delta\theta_{kj}|) - \beta\{J_{ee,kj}(t) + \alpha \exp(-\gamma|\Delta\theta_{kj}|) - \hat{J}_{Hebb}\} \quad \text{(S1)}.$$

Assuming quasi-steady J_{ee} distribution, $J_{ee,kj}(t + \Delta t) = J_{ee,kj}(t)$. If α is small enough, homeostatic set point attained after Hebbian potentiation $\hat{J}_{Hebb}$ can be approximated as that of before Hebbian potentiation, $\hat{J}_{Hebb} \approx \hat{J}_{ee,jk}$. Under these assumptions, the equation simplifies to:

$$\alpha(1 - \beta) \exp(-\gamma|\Delta\theta_{kj}|) - \beta(J_{ee,kj} - \hat{J}_{ee,kj}) = 0 \quad \text{(S2)}.$$

Thus,

$$J_{ee,kj} = \hat{J}_{ee,kj} + c \exp(-\gamma|\Delta\theta_{kj}|) \quad \text{(S3)}.$$

$$c = \frac{\alpha(1 - \beta)}{\beta}$$

This corresponds to Eq. 1 in the main text. The first term, $\hat{J}_{ee,kj}$, represents the homeostatic set point characterized by a J_{ee} distribution with mean $\hat{\mu}_{Jee}$ and variance $\hat{\sigma}_{Jee}$. This second term is the Hebbian potentiation component with a coefficient of $c = \alpha(1 - \beta)/\beta$. Repeating this update, the J_{ee} distribution gradually reflects the influence of the $\Delta\theta$-dependent Hebbian term.

We defined the probability density function (PDF) of the phases $f_\theta(\theta)$ assuming enough large number of units:

$$\int_0^{2\pi} f_\theta(\theta)d\theta = 1 \quad \text{(S4)}$$

The distribution of the phase difference, $f_{\Delta\theta}(\Delta\theta)$, between two independent phases is given by the circular autocorrelation of the phase distribution:

$$f_{\Delta\theta}(\Delta\theta) = \int_0^{2\pi} f_\theta(\theta)f_\theta(\theta + \Delta\theta)d\theta \quad (-\pi \leq \Delta\theta < \pi) \quad \text{(S5)}$$

In a fully desynchronized state, the phase distribution is uniform, $f(\theta) = 1/2\pi$. In this case, Eq. S5 becomes:

$$f_{\Delta\theta}(\Delta\theta) = \int_0^{2\pi} \frac{1}{2\pi} \cdot \frac{1}{2\pi} d\theta = \frac{1}{2\pi} \quad \text{(S6)}$$

This shows that the distribution of phase differences is also uniform. It follows that the distribution of the absolute phase difference, $x = |\Delta\theta| \sim \text{Unif}(0, \pi)$, is a uniform distribution over [0,π], with a PDF:

$$f_{|\Delta\theta|}(x) = \frac{1}{\pi} \quad \text{(S7)}$$

Here, we evaluate $E[J_{ee}]$ and $S[J_{ee}]$ to calculate $\hat{J}_{ee,kj}$. We analyze the distribution of the Hebbian component from Eq. S3, $y = c \exp(-\gamma x)$, which is a transformation of the random variable x. Using this, the inverse transformation is:

$$x = -\frac{1}{\gamma} \log\frac{y}{c} \quad \text{(S9)}$$

The absolute value of the Jacobian of this transformation is:

$$\left|\frac{dx}{dy}\right| = \frac{1}{\gamma y} \quad \text{(S10)}$$

The relationship between the PDFs is given by $f_{|\Delta\theta|}(x)dx = f_{Jee}(y)dy$. Substituting the known quantities, we find the PDF for x:

$$f_{Jee}(y) = f_{|\Delta\theta|}(x)\left|\frac{dx}{dy}\right| = \frac{1}{\gamma\pi y} \quad (y_{min} = ce^{-\gamma\pi} \leq y \leq y_{max} = c) \quad \text{(S12)}$$

Assuming the statistics of the full J_{ee} distribution are dominated by this Hebbian component in the desynchronized state, The mean $E[J_{ee}]$ and variance $Var[J_{ee}]$ is:

$$E[J_{ee}] = \int_{y_{min}}^{y_{max}} y f_{Jee}(y)dy = \int_{ce^{-\gamma\pi}}^{c} y \frac{1}{\gamma\pi y} dy = \frac{c(1 - e^{-\pi\gamma})}{\gamma\pi} \quad \text{(S13)}$$

$$Var[J_{ee}] = E[J_{ee}^2] - (E[J_{ee}])^2 = \int_{ce^{-\gamma\pi}}^{c} y^2 \frac{1}{\gamma\pi y} dy - (E[J_{ee}])^2 = \frac{c^2(1 - e^{-2\pi\gamma})}{2\gamma\pi} - \left\{\frac{c(1 - e^{-\pi\gamma})}{\gamma\pi}\right\}^2 \quad \text{(S14)}$$

Thus,

$$\hat{J}_{ee,kj} = \hat{\sigma}_{Jee}\left(\frac{j - E[J_{ee}]}{S[J_{ee}]}\right) + \hat{\mu}_{Jee}$$

$$\frac{x - E[J_{ee}]}{S[J_{ee}]} = \left[\frac{1}{\pi\gamma c \exp(-\gamma|\Delta\theta|)} - \frac{c(1 - e^{-\pi\gamma})}{\gamma\pi}\right]\Bigg/\left[\frac{c^2(1 - e^{-2\pi\gamma})}{2\gamma\pi} - \left\{\frac{c(1 - e^{-\pi\gamma})}{\gamma\pi}\right\}^2\right]^{\frac{1}{2}} \qquad (S15)$$

Next, we considered the change in the phase difference for a pair of excUnits. Given Eq. M9 and S3, we calculate $d\Delta\theta_{kj}/dt = d\theta_k/dt - d\theta_j/dt$.

$$\frac{d\Delta\theta_{kj}}{dt} = \Delta\omega - 2\left\{\hat{J}_{ee,kj} + c\exp\left(-\gamma|\Delta\theta_{kj}|\right)\right\}\sin\left(\Delta\theta_{kj}\right) + F_{exc}^{(-kj)} - F_{inh} \qquad (S16)$$

, where natural frequency difference $\Delta\omega = \omega_k - \omega_j$, repulsive force from the inhUnits $F_{inh} = -K_{ie}R_{inh}\{\sin(\theta_k - \psi_{inh}) - \sin(\theta_j - \psi_{inh})\}$, and attractive force from the excUnits except unit k and j, $F_{exc}^{(-kj)}$. Under a mean-field approximation for sufficiently large number of units, this excitatory force term can be written as:

$$F_{exc}^{(-kj)} = -\sum_{l\neq k,j}\left\{J_{ee,kl}\sin(\theta_k - \theta_l) - J_{ee,jl}\sin(\theta_j - \theta_l)\right\}$$

$$\approx -J'^{(-kj)}_{ee}R_{exc}^{(-kj)}\left\{\sin\left(\theta_k - \psi_{exc}^{(-kj)}\right) - \sin\left(\theta_j - \psi_{exc}^{(-kj)}\right)\right\} \qquad (S17)$$

, where $J'^{(-kj)}_{ee}$ represents the effective attractive interaction strength on the pair from other excUnits. This system also reaches a steady state where $d\Delta\theta_{kj}/dt = 0$. Under this condition,

$$-\Delta\omega + F_{inh} - F_{exc}^{(-kj)} = -2\left\{\hat{J}_{ee,kj} + c\exp\left(-\gamma|\Delta\theta_{kj}|\right)\right\}\sin\left(\Delta\theta_{kj}\right) \qquad (S18)$$

This corresponds to Eq. 2 in the main text. The right-hand side represents the interaction term between the excUnit k and j, while the left-hand side of this equation comprises all other influences, namely the natural frequency difference among the pair and the net mean-field force from the other units. A stable fixed point for Δθkj exists where the functions for the left and right-hand sides intersect (SI Appendix, Fig. S3). In the desynchronized state, the order parameters Rexc, Rinh are close to zero, limiting the influence of F_{inh} and $F_{exc}^{(-kj)}$. However, in the bistable state, these order parameters fluctuate periodically. This causes these forces to become influential, and as a result, the Jee distribution is no longer in a quasi-steady state. Furthermore, the interaction term $\sin(\theta_k - \psi) - \sin(\theta_j - \psi)$ in F_{inh} and $F_{exc}^{(-kj)}$ is enhanced when the mean phase ψ lies on the shorter arc between θ_k and θ_j, but is weakened when it lies on the longer arc.

Fig. S1. Cosine Hebbian plasticity rule

(A) Cosine rule for the Hebbian potentiation plasticity. The J_{ee} weight of each pair of units is potentiated based on the phase difference between the units. (B) J_{ee} distribution using the cosine Hebbian plasticity rule. (C) Phase and phase difference distributions for each plasticity setting. Left panels: Polar plots showing the phases of excUnits (warm colors) and inhUnits (cool colors). Center panels: Histograms of the excUnit phases. Right panels: Histograms of the phase differences between pairs of excUnits.

Fig. S2. Effect or Hebbian learning rate and J_{ee} variance for the synchronization

(A) $E[R_{exc}]_t$ as a function of K_i at different learning rates under Hebbian plasticity (α) settings
(B) $E[R_{exc}]_t$ as a function of K_i for different σ_{Jee} settings. The thick black lines are the same as those in Figure 2.

Fig. S3. Interaction function and equilibrium

The thick solid line indicates interaction function between excUnit pair $-2\left\{\hat{J}_{ee} + c\exp(-\gamma|\Delta\theta|)\right\}\sin(\Delta\theta)$. The thin solid line is total interaction strength from the other factors $-\Delta\omega + F_{inh} - F_{exc}^{(-kj)}$. The dot line indicates pure sine curve $\hat{\mu}_{Jee}\sin(\Delta\theta)$ for comparison. The black and white points indicate stable and unstable equilibrium, respectively. The parameter setting is α=0.01, β=0.5, γ=2, $\hat{\mu}_{Jee}$ =0.0125 (μ_{Jee}/N_{exc}= 1/80), $\hat{\sigma}_{Jee}$ = 0.00125 (σ_{Jee}/N_{exc}= 0.1/80) (Fig 1F magenta). $\Delta\omega$ = -0.01, $F_{inh} - F_{exc}^{(-kj)} = -0.2\sin\Delta\theta$.

Movie S1 (separate file) Synchronized state: $\mu_{Jee} = 1$, $\sigma_{Jee} = 0.1$, $K_{ei}=3$, $K_{ie}=0.1$, $K_{ii}=0.1$, $\gamma=2$, $\alpha=0.01$, $\beta=0.5$; playback speed: ×1

Movie S2 (separate file) Desynchronized state: $\mu_{Jee} = 1$, $\sigma_{Jee} = 0.1$, $K_{ei}=3$, $K_{ie}=1.5$, $K_{ii}=1.5$, $\gamma=2$, $\alpha=0.01$, $\beta=0.5$; playback speed: ×1

Movie S3 (separate file) Bistable state1: $\mu_{Jee} = 1$, $\sigma_{Jee} = 0.1$, $K_{ei}=3$, $K_{ie}=0.8$, $K_{ii}=0.8$, $\gamma=2$, $\alpha=0.01$, $\beta=0.5$; playback speed: ×1

Movie S4 (separate file) Bistable state2: $\mu_{Jee} = 1$, $\sigma_{Jee} = 0.1$, $K_{ei}=1$, $K_{ie}=1$, $K_{ii}=1$, $\gamma=2$, $\alpha=0.01$, $\beta=0.5$; playback speed: ×1

Movie S5 (separate file) Cyclic inhibitory modulation, K_i, playback speed ×20