Structural robustness and temporal vulnerability of the starvation-responsive metabolic network in liver of healthy and obese mice

Keigo Morita[1,2], Atsushi Hatano[1,3,4], Toshiya Kokaji[1,5], Hikaru Sugimoto[6],

Takaho Tsuchiya[7,8], Haruka Ozaki[7,8], Riku Egami[9], Dongzi Li[1], Akira Terakawa[1],

Satoshi Ohno[1,2,10], Hiroshi Inoue[11], Yuka Inaba[11], Yutaka Suzuki[9],

Masaki Matsumoto[3], Masatomo Takahashi[12], Yoshihiro Izumi[12], Takeshi Bamba[12],

Akiyoshi Hirayama[13], Tomoyoshi Soga[13, 14], and Shinya Kuroda[1,2,6,9,15, *]

[1]Department of Biological Sciences, Graduate School of Science, The University of

Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan.

[2]Molecular Genetics Research Laboratory, Graduate School of Science, The University

of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo, Japan.

[3]Department of Omics and Systems Biology, Graduate School of Medical and Dental

Sciences, Niigata University, 757 Ichibancho, Asahimachi-dori, Chuo-ku, Niigata City,

Niigata 951-8510, Japan.

[4]Laboratory for Integrated Cellular Systems, RIKEN Center for Integrative Medical

Sciences, 1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama, Kanagawa, 230-0045, Japan

[5]Data Science Center, Nara Institute of Science and Technology, 8916-5 Takayama,

Ikoma, Nara, 630-0192, Japan.

[6]Department of Biochemistry and Molecular Biology, Graduate School of Medicine,

The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan.

[7]Bioinformatics Laboratory, Institute of Medicine, University of Tsukuba, Ibaraki,

305-8575, Japan.

[8]Center for Artificial Intelligence Research, University of Tsukuba, Ibaraki, 305-8577,

Japan.

26   [9]Department of Computational Biology and Medical Sciences, Graduate School of

27   Frontier Sciences, The University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa, Chiba 277-

28   8562, Japan.

29   [10]Department of AI Systems Medicine, M&D Data Science Center, Tokyo Medical and

30   Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8510, Japan

31   [11]Metabolism and Nutrition Research Unit, Institute for Frontier Science Initiative,

32   Kanazawa University, 13-1 Takaramachi, Kanazawa, Ishikawa, 920-8641, Japan

33   [12]Division of Metabolomics, Medical Research Center for High Depth Omics, Medical

34   Institute of Bioregulation, Kyushu University, Fukuoka 812-8582, Japan

35   [13]Institute for Advanced Biosciences, Keio University, 246-2 Mizukami, Kakuganji,

36   Tsuruoka, Yamagata 997-0052, Japan

37   [14]Human Biology-Microbiome-Quantum Research Center (WPI-Bio2Q), Keio

38   University, Tokyo 108-8345, Japan

39   [15]Lead contact

40   *Correspondence to: skuroda@bs.s.u-tokyo.ac.jp (S.K.)

Abstract

Adaptation to starvation is a multi-molecular and temporally ordered metabolic process

to optimize fuel usage, that could be impaired in obesity. However, how the healthy

liver adapts to starvation by temporally coordinating various molecules, and how the

process is disrupted in obesity have not been elucidated. Here, we measured time course

multi-omic data in the liver of wild-type (WT) and leptin-deficient obese (*ob*/*ob*) mice

during starvation and constructed the starvation-responsive metabolic network. The hub

molecules of the network in WT mice, including ATP and AMP, were not responsive in

ob/*ob* mice, however the network structure was maintained in *ob*/*ob* mice. The

molecules in the network were temporally ordered through metabolic process driven by

the hub molecules and co-regulated, both were disrupted in *ob*/*ob* mice. Thus, the

starvation-responsive metabolic network is structurally robust, but temporally

vulnerable by the loss of responsiveness of the hub molecules, including ATP and AMP,

in obesity.

Introduction

Adaptation to starvation is one of the most important functions for life to maintain systemic homeostasis. The adaptation is a multi-molecular and temporally ordered process. Various molecules, including metabolites, lipids, metabolic enzymes, transcription factors (TFs), and protein kinases, are involved in this process[1–3]. These molecules coordinate metabolic processes, such as glycogen degradation, gluconeogenesis, and fatty acid degradation, in a temporally ordered manner[1,4–6], important for optimizing fuel usage to keep metabolic homeostasis. However, how the liver, a central organ for metabolism[7,8], coordinates various molecules in a temporally ordered manner during starvation at a multi-omic scale has not been elucidated. A multi-omic measurement at precise time intervals during starvation has not been conducted in mammals, although omic measurements during starvation at a limited number of time intervals have been conducted in mammals[9–19] and time course omic measurements have been conducted only in microorganisms[20–23] or worms[24].

Obesity is a pathological status that affect various metabolic processes[25–29]. Adaptation to starvation is also impaired by obesity[30], but the global landscape of the effects at a multi-omic scale has not been elucidated. The differences of adaptation to starvation between healthy and obese state are clinically important, because obesity is a major therapeutic target of the calorie-restricted diet[31], which are known to have the potential to improve health[3,32,33]. However, differential effects of calorie-restricted diet between healthy and obese individuals have not been fully elucidated[34]. Because starvation is an important factor for the calorie-restricted diet[31], clarifying the differences of adaptation to starvation between healthy and obese mice is informative

for establishing an efficient therapeutic strategy of calorie-restricted diet for obese

patients.

Analysis of properties of biological network is a powerful approach to

understand such a multi-molecular process like adaptation to starvation[35–38]. We have

developed trans-omic analysis, an analytical workflow for addressing multi-molecular

processes[39–41]. With this approach, we can integrate multi-omic data into a trans-omic

network, offering a global landscape of biochemical reactions. Recently, we showed that

analysis of the structural properties of the trans-omic network aids in their interpretation

during oral glucose ingestion[42]. A network is composed of nodes and edges: nodes are

entities such as molecules; edges are relationships between nodes such as regulatory

events. Network structure is how the nodes are connected by edges. Biological networks,

such as metabolic networks[43–45], are often scale-free, which means that a small number

of "hub" nodes interact with many other nodes, and large number of "leaf" nodes

interact with a few other nodes[46]. We showed that the trans-omic network is scale-free

and structural properties of the networks were different between healthy and obese mice

during oral glucose ingestion[42]. However, the relationships between the structural

properties and temporal orders of the trans-omic network has not been investigated, as

well as temporal co-regulations, the correlated temporal patterns of nodes in a biological

network. Because adaptation to starvation is a temporally ordered multi-molecular

process, integrating the structural and temporal analysis of the trans-omic network

would offer mechanistic insights of adaptation to starvation.

In this study, we measured multi-omic data, constructed a trans-omic network,

and conducted structural and temporal analyses of the network in the liver of wild-type

(WT) and leptin-deficient obese (*ob*/*ob*) mice, a commonly used model of obesity

caused by overeating due to loss of leptin[47]. Through structural analyses, we showed

that the starvation-responsive metabolic network is a scale-free network and that the

structural properties of the network were not disrupted in *ob*/*ob* mice. We integrated

structural and temporal analyses of the trans-omic network. We found that the network

was temporally ordered by hub nodes, such as ATP and AMP, in WT mice, but the

temporal order was disrupted in *ob*/*ob* mice, as well as temporal co-regulations. Thus,

the starvation-responsive metabolic network was well-structured, temporally ordered,

and co-regulated in healthy liver. The network was structurally robust but temporally

vulnerable to obesity.

The focus here is adaptive responses to starvation, rather than direct

comparisons between WT and *ob*/*ob* mice in a steady state. We have conducted such

comparative studies previously for fasting and fed states[48,49]. Note that identical data of

0 h starvation have been already published in one of our studies[49]. Additionally, we

focus on the properties of the trans-omic network and the subnetwork that represents the

starvation-responsive metabolic network; analysis of individual metabolic pathways

will be conducted in a separate study.

Results

Workflow for the construction and analysis of a global starvation-responsive trans-

omic network in liver of WT and *ob*/*ob* mice

To describe the temporal dynamics of various molecules during adaptation to starvation

in healthy liver and pathological alterations caused by obesity, we constructed and

analyzed a global starvation-responsive trans-omic network in liver of WT and *ob*/*ob*

mice (Fig. 1). We collected multi-omic data from mice starved for 0, 2, 4, 6, 8, 12, 16,

and 24 hours (Fig. 1a). We measured the metabolome, lipidome, FFA & acyls (free fatty

acids, acylcarnitine, and acyl-CoA), transcriptome, proteome, and phosphoproteome in

liver. We also measured the metabolome in plasma.

Using the collected data, we conducted a single-omic layer analysis (Fig. 1b,

and c). We identified starvation-responsive molecules and classified them into 4 groups

in each layer (Fig. 1b): "common," those that exhibited the same response in both WT

and *ob*/*ob* mice; "WT-specific," those that occurred only in WT mice; "*ob*/*ob*-specific,"

those that occurred only in *ob*/*ob* mice; and "opposite," those that exhibited opposing

responses between WT and *ob*/*ob* mice. These results formed a single omic layer (Fig.

1c). Note that phosphorylation is a process occurred on a phosphosite, not a molecule,

but we refer phosphorylation events as molecules as well as metabolites, lipids, FFA &

acyls, proteins, and transcripts for simplicity.

The starvation-responsive molecules cooperatively regulate metabolic reactions

to adapt to starvation across the multi-omic layers (Fig. 1d). Metabolites, lipids, and

FFA & acyls serve as substrates, products, and allosteric regulators for metabolic

reactions. Enzyme abundance and activity regulate the metabolic reactions. TFs control

enzyme abundance through the gene regulatory network. Enzyme activity is regulated

through the phosphorylation network controlled by protein kinases. Using bioinformatic

platforms (see Methods), we integrated single omic layers along the regulatory network

of metabolic reactions and constructed a global starvation-responsive trans-omic

network (Fig. 1e).

149   Metabolic enzymes, metabolites, lipids, and FFA & acyls, are direct, major and

fine regulators of metabolism[40,50,51]. Therefore, we focused on the starvation-responsive

metabolic network, that is the sub-network of the global starvation-responsive trans-

omic network (Fig. 1d and e, dotted orange outline) to investigate structural and

temporal properties of the network. The starvation-responsive metabolic network

contains the direct regulators of metabolic reactions: metabolite, lipid, FFA & acyls,

enzyme_protein (protein abundances of metabolic enzymes), and enzyme_PTM

(phosphorylation of metabolic enzymes). We investigated structural properties of this

starvation-responsive metabolic network (Fig. 1f), the temporal order of changes in the

network (Fig. 1g), and co-regulation among molecules (Fig. 1h).

Single-omic layer analysis: identification of starvation-responsive molecules and their regulators

We measured seven omic layers: metabolome, lipidome, FFA & acyls, transcriptome, proteome, and phosphoproteome in the liver and metabolome in plasma (see Fig. S1 for representative examples and Table S1 for complete data) and performed a single-omic layer analysis in WT and *ob*/*ob* mice (Fig. 2). We conducted one-way ANOVA tests (ANOVA-like testing of edgeR[52–54] for transcriptome) in WT and *ob*/*ob* mice respectively, and defined molecules that showed a *P* value less than 0.025 as starvation-responsive molecules (Fig. 2a; Table S2), whose false discovery rates were less than 0.1 except for phosphoproteome data (Table 1). Based on the area under the curve (AUC) of the log2-transformed time courses (Fig. 2b), starvation-responsive molecules with positive AUCs were classified as increased molecules (orange), and those with negative AUCs were classified as decreased molecules (purple). Using these classifications, we defined four classes of molecules: common (changing similarly in both WT and *ob*/*ob* mice, green), WT-specific (blue), *ob*/*ob*-specific (red), and opposite (changing in opposite directions in WT and *ob*/*ob* mice, magenta) (Fig. 2c).

We identified starvation-responsive molecules in the single-omic layer analyses of metabolites (Fig. 2d), lipids (Fig. 2e; Table 2), FFA & acyls (Fig. 2f), plasma metabolites (Fig. 2g), transcripts (Fig. 2h), proteins (Fig. 2i), and phosphorylation events (Fig. 2j). The common group comprised more than one-third of the total responsive metabolites in both liver (Fig. 2d) and plasma (Fig. 2g), FFA & acyls (Fig. 2f), and transcripts (Fig. 2h). Most starvation-responsive lipids were in the *ob*/*ob*-specific group (Fig. 2e). More than half of responsive proteins (Fig. 2i) and phosphorylation events (Fig. 2j) were WT-specifically responsive.

KEGG pathway enrichment analysis (Table S3) revealed pathways that were

associated with the starvation-responsive changes in both the transcriptome (Table 3)

and proteome (Table 4), but this analysis did not find any pathways that were enriched

in the phosphoproteome data (Table S3). For example, KEGG pathways such as "fatty

acid degradation" and "PPAR signaling pathway" were enriched in both transcriptome

(Table 3) and proteome (Table 4) in WT mice.

We inferred changes in the activities of following regulators of the starvation-

responsive molecules (Fig. 2k, see Methods): TFs (Fig, 2l; Table 5; Table S4), protein

kinases (Fig. 2m; Table 6; Table S4), and signaling pathways (Fig. 2n; Table 7) based on

the regulatory network obtained from bioinformatic platforms and measured omic data.

Note that signaling pathway layer (Fig. 2n; Table 7) contains significantly enriched

signaling pathways in any of the transcriptome (Table 3) or proteome data (Table 4). We

classified the activities of these regulators as common, WT-specific, *ob*/*ob*-specific, or

opposite (Fig. 2k). The common group comprised more than half of the total responsive

TFs (Fig. 2l; Table 5), including Pparα and Hnf4α, well-established for promoting lipid

degradation[2]. Note that some results were not consistent with the previous studies such

as failure of identification of Foxo1 and unexpected identification of Creb1 as down-

regulated TFs (see Discussion). "PKC_group" kinases including Prkce, Prkcd and Prkcz,

master regulators of liver metabolism[55–57], were commonly downregulated (Fig. 2m;

Table 6). The "PPAR signaling pathway" was inferred as having up-regulated activity

(Fig. 2n; Table 7) commonly in WT and *ob*/*ob* mice.

Construction of the global starvation-responsive trans-omic network by

integrating multi-omic data

Using the starvation-responsive molecules and their regulators (Fig. 2), we constructed

a global starvation-responsive trans-omic network (Fig. 3a; Table S5). The network

consists of nodes, layers, and edges. Nodes represent measured molecules (Figs 2d-j),

inferred activities of the TFs, protein kinases, and signaling pathways (Figs. 2l, m, and

n), metabolic reactions defined using KEGG[58–60], and transport activities from TCDB[61]

(see Methods). Based on chemical properties, biological functions, or both, nodes were

clustered into layers (Table 8). For example, the enzyme_proten layer includes proteins

(chemical property) that are metabolic enzymes (biological functions). Enzyme_PTM

layer represents phosphorylation (chemical property) of metabolic enzymes (biological

functions). Metabolite, lipid, and FFA & acyls layers are assigned based on the chemical

properties of the molecules. Edges are the regulatory connections between nodes,

connected based on the regulatory relationships among omic layers (Fig. S2), using

bioinformatic platforms (see Methods). For example, edges from metabolite, lipid, and

FFA & acyls layers to metabolic reaction layer represent the relationships among

metabolites, lipids, and FFA & acyls— as substrates, products, and allosteric

regulators— to metabolic reactions. Edges from enzyme_protein and enzyme_PTM to

the metabolic reaction layer represent the regulations between metabolic enzymes and

metabolic reactions mediated by enzyme abundance (enzyme_protein) and

phosphorylation (enzyme_PTM). Colors of nodes and edges represent the type of

response: green, common in both WT and *ob*/*ob* mice; blue, specific to WT mice; red,

specific to *ob*/*ob* mice; magenta, opposite responses between WT and *ob*/*ob* mice (Fig.

3a, b).

We showed the proportions of the commonly, WT-specifically, *ob*/*ob*-

specifically, and oppositely responsive nodes and edges in the global starvation-

responsive trans-omic network (Fig 3b). In this network, pie charts represent the

proportions of the classes of responses of nodes in the omic layers, and bar plots

between pie charts represent proportions of regulatory responses of edges between omic

layers. For clear visualization, pie charts for signaling_activity and TF_activity are

displayed twice.

We assessed the proportions of the global starvation-responsive trans-omic

network (Fig. 3b). Among the starvation-responsive metabolites ("metabolite" in Fig.

3b), less than half changes (47 out of 122 nodes) were specific to WT mice, but more

than half regulatory connections between the metabolites ("metabolite" in Fig. 3b) and

the metabolic reactions ("metabolic reaction" in Fig. 3b) with 604 out of 1,008 substrate

and product relationships WT-specific and 828 out of 1,538 allosteric regulatory

relationships WT-specific. Among the starvation-responsive lipids ("lipid" in Fig. 3b),

but not the FFA & acyls ("FFA & acyls" in Fig. 3b), most changes (8 out of 10) were

specific to *ob*/*ob* mice. This enrichment in the *ob*/*ob*-specific response also occurred

among the substrate, product, and allosteric regulatory connections between the lipids

("lipid" in Fig. 3b) and the metabolic reactions ("metabolic reaction" in Fig. 3b) with 43

out of 48 substrate and product relationships *ob/ob*-specific and 160 out of 163

allosteric regulator relationships *ob/ob* specific. However, most of the starvation-

responsive metabolic reactions ("metabolic reaction" in Fig. 3b) were commonly

responsive in WT and to *ob*/*ob* mice (493 out of 702).

In the enzyme_protein layer ("enzyme_protein" in Fig. 3b), most nodes were in

the WT-specific class (130 of the 203, 64%) and only 17 of the 203 (8.4%) starvation-

responsive nodes responded in common between WT and *ob/ob* mice. However, more

than half of all the starvation-responsive nodes (587 out of 1144; 51.3%) exhibited a

common response in the enzyme_mRNA layer ("enzyme_mRNA" in Fig. 3b). This

disconnect between the responses at the mRNA and protein levels indicated that the

abundance of the metabolic enzymes is determined processes occurring after

transcription, such as the rate of translation and protein degradation, that are known to

be downregulated in *ob*/*ob* mice[62,63]. Among most of the nodes and edges related to

phosphorylation ("kinase_protein", "kinase_PTM", "kinase_activity", and

"enzyme_PTM" in Fig. 3b), WT-specific groups occurred most often compared to the

common, *ob*/*ob*-specific, and opposite groups. These results suggested a decrease in

regulation by phosphorylation in response to starvation in *ob*/*ob* mice.

Compared with nodes in the other classes of responses, nodes with common

responses were most abundant in several layers related to inter-organ metabolic cycles

("plasma metabolite", "transporter_mRNA", "transporter_protein", and

"transport_activity" in Fig. 3b).

Taken together, the regulatory connections from metabolite, lipid,

enzyme_protein, and enzyme_PTM layers to metabolic reactions were different

between WT and *ob/ob* mice. The inter-organ metabolic cycle was maintained.

Structural properties of the starvation-responsive metabolic network

We focused on the sub-network of the global starvation-responsive trans-omic network

that directly regulate metabolic reactions, defined as the starvation-responsive metabolic

network (Fig. 4). This network contains distinct regulatory connections in the global

starvation-responsive trans-omic network between WT and *ob*/*ob* mice (Fig. 3b). To

visualize the differences between WT and *ob/ob* mice, we show the starvation-

responsive metabolic network separately for WT (Fig. 4a) and *ob/ob* (Fig. 4b) mice. For

each genotype, the networks include the layers for metabolite, lipid, FFA & acyls,

enzyme_protein, and enzyme_PTM (Table S6) and their relationship to the metabolic

reaction layer (Fig. S2a). The edges connecting metabolite, lipid, and FFA & acyls

layers to the metabolic reaction layer represent their roles as substrates, products, and

allosteric regulators of the metabolic reactions.

We analyzed the structural properties of the starvation-responsive metabolic

networks in WT and *ob/ob* mice (Fig. 5). We calculated the number of edges connected

to each node (called "degree") and then assessed the degree distribution of the network

(Fig. 5a), by a log-log scatter plot. We found that nodes with large degree were rare,

while nodes with small degree were abundant. This type of network is called a scale-

free network, and nodes with large degree are called "hub", while nodes with small

degree are called "leaves" (Fig. 5b). Degree distribution of a scale-free network can be

approximated by a power-law[35,46,64] defined as:

$$n(k) \propto k^{-\gamma},$$

where $n(k)$ denotes the number of nodes with a degree of k, and γ is a constant called

the scaling parameter. The scaling parameter represents the heterogeneity of the degree

distribution[65] with a small scaling parameter representing a large heterogeneity of

degree distribution of the network. From the calculated scaling parameter for each

genotype's network, we found that the starvation-responsive network for WT and *ob/ob*

mice had a similar amount of heterogeneity in the degree distribution (Fig. 5a). Note

that scaling parameters should be $2 < \gamma < 3$ according to the strict definition[64,66], but

we call this type of network as a scale-free network in this manuscript for simplicity.

We defined the nodes that had a degree in the top 2% among the responsive

nodes in each genotype as hub nodes (Fig. 5c and d), that regulated multiple metabolic

pathways (Fig. S3). We identified 20 hub nodes in the network of WT mice and 15 hub

nodes in the network of *ob/ob* mice (Fig. 5c, d and labeled on Fig. 4a, b). In WT mice,

15 of the 20 were WT-specific (Fig. 4a; Fig. 5c). In *ob/ob* mice, 7 hub nodes out of 15

were *ob/ob*-specific and 3 of the 7 were lipids (Fig. 4b; Fig. 5d). The hub nodes with the

highest degrees were ATP (degree >200) and AMP (degree >100) in the network of WT

mice (Fig. 5c); whereas all hub nodes in the network of *ob/ob* mice had degrees less

than 100 with reduced glutathione (degree 81) and NADH (degree 69) (Fig. 5d). Thus,

the starvation-responsive metabolic network in *ob/ob* mice lacked the most highly

connected hub nodes such as ATP and AMP.

To evaluate how the loss of responsiveness of the highly connected hub nodes

in *ob/ob* mice, we analyzed the structural properties of the starvation-responsive

metabolic network (see method in detail). Cluster is a set of nodes connected through

edges regardless of the distances. The largest cluster is the cluster that includes the

largest number of nodes in a network. Fragmentation breaks the network into isolated

clusters and reduce the relative size of the largest cluster (Fig. 5e), that is the number of

nodes in the largest cluster divided by the total number of the responsive nodes in each

genotype. Loss of communicability among nodes, which increases the average number

of edges connecting two nodes, defined as average path length (Fig. 5f)[67]. In the

starvation-responsive metabolic networks of WT and *ob*/*ob* mice, the relative sizes of

the largest cluster were similar (Fig. 5g), indicating that the network was not

fragmentated in *ob*/*ob* mice. Additionally, the average path lengths were similar in the

networks of the WT and *ob*/*ob* mice (Fig. 5h), indicating that communicability was

maintained in *ob*/*ob* mice. Collectively, all structural properties of the starvation-

responsive metabolic networks, the scaling parameter (Fig. 5b), the relative size of the

largest cluster (Fig. 5g), and the average path length (Fig. 5h) were similar between WT

and *ob*/*ob* mice. These results indicated that the starvation-responsive metabolic

network was structurally maintained in *ob*/*ob* mice, despite the loss of responsiveness of

highly connected hub nodes such as ATP and AMP.

The maintenance of the structural properties of the starvation-responsive

metabolic network in *ob*/*ob* mice seems counter-intuitive because it has been known

that the loss of hub nodes in a scale-free network drastically alter the structural

properties[67]. Taking a close look at the degree distribution (Fig. 5a, S4), the number (y-

axis of Fig. 5a) of nodes with small degree (log10-transformed degree < 0.5 in the x-

axis of Fig. 5a) were smaller in *ob*/*ob* mice than in WT mice, but the number and of

nodes (y-axis of Fig. 5a) with log10-transformed degree ranging from 0.5 to 2 (x-axis of

Fig. 5a) were comparable between WT and *ob*/*ob* mice (Fig. 5a, see Fig. S4 in detail).

Totally, the percentages of nodes with relatively large degrees (log10-transformed

degree > 1) were similar between WT and *ob*/*ob* mice (Fig. S4). The maintenance of

nodes with relatively large degrees could maintain the structural properties of the

starvation-responsive metabolic network in *ob*/*ob* mice.

Temporal order of the starvation-responsive metabolic network in WT mice and its

disruption in *ob/ob* mice

Adaptation to starvation is a temporally well-ordered metabolic process[1,4–6]. To explore

the relationships between the network structure and the temporal order of the starvation-

induced changes in the starvation-responsive metabolic network, we defined the time

when the response reached half of the maximum amplitude of the responses as *t*-half [40].

A small *t*-half represents rapid responsiveness, and a large *t*-half represents slow

responsiveness. We analyzed molecules [metabolites, lipids, FFA & acyls, proteins

(enzyme_protein), and phosphorylation events (enzyme_PTM)] in the starvation-

responsive metabolic networks, because *t*-half can be defined only for molecules not for

metabolic reactions. We refer to hub nodes as hub molecules because all the hub nodes

were molecules.

   We investigated the relationships between network structure and *t*-half in WT

and *ob/ob* mice. We found a negative correlation between *t*-half and degree of

molecules in WT mice (Fig. 6a), indicating that the molecules connected with many

edges respond quickly to starvation. In addition, 9 out of 20 hub molecules, including

ATP, were essential for the negative correlation with *t*-half in WT mice (Fig. S5, see

Methods), suggesting a dominant role of hub molecules in coordinating the temporal

order of the starvation-responsive metabolic network. By contrast, no significant

correlation between *t*-half and degree of molecules was observed in *ob/ob* mice (Fig.

6b). The *t*-half values of *ob/ob*-specific hub molecules were larger than those of WT-

specific hub molecules (Fig. 6c), indicating a slower response of the *ob/ob*-specific hub

molecules.

Glycogen, carbohydrates, amino acids, and lipids all serve as energy sources

during starvation in a temporally ordered manner[1,4–6]. We used *t*-half analysis to

evaluate how the temporal order of use of these various sources of cellular energy

differs during starvation in WT and *ob/ob* mice. We used the KEGG database[58–60] to

classify molecules into the following metabolic pathways: "Starch and sucrose

metabolism" (containing glycogen and glycogen metabolic enzymes),

"Glycolysis/gluconeogenesis", "Arginine biosynthesis" (containing the urea cycle that

could indicate amino acid degradation), and "Fatty acid degradation". We calculated *t*-

half values for the molecules in each of the metabolic pathways for the starvation-

responsive metabolic network in WT (Fig. 6d) and *ob*/*ob* mice (Fig. 6e). In WT mice,

we observed significant differences in *t*-half values of nodes among the investigated

metabolic pathways (p<0.001, Kruskal-Wallis test; Fig. 6d), indicating that the

metabolic processes are temporally ordered across the metabolic pathways in WT mice.

Among them, "Starch and sucrose metabolism" responded to starvation the fastest, and

"Fatty acid degradation" responded to starvation the slowest, which is consistent with

glycogen as metabolic fuel for the acute phase of starvation and fatty acids as fuel for

the late stage of starvation[1,4–6]. By contrast, no significant differences in *t*-half values

were observed in *ob*/*ob* mice (p=0.495, Kruskal-Wallis test; Fig. 6e), indicating that the

temporal order among the metabolic pathways was disrupted in *ob*/*ob* mice.

In addition, we compared *t*-half values of molecules in each omic layer in the

starvation-responsive metabolic network in WT (Fig. 6f) and *ob*/*ob* mice (Fig. 6g). In

WT mice, we observed significant differences in *t*-half values of molecules among the

omics layers (p<0.001, Kruskal-Wallis test; Fig. 6f), indicating that adaptation to

starvation is temporally ordered across omic layers in WT mice. Metabolites responded

to starvation fastest, indicating that metabolic reactions are initially regulated by

metabolites through the effect of starvation-induced changes in substrates and products

and metabolite-mediated allosteric regulators. Phosphorylation (enzyme_PTM), protein

abundance (enzyme_protein), and lipids responded next. The FFA & acyls layer

responded the slowest, consistent with fatty acid degradation important for the late

stages of adaptation to starvation[5]. By contrast, no significant differences in t-half

values were observed in *ob*/*ob* mice (p=0.197, Kruskal-Wallis test; Fig. 6g), indicating

that the temporal order of the starvation response by the omic layers was disrupted.

398   Taken together, these results indicated that hub molecules coordinate the

temporal order of the responses of molecules, metabolic pathways, and omic layers in

the starvation-responsive metabolic network in WT mice. In *ob*/*ob* mice, the temporal

order of the starvation response throughout the network was disrupted.

Co-regulation among molecules in the starvation-responsive metabolic network

and disruption in *ob*/*ob* mice

Molecules regulated by common regulatory mechanisms show positively or negatively

correlated time courses, defined as co-regulation (Fig. 7a). Thus, positively or

negatively correlated time courses represent co-regulations, which could be disrupted by

the disrupted temporal order of metabolites, lipids, FFA & acyls, protein abundance, and

phosphorylation events in *ob*/*ob* mice (Fig. 6). As an index of temporal co-regulation

(Fig. 7a), we calculated the pairwise Pearson's correlation coefficients for time courses

of each pair of molecules (including phosphorylation events) in the starvation-

responsive metabolic network in WT and *ob*/*ob* mice, respectively (Fig. 7b). The

distribution of correlation coefficients in WT mice was bimodal with peaks at around

0.75 and -0.75, indicating both positive or negative co-regulation occurred among

molecules. By contrast, the distribution of correlation coefficients was unimodal with a

broad peak around 0 in *ob*/*ob* mice, indicating that co-regulation was disrupted in *ob*/*ob*

mice.

To obtain a qualitative landscape of co-regulations at a molecular level, we

constructed networks of molecules that exhibited strong co-regulations (the absolute

value of Pearson's correlation coefficient > 0.9, see Fig. S6) among metabolites, lipids,

FFA & acyls, enzyme_protein, and enzyme_PTM in WT mice (Fig. 7c) and *ob/ob* mice

(Fig. 7d), named as the correlation metabolic network. Note that the correlation

metabolic network is data-dependent and the network structure highly depends on the

threshold for the absolute value of correlation coefficient (Fig. S6).

The correlation metabolic network in WT mice included a single large cluster

comprised of 275 molecules, representing molecules with various biological functions

such as carbohydrate metabolism and fatty acid degradation (Fig. 7c), indicating that

most of the responsive molecules in WT mice are co-regulated. ATP and AMP, the rapid

hub molecules in the starvation-responsive metabolic network (Fig. 5 and 6) were

included in the large cluster, suggesting that ATP and AMP formed this large cluster by

regulating the large number of metabolic reactions. By contrast, the network in *ob/ob*

mice was fragmentated with some clusters containing functionally related molecules

(Fig. 7d). Taken together, temporal co-regulation of metabolism in response to

starvation was disrupted in the *ob/ob* mice.

Discussion

We measured multi-omic data during starvation in WT and *ob*/*ob* mice (Fig. 2) and

constructed a global starvation-responsive trans-omic networks by integrating these data

(Fig. 3). We also showed that the starvation-responsive metabolic network (Fig. 4), a

sub network of the global starvation-responsive trans-omic network, is a scale-free

network. The hub molecules of the networks, regulating many metabolic reactions, were

energy indicators, such as ATP and AMP in WT mice, and less highly connected hub

molecules, such as lipids in *ob*/*ob* mice (Fig. 5). Despite the loss of the responsiveness

of the hub energy indicators in *ob*/*ob* mice, structural properties of the network in *ob*/*ob*

mice were maintained (Fig. 5), likely through the maintenance of nodes with relatively

large degrees (Fig. S4). By contrast, the temporal properties including the temporal

order (Fig. 6) and the temporal co-regulation of molecules were disrupted in *ob*/*ob* mice

(Fig. 7).

The maintenance of the structural properties of the starvation-responsive

metabolic network in *ob*/*ob* mice (Fig. 5) suggests the robustness of the metabolic

networks against pathological perturbations. Robustness of the structural properties of

metabolic networks have also been investigated in cancers[44,45]. Structural properties of

metabolic networks based on the tissue-specific gene expression were similar between

15 cancers and corresponding normal tissues[44]. Metabolic network based on flux

correlations showed similar structural properties between normal and cancer tissues[45].

These two studies suggest that structural properties of metabolic networks in cancer are

similar to those of normal tissues, despite profound differences in gene expression

patterns of cancer and non-cancer cells[68]. These studies, together with our results,

indicate that structural properties of metabolic networks are robust among various

metabolic perturbations including starvation in healthy or obese conditions and cancers.

By contrast to the robust structural properties (Fig. 5), temporal properties were

maintained depending on the hub molecules including energy indicators such as ATP

and AMP in WT mice, and disrupted in *ob*/*ob* mice (Figs. 6 and 7). These results

highlight the significance of the highly connected hub molecules for temporal properties

of the starvation-responsive metabolic networks. If a small number of hub molecules

regulate the entire network, nodes in the network would be uniformly regulated,

resulting in a temporally ordered and co-regulated network. Thus, the starvation-

responsive metabolic network is temporally disrupted in *ob*/*ob* mice, that lost the

responsiveness of the highly connected hub molecules such as ATP and AMP.

If the metabolic network is temporally vulnerable, the structural robustness

observed in starvation (several hours) and cancers (~years) could be disrupted in a rapid

metabolic shift such as glucose ingestion because of the lack of time for rescuing the

structural properties. Previously, we conducted a trans-omic analysis of liver in WT and

ob/*ob* mice during oral glucose ingestion, in which we measured metabolome, lipidome,

and transcriptome[40,42]. Small fraction of the responsive metabolites (15 out of 88,

17.0%) and transcripts (732 out of 5,091, 14.4%) was common between WT and *ob*/*ob*

mice during oral glucose ingestion, while more than one-third of the responsive

molecules were common during starvation (Fig. 2)[40]. Scaling parameters of the trans-

omic networks, representing the heterogeneity of the network, were different during oral

glucose ingestion but similar during starvation between WT and *ob*/*ob* mice[42]. Taken

together, glucose ingestion requires distinct responsive molecules and heterogeneities of

trans-omic network between WT and *ob*/*ob* mice, suggesting metabolic network is more

483 vulnerable to temporally rapid perturbations (oral glucose ingestion) than slow

484 (starvation) or chronic (cancer) perturbations.

485  By evaluating the temporal order among omic layers during starvation (Fig. 6),

486 we found that metabolite layer was the fastest even compared to the enzyme_PTM layer,

487 which is composed of protein phosphorylation events (Fig. 6f). Our previous study

488 showed that rapid metabolic regulation by metabolites also occurs in the liver of WT

489 mice during oral glucose ingestion[40]. However, we did not measure the

490 phosphoproteome in that study[40], thus we cannot ascertain if metabolite regulation

491 precedes phosphorylation-mediated regulation. Because here we measured both the

492 metabolome and phosphoproteome, our data indicated that metabolites take the

493 initiating role in controlling the trans-omic layers during starvation in liver.

494  One of the limitations of this study is the accuracy of the inference of regulator

495 activities, including TFs and protein kinases. For inference of TF activity, we failed to

496 identify Foxo1, which is master regulator of liver metabolism and is necessary for

497 upregulation of glucose production during starvation[69]. We can explain this discrepancy

498 because we removed data for Foxo1 in liver from the data obtained from the ChIP-Atlas

499 database[70,71] as not meeting our quality check process for inference input (see Methods).

500 Another example, Creb1, which has been known to be upregulated during starvation[2],

501 was classified as a down-regulated TF (Table 5), probably due to the failure of detection

502 in phosphorylation, that is important for Creb1 activity[2]. For kinase inference, we failed

503 both mTorc1 and Ampk as responsive protein kinases, both of which are important

504 during starvation[72,73]. At present, protein kinases were identified for less than 5% of

505 phosphorylation events[74], thus the prediction algorithm could be inaccurate. These

limitations could be solved by integrating recently developed tool such as CollectTRI-

based database[75], The Kinase Library[76], and OmniPath[77] to trans-omic analyses.

Another limitation of our study is that we only used one model of obesity. By

only using *ob*/*ob* mice, we cannot exclude the possibility that the results are caused by

the effect of leptin deficiency[78]. We compared our results with the previous study using

high fat diet (HFD) as a model for obesity[30]. We observed similar changes in metabolic

enzymes and in glycogen during starvation in WT and *ob/ob* mice (Fig. S1) as were

reported for both lean and HFD mice[30]. Thus, our results are related to obesity not

leptin-deficiency itself.

In addition, our experimental condition is not optimized for calorie-restricted

diet, because we focused on adaptation to starvation. Considering future therapeutic

application of our results, another study targeting calorie-restricted diet is necessary.

Despite several limitations, our present study successfully described the global

landscape of adaptation to starvation in WT and *ob*/*ob* mice and suggested that the

highly connected hub energy indicators temporally coordinate the starvation-responsive

metabolic network in the healthy condition and that these critical hubs are no longer

starvation-responsive in obesity, resulting in temporal disruption of the starvation-

responsive metabolic network. Robustness of the structures against pathological

perturbations could be a general nature of metabolic networks, combining with the

previous studies in cancer[44,45]. Vulnerability of temporal properties of the network in

ob/*ob* mice could be helpful to establish an efficient protocol for calorie-restricted diet,

whose major therapeutic target is obesity.

 FIGURE TITLES AND LEGENDS

Fig. 1: Workflow for the construction and analysis of a global starvation-

responsive trans-omic network in liver of WT and *ob*/*ob* mice

(a) Experimental design. WT and *ob*/*ob* mice were starved for the indicated hours,

followed by the collection of liver and plasma samples. These samples were subjected

to multi-omic measurements. It should be noted that metabolome was basically

measured by CE-MS, but IC-QEMS was used for quantification of F1,6BP and F2,6BP.

Enzymatic assay kit was used for glycogen. (b) Classification of molecules into 4

groups: commonly responsive in both WT and *ob*/*ob* mice (common, green); WT-

specifically (WT-specific, blue); *ob*/*ob*-specifically (*ob/ob*-specific, red); oppositely

(opposite, magenta) responsive molecules between WT mice and *ob*/*ob* mice. (c)

Schema for a single-omic layer composed of responsive molecules classified as in (b).

(d) Schema of regulatory input into the metabolic reactions. TFs control metabolic

enzyme abundance through gene regulation. Protein kinases control metabolic enzyme

activities through phosphorylation. The metabolites, FFA & acyls, and lipids act as

substrates, products, and allosteric regulators in metabolic reactions. (e) Schematic

representation of the global starvation-responsive trans-omic network consisting of the

indicated layers. The subset of the network that is defined as the starvation-responsive

metabolic network is outlined in orange. (f) Structural properties of the starvation-

responsive metabolic network in WT and *ob/ob* mice. (g) Temporal order of the

starvation-responsive metabolic network in WT and *ob*/*ob* mice. (h) Strength of co-

regulation among molecules in the starvation-responsive metabolic network in WT and

ob/*ob* mice.

Fig. 2: Single-omic layer analysis during starvation

(a-c) Identification and classification of starvation-responsive molecules. (a) One-way

ANOVA tests ($P < 0.025$) defined starvation-responsive molecules. (b) Area under the

curve (AUC) of log2-transformed mean time courses classified molecules into increased

(AUC > 0, orange) and decreased (AUC < 0, purple) classes. (c) Classification of

starvation-responsive molecules: common in both WT and *ob*/*ob* (either increased or

decreased in both, green), WT-specific (blue), *ob*/*ob*-specific (red), or opposite

(increased in WT and decreased in *ob*/*ob*, or vice versa, magenta). (d-j) The number of

molecules and distribution in each group shown as pie charts colored as defined in (c).

Increased or decreased molecules are indicated by the orange or purple arcs,

respectively. The number and the ratio of significantly changed molecules of the

molecules in each category are shown. Responsive metabolites (d), lipids (e), and FFA

& acyls (f) in liver. Responsive plasma metabolites (g). Responsive transcripts (h),

proteins (i), and phosphorylation events (j) in liver. (k) Strategy for inferring activity of

upstream regulators and signaling pathways. Briefly, we constructed regulatory

networks from database and identified increased and decreased downstream molecules.

In the case of positive regulators, we assumed that regulator activity is up-regulated if

most of the downstream molecules increased and vice versa (see method in detail). Up-

regulated or down-regulated regulators were inferred and classified them into common

(green), WT-specific (blue), *ob*/*ob*-specific (red), or opposite (magenta) groups. (l-n)

The number and distribution of these inferred TFs (l), protein kinases (m), or signaling

pathways (n) are shown in pie charts.

Fig. 3: Global starvation-responsive trans-omic network.

(a) Global starvation-responsive trans-omic network. Nodes in the layers represent

metabolic reactions, transport activities, measured molecules, measured

phosphorylation events, and the inferred activities of TF, kinase, and signaling pathway.

Edges between the nodes represent regulatory connections identified using

bioinformatic platforms. Colors represent the responsiveness: green indicates common

responses both in WT and *ob/ob* mice; blue, specific to WT mice; red, specific to *ob/ob*

mice; magenta, opposite responses between the WT and *ob/ob* mice. For metabolic

reaction and transport_activity layers, the classes were defined by the existences of

regulations regardless of the regulatory directions (activation or inhibition): green

indicates the node is regulated in both WT and *ob/ob* mice; blue, specific to WT mice;

red, specific to *ob/ob* mice. (b) The number and proportions of nodes and edges

classified as common (green), WT-specific (blue), *ob/ob*-specific (red), and opposite

(magenta) in the global starvation-responsive trans-omic network. Pie charts represent

the distribution of nodes in the omic layers, and bar plots between pie charts represent

the distribution of edges between omic layers. Type of regulation into the metabolic

reactions for metabolite, FFA& acyls, and lipid are indicated as allosteric regulation (A)

and substrate and product (S & P). The starvation-responsive metabolic network is

outlined in orange, and layer names included in the starvation-responsive metabolic

networks are bolded.

Fig. 4: Starvation-responsive metabolic network

Projected views of the starvation-responsive metabolic networks in WT (a) and *ob/ob*

mice (b). The network is composed of direct regulators of metabolic reactions:

metabolites, lipids, FFA & acyls, enzyme_proteins, and enzyme_PTMs. Colors

represent the responsiveness: green indicates common responses both in WT and *ob*/*ob*

mice; blue, specific to WT mice; red, specific to *ob*/*ob* mice; magenta, opposite

responses between the two genotypes. Hub nodes, which will be defined in Fig. 5, are

labeled.

Fig. 5: Structural properties of the starvation-responsive metabolic network

(a) Degree distributions of the starvation-responsive metabolic network in WT (blue)

and *ob*/*ob* (red) mice. The x-axis represents the log10-transformed degree (k); the y-axis

represents the log10-transformed number of nodes with degree of k. Lines show the

linear regression of the degree distributions. The slope of the lines (γ), which will be

defined as scaling parameters, and the 95% credible intervals are shown. (b) Diagram of

a scale-free network, where $n(k)$ denotes the number of nodes with degree of k, and γ

is a constant called a scaling parameter. Circles represent nodes; lines represent edges.

Light blue represents the hub nodes; yellow represents leaf nodes. (c and d) Hub nodes

with the top 2% highest degree in WT (c) and *ob*/*ob* mice (d). The x-axis represents hub

nodes. The y-axis represents degrees. Colors represent the responsiveness: green

indicates common responses both in WT and *ob*/*ob* mice; blue, specific to WT mice; red,

specific to *ob*/*ob* mice; magenta, opposite responses between WT and *ob*/*ob* mice. (e)

Illustration of fragmentation of a network through loss of hub nodes. The largest cluster

is orange. (f) Illustration of how communicability of a network influences average path

length (average length of the shortest paths between all the pairs of nodes). (g) The

relative size of the largest cluster of the starvation-responsive metabolic network in WT

and *ob*/*ob* mice, calculated as the number of nodes in the largest cluster of the network

divided by the total number of responsive nodes in the network for each genotype. (h)

The average path length of the starvation-responsive metabolic network in WT and

ob/*ob* mice.

Fig. 6: Temporal order of the starvation-responsive metabolic network in WT mice

and its disruption in *ob*/*ob* mice

(a and b) Scatter plot for the log10-transformed degree (x-axis) and *t*-half (y-axis) in

WT (a) and *ob*/*ob* (b) mice. Dots represent the molecules (including phosphorylation

events) in the starvation-responsive metabolic network and colors denote the

responsiveness: green indicates common responses both in WT and *ob*/*ob* mice; blue,

specific to WT mice; red, specific to *ob*/*ob* mice; magenta, opposite responses between

WT and *ob*/*ob* mice. Spearman's correlation coefficient (Rho) and *P* values are shown.

Hub molecules for each genotype are labeled (see Fig. 5c and d). (c) Values of *t*-half for

the WT-specific and *ob*/*ob*-specific hub molecules (see Fig. 5c and d). The *P* value of

Wilcoxon's rank sum test is shown. (d and e) Distributions of *t*-half values of molecules

(metabolites, lipids, FFA & acyls, proteins, and phosphorylation events) in the indicated

metabolic pathways in WT (d) and *ob*/*ob* mice (e). The data are ordered along the x-axis

by the median *t*-half (red line) with each molecule represented by a dot. *P* values of the

Kruskal-Wallis test are shown above the panels and those of the Dunn-Sidak test are

shown above the lines indicating the pairs of metabolic pathways with significant

differences. (f and g) Distributions of *t*-half values in each omic layer in the starvation-

responsive metabolic network in WT (f) and in *ob*/*ob* (g) mice. The data are ordered

along the x-axis by the median *t*-half (red line). *P* values of the Kruskal-Wallis test are

shown above the panels and those of the Dunn-Sidak test are shown above the lines

indicating the pairs of omic layers with significant differences.

Fig. 7: Co-regulation among molecules in the starvation-responsive metabolic

network and disruption in *ob*/*ob* mice

(a) Schematic representation of the analyses for co-regulation of the molecules in the

starvation-responsive metabolic network. Time courses of molecules sharing a common

regulatory mechanism could be positively or negatively correlated, defined as co-

regulation. Pairwise Pearson's correlation coefficients were calculated for the time

courses of molecules that showed significant responses during starvation in the

starvation-responsive metabolic network in WT and *ob*/*ob* mice. Positive correlation

represents positive co-regulation and negative correlation represents negative co-

regulation. (b) Distribution of the pairwise Pearson's correlation coefficients for time

courses of responsive molecules in the starvation-responsive metabolic network in WT

and *ob*/*ob* mice. The x-axis represents correlation coefficients; the y-axis represents the

number of pairs of molecules examined for correlation. Pairs with an absolute value of

correlation coefficient > 0.9 were connected by edges in the correlation metabolic

network (see Fig. S6). (c and d) The correlation metabolic network during starvation in

WT (c) and *ob*/*ob* mice (d). Orange indicates the largest cluster; all other clusters are

dark green. Individual orphan nodes with no connections are shown only by numbers.

 TABLE TITLES AND LEGENDS

 Table 1: Maximum Q-values for the omic layers of the starvation-responsive

 molecules.

Maximum Q-values for each omic layer for molecules with P-values that were less than

0.025. All the maximum Q-values were less than 0.1 except for phosphoproteome,

suggesting that false discovery rates of the starvation-responsive molecules were less

than 0.1 in all the omic layers except for phosphoproteome.

Table 2: Lipid classes measured.

Measured lipid classes and their abbreviations.

Table 3: KEGG pathway enrichment analysis of the starvation-responsive

transcripts

KEGG pathways with Q-value < 0.1 are shown.

Table 4: KEGG pathway enrichment analysis of the starvation-responsive proteins

KEGG pathways with Q-value < 0.1 are shown.

Table 5: Up- and down-regulated TFs during starvation

Up/Down-regulated TFs during starvation. 'Up', 'Down', and 'NE' represent up-

regulation, down-regulation, and no enrichment, respectively.

Table 6: Up- and down-regulated protein kinase groups during starvation

Up/Down-regulated protein kinase groups during starvation. 'Up', 'Down', and 'NE'

represent up-regulation, down-regulation, and no enrichment, respectively.

Table 7: Up- and down-regulated signaling pathways during starvation

Up/Down-regulated signaling pathways during starvation. 'Up', 'Down', and 'NE'

represent up-regulation, down-regulation, and no enrichment, respectively.

Table 8: Omic layers in the global starvation-responsive metabolic network

Classification of molecules or inferred regulators into omic layers, biological function,

or chemical property.

Materials and Methods

Mouse experiments

All the animal experiments were conducted after approval by the animal ethics

committee of the University of Tokyo. Ten-week-old C57BL/6 WT and *ob/ob* male

mice were purchased from Japan SLC Inc. Mice were starved for 0, 2, 4, 6, 8, 12, 16, 24

hours, starting at approximately 18:00. For omic measurement, mice were euthanized

by cervical dislocation, blood was collected from retro-orbital sinus into tubes

containing 0.5 mg of EDTA, and liver was dissected and immediately frozen in liquid

nitrogen. Blood was centrifuged at $2,300 \times g$ for 15 min at 4°C to separate plasma.

Plasma was frozen in liquid nitrogen and stored at -80°C. The frozen liver was crushed

with dry ice to a fine powder with a Wonder Blender (WB-1, OSAKA CHEMICAL CO.,

LTD). Blood glucose was determined using ACCU-CHECK (Roche).

Metabolomic analysis

Metabolomic analysis was conducted as previously described[40]. For liver, total

metabolites were extracted with methanol:chroloform:water (2.5:2.5:1) along with total

proteins. Approximately 40 mg of liver was homogenized in 500 µl of ice-cold

methanol containing internal standards [20 µM l-methionine sulfone (Wako), 2-

morpholinoethanesulfonic acid, monohydrate (Dojindo), and d-camphor-10-sulfonic

acid (Wako)] for normalization of peak intensities of mass spectrometry (MS) among

runs. Next, 500 µL of chloroform and then 200 µL of water were added and mixed.

After centrifugation at $4,600 \times g$ for 15 min at 4°C, the aqueous layer was filtered

through a 5 kDa cutoff filter (UFC3LCCNB-HMT, Human Metabolome Technologies)

to remove protein contamination. Simultaneously, proteins were precipitated from the

interphase and organic layers by addition of 800 μL of ice-cold methanol followed by

centrifugation at $12,000 \times g$ for 15 min at 4ºC. The pellet was washed with 1 mL of ice-

cold 80% (v/v) methanol and resuspended in 500 μL water, followed by sonication with

Bioruptor (UCW-310, COSMO BIO). By addition of 500 μL of 2% SDS and 100 mM

Tris-HCl pH8.8, the pellet was lysed for 1 hour at 4°C with gentle mixing followed by

sonication using Handy Sonic (UR-20P, TOMY). The concentration of total protein was

determined with the bicinchoninic acid (BCA) assay (23227, Thermo Fisher Scientific)

and used for normalization of concentration among samples.

For plasma, 40 μL of the samples were used to extract metabolites with

methanol:chloroform:water (2.5:2.5:1). The samples were dissolved in 400 μL of ice-

cold methanol containing the internal standards [20 μM l-methionine sulfone (Wako), 2-

morpholinoethanesulfonic acid, monohydrate (Dojindo), and d-camphor-10-sulfonic

acid (Wako)], followed by addition of 400 μL of chloroform, and then 120 μL of water.

After centrifugation at $10,000 \times g$ for 3 min at 4 ºC, the aqueous layer was filtered

through a 5 kDa cutoff filter (UFC3LCCNB-HMT, Human Metabolome Technologies)

to remove protein contamination.

The filtrate was lyophilized and suspended in 50 μL of water containing the

reference compounds [200 μM each of trimesate (206-03641, Wako) and 3-

aminopyrrolidine (404624, Sigma-Aldrich)]. Metabolites were quantified using an

Agilent 1600 Capillary Electrophoresis system (Agilent technologies), a G1603A

Agilent CE-MS adapter kit, and a G1607A Agilent CE electrospray ionization (ESI)–

MS sprayer kit. For cationic compounds, a fused silica capillary [50 μm internal

diameter (i.d.) $\times$ 100 cm] was used with 1 M formic acid as the electrolyte[79].

Methanol/water (50%, v/v) containing 0.01 μM hexakis (2,2-difluoroethoxy)

phosphazene was used as sheath liquid and delivered at 10 µl/min. ESI–time-of-flight

(TOF) MS was conducted in positive ion mode, and the capillary voltage was set to 4

kV. Each acquired spectrum was recalibrated automatically using the masses of the

reference standards [^{13}C isotopic ion of a protonated methanol dimer (2 MeOH + H)]$^{+}$,

mass/charge ratio (m/z) 66.0631 and [hexakis(2,2-difluoroethoxy)phosphazene + H]$^{+}$,

m/z 622.0290. For identification of metabolites, relative migration times of all peaks to

the reference compound 3-aminopyrrolidine were calculated, followed by comparison

of their m/z values and relative migration times to the metabolite standards. For

quantification of the metabolites, peak areas were compared to calibration curves

generated using internal standardization techniques with methionine sulfone. The other

parameters and conditions were the same as previously described[80]. For anionic

metabolites, COSMO (+) (chemically coated with cationic polymer) capillary (50 µm

i.d. by 105 cm) (Nacalai Tesque, Kyoto, Japan) was used with a 50 mM ammonium

acetate solution (pH 8.5) as the electrolyte. Methanol/5 mM ammonium acetate (50%,

v/v) containing 0.01 µM hexakis (2,2-difluoroethoxy) phosphazene was used as sheath

liquid and delivered at 10 µl/min. ESI-TOFMS was conducted in negative ion mode,

and the capillary voltage was set to 3.5 kV. Trimesate and d-camphor-10-sulfonic acid

were used as the reference and the internal standards, respectively. The other parameters

and conditions were same as described previously[81]. The Agilent MassHunter software

(Agilent technologies) was used for data analysis[79–81].

F1,6P and F2,6P were measured separately using IC-QEMS[82]. A Dionex

IonPac AS11-HC-4 µm column (250 × 0.4 mm, 4 µm; Thermo Fisher Scientific) was

used at 35°C to separate metabolite. We used KOH as an eluent at the speed of 0.02

mL/min. The gradient was as follows: 1 mM from 0 to 2 min, 20 mM at 16 min, 100

mM at 35 min. As a sheath solution, isopropanol containing 0.1% acetic acid was used

at the speed of 5 µl/min. MS was performed in the ESI negative-ion mode. The

parameters were as follows: sheath gas, 20 (arbitrary units); auxiliary gas, 10 (arbitrary

units); spray voltage, 4.0 kV; capillary temperature, 300°C; S-lens, 50 (arbitrary units).

Data acquisition was conducted in full MS scan mode. Parameters of the scanning were

as follows: resolution, 70,000; auto-gain control target, 3×10^6; maximum ion injection

time, 100 ms; scan range, 70–1,000 m/z.

Lipidomic analysis

Liver samples were prepared for lipid extraction using the Bligh and Dyer's method[83]

with minor modifications. Briefly, the lipids and their related metabolites were extracted

from frozen and crushed liver tissues (~50 mg) with 1 ml of cold methanol. The samples

were vigorously mixed for 1 min and sonicated for 5 min. The extracts were then

centrifuged at 16,000 $\times g$ for 5 min at 4°C, and the resultant supernatant was collected.

Protein concentrations in the pellet were determined using the bicinchoninic acid (BCA)

assay (23227, Thermo Fisher Scientific). The collected supernatant (180 µL) was mixed

with 200 µL of chloroform, 150 µL of water, 10 µL of internal standard (IS) A (Mouse

SPLASH Lipidomix Mass Spec Standard, Avanti Polar Lipids Inc., Alabaster, AL,

USA) containing 1.0 nmol phosphatidylcholine (PC) 15:0–18:1 (d_7), 0.070 nmol

phosphatidylethanolamine (PE) 15:0–18:1 (d_7), 0.20 nmol phosphatidylserine (PS)

15:0–18:1 (d_7), 0.050 nmol phosphatidylglycerol (PG) 15:0–18:1 (d_7), 0.20 nmol

phosphatidylinositol (PI) 15:0–18:1 (d_7), 0.10 nmol phosphatidic acid (PA) 15:0–18:1

(d_7), 0.45 nmol lysophosphatidylcholine (LPC) 18:1 (d_7), 0.020 nmol

lysophosphatidylethanolamine (LPE) 18:1 (d_7), 2.5 nmol cholesteryl ester (CE) 18:1

(d$_7$), 0.15 nmol diacylglycerol (DG) 15:0–18:1 (d$_7$), 0.35 nmol triacylglycerol (TG)

15:0–18:1 (d$_7$)–15:0, and 0.20 nmol sphingomyelin (SM) d18:1–18:1 (d$_9$), 10 µL of IS

B (Avanti Polar Lipids Inc.) containing 1.0 nmol monoacylglycerol (MG) 18:1 (d$_7$),

0.10 nmol ceramide (Cer) d18:1 (d$_7$)–15:0, 0.10 nmol hexosylceramide (HexCer) d18:1

(d$_7$)–18:1, 1.1 nmol free fatty acid (FA) 16:0 (^{13}C$_{16}$), and 3.0 nmol cholesterol (d$_7$), and

10 µL of IS C (Merck, Darmstadt, Germany) containing 0.40 nmol acylcoenzyme A

(acyl-CoA) 2:0 (^{13}C^2). The aqueous and organic layers were separated by vortexing and

subsequent centrifugation at 16,000 ×g and 4°C for 5 min.

The aqueous (upper) layer (250 µL) was transferred to a clean tube for analysis

by a liquid chromatography (Nexera X2 UHPLC system, Shimadzu Co., Kyoto, Japan)

with a metal-free peek-coated InertSustain C18 column (2.1 mm i.d. × 150 mm, 3 µm

particle size, GL Sciences Inc., Tokyo, Japan) coupled with a Q Exactive, high-

performance benchtop quadrupole Orbitrap high-resolution tandem mass spectrometer

(Thermo Fisher Scientific) (C18-LC/MS) for acyl-CoAs and acyl-carnitines[84]. After the

aqueous layer extracts were evaporated under vacuum, dried extracts were stored at

−80°C until use for C18-LC/MS analysis. Prior to analysis, the dried aqueous layer was

reconstituted in 50 µL of water.

The organic (lower) layer (140 µL) was transferred to a clean tube for analysis

by supercritical fluid chromatography (SFC, Nexera UC system, Shimadzu Co.) with an

ACQUITY UPC^2 HSS C18 column (3.0 mm i.d. × 100 mm, 1.8 µm particle size, Waters,

Milford, MA, USA) coupled with a triple quadrupole mass spectrometry (TQMS,

LCMS-8060, Shimadzu Co.) (C18-SFC/MS/MS) for FAs, TGs, and CEs[84] and an SFC

(Shimadzu Co.) with a ACQUITY UPC2 Torus diethylamine (DEA) (3.0 mm i.d. × 100

mm, 1.7 µm particle size, Waters) coupled with a TQMS (DEA-SFC/MS/MS) for PCs,

PEs, PSs, PGs, PIs, PAs, LPCs, LPEs, MGs, DGs, SMs, cholesterol, Cers, and

HexCers[84,85]. The liver lipid extract (140 µL) was diluted to a final volume of 200 µL

with a solution of methanol for C18-SFC/MS/MS and DEA-SFC/MS/MS analysis.

Transcriptomic analysis

Total RNAs were extracted from approximately 10 mg of liver using RNeasy Mini Kit

(74106, QIAGEN) and QIAshredder (79656, QIAGEN). The amount and quality of

RNAs were assessed using Nanodrop (Thermo Fisher Scientific) and the 2100

Bioanalyzer (Agilent Technologies). The cDNA libraries were prepared using the

TruSeq Stranded mRNA Kit (Illumina, San Diego, CA, USA). The resulting cDNAs

were subjected to 150 base pair (bp) paired-end sequencing on an Illumina

NovaSeq6000 Platform (Illumina). Sequences were aligned to the reference genome of

mice derived from the Ensembl database[86,87] (GRCm38/mm10, Ensembl release 97)

using the STAR software package (v.2.5.3a)[88]. The RSEM tool[89] (v.1.3.0) was used to

estimate gene expression levels from the aligned sequences. Resultant read counts were

used as inputs for edgeR software[52]. For other analyses, transcripts per kilobase million

(TPM) were used.

Proteomic analysis

Total proteins extracted during sample preparation of metabolomic analysis were

subjected to the following proteomic analysis. For standardization among runs of

measurement, proteins from frozen liver powder of Lys(6)-SILAC mice (Silantes) were

also extracted with methanol/chloroform extraction. Protein concentration of the lysates

was determined using the bicinchoninic acid (BCA) assay (23227, Thermo Fisher

Scientific) and calibrated to 1 mg/mL. Then, 50 µg each of non-labeled and labeled

lysates were mixed for peptide preparation. Cysteine residues were blocked by 2 mM

Tris (2-carboxyethyl) phosphine hydrochloride (TCEP) (Thermo Fisher Scientific) at

37°C for 30 min followed by alkylation with 10 mM 2-iodoacetamide (IAA) at room

temperature for 30 min. The proteins were precipitated with acetone for 3 hours at

−30°C and the resulting pellet was dispersed in 50 mM triethylammonium bicarbonate

by ultrasonic treatment (three times for 30 s with intervals of 30 s) with a Bioruptor

(Diagenode). The protein suspension was subjected to digestion with lysyl

endopeptidase (Wako) for 16 h at 37°C. Resulting peptides were centrifuged at 15,000 ×

g for 15 min at 4°C and subjected to C18-StageTip purification[90]. Peptides were

resolved in 50 µL 3% acetonitrile (can)-0.1% foromic acid prior to MS analysis.

All samples were analyzed with Q Exactive HF-X (Thermo Fisher Scientific)

instrument equipped with an UltiMate 3000 RSLCnano LC System (Thermo Fisher

Scientific) via a nano-electrospray source with a column oven set at 50°C (AMR Inc.).

Peptides were directly injected onto a 75 μm × 25 cm PicoFrit emitter (New Objective,

Woburn, MA, USA) packed in-house with C18 core-shell particles (CAPCELL CORE

MP 2.7 μm, 160 Å material; Osaka Soda Co., Ltd., Osaka, Japan) with a linear gradient

of 7–42% B for 76 min, 42-80% B for 5 min, and 80% B for 9 min at a flow rate of 100

nL/min, where A is 0.1% formic acid and B is 0.1% formic acid and 80% acetonitrile.

MS data acquisition was performed in data-independent acquisition (DIA) mode. For

quantification in the DIA mode, MS1 spectra were collected in the range of m/z 423-865

at 60,000 resolution to set an AGC target of 3×10^6. For MS2 spectra, collections were

conducted in the range of $> m/z$ 200 at 30,000 resolution to set an AGC target of 3×10^6.

The isolation width was configured to 8 Th with stepped normalized collision energies

of 20, and 23%. Through the optimization by Skyline 4.1[91], isolation window patterns in

m/z 428-860 were used as window placements. In the DIA mode for a spectral library,

the pool of all samples was analyzed by using the above measurements and the gas-

phase fractionation method. In the gas-phase fraction method, six MS ranges (m/z 426-

502, 498-574, 570-646, 642-718, 714-790, and 786-862) were used, and each was

measured by overlapping DIA[92,93]. MS1 spectra were collected at 120,000 resolution to

set an AGC target of 3×10^6. MS2 spectra were collected in the range of $> m/z$ 200 at

60,000 resolution to set an AGC target of 3×10^6. The isolation width was set to 2 Th

with stepped normalized collision energies of 20 and 23%. Isolation window patterns in

the ranges of m/z 428-500, 500-572, 572-644, 644-716, 716-788, and 788-860 were

used as window placements optimized by Skyline 4.1.

The spectra library was generated by searching MS data in the library against

the mouse Ensembl release 97 database using Scaffold DIA (Proteome Software Inc.,

Portland OR, USA). The parameters were as follows: experimental data search enzyme,

LysC/P; maximum missed cleavage sites, 1; precursor mass tolerance, 6 ppm; fragment

mass tolerance, 8 ppm; and static modification, cysteine carbamidomethylation. The

threshold for identification of peptides was a peptide false discovery rate (FDR) < 1%.

To identify peptides derived from Lys(6)-SILAC mice, a library of identified peptides

with a mass increase 6.020129 Da on Lys was built and added to the library obtained

above. All raw data was processed with DIA-NN (ver. 1.8) using library search mode

using the following parameters: reannotation, Mus_musculus.GRCm38.97.pep.all.fa;

modification, UniMod188 (6.020129), and peak-translation. Using the output from

DIA-NN software, the SILAC ratios of all assigned fragment ion pairs were calculated,

and the median of the SILAC ratios of each protein was used for protein quantification.

Proteins assigned to a single ensemble gene ID were used in the following analyses. The

SILAC ratios of all the proteins were normalized by the median of the SILAC ratio in

each biological sample.

Phosphoproteomic analysis

Total proteins extracted during sample preparation of metabolomic analysis were

subjected to the following phosphoproteomic analysis. As described above, 2 mg of

protein solution was subjected to reduction and alkylation by TCEP and IAA,

respectively. Then, proteins were precipitated in three volumes of ice-cold acetone and

centrifuged at $16,000 \times g$ for 2 h. The pellet was rinsed with 90% acetone followed by

digestion with 30 µg of trypsin in 600 µl of 100 mM ammonium bicarbonate for 16 h.

Thirty-five µl of 10% trifluoroacetic acid (TFA) and 450 µl of ACN were added to the

protein digests (1 mg/300 µl). The mixture was centrifuged at $16,000 \times g$ and the

supernatant was collected into a new tube. Phosphopeptides were enriched using Fe^{3+}-

IMAC[94]. NTA-agarose resin (1 mL) (Qiagen) was washed twice with 3 ml of distilled

water and 2 ml of 100 mM $FeCl_3$ in 0.1% acetic acid was added. Following a wash with

2 ml of 0.1% TFA, 2% ACN and 0.1% TFA, 60% ACN, the resin was stored in 1 ml of

0.1% TFA, 60% ACN (Fe^{3+}-IMAC agarose resin). C18-StageTip (3M) packed with 75

µl of Fe^{3+}-IMAC agarose resin was equilibrated with 200 µl of 0.1% TFA, 60% ACN,

then the protein digests were loaded onto the IMAC/C18-StageTip and the loaded

system was washed twice with 200 µL of 0.1% TFA, 60% ACN. Following

equilibration with 200 µl of 0.1% TFA, 2% ACN, 200 µl of 1% inorganic phosphate

was added to elute phosphopeptides from the Fe^{3+}-IMAC agarose resin and bind them

to the C18-StageTip. Following a wash with 200 µL of 0.1% TFA, 2% ACN, the

peptides were eluted with 0.1% TFA, 60%ACN from the IMAC/C18-StageTip. The

resulting peptides were dried, reconstituted in 3% ACN with 0.1% formic acid, and

subjected to MS analysis.

All samples were analyzed with Orbitrap Exploris 480 MS (Thermo Fisher

Scientific) instrument equipped with an UltiMate 3000 RSLCnano LC System (Thermo

Fisher Scientific) via a nano-electrospray source with a column oven set at 60°C (AMR

Inc.). Peptides were directly applied onto a 75 μm × 120 mm column (Nikkyo Technos

Co.,Ltd) with a linear gradient of 5–32% B for 70 min and 32-65% B for 10 min at a

speed of 200 nL/min, where A is 0.1% formic acid and B is 0.1% formic acid and 80%

acetonitrile.

MS data were acquired in overlapping data-independent acquisition (DIA)

mode[92,93]. For quantification in the DIA mode, MS1 spectra were collected in the range

of m/z 390-1010 at 30,000 resolution to set an AGC target of 3×10^6. MS2 spectra were

collected in the range of $> m/z$ 200 at 30,000 resolution to set an AGC target of 3×10^6,

and stepped normalized collision energies of 22, 26, and 30%. The isolation width for

MS2 was set to 10 Th and overlapping window patterns in m/z 400-1000 were used

window placements optimized by Skyline v4.1[91]. In the DDA-MS mode for a spectral

library, the pool of all samples was analyzed by using the gas-phase fractionation

method with MS ranges of m/z 395-555, 545-705, 695-1005, and 390-1010. MS1

spectra were collected at 120,000 resolution to set an AGC target of 3×10^6. Within the

MS ranges of m/z 395-555, 545-705, 695-1005, the 20 most intense ions with charge

states of 2^+ to 5^+ more than 5.0×10^3 were fragmented by HCD with normalized

collision energies of 22, 26, and 30%, and MS2 spectra were collected in the range of $>$

m/z 200 at 60,000 resolution to set an AGC target of 5×10^5. Within the MS ranges of

m/z 390-1010, the 40 most intense ions with charge states of 2^+ to 5^+ that exceeded $5.0 \times$

10^3 were fragmented by HCD with normalized collision energies of 22, 26, and 30%,

and MS2 spectra were collected in the range of > *m/z* 200 at 30,000 resolution to set an

AGC target of 2×10^5.

The spectra library was built by searching MS data in the library against the

mouse Ensembl database (GRCm38/mm10, Ensembl release 97) using Proteome

Discoverer v2.3 (Thermo Fisher Scientific). The setting parameters were as follows:

Fragmentation, HCD; Precursor Tolerance, 8 ppm; Fragment Tolerance, 0.02 Da;

Digestion Enzyme, Trypsin; Max Missed Cleavages, 1; Variable Modification, Phospho

[S, T, Y]; Fixed Modification, Carbamidomethylation [C]; Site Probability Threshold,

75; Peptide FDR, < 1%. Unique phosphopeptides were subjected to the following

analyses. To determine phosphorylation at each phosphosite, the intensities of all signals

of quantified phosphopeptides that include the phosphosite of interest were summed.

Glycogen content assay

Glycogen content in liver was measured using Amplex Red Glucose/Glucose Oxidase

Assay Kit glucose assay (Thermo Fisher Scientific) after digestion of glycogen to

glucose[40,95]. Approximately 20 mg of crushed tissue samples was incubated for 1 hour

at 95°C in 1 mL of 30% (w/v) KOH solution. The lysate (50 μL) was collected into

another 1.5 mL tube and neutralized by 15.3 μL of glacial acetic acid. Total protein

concentration of the liver digest was determined using the BCA assay (23227, Thermo

Fisher Scientific) and the concentration of protein was adjusted to 1 μg/μL. Lipids were

removed by the Bligh and Dyer method and glycogen was extracted. Ice-cold methanol

(120 μL), 50 μL of chloroform, 10 μL 1% (w/v) of linear polyacrylamide, and 70 μL of

water were added to the liver digest (50 µL) and mixed. The mixture was incubated for

30 min on ice and centrifuged at $12,000 \times g$ to remove the aqueous layer. The glycogen

was precipitated by adding 200 µL of methanol, followed by centrifugation at $12,000 \times$

g for 30 min at 4°C. The pellet was washed with ice-cold 80% (v/v) methanol and dried

completely. Glycogen pellets were resuspended in 20 µL of amyloglucosidase (0.1

mg/mL; Sigma-Aldrich) in 50 mM sodium acetate buffer and incubated for 2 hours at

55°C. The glycogen content was determined using the Amplex Red Glucose/Glucose

Oxidase Assay Kit glucose assay (Thermo Fisher Scientific) following the

manufacture's protocol.

Identification of starvation-responsive molecules

We analyzed molecules or phosphorylation events that were quantified in more than half

of replicates in both WT and *ob*/*ob* mice at all the time points after food removal (Fig. 2

and Fig. S1). For every molecule, one-way-ANOVA tests were conducted in data for

WT or *ob*/*ob* mice. Molecules with $P < 0.025$ were defined as starvation-responsive

molecules. False discovery rate (FDR) of the responsive molecules was less than 0.1

except for phosphoproteome data (Table 1). FDR was calculated using the Storey's

procedure[96] except for the lipidomic and FFA & acyls layers, including molecules in the

lipid and FFAs & acyls layers. For lipidomic and FFA & acyls analyses, Benjamini-

Hochberg procedure was conducted[97], because the number of measured molecules were

small in these omic layers (16 for lipidome and 41 for FFA & acyls). For transcriptomic

analysis, the likelihood ratio test in the edgeR package (version 3.34.1)[52] of the R

language (version 4.1.1) was used to identify significantly changed molecules during

starvation. Mean time courses for each molecule in WT and *ob*/*ob* mice were log2-

transformed relative to the geometric mean of the values of WT and *ob/ob* mice before

food deprivation (0 h). Using these log2-transformed time courses, the area under the

curve (AUC) was calculated as the area between the time course curves during

starvation and the baseline value at 0 h (before food deprivation). Molecules with

positive AUC and *P* value < 0.025 were defined as increased molecules, and molecules

with negative AUC and *P* value < 0.025 were defined as decreased molecules.

Pathway enrichment analysis

We conducted pathway enrichment analysis for increased or decreased transcripts

(Table 3; Table S3), proteins (Table 4; Table S3), and phosphoproteins (Table S3). From

KEGG database[58–60], we extracted gene sets of pathways in "Metabolism", "Genetic

Information Processing", "Environmental Information Processing", "Cellular

Processes", and "Organismal Systems". Note that "Global/overview pathways" were

removed. The enrichment of increased or decreased molecules in each pathway was

determined using one-tailed Fisher's exact test followed by FDR correction by the

Benjamini-Hochberg procedure[97]. Transcripts, proteins, and phosphoproteins quantified

in more than half of the replicates in WT and *ob/ob* mice were used as background.

Pathways with *Q*-values < 0.1 were identified significantly enriched.

1000

Inference for activities of TFs

We inferred the TF activities using transcriptomic data and ChIP-Atlas[70,71] (Table 5;

Table S4). ChIP-Atlas includes a database curating publicly available ChIP-seq

experiments, that can be used for inference of the activities of TFs[98]. The list for all the

ChIP-seq experiments was downloaded (https://github.com/inutano/chip-

atlas/wiki#downloads_doc) and all the experimental IDs of mm10 were extracted. We

focused on the ChIP-seq experiments conducted using "liver" or "hepatocytes" in mice.

Using this list of experimental IDs of mm10, we scraped "Target Genes" information

that includes genes whose transcription start sites (TSSs) were identified as binding

partners with the bait proteins (TFs) (https://chip-

atlas.dbcls.jp/data/mm10/target/**Protein.Distance**.tsv). The "Distance" was set as 1 kb

from TSS. We set the threshold for statistical significance of values calculated by peak-

caller MACS2 as 50 and identified target genes for each ChIP-seq experiment. To keep

quality of the inference, we removed ChIP-seq experiments that detected less than 50

target genes. Through these processes, we obtained a network between bait proteins

(TFs) for each ChIP-seq experiment and "Target Genes" with TSSs that were bound to

the bait proteins (TFs). Note that this network is a network of ChIP-seq experiments and

transcripts, not TFs and transcripts. In other words, ChIP-seq experiments for the

identical TFs were not combined at this step.

Using this network, we conducted TF enrichment analyses for increased or

decreased transcripts during starvation in WT and *ob/ob* mice. The enrichment of

increased or decreased transcripts in each ChIP-seq experiment was determined using

one-tailed Fisher's exact test followed by FDR correction conducted by the Benjamini-

Hochberg procedure[97]. Transcripts quantified in more than half of the replicates in both

WT and *ob/ob* mice were used as background. ChIP-seq experiments with Q-values less

than 0.1 were identified as significantly enriched ChIP-seq experiments. TFs for which

significant enrichment was observed in at least one ChIP-seq experiment were identified

as candidate TFs that changed their activities during starvation.

We further screened the candidate TFs to define the up-regulated or down-

regulated TFs using transcriptomic, proteomic, and phosphoproteomic data. It should be

noted that TFs that are registered as negative regulators in Transcriptional Repressor

Complex (GO_0017053) were defined as negative regulators, and other TFs were

regarded as activators in this study. If a transcriptional activator was enriched in

increased transcripts, and any of the transcriptomic, proteomic, or phosphoproteomic

data of the TF increased and none of the transcriptomic nor proteomic data decreased,

the TF was identified as an up-regulated TF. If a transcriptional activator was enriched

in decreased transcripts, and any of the transcriptomic, proteomic, or phospho-

proteomic data of the TF decreased and none of the transcriptomic nor proteomic data

increased, the TF was identified as a down-regulated TF. If a transcriptional inhibitor

was enriched in increased transcripts, and any of the transcriptomic, proteomic, or

phosphoproteomic data of the TF decreased and none of the transcriptomic nor

proteomic data increased, the TF was identified as an up-regulated TF. If a

transcriptional inhibitor was enriched in decreased transcripts, and any of the

transcriptomic, proteomic, or phosphoproteomic data of the TF increased and none of

the transcriptomic nor proteomic data decreased, the TF was identified as a down-

regulated TF.

Downstream genes of the inferred TFs were defined based on the gene

regulatory network from the ChIP-Atlas database. For each up-regulated TF, increased

transcripts whose TSS bind to the TF in at least one ChIP-experiment were defined as

downstream transcripts of the TF. For each down-regulated TF, decreased transcripts

whose TSS bind to the TF in at least one ChIP-experiment were defined as downstream

transcripts of the TF.

1053

Inference for activities of protein kinases

1054

We inferred the protein kinase activities using phosphoproteomic data and NetPhorest[99]

(Table 6; Table S4). We constructed a kinase regulatory network using NetPhorest[99], a

protein sequence-based algorithm to infer kinase-substrate relationships. The inputs for

NetPhorest were mouse protein sequences that were associated with ENSMUSP in

FASTA format obtained from Mus_musculus.GRCm38.97.pep.all.fa in Ensembl

database[86,87], and a list for phosphorylated residues measured in our phosphoproteomic

data. The sequences in FASTA format were handled using Seqkit tool[100]. With the

NetPhorest algorithm, we obtained posterior probabilities that a residue was

phosphorylated by protein kinase groups. We selected protein kinase groups with

posterior probability value more than 0.1 for each residue. Protein kinases contained in

protein kinase groups were obtained from NetworKIN[101]. We converted the NetworKIN

kinase names to Ensembl IDs using Ensembl databases, STRING data in

NetWorKIN_release3.0, and manual curation (Table S9).

We obtained a regulatory network among transcripts, proteins, phosphorylation,

and activities of protein kinases and phosphorylation of substrate proteins. Using this

network, we conducted kinase enrichment analyses for increased or decreased

phosphorylation events during starvation in WT and *ob*/*ob* mice. The enrichment of

increased or decreased phosphorylation events in each kinase group was determined

using one-tailed Fisher's exact test followed by FDR correction conducted by the

Benjamini-Hochberg procedure[97]. Phosphorylation events quantified in more than half

of the replicates in WT and *ob*/*ob* mice were used as background. Protein kinase groups

with Q-values less than 0.1 were identified significantly enriched protein kinases.

We screened the candidate protein kinases to define the upregulated or

downregulated protein kinases using transcriptomic, proteomic, and phosphoproteomic

data. If a protein kinase was enriched in increased phosphorylation events, and any of

the transcriptomic, proteomic, or phosphoproteomic data of the protein kinase increased

and none of the transcriptomic nor proteomic data decreased, the protein kinase was

identified as an up-regulated protein kinase. If a protein kinase was enriched in

decreased phosphorylation events, and any of the transcriptomic, proteomic, or

phospho-proteomic data of the protein kinase decreased and none of the transcriptomic,

proteomic, nor phosphoproteomic data increased, the protein kinase was identified as a

downregulated protein kinase.

1087

Inference for activities of signaling pathways

Pathways in KEGG[58–60] database that contain "signaling" in their pathway names were

defined as signaling pathways. We identified up-regulated or down-regulated signaling

pathways that were enriched in increased or decreased transcripts, proteins, or

phosphorylation events at least one omics layer of transcriptomic, proteomic, or

phosphoproteomic data (Table 7; Table S3).

1094

Construction of allosteric regulatory network

Allosteric regulation of metabolic reactions by metabolites was determined using

BRENDA database (https://www.brenda-enzymes.org/download.php)[102]. Activating and

inhibiting compounds in mammals were extracted from brenda_2023_1.txt. Note that

the taxonomy list of mammals was obtained from NCBI

(https://ftp.ncbi.nlm.nih.gov/pub/taxonomy/new_taxdump/)[103]. Compound names in

brenda_2023_1.txt were transformed to KEGG ID via InChI key or using compound

names in KEGG (ftp.kegg.net/kegg/ligand/compound.tar.gz)[58–60] or HMDB

(https://hmdb.ca/downloads, ver. 5)[104]. InChI keys of BRENDA ligands were obtained

from the search results of "Search Ligands" function in the classic view of BRENDA

without any query terms (Table S10). Through these processes, we obtained the table of

the pairs of EC numbers and KEGG compound IDs.

1107

Construction of transport network

We used the Transporter Classification Database (TCDB, https://tcdb.org/)[61] to identify

transporters and substrates. Transport_activity was defined by the substrates of

transports. To define substrates of transports, TC2ChEBI.txt in TCDB was processed as

follows. TC2ChEBI.txt includes ChEBI IDs, that is a hierarchical database and linked to

each other with "is_a" function (for example, anion (CHEBI: 22563) **is a** ion (CHEBI:

24870))[105]. To focus on specific substrates, we selected ChEBI IDs that include less than

50 ChEBI IDs that can be traced by "is_a" function. The selected ChEBI IDs were

transformed to KEGG compound IDs using KEGG

(ftp.kegg.net/kegg/ligand/compound.tar.gz, downloaded on 13rd/12/2022)[58–60] or

HMDB (https://hmdb.ca/downloads, ver. 5)[104], directly or via compound names.

For transporter protein, we selected transporters that are expressed in human or

mice using Uniprot database

(https://ftp.uniprot.org/pub/databases/uniprot/knowledgebase/complete/)[106], using

taxonomy list of mammals in NCBI

(https://ftp.ncbi.nlm.nih.gov/pub/taxonomy/new_taxdump/)[103]. Among them, we further

screened the transporters localizing on cell membrane, by selecting transporters that

include "Apical cell membrane", "Apicolateral cell membrane", "Basal cell membrane",

"Basolateral cell membrane", "Cell membrane", or "Lateral cell membrane" in

SUBCELLULARLOCATION in Uniprot database with at least one evidence in

ECO:0000250, ECO:0000269, ECO:0000303, or ECO:0000305. Uniprot accessions

were transformed to Entrez IDs by Uniprot database. Entrez IDs for human were

translated to Entrez IDs for mice using gene ortholog table in NCBI database

(gene_orthologs, https://ftp.ncbi.nlm.nih.gov/gene/DATA/)[103].

Combining information of substrates and transporter proteins, we obtained a

table of combinations of TCDB IDs, transporter Entrez IDs and KEGG compound IDs.

Note that the transport_activities were defined by the name of substrates of the transport.

1135

Construction of the global starvation-responsive trans-omics network

To construct the global starvation-responsive trans-omics network, nodes were grouped

into layers based on the chemical properties, biological functions, or both (Table 8).

Layers were defined as follows:

• The signaling_activity layer includes signaling pathways of KEGG database[58–60],

that contains "signaling" in their pathway names.

• The TF_activity, TF_mRNA, TF_protein, and TF_PTM layers include activities,

transcripts, proteins, and phosphorylations of all the TFs investigated, those

were included in "liver" or "hepatocytes" data in ChIP-Atlas database[70,71].

• The kinase_activity layer contains all the activities of the protein kinase groups

in the NetPhorest database[99].

• The kinase_mRNA, kinase_protein, and kinase_PTM layers include transcripts,

proteins, and phosphorylations of the protein kinases that belong to the kinase

groups in the NetPhorest database, respectively.

• Metabolic enzymes were defined using KEGG Pathway information

(ftp.kegg.net\kegg\pathway\organisms\mmu.tar.gz and

ftp.kegg.net\kegg\pathway\pathway.list). The enzymes that belong to the

pathways that are classified in "Metabolism" were defined as metabolic enzymes,

except for Slc33a1 (mmu:11416), a transporter.

- The enzyme_mRNA, enzyme_protein, and enzyme_PTM layers include

   transcripts, proteins, and phosphorylations of metabolic enzymes, respectively.

- The metabolite (in the liver), lipid, FFA & acyls, and plasma metabolite layers

   include all the compounds measured in our metabolomic, lipidomic, and the FFA

   & acyls measurements.

- The transport_activity layer includes all the possible transport activities, that is

   defined using substrates of the transports in TCDB.

- The transporter_mRNA, transporter_protein, and transporter_PTM layers

   include all the measured transcripts, proteins, and phosphorylations of

   transporters regulating transport_activities, respectively.

- The metabolic reaction layer includes EC numbers in the KEGG database[58–60]

   (ftp.kegg.net\kegg\genes\organisms\mmu\mmu_link.tar.gz) that metabolic

   enzymes belong to.

   Regulatory connections between nodes were determined using databases. The

following databases or assumptions were used to infer the edges (regulatory

connections):

- KEGG database (ftp.kegg.net\kegg\ligand\enzyme.tar.gz)[58–60] was used to infer

   the substrate and product regulation from metabolites, lipids, and FFA & acyls to

   metabolic reactions.

- The BRENDA[102] database was used to infer the allosteric regulatory

1175   connections from metabolites, lipids, and FFA & acyls to metabolic reactions.

- 1176 Regulation from enzyme_protein and enzyme_PTM to metabolic reactions were
1177   defined using KEGG database[58–60].

- 1178 Regulation of transport_activities was assumed to be mediated by the plasma
1179   metabolites, transporter_proteins, and transporter_PTMs.

- 1180 TCDB was used to determine regulation of metabolites in the liver by
1181   transport_activity[61].

- 1182 To determine regulatory connections between the TF_mRNA, TF_protein,
1183   TF_PTM and TF_activity layers, gene symbols of the bait in ChIP-Atlas
1184   database (TFs) were translated to ENSMUSG IDs using Ensembl database (ver.
1185   97)[86,87].

- 1186 ChIP-Atlas[70,71] was used to determine regulatory connections between the
1187   TF_activity, and enzyme_mRNA or transporter_mRNA layers.

- 1188 Regulation between the enzyme_mRNA and enzyme_protein, and
1189   transporter_mRNA and transporter_protein, were established by aligning Entrez
1190   IDs[103] of the metabolic enzymes and transporters.

- 1191 For protein kinase and phosphorylations, the kinase_mRNA, kinase_protein, and
1192   kinase_PTM layers were assumed to regulate the kinase_activity layer.

- 1193 NetPhorest[99] was used to determine regulation between the kinase_activity layer
1194   and the enzyme_PTM layer.

- 1195 Signaling pathways were assumed to regulate TF_mRNA, TF_protein, TF_PTM,
kinase_mRNA, kinase_protein, kinase_PTM, enzyme_mRNA, enzyme_protein,
enzyme_PTM, transporter_mRNA, transporter_protein, and transporter_PTM.
The regulatory relationships were defined using KEGG database. Note that

signaling pathways and downstream molecules were connected regardless of

increase or decrease of the molecules.

For phosphorylation-related regulation, phosphorylation events curated as

"activity, inhibited", "enzymatic activity, inhibited", "protein degradation", "receptor

desensitization, induced", "receptor inactivation, induced", "receptor internalization,

induced", or "receptor recycling, inhibited" in PhosphositePlus

(https://www.phosphosite.org/staticDownloads)[107] were assumed to negatively regulates

their downstream nodes.

1207

Construction and structural analysis of the starvation-responsive metabolic

network

The starvation-responsive metabolic network is a sub-network of the global starvation-

responsive metabolic network containing metabolic reactions and their direct regulators,

including metabolite, lipid, FFA & acyls, enzyme_protein and enzyme_PTM layers

(Table S6). The network was regarded as directed graph for the calculation of degree

and degree distribution. The degree of each node (k) was counted by the centrality

function in MATLAB, with the 'outdegree' option specified, and degree distribution

($n(k)$) was determined. Scaling parameter was inferred by fitlm function in MATLAB.

Hub nodes were defined as responsive nodes within the top 2% based on highest

degrees.

The network was regarded as an undirected graph for the calculation of the

following structural properties. Because the starvation-responsive metabolic network is

composed of metabolic reactions and their direct regulators, all the edges are directed to

metabolic reactions and no edge starts from metabolic reactions. In this case,for the

calculation of the cluster sizes, only weakly connected components are generated. For

simplicity, we regarded the starvation-responsive metabolic network as an undirected

graph. In addition, the average path length would be 1 if we regard the starvation-

responsive metabolic network as a directed graph, because the regulators of the

metabolic reactions (metabolites, lipids, FFA & acyls, proteins and phosphorylation

events) regulate metabolic reactions with a single edge, and no edge starts from the

metabolic reactions.

The conncomp function in MATLAB was used to determine the maximum

cluster size. Distances between nodes were calculated by the distances function in

MATLAB and the average path length of the graphs were determined. Note that

distances among nodes without any connection were removed for calculation of the

average path length.

To connect hub nodes to metabolic pathways (Fig. S3), metabolic reactions

were classified into metabolic pathways in carbohydrate metabolism, lipid metabolism,

nucleotide metabolism, and amino acid metabolism using KEGG database[58–60]. Then the

number of metabolic reactions in each pathway that was regulated by the hub nodes was

counted. The number of metabolic reactions were log2-transformed to display in the

heatmap. Note that some metabolic reactions overlap among pathways because KEGG

database[58–60] allow overlaps of metabolic reactions among pathways.

1242

Calculation of *t*-half and temporal order analysis

The index of the temporal rate of response (*t*-half) was defined as the time when the

responses to starvation reached the half of the maximum absolute values of the gain of

the responses. We defined *t*-half for the responsive nodes not for metabolic reactions.

Calculated t-half values were analyzed by the following procedures. The

correlation between t-half and log10-transformed degree was determined by Spearman's

correlation coefficient. Comparison between t-half values of WT-specific and *ob/ob*-

specific hub molecules were conducted by Wilcoxon rank sum test. The classification of

starvation-responsive metabolites, FFA & acyls, and lipids into metabolic pathways was

conducted using KEGG database[58–60] for comparison of t-half values among metabolic

pathways. Note that some molecules are included in multiple metabolic pathways.

Comparisons of t-half values among metabolic pathways and omic layers were

conducted using Kruskal-Wallis test followed by Dunn-Sidak test.

To identify the essential molecules for negative correlation between log10-

transformed degree and t-half in the starvation-responsive metabolic network in WT

mice, leave-one-out correlation and cumulative leave-one-out correlation were

calculated for each molecule. The leave-one-out correlation of molecule A was defined

as correlation coefficient between the log10-transformed degree and t-half for all the

responsive molecules in the starvation-responsive metabolic network except for A. The

cumulative leave-one-out correlation of molecule A was defined as correlation

coefficient between the log10-transformed degree and t-half of the responsive molecules

in the starvation-responsive metabolic network with leave-one-out correlations that

were smaller (the contribution to the negative correlation between log10-transformed

degree and t-half is smaller) than A.

1267

Analysis of co-regulation among molecules in the starvation-responsive metabolic

network and construction of correlation metabolic network

Pairwise Pearson's correlation coefficients were calculated among time courses of

significantly changed molecules [metabolites, lipids, FFA & acyls, protein

(enzyme_protein), and phosphorylation events (enzyme_PTM)] in the starvation-

responsive metabolic network in WT and *ob*/*ob* mice.

Based on these correlations, pairs of molecules whose Pearson's correlation

coefficients are more than the threshold were connected by edges, resulting in

correlation networks of the responsive molecules in the starvation-responsive metabolic

network in WT and *ob*/*ob* mice, named as correlation metabolic networks. We tested

several thresholds for the absolute value of the Pearson's correlation coefficients (from

0.3 to 0.9). For each tested threshold, degree of each node was calculated using

centrality function in MATLAB with the 'degree' option specified, and then degree

distribution ($n(k)$) was determined. The scaling parameter was inferred by fitlm function

in MATLAB for each threshold. In addition, the number of the clusters (Fig. S6b), the

relative size of the large cluster (Fig. S6c) and the average path length (Fig. S6d) were

calculated using MATLAB (conncomp function for the number of the clusters and the

relative size of the largest cluster and distances function for the average path length).

1286

Implementation

Statistical test, enrichment analysis, construction of trans-omic network, and network

analysis were conducted using MATLAB 2021a (The MathWorks Inc.). For edgeR, R

language (version 4.1.1) was used. For handling of database, MATLAB, Python 3.8, R

language, and Cygwin (version 3.2.0-1) were used. Transcriptomic analysis obtaining

count and TPM data from sequences and running of NetPhorest software were

conducted on the supercomputer system from the National Institute of Genetics in Japan.

Trans-omic networks were visualized in Graph Modeling Language formats using

Python 2.7 and Visualization and Analysis of Networks containing Experimental Data

(VANTED)[108].

1297

Acknowledgments:

We thank Maki Ohishi and Ayano Ueno (Keio University) for their expertise and

assistance with metabolome analysis using CE-MS. We thank Maiko Goto (Kyushu

University) for technical assistance with lipidome analysis using SFC-MS/MS and LC-

MS. We also thank Kazusa DNA Research Institute for conducting the proteomic and

phosphoproteomic measurements. We also thank our laboratory members for critically

reading this manuscript and technical assistance with the experiments. The

computational analysis of this work was performed in part with support of the

supercomputer system of the National Institute of Genetics (NIG), Research

Organization of Information and Systems (ROIS). The authors thank Nancy R. Gough

(BioSerendipity, LLC) for editorial assistance and critical input.

1309

1310   **Conflict of Interest.** The authors declare no competing interests.

1311

Funding and Assistance.

This study was supported by the Japan Society for the Promotion of Science (JSPS)

KAKENHI grant numbers JP17H06299, JP17H06300, JP18H03979, JP21H04759, and

JP23H04939 to S.K.); CREST, the Japan Science and Technology Agency (JST)

(JPMJCR2123 to S.K., Y. Inaba, and T.S.); The Uehara Memorial Foundation (to S.K.);

K.M. receives funding from a Grant-in-Aid for Early-Career Scientists (JP21K15342).

A.Hatano receives funding from a Grant-in-Aid for Early-Career Scientists

(JP22K15034). T.K. receives funding from a Grant-in-Aid for Early-Career Scientists

(JP21K16349). T.T. receives funding from a Grant-in-Aid for Early-Career Scientists

(JP20K19915). H.O. receives funding from a Grant-in-Aid for Early-Career Scientists

(JP22K17992). S.O. receives funding from a Grant-in-Aid for Early-Career Scientists

(JP21K14467). Y. Inaba also receives AMED-PRIME (JP23gm6910002). This work

was also supported by the Japan Society for the Promotion of Science (JSPS)

KAKENHI grant number JP22H04925 (PAGS); in part by the MEXT Cooperative

Research Project Program, Medical Research Center Initiative for High Depth Omics,

and CURE:JPMXP1323015486 for MIB, Kyushu University; and AMED Grant

Number JP21zf0127001 (T.S.), JST, MEXT KAKENHI Grant Number

JP23H04946(T.S.) and World Premier International Research Center Initiative (WPI),

Human Biology-Microbiome-Quantum Research Center (Bio2Q) (T.S.), MEXT, Japan.;

and JST FOREST Program (Grant Number JPMJFR2052, to A. Hirayama).

1332

Author contributions: K.M., A. Hatano, T.K., H.S, S.O, and S.K. designed the project.

K.M., A. Hatano, and M.M. conducted animal experiments and sample preparation for

omic measurements. K.M. conducted single-omic analysis, trans-omic analysis, and

network structural and temporal analyses. T.S. and A. Hirayama performed

metabolomic analysis using CE-MS and IC-QEMS. Y. Izumi., M.T., and T.B. performed

lipidomic analysis using LC-MS and SFC-MS/MS. Y.S. performed transcriptomic

analysis using RNA-seq. K.M., T.T., R.E., and H.O. analyzed the RNA-seq data. K.M.,

T.K., H.S., D.L., and A.T. conducted database for trans-omic analysis. The writing

group consisted of K.M., Y. Inaba, H.I., and S.K. The study was conceived and

supervised by K.M. and S.K.

1343

Data and code availability. Sequencing data have been deposited in the DNA Data

Bank of Japan Sequence Read Archive (DRA) (www.ddbj.nig.ac.jp/) with accession no.

DRR571760-DRR571839. Proteome and phosphoproteome data were deposited in

jPOSTrepo[109] (https://repository.jpostdb.org/) with accession no, JPST003082

(proteome) and JPST003130 (phosphoproteome). All other data are present in the paper

or the Supplementary Materials. The code used for the analysis in this paper is available

upon request.

1351

 SUPPLEMENTARY FIGURE TITLES AND LEGENDS

Fig. S1: Time courses of responsive molecules during starvation.

The x-axis represents the starvation time (hours) and the y-axis represents the

concentration of the indicated molecule: ng/µg-protein for glycogen, nmol/mg-protein

for metabolite (F6P, alanine, 3-hydroxybutyrate, ATP, AMP); pmol/mg-protein for lipids

(PS, PE, PI) and FFA & acyls (Acyl-CoA 16:0); arbitrary units for proteins (Cpt1a, Pck1,

G6pc) and phosphorylation (S775 of Pfkl). TPM for transcripts (*Cpt1a*, *Pck1*, *G6pc*).

Blue lines are the responses to starvation in WT mice, and red lines are those in *ob/ob*

mice. Solid lines represent that the responses were significant; dotted lines represent no

significance. Data are shown as the mean and SEM. Dots represent the data from

individual mice.

1363

Fig. S2: Regulatory relationships in the starvation-responsive trans-omic network

We assumed that the starvation-responsive molecules cooperatively regulate metabolic

reactions (see also Fig. 1d).

(a) Metabolic network. The enzyme_protein (protein abundance of metabolic enzymes)

and enzyme_PTM (phosphorylation events on metabolic enzymes) regulate metabolic

reactions (identified with KEGG). Metabolites, lipids, and FFA & acyls regulate

metabolic reactions as substrates, products (identified with KEGG), and allosteric

regulators (identified by BRENDA). (b) Transport network identified with TCDB. The

transporter_protein (protein abundance of transporters), transporter_PTM

(phosphorylation events on transporters), and plasma metabolites regulate

transport_activity (activities of transport). Transport_activity regulates metabolites in

the liver, thereby regulating metabolic reactions in the liver. (c) Gene regulatory

network identified with ChIP-Atlas and Ensembl. The TF_mRNA (mRNA abundance of

TFs), TF_protein (protein abundance of TFs), and TF_PTM (phosphorylation events on

TFs) regulate TF_activity (activities of TFs). TF_activity regulates enzyme_mRNA

(mRNA abundance of metabolic enzymes) and transporter_mRNA (mRNA abundance

of transporters). Enzyme_mRNA and transporter_mRNA regulate enzyme_protein and

transporter_protein, respectively, and were aligned with Ensembl database. (d)

Phosphorylation network identified by NetPhorest. The kinase_mRNA (mRNA

abundance of kinases), kinase_protein (protein abundance of kinase), and kinase_PTM

(phosphorylation events on kinases) regulate kinase_activity (activities of kinases). The

kinase_activity regulates enzyme_PTM and transporter_PTM.

1386

Fig. S3: Metabolic pathways regulated by the hub nodes

Heatmaps representing the log2-transformed number of regulated metabolic reactions in

each metabolic pathway (x-axis) regulated by the hub nodes (y-axis) in the starvation-

responsive metabolic network in WT mice (a) and *ob*/*ob* mice (b). Colors of the name of

the hub nodes represent the responsiveness: green indicates common responses both in

WT and *ob*/*ob* mice; blue, specific to WT mice; red, specific to *ob*/*ob* mice; magenta,

opposite responses between WT and *ob*/*ob* mice. KEGG metabolic pathways (x-axis) in

carbohydrate metabolism (olive green), lipid metabolism (orange), nucleotide

metabolism (brown), and amino acid metabolism (light blue) are displayed.

1396

Fig. S4: Maintenance of the number of nodes with relatively large degrees in *ob*/*ob*

mice

The number of nodes (a-e) and the percentage of nodes to the number of responsive

nodes in each genotype (f-j), whose log10-transformed degree are as indicated. The

number of nodes whose log10-transformed degree is between 0.5 to 1 (b) and 1 to 1.5

(c) are comparable between WT and *ob/ob* mice, and the percentage of nodes to the

number of the responsive nodes were larger in *ob/ob* mice for nodes whose log10-

transformed degree is between 0.5 to 1 (g) and 1 to 1.5 (h). (k)Percentages of nodes

whose log10-transformed degree is more than 1.

1406

Fig. S5: Identification of molecules essential for the negative correlation between

degree and *t*-half in WT mice

(a) Schema for leave-one-out correlations. Log10-transformed degree negatively

correlated with *t*-half in WT mice (center, "Rho of full"). The leave-one-out correlation

of molecule A (left, "Rho without A") was defined as correlation coefficient between

log10-transformed degree and *t*-half that was calculated for all the molecules except for

A. The larger leave-one-out correlation (left, "Rho without A") represents the larger

contribution to the negative correlation between log10-transformed degree and *t*-half.

By contrast, the smaller leave-one-out correlation (right, "Rho without B") represents

the small contribution to the negative correlation between log10-transformed degree and

t-half. (b) The cumulative leave-one-out correlation of molecule A (right) was defined

by the correlation coefficient between log10-transformed degree and *t*-half that was

calculated for the molecules with leave-one-out correlation that were smaller than A

(indicated in light blue). (c) Leave-one-out correlations (gray bars), cumulative leave-

one-out correlations (black dots), and *P*-values of the cumulative leave-one-out

correlations (black solid line) of the indicated molecules. Here, we displayed molecules

with *P*-values of the cumulative leave-one-out correlations that were less than 0.05 and

with leave-one-out correlations that were less than other molecules.

1425

Fig. S6: The correlation metabolic network during starvation

To construct correlation metabolic networks during starvation, we tested several

thresholds for absolute values of correlation coefficients (|Rho|) ranging from 0.3 to 0.9

and visualized the degree distributions by the scatter plots (a). The x-axis represents

$\log_{10}(k)$, the log10-transformed degree; the y-axis represents $\log_{10} n(k)$, the log10-

transformed number of nodes with degree of k. Dots denote the degree distribution of

the indicated networks and lines show the linear-regression of the degree distribution.

The scaling parameters (γ) and the 95% credible intervals are shown. Data for WT (top

panels) and *ob*/*ob* mice (bottom panels) are shown. The number of clusters (b), relative

size of the largest cluster (c) and the average path length (d) for each threshold are also

shown. The lower threshold than 0.8 for the absolute value of correlation coefficients

resulted in formation of a single cluster (b). The higher threshold (0.8 or 0.9) for the

absolute value of correlation coefficients resulted in more scale-free networks. To

identify the qualitative differences between WT and *ob*/*ob* mice clearly, we used 0.9 as a

threshold for the absolute value of correlation coefficients in Fig. 7c and d. Using 0.9 as

a threshold for the absolute value of correlation coefficients, the relative size of the

largest cluster was smaller (c) and the average path length was larger in *ob*/*ob* mice than

in WT mice (d).

1444

TITLES AND LEGENDS FOR SUPPLEMENTARY TABLES

Table S1_analyzed_data.xlsx

This file includes all the analyzed data, molecules [metabolites, lipids, FFA & acyls,

proteins (enzyme_protein), and phosphorylation events (enzyme_PTM)] that were

quantified in more than half of replicates in both WT and *ob/ob* mice at all the time

points after food removal. Raw transcriptome data, raw proteome data, and raw

phosphoproteome data are uploaded on public repositories (see data and code

availability section). All measured metabolome, lipidome, and FFA & acyls data are

included in Table S8. The columns represent biological samples (S25-S104), whose

biological conditions are summarized in 'Sample_information' tab. The rows represent

molecules or phosphorylation events. The units were nmol/mg-protein for metabolites

except for glycogen, ng/μg-protein for glycogen, μM for plasma metabolites, pmol/mg-

protein for lipids and FFA & acyls, TPM for transcripts, and arbitrary units for proteins

and phosphorylation events.

1459

TableS2_mean_and_properties.xlsx

This file includes all the mean values of each time point in each genotype, properties of

the time courses (AUC and *t*-half), and results of statistical tests. The columns represent

timepoints and genotypes for mean values and name of properties of the time courses

(AUC and *t*-half) and results of statistical test (*P*-values, *Q*-values, and changes). The

rows represent molecules. The units for mean values were nmol/mg-protein for

metabolites except for glycogen, ng/μg-protein for glycogen, μM for plasma metabolites,

pmol/mg-protein for lipids and FFA & acyls, TPM for transcripts, and arbitrary units for

proteins and phosphorylation events. Abbreviations for WT_change and *ob/ob*-change

columns were as follows: Up, Down, and NS represent increase, decrease and no

significant changes.

1471

Table S3_KEGGPathwayEnrichment.xlsx

This file includes all the results of Fisher's exact test in KEGG Pathway Enrichment

Analyses for transcriptome, proteome, and phosphoproteome data. The Entry, Name,

Class, and Subclass are displayed for the information of each pathway. P-values, Q-

values, and odds ratio for Fisher's exact test, and significantly changed downstream

molecules are listed.

1478

Table_S4_TF_kinase_enrichment.xlsx

This file includes all the results of TF/kinase enrichment analysis for increased or

decreased transcripts/phosphorylation events in WT and *ob/ob* mice, respectively. For

TF, Fisher's exact tests were conducted for each ChIP-seq experiment. The table

contains experimental IDs, genome, type of the experiment, name of TFs, organ, and

cell for the information of ChIP-experiments. Also, P-values, Q-values, and odds ratio

for Fisher's exact test, and significantly changed downstream transcripts are displayed.

For protein kinases, names of kinase groups, ENSMUSG IDs of kinases, and names of

kinases were displayed. Also, P-values, Q-values, and odds ratio for Fisher's exact test,

and significantly changed targeted phosphorylated proteins (by ENSMUSG) are listed.

1489

Table_S5_the_global_starvation_responsive_transomic_NW.xlsx

This file includes all the information of nodes and edges of the global starvation-

responsive trans-omic network. The node tab includes information of node; IDs, names,

omic layers, changes in WT and *ob*/*ob* mice, and comparisons between them (common

responses both in WT and *ob*/*ob* mice; specific to WT mice; specific to *ob*/*ob* mice;

opposite responses between the two genotypes). The edge tab includes the layers and

IDs of the source and target nodes, the type of regulation (Act: activation or Inh:

inhibition), and the presence or absence of the regulation in WT and *ob*/*ob* mice (TRUE

or FALSE).

1499

Table_S6_stv_resp_metab_NW.xlsx

This file includes all the information of nodes and edges of the starvation-responsive

metabolic network. The node tab includes information of node: names, IDs, omic layers,

significance of responses to starvation in WT and *ob*/*ob* mice (TRUE or FALSE), AUC

of the time courses in WT and *ob*/*ob* mice, *t*-half in WT and *ob*/*ob* mice, and degrees

and cluster IDs of the background network (network of the database) and starvation-

responsive metabolic networks in WT and *ob*/*ob* mice. The edge tab includes the layers

and IDs of the source and target nodes, the type of regulation (Act: activation or Inh:

inhibition), and the presence or absence of the regulation in WT and *ob*/*ob* mice (TRUE

or FALSE).

1510

Table_S7_cor_metab_NW.xlsx

This file includes all the pairwise correlation coefficients and *P*-values between

molecules in the starvation-responsive metabolic networks. Each tab represents the type

of values (correlation coefficients or *P*-values) in WT or *ob*/*ob* mice. Both the columns

and rows represent responsive molecules in the starvation-responsive metabolic network

in WT and *ob*/*ob* mice. The pairwise correlation coefficients and *P*-values are displayed

as a matrix in each tab.

1518

Table S8_all_metabolome_lipidome_FFA_Acyls.xlsx

This file includes all data of metabolome, lipidome (including FFA), acyl-carnitines,

and acyl-CoAs. The columns represent biological samples (S25-S104), whose

biological conditions are summarized in 'Sample_information' tab. The rows represent

molecules. The units were nmol/mg-protein for metabolites, pmol/mg-protein for lipids

including FFA, acyl-carnitine, and acyl-CoA. Most metabolites were measured by CE-

MS, while F1,6P and F2,6P were separately measured by IC-QEMS. F1,6P was

quantified both in CE-MS and IC-QEMS. We used values of IC-QEMS in this study.

1527

1528   **Table S9_table_for_kinase_name_groups_ENSMUSGs**

1529   Kinase names, kinase groups and ENSMUSGs of protein kinases were displayed in this

1530   table.

1531

1532   **Table S10_Ligands_and_InChI_Key_in_BRENDA**

1533   InChI keys of BRENDA ligands obtained from the search results of "Search Ligands"

1534   function in the classic view of BRENDA without any query terms.

1535

1.   Cahill, G. F., Jr. Fuel metabolism in starvation. *Annu. Rev. Nutr.* **26**, 1–22 (2006).

2.   Goldstein, I. & Hager, G. L. Transcriptional and Chromatin Regulation during

Fasting - The Genomic Era. *Trends Endocrinol. Metab.* **26**, 699–710 (2015).

3.   Hofer, S. J., Carmona-Gutierrez, D., Mueller, M. I. & Madeo, F. The ups and downs

of caloric restriction and fasting: from molecular effects to clinical application.

EMBO Mol. Med. **14**, e14418 (2022).

4.   Wang, T., Hung, C. C. Y. & Randall, D. J. The comparative physiology of food

deprivation: from feast to famine. *Annu. Rev. Physiol.* **68**, 223–251 (2006).

5.   Geisler, C. E., Hepler, C., Higgins, M. R. & Renquist, B. J. Hepatic adaptations to

maintain metabolic homeostasis in response to fasting and refeeding in mice. *Nutr.*

Metab.   **13**, 62 (2016).

6.   Anton, S. D. *et al.* Flipping the Metabolic Switch: Understanding and Applying the

Health Benefits of Fasting. *Obesity*   **26**, 254–268 (2018).

7.   Ghazalpour, A. *et al.* Genetic regulation of mouse liver metabolite levels. *Mol. Syst.*

Biol. **10**, 730 (2014).

8.   Owen, O. E., Felig, P., Morgan, A. P., Wahren, J. & Cahill, G. F., Jr. Liver and

kidney metabolism during prolonged starvation. *J. Clin. Invest.* **48**, 574–583 (1969).

9. Pietrocola, F. *et al.* Metabolic effects of fasting on human and mouse blood in vivo.

Autophagy **13**, 567–578 (2017).

10. Defour, M., Hooiveld, G. J. E. J., van Weeghel, M. & Kersten, S. Probing metabolic

memory in the hepatic response to fasting. *Physiol. Genomics* **52**, 602–617 (2020).

11. Thøgersen, R. *et al.* Hepatic PGC-1α is not essential for fasting-induced

cytochrome p450 regulation in mouse liver. *Biochem. Pharmacol.* **172**, 113736

(2020).

12. Rennert, C. *et al.* The Diurnal Timing of Starvation Differently Impacts Murine

Hepatic Gene Expression and Lipid Metabolism - A Systems Biology Analysis

Using Self-Organizing Maps. *Front. Physiol.* **9**, 1180 (2018).

13. Régnier, M. *et al.* Insights into the role of hepatocyte PPARα activity in response to

fasting. *Mol. Cell. Endocrinol.* **471**, 75–88 (2018).

14. Schupp, M. *et al.* Metabolite and transcriptome analysis during fasting suggest a

role for the p53-Ddit4 axis in major metabolic tissues. *BMC Genomics* **14**, 758

(2013).

15. Lenaerts, K. *et al.* Starvation induces phase-specific changes in the proteome of

mouse small intestine. *J. Proteome Res.* **5**, 2113–2122 (2006).

16. Steinhauser, M. L. *et al.* The circulating metabolome of human starvation. *JCI*

Insight **3**, (2018).

17. Rubio-Aliaga, I. *et al.* Metabolomics of prolonged fasting in humans reveals new

catabolic markers. *Metabolomics* **7**, 375–387 (2011).

18. Pietzner, M. *et al.* Systemic proteome adaptions to 7-day complete caloric

restriction in humans. *Nature Metabolism* 1–14 (2024).

19. Wasselin, T. *et al.* Exacerbated oxidative stress in the fasting liver according to fuel

partitioning. *Proteomics* **14**, 1905–1921 (2014).

20. Brauer, M. J. *et al.* Conservation of the metabolomic response to starvation across

two divergent microbes. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 19302–19307 (2006).

21. Bergdahl, B., Heer, D., Sauer, U., Hahn-Hägerdal, B. & van Niel, E. W. Dynamic

metabolomics differentiates between carbon and energy starvation in recombinant

Saccharomyces cerevisiae fermenting xylose. *Biotechnol. Biofuels* **5**, 34 (2012).

22. Link, H., Fuhrer, T., Gerosa, L., Zamboni, N. & Sauer, U. Real-time metabolome

profiling of the metabolic switch between starvation and growth. *Nat. Methods* **12**,

1091–1097 (2015).

23. Peng, D. *et al.* Atg9-centered multi-omics integration reveals new autophagy

regulators in Saccharomyces cerevisiae. *Autophagy* **17**, 4453–4476 (2021).

24. Harvald, E. B. *et al.* Multi-omics Analyses of Starvation Responses Reveal a

Central Role for Lipoprotein Metabolism in Acute Starvation Survival in C. elegans.

Cell Syst **5**, 38-52.e4 (2017).

25. Goodman, M. N. *et al.* Starvation in the rat. II. Effect of age and obesity on protein

sparing and fuel metabolism. *Am. J. Physiol.* **239**, E277–E286 (1980).

26. Soltis, A. R. *et al.* Hepatic Dysfunction Caused by Consumption of a High-Fat Diet.

Cell Rep. **21**, 3317–3328 (2017).

27. Stöckli, J. *et al.* Metabolomic analysis of insulin resistance across different mouse

strains and diets. *J. Biol. Chem.* **292**, 19135–19145 (2017).

28. Nelson, M. E. *et al.* Systems-level analysis of insulin action in mouse strains

provides insight into tissue- and pathway-specific interactions that drive insulin

resistance. *Cell Metab.* **0**, (2022).

29. Fazakerley, D. J. *et al.* Phosphoproteomics reveals rewiring of the insulin signaling

network and multi-nodal defects in insulin resistance. *Nat. Commun.* **14**, 1–20

(2023).

30. Stern, J. H. *et al.* Obesity dysregulates fasting-induced changes in glucagon

secretion. *J. Endocrinol.* **243**, 149–160 (2019).

31. Pak, H. H. *et al.* Fasting drives the metabolic, molecular and geroprotective effects

of a calorie-restricted diet in mice. *Nat Metab* **3**, 1327–1341 (2021).

32. Parkhitko, A. A., Filine, E., Mohr, S. E., Moskalev, A. & Perrimon, N. Targeting

metabolic pathways for extension of lifespan and healthspan across multiple species.

Ageing Res. Rev. **64**, 101188 (2020).

33. Mihaylova, M. M. *et al.* When a calorie is not just a calorie: Diet quality and timing

as mediators of metabolism and healthy aging. *Cell Metab.* (2023)

doi:10.1016/j.cmet.2023.06.008.

34. Magkos, F. Is calorie restriction beneficial for normal-weight individuals? A

narrative review of the effects of weight loss in the presence and absence of obesity.

Nutr. Rev. **80**, 1811–1825 (2022).

35. Barabási, A.-L. & Oltvai, Z. N. Network biology: understanding the cell's

functional organization. *Nat. Rev. Genet.* **5**, 101–113 (2004).

36. Halu, A., Mondragón, R. J., Panzarasa, P. & Bianconi, G. Multiplex PageRank.

PLoS One **8**, e78293 (2013).

37. Chella Krishnan, K. *et al.* Integration of Multi-omics Data from Mouse Diversity

Panel Highlights Mitochondrial Dysfunction in Non-alcoholic Fatty Liver Disease.

Cell Syst **6**, 103-115.e7 (2018).

38. Halu, A., De Domenico, M., Arenas, A. & Sharma, A. The multiplex network of

human diseases. *NPJ Syst Biol Appl* **5**, 15 (2019).

39. Yugi, K. *et al.* Reconstruction of insulin signal flow from phosphoproteome and

metabolome data. *Cell Rep.* **8**, 1171–1183 (2014).

40. Kokaji, T. *et al.* Transomics analysis reveals allosteric and gene regulation axes for

altered hepatic glucose-responsive metabolism in obesity. *Sci. Signal.* **13**, (2020).

41. Kokaji, T. *et al.* In vivo transomic analyses of glucose-responsive metabolism in

skeletal muscle reveal core differences between the healthy and obese states. *Sci.*

Rep. **12**, 13719 (2022).

42. Sugimoto, H. *et al.* iTraNet: A Web-Based Platform for integrated Trans-Omics

Network Visualization and Analysis. *bioRxiv* 2023.11.30.569499 (2023)

doi:10.1101/2023.11.30.569499.

43. Ravasz, E., Somera, A. L., Mongru, D. A., Oltvai, Z. N. & Barabási, A. L.

Hierarchical organization of modularity in metabolic networks. *Science* **297**, 1551–

1555 (2002).

44. Asgari, Y., Salehzadeh-Yazdi, A., Schreiber, F. & Masoudi-Nejad, A. Controllability

in cancer metabolic networks according to drug targets as driver nodes. *PLoS One* **8**,

e79397 (2013).

45. Schwartz, J.-M., Otokuni, H., Akutsu, T. & Nacher, J. C. Probabilistic

controllability approach to metabolic fluxes in normal and cancer tissues. *Nat.*

Commun. **10**, 2725 (2019).

46. Barabasi, A. L. & Albert, R. Emergence of scaling in random networks. *Science* **286**,

509–512 (1999).

47. Lindström, P. The physiology of obese-hyperglycemic mice [ob/ob mice].

ScientificWorldJournal **7**, 666–685 (2007).

48. Egami, R. *et al.* Trans-omic analysis reveals obesity-associated dysregulation of

inter-organ metabolic cycles between the liver and skeletal muscle. *iScience* **24**,

102217 (2021).

49. Bai, Y. *et al.* Trans-omic analysis reveals opposite metabolic dysregulation between

feeding and fasting in liver associated with obesity. *iScience* **27**, 109121 (2024).

50. Fendt, S.-M. *et al.* Tradeoff between enzyme and metabolite efficiency maintains

metabolic homeostasis upon perturbations in enzyme capacity. *Mol. Syst. Biol.* **6**,

356 (2010).

51. Lorendeau, D., Christen, S., Rinaldi, G. & Fendt, S.-M. Metabolic control of

signalling pathways and metabolic auto-regulation. *Biol. Cell* **107**, 251–272 (2015).

52. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package

for differential expression analysis of digital gene expression data. *Bioinformatics*

26, 139–140 (2010).

53. McCarthy, D. J., Chen, Y. & Smyth, G. K. Differential expression analysis of

multifactor RNA-Seq experiments with respect to biological variation. *Nucleic*

Acids Res. **40**, 4288–4297 (2012).

54. Chen, Y., Lun, A. T. L. & Smyth, G. K. From reads to genes to pathways:

differential expression analysis of RNA-Seq experiments using Rsubread and the

edgeR quasi-likelihood pipeline. *F1000Res.* **5**, 1438 (2016).

55. King, G. L. & Das-Evcimen, N. Role of protein kinase C in diabetic complications.

Expert Rev. Endocrinol. Metab. **5**, 77–88 (2010).

56. Newton, A. C. Protein kinase C: perfectly balanced. *Crit. Rev. Biochem. Mol. Biol.*

53, 208–230 (2018).

57. Bezy, O. *et al.* PKCδ regulates hepatic insulin sensitivity and hepatosteatosis in

mice and humans. *J. Clin. Invest.* **121**, 2504–2517 (2011).

58. Kanehisa, M., Goto, S., Sato, Y., Furumichi, M. & Tanabe, M. KEGG for

integration and interpretation of large-scale molecular data sets. *Nucleic Acids Res.*

40, D109-14 (2012).

59. Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y. & Morishima, K. KEGG: new

perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res.* **45**,

D353–D361 (2017).

60. Kanehisa, M., Furumichi, M., Sato, Y., Kawashima, M. & Ishiguro-Watanabe, M.

KEGG for taxonomy-based analysis of pathways and genomes. *Nucleic Acids Res.*

51, D587–D592 (2023).

61. Saier, M. H. *et al.* The Transporter Classification Database (TCDB): 2021 update.

Nucleic Acids Res. **49**, D461–D467 (2021).

62. Yang, L., Li, P., Fu, S., Calay, E. S. & Hotamisligil, G. S. Defective hepatic

autophagy in obesity promotes ER stress and causes insulin resistance. *Cell Metab.*

11, 467–478 (2010).

63. Fu, S. *et al.* Polysome profiling in liver identifies dynamic regulation of

endoplasmic reticulum translatome by obesity and fasting. *PLoS Genet.* **8**,

e1002902 (2012).

64. Atiia, A. A., Hopper, C., Inoue, K., Vidal, S. & Waldispühl, J. Computational

intractability law molds the topology of biological networks. *Applied Network*

Science **5**, 1–22 (2020).

65. Liu, Y.-Y., Slotine, J.-J. & Barabási, A.-L. Controllability of complex networks.

Nature **473**, 167–173 (2011).

66. Broido, A. D. & Clauset, A. Scale-free networks are rare. *Nat. Commun.* **10**, 1017

(2019).

67. Albert, R., Jeong, H. & Barabasi, A. L. Error and attack tolerance of complex

networks. *Nature* **406**, 378–382 (2000).

68. Saez-Rodriguez, J. *et al.* Comparing signaling networks between normal and

transformed hepatocytes using discrete logical models. *Cancer Res.* **71**, 5400–5411

(2011).

69. Haeusler, R. A. *et al.* Integrated control of hepatic lipogenesis versus glucose

production requires FoxO transcription factors. *Nat. Commun.* **5**, 5190 (2014).

70. Oki, S. *et al.* ChIP-Atlas: a data-mining suite powered by full integration of public

ChIP-seq data. *EMBO Rep.* **19**, (2018).

71. Zou, Z., Ohta, T., Miura, F. & Oki, S. ChIP-Atlas 2021 update: a data-mining suite

for exploring epigenomic landscapes by fully integrating ChIP-seq, ATAC-seq and

Bisulfite-seq data. *Nucleic Acids Res.* **50**, W175–W182 (2022).

72. Gowans, G. J. & Hardie, D. G. AMPK: a cellular energy sensor primarily regulated

by AMP. *Biochem. Soc. Trans.* **42**, 71–75 (2014).

73. Liu, G. Y. & Sabatini, D. M. mTOR at the nexus of nutrition, growth, ageing and

disease. *Nat. Rev. Mol. Cell Biol.* **21**, 183–203 (2020).

74. Needham, E. J., Parker, B. L., Burykin, T., James, D. E. & Humphrey, S. J.

Illuminating the dark phosphoproteome. *Sci. Signal.* **12**, (2019).

75. Müller-Dott, S. *et al.* Expanding the coverage of regulons from high-confidence

prior knowledge for accurate estimation of transcription factor activities. *Nucleic*

Acids Res. **51**, 10934–10949 (2023).

76. Johnson, J. L. *et al.* An atlas of substrate specificities for the human

serine/threonine kinome. *Nature* (2023) doi:10.1038/s41586-022-05575-3.

77. Türei, D., Korcsmáros, T. & Saez-Rodriguez, J. OmniPath: guidelines and gateway

for literature-curated signaling pathway resources. *Nat. Methods* **13**, 966–967

(2016).

78. Davis, R. C. *et al.* Systems genetics of susceptibility to obesity-induced diabetes in

mice. *Physiol. Genomics* **44**, 1–13 (2012).

79. Soga, T. & Heiger, D. N. Amino acid analysis by capillary electrophoresis

electrospray ionization mass spectrometry. *Anal. Chem.* **72**, 1236–1241 (2000).

80. Soga, T. *et al.* Differential metabolomics reveals ophthalmic acid as an oxidative

stress biomarker indicating hepatic glutathione consumption. *J. Biol. Chem.* **281**,

16768–16776 (2006).

81. Soga, T. *et al.* Metabolomic profiling of anionic metabolites by capillary

electrophoresis mass spectrometry. *Anal. Chem.* **81**, 6165–6174 (2009).

82. Hirayama, A. *et al.* The use of a double coaxial electrospray ionization sprayer

improves the peak resolutions of anionic metabolites in capillary ion

chromatography-mass spectrometry. *J. Chromatogr. A* **1619**, 460914 (2020).

83. Bligh, E. G. & Dyer, W. J. A rapid method of total lipid extraction and purification.

Can. J. Biochem. Physiol. **37**, 911–917 (1959).

84. Fushimi, T. *et al.* Dynamic Metabolome Analysis Reveals the Metabolic Fate of

Medium-Chain Fatty Acids in AML12 Cells. *J. Agric. Food Chem.* **68**, 11997–

12010 (2020).

85. Takeda, H. *et al.* Widely-targeted quantitative lipidomics method by supercritical

fluid chromatography triple quadrupole mass spectrometry. *J. Lipid Res.* **59**, 1283–

1293 (2018).

86. Flicek, P. *et al.* Ensembl 2014. *Nucleic Acids Res.* **42**, D749-55 (2014).

87. Cunningham, F. *et al.* Ensembl 2015. *Nucleic Acids Res.* **43**, D662-9 (2015).

88. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–

21 (2013).

89. Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq

data with or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

90. Rappsilber, J., Mann, M. & Ishihama, Y. Protocol for micro-purification,

enrichment, pre-fractionation and storage of peptides for proteomics using

StageTips. *Nat. Protoc.* **2**, 1896–1906 (2007).

91. MacLean, B. *et al.* Skyline: an open source document editor for creating and

analyzing targeted proteomics experiments. *Bioinformatics* **26**, 966–968 (2010).

92. Kawashima, Y. *et al.* Optimization of Data-Independent Acquisition Mass

Spectrometry for Deep and Highly Sensitive Proteomic Analysis. *Int. J. Mol. Sci.*

20, (2019).

93. Amodei, D. *et al.* Improving Precursor Selectivity in Data-Independent Acquisition

Using Overlapping Windows. *J. Am. Soc. Mass Spectrom.* **30**, 669–684 (2019).

94. Hatano, A. & Matsumoto, M. Sample preparation for phosphopeptides enrichment

using Fe^{3+}-IMAC. *Journal of Proteome Data and Methods* **2**, 3 (2020).

95. Noguchi, R. *et al.* The selective control of glycolysis, gluconeogenesis and

glycogenesis by temporal insulin patterns. *Mol. Syst. Biol.* **9**, 664 (2013).

96. Storey, J. D. A Direct Approach to False Discovery Rates. *J. R. Stat. Soc. Series B*

Stat. Methodol. **64**, 479–498 (2002).

97. Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: A practical and

powerful approach to multiple testing. *J. R. Stat. Soc.* **57**, 289–300 (1995).

98. Garcia-Alonso, L., Holland, C. H., Ibrahim, M. M., Turei, D. & Saez-Rodriguez, J.

Benchmark and integration of resources for the estimation of human transcription

factor activities. *Genome Res.* **29**, 1363–1375 (2019).

99. Miller, M. L. *et al.* Linear motif atlas for phosphorylation-dependent signaling. *Sci.*

Signal. **1**, ra2 (2008).

100.   Shen, W., Le, S., Li, Y. & Hu, F. SeqKit: A Cross-Platform and Ultrafast

Toolkit for FASTA/Q File Manipulation. *PLoS One* **11**, e0163962 (2016).

101.   Linding, R. *et al.* NetworKIN: a resource for exploring cellular

phosphorylation networks. *Nucleic Acids Res.* **36**, D695-9 (2008).

102.   Chang, A. *et al.* BRENDA, the ELIXIR core data resource in 2021: new

developments and updates. *Nucleic Acids Res.* **49**, D498–D508 (2021).

103.   Sayers, E. W. *et al.* Database resources of the national center for biotechnology

information. *Nucleic Acids Res.* **50**, D20–D26 (2022).

104.   Wishart, D. S. *et al.* HMDB 5.0: the Human Metabolome Database for 2022.

Nucleic Acids Res. **50**, D622–D631 (2022).

105.   Hastings, J. *et al.* ChEBI in 2016: Improved services and an expanding

collection of metabolites. *Nucleic Acids Res.* **44**, D1214-9 (2016).

106.   UniProt Consortium. UniProt: the Universal Protein Knowledgebase in 2023.

Nucleic Acids Res. **51**, D523–D531 (2023).

107.   Hornbeck, P. V. *et al.* PhosphoSitePlus, 2014: mutations, PTMs and

recalibrations. *Nucleic Acids Res.* **43**, D512-20 (2015).

108.   Junker, B. H., Klukas, C. & Schreiber, F. VANTED: a system for advanced data

analysis and visualization in the context of biological networks. *BMC*

Bioinformatics **7**, 109 (2006).

109.   Okuda, S. *et al.* jPOSTrepo: an international standard data repository for

proteomes. *Nucleic Acids Res.* **45**, D1107–D1111 (2017).

Fig. 1: Workflow for the construction and analysis of a global starvation-responsive trans-omic network in liver of WT and *ob/ob* mice

(a) Experimental design. WT and *ob/ob* mice were starved for the indicated hours, followed by the collection of liver and plasma samples. These samples were subjected to multi-omic measurements. It should be noted that metabolome was basically measured by CE-MS, but IC-QEMS was used for quantification of F1,6BP and F2,6BP. Enzymatic assay kit was used for glycogen. (b) Classification of molecules into 4 groups: commonly responsive in both WT and *ob/ob* mice (common, green); WT-specifically (WT-specific, blue); *ob/ob*-specifically (*ob/ob*-specific, red); oppositely (opposite, magenta) responsive molecules between WT mice and *ob/ob* mice. (c) Schema for a single-omic layer composed of responsive molecules classified as in (b). (d) Schema of regulatory input into the metabolic reactions. TFs control metabolic enzyme abundance through gene regulation. Protein kinases control metabolic enzyme activities through phosphorylation. The metabolites, FFA & acyls, and lipids act as substrates, products, and allosteric regulators in metabolic reactions. (e) Schematic representation of the global starvation-responsive trans-omic network consisting of the indicated layers. The subset of the network that is defined as the starvation-responsive metabolic network is outlined in orange. (f) Structural properties of the starvation-responsive metabolic network in WT and *ob/ob* mice. (g) Temporal order of the starvation-responsive metabolic network in WT and *ob/ob* mice. (h) Strength of co-regulation among molecules in the starvation-responsive metabolic network in WT and *ob/ob* mice.

Fig. 2: Single-omic layer analysis during starvation

(a-c) Identification and classification of starvation-responsive molecules. (a) One-way ANOVA tests ($P < 0.025$) defined starvation-responsive molecules. (b) Area under the curve (AUC) of log2-transformed mean time courses classified molecules into increased (AUC > 0, orange) and decreased (AUC < 0, purple) classes. (c) Classification of starvation-responsive molecules: common in both WT and *ob/ob* (either increased or decreased in both, green), WT-specific (blue), *ob/ob*-specific (red), or opposite (increased in WT and decreased in *ob/ob*, or vice versa, magenta). (d-j) The number of molecules and distribution in each group shown as pie charts colored as defined in (c). Increased or decreased molecules are indicated by the orange or purple arcs, respectively. The number and the ratio of significantly changed molecules of the molecules in each category are shown. Responsive metabolites (d), lipids (e), and FFA & acyls (f) in liver. Responsive plasma metabolites (g). Responsive transcripts (h), proteins (i), and phosphorylation events (j) in liver. (k) Strategy for inferring activity of upstream regulators and signaling pathways. Briefly, we constructed regulatory networks from database and identified increased and decreased downstream molecules. In the case of positive regulators, we assumed that regulator activity is up-regulated if most of the downstream molecules increased and vice versa (see method in detail). Up-regulated or down-regulated regulators were inferred and classified them into common (green), WT-specific (blue), *ob/ob*-specific (red), or opposite (magenta) groups. (l-n) The number and distribution of these inferred TFs (l), protein kinases (m), or signaling pathways (n) are shown in pie charts.

Fig. 3: Global starvation-responsive trans-omic network.

(a) Global starvation-responsive trans-omic network. Nodes in the layers represent metabolic reactions, transport activities, measured molecules, measured phosphorylation events, and the inferred activities of TF, kinase, and signaling pathway. Edges between the nodes represent regulatory connections identified using bioinformatic platforms. Colors represent the responsiveness: green indicates common responses both in WT and *ob/ob* mice; blue, specific to WT mice; red, specific to *ob/ob* mice; magenta, opposite responses between the WT and *ob/ob* mice. For metabolic reaction and transport_activity layers, the classes were defined by the existences of regulations regardless of the regulatory directions (activation or inhibition): green indicates the node is regulated in both WT and *ob/ob* mice; blue, specific to WT mice; red, specific to *ob/ob* mice. (b) The number and proportions of nodes and edges classified as common (green), WT-specific (blue), *ob/ob*-specific (red), and opposite (magenta) in the global starvation-responsive trans-omic network. Pie charts represent the distribution of nodes in the omic layers, and bar plots between pie charts represent the distribution of edges between omic layers. Type of regulation into the metabolic reactions for metabolite, FFA& acyls, and lipid are indicated as allosteric regulation (A) and substrate and product (S & P). The starvation-responsive metabolic network is outlined in orange, and layer names included in the starvation-responsive metabolic networks are bolded.

Fig. 4 Starvation-responsive metabolic network

Projected views of the starvation-responsive metabolic networks in WT (a) and *ob/ob* mice (b). The network is composed of direct regulators of metabolic reactions: metabolites, lipids, FFA & acyls, enzyme_proteins, and enzyme_PTMs. Colors represent the responsiveness: green indicates common responses both in WT and *ob/ob* mice; blue, specific to WT mice; red, specific to *ob/ob* mice; magenta, opposite responses between the two genotypes. Hub nodes, which will be defined in Fig. 5, are labeled.

Fig. 5: Structural properties of the starvation-responsive metabolic network

(a) Degree distributions of the starvation-responsive metabolic network in WT (blue) and *ob/ob* (red) mice. The x-axis represents the log10-transformed degree (k); the y-axis represents the log10-transformed number of nodes with degree of k. Lines show the linear regression of the degree distributions. The slope of the lines (γ), which will be defined as scaling parameters, and the 95% credible intervals are shown. (b) Diagram of a scale-free network, where n(k) denotes the number of nodes with degree of k, and γ is a constant called a scaling parameter. Circles represent nodes; lines represent edges. Light blue represents the hub nodes; yellow represents leaf nodes. (c and d) Hub nodes with the top 2% highest degree in WT (c) and *ob/ob* mice (d). The x-axis represents hub nodes. The y-axis represents degrees. Colors represent the responsiveness: green indicates common responses both in WT and *ob/ob* mice; blue, specific to WT mice; red, specific to *ob/ob* mice; magenta, opposite responses between WT and *ob/ob* mice. (e) Illustration of fragmentation of a network through loss of hub nodes. The largest cluster is orange. (f) Illustration of how communicability of a network influences average path length (average length of the shortest paths between all the pairs of nodes). (g) The relative size of the largest cluster of the starvation-responsive metabolic network in WT and *ob/ob* mice, calculated as the number of nodes in the largest cluster of the network divided by the total number of responsive nodes in the network for each genotype. (h) The average path length of the starvation-responsive metabolic network in WT and *ob/ob* mice.

Fig. 6: Temporal order of the starvation-responsive metabolic network in WT mice and its disruption in *ob/ob* mice

(a and b) Scatter plot for the log10-transformed degree (x-axis) and *t*-half (y-axis) in WT (a) and *ob/ob* (b) mice. Dots represent the molecules (including phosphorylation events) in the starvation-responsive metabolic network and colors denote the responsiveness: green indicates common responses both in WT and *ob/ob* mice; blue, specific to WT mice; red, specific to ob/ob mice; magenta, opposite responses between WT and *ob/ob* mice. Spearman's correlation coefficient (Rho) and *P* values are shown. Hub molecules for each genotype are labeled (see Fig. 5c and d). (c) Values of t-half for the WT-specific and ob/ob-specific hub molecules (see Fig. 5c and d). The P value of Wilcoxon's rank sum test is shown. (d and e) Distributions of t-half values of molecules (metabolites, lipids, FFA & acyls, proteins, and phosphorylation events) in the indicated metabolic pathways in WT (d) and *ob/ob* mice (e). The data are ordered along the x-axis by the median *t*-half (red line) with each molecule represented by a dot. *P* values of the Kruskal-Wallis test are shown above the panels and those of the Dunn-Sidak test are shown above the lines indicating the pairs of metabolic pathways with significant differences. (f and g) Distributions of t-half values in each omic layer in the starvation-responsive metabolic network in WT (f) and in *ob/ob* (g) mice. The data are ordered along the x-axis by the median *t*-half (red line). *P* values of the Kruskal-Wallis test are shown above the panels and those of the Dunn-Sidak test are shown above the lines indicating the pairs of omic layers with significant differences.

Fig. 7: Co-regulation among molecules in the starvation-responsive metabolic network and disruption in *ob/ob* mice

(a) Schematic representation of the analyses for co-regulation of the molecules in the starvation-responsive metabolic network. Time courses of molecules sharing a common regulatory mechanism could be positively or negatively correlated, defined as co-regulation. Pairwise Pearson's correlation coefficients were calculated for the time courses of molecules that showed significant responses during starvation in the starvation-responsive metabolic network in WT and *ob/ob* mice. Positive correlation represents positive co-regulation and negative correlation represents negative co-regulation. (b) Distribution of the pairwise Pearson's correlation coefficients for time courses of responsive molecules in the starvation-responsive metabolic network in WT and *ob/ob* mice. The x-axis represents correlation coefficients; the y-axis represents the number of pairs of molecules examined for correlation. Pairs with an absolute value of correlation coefficient > 0.9 were connected by edges in the correlation metabolic network (see Fig. S6). (c and d) The correlation metabolic network during starvation in WT (c) and *ob/ob* mice (d). Orange indicates the largest cluster; all other clusters are dark green. Individual orphan nodes with no connections are shown only by numbers.