A computational approach to understanding effort-based decision-making in depression

Vincent Valton, PhD*[1]; Anahit Mkrtchian, PhD*[2]; Madeleine Moses-Payne, PhD[3]; Alan Gray, MSc[1]; Karel Kieslich, MUDr[1]; Samantha VanUrk, MSc[1]; Veronika Samborska, PhD[1]; Don Chamith Halahakoon, DPhil[1]; Sanjay G. Manohar, PhD[4]; Peter Dayan, PhD[5]; Masud Husain, DPhil[4] & Jonathan P. Roiser, PhD[1]

[1] Institute of Cognitive Neuroscience, University College London, London, UK

[2] Division of Psychiatry and Max Planck Centre for Computational Psychiatry and Ageing Research, Queen Square Institute of Neurology, University College London, London, UK

[3] Department of Clinical, Educational and Health Psychology, University College London, London, UK

[4] Nuffield Department of Clinical Neurosciences and Department of Experimental Psychology, Oxford University, Oxford, UK

[5] Max Planck Institute for Biological Cybernetics and the University of Tübingen, Tübingen, Germany

* These authors contributed equally to the manuscript

Correspondence
Anahit Mkrtchian
UCL Division of Psychiatry
149 Tottenham Ct Rd
London
W1T 7NF
UK

Jonathan Roiser
UCL Institute of Cognitive Neuroscience
17 Queen Square
London
WC1N 3AZ
UK

Word count: 2980

Running Title: Effort-based decision-making in depression

Keywords: Effort-based decision-making; Depression; Computational Psychiatry; Motivation; Anhedonia

Key Points

Question: What are the computational mechanisms underlying altered motivation in depression?

Findings: In a case-control study, we find lower willingness to exert effort in both currently symptomatic and remitted depressed individuals (all unmedicated), compared with healthy volunteers. Detailed computational analysis revealed that this difference was driven by a lower overall effort acceptance bias, but not altered sensitivity to either reward or effort.

Meaning: Lower willingness to engage in effort in depression is driven by an overall bias against exertion, which may represent a trait-like factor, presenting a potentially valuable target for treatment and prevention.

Abstract

Importance: Motivational dysfunction is a core feature of depression, and can have debilitating effects on everyday function. However, it is unclear which disrupted cognitive processes underlie impaired motivation, and whether impairments persist following remission. Decision-making concerning exerting effort to obtain rewards offers a promising framework for understanding motivation, especially when examined with computational tools which can offer precise quantification of latent processes.

Objective: To understand the computational mechanisms driving motivational dysfunction in depression.

Design, Setting, and Participants: We conducted two studies: a Pilot study in healthy volunteers (N=67, 66% female, mean[SD] age=28.45[9.88]) to validate our computational model, before applying it in a Case-control study including current (N=41, 71% female, mean[SD] age=30.24[11.57]) and remitted (N=46, 63% female, mean[SD] age=26.91[7.06]) unmedicated depressed individuals, and healthy volunteers with (N=36, 64% female, mean[SD] age=26.06[8.19]) and without (N=57, 68% female, mean[SD] age=26.70[8.14]) a family history of depression. The Pilot study data was collected during 2015 and the Case-control study data was collected between 2015 and 2019.

Exposures: Effort-based decision-making was assessed using the Apple Gathering Task, in which participants decide whether to exert effort via a grip-force device to obtain varying levels of reward; effort levels were individually calibrated and varied parametrically.

Main Outcome and Measures: The probability to accept offers as a function of reward and effort levels was examined. A comprehensive Bayesian computational analysis was implemented to examine the precise computational mechanisms influencing decision-making.

Results: Four fundamental computational mechanisms that drive patterns of effort-based decisions, which replicated across samples, were identified: overall bias to accept effort

challenges; reward sensitivity; and linear and quadratic effort sensitivity. Traditional model-agnostic analyses showed that both depressed groups had a lower willingness to exert effort than control participants. In contrast with previous findings, computational analysis revealed that this difference was primarily driven by lower effort acceptance bias, but not altered effort or reward sensitivity.

Conclusion and Relevance: This work provides insight into the computational mechanisms underlying motivational dysfunction in depression. Lower willingness to exert effort could represent a trait-like factor contributing to both symptoms and risk of relapse, and might represent a fruitful target for treatment and prevention.

Introduction

Motivational impairment is common in depression,[1-3] closely linked to anhedonia, decision-making difficulties and fatigue, which together predict poorer treatment outcome[4] and lower quality of life[5]. Although anhedonia—the loss of interest or pleasure in previously enjoyable activities—is a cardinal symptom of depression, in-the-moment pleasurable experience appears to be relatively preserved in depressed individuals[6-11]; by contrast, reward learning and decision-making have repeatedly been shown to be impaired in depression[12], consistent with disrupted motivation.

A consistent theme emerging from this body of work is that *effort-based decision-making for reward* offers a promising lens through which motivational dysfunction can be understood. A ubiquitous finding is that the willingness to engage in effort (physical or mental) depends on both the perceived reward magnitude, and the discounting of that reward by the effort required to obtain it[13]. While numerous studies have examined effort discounting in healthy individuals[13-18], and contemporary neurocognitive models of depression suggest that lower willingness to exert effort drives depressive symptoms related to motivation[19-21], few studies have investigated these processes in patients experiencing motivational symptoms.

The first systematic attempt to examine effort-based decision-making in depression (i.e., measuring *explicit choices* to exert effort, as opposed to the degree of exertion) was reported by Treadway and colleagues (2012), who compared performance on the *Effort Expenditure for Rewards Task* (EEfRT[22]) between currently depressed individuals and healthy volunteers. Drawing on an extensive literature in rodents[23], the EEfRT requires participants to choose between a "hard" versus an "easy" task (quick versus slow button pressing), with more reward delivered for hard choices, with varying levels of probabilities. Treadway and

colleagues reported that depressed participants were less willing to choose the hard task, and less sensitive to both reward and probability. Since this initial study, several others have reported comparable results in currently depressed individuals[24-26], although with discrepant results for remitted depression[25,27].

Although these studies provided an important foundation for advancing our understanding of disrupted motivation in depression, several design features complicate their interpretation. First, most studies, including those examining remitted depression, recruited individuals taking antidepressant medication, which represents a potentially important confound given that antidepressants are known to blunt neural reward processing[28]. Second, the exertion level required to obtain reward was typically not calibrated individually; making the hard task easier for some participants than others. Third, the inclusion of different probability conditions (with no deterministic condition) raises the possibility of an interaction between probability and effort discounting. Fourth, effort levels are not varied parametrically in the EEfRT, and no high-reward/low-effort or low-reward/high-effort options are included. Consequently, individual differences in choices of the high-reward/high-effort option could be driven by sensitivity to either reward or effort.

Another effort-based decision task, the Apple Gathering Task (AGT[29]), offers several key advantages: a grip-strength device is used for exertion[30,31], with force level calibrated to each participant; reward and effort are varied independently and parametrically; and outcomes are deterministic. Studies using the AGT have reported that patients with Parkinson's disease (PD; in which depression is very common[32]) were less willing to exert effort for low rewards[33], especially those with pronounced apathy[34]. Other studies found that experimentally induced inflammation increased sensitivity to effort in healthy volunteers[35],

and that, relative to matched controls, patients with treatment-resistant schizophrenia were both less willing to exert effort overall and less sensitive to reward[36]. However, no study to date has reported behaviour on the AGT in depression.

Over the past decade, computational modelling has provided important insights into the cognitive processes underlying disrupted motivation[37]. A computational approach systematically evaluates competing models to identify the latent cognitive processes governing behaviour. Computational modelling enables the precise measurement of cognitive processes by estimating specific parameters, offering deeper insights than traditional (model-agnostic) measures of performance, often with superior psychometric properties[38,39].

Therefore, in the present study we performed a computational analysis of AGT performance to provide insights into the cognitive processes underlying effort-based decision-making in depression. To address the question of whether altered effort-based decision-making is merely a consequence of ongoing depressive symptoms, or if it plays a causal role in their development, we studied four groups, all unmedicated: currently depressed individuals (MDD); individuals remitted from depression (REM); healthy individuals with a depressed close relative, but no personal history (REL, who are known to be at elevated risk); and healthy individuals without any personal or family history of psychiatric diagnosis (CTR).

Drawing on contemporary neurocognitive models of depression[19-21,40], we hypothesised that lower willingness to exert effort drives depressive symptoms, especially those related to motivation. We predicted that the MDD group would accept fewer offers to exert effort relative to the CTR group, and that this would be driven by lower sensitivity to reward and greater sensitivity to effort. Including the REL and REM groups allowed us to examine

whether effort-based decision-making represents a risk factor for depression, and we predicted that a similar pattern of behaviour would be observed in these two groups.

Materials and Methods

Participants

Two studies were conducted, an initial pilot with healthy volunteers (HVs; "Pilot"), and a study including four groups ("Case-control").

Pilot study

Only HVs (N=102) completed this study. The final sample consisted of 67 participants (see Supplemental Information [SI] for exclusions).

Case-control study

Sixty-two HVs without a family history of depression (CTR; independent from the Pilot study), 38 HVs with a depressed first-degree relative (REL), 50 remitted depressed participants (REM), and 51 currently depressed participants (MDD) completed the Case-control study. The final sample included 57 CTR, 36 REL, 46 REM and 41 MDD participants (see SI for exclusions). With this sample size, at alpha=0.05, we had 80% power to detect effect sizes of at least f=0.25 (medium effect size).

Experimental procedure

For the Pilot study eligibility was assessed using a structured telephone interview. For the Case-control study, participants attended a screening session, completing the Mini International Neuropsychiatric Interview[41], the Family Interview for Genetic Studies[42] and symptom questionnaires before returning for cognitive testing in a separate session.

In both studies, participants completed the digit span forwards and backwards[43], the Wechsler Test of Adult Reading (WTAR[44]), the Apathy Evaluation Scale (AES[45]), the Beck Depression Inventory (BDI-II[46]), the Dysfunctional Attitudes Scale (DAS-SF1-2[7]), the Life Orientation Test-Revisited (LOTR[47]), the Snaith-Hamilton Pleasure Scale (SHAPS[48]), the State-Trait Anxiety Inventory for Adults (STAI[49]), and the Temporal Experience of Pleasure Scale (TEPS[50]). In the Pilot study, participants completed the Chapman Physical Anhedonia Scale (CPAS[51]). In the Case-control study, participants completed the HAM-D.

Apple Gathering Task (AGT)

The AGT measures willingness to exert physical effort for reward[33]. Before the main task, participants performed a six-trial calibration phase, squeezing a hand-dynamometer as hard as possible with their non-dominant hand to fill an on-screen gauge. The peak force from the final three trials determined their maximum voluntary contraction (MVC), setting comparable effort levels across subjects. Participants then attempted four effort levels (20%, 40%, 60% and 80% of MVC).

On each trial in the main task, participants saw a tree containing a number of apples representing the available reward, and a bar on the tree trunk representing the force required to obtain them (Figure 1A; see SI). There were four levels of reward (3, 6, 9, or 12 apples) and effort (20%, 40%, 60% and 80%). Each reward/effort combination was repeated five times, resulting in a total of 80 trials. Participants could accept or refuse offers. For refused offers, 'no response required' was displayed, followed by the next decision. An accepted offer required squeezing the hand-dynamometer at or above the effort level to win points, followed by feedback. The next trial started immediately after the feedback. To mitigate fatigue, the exertion phase was omitted and "No response required" appeared on 25% of accepted trials.

Statistical Analysis

Repeated-measures analysis of variance (ANOVA) was used to examine how acceptance rates varied as a function of reward and effort. For the Case-control study, group was added as a between-subjects measure. Greenhouse-Geisser correction was applied when sphericity violations occurred. Acceptance rates were arcsine transformed prior to analysis to satisfy Gaussian assumptions. Decision reaction times, success rates and fatigue effects were also analysed (see SI). Covariates in analyses (age, see below) were mean corrected.

Questionnaire data was synthesized using exploratory factor analysis on the total score of each questionnaire using Promax rotation (see SI).

Computational Analysis

Computational analysis was performed to compare competing hypotheses of reward and effort contributions to decision-making, and to estimate participant-level parameters. All models were implemented using hierarchical Bayesian estimation in Stan[52]. Participants in the Pilot study were fit under the same group-level priors. For the Case-control study, all participants were fit under the same prior for parameters of no interest and separate group-level priors were used for the parameter of interest (one parameter at a time; see SI; Figure S1). Sensitivity analyses with separate group priors did not materially impact the estimated parameters or group comparison results (Figure S2).

Winning model

Seventy models of varying complexity were compared to select the winning model (see SI). The winning model in both studies was a four-parameter model including two effort

sensitivity terms (linear: LinE - more negative=greater subjective effort, and quadratic: E^2 – effort cost increases disproportionately with increasing effort when $E^2<0$), a linear reward sensitivity term (LinR; more positive=greater subjective reward), and an acceptance bias term (K) representing the overall tendency to accept offers independent of reward or effort (higher=more likely to accept; Figure S1). In the Case-control study, owing to high trade-off between the LinE and E^2 parameters and limited LinE range, the LinE term was constrained to that of the average value estimated from the Pilot study (ConstE=-15; see SI). Together these parameters could accurately recapitulate both the group-level pattern of choices and individual differences in acceptance rates (Figure S3; Pilot: r=0.997; Case-control: r=0.999). All parameters showed high recoverability (Figure S4).

Results

Participant characteristics are presented in Table 1.

Model-agnostic AGT analysis

In preliminary analyses we found that older individuals accepted more offers overall (Pearson r=0.20, p=0.014), and therefore age was included as a covariate in all Case-control analyses. We observed wide variability in overall acceptance rates, ranging from 35-100% in the Pilot and 40-100% in the Case-control study (Figure 1C&E). As expected, acceptance rates decreased significantly as effort increased (Pilot: $F(1.57,103.77)=120.19$, $p<0.001$; Case-control: $F(1.75,306.31)=351.79$, $p<0.001$), and increased significantly as reward increased (Pilot: $F(1.63,107.57)=92.02$, $p<0.001$; Case-control: $F(1.65,288.06)=325.45$, $p<0.001$). There was a significant reward-by-effort interaction in both studies, driven by particularly high acceptance rates for high-reward/low-effort trials (Pilot: $F(4.25,280.88)=16.42$, $p<0.001$; Case-control: $F(4.95,866.79)=90.13$, $p<0.001$; Figure 1B&D).

In the Case-control study we detected a significant main effect of group ($F(3,175)=3.26$, $p=0.023$; Figure 2), but all interactions with group were non-significant. Post-hoc tests revealed significantly lower acceptance rates for the REM and MDD groups compared to REL (REM vs REL: Cohen's $d=0.63$, $p=0.007$; MDD vs REL: $d=0.41$, $p=0.015$; reported effect sizes exclude covariates).

There were no significant group differences in success rates, decision RTs, MVCs or either immediate or cumulative fatigue (all $p>0.05$, see SI; Figure S5&S6). Importantly, success rate at the highest effort level was above 80% across all groups.

Questionnaire factor analysis

Factor analysis was performed on questionnaire measures for both studies. Despite minor differences in measures (CPAS and the short form of the DAS was only included in the Pilot; HAM-D was only included in the Case-control study), a similar four-factor solution was obtained in both studies (Table S1). The factors were (named according to the scales displaying the highest loadings (>0.3)): "Low-mood", with loadings for BDI-II, HAM-D, LOTR and STAI; "Apathy", mostly comprising the AES sub-scales; "Hedonia", including the SHAPS and TEPS (owing to scoring conventions these load with opposite signs), and (in the Pilot) the CPAS; and "Dysfunctional Attitudes", comprising the two DAS sub-scales.

Computational AGT analysis

Model comparison

The winning model in both studies was a derivative of a four-parameter model including an acceptance bias parameter (K), linear reward (LinR) and effort (LinE) sensitivity parameters,

and a quadratic effort parameter (E^2; Figure 3; Figures S7&8). In the Case-control study, the LinE term was set at ConstE=-15 (see Materials and Methods and SI).

Case-control comparisons

Participant-level parameters were compared between the groups including age as a covariate (Figure 3). The K (acceptance bias) parameter differed significantly between the groups ($F_{(3,175)}=3.19$, p=0.025). Post-hoc tests revealed that this was driven by the REM (d=0.61, p=0.006) and MDD (d=0.41, p=0.032) groups having a lower acceptance bias than the REL group, and the REM group having a lower acceptance bias than the HC group (d=0.41, p=0.04). Planned comparisons revealed that the combined REM+MDD group had a lower acceptance bias than the combined REL+CTR group (d=0.39, $F_{(1,177)}=8.26$, p=0.005). However, the combined REL+REM+MDD group did not differ from the CTR group (d=0.16, p=0.276). No other parameter differed significantly between groups (LinR: $F_{(3,175)}=0.33$, p=0.806; E^2: $F_{(3,175)}=0.13$, p=0.942).

The above analysis suggests that lower acceptance rates in the depression groups were driven by a lower overall willingness to exert effort, not alterations in reward or effort sensitivity.

Associations between parameters and symptoms

Pilot study

In the Pilot study, LinR correlated negatively with Low Mood (r=-0.344, p=0.004; Figure 4), suggesting that participants with greater anxiety/depression perceived rewards as less valuable. The E^2 parameter correlated negatively with Low Mood (r=-0.25, p=0.04) and positively with Hedonia (r=0.248, p=0.043), suggesting that participants with greater

anxiety/depression and anhedonia perceived increasing levels of effort as disproportionately costly.

Case-control study

In the Case-control study, no significant associations between symptom factors and computational parameters were observed, either in the combined CTR+REL+REM sample or in the MDD group alone. In the combined CTR+REL group, surprisingly (and in contrast to the corresponding result in the Pilot study) LinR correlated with Low Mood (r=0.248, p=0.016; Figure 4).

Discussion

Impaired motivation is a hallmark of depression, but the underlying cognitive and computational processes are poorly understood. Consistent with our hypotheses, we found that depression is characterised by a lower willingness to engage in effort. Our computational analyses suggest that this difference was not driven by altered effort or reward sensitivity, but rather by a lower overall tendency to avoid exerting effort, reflected in the acceptance bias parameter. The presence of lower acceptance bias in both the remitted and currently depressed groups suggests that this decision-making process could represent a trait-like feature, as proposed by neurocognitive models of depression[19-21,40], or possibly a "scar" effect of having previously experienced depression[53].

These findings align with previous work demonstrating lower willingness to exert effort for reward in depression[22,24-26,54] but challenge a key assumption that mis-calibrated reward or effort valuation drives this behaviour[18,55,56]. By examining a large space of computational models, we instead suggest that a different process is at play: acceptance bias. While

surprising and not entirely straightforward to interpret, this result clarifies prior findings from tasks that were unable to disambiguate between these factors. One possibility is that lower acceptance bias could be driven by lower confidence in being able to achieve the required effort, despite individual calibration. This would indicate that metacognitive processes might play a role in motivational impairment. Indeed, previous studies have identified that transdiagnostic symptoms of apathy, poor self-esteem and low mood are associated with low confidence in perceptual decisions[57,58], independent of accuracy, mirroring our findings. This could plausibly be related to low global expectations of success in depression[59-61]. It is also possible that fatigue may influence acceptance bias, although we did not detect any group differences in sensitivity analyses examining task-related fatigue.

An important goal of neuroscientific research in depression is to determine whether cognitive disruptions are simply a consequence of depressive symptoms, or whether they contribute causally to their development[62]. We observed a similar pattern of effort-based decisions for reward in both current and remitted depressed groups. This suggests that a general bias against exerting effort might represent a core feature of depression. However, we did not observe a similar pattern in never-depressed first-degree relatives. Therefore, an alternative interpretation might be that lower willingness to exert effort is a consequence rather than a cause of depression, which does not recover after symptoms remit (akin to a 'scar' effect). A third possibility is that the first-degree relatives we tested were not at elevated risk for depression, but were instead actually resilient: depression often presents early in life and our REL sample was older (mean age~26) than the peak onset of depressive symptoms[63,64]. Thus, many individuals in this group might never go on to develop depression. Future studies should recruit younger samples and test these predictions longitudinally.

If confirmed as a risk factor for depression, lower overall willingness to exert effort may represent a fruitful target for intervention. Such targets are sorely needed for motivational symptoms, which are particularly difficult to treat and constitute an area of unmet clinical need[4,65]. Treatments that can boost engagement in activities through altering effort acceptance bias could be particularly effective in treating or even preventing depression. Elements of behavioural activation therapy, such as activity scheduling, might play such a role, as these require acting on pre-planned commitments rather than internal states[66]. Interestingly, this component has recently been shown to alter effort processing in healthy participants, offering proof-of-concept of this idea[67].

Limitations

While our model of effort-based decision-making replicated across studies, relationships with specific symptom factors were inconsistent, potentially owing to the limited sample size of the Case-control study. Larger studies are needed to confirm these associations. An important strength of the effort task we employed is its individually tailored effort levels, which resulted in >80% average success rates at all effort levels. Nonetheless, it remains possible that occasional failures to obtain reward following exertion could have influenced decisions in some participants. Importantly, however, there was no group difference in success rates—or in effort sensitivity—making it unlikely that the lower effort acceptance bias we observed in depression was influenced by success rates.

Conclusion

This study advances our understanding of motivational impairment in depression by identifying low effort acceptance bias, independent of reward or effort sensitivity, as a

computational feature of both current and remitted depression. These results raise the possibility that interventions that boost this bias could improve treatment outcomes.

Acknowledgments and Disclosures

This research was funded in whole, or in part, by a Wellcome Trust grant (101798/Z/13/Z) to JPR. For the purpose of Open Access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission. AM was supported by a Wellcome Trust grant (221826/Z/20/Z), PD was funded by the Max Planck Society and the Humboldt Foundation, SGM was supported by an MRC Clinician Scientist Fellowship MR/P00878/X and by the National Institute of Healthcare Research (NIHR) Oxford Biomedical Research Centre (BRC). The authors report no biomedical financial interests or potential conflicts of interest.

References

1.   Pelizza L, Ferrari A. Anhedonia in schizophrenia and major depression: state or trait? *Ann Gen Psychiatry*. Oct 8 2009;8:22. doi:10.1186/1744-859x-8-22

2.   Yuen GS, Bhutani S, Lucas BJ, et al. Apathy in late-life depression: common, persistent, and disabling. *Am J Geriatr Psychiatry*. May 2015;23(5):488-94. doi:10.1016/j.jagp.2014.06.005

3.   Fervaha G, Foussias G, Takeuchi H, Agid O, Remington G. Motivational deficits in major depressive disorder: Cross-sectional and longitudinal relationships with functional impairment and subjective well-being. *Comprehensive Psychiatry*. 2016/04/01/ 2016;66:31-38. doi:https://doi.org/10.1016/j.comppsych.2015.12.004

4.   Uher R, Perlis RH, Henigsberg N, et al. Depression symptom dimensions as predictors of antidepressant treatment outcome: replicable evidence for interest-activity symptoms. *Psychological Medicine*. 2012;42(5):967-980. doi:10.1017/S0033291711001905

5.   Spijker J, Bijl RV, De Graaf R, Nolen WA. Determinants of poor 1-year outcome of DSM-III-R major depression in the general population: results of the Netherlands Mental Health Survey and Incidence Study (NEMESIS). *Acta Psychiatrica Scandinavica*. 2001;2:122-130. doi:https://doi.org/10.1034/j.1600-0447.2001.103002122.x

6.   Amsterdam JD, Settle RG, Doty RL, Abelman E, Winokur A. Taste and smell perception in depression. *Biol Psychiatry*. Dec 1987;22(12):1481-5. doi:10.1016/0006-3223(87)90108-9

7.   Beevers CG, Strong DR, Meyer B, Pilkonis PA, Miller IR. Efficiently assessing negative cognition in depression: an item response theory analysis of the Dysfunctional Attitude Scale. *Psychol Assess*. Jun 2007;19(2):199-209. doi:10.1037/1040-3590.19.2.199

8.   Bylsma LM, Taylor-Clift A, Rottenberg J. Emotional reactivity to daily events in major and minor depression. *J Abnorm Psychol*. Feb 2011;120(1):155-67. doi:10.1037/a0021662

9.   Chentsova-Dutton Y, Hanley K. The effects of anhedonia and depression on hedonic responses. *Psychiatry Res*. Sep 30 2010;179(2):176-80. doi:10.1016/j.psychres.2009.06.013

10.   Dichter GS, Smoski MJ, Kampov-Polevoy AB, Gallop R, Garbutt JC. Unipolar depression does not moderate responses to the Sweet Taste Test. *Depress Anxiety*. Sep 2010;27(9):859-63. doi:10.1002/da.20690

11.   Peeters F, Nicolson NA, Berkhof J, Delespaul P, deVries M. Effects of daily events on mood states in major depressive disorder. *J Abnorm Psychol*. 2003;112:203–211.

12.   Halahakoon DC, Kieslich K, O'Driscoll C, Nair A, Lewis G, Roiser JP. Reward-Processing Behavior in Depressed Participants Relative to Healthy Volunteers: A Systematic Review and Meta-analysis. *JAMA Psychiatry*. Dec 1 2020;77(12):1286-1295. doi:10.1001/jamapsychiatry.2020.2139

13.   Chong TT, Apps M, Giehl K, Sillence A, Grima LL, Husain M. Neurocomputational mechanisms underlying subjective valuation of effort costs. *PLoS Biol*. Feb 2017;15(2):e1002598. doi:10.1371/journal.pbio.1002598

14.   Bonnelle V, Manohar S, Behrens T, Husain M. Individual Differences in Premotor Brain Systems Underlie Behavioral Apathy. *Cereb Cortex*. Feb 2016;26(2):807-819. doi:10.1093/cercor/bhv247

15.   Croxson PL, Walton ME, O'Reilly JX, Behrens TE, Rushworth MF. Effort-based cost-benefit valuation and the human brain. *J Neurosci*. Apr 8 2009;29(14):4531-41. doi:10.1523/jneurosci.4515-08.2009

16.   Klein-Flügge MC, Kennerley SW, Friston K, Bestmann S. Neural Signatures of Value Comparison in Human Cingulate Cortex during Decisions Requiring an Effort-Reward

Trade-off. *The Journal of Neuroscience*. 2016;36(39):10002-10015. doi:10.1523/jneurosci.0292-16.2016

17. Kurniawan IT, Guitart-Masip M, Dayan P, Dolan RJ. Effort and valuation in the brain: the effects of anticipation and execution. *J Neurosci*. Apr 3 2013;33(14):6160-9. doi:10.1523/jneurosci.4777-12.2013

18. Pessiglione M, Vinckier F, Bouret S, Daunizeau J, Le Bouc R. Why not try harder? Computational approach to motivation deficits in neuro-psychiatric diseases. *Brain*. Mar 1 2018;141(3):629-650. doi:10.1093/brain/awx278

19. Eshel N, Roiser JP. Reward and punishment processing in depression. *Biol Psychiatry*. Jul 15 2010;68(2):118-24. doi:10.1016/j.biopsych.2010.01.027

20. Roiser JP, Elliott R, Sahakian BJ. Cognitive mechanisms of treatment in depression. *Neuropsychopharmacology*. Jan 2012;37(1):117-36. doi:10.1038/npp.2011.183

21. Pizzagalli DA. Toward a Better Understanding of the Mechanisms and Pathophysiology of Anhedonia: Are We Ready for Translation? *Am J Psychiatry*. Jul 2022;179(7):458-469. doi:10.1176/appi.ajp.20220423

22. Treadway MT, Buckholtz JW, Schwartzman AN, Lambert WE, Zald DH. Worth the 'EEfRT'? The effort expenditure for rewards task as an objective measure of motivation and anhedonia. *PLoS One*. Aug 12 2009;4(8):e6598. doi:10.1371/journal.pone.0006598

23. Salamone JD, Yohn SE, López-Cruz L, San Miguel N, Correa M. Activational and effort-related aspects of motivation: neural mechanisms and implications for psychopathology. *Brain*. May 2016;139(Pt 5):1325-47. doi:10.1093/brain/aww050

24. Yang XH, Huang J, Lan Y, et al. Diminished caudate and superior temporal gyrus responses to effort-based decision making in patients with first-episode major depressive disorder. *Prog Neuropsychopharmacol Biol Psychiatry*. Jan 4 2016;64:52-9. doi:10.1016/j.pnpbp.2015.07.006

25. Yang XH, Huang J, Zhu CY, et al. Motivational deficits in effort-based decision making in individuals with subsyndromal depression, first-episode and remitted depression patients. *Psychiatry Res*. Dec 30 2014;220(3):874-82. doi:10.1016/j.psychres.2014.08.056

26. Zou YM, Ni K, Wang YY, et al. Effort-cost computation in a transdiagnostic psychiatric sample: Differences among patients with schizophrenia, bipolar disorder, and major depressive disorder. *Psych J*. Apr 2020;9(2):210-222. doi:10.1002/pchj.316

27. Berwian IM, Wenzel JG, Collins AGE, et al. Computational Mechanisms of Effort and Reward Decisions in Patients With Depression and Their Association With Relapse After Antidepressant Discontinuation. *JAMA Psychiatry*. May 1 2020;77(5):513-522. doi:10.1001/jamapsychiatry.2019.4971

28. McCabe C, Mishor Z, Cowen PJ, Harmer CJ. Diminished neural processing of aversive and rewarding stimuli during selective serotonin reuptake inhibitor treatment. *Biol Psychiatry*. Mar 1 2010;67(5):439-45. doi:10.1016/j.biopsych.2009.11.001

29. Bonnelle V, Veromann KR, Burnett Heyes S, Lo Sterzo E, Manohar S, Husain M. Characterization of reward and effort mechanisms in apathy. *J Physiol Paris*. Feb-Jun 2015;109(1-3):16-26. doi:10.1016/j.jphysparis.2014.04.002

30. Cléry-Melin ML, Schmidt L, Lafargue G, Baup N, Fossati P, Pessiglione M. Why don't you try harder? An investigation of effort production in major depression. *PLoS One*. 2011;6(8):e23178. doi:10.1371/journal.pone.0023178

31. Schmidt L, d'Arc BF, Lafargue G, et al. Disconnecting force from money: effects of basal ganglia damage on incentive motivation. *Brain*. May 2008;131(Pt 5):1303-10. doi:10.1093/brain/awn045

32. Costello H, Husain M, Roiser JP. Apathy and Motivation: Biological Basis and Drug Treatment. *Annu Rev Pharmacol Toxicol*. Jan 23 2024;64:313-338. doi:10.1146/annurev-pharmtox-022423-014645

33.   Chong TT, Bonnelle V, Manohar S, et al. Dopamine enhances willingness to exert effort for reward in Parkinson's disease. *Cortex*. Aug 2015;69:40-6. doi:10.1016/j.cortex.2015.04.003

34.   Le Heron C, Plant O, Manohar S, et al. Distinct effects of apathy and dopamine on effort-based decision-making in Parkinson's disease. *Brain*. 2018;141(5):1455-1469. doi:10.1093/brain/awy110

35.   Draper A, Koch RM, van der Meer JW, et al. Effort but not Reward Sensitivity is Altered by Acute Sickness Induced by Experimental Endotoxemia in Humans. *Neuropsychopharmacology*. Apr 2018;43(5):1107-1118. doi:10.1038/npp.2017.231

36.   Saleh Y, Jarratt-Barnham I, Petitet P, Fernandez-Egea E, Manohar SG, Husain M. Negative symptoms and cognitive impairment are associated with distinct motivational deficits in treatment resistant schizophrenia. *Mol Psychiatry*. Aug 25 2023;doi:10.1038/s41380-023-02232-7

37.   Adams RA, Huys QJ, Roiser JP. Computational Psychiatry: towards a mathematically informed understanding of mental illness. *J Neurol Neurosurg Psychiatry*. Jan 2016;87(1):53-63. doi:10.1136/jnnp-2015-310737

38.   Karvelis P, Paulus MP, Diaconescu AO. Individual differences in computational psychiatry: A review of current challenges. *Neuroscience & Biobehavioral Reviews*. 2023/05/01/ 2023;148:105137. doi:https://doi.org/10.1016/j.neubiorev.2023.105137

39.   Mkrtchian A, Valton V, Roiser JP. Reliability of Decision-Making and Reinforcement Learning Computational Parameters. *Computational Psychiatry*. 2023;doi:10.5334/cpsy.86

40.   Pringle A, McCabe C, Cowen PJ, Harmer CJ. Antidepressant treatment and emotional processing: can we dissociate the roles of serotonin and noradrenaline? *J Psychopharmacol*. Aug 2013;27(8):719-31. doi:10.1177/0269881112474523

41.   Sheehan DV, Lecrubier Y, Harnett Sheehan K, et al. The validity of the Mini International Neuropsychiatric Interview (MINI) according to the SCID-P and its reliability. *European Psychiatry*. 1997/01/01/ 1997;12(5):232-241. doi:https://doi.org/10.1016/S0924-9338(97)83297-X

42.   Maxwell ME. Family Interview for Genetic Studies (FIGS): a manual for FIGS. *Bethesda, MD: Clinical Neurogenetics Branch, Intramural Research Program, National Institute of Mental Health*. 1992;

43.   Wechsler D. WMS-III Administration and Scoring Manual. San Antonio, TX: The Psychological Corporation. Harcourt Brace & Co; 1997.

44.   Holdnack H. *Wechsler test of adult reading: WTAR*. The Psychological Corporation; 2001.

45.   Marin RS, Biedrzycki RC, Firinciogullari S. Reliability and validity of the Apathy Evaluation Scale. *Psychiatry Res*. Aug 1991;38(2):143-62. doi:10.1016/0165-1781(91)90040-v

46.   Beck AT, Steer RA, Brown GK. Beck depression inventory. 1996;

47.   Scheier MF, Carver CS, Bridges MW. Distinguishing optimism from neuroticism (and trait anxiety, self-mastery, and self-esteem): a reevaluation of the Life Orientation Test. *J Pers Soc Psychol*. Dec 1994;67(6):1063-78. doi:10.1037//0022-3514.67.6.1063

48.   Snaith RP, Hamilton M, Morley S, Humayan A, Hargreaves D, Trigwell P. A scale for the assessment of hedonic tone the Snaith-Hamilton Pleasure Scale. *Br J Psychiatry*. Jul 1995;167(1):99-103. doi:10.1192/bjp.167.1.99

49.   Spielberger CD. Manual for the State-Trait Anxiety Inventory (self-evaluation questionnaire). *(No Title)*. 1970;

50.   Gard DE, Gard MG, Kring AM, John OP. Anticipatory and consummatory components of the experience of pleasure: a scale development study. *Journal of research in personality*. 2006;40(6):1086-1102.

51.   Chapman LJ, Chapman JP, Raulin ML. Scales for physical and social anhedonia. *J Abnorm Psychol*. Aug 1976;85(4):374-82. doi:10.1037//0021-843x.85.4.374

52.   Carpenter B, Gelman A, Hoffman MD, et al. Stan: A Probabilistic Programming Language. *J Stat Softw*. 2017;76doi:10.18637/jss.v076.i01

53.   Rohde P, Lewinsohn PM, Seeley JR. Are people changed by the experience of having an episode of depression? A further test of the scar hypothesis. *J Abnorm Psychol*. Aug 1990;99(3):264-71. doi:10.1037//0021-843x.99.3.264

54.   Treadway MT, Bossaller NA, Shelton RC, Zald DH. Effort-based decision-making in major depressive disorder: a translational model of motivational anhedonia. *J Abnorm Psychol*. Aug 2012;121(3):553-8. doi:10.1037/a0028813

55.   Husain M, Roiser JP. Neuroscience of apathy and anhedonia: a transdiagnostic approach. *Nat Rev Neurosci*. Aug 2018;19(8):470-484. doi:10.1038/s41583-018-0029-9

56.   Treadway MT, Zald DH. Reconsidering anhedonia in depression: lessons from translational neuroscience. *Neurosci Biobehav Rev*. Jan 2011;35(3):537-55. doi:10.1016/j.neubiorev.2010.06.006

57.   Moses-Payne ME, Rollwage M, Fleming SM, Roiser JP. Postdecision Evidence Integration and Depressive Symptoms. *Front Psychiatry*. 2019;10:639. doi:10.3389/fpsyt.2019.00639

58.   Rouault M, Seow T, Gillan CM, Fleming SM. Psychiatric Symptom Dimensions Are Associated With Dissociable Shifts in Metacognition but Not Task Performance. *Biol Psychiatry*. Sep 15 2018;84(6):443-451. doi:10.1016/j.biopsych.2017.12.017

59.   Seow TXF, Rouault M, Gillan CM, Fleming SM. How Local and Global Metacognition Shape Mental Health. *Biological Psychiatry*. 2021/10/01 2021;90(7):436-446. doi:https://doi.org/10.1016/j.biopsych.2021.05.013

60.   Stephan KE, Manjaly ZM, Mathys CD, et al. Allostatic Self-efficacy: A Metacognitive Theory of Dyshomeostasis-Induced Fatigue and Depression. *Front Hum Neurosci*. 2016;10:550. doi:10.3389/fnhum.2016.00550

61.   Huys QJM, Daw ND, Dayan P. Depression: A Decision-Theoretic Analysis. *Annual Review of Neuroscience*. 2015;38(Volume 38, 2015):1-23. doi:https://doi.org/10.1146/annurev-neuro-071714-033928

62.   Halahakoon DC, Lewis G, Roiser JP. Cognitive Impairment and Depression-Cause, Consequence, or Coincidence? *JAMA Psychiatry*. Mar 1 2019;76(3):239-240. doi:10.1001/jamapsychiatry.2018.3631

63.   Lewinsohn PM, Clarke GN, Seeley JR, Rohde P. Major Depression in Community Adolescents: Age at Onset, Episode Duration, and Time to Recurrence. *Journal of the American Academy of Child & Adolescent Psychiatry*. 1994/07/01 1994;33(6):809-818. doi:https://doi.org/10.1097/00004583-199407000-00006

64.   Solmi M, Radua J, Olivola M, et al. Age at onset of mental disorders worldwide: large-scale meta-analysis of 192 epidemiological studies. *Molecular Psychiatry*. 2022/01/01 2022;27(1):281-295. doi:10.1038/s41380-021-01161-7

65.   McMakin DL, Olino TM, Porta G, et al. Anhedonia predicts poorer recovery among youth with selective serotonin reuptake inhibitor treatment–resistant depression. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2012;51(4):404-411.

66.   Martell CR, Dimidjian S, Herman-Dunn R. *Behavioral Activation for Depression: A Clinician's Guide*. Guilford Press; 2013.

67.   Norbury A, Hauser T, Fleming S, Dolan R, Huys QJM. Different components of cognitive-behavioural therapy affect specific cognitive mechanisms. PsyArXiv; 2023.

Figure 1: Apple Gathering Task (AGT) and acceptance rates. (A) On each trial, participants are given a different offer comprising a number of apples (3, 6, 9, or 12 apples) for a given effort cost (20%, 40%, 60% or 80% of their maximum grip strength). Participants can either accept the offer or refuse the offer. If the offer is accepted, participants need to squeeze the gripper to the required effort level (or above) for 3 seconds in order to win the apples on this trial. For refused offers, 'no response required' was displayed, followed by the next decision. (B) Average acceptance rates as a function of reward level (number of points) and effort level (% MVC) for the Pilot study. (C) Distribution of the number of accepted offers (out of 80) in the Pilot. (D) Average acceptance rates as a function of reward level and effort level across all groups in the Case-control study. (E) Distribution of the number of accepted offers (out of 80) across all groups in the Case-control study. Continuous lines denote the kernel density estimate from the data, dashed lines denote the mean, and dotted lines represent ± 1 standard deviation from the mean.

Figure 2: Acceptance rates for the Case-control study. (A) Average acceptance rate as a function of reward level (points) and effort level (% MVC) for the control (CTR), first degree relatives (REL), patients with current depression (MDD), and remitted depression (REM) group. (B) Distribution of the number of accepted offers for each group. Continuous lines denote the kernel density estimate from the data, dashed lines denote the mean, and dotted lines represent ± 1 standard deviation from the mean. (C) Overall probability to accept offers for each group. Black dots represent the mean and error bars represent the standard error of the mean. Note that raw data is presented but analyses were conducted on arcsine transformed data.

Figure 3: Estimated model parameters for the Pilot (A-D) and Case-control (E-G) study. Figures are showing violin and boxplots as well as the mean (plus sign) and median (notch) for (A) estimated intercept/bias (K), (B) reward sensitivity (LinR), (C) linear effort (LinE), and (D) quadratic effort sensitivity (E^2) parameter values from the winning model in the pilot study. Data are shown for estimated (E) intercept/bias (K), (F) reward sensitivity (LinR), and (G) quadratic effort (E^2) sensitivity parameter values from the winning model in the Case-control study. CTR: Control group; REL: First-degree relative group, MDD: Current depression group; REM: Remitted depression group. *Denotes significance at p<0.05.

Figure 4: Correlations between computational parameters and symptom factors. (A) Correlation between the Low Mood factor and the linear reward sensitivity (LinR) parameter in the pilot study. (B) Correlation between the Low Mood factor and the quadratic effort sensitivity (E^2) parameter in the pilot study. (C) Correlation between the Hedonia factor and the quadratic effort sensitivity (E^2) parameter in the pilot study. (D) Correlation between the Low Mood factor and the linear reward sensitivity (LinR) parameter in the case-control study for the CTR+REL group only.

Table 1: Participant characteristics.

Sample	Pilot	Case-Control				
	HV (N=67)	CTR (N=57)	REL (N=36)	REM (N=46)	MDD (N=41)	Statistics (Case-control)
Age	28.45 (9.88)	26.70 (8.14)	26.06 (8.19)	26.91 (7.06)	30.24 (11.57)	$F_{(3,176)}=1.87, p>0.05$
Gender (M/F)	23/44	18/39	13/23	17/29	12/29	$X_{(3)}=0.78, p>0.05$
Digit Span forward	9.86 (1.90)	9.35 (1.86)	9.28 (1.97)	9.41 (1.85)	9.07 (2.08)	$F_{(3,176)}=0.26, p>0.05$
Digit Span backward	8.20 (2.36)	7.84 (2.23)	7.06 (2.27)	8.04 (2.28)	7.44 (2.31)	$F_{(3, 176)}=1.54, p>0.05$
IQ (WTAR)	109.79 (11.47)	111.91 (7.43)	113.15 (7.65)	116.14 (8.08)	114.51 (9.95)	$F_{(3,172)}=2.33, p>0.05$
Years Education	--	16.51 (2.82)	17.19 (2.97)	17.04 (2.82)	15.68 (2.31)	$F_{(3, 176)}=2.51, p>0.05$
HAM-D	--	0.58 (1.05)	1.06 (1.35)	1.35 (1.98)	17.00 (5.40)	--
BDI-II	3.39 (4.40)	2.09 (2.71)	2.78 (4.08)	5.00 (5.22)	28.78 (7.95)	--
STAI Trait	34.96 (9.18)	32.88 (7.58)	34.86 (11.63)	39.07 (9.50)	63.46 (8.10)	--
SHAPS	24.34 (5.98)	21.93 (5.39)	21.75 (4.75)	23.30 (4.59)	37.20 (5.52)	--
TEPS-A	46.99 (7.00)	44.83 (7.11)	46.03 (7.85)	43.63 (6.69)	30.05 (8.49)	--
TEPS-C	36.60 (6.79)	36.04 (5.95)	37.78 (6.16)	37.30 (6.85)	25.90 (7.46)	--
AES Cognitive Apathy	12.33 (2.94)	10.33 (2.53)	10.00 (2.16)	11.18 (3.11)	17.93 (3.75)	--
AES Behavioural Apathy	7.60 (1.72)	7.12 (1.51)	7.25 (1.95)	8.07 (2.38)	13.39 (2.38)	--
AES Emotional Apathy	3.52 (1.13)	3.53 (1.10)	3.50 (1.11)	3.84 (1.33)	5.88 (1.54)	--
AES Other Apathy	5.15 (1.59)	4.95 (1.46)	4.86 (1.55)	5.29 (1.62)	8.46 (1.90)	--
Chapman Physical Anhedonia	12.72 (8.06)	--	--	--	--	--
DAS Perfectionism	15.40 (4.27)	41.32 (13.25)	38.28 (13.56)	48.76 (12.83)	64.27 (17.46)	--
DAS Social Approval	9.24 (2.63)	31.82 (7.29)	33.03 (7.32)	35.63 (9.22)	41.44 (10.26)	--
LOTR Optimism	15.60 (4.70)	16.26 (3.93)	16.67 (4.50)	14.26 (4.20)	7.54 (4.48)	--

HV: Healthy Volunteers (in the Pilot study); CTR: Control Participants; REL: Participants with at a first-degree relative with Depression; REM: Remitted depressed participants; MDD; Currently depressed participants; WTAR: Wechsler Test of Adult Reading; HAM-D: Hamilton Depression Rating Scale; BDI-II: Beck Depression Inventory; STAI: State Trait Anxiety Inventory; SHAPS: Snaith Hamilton Pleasure Scale; TEPS-A: Temporal Experience of Pleasure Scale - Anticipatory subscale; TEPS-C: Temporal Experience of Pleasure Scale – Consummatory; AES: Apathy Evaluation Scale; DAS: Dysfunctional Attitudes Scale; LOTR: Life Orientation Test-Revised. The DAS short form scales were used in the Pilot. Brackets represent standard deviations. In the Pilot study, there was missing data for the digit span forward (N=2), digit span backward (N=3) and IQ (N=4). IQ data were missing for four Case-control participants (REM N=2, REL N=2).