Manuscript Title: The DNA Damage Checkpoint Targets the Kinetochore for Relocation of Collapsed Forks to the Periphery

Running Title: DRC targets Cep3 for fork relocation

Keywords: Nuclear Pore, Collapsed Fork, DNA Replication Checkpoint, Damage Induced Microtubule, DNA Repair, Nuclear Periphery, Rad53, Dun1, TNR, Non-B DNA, Huntington's Disease, CAG Repeat

Authors: Tyler Maclay[*,1], Jenna Whalen[*,1], Matthew Johnson[1] & Catherine H. Freudenreich[1]
[1]Department of Biology, Tufts University, Medford, MA 02155
[*]Authors contributed equally

Abstract

Hairpin forming expanded CAG/CTG repeats pose significant challenges to DNA replication which can lead to replication fork collapse. Long CAG/CTG repeat tracts relocate to the nuclear pore complex to maintain their integrity. Forks impeded by DNA structures are known to activate the DNA damage checkpoint, thus we asked whether checkpoint proteins play a role in relocation of collapsed forks to the nuclear periphery in *S. cerevisiae*. We show that relocation of a $(CAG/CTG)_{130}$ tract is dependent on activation of the Mrc1/Rad53 replication checkpoint. Further, checkpoint-mediated phosphorylation of the kinetochore protein Cep3 is required for relocation, implicating detachment of the centromere from the spindle pole body. Activation of this pathway leads to DNA damage-induced microtubule recruitment to the repeat. These data suggest a role for the DNA replication checkpoint in facilitating movement of collapsed replication forks to the nuclear periphery by centromere release and microtubule-directed motion.

Highlights

- The DNA damage checkpoint is needed for relocation of a structure-forming CAG repeat tract to the nuclear pore complex
- The importance of Mrc1 (hClaspin) implicates fork uncoupling as the initial checkpoint signal
- Phosphorylation of the Cep3 kinetochore protein by Dun1 kinase modulates centromere release, which is critical for collapsed fork repositioning
- Damage-inducible nuclear microtubules colocalize with the CAG repeat locus and are required for relocalization
- Establishes a new role for the DNA replication and DNA damage checkpoint response to trigger repositioning of collapsed forks within the nucleus.

Introduction

The nuclear periphery (NP) has emerged as an important site for the maintenance of genome stability when normal replication is impaired. Several types of replication stress have been shown to provoke relocation to the nuclear periphery in both yeast and human cells. These include drugs known to stall or collapse replication forks such as hydroxyurea combined with methyl methanesulfonate (MMS) in yeast (Nagai et al., 2008), expanded CAG repeats that form

secondary structures (Su et al., 2015), the polymerase inhibitor aphidicolin in human cells
(Lamm et al., 2020), protein mediated stalls by replication fork barriers (RFBs) (Horigome et al.,
2019; Kramarz et al., 2020), and replication stalls within telomeres (Aguilera et al., 2020;
Pinzaru et al., 2020). Some persistent types of DNA damage such as double-strand breaks
(DSBs) lacking a readily available template for repair or in sequestered heterochromatin domains
and eroded telomeres also relocate the nuclear periphery (for review, see (Whalen and
Freudenreich, 2020; Lamm et al., 2021; Gasser and Stutz, 2023)).
We previously elucidated a detailed pathway for how collapsed forks caused by an
expanded CAG/CTG tract of 130 repeat units (abbreviated $(CAG)_{130}$) relocates to the *S.*
cerevisiae nuclear pore complex (NPC) in late S phase in a manner dependent on replication (Su
et al., 2015; Whalen et al., 2020). CAG/CTG repeats over 35 repeat units have a propensity to
form hairpin structures that can impair replication leading to fork collapse (Usdin et al., 2015;
Khristich and Mirkin, 2020). Defects in relocation to the NPC result in increased chromosome
end loss and repeat instability, implicating the pathway in protecting against fork breakage and
inaccurate repair (Su et al., 2015). This pathway has many similarities with pathways that have
been described for persistent or heterochromatic DSBs and eroded telomeres, but also several
differences (reviewed in (Whalen and Freudenreich, 2020)). We determined that relocation of
collapsed forks caused by CAG repeats was dependent on resection, the binding and
SUMOylation of key fork associated repair proteins (RPA, Rad52, Rad59, Smc5) by the Smc5-6
associated SUMO ligase Mms21, and interaction with the SUMO interacting motifs (SIMs) of
Slx5 which targets the collapsed fork to the NPC (Whalen et al., 2020).
In addition to the SUMOylation requirements, we wanted to investigate other processes
that could be involved in relocation of collapsed forks caused by CAG repeats to the NPC. One
important pathway known to respond to stalled and collapsed forks is the DNA damage
response. The DNA damage response can be subdivided into two pathways: the DNA replication
checkpoint (DRC) and the DNA damage checkpoint (DDC), defined by their mediator proteins
Mrc1 or Rad9, respectively. Our previous findings showed that expanded CAG repeats activate
the DDC (Sundararajan and Freudenreich, 2011), and long range resection is required for
$(CAG)_{130}$ relocation (Whalen et al., 2020), which would produce ssDNA to stimulate DDC
activation. In addition, members of the DDC and the DRC were important for maintaining repeat
stability and preventing chromosome breaks at long CAG tracts (Lahiri et al., 2004;
Freudenreich and Lahiri, 2004; Gellon et al., 2019).
The checkpoint is composed of several sensor and mediator components that activate
downstream effectors, reviewed in (Hustedt et al., 2013; McClure et al., 2022) and summarized
in (Figure 1A). Sensor proteins identify processed DNA lesions, such as resected DNA
substrates, and then recruit other proteins to the damage site. In *S. cerevisiae*, the sensor Ddc2
(hATRIP) interacts with RPA bound to ssDNA to recruit the kinase Mec1 (ATR). Mec1 is also
activated by the PCNA-like 9-1-1 clamp (Ddc1, Rad17, Mec3 in budding yeast) which is loaded
by the Rad24-RFC complex onto ds/ssDNA junctions. Downstream effector proteins such as
Rad53 (hChk2) and Chk1 elicit the DNA damage response through further phosphorylation
signaling. At the replication fork there is an additional Rad53 activation mechanism involving
the mediator, Mrc1 (hClaspin), which travels with the replisome and is phosphorylated upon
replisome uncoupling (Katou et al., 2003; Lou et al., 2008). Another pathway more specific to
DSBs involves Tel1 (hATM), which senses DNA ends through its interaction with the yeast
MRX (hMRN) complex. Finally, the yeast mediator protein Rad9 (h53BP1) amplifies Rad53
phosphorylation. Rad9 is recruited to histone modifications including methylated H3 (H3-K79)

and damage-specific H2A-S129 phosphorylation, which can occur at either collapsed forks or
DSBs. Due to the different DNA damage sensors, identification of the checkpoint pathway that is
activated can give insight into the lesion or lesions that are created by a particular type of DNA
damage.
The relocation of persistent DSBs in *S. cerevisiae* and heterochromatic DSBs in
Drosophila required both Mec1/Tel1 and ATR/ATM, respectively (Nagai et al., 2008; Khadaroo
et al., 2009; Chiolo et al., 2011). In human U2OS and IMR90 cells that were treated with
aphidicolin to inhibit polymerases, ATR inhibition prevented the localization of stalled
replication forks to the nuclear periphery (Lamm et al., 2020). In contrast, single deletion of
either Mec1 or Tel1 did not prevent the relocation of an expanded $(CAG)_{130}$ repeat to the NPC
(Su et al., 2015), leaving it an open question whether the checkpoint is involved in the relocation
of DNA damage caused by DNA structures to the nuclear periphery, and if so what pathway is
involved. In yeast, Mec1/ATR is not only required for DSB relocation, but has been linked to
increasing chromatin mobility in response to DSBs (Dion et al., 2012). Yet an expanded
$(CAG)_{130}$ repeat not only failed to increase mobility of a chromosome harboring this fork-stalling
element, it decreased mobility due to the added constraint of NPC interaction (Su et al., 2015).
Therefore, it was not clear whether the checkpoint pathway would be involved in relocation of
DNA structure-mediated collapsed forks to the NPC, and if so, in what manner.
In this study, we show that the activation of the effector kinase Rad53 by phosphorylation
of the replisome-associated Mrc1 protein is needed for the relocation of a $(CAG)_{130}$ tract to the
nuclear periphery. We identified that the Rad53 target Dun1 is also required for relocation.
Through investigating several Dun1 substrates, we determined that the phosphorylation of the
kinetochore protein Cep3 at serine 575 is the important target to allow for repositioning of the
DNA structure-induced collapsed fork in the nucleus. We investigated several possible
mechanisms by which Cep3 phosphorylation could act and found an important factor was the
disruption of centromere attachment to the spindle pole body (the yeast microtubule organizing
center which is functionally equivalent to the mammalian centrosome). In contrast, Cep3
phosphorylation-dependent increases in chromosome movement were not detectable. Finally, we
show that damage-induced microtubules (DIMs) (Oshidari et al., 2018) are formed in response to
damage at the CAG tract, colocalize with the CAG tract locus, and are dependent on this
checkpoint pathway. We speculate that the phosphorylation of Cep3-S575 allows for relocation
of collapsed forks to the NPC by relieving centromere constraint and supporting DIM-directed
motion towards the nuclear periphery.
126
127
128 **Results**
129
Relocation of a $(CAG)_{130}$ tract to the nuclear pore complex depends on the activation of
Rad53 and Mrc1.
We used a previously described zoning assay (Su et al., 2015) to visualize the CAG
repeat in relation to the nuclear periphery. Briefly, 130 CAG/CTG repeats are integrated on
chromosome VI 6.4 kb away from a LacO array which can be visualized by binding of GFP-
LacI; in some experiments the repeat was placed 6.4 kb from a TetO/mCherry-TetR array
(Figure 1B). GFP-Nup49 was used to mark NPCs to allow for visualization of the nuclear
periphery. The nucleus was divided into three equal areas to determine the frequency of CAG
repeat occupancy in each zone (see Methods). Zone 1 represents the nuclear periphery (where

the NPCs are located), zone 3 is the middle of the nucleus, and zone 2 is the area in between
(Figure 1B). The presence of the $(CAG)_{130}$ tract caused an increase in the occupancy of the
nearby tagged locus in zone 1, from 31% in the $(CAG)_0$ control to 48% for the chromosome
containing $(CAG)_{130}$ (Fig. 1C). The distribution between zones 1, 2 and 3 was consistent
between both wild-type $(CAG)_{130}$ strains, whether the LacO/GFP-LacI or TetO/mCherry-TetR
array was used to follow the CAG repeat locus (Figure S1).
145   Previously, it was shown that single deletions in Mec1 or Tel1 sensor kinases (ATR and
ATM in mammals, respectively) do not affect relocation of the $(CAG)_{130}$ tract to the nuclear
periphery during S phase (Su et al., 2015). Simultaneous disruption of Mec1 and Tel1 could not
be tested by the zoning assay due to severely abnormal nuclear morphology. As there are many
factors involved in the activation of the DNA damage response that can act redundantly (Figure
1A), we investigated if there was any role for the checkpoint in CAG tract relocation by testing
the effector kinases. The two effector kinases of the yeast DNA damage checkpoint are Chk1 and
Rad53 (reviewed in (Hustedt et al., 2013)). Deletion of Chk1 showed no significant decrease in
the frequency the CAG tract was found at the nuclear periphery (Figure 1C). However, deletion
of Rad53 (viable with the added deletion of Sml1 (Zhao et al., 1998)) did result in a significant
decrease in $(CAG)_{130}$ occupancy in zone 1. This result supports that the DNA damage response is
required for relocation. This was further confirmed by testing $rad53^{K227A}$, a kinase-defective
mutant of Rad53 that is checkpoint-deficient (Fay et al., 1997). $rad53^{K227A}$ showed the same
significant decrease in relocation as $rad53\Delta sml1\Delta$ (Figure 1C). We conclude that activation of
the DNA damage response and the kinase activity of Rad53 is needed for relocation of the
collapsed fork to the NPC.
161   There are multiple sensor and mediator proteins that can activate Rad53 in response to
different types of DNA damage. We had previously shown that long CAG tracts were especially
prone to breakage and exhibited increased fork stalling in the absence of the Mrc1 protein
(Gellon et al., 2019), which couples polymerase epsilon to the CMG helicase. Mrc1 contains
multiple SQ/TQ motifs that are targeted by kinases of the PIKK family, such as Mec1 (hATR),
Tel1 (hATM) or Rad53 (hChk2) for phosphorylation upon fork uncoupling (Naylor et al., 2009;
Osborn and Elledge, 2003). The Mrc1^{AQ} mutant, with its 17 SQ/TQ sites mutated to alanine
(Osborn and Elledge, 2003), had a modest effect on chromosome fragility but exhibited
increased repeat instability (Gellon et al., 2019), suggesting a possible role for Mrc1 in sensing
forks stalled at the $(CAG)_{130}$ tract. The deletion of Mrc1 had severely abnormal nuclear
morphology precluding zoning analysis. However, complementation with the checkpoint-
deficient Mrc1^{AQ} mutant abrogated these abnormal morphologies and showed a significant
reduction in relocation to the nuclear periphery to a similar level as a Rad53 mutant (Figure 1C).
Thus, the checkpoint signal relevant for repositioning to the NPC is channeled through Mrc1,
which suggests that the signal originates from an uncoupled replication fork.
176
177   **Mec1 and Rad9 act redundantly to sense damage at the $(CAG)_{130}$ tract and signal**
178   **repositioning within the nucleus.**
179   Though single mutations in either of the main checkpoint sensor kinases, Mec1 and Tel1,
180   did not result in a decrease in relocation to the nuclear periphery (see Figure 1D) the presence of
181   the Mre11 protein, which senses and processes DSB ends, is required (Whalen et al., 2020).
182   Deletion of the Rad9 mediator protein (h53BP1 ortholog), which acts downstream of Tel1 to
183   facilitate Rad53 autophosphorylation (Emili, 1998; Gilbert et al., 2001), showed no effect on the
184   relocation of the repeat to the nuclear periphery (Figure 1D). Similarly, deletion of the 9-1-1

clamp component Ddc1 (9-1-1 activates Mec1) or the clamp loader Rad24 also had no effect on the percent of the CAG locus in zone 1 (Figure 1D). These results were somewhat surprising given the important role of Mrc1, which is phosphorylated by Mec1, and our previous findings that both Mec1 and Rad9 were important for preventing breaks and instability at expanded CAG tracts (Lahiri et al., 2004). Since there is a well-described redundancy between these sensing pathways (reviewed in (Hustedt et al., 2013)), we explored whether there was redundancy in the sensing of lesions to signal relocation of the $(CAG)_{130}$ to the nuclear periphery. Deletion of both Mec1 and Rad9 resulted in a decrease in relocation of the $(CAG)_{130}$ to the nuclear periphery (Figure 1D). In addition, co-disruption of Rad9 and Rad24 also led to a decrease in relocation to the NP, and deletion of all 3 genes ($mec1\Delta rad24\Delta rad9\Delta$) led to the lowest % zone 1 occupancy (Figure 1D). These effects were not made worse by additional disruption of Tel1 (Figure 1D), suggesting that the role of Tel1 is not independent of Rad9 in this situation. We previously showed that both H2A-S129 and H2A-T126 phosphorylation could be detected at a $(CAG)_{155}$ tract 40 minutes into S phase (House et al., 2014, 2019), which is shortly before relocation occurs (Whalen et al., 2020) and could serve to recruit Rad9. Importantly, DNA damage dependent phosphorylation of H2A requires either Mec1 or Tel1 (Downs et al., 2000; Shroff et al., 2004). So, in the absence of Mec1, the recruitment of Rad9 would be expected to be downstream of Tel1 activity. Thus, our data indicate that there are multiple ways to activate Rad53 to enable relocation of the $(CAG)_{130}$ tract: a Tel1/Rad9 mediated pathway, or a Rad24/Mec1 pathway, acting redundantly with the Rad9 pathway. In addition, Mrc1 phosphorylation, which facilitates recruitment of Rad53 (Alcasabas et al., 2001), is an important mediator of Rad53 activation at the CAG-stalled fork.

Altogether, this analysis suggests that there may be multiple types of damage at the CAG tract that can provoke the checkpoint response and are important for nuclear repositioning. Based on our data, repeat mediated checkpoint activation and relocation to the NPC could be due to an uncoupled fork (sensed by Mrc1), a region of single-stranded DNA (sensed by the Ddc2-Mec1 and 9-1-1 complexes), or a DSB end (sensed by the MRX-Tel1-Rad9 pathway). Single-strand DNA could occur at an uncoupled fork, at a reversed fork, or at a post-replicative gap, and a DSB end could result either from fork reversal or fork breakage/cleavage. All of these signals would be present at an uncoupled fork that transitions to either a reversed or broken fork, which we refer to collectively as "collapsed forks" (Figure 1E). Alternatively, there could be a mixture of possible species that occur in different cells.

Phosphorylation of Cep3 Serine 575 by Dun1 is needed for relocation of collapsed forks to the nuclear pore complex.

After determining that activation of the transducing kinase Rad53 by the DNA replication checkpoint is required for $(CAG)_{130}$ relocation, we investigated what targets of Rad53 are needed. Dun1 is a well characterized target of Rad53 (Figure 2A) (Bashkirov et al., 2003; Nordlund and Reichard, 2006; de la Torre Ruiz and Lowndes, 2000; Zhou and Elledge, 1993). A $dun1\Delta$ showed a significant decrease in CAG tract relocation to the N.P. similar to $rad53\Delta sml1\Delta$ and $rad53^{K227A}$ mutants (Figure 2B). Therefore, Rad53 activation of Dun1 is a crucial step in the pathway leading to repositioning of collapsed forks to the NPC.

Next, we investigated what targets of Dun1 are required for relocation (Figure 2A). Dun1 is required for the phosphorylation of the Rad51 associated repair protein Rad55 (Bashkirov et al., 2003). Phosphorylation of Rad55 allows for it to dimerize with Rad57 and effectively stabilize Rad51 binding to promote more successful strand invasion (Fortin and Symington,

2002). Therefore, Dun1 plays a direct role in recombination mediated repair. However, neither
rad55Δ nor *rad55Δrad57Δ* mutants showed a decrease in the percentage of CAG tract loci detect
in zone 1 (Figure 2B). Dun1 has also been well studied as an activator of ribonucleotide
reductase (RNR) activity and functions to regulate nucleotide metabolism in response to
replicative stress. Dun1 inhibits Rfx1, Sml1, and Dif1 through phosphorylation which in turn
prevents them from downregulating nucleotide pools (Huang et al., 1998; Wu and Huang, 2008;
Zhao and Rothstein, 2002). Since Dun1 inhibits Sml1 activity, in order to determine if Dun1's
role in relocation is through maintaining nucleotide pools, Sml1 was deleted in the *dun1Δ* strain.
If the deletion of Sml1 rescues the defect in CAG tract repositioning in the *dun1Δ* strain, it would
indicate that this activity of Dun1 is required. However, *dun1Δsml1Δ* showed a similar percent S
phase zone 1 occupancy as *dun1Δ* strains (Figure 2B). We conclude that the role for Dun1 in
relocation to the NP is neither to promote recombination mediated repair nor to increase
nucleotide pools.
Another phosphorylation target of Dun1 is the kinetochore protein Cep3 (Strecker et al.,
2016). Cep3 is a DNA-binding protein that recognizes the centromere sequence as a member of
the CBF3 kinetochore complex (Lechner and Carbon, 1991; Lechner, 1994). Phosphorylation of
Cep3 at serine 575 (S575) occurs in response to an induced DSB or zeocin-induced DNA
damage in a manner dependent on both Rad53 and Dun1 (Strecker et al., 2016). The yeast
kinetochore interacts with the spindle pole body (SPB), the yeast microtubule organizing center
which is functionally equivalent to the mammalian centrosome. Cep3-S575 phosphorylation has
been reported to allow for an increase in the dynamics of the SPB-CEN attachment after DSB
induction on a different chromosome (Strecker et al., 2016). Mutation of serine 575 to a non-
phosphorylatable alanine residue resulted in a significant decrease in the CAG locus occupancy
in zone 1, showing that Cep3 phosphorylation is required for relocation of the CAG tract to the
NPC (Figure 2B).
To confirm that Cep3 phosphorylation at S575 is required for relocation we constructed a
phosphomimetic mutant of Cep3 by mutating serine 575 to glutamic acid. The *cep3^{S575E}*
mutation bypassed the need for Dun1 as the *dun1Δcep3^{S575E}* mutant showed a normal level of
CAG locus S phase occupancy in zone 1 (Figure 2C). We conclude that phosphorylation of
Cep3-S575 mediated by Dun1 is needed for relocation of collapsed forks to the NPC.
To further address the role of Cep3 phosphorylation we assessed the timing of CAG locus
relocation to the NP in *cep3^{S575E}* cells over the course of an S-phase. As previously reported, in
wild-type cells the $(CAG)_{130}$ tract is rarely at the NPC in early S-phase and is maximal in late S
phase at 60 min post release from G1 (Figure 2D) (Whalen et al., 2020). However Rad52 is
found at the repeat locus 40 minutes into S phase (Su et al., 2015), suggesting that fork collapse
has occurred by then and that there is a delay between the occurrence of damage and anchoring
at the pore. The *cep3^{S575E}* mutation led to a significant increase in zone 1 occupancy of the
$(CAG)_{130}$ repeat tract at the 40- and 50-minute time points. (Figure 2D). This suggests that the
cep3^{S575E} phosphomimetic is bypassing the normal signaling pathway, priming the fork for
repositioning as soon as enough RPA-coated ssDNA has occurred to recruit Rad52.
271
Phosphorylation of Cep3-S575 allows for relocation of collapsed forks to the NPC by
modulating kinetochore-SPB attachment, but not by increasing global chromosome
movement or activating the spindle assembly checkpoint.
It has been shown that forcing transcription through the centromere with a galactose-
inducible promoter causes detachment of the centromere from the SPB (Collins et al., 2005; Hill

and Bloom, 1987). Cep3-S575 phosphorylation has been suggested to relieve attachment of
centromeres to the SPB (Strecker et al., 2016), though the $cep3^{S575A}$ mutation does not increase
SPB-CEN distance or position with respect to the nuclear periphery (Strecker et al., 2016;
Cheblal et al., 2020). We reasoned that if centromere attachment to the SPB was preventing
relocation to the NPC in the $cep3^{S575A}$ mutant, then forcing transcription through the centromere
and thus interrupting that attachment could allow for relocation of the repeat tract in the $cep3^{S575A}$
mutant. We integrated the *GAL1* promoter upstream of the centromere on chromosome 6
(*pGAL1-CEN6*) (Pobiega and Marcand, 2010), which is the chromosome containing the
$(CAG)_{130}$ tract. Upon galactose induction and transcription through the centromere, the reduced
zone 1 occupancy of the CAG locus in the $cep3^{S575A}$ strain was rescued to normal wild-type
levels (Figure 3A). We conclude that phosphorylation of Cep3-S575 may allow for relocation of
collapsed forks to the NPC by relieving centromere attachment to the SPB.
Detachment of the centromere from the SPB has a number of consequences. It has been
shown that inactivating a centromere can increase the mobility of that chromosome (Strecker et
al., 2016). It is also known that induced DSBs increase both local (Dion et al., 2012) and global
(Miné-Hattab and Rothstein, 2012) chromosome mobility (reviewed in (García Fernández and
Fabre, 2022)). This DSB-induced mobility was shown to be dependent on Cep3 phosphorylation
in one study (Strecker et al., 2016), but not another (Cheblal et al., 2020). We have previously
published that the $(CAG)_{130}$ tract, in contrast to a DSB, does not increase either local or global
chromosome mobility, and instead causes decreased locus mobility relative to chromosomes
lacking the repeat tract (Su et al., 2015). However, those data were collected from
unsynchronized cells in S phase, and the locus likely becomes constrained only after anchoring
to the NPC. We reasoned that an increase in chromosome mobility due to a collapsed fork at the
$(CAG)_{130}$ tract and subsequent Cep3-S575 phosphorylation could be transient. The anchoring at
the NPC occurs in a distinct time window in late S phase, between about 55-75 minutes after
release from G1 and peaking at ~60 minutes (Figure 2D and (Whalen et al., 2020)), therefore any
chromosome VI movement could be confined to a short time period preceding NPC anchoring.
The radius of constraint, a measurement of locus mobility (Dion and Gasser, 2013), was
measured for the $(CAG)_{130}$ locus in cells that were synchronously released from G1 phase over a
100 minute time course segmented into 5 minute intervals. During each 5 minute interval,
images were acquired every 1.5 seconds, closely mirroring the conditions used in (Strecker et al.,
2016), and MSD curves for each 5 minute window were plotted to obtain a radius of constraint.
An example of the MSD curve for the first 100 seconds of the 60-minute timepoint is depicted
(Figure 3B). For each timepoint, the radius of constraint was calculated based on the plateau
reached within $\Delta t = 100s$ (Figure 3C; Table S3). No difference in mobility of chromosome VI
adjacent to the $(CAG)_{130}$ tract was observed between wild-type and $cep3^{S575A}$ cells (Figure 3C).
Though a fast directed motion may not have been detected by this method, this result argues
against a local increase in chromosome movement triggered by Cep3 phosphorylation in
response to a collapsed fork.
Another driver of DSB mobility is local nucleosome depletion (Cheblal et al., 2020).
Loss of Nhp6, encoded in two loci Nhp6a and Nhp6b, has been shown to reduce histone
occupancy on DNA (Celona et al., 2011), and this leads to increased chromosome mobility
(Hauer et al., 2017; Cheblal et al., 2020).To test the mobility hypothesis in a different way, we
reasoned that if increasing chromosome mobility by phosphorylating Cep3^{S575} was a requirement
for $(CAG)_{130}$ relocation, then causing global nucleosome depletion by deletion of Nhp6a and
Nhp6b could rescue the $cep3^{S575A}$ relocation defect. Deletion of Nhp6a and Nhp6b (*nhp6ΔΔ*) did

not cause a decrease in relocation, and when combined with $cep3^{S575A}$ did not rescue the
relocation defect of those cells (Figure 3D). Altogether, we conclude that an increase in
chromosome mobility does not explain the importance of Cep3 phosphorylation for relocation of
the $(CAG)_{130}$ tract to the NP.
Centromeres unattached to the spindle can activate the spindle assembly checkpoint
(SAC) via Mad2 (Hoyt et al., 1991), and phosphorylation of Cep3 has been shown to activate the
SAC (Strecker et al., 2016). Therefore, we tested whether activation of the SAC by Mad2 is
required for relocation, however there was no relocation defect in $mad2\Delta$ cells (Figure 3D). We
conclude that the role of Cep3 phosphorylation in collapsed fork repositioning to the NP is not
due to a defect in SAC activation.
333
Expanded CAG repeat loci associate with DIMs in a Dun1 and Cep3-S575
phosphorylation-dependent manner.
In yeast, microtubule structures termed "Damage-Inducible Microtubules" or "DIMs"
form in response to double stranded breaks in a Rad9 checkpoint-dependent manner (Oshidari et
al., 2018). Interestingly, inactivating a centromere is sufficient to drive formation of DNA
damage-inducible microtubules (DIMs) (Oshidari et al., 2018). Since our data supported that
centromere inactivation and Cep3 phosphorylation were important to allow relocation of the
CAG locus to the NP, but not through increasing chromosome mobility, we wondered whether
DIM formation might be involved. We reasoned that if DIMs were important for relocation to
the periphery, then the formation of microtubules in general would be important. To test this,
cells were treated for 30 minutes with either 15 μg/mL of the tubulin depolymerizing drug
nocodazole or an equivalent amount of the vehicle (DMSO) and the zoning assay was
performed. Addition of nocodazole significantly reduced the percentage of the $(CAG)_{130}$ tract
localized to the nuclear periphery in mid-to-late S phase cells (Figure 4A). It is known that the
minus end-directed microtubule motor Kar3 promotes, though is not strictly required, for DIM
formation (Oshidari et al., 2018). Therefore, we tested whether deletion of Kar3 had an effect on
the frequency of $(CAG)_{130}$ locus association with the NP. In $kar3\Delta$ cells in S phase, relocation of
the repeat to zone 1 was reduced (though not eliminated; Figure 4B), in line with the reported
$kar3\Delta$ effect on DIMs.
To determine if DIMs were being recruited to the CAG repeat locus, we integrated an
additional copy of the tubulin subunit Tub1 tagged with the YFP variant mVenus at the *TUB1*
genomic locus (Markus et al., 2015). GFP-LacI was replaced with CFP-LacI and GFP-Nup49
with Nup49-mCherry to allow simultaneous visualization of all 3 nuclear components (Figure
4C). Consistent with previous data (Oshidari et al., 2018), treatment with 0.03% MMS caused
DIM formation in 62% of S-phase cells (Figure 4D). To test whether DIMs were also induced by
fork collapse, they were measured in cells treated with 0.03%MMS + 0.2M Hydroxyurea for 60
minutes, which causes genome-wide fork collapse and relocation of marked chromosome loci to
the nuclear periphery (Nagai et al., 2008; Whalen et al., 2020). Under fork collapsing conditions,
87% of mid-to-late S phase cells contained a DIM (Figure 4D). We then tested whether a single
expanded $(CAG)_{130}$ repeat locus was sufficient to cause DIM formation in cells. We observed
that there is an increase in prevalence of DIM formation in cells with an expanded CAG repeat to
36% compared to 18% for the no-tract control (Figure 4E). The majority of these DIMs (74%)
colocalized with the repeat locus (Figure 4E; see Figures 4C and S2 examples). DIM formation
was dependent upon Dun1 and Cep3 phosphorylation as deletion of Dun1 or mutation to
$cep3^{S575A}$ caused a significant reduction of DIMs (Figure 4E). The repeat tract-induced DIMs

were specifically affected in these mutants and were reduced to 7% in the *cep3*^{S575A} strain (Figure
4E). We conclude that there is a checkpoint-dependent induction of DIMs and recruitment of a
DIM to the site of fork collapse. These DIMs appear to be required to mediate relocation of the
collapsed fork to the NPC as deletion of *KAR3* or treatment with nocodazole, both of which
perturb DIM formation, led to a decrease in the frequency of relocation of the CAG tract to the
nuclear periphery.
375
376   **Discussion**
377   In this study we aimed to determine if the DNA damage checkpoint response plays a role
in the relocation of collapsed forks to the nuclear pore complex (NPC). We previously showed
that long CAG/CTG repeat tracts activate the DNA damage response and that the fork protection
complex containing Mrc1 is crucial for preventing CAG fragility and instability and promoting
fork progression through the repeat (Lahiri et al., 2004; Sundararajan and Freudenreich, 2011;
Gellon et al., 2019). Other systems showed varying checkpoint requirements for relocation of
chromosomes with persistent DSBs, aphidicolin stalled forks or eroded telomeres (Nagai et al.,
2008; Khadaroo et al., 2009; Chiolo et al., 2011; Lamm et al., 2020). It was unknown what, if
any, role the checkpoint played in the relocation of stalled/collapsed forks caused by expanded
CAG repeats to the nuclear periphery. Even for types of damage where a requirement for
checkpoint proteins had been established, the mechanism by which the checkpoint pathway
mediates repositioning of damaged chromosomal loci had not been elucidated.
389   Our results indicate that the Mrc1/Rad53 axis is required for the repositioning of CAG
repeats to the NPC in mid-to-late S phase. The (CAG)$_{130}$ repeat forms hairpin structures that
interfere with DNA replication (Pelletier et al., 2003; Viterbo et al., 2016) and CAG repeats have
similar relocation requirements as cells which have been treated with both HU and MMS which
causes fork collapse (Nagai et al., 2008; Su et al., 2015; Whalen et al., 2020), thus it was
speculated that the relocation event was a consequence of a collapsed fork (Su et al., 2015;
Whalen et al., 2020). Since the phosphorylation of Mrc1 is required for the efficient sensing and
stabilizing of stalled or uncoupled forks and the checkpoint activity of Mrc1 is critical for the
relocation of the expanded repeat to the nuclear periphery, this implies that a stalled or
uncoupled fork is the initial lesion being sensed and then processed for relocation to the NPC.
399   We found that there is redundancy in the sensing of the lesion caused by the (CAG)$_{130}$
tract, suggesting that by-products of fork uncoupling such as ssDNA or an exposed DSB end
caused by fork reversal or breakage also come into play. It was revealing that both the Mec1 and
Rad9 pathways needed to be inactivated in order to abolish relocation to the NP. Long-range
resection and RPA SUMOylation are required for CAG repeat relocation to the NPC (Whalen et
al., 2020), thus we would expect RPA-coated ssDNA, which recruits Mec1, to be available prior
to relocation. Rad9, which is activated by both Mec1 and Tel1, was required only when Mec1
and/or the 9-1-1 complex are compromised, indirectly suggesting a role for Tel1. Tel1 has been
shown to bind to the MRX complex (Cassani et al., 2016), which binds DSB ends and is required
for relocation (Whalen et al., 2020). These results imply that either ssDNA or a DSB end can
activate the pathway needed for repositioning, suggesting that there could be a transition from
the initial uncoupled fork to a reversed or broken fork that provides a DSB end (Figure 1E). This
redundancy in signaling leads to a robust mechanism to activate the Rad53 checkpoint in
response to the initial fork stall caused by an expanded CAG repeat.
413   Rad53 is the master regulator of the checkpoint in *S. cerevisiae* and has multiple roles in
the response to stalled or collapsed replication forks (Santocanale and Diffley, 1998; Zhao et al.,

1998; Herzberg et al., 2006; McClure and Diffley, 2021). One well-studied target of Rad53 is the
kinase Dun1 (Bashkirov et al., 2003), which we determined was needed for relocation (Figure
2B). It is important to note that while Dun1 is a required target of Rad53 for relocation, it may
not be the only target as the kinase-deficient $rad53^{K227A}$ mutant is proficient for Dun1 activation
(Hoch et al., 2013), yet does not relocate. Dun1 phosphorylates several downstream targets
including the recombination factor Rad55 (Bashkirov et al., 2003), RNR inhibition factors (Crt1,
Sml1, and Dif1 reviewed in (Nordlund and Reichard, 2006)), and the kinetochore protein Cep3
(Strecker et al., 2016). Of these targets, the phosphorylation of Cep3 was required for relocation
(Figure 2C).
The yeast centromere is only ~120bp in length and has three sub-sections: CDEI, CDEII,
and CDEIII. Kinetochore proteins at each centromere bind to a single microtubule, which is
attached to the spindle pole body (SPB) throughout most of the cell cycle (Heun et al., 2001).
Cep3 has a zinc finger DNA binding domain that interacts with the CDEIII region as a member
of the CBF3 complex (reviewed in (Biggins, 2013)). Phosphorylation of the kinetochore has
been widely studied and has varying functions (reviewed in (Funabiki and Wynne, 2013; Klemm
et al., 2021). For example, phosphorylation of the kinetochore proteins, such as Dam1, by Aurora
B kinase (Ipl1) destabilizes erroneous microtubule attachments (Tanaka et al., 2002; Pinsky et
al., 2006). However, the consequences of Cep3 phosphorylation have been comparatively poorly
described. It was previously suggested that Cep3 phosphorylation led to a release of centromere
constraint to allow the increased global mobility of chromosomes observed in response to an
HO-induced DSB at the *MAT* locus in the G2/M phase of the cell cycle (Strecker et al., 2016).
However, another group was unable to show a Cep3-dependent increased radius of constraint
(Cheblal et al., 2020) using similar conditions. Our results indicate that Cep3 phosphorylation
may act through another mechanism besides increasing local or global chromatin mobility, at
least in response to replication fork collapse. We propose that the DNA damage dependent
phosphorylation of Cep3 is causing centromere release from the SPB attachment normally
present during S phase. Consistent with this model, the requirement for Cep3 phosphorylation
can be obviated by centromere inactivation (Figure 3A).
DIMs form in response to DNA damage induced by MMS, zeocin, camptothecin, or I-
SceI-induced DSBs (Oshidari et al., 2018). With this work, we've now shown that while DIM
formation is common after treatment with MMS, which stalls forks, it is dramatically increased
after treatment with HU + MMS, which has been shown to be fork collapsing. DIM formation
even occurs as a result of a single fork collapse caused by an expanded CAG repeat tract.
Interestingly, much like how CEN inactivation eliminates the need for Cep3 phosphorylation for
relocation, CEN inactivation can cause spontaneous formation of DIMs (Oshidari et al., 2018).
This suggests that while the initial damage (whether a DSB or uncoupled/reversed/broken fork)
is responsible for the centromeric release from the SPB through Cep3 phosphorylation, the DIM
may form as a result of that release rather than directly responding to the damage. Thus, we
hypothesize that release of the centromere from the SPB triggers the formation of a DIM.
The microtubule motor protein Kar3 has been shown to promote the capture and
directional movement of damaged loci by DIMs (Oshidari et al., 2018). When centromeres are
inactivated, kinetochores that were previously attached to the microtubule plus end reposition
and bind the microtubule side-on (Tanaka et al., 2007, 2005). Kar3, as a minus end directed
motor, is involved in repositioning the centromere on the microtubule, by hindering tethering of
the kinetochore to the plus end of the microtubule (Tanaka et al., 2005; Tytell and Sorger, 2006;
Tanaka et al., 2007). During this kinetochore release and repositioning process, the damaged

repeat locus could be freed up to move to the NPC, for example, by allowing the centromere to
slide along the DIM. If this is the case, then we might expect that the presence of Kar3 could
increase the duration of this "relocation permissive" state by antagonizing reattachment of the
kinetochore to the microtubule plus end, which could explain why a *kar3Δ* reduces but does not
completely abrogate relocation to the NPC. What remains unclear is whether the microtubule
also interacts directly with the collapsed fork, or if the DIM allows for repositioning of the lesion
in an indirect manner. The former is an attractive hypothesis due to the increased colocalization
of the repeat locus with the DIM (Figure 4E). It has recently been shown that DIMs serve as sites
of repair center nucleation, with liquid droplets of Rad52 colocalizing with DIMs (Oshidari et
al., 2020), and we've previously shown that many repair proteins, including Rad52 (Su et al.,
2015) and RPA (Whalen et al., 2020) are recruited to the site of the collapsed fork prior to
relocation. These factors have been shown to become SUMOylated, and their SUMOylation has
been shown to be critically required for relocation to the NPC through an interaction with Slx5/8
(Whalen et al., 2020). Slx5/8 in turn is critical for anchoring of damaged DNA at the NPC
(reviewed in (Whalen and Freudenreich, 2020)). SUMOylation has also been shown to promote
condensation of repair factors (Keiten-Schmitz et al., 2020). Possibly, SUMOylation or factors
within SUMOylation-induced repair centers could play a role in the association between the
collapsed fork at the CAG tract and the DIM.
Thus, we propose that after fork collapse at the CAG repeat, the DNA replication
checkpoint results in the phosphorylation of Cep3. This phosphorylation causes centromere
release from the SPB, which promotes DIM formation. The DIM allows movement of the
collapsed fork locus to the periphery in a manner yet to be fully elucidated, but which may
involve interaction of the damaged locus with the DIM. Once at the periphery, interactions
between SUMOylated repair factors, Slx5/8, and the nucleoporins tether the locus to the NPC for
repair or fork restart (Figure 4F).
There is prior evidence in other types of eukaryotic cells for a physical mechanism by
which movement of lesions to repair centers within the nucleus occurs. Heterochromatic DSBs
in *Drosophila* require nuclear F-actin and myosin for directed movement to the nuclear periphery
(Caridi et al., 2018). In human U2OS cells the relocation of collapsed forks caused by
aphidicolin is dependent on the polymerization of nuclear F-actin, and late-replication foci were
observed to move along actin filaments to the nuclear periphery (Lamm et al., 2020). Further it's
been shown that recruitment of nuclear actin in response to damage can control pathway choice
in response to replication stress (Palumbieri et al., 2023). It is unclear if yeast have equivalent
nuclear actin filaments or if directed movement to the NP would be detectable in the smaller
yeast nucleus. However, in yeast, the kinesin-14 motor protein complex is required for
movement of telomeric DSBs from their usual attachment site on the nuclear membrane to NPCs
and these were observed to move along DIMs (Oshidari et al., 2018; Chung et al., 2015). Our
results suggest that collapsed forks may also utilize DIMs to facilitate their movement to the
yeast NPC. In human cells, it was recently reported that cytoplasmic microtubules that form in
response to DSBs push on the nuclear envelope to form nuclear envelope tubules (NETs) that
elongate into the nucleus and bring the nuclear envelope into the proximity of DSBs within the
nuclear interior, and the DSBs may also move toward the dsbNETs (Shokrollahi et al., 2024).
The formation of these dsbNETs serves to mobilize repair proteins to sites of DSBs and is
dependent on the ATM/ATR/DNA-PK checkpoint response (Shokrollahi et al., 2024). Thus, the
involvement of microtubules in mobilizing a repair response seems to be conserved, though the
details of where in the nucleus the damage-inducible microtubules form and how they interact

with the damaged locus may differ between organisms. It will be interesting to elucidate if DIMs are utilized in multicellular eukaryotic cells or if their function has been replaced by a combination of nuclear actin filaments and dsbNETs in these larger nuclei.

In summary, our results illuminate a new role for the DNA replication and DNA damage checkpoint response, which is to phosphorylate a kinetochore component to allow for repositioning of a collapsed fork within the nucleus. Since this repositioning is important for repair or fork restart, we identify it as a crucial new role for the DNA damage checkpoint response in *S. cerevisiae*. Repositioning of damaged loci to a repair center is present across many organisms. Where it has been studied, the checkpoint is critical to facilitate this relocation, but the targets involved have not been elucidated. Given the importance of recovery from replication fork collapse, a mechanism to allow centromere release to facilitate damaged locus repositioning might be especially critical at regions with fork-stalling DNA structures and for cancerous or aging cells experiencing high levels of replication stress.

Acknowledgements
We would like to thank Stéphane Marcand for the gift of the *pGAL1*-CEN6 plasmid and Wei-Lih Lee for the *p*HIS3 *p*:Venus-Tub1+3'UTR::LEU2 plasmid.

Funding
This work was funded by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number R35 GM144215 to CHF.

Author Contributions
JW: Conceptualization, Investigation, Analysis, Supervision, Writing
TM: Conceptualization, Investigation, Analysis, Software, Writing
MJ: Investigation, Analysis
CHF: Supervision, Resources, Writing, Administration, Conception

References

Aguilera, P., J. Whalen, C. Minguet, D. Churikov, C. Freudenreich, M.-N. Simon, and V. Géli. 2020. The nuclear pore complex prevents sister chromatid recombination during replicative senescence. *Nat. Commun.* 11:160. doi:10.1038/s41467-019-13979-5.

Alcasabas, A.A., A.J. Osborn, J. Bachant, F. Hu, P.J.H. Werler, K. Bousset, K. Furuya, J.F.X. Diffley, A.M. Carr, and S.J. Elledge. 2001. Mrc1 transduces signals of DNA replication stress to activate Rad53. *Nat. Cell Biol.* 3:958–965. doi:10.1038/ncb1101-958.

Bashkirov, V.I., E.V. Bashkirova, E. Haghnazari, and W.-D. Heyer. 2003. Direct Kinase-to-Kinase Signaling Mediated by the FHA Phosphoprotein Recognition Domain of the Dun1 DNA Damage Checkpoint Kinase. *Mol. Cell. Biol.* 23:1441–1452. doi:10.1128/MCB.23.4.1441-1452.2003.

Biggins, S. 2013. The Composition, Functions, and Regulation of the Budding Yeast Kinetochore. *Genetics*. 194:817–846. doi:10.1534/genetics.112.145276.

Caridi, C.P., C. D'Agostino, T. Ryu, G. Zapotoczny, L. Delabaere, X. Li, V.Y. Khodaverdian, N.
Amaral, E. Lin, A.R. Rau, and I. Chiolo. 2018. Nuclear F-actin and myosins drive
relocalization of heterochromatic breaks. *Nature*. 559:54–60. doi:10.1038/s41586-018-
0242-8.

Cassani, C., E. Gobbini, W. Wang, H. Niu, M. Clerici, P. Sung, and M.P. Longhese. 2016. Tel1
and Rif2 Regulate MRX Functions in End-Tethering and Repair of DNA Double-Strand
Breaks. *PLOS Biol.* 14:e1002387. doi:10.1371/journal.pbio.1002387.

Celona, B., A. Weiner, F.D. Felice, F.M. Mancuso, E. Cesarini, R.L. Rossi, L. Gregory, D.
Baban, G. Rossetti, P. Grianti, M. Pagani, T. Bonaldi, J. Ragoussis, N. Friedman, G.
Camilloni, M.E. Bianchi, and A. Agresti. 2011. Substantial Histone Reduction Modulates
Genomewide Nucleosomal Occupancy and Global Transcriptional Output. *PLOS Biol.*
9:e1001086. doi:10.1371/journal.pbio.1001086.

Cheblal, A., K. Challa, A. Seeber, K. Shimada, H. Yoshida, H.C. Ferreira, A. Amitai, and S.M.
Gasser. 2020. DNA Damage-Induced Nucleosome Depletion Enhances Homology
Search Independently of Local Break Movement. *Mol. Cell*. 80:311-326.e4.
doi:10.1016/j.molcel.2020.09.002.

Chiolo, I., A. Minoda, S.U. Colmenares, A. Polyzos, S.V. Costes, and G.H. Karpen. 2011.
Double-Strand Breaks in Heterochromatin Move Outside of a Dynamic HP1a Domain to
Complete Recombinational Repair. *Cell*. 144:732–744. doi:10.1016/j.cell.2011.02.012.

Chung, D.K.C., J.N.Y. Chan, J. Strecker, W. Zhang, S. Ebrahimi-Ardebili, T. Lu, K.J. Abraham,
D. Durocher, and K. Mekhail. 2015. Perinuclear tethers license telomeric DSBs for a
broad kinesin- and NPC-dependent DNA repair process. *Nat. Commun.* 6:7742.
doi:10.1038/ncomms8742.

Collins, K.A., A.R. Castillo, S.Y. Tatsutani, and S. Biggins. 2005. De Novo Kinetochore
Assembly Requires the Centromeric Histone H3 Variant. *Mol. Biol. Cell*. 16:5649–5660.
doi:10.1091/mbc.E05-08-0771.

Dion, V., and S.M. Gasser. 2013. Chromatin Movement in the Maintenance of Genome Stability.
Cell. 152:1355–1364. doi:10.1016/j.cell.2013.02.010.

Dion, V., V. Kalck, C. Horigome, B.D. Towbin, and S.M. Gasser. 2012. Increased mobility of
double-strand breaks requires Mec1, Rad9 and the homologous recombination
machinery. *Nat. Cell Biol.* 14:502–509. doi:10.1038/ncb2465.

Downs, J.A., N.F. Lowndes, and S.P. Jackson. 2000. A role for Saccharomyces cerevisiae
histone H2A in DNA repair. *Nature*. 408:1001–1004. doi:10.1038/35050000.

Emili, A. 1998. MEC1-Dependent Phosphorylation of Rad9p in Response to DNA Damage.
Mol. Cell. 2:183–189. doi:10.1016/S1097-2765(00)80128-8.

Fay, D.S., Z. Sun, and D.F. Stern. 1997. Mutations in SPK1/RAD53 that specifically abolish
checkpoint but not growth-related functions. *Curr. Genet.* 31:97–105.
doi:10.1007/s002940050181.

Fortin, G.S., and L.S. Symington. 2002. Mutations in yeast Rad51 that partially bypass the
requirement for Rad55 and Rad57 in DNA repair by increasing the stability of Rad51–
DNA complexes. *EMBO J.* 21:3160–3170. doi:10.1093/emboj/cdf293.

Freudenreich, C.H., and M. Lahiri. 2004. Structure-Forming CAG/CTG Repeat Sequences are
Sensitive to Breakage in the Absence of Mrc1 Checkpoint Function and S-Phase
Checkpoint Signaling: Implications for Trinucleotide Repeat Expansion Diseases. *Cell*
Cycle. 3:1370–1374. doi:10.4161/cc.3.11.1246.

Funabiki, H., and D.J. Wynne. 2013. Making an effective switch at the kinetochore by
phosphorylation and dephosphorylation. *Chromosoma*. 122:135–158.
doi:10.1007/s00412-013-0401-5.

García Fernández, F., and E. Fabre. 2022. The Dynamic Behavior of Chromatin in Response to
DNA Double-Strand Breaks. *Genes*. 13:215. doi:10.3390/genes13020215.

Gasser, S.M., and F. Stutz. 2023. SUMO in the regulation of DNA repair and transcription at
nuclear pores. *FEBS Lett.* 597:2833–2850. doi:10.1002/1873-3468.14751.

Gellon, L., S. Kaushal, J. Cebrián, M. Lahiri, S.M. Mirkin, and C.H. Freudenreich. 2019. Mrc1
and Tof1 prevent fragility and instability at long CAG repeats by their fork stabilizing
function. *Nucleic Acids Res.* 47:794–805. doi:10.1093/nar/gky1195.

Generoso, W.C., M. Gottardi, M. Oreb, and E. Boles. 2016. Simplified CRISPR-Cas genome
editing for Saccharomyces cerevisiae. *J. Microbiol. Methods*. 127:203–205.
doi:10.1016/j.mimet.2016.06.020.

Gilbert, C.S., C.M. Green, and N.F. Lowndes. 2001. Budding Yeast Rad9 Is an ATP-Dependent
Rad53 Activating Machine. *Mol. Cell*. 8:129–136. doi:10.1016/S1097-2765(01)00267-2.

Hauer, M.H., A. Seeber, V. Singh, R. Thierry, R. Sack, A. Amitai, M. Kryzhanovska, J.
Eglinger, D. Holcman, T. Owen-Hughes, and S.M. Gasser. 2017. Histone degradation in
response to DNA damage enhances chromatin dynamics and recombination rates. *Nat.*
Struct. Mol. Biol. 24:99–107. doi:10.1038/nsmb.3347.

Hediger, F., F.R. Neumann, G. Van Houwe, K. Dubrana, and S.M. Gasser. 2002. Live Imaging
of Telomeres: yKu and Sir Proteins Define Redundant Telomere-Anchoring Pathways in
Yeast. *Curr. Biol.* 12:2076–2089. doi:10.1016/S0960-9822(02)01338-6.

Herzberg, K., V.I. Bashkirov, M. Rolfsmeier, E. Haghnazari, W.H. McDonald, S. Anderson,
E.V. Bashkirova, J.R. Yates, and W.-D. Heyer. 2006. Phosphorylation of Rad55 on
Serines 2, 8, and 14 Is Required for Efficient Homologous Recombination in the
Recovery of Stalled Replication Forks. *Mol. Cell. Biol.* 26:8396–8409.
doi:10.1128/MCB.01317-06.

Heun, P., T. Laroche, M.K. Raghuraman, and S.M. Gasser. 2001. The Positioning and Dynamics
of Origins of Replication in the Budding Yeast Nucleus. *J. Cell Biol.* 152:385–400.

Hill, A., and K. Bloom. 1987. Genetic manipulation of centromere function. *Mol. Cell. Biol.*
7:2397–2405.

Hoch, N.C., E.S.-W. Chen, R. Buckland, S.-C. Wang, A. Fazio, A. Hammet, A. Pellicioli, A.
Chabes, M.-D. Tsai, and J. Heierhorst. 2013. Molecular basis of the essential s phase
function of the rad53 checkpoint kinase. *Mol. Cell. Biol.* 33:3202–3213.
doi:10.1128/MCB.00474-13.

Horigome, C., E. Unozawa, T. Ooki, and T. Kobayashi. 2019. Ribosomal RNA gene repeats
associate with the nuclear pore complex for maintenance after DNA damage. *PLOS*
Genet. 15:e1008103. doi:10.1371/journal.pgen.1008103.

House, N.C., E.J. Polleys, I. Quasem, M. De la Rosa Mejia, C.E. Joyce, O. Takacsi-Nagy, J.E.
Krebs, S.M. Fuchs, and C.H. Freudenreich. 2019. Distinct roles for S. cerevisiae H2A
copies in recombination and repeat stability, with a role for H2A.1 threonine 126. *eLife*.
8:e53362. doi:10.7554/eLife.53362.

House, N.C.M., J.H. Yang, S.C. Walsh, J.M. Moy, and C.H. Freudenreich. 2014. NuA4 Initiates
Dynamic Histone H4 Acetylation to Promote High-Fidelity Sister Chromatid
Recombination at Postreplication Gaps. *Mol. Cell*. 55:818–828.
doi:10.1016/j.molcel.2014.07.007.

Hoyt, M.A., L. Totis, and B.T. Roberts. 1991. S. cerevisiae genes required for cell cycle arrest in
response to loss of microtubule function. *Cell*. 66:507–517. doi:10.1016/0092-
8674(81)90014-3.

Huang, M., Z. Zhou, and S.J. Elledge. 1998. The DNA Replication and Damage Checkpoint
Pathways Induce Transcription by Inhibition of the Crt1 Repressor. *Cell*. 94:595–605.
doi:10.1016/S0092-8674(00)81601-3.

Hustedt, N., S.M. Gasser, and K. Shimada. 2013. Replication Checkpoint: Tuning and
Coordination of Replication Forks in S Phase. *Genes*. 4:388–434.
doi:10.3390/genes4030388.

Katou, Y., Y. Kanoh, M. Bando, H. Noguchi, H. Tanaka, T. Ashikari, K. Sugimoto, and K.
Shirahige. 2003. S-phase checkpoint proteins Tof1 and Mrc1 form a stable replication-
pausing complex. *Nature*. 424:1078–1083. doi:10.1038/nature01900.

Keiten-Schmitz, J., K. Schunck, and S. Müller. 2020. SUMO Chains Rule on Chromatin
Occupancy. *Front. Cell Dev. Biol.* 7. doi:10.3389/fcell.2019.00343.

Khadaroo, B., M.T. Teixeira, P. Luciano, N. Eckert-Boulet, S.M. Germann, M.N. Simon, I.
Gallina, P. Abdallah, E. Gilson, V. Géli, and M. Lisby. 2009. The DNA damage response
at eroded telomeres and tethering to the nuclear pore complex. *Nat. Cell Biol.* 11:980–
987. doi:10.1038/ncb1910.

Khristich, A.N., and S.M. Mirkin. 2020. On the wrong DNA track: Molecular mechanisms of
repeat-mediated genome instability. *J. Biol. Chem.* 295:4134–4170.
doi:10.1074/jbc.REV119.007678.

Klemm, C., P.H. Thorpe, and G. Ólafsson. 2021. Cell-cycle phospho-regulation of the
kinetochore. *Curr. Genet.* 67:177–193. doi:10.1007/s00294-020-01127-2.

Kramarz, K., K. Schirmeisen, V. Boucherit, A. Ait Saada, C. Lovo, B. Palancade, C.
Freudenreich, and S.A.E. Lambert. 2020. The nuclear pore primes recombination-
dependent DNA synthesis at arrested forks by promoting SUMO removal. *Nat. Commun.*
11:5643. doi:10.1038/s41467-020-19516-z.

Lahiri, M., T.L. Gustafson, E.R. Majors, and C.H. Freudenreich. 2004. Expanded CAG Repeats
Activate the DNA Damage Checkpoint Pathway. *Mol. Cell.* 15:287–293.
doi:10.1016/j.molcel.2004.06.034.

Lamm, N., M.N. Read, M. Nobis, D. Van Ly, S.G. Page, V.P. Masamsetti, P. Timpson, M. Biro,
and A.J. Cesare. 2020. Nuclear F-actin counteracts nuclear deformation and promotes
fork repair during replication stress. *Nat. Cell Biol.* 22:1460–1470. doi:10.1038/s41556-
020-00605-6.

Lamm, N., S. Rogers, and A.J. Cesare. 2021. Chromatin mobility and relocation in DNA repair.
Trends Cell Biol. 31:843–855. doi:10.1016/j.tcb.2021.06.002.

Lechner, J. 1994. A zinc finger protein, essential for chromosome segregation, constitutes a
putative DNA binding subunit of the Saccharomyces cerevisiae kinetochore complex,
Cbf3. *EMBO J.* 13:5203–5211. doi:10.1002/j.1460-2075.1994.tb06851.x.

Lechner, J., and J. Carbon. 1991. A 240 kd multisubunit protein complex, CBF3, is a major
component of the budding yeast centromere. *Cell.* 64:717–725. doi:10.1016/0092-
8674(91)90501-o.

Lou, H., M. Komata, Y. Katou, Z. Guan, C.C. Reis, M. Budd, K. Shirahige, and J.L. Campbell.
2008. Mrc1 and DNA Polymerase ☐ Function Together in Linking DNA Replication and
the S Phase Checkpoint. *Mol. Cell.* 32:106–117. doi:10.1016/j.molcel.2008.08.020.

Markus, S.M., S. Omer, K. Baranowski, and W.-L. Lee. 2015. Improved Plasmids for
Fluorescent Protein Tagging of Microtubules in Saccharomyces cerevisiae. *Traffic Cph.*
Den. 16:773–786. doi:10.1111/tra.12276.

McClure, A.W., B. Canal, and J.F.X. Diffley. 2022. A DNA replication fork-centric view of the
budding yeast DNA damage response. *DNA Repair.* 119:103393.
doi:10.1016/j.dnarep.2022.103393.

McClure, A.W., and J.F. Diffley. 2021. Rad53 checkpoint kinase regulation of DNA replication
fork rate via Mrc1 phosphorylation. *eLife.* 10:e69726. doi:10.7554/eLife.69726.

Meister, P., L.R. Gehlen, E. Varela, V. Kalck, and S.M. Gasser. 2010. Chapter 21 - Visualizing
Yeast Chromosomes and Nuclear Architecture. *In* Methods in Enzymology. Academic
Press. 535–567.

Miné-Hattab, J., and R. Rothstein. 2012. Increased chromosome mobility facilitates homology
search during recombination. *Nat. Cell Biol.* 14:510–517. doi:10.1038/ncb2472.

Nagai, S., K. Dubrana, M. Tsai-Pflugfelder, M.B. Davidson, T.M. Roberts, G.W. Brown, E.
Varela, F. Hediger, S.M. Gasser, and N.J. Krogan. 2008. Functional Targeting of DNA
Damage to a Nuclear Pore–Associated SUMO-Dependent Ubiquitin Ligase. *Science*.
322:597–602. doi:10.1126/science.1162790.

Naylor, M.L., J. Li, A.J. Osborn, and S.J. Elledge. 2009. Mrc1 phosphorylation in response to
DNA replication stress is required for Mec1 accumulation at the stalled fork. *Proc. Natl.*
Acad. Sci. U. S. A. 106:12765–12770. doi:10.1073/pnas.0904623106.

Nordlund, P., and P. Reichard. 2006. Ribonucleotide Reductases. *Annu. Rev. Biochem.* 75:681–
706. doi:10.1146/annurev.biochem.75.103004.142443.

Osborn, A.J., and S.J. Elledge. 2003. Mrc1 is a replication fork component whose
phosphorylation in response to DNA replication stress activates Rad53. *Genes Dev.*
17:1755–1767. doi:10.1101/gad.1098303.

Oshidari, R., R. Huang, M. Medghalchi, E.Y.W. Tse, N. Ashgriz, H.O. Lee, H. Wyatt, and K.
Mekhail. 2020. DNA repair by Rad52 liquid droplets. *Nat. Commun.* 11:695.
doi:10.1038/s41467-020-14546-z.

Oshidari, R., J. Strecker, D.K.C. Chung, K.J. Abraham, J.N.Y. Chan, C.J. Damaren, and K.
Mekhail. 2018. Nuclear microtubule filaments mediate non-linear directional motion of
chromatin and promote DNA repair. *Nat. Commun.* 9:2567. doi:10.1038/s41467-018-
05009-7.

Palumbieri, M.D., C. Merigliano, D. González-Acosta, D. Kuster, J. Krietsch, H. Stoy, T. von
Känel, S. Ulferts, B. Welter, J. Frey, C. Doerdelmann, A. Sanchi, R. Grosse, I. Chiolo,
and M. Lopes. 2023. Nuclear actin polymerization rapidly mediates replication fork
remodeling upon stress by limiting PrimPol activity. *Nat. Commun.* 14:1–15.
doi:10.1038/s41467-023-43183-5.

Pelletier, R., M.M. Krasilnikova, G.M. Samadashwily, R. Lahue, and S.M. Mirkin. 2003.
Replication and Expansion of Trinucleotide Repeats in Yeast. *Mol. Cell. Biol.* 23:1349–
1357. doi:10.1128/MCB.23.4.1349-1357.2003.

Pike, B.L., A. Hammet, and J. Heierhorst. 2001. Role of the N-terminal Forkhead-associated
Domain in the Cell Cycle Checkpoint Function of the Rad53 Kinase*210. *J. Biol. Chem.*
276:14019–14026. doi:10.1074/jbc.M009558200.

728 Pinsky, B.A., C. Kung, K.M. Shokat, and S. Biggins. 2006. The Ipl1-Aurora protein kinase
729    activates the spindle checkpoint by creating unattached kinetochores. *Nat. Cell Biol.*
730    8:78–83. doi:10.1038/ncb1341.

731 Pinzaru, A.M., M. Kareh, N. Lamm, E. Lazzerini-Denchi, A.J. Cesare, and A. Sfeir. 2020.
732    Replication stress conferred by POT1 dysfunction promotes telomere relocalization to the
733    nuclear pore. *Genes Dev.* 34:1619–1636. doi:10.1101/gad.337287.120.

734 Pobiega, S., and S. Marcand. 2010. Dicentric breakage at telomere fusions. *Genes Dev.* 24:720–
735    733. doi:10.1101/gad.571510.

736 Santocanale, C., and J.F.X. Diffley. 1998. A Mec1- and Rad53-dependent checkpoint controls
737    late-firing origins of DNA replication. *Nature.* 395:615–618. doi:10.1038/27001.

738 Shokrollahi, M., M. Stanic, A. Hundal, J.N.Y. Chan, D. Urman, C.A. Jordan, A. Hakem, R.
739    Espin, J. Hao, R. Krishnan, P.G. Maass, B.C. Dickson, M.P. Hande, M.A. Pujana, R.
740    Hakem, and K. Mekhail. 2024. DNA double-strand break–capturing nuclear envelope
741    tubules drive DNA repair. *Nat. Struct. Mol. Biol.* 1–12. doi:10.1038/s41594-024-01286-
742    7.

743 Shroff, R., A. Arbel-Eden, D. Pilch, G. Ira, W.M. Bonner, J.H. Petrini, J.E. Haber, and M.
744    Lichten. 2004. Distribution and Dynamics of Chromatin Modification Induced by a
745    Defined DNA Double-Strand Break. *Curr. Biol. CB.* 14:1703–1711.
746    doi:10.1016/j.cub.2004.09.047.

747 Strecker, J., G.D. Gupta, W. Zhang, M. Bashkurov, M.-C. Landry, L. Pelletier, and D. Durocher.
748    2016. DNA damage signalling targets the kinetochore to promote chromatin mobility.
749    *Nat. Cell Biol.* 18:281–290. doi:10.1038/ncb3308.

750 Su, X.A., V. Dion, S.M. Gasser, and C.H. Freudenreich. 2015. Regulation of recombination at
751    yeast nuclear pores controls repair and triplet repeat stability. *Genes Dev.* 29:1006–1017.
752    doi:10.1101/gad.256404.114.

753 Sundararajan, R., and C.H. Freudenreich. 2011. Expanded CAG/CTG Repeat DNA Induces a
754    Checkpoint Response That Impacts Cell Proliferation in Saccharomyces cerevisiae. *PLoS*
755    *Genet.* 7:e1001339. doi:10.1371/journal.pgen.1001339.

756 Tanaka, K., E. Kitamura, Y. Kitamura, and T.U. Tanaka. 2007. Molecular mechanisms of
757    microtubule-dependent kinetochore transport toward spindle poles. *J. Cell Biol.* 178:269–
758    281. doi:10.1083/jcb.200702141.

759 Tanaka, T.U., N. Rachidi, C. Janke, G. Pereira, M. Galova, E. Schiebel, M.J.R. Stark, and K.
760    Nasmyth. 2002. Evidence that the Ipl1-Sli15 (Aurora kinase-INCENP) complex
761    promotes chromosome bi-orientation by altering kinetochore-spindle pole connections.
762    *Cell.* 108:317–329. doi:10.1016/s0092-8674(02)00633-5.

763 Tanaka, T.U., M.J.R. Stark, and K. Tanaka. 2005. Kinetochore capture and bi-orientation on the
764    mitotic spindle. *Nat. Rev. Mol. Cell Biol.* 6:929–942. doi:10.1038/nrm1764.

765 de la Torre Ruiz, M.-A., and N.F. Lowndes. 2000. DUN1 defines one branch downstream of
766   RAD53 for transcription and DNA damage repair in Saccharomyces cerevisiae. *FEBS*
767   *Lett.* 485:205–206. doi:10.1016/S0014-5793(00)02198-0.

768 Tytell, J.D., and P.K. Sorger. 2006. Analysis of kinesin motor function at budding yeast
769   kinetochores. *J. Cell Biol.* 172:861–874. doi:10.1083/jcb.200509101.

770 Usdin, K., N.C.M. House, and C.H. Freudenreich. 2015. Repeat instability during DNA repair:
771   Insights from model systems. *Crit. Rev. Biochem. Mol. Biol.* 50:142–167.
772   doi:10.3109/10409238.2014.999192.

773 Viterbo, D., G. Michoud, V. Mosbach, B. Dujon, and G.-F. Richard. 2016. Replication stalling
774   and heteroduplex formation within CAG/CTG trinucleotide repeats by mismatch repair.
775   *DNA Repair.* 42:94–106. doi:10.1016/j.dnarep.2016.03.002.

776 Whalen, J.M., N. Dhingra, L. Wei, X. Zhao, and C.H. Freudenreich. 2020. Relocation of
777   Collapsed Forks to the Nuclear Pore Complex Depends on Sumoylation of DNA Repair
778   Proteins and Permits Rad51 Association. *Cell Rep.* 31:107635–107635.
779   doi:10.1016/j.celrep.2020.107635.

780 Whalen, J.M., and C.H. Freudenreich. 2020. Location, Location, Location: The Role of Nuclear
781   Positioning in the Repair of Collapsed Forks and Protection of Genome Stability. *Genes.*
782   11:635. doi:10.3390/genes11060635.

783 Wu, X., and M. Huang. 2008. Dif1 Controls Subcellular Localization of Ribonucleotide
784   Reductase by Mediating Nuclear Import of the R2 Subunit. *Mol. Cell. Biol.* 28:7156–
785   7167. doi:10.1128/MCB.01388-08.

786 Zhao, X., E.G.D. Muller, and R. Rothstein. 1998. A Suppressor of Two Essential Checkpoint
787   Genes Identifies a Novel Protein that Negatively Affects dNTP Pools. *Mol. Cell.* 2:329–
788   340. doi:10.1016/S1097-2765(00)80277-4.

789 Zhao, X., and R. Rothstein. 2002. The Dun1 checkpoint kinase phosphorylates and regulates the
790   ribonucleotide reductase inhibitor Sml1. *Proc. Natl. Acad. Sci.* 99:3746–3751.
791   doi:10.1073/pnas.062502299.

792 Zhou, Z., and S.J. Elledge. 1993. DUN1 encodes a protein kinase that controls the DNA damage
793   response in yeast. *Cell.* 75:1119–1127. doi:10.1016/0092-8674(93)90321-G.

<u>Figure Legends</u>

Figure 1: Relocation of (CAG)$_{130}$ locus to the nuclear periphery depends on the activation
of the DNA replication or DNA damage checkpoint. A) Flow chart of the DNA damage
checkpoint. B) Yeast chromosome VI containing integrated 130 CAG/CTG repeat tract
((CAG)$_{130}$) and a *lacO* or *tetO* array that binds the LacI-GFP or TetR-RFP protein, respectively.

Mid/late S phase cells were imaged, and the location of the foci were scored into one of three
zones of equal area, using the GFP-Nup49 signal to mark the nuclear periphery. Representative
images are shown. C) Percent of cells with LacO loci in zone 1 for mid/late S-phase cells. Cells
have either $(CAG)_0$ or $(CAG)_{130}$ as marked. D) Percent of zone 1 foci for mid/late S-phase cells
for indicated strains. All strains have $(CAG)_{130}$. E) Model illustrating activation of the DRC in
response to an uncoupled fork (left) or activation of the DDC following fork reversal and
recruitment of the MRX complex (right). (*) P≤0.05, (**) P≤0.01 (***) P≤0.001 compared with
wild-type $(CAG)_{130}$ strain by Fisher's exact test. More than 100 cells were analyzed per strain.
See Table S1 for the exact number of cells analyzed, percentages, and P-values.

Figure 2: Phosphorylation of Cep3^{S575} by Dun1 is needed for relocation of $(CAG)_{130}$ to the
nuclear periphery. A) Diagram of Dun1 targets. The colors used for mutants in this figure
correspond with colors used in this diagram. B) Percent of zone 1 foci for $(CAG)_{130}$ mid/late S-
phase cells for strains deleted for Dun1 or its targets. C) Percent of zone 1 foci for $(CAG)_{130}$
mid/late S-phase cells for a phosphomimetic mutant of Cep3 (*cep3^{S575E}*) individually and in the
dun1Δ mutant. D) Percentage of zone 1 foci for $(CAG)_{130}$ cells at indicated time points after
release from alpha factor arrest in wild-type and *cep3^{S575E}* strains. See Table S2 for the exact
number of cells analyzed, percentages, and P-values. (*) P≤0.05, (**) P≤0.01 (***) P≤0.001
compared with wild-type by Fisher's exact test. (^^^) P ≤ 0.001 compared as indicated by
Fisher's Exact Test. More than 100 cells were analyzed per strain. See Table S1 for the exact
number of cells analyzed, percentages, and P-values.

Figure 3: Cep3 phosphorylation allows for $(CAG)_{130}$ relocation through release from
centromeric constraint. A) Percent zone 1 foci for $(CAG)_{130}$ mid/late S-phase cells for
cep3^{S575A} and *P_{GAL1}-CEN6 cep3^{S575A}* strains. Cultures were grown in YP-lactate media then split
into media containing either glucose or galactose to induce transcription through CEN6. B) MSD
track for Cep3^{S575A} and WT strains containing $(CAG)_{130}$. Cells were imaged every 1.5 seconds
for 5 minutes, first Δt =[0,100] seconds shown. Data was captured 60 minutes post-release from
alpha factor arrest. The lines represent the 95% Confidence intervals of the mean square
displacement at a particular timepoint. C) Summary of radii of constraint calculated for WT and
cep3^{S575A} over 100 minutes post release from alpha factor arrest. Points represent calculated
Radius of constraint for a 5-minute interval imaged every 1.5s as in B, error bars are standard
error of the calculated radius of constraint., specific details in Table S3. D) Percent of zone 1 foci
for $(CAG)_{130}$ mid/late S-phase cells >100 cells per strain analyzed, see Table S1 for exact
values(***) P≤0.001 compared with wild-type by Fisher's exact test. (^^^) P= 0.0004 by
Fisher's exact test.

Figure 4: Damage-Inducible Microtubules associate with expanded CAG repeats in a Dun1
and Cep3 phosphorylation-dependent manner. A) Percentage of zone 1 $(CAG)_{130}$ cells in
mid/late S-phase. Cells were treated with either 15µg/mL nocodazole or an equivalent volume of
DMSO (0.1%). B) Percent zone 1 $(CAG)_{130}$ cells for cells lacking Kar3. C) Example images of
CFP-LacI colocalizing with DIMs. Bottom left image is a cell without a DIM for comparison.
The yellow channel was contrast-adjusted for clarity, raw images can be found in Figure S2A. D)
Fraction of mid-late S phase cells (no CAG tract strain) treated as indicated with at least 1 DIM.
E) Fraction of DIM containing cells of the indicated genotype. DIMs, if present, were scored as

colocalized or not colocalized with the CFP-LacI focus. (*) P≤0.05, (***) P≤0.001, (****) P ≤
0.0001 compared with wild-type by Fisher's exact test (zoning assay) or chi-square test (DIM
assay). See Table S1 (for zoning) or Table S4 (for DIM) for the exact number of cells analyzed,
percentages, and P-values. F) Proposed model for the relocation of a collapsed fork at the CAG
repeat to the nuclear pore complex. Collapsed or uncoupled forks at a CAG repeat result in a
Mrc1 phosphorylation followed by Rad53 activation. Rad53 activated Dun1 leads to
phosphorylation of Cep3. This phosphorylation alters the SPB-CEN attachment, leading to the
formation of DIMs. This may allow repositioning of the centromere on the DIM, for example in
a side-on attachment to permit relocation of the repeat to the NPC. Created with BioRender.com
855
856
857 **Figure S1: Comparison between wild-type strains.** Percent of zone 1 foci for $(CAG)_{130}$
858 mid/late S-phase cells for wild-type cells with the LacO/GFP-LacI array or TetO/mCherry-TetR
859 array ~6.4kb away from $(CAG)_{130}$ near ARS0607 on Chr VI. The number of cells analyzed per
860 strain were 157 and 162 respectively.
861
862 **Figure S2: DIM Examples:** A) Example images of Damage-inducible microtubules. Defined as
863 monopolar microtubule extrusions during S-phase which exhibit rapidly decaying fluorescence
864 intensity. Cell lacking a DIM shown for context. B) Example movie of a DIM associating with
865 an expanded CAG repeat locus. Video acquired at 0.2 frame/s.
866
867
<u>Method Details</u>
Yeast strains and genetic manipulation
Standard procedures were used in cell growth and medium preparation. All *S. cerevisiae* strains
were made in the W303 (RAD5+) background and are listed in Table S5. Point mutant alleles
have been previously characterized: $rad53^{K227A}$(Pike et al., 2001), $cep3^{S575A}$(Strecker et al.,
2016). The point mutants were generated using CRISPR-Cas9 methods as described
previously(Generoso et al., 2016). See Table S6 for primers used to create each mutation.
Correct mutation was screened by both restriction enzyme digest and sequencing. The P_{GAL1}-
CEN6 construct has been previously characterized (Pobiega and Marcand, 2010). Centromere
inactivation was confirmed by plating cells onto galactose containing media and observing a
decrease in viability compared to glucose. Venus-Tub1 was introduced as an additional copy
using plasmid pHIS3p:Venus-Tub1+3'UTR::LEU2, which was a gift from Wei-Lih Lee (Markus
et al., 2015).
881
Microscopy
Colonies were checked for 130-CAG repeats by PCR with primers flanking the repeat
(CTGRev2/T720 or 2StepCAG_pAG32-F/R). PCR products separated on a 2% agarose gel
(MetaPhor, Lonza). Cells from colonies with the correct repeat length were inoculated into YC
media, grown overnight at 30C with shaking, diluted back to OD=0.2 and grown to
approximately 5×10^6 cells per mL in YC media. Cells were fixed with 4% paraformaldehyde
and washed 3 times with 1X PBS. 1.4% YC agar was used to make agar pads on depression
slides and 5 µL of cells were added on top with a coverslip. Z-stack images were taken using a
Zeiss AX10 fluorescent microscope and/or a Delta Vision Ultra High-Resolution using an
Olympus 100X with 1.45NA. Step interval size was 0.15-0.2um and approximately 25 Z-planes

were taken per field of cells. Exposure time was approximately DIC: 100ms; GFP: 300ms and
mCherry: 500ms (for strains with mCherry-TetR). Images were deconvolved, and three-zoning
criteria was used to evaluate the location of the GFP foci for mid/late S-phase cells with the
ImageJ point picker program as described in (Meister et al., 2010). Only the middle two-thirds of
the stacks were used for analysis. Mid/Late S-phase cells were determined by yeast morphology
using bud size criteria of 2/3rd or less, and at least 15% the size of the mother cell (Hediger et al.,
2002). For drug treatments, cells were grown to approximately 1 x 10^7, split into 2 cultures and
grown for 30 minutes in either 0.1% dimethyl sulfoxide (DMSO) or 15ug/mL nocodazole. The
cells were then fixed and imaged as described above. For DIM experiments, cells were prepared
as described above and imaged with the filters (and exposure): Cyan (200ms), Yellow (200ms),
Red (200ms). Cells were scored for colocalization determined by any overlap (not inclusive of
"touching") of the repeat and the nuclear periphery, and of the repeat with DIMs. DIMs were
defined as monopolar microtubule structures emanating from the SPB during mid-late S phase
(See Figure 4C for examples).
906
P_{GAL1}-**CEN6 Inactivation**
Colonies were checked for 130-CAG repeats by PCR with primers flanking the repeat
(CTGRev2/T720). Cells from colonies with the correct repeat length were grown overnight in
synthetic complete media +2% glucose. Cultures were spun down and resuspended in YEP-
lactate media to an OD$_{600}$ of 0.3 and grown overnight. Cultures were spun down, split in half,
and resuspended in synthetic complete media +2% galactose or +2% glucose to an OD$_{600}$ of 0.3.
Cultures were grown for 3 hours followed by the zoning assay protocol as outlined above.
914
MSD Analysis
Colonies were checked for 130-CAG repeats by PCR with primers flanking the repeat
(2StepCAG_pAG32-F/R). Cells with correct repeat length were grown to approximately OD$_{600}$
of 0.5 before being transferred to a concanavalin A-coated chamber slide for attachment. From
there, cells were synchronized by addition of alpha factor (1µM final concentration) then
released by washing 3X with YPD media. A single field of cells was imaged with DeltaVision
Ultra High Resolution microscope using a 100X objective using optical axis integration every
1.5s for 5 minutes. This was repeated using cells in different areas of the slide until a total of 100
minutes of timepoints were taken. Exposure was: Red 400ms, Green 400ms, Pol 50ms (every
50th frame). Images were analyzed with CellProfiler, tracking the LacO/LacI-mCherry focus and
correcting for nuclear translation by tracking Nup49-GFP. Tracked foci were quantified using
MSD $= < [\![(x^{\rightarrow} - (x\_dt)^{\rightarrow})]\!]^2 >$ and radius of constraint = sqrt(5/4 x MSD plateau) (Dion &
Gasser, 2013). The plateau was calculated by regression in Graphpad PRISM software using the
first 100 Δt timepoints.
929
930 **Quantification and statistical analysis**
931 Prism software was used to calculate statistical significance. For microscopy experiments
932 Fisher's exact test (Relocation experiments) or chi square test (DIM experiments) were used.
933
934 **Data and software availability-** All raw data is available in the supplemental tables

A

Contrast Adjust

| | CFP-LacI | mVenus-Tub1 | Nup49-mCherry | Merge | mVenus-Tub1 | Merge |

DIM, Repeat @ NPC

No DIM

DIM, Repeat @ NPC

DIM, Repeat not @ NPC

DIM, Repeat not @ NPC

DIM, Repeat not @ NPC

1 µm

B

t=0   t=5s   t=10s   t=15s   t=20s   t=25s

Raw

Contrast adjusted

1 µm